

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

## Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

#### **About Google Book Search**

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/



Это цифровая коиия книги, хранящейся для иотомков на библиотечных иолках, ирежде чем ее отсканировали сотрудники комиании Google в рамках ироекта, цель которого - сделать книги со всего мира достуиными через Интернет.

Прошло достаточно много времени для того, чтобы срок действия авторских ирав на эту книгу истек, и она иерешла в свободный достуи. Книга иереходит в свободный достуи, если на нее не были иоданы авторские ирава или срок действия авторских ирав истек. Переход книги в свободный достуи в разных странах осуществляется ио-разному. Книги, иерешедшие в свободный достуи, это наш ключ к ирошлому, к богатствам истории и культуры, а также к знаниям, которые часто трудно найти.

В этом файле сохранятся все иометки, иримечания и другие заииси, существующие в оригинальном издании, как наиоминание о том долгом иути, который книга ирошла от издателя до библиотеки и в конечном итоге до Вас.

### Правила использования

Комиания Google гордится тем, что сотрудничает с библиотеками, чтобы иеревести книги, иерешедшие в свободный достуи, в цифровой формат и сделать их широкодостуиными. Книги, иерешедшие в свободный достуи, иринадлежат обществу, а мы лишь хранители этого достояния. Тем не менее, эти книги достаточно дорого стоят, иоэтому, чтобы и в дальнейшем иредоставлять этот ресурс, мы иредириняли некоторые действия, иредотвращающие коммерческое исиользование книг, в том числе установив технические ограничения на автоматические заиросы.

Мы также иросим Вас о следующем.

- Не исиользуйте файлы в коммерческих целях. Мы разработали ирограмму Поиск книг Google для всех иользователей, иоэтому исиользуйте эти файлы только в личных, некоммерческих целях.
- Не отиравляйте автоматические заиросы.

Не отиравляйте в систему Google автоматические заиросы любого вида. Если Вы занимаетесь изучением систем машинного иеревода, оитического расиознавания символов или других областей, где достуи к большому количеству текста может оказаться иолезным, свяжитесь с нами. Для этих целей мы рекомендуем исиользовать материалы, иерешедшие в свободный достуи.

- Не удаляйте атрибуты Google.
  - В каждом файле есть "водяной знак" Google. Он иозволяет иользователям узнать об этом ироекте и иомогает им найти доиолнительные материалы ири иомощи ирограммы Поиск книг Google. Не удаляйте его.
- Делайте это законно.
  - Независимо от того, что Вы исиользуйте, не забудьте ироверить законность своих действий, за которые Вы несете иолную ответственность. Не думайте, что если книга иерешла в свободный достуи в США, то ее на этом основании могут исиользовать читатели из других стран. Условия для иерехода книги в свободный достуи в разных странах различны, иоэтому нет единых иравил, иозволяющих оиределить, можно ли в оиределенном случае исиользовать оиределенную книгу. Не думайте, что если книга иоявилась в Поиске книг Google, то ее можно исиользовать как угодно и где угодно. Наказание за нарушение авторских ирав может быть очень серьезным.

## О программе Поиск кпиг Google

Muccus Google состоит в том, чтобы организовать мировую информацию и сделать ее всесторонне достуиной и иолезной. Программа Поиск книг Google иомогает иользователям найти книги со всего мира, а авторам и издателям - новых читателей. Полнотекстовый иоиск ио этой книге можно выиолнить на странице http://books.google.com/

The Lopkins Library presented to the Celand Stanford Junior Amversity by Cimuthy Kupkins.



at and



# THE ZOOLOGICAL RECORD.

-----

THE object of the Zoological Record is to give, by means of an annual Volume, complete lists of the Works and Publications relating to Zoology in all its branches that have appeared during the year preceding the date of the Volume; together with full information as to the points they deal with, arranged in such a manner as to serve as an Index to the literature of Zoology in all parts of the globe, and thus to form a repertory which will retain its value for the Student in future years.

The 'Zoological Record' is published for the Society by Messrs. Gurney and Jackson at the price of 80s. per volume. But all Members of the Zoological Society of London have the privilege of receiving it, including the cost of delivery (within the United Kingdom), at a subscription price of 20s. per annum. This Subscription is due on the 1st of July in every year, but the privilege of Subscription is forfeited unless the amount be paid before the 1st of December following.

The Zoological Society, having purchased the entire stock of the 'Zoological Record,' are able to supply complete sets of the first twenty-two volumes at the price of £5 10s. Volumes of any single year (exclusive of the last two volumes and Vol. 6) can likewise be supplied at 10s, per volume.

Members of the Society wishing to subscribe to the 'Record' are requested to apply at this office for a Form, to be returned when filled up and signed by the subscriber. In order to facilitate the payment of the subscription, a Banker's Order Form is also furnished to those who prefer that mode of payment. This order, when filled up and signed, should be sent to the Society's office for registration; it will then be sent to the Agents named therein.

Learned Societies and Institutions and members of the former Zoological Record Association are permitted to subscribe to the 'Record' on the same conditions as are accorded to Members of the Zoological Society.

P. L. SCLATER,

Secretary.

Oct. 1st, 1893.

ZOOLOGICAL SOCIETY OF LONDON,

3, HANOVER SQUARE, W.

### LIST OF VOLUMES

OF THE

## 'ZOOLOGICAL RECORD.'

The Record of Zoological Literature, 1864. Volume First. Edited by Albert C. L. G. GUNTHER, M.A., M.D., Ph.D., F.Z.S., &c. London, 1865. Price 10s.

The Record of Zoological Literature, 1865. Volume Second. Edited by Albert C. L. G. GUNTHER, M.A., M.D., Ph.D., F.Z.S., &c. London, 1866. Price 10s.

The Record of Zoological Literature, 1866. Volume Third. Edited by Albert C. L. G. GÜNTHER, M.A., M.D., Ph.D., F.R.S., F.Z.S., &c. London, 1867. Price 10s.

The Record of Zoological Literature, 1867. Volume Fourth. Edited by Albert C. L. G. Gunther, M.A., M.D., Ph.D., F.R.S., F.Z.S., &c. London, 1868. Price 10s.

The Record of Zoological Literature, 1868. Volume Fifth. Edited by Albert C. L. G. GÜNTHER, M.A., M.D., Ph.D., F.R.S., F.Z.S., &c. London, 1869. Price 10s.

The Record of Zoological Literature, 1869. Volume Sixth. Edited by Albert C. L. G. GUNTHER, M.A., M.D., Ph.D., F.R.S., F.Z.S., &c. London, 1870. Price 80s.

The Zoological Record for 1870; being Volume Seventh of the Record of Zoological Literature. Edited by ALFRED NEWTON, M.A., F.R.S., Professor of Zoology and Comparative Anatomy in the University of Cambridge, F.L.S., V.P.Z.S, &c. London, 1871. Price 10\*.

The Zoological Record for 1871; being Volume Eighth of the Record of Zoological Literature. Edited by ALFRED NEWTON, M.A., F.R.S., Professor of Zoology and Comparative Anatomy in the University of Cambridge, F.L.S., V.P.Z.S., &c. London, 1873. Price 10s.

The Zoological Record for 1872; being Volume Ninth of the Record of Zoological Literature. Edited by Alfred Newton, M.A., F.R.S., Professor of Zoology and Comparative Anatomy in the University of Cambridge, F.L.S., V.P.Z.S., &c. London, 1874. Price 10s.

The Zoological Record for 1873; being Volume Tenth of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., Librarian to the Royal Geographical Society. London, 1875. Price 10s.

The Zoological Record for 1874; being Volume Eleventh of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., M.E.S., Editor Ent. M. Mag., Librarian to the Royal Geographical Society. London, 1876. Price 10s.

The Zoological Record for 1875; being Volume Twelfth of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., M.E.S., Editor Ent. M. Mag., Librarian to the Royal Geographical Society. London, 1877. Price 10s.

The Zoological Record for 1876; being Volume Thirteenth of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., M.E.S., Editor Ent. M. Mag., Librarian to the Royal Geographical Society. London, 1878. Price 10s.

The Zoological Record for 1877; being Volume Fourteenth of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., M.E.S., Editor Ent. M. Mag., Librarian to the Royal Geographical Society. London, 1879. Price 10s.

The Zoological Record for 1878; being Volume Fifteenth of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., M.E.S., Editor Ent. M. Mag., Librarian to the Royal Geographical Society. London, 1880. Price 10s.

The Zoological Record for 1879; being Volume Sixteenth of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., M.E.S., Editor Ent. M. Mag., Librarian to the Royal Geographical Society. London, 1881. Price 10s.

The Zoological Record for 1880; being Volume Seventeenth of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., M.E.S., Editor Ent. M. Mag., Librarian to the Royal Geographical Society. London, 1881. Price 10s.

The Zoological Record for 1881; being Volume Eighteenth of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., M.E.S., Editor Ent. M. Mag., Librarian to the Royal Geographical Society. London, 1882. Price 10s.

The Zoological Record for 1882; being Volume Nineteenth of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., M.E.S., Editor Ent. M. Mag., Librarian to the Royal Geographical Society. London, 1888. Price 10s.

The Zoological Record for 1883; being Volume Twentieth of the Record of Zoological Literature. Edited by Edward Caldwell Rye, F.Z.S., M.E.S., Editor Ent. M. Mag., Librarian to the Royal Geographical Society. London, 1884. Price 10s. The Zoological Record for 1884; being Volume the Twenty-first of the Record of Zoological Literature. Edited by F. Jeffrey Bell, M.A., Sec. R.M.S., Professor of Comparative Anatomy and Zoology in King's College, London. London, 1885. Price 10s.

The Zoological Record for 1885; being Volume the Twenty-second of the Record of Zoological Literature. Edited by F. Jeffrey Bell, M.A., Sec. R.M.S., Professor of Comparative Anatomy and Zoology in King's College, London. London, 1886. Price 10s.

The Zoological Record for 1886; being Volume the Twenty-third of the Record of Zoological Literature. Edited by Frank E. Beddard, M.A., F.Z.S., Prosector and Davis Lecturer to the Zoological Society of London. London, 1887. Price 10s.

The Zoological Record for 1887; being Volume the Twenty-fourth of the Record of Zoological Literature. Edited by Frank E. Beddard, M.A., F.Z.S., Prosector and Davis Lecturer to the Zoological Society of London. London, 1888. Price 10s.

The Zoological Record for 1888; being Volume the Twenty-fifth of the Record of Zoological Literature. Edited by Frank E. Beddard, M.A., F.Z.S., Prosector and Davis Lecturer to the Zoological Society of London. London, 1890. Price 10s.

The Zoological Record for 1889; being Volume the Twentysixth of the Record of Zoological Literature. Edited by Frank E. Beddard, M.A., F.Z.S., Prosector and Davis Lecturer to the Zoological Society of London. London, 1890. Price 10s.

The Zoological Record for 1890; being Volume the Twenty-seventh of the Record of Zoological Literature. Edited by Frank E. Beddard, M.A., F.Z.S., Prosector and Davis Lecturer to the Zoological Society of London. London, 1892. Price 10s.

The Zoological Record for 1891; being Volume the Twenty-eighth of the Record of Zoological Literature. Edited by David Sharp, Esq., M.D., F.R.S., F.Z.S. London, 1892. Price 80s.

The Zoological Record for 1892; being Volume the Twentyninth of the Record of Zoological Literature. Edited by David Sharp, Esq., M.D., F.R.S., F.Z.S. London, 1898. Price 80s.

These publications may be obtained at the SOCIETY'S OFFICE (3 Hanover Square, W.), of Messrs. Gurney and Jackson (Paternoster Row, E.C.), or through any bookseller.

# ZOOLOGICAL RECORD,

VOLUME THE TWENTY-NINTH.

BEING

## RECORDS OF ZOOLOGICAL LITERATURE

RELATING CHIEFLY TO THE YEAR

## 1892.

By J. A. Thomson, R. Lydekker, R. Bowdler Sharpe, G. A. Boulenger, W. A. Herdman, B. B. Woodward, R. I. Pocock, D. Sharp, F. A. Bather, Florence Buchanan, S. J. Hickson, and R. Hanitsch.

#### EDITED BY

D. SHARP, M.A., F.R.S., F.Z.S., &c., curator in zoology, university museum of zoology and comparative anatomy, cambridge.

Explorate solum: sic fit via certior ultrà.

LONDON:
GURNEY & JACKSON, PATERNOSTER ROW.
M.DCCC.XCIII.



141394

Communications, Papers, and Memoirs intended for this work should be addressed to "THE EDITOR of the Zoological Record, Zoological Society, 3, Hanover Square, London, W." It is earnestly requested that in the case of separately-printed copies of papers so forwarded, the original pagination be indicated.

The Editor thanks those Societies and individuals who have forwarded works and papers for the purposes of the Record, and suggests that Societies publishing Zoological Papers would much facilitate the work of the Recorders by forwarding a separate copy of each Zoological Memoir they publish to the Editor for the use of the Recorder.

## PREFACE.

Owing to the punctuality of most of the Recorders the twentyninth volume of the Zoological Record will be in the hands of the subscribers at a date that may, it is hoped, be considered satisfactory.

The names of three new Recorders appear on the title page, and the Editor feels it to be a duty, as well as a pleasure, to thank all the Contributors, old and new, for their assistance. The three new Recorders, Miss Florence Buchanan, Mr. F. O. Bather, and Dr. R. Hanitsch, have carried out their work in an efficient manner, though it is much to be regretted that Mr. Bather was prevented by a weakness of the eyes from completing his examination of the literature relating to *Echinodermata*. Mr. J. A. Thomson, being desirous of facilitating reference in the very extensive and difficult department he deals with, has drawn up a complete Subject Record in addition to a separate List of Works and Essays.

On comparing this twenty-ninth volume with those of the earlier part of the series, it will be noticed that the scope of the Record has become much enlarged. The Zoology of our days is an aggregate of a number of special departments of observation and research; hence, it is no easy task to furnish a complete Record, and at the same time to arrange it so as to meet the wishes of the various persons who may consult it from different points of view. In order to make the work as widely useful as possible, it appears to be necessary to give in each Record an Index of Special Subjects in addition to the List of Titles and the taxonomical arrangement according to genera; and this the Editor hopes may prove to be

feasible. Abstracts of, and Reports on, the Memoirs recorded will, as a consequence, be reduced to the narrowest dimensions.

As completeness should be the most indispensable of the characteristics of the work, there is in each of the Records one department it is impossible to speak of with satisfaction, viz.: Palæontology. When the Record was established it was not intended to include this subject. At the present time some of the Recorders examine the Geological literature of each year, and give a nearly complete epitome of the Palæontology they meet with, while others are content with a less exhaustive treatment of the subject.

To make the Palæontology complete, and to give a stratigraphical Index to it in each of the eighteen Records of which our volume consists, is a course that would add considerably to the labours of the Recorders and somewhat to the extent of the Volume, and cannot, therefore, in the present circumstances of the Record, be adopted. Indeed, it would be preferable for the Palæontologists to undertake a separate Record, and if no other satisfactory arrangement can be made, it may be hoped that they will do so, and thus relieve the Zoological Recorders from a task which they only imperfectly carry out.

D. SHARP.

CAMBRIDGE, Sept. 1, 1893.

## LIST OF THE ABBREVIATED TITLES

#### OF THE PRINCIPAL

# JOURNALS AND OF THE TRANSACTIONS OF LEARNED SOCIETIES

#### THAT CONTAIN ZOOLOGICAL PAPERS.

- Abh. Ak. Berl.—Abhandlungen der königlich Akademie der Wissenschaften zu Berlin. (Also SB.)
- Abh. Bayer. Ak.—Abhandlungen der mathematisch-physikalischen Classe der k. Bayerischen Akademie der Wissenschaften (Munich). (Also SB.)
- Abh. Böhm. Ges. Abhandlungen der mathematisch-naturwissenschaftlichen Classe der k. Böhmischen Gesellschaft der Wissenschaften (Prague). (Called also Rozpravy trídy mathematicko-přírodovědecké královske české společnosti nauk.) (Also SB.)
- Abh. Ges. Götting.—Abhandlungen der k. Gesellschaft der Wissenschaften zu Göttingen.
- Abh. Ges. Hulle—Abhandlungen der naturforschenden Gesellschaft in Halle. (Also Ber.)
- Abh. Ges. Isis—Abhandlungen der naturwissenschaftlichen Gesellschaft 'Isis' in Dresden. (See SB.)
- Abh. Ges. Königsb.—Abhandlungen der k. physikalisch-ökonomischen Gesellschaft in Preussen (Königsberg). (Also SB.)
- Abh. naturf. Ges. Görlitz—Abhandlungen der naturforschenden Gesellschaft zu Görlitz.
- Abh. naturh. Ges. Nürnberg—Abhandlungen der naturhistorischen Gesellschaft zu Nürnberg (Nürnberg).
- Abh. Sächs. Ges.—Abhandlungen der k. Sächsischen Gesellschaft der Wissenschaften (Leipzig). (Also Ber.)
- Abh. Schw. pal. Ges.—Abhandlungen der Schweizerischen paläontographischen Gesellschaft (Båle).
- Abh. Senck. Ges.—Abhandlungen herausgegeben von der Senckenbergischen naturforschenden Gesellschaft (Frankfort). (Also Ber.)

- Abh. Ver. Brem.—Abhandlungen herausgegeben vom naturwissenschaftlichen Verein zu Bremen.
- Abh. Ver. Hamb.—Abhandlungen aus dem Gebiete der Naturwissenschaften herausgegeben vom naturw. Verein in Hamburg.
- Abh. zool. Mus. Dresden—Abhandlungen und Berichte des k. zoologischen, etc., Museums in Dresden.
- Act. Ac. Bordeaux—Actes de l'Académie nationale des Sciences, Belles Lettres et Arts de Bordeaux.
- Act. Ac. Córdob.—Actas de l'Academia nacional de Ciencias en Córdoba (Buenos Ayres).
- Act. Lund.—Acta Universitatis Lundensis (Lund). (Called also Lunds Universitets Årsskrift.)
- Act. Soc. Chili-Actes de la Société scientifique du Chili.
- Act. Soc. Fenn.—Acta Societatis Scientiarum Fennica (Helsingfors).
- Act. Soc. Helv.—Actes de la Société Helvétique des Sciences naturelles. (Called also Verhandl. d. schweiz. Naturforsch., Gesells.)
- Act. Soc. Jura.—Actes de la Société Jurassienne d'Émulation,
- Act. Soc. L. Bord.—Actes de la Société Linnéenne de Bordeaux. (Also Comptes rendus.)
- Act. Upsala.—Acta Universitatis Upsalensis, (Called also Upsala Universitets Årsskrift.)
- Agric. Gaz. N.S.W. Agricultural Gazette of New South Wales (Sydney).
- Am. Geol.—The American Geologist (Calvin et alii: Minneapolis).
- Am. J. Sci.—American Journal of Science (New Haven).
- Am. Micr. J.—American Monthly Microscopical Journal (Washington).
- Am. Nat.-The American Naturalist (Philadelphia).
- Anat. Anz.—Anatomischer Anzeiger (Bardeleben: Jena).
- An. Mus. B. Aires—Anales del Museo nacional, Buenos Aires.
- An. Mus. Costa Rica—Anales del Museo nacional, Republica de Costa Rica (San José).
- An. Mus. La Plata—Anales del Museo, La Plata (Buenos Ayres).
- An. Mus. nuc. Mexico-Anales del Museo nacional de México.
- An. Soc. Arg.—Anales de la Sociedad Cientifica Argentina (Buenos Ayres).
- An. Soc. Esp.—Anales de la Sociedad Española de Historia Natural (Madrid).
- Ann. Ent. Belg.—Annales de la Société Entomologique de Belgique (Brussels).
- Ann. Fac. Marseille—Annales de la Faculté des Sciences de Marseille, publiées sous les auspices de la municipalité (Marseille & Paris).
- Ann. Géol. univ. Paris—Annuaire Géologique universel, Revue de Géologie et Paléontologie (Carey & Douvillé).
- Ann. Hofmuseum Wien—Annalen des k. k. naturhistorischen Hofmuseums (von Hauer: Vienna).

- Ann. Mal.—Annales de Malacologie (Servain : Paris).
- Ann. Micrograph.—Annales de Micrographie spécialement consacrées à la Bactériologie aux Protophytes et aux Protozoaires (Miquel : Paris).
- Ann. Mus. Belg.—Annales du Musée royal d'Histoire Naturelle de Belgique (Brussels).
- Ann. Mus. Genov.—Annali del Museo civico di Storia Naturale di Genova (Genoa).
- Ann. Mus. Marseille—Annales du Musée d'Histoire Naturelle de Marseille.
  Zoologie (Marion: Marseilles).
- Ann. N. H.—Annals and Magazine of Natural History (London).
- Ann. N. York Ac.—Annals of the New York Academy of Sciences.
  (Also Trans.)
- Ann. Queensland Mus.—Annals of the Queensland Museum (Brisbane).
- Ann. Sci. Géol.—Annales des Sciences Géologiques (Hébert & Milne-Edwards : Paris).
- Ann. Sci. Nat.—Annales des Sciences Naturelles. Zoologie. (Paris).
- Ann. Scot. N. H .- The Annals of Scottish Natural History.
- Ann. Soc. Agric. Lyon—Annales de la Société d'Agriculture, Histoire Naturelle, et Arts utiles de Lyon (Lyons & Paris).
- Ann. Soc. Belg. Micr.—(Also Bull. Soc. Belg. Micr., q.v.)
- Ann. Soc. Brux.—Annales de la Société Scientifique de Bruxelles (Brussels).
- Ann. Soc. Char.—Annales de la Société des Sciences Naturelles de la Charente Inférieure (= Academie de la Rochelle).
- Ann. Soc. Ent. Fr.—Annales de la Société Entomologique de France (Paris).
- Ann. Soc. Géol. Belg.—Annales de la Société Géologique de Belgique (Liége).
- Ann. Soc. Géol. Nord—Annales de la Société Géologique du Nord (Lille).
  (Also Mémoires.)
- Ann. Soc. L. Lyon (n.s.)—Annales de la Société Linnéenne de Lyon. Nouvelle série.
- Ann. Soc. Mal. Belg.—Annales de la Société Malacologique de Belgique (Brussels).
- Ann. Univ. Lyon-Annales de l'université de Lyon (Paris).
- Ann. Univ. Toscane—Annali delle Università Toscane (Pisa).
- Ant. Annual—The Antananarivo Annual and Madagascar Magazine (Sibree: Antananarivo).
- Anz. Ak. Wien—Anzeiger der mathematisch-naturwissenschaftlichen Classe der k. Akademie der Wissenschaften zu Wien (Vienna).
- Appalachia—Appalachia: the Journal of the Appalachian Mountain Club (Boston).
- Arb. Inst. Würzb.—Arbeiten aus dem zoologisch-zootomischen Institute in Würzburg.

- Arb. z. Inst. Wien—Arbeiten aus dem zoologischen Institute der Universität Wien (Vienna).
- Arb. z. Inst. Graz—Arbeiten aus dem zoologischen Institute zu Graz (Leipsic).
- Arch. Anat. Phys.—Archiv für Anatomie und Physiologie (His, Braune, & Du Bois Reymond : Leipzig).
- Arch. Biol.—Archives de Biologie (Van Beneden & Van Bambeke : Ghent).
- Arch. d'Anthrop.—Archives d'Anthropologie (Toulouse).
- Arch. f. Anthrop.—Archiv für Anthropologie: Zeitschrift für Naturgeschichte und Urgeschichte des Menschen (Brunswick).
- Arch. f. Math. og Naturr.—Archiv for Mathematik og Naturvidenskab (Lie, G. O. Sars: Kristiania).
- Arch. f. Nat.-Archiv für Naturgeschichte. Neue Folge (Berlin).
- Arch. f. Thierheilk.—Archiv für Thierheilkunde.
- Arch. ges. Phys.—Archiv für die gesammte Physiologie des Menschen und der Thiere (Pflüger : Bonn).
- Arch. Ital. Biol.—Archives Italiennes de Biologie; Revues, Résumés, Reproductions des travaux scientifiques Italieus (Emery & Mosso: Turin).
- Arch. mikr. Anat .-- Archiv für mikroskopische Anatomie (Bonn).
- Arch. Miss. Sci.—Archives des Missions Scientifiques et Litéraires (Paris).
- Arch. Mus. Lyon-Archives du Muséum d'Histoire Naturelle de Lyon.
- Arch. Mus. R. Jan.—Archivos do Museu nacional do Rio de Janeiro.
- Arch. Mus. Teyl. -Archives du Musée Teyler (Haarlem).
- Arch. Nat. Liv.—Archiv für die Naturkunde Liv-, Esth-, und Kurlands (Dorpat).
- Arch. naturw. Landesforsch. Böhmen—Archiv für naturwissenschaftliche Landesdurchforschung von Böhmen (Prag).
- Arch. Néerl.—Archives Néerlandaises des Sciences Exactes et Naturelles . (Bosscha: Haarlem).
- Arch. Phys.—Archives de Physiologie Normale et Pathologique (Brown-Séquard : Paris).
- Arch. Sci. Nat.—Archives des Sciences Physiques et Naturelles (Geneva).
- Arch. Ver. Mecklenb.—Archiv des Vereins der Freunde der Naturgeschichte in Mecklenburg.
- Arch. Zeeuwsch Genoots. Wetensch.—Archief vroegere en latere Mededeelingen voornamelijk in Betrekking tot Zeeland uitgegeven door het Zeeuwsch Genootschap der Wetenschappen (Middelburg).
- Arch. Z. expér.—Archives de Zoologie expérimentale et générale (Lacaze-Duthiers: Paris).
- Atti Acc. Gioen.—Atti dell' Accademia Gioenia di Scienze Naturali (Catania).

- Atti Acc. Napoli—Atti della B. Accademia delle Scienze Fisiche e Matematiche.
- Atti Acc. Palermo—Atti della R. Accademia di Scienze, Lettere é Belle Arti di Palermo (Palermo).
- Atti Acc. Pontaniana-Atti dell' Accademia Pontaniana (Naples).
- Atti Acc. Pontif. Lincei.—Atti dell' Accademia Pontificia de' nuovi Lincei.
- Atti Acc. Tor .- Atti della R. Accademia delle Scienze di Torino (Turin).
- Atti Ist. Nap.—Atti del R. Istituto d'incorraggiamento alle Scienze Naturali Economichi e Technologiche, &c., di Napoli (Naples).
- Atti Ist. Venet.—Atti del R. Istituto Veneto di Scienze, Lettere et Arti, &c. (Venice).
- Atti [Mem. Rend.] Acc. Rom.—Atti [Memorie: Rendiconti] della R. Accademia dei Lincei (Rome).
- Atti [Mem.] Soc. Tosc.—Atti [Memorie] della Società Toscana di Scienze Naturali residente in Pisa.
- Atti [Rend. Mem.] Soc. Mod.—Atti [Memorie: Rendiconti] della Società dei Naturalisti di Modena.
- Atti Soc. Ital.—Atti della Società Italiana di Scienze Naturali (Milan).
- Atti Soc. Ligust.—Atti della Società Ligustica di scienze naturali e geografiche (Genova).
- Atti Soc. Ven.-Trent.—Atti della Società Veneto-Trentina di Scienze Naturali residente in Padova (Padua). (Also Bull.)
- Atti Univ. Genova-Atti della R. Università di Genova.
- Auk—The Auk. A Quarterly Journal of Ornithology. (Continuation of the Bulletin of the Nuttall Ornithological Club.)
- Ausland-Das Ausland (Stuttgart).
- Beitr. Morphol. Morphog.—Beiträge zur Morphologie und Morphogenie (Gerlach: Stuttgart).
- Beitr. Pal. Oesterr.-Ung. Beiträge zur Paläontologie Oesterreich-Ungarn's und des Orients (Mojsisovics & Neumayr: Vienna).
- Beitr. Russ. Reiches (2)—Beiträge zur Kenntniss des Russichen Reiches und der angrerzenden Länder Asiens. Neue Folge (Schrenck & Maximowics: St. Petersburg).
- Ber. deuts. botan. Ges.—Bericht der deutschen botanischen Gesellschaft (Berlin).
- Ber. deutsch. orn. Ges.—Bericht der allgemeines deutsche ornithologische Gesellschaft zu Berlin.
- Ber. Ges. Freiburg—Berichte der naturforschenden Gesellschaft zu Freiburg (Freiburg, i Br.).
- Bergens Mus. Aarsber Bergens Museum Aarsberetning (Bergen).
- Ber. Ges. Chemn.—Bericht der naturwissenschaftlichen Gesellschaft zu Chemnitz.

- Ber. Ges. Halle—Bericht über die Sitzungen der naturforschenden Gesellschaft zu Halle. (Also Abhandl.)
- Ber. Ges. Hanau = JB. wetter, Ges.
- Ber. Komm. wiss. Unters. deutsch. Meere—Bericht der Kommission zur Untersuchung der deutschen Meere.
- Ber. Naturf. Ärtste-Ämtliche Bericht deutscher Naturforscher und Ärtste.
- Ber. naturf. Ges. Bamberg—Bericht der naturforschenden Gesellschaft in Bamberg.
- Ber. naturhist. Mus. Hamburg—Bericht des naturhistorischen Museums in Hamburg.
- Ber. natur. Ver. Passau-Bericht des naturhistorischen Vereins in Passau.
- Ber. naturw. Ver. Regensburg Bericht der naturwissenschaftlichen Vereins in Regensburg. (Formerly CB.).
- Ber. Oberhess. Ges.—Bericht der Oberhessischen Gesellschaft für Naturund Heilkunde (Giessen).
- Ber. Offenb. Ver.—Bericht über die Thätigkeit des Offenbacher Vereins für Naturkunde (Offenbach-on-the-Main).
- Ber. Sächs. Ges.—Bericht ueber die Verhandlungen der königlichen Sächs. Gesellschaft der Wissenschaft in Leipzig. (Also Abhandl.)
- Ber. Senck. Ges.—Bericht der Senckenbergische naturforschende Gesellschaft im Frankfurt am Main. (Also Abhandl.)
- Ber. St. Gall. Ges.—Bericht über die Thätigkeit der St. Gallischen naturwissenschaftlichen Gesellschaft (St. Gallen).
- Ber. Ver. Cassel-Berichte des Vereins für Naturkunde zu Cassel.
- Ber. Ver. Pass. Bericht der naturwissenschaftlichen Vereins in Passau.
- Ber. Ver. Schwaben—Bericht der naturwissenschaftlichen Vereins für Schwaben und Neuburg (a. V.), früher naturhistorischen Vereins Augsburg (Augsburg).
- Berl. Monats.- Berliner Monatchefte.
- B. E. Z.—Berliner Entomologische Zeitschrift.
- Bibl. haut. études.—Bibliothèque de l'école des hautes études. Section des Sciences Naturelles (Paris).
- Bibl. univ.—Bibliothèque universelle et Revue Suisse (Geneva). (See Arch. Sci. Nat.)
- Bibl. Zool.—Bibliotheca Zoologica (Leipsic).
- Bidr. Finl. Nat.—Bidrag till Kännedom af Finlands Natur och Folk (Helsingfors).
- Bik. Sv. Ak. Handl.—Bihang till K. Svenska Vetenskaps-Akademiens Handlingar (Stockholm).
- Bijdr. Dierk.—Bijdragen tot de Dierkunde (Amsterdam).
- Biol. Centralbl.—Biologisches Centralblatt (Rosenthal: Erlangen).
- Biol. Centr. Am.—Biologia Centrali Americana (Godman & Salvin: London).

- Biol. Fören.—Biologiska Föreningens Förhandlingar. Verhandlungen des biologischen Vereins in Stockholm. (Figerstedt: Stockholm.)
- Bol. Ac. Arg.—Boletin de la Academia nacional de Ciencias de la Republica Argentina (Cordoba).
- Bol. yeol. S. Paulo—Boletim da Commissão geographica e geologica da Provincia de S. Paulo.
- Bol. Mus. la Plata-Boletin del Museo la Plata (Buenos Ayres).
- Bol. Com. Geol.—Bollettino del R. Comitato Geologico d'Italia (Roma).
- Boll. Mus. Zool. Anat. Comp. Torino—Bollettino dei Musei di Zoologia ed Anatomia comparata della R. Università di Torino.
- Boll. scient.—Bollettino scientifico (Maggi, Zoja, & Giovanni: Pavia).
- Boll. Soc. Adr.—Bollettino della Società Adriatica di Scienze naturali (Trieste).
- Boll. Sec. geol. Ital.—Bollettino della Società geologica Italiana (Rome).
- Boll. Soc. Nat. Napoli Bollettino della Società di Naturalisti in Napoli. (Formerly Rivista italiana di scienze naturali [Circolo degli aspiranti naturalisti].)
- Boll. Soc. Rom. Stud. Zool.—Bollettino della Società Romana per gli Studi Zoologici.
- Bot. Z.—Botanische Zeitung (Halle).
- Brit. Nat.—The British Naturalist (Robson: London).
- Bull. Ac. Belg.—Bulletin de l'Académie royale des Sciences, des Lettres, et des Beaux Arts de Belgique (Brussels). (Also Mem.)
- Bull. Ac. Cracovie—Bulletin international de l'Académie des sciences de Cracovie (Cracovie).
- Bull. Ac. Hippone—Bulletin de l'Académie d'Hippone (Bône).
- Bull. Am. Mus. Nat. Hist.—Bulletin of the American Museum of Natural History (New York).
- Bull. Brit. Ornith. Club-Bulletin of the British Ornithologists' Club.
- Bull. Brookville Soc.—Bulletin of the Brookville Society of Natural History (Brookville, Indiana, U.S.A.).
- Bull. Buff. Soc.—Bulletin of the Buffalo Society of Natural Sciences.
- Bull. Bussey Inst.—Bulletin of the Bussey Institution (Boston).
- Bull. Cornell Univ.—Bulletin of the Cornell University. (Ithaca.)
- Bull. Denison Univ.—Bulletin of the Scientific Laboratories of Denison Universities (Granville, Ohio).
- Bull. Dep. Agric. Ent.—U.S. Department of Agriculture. Division of Entomology. Bulletin (Washington).
- Bull. Ent. Ital.—Bullettino della Società Entomologica Italiana (Florence).
- Bull. Ess. Inst.—Bulletin of the Essex Institute (Salem, U.S.A.).
- Bull. Geol. Soc. Amer.—Bulletin of the Geological Society of America (New York).

- Bull. Illin. Lub. N. H.—Bulletin of the Illinois State Laboratory of Natural History. (Champaign, Illinois.)
- Bull. Inst. Nat. Genevois-Bulletin de l'Institut nationale Genevois.
- Bull. Lab. Iowa—Bulletin from the Laboratories of Natural History of the State University of Iowa (Iowa city).
- Bull. Minnesota Acad.—Bulletin of the Minnesota Academy of Natural Sciences.
- Bull. Mosc.—Bulletin de la Société impériale des Naturalistes de Moscou (Menzbier : Moscow).
- Bull. Mus. Belg.—Bulletin du Musée royal d'Histoire Naturelle de Belgique (Brussels).
- Bull. Mus. C. Z.—Bulletin of the Museum of Comparative Zoology of Harvard College (Cambridge, U.S.A.).
- Bull. Nat. Hist. Soc. New Brunswick—Bulletin of the Natural History Society of New Brunswick (St. John's, N.B.).
- Bull. New York Mus. Nat. Hist.—Bulletin of the New York State Museum of Natural History (Albany).
- Bull. Ohio Exp. Station—Bulletin of the Ohio Agricultural Experiment Station, Technical Series (Columbus).
- Bull. Pétersb.—Bulletin de l'Académie impériale des Sciences de St. Pétersbourg.
- Bull. Phil. Soc. Wash.—Bulletin of the Philosophical Society of Washington.
- Bull. Sci. Fr. Belg.—Bulletin Scientifique de la France et de la Belgique (Giard: Paris).
- Bull. Soc. Ac. Brest—Bulletin de la Société Academique de Brest.
- Bull. Soc. Acclim.—Bulletin mensuel de la Société nationale d'Acclimatation de Paris (Paris).
- Bull. Soc. Angers—Bulletin de la Société d'Études scientifiques d'Angers (Angers).
- Bull. Soc. Anthrop. Lyon—Bulletin de la Société d'Anthropologie de Lyon.
- Bull. Soc. Anthrop. Paris—Bulletin de la Société d'Anthropologie de Paris.
- Bull. Soc. Aude—Bulletin de la Société d'Études scientifiques de l'Aude (Carcassoune).
- Bull. Soc. Autun-Société d'Histoire Naturelle d'Autun.
- Bull. Soc. Belg. Micr.—Bulletin de la Société Belge de Microscopie (Brussels). (Also Annales.)
- Bull. Soc. Béziers—Bulletin de la Société d'Étude des Sciences Naturelles de Béziers. Comptes rendus des Séances (Béziers).
- Bull. Soc. Borda-Dax—Bulletin de la Société de Borda (Dax).
- Bull. Soc. Colm. Bulletin de la Société d'Histoire Naturelle de Colmar.

- Bull. Soc. d'Elbeuf.—Bulletin de la Société d'étade des sciences naturelles d'Elbeuf.
- Bull. Soc. Dinan.—Bulletin de la Société des Naturalistes Dinantais (Dinant).
- Bull. Soc. Ent. Suisse-See MT. Schw. ent. Ges.
- Bull. Soc. Ent. Fr.—Bulletin des séances de la Société Entomologique de France (Paris) (See Ann.).
- Bull. Soc. Finistère—Bulletin de la Société d'Etudes scientifiques du Finistère (Morlaix).
- Bull. Soc. Geogr.—Bulletin de la Société de Geographie (Paris).
- Bull. Soc. Géol.—Bulletin de la Société Géologique de France (Paris).
- Bull. Soc. Isère—Bulletin de la Société de statistique des sciences naturelles et des arts industriels du département de l'Isère (Grenoble).
- Bull. Soc. L. Bruxelles—Bulletin de la Société royale Linuéenne de Bruxelles,
- Bull. Soc. L. Nord France—Bulletin de la Société Linnéenne du Nord de la France (Amiens).
- Bull. Soc. L. Norm.—Bulletin de la Société Linnéenne de Normandie (Caen).
- Bull. Soc. Mal. Fr.—Bulletin de la Société Malacologique de France (Paris).
- Bull. Soc. Mal. Ital.—Bullettino della Società Malacologica Italiana (Pisa).
- Bull. Soc. Metz—Bulletin de la Société d'Histoire Naturelle de Metz (formerly du Département de la Moselle).
- Bull. Soc. Murith.—Bulletin des travaux de la Société Murithienne du Valais (Neufchâtel).
- Bull. Soc. Nancy-Bulletin de la Société des Sciences de Nancy (Paris).
- Bull. Soc. Neuchatel—Bulletin de la Société des Sciences Naturelles de Neuchatel.
- Bull. Soc. Nimes—Bulletin de la Société d'étude des sciences naturelles de Nimes (Nimes).
- Bull. Soc. Ouest France—Bulletin de la Société des Sciences Naturelles de l'Ouest de la France (Paris).
- Bull. Soc. Philom.—Bulletin de la Société Philomathique de Paris.
- Bull. Soc. Rouen—Bulletin de la Société des Amis des Sciences Naturelles de Rouen (Rouen).
- Bull. Soc. Saone—Bulletins de la Société des Sciences Naturelles de Saoneet-Loire (Chalons sur Saone.) (Also Mem.)
- Bull. Soc. Saroie—Bulletin de la Société d'Histoire Naturelle de Savoie (Chambèry).
- Bull. Soc. Sci. Phys. Nat. Toulouse—Bulletin de la Société des Sciences
  Physiques et Naturelles de Toulouse (Toulouse).
- Bull. Soc. Isère—Bulletin de la Société de Statistique des Sciences Naturelles, &c., du Département de l'Isère (Grenoble).

- Bull. Soc. Toulouse—Bulletin de la Société d'Histoire Naturelle de Toulouse.
- Bull. Soc. Vaud.—Bulletin de la Société Vaudoise des Sciences Naturelles (Lausanne).
- Bull. Soc. Ven.-Trent.—Bullettino della Società Veneto-Trentina di Scienze Naturali (Padua). (Also Atti.)
- Bull. Soc. Yonne—Bulletin de la Société des Sciences Historiques et Naturelles de l'Yonne (Auxerre).
- Bull. Soc. Z. Fr.—Bulletin de la Société Zoologique de France (Paris).
- Bull. U. S. Fish Comm.—Bulletin of the United States Fish Commission (Washington).
- Bull. U. S. Geol. Surv.—Bulletin of the United States Geological Survey (Washington).
- Bull. U. S. Nat. Mus.—Bulletin of the United States National Museum (Washington). (Also Proceedings.)
- Bull. Washb. Coll.—Bulletin of the Washburn Laboratory of Natural History (Topeka, Kansas).
- Canad. Ent.—Canadian Entomologist (Saunders: Montreal).
- Can. Rec.—Canadian Record of Science.
- Cardiff Nat. Soc.—Cardiff Naturalists' Society. Report and Transactions (Cardiff).
- CB. Bakt. Parasit.—Centralblatt für Bakteriologie und Parasitenkunde (Uhlworn: Cassel).
- CB. Ges. Anthrop.—Correspondenzblatt der deutschen Gesellschaft für Anthropologie, &c. (Brunswick).
- CB. Iris—Correspondenz-Blatt des entomologischen Vereins Iris zu Dresden.
- CB. med. Wiss.—Centralblatt für die medicinischen Wissenschaften (Berlin).
- CB. Ver. Regensb.—(Now Ber.)
- CB. Ver. Rheinl.—Correspondenz-Blatt des naturhistorischen Vereins der preussischen Rheinlande und Westphalens (Bonn). (Also Verh. & SB.)
- CB. Ver. Riga Correspondenzblatt des Naturforscher-Vereins zu Riga.
- Cellule—La Cellule. Recueil de Cytologie et d'Histologie Générale (Carnoy, Gilson, & Denys: Lierre & Gand).
- Circ. deutsch. Fisch. Ver.—Circulare des deutschen Fischerei-Vereins (Berlin).
- Comm. Ateneo Brescia-Commentari dell' Ateneo di Brescia.
- Conch. Mittheil.—Conchologische Mittheilungen (Martens: Cassel).
  - 3. M. Mus. Geol. Princeton—Contributions from the E. M. Museum vology, Princeton College, U.S.A. (Scott & Osborn).

- C.R.—Comptes rendus des Séances hebdomadaires de l'Académie des Sciences (Paris).
- C.R. Ass. Fr. Sci.—Compte-rendu de l'Association Français pour l'avancement des Sciences.
- C.R. Ent. Belg.—Comptes rendus des Séances de la Société Entomologique de Belgique (Brussels).
- C.R. Soc. Biol.—Comptes rendus hebdomadaires des Séances et Mémoires de la Société de Biologie (Paris).
- C.R. Soc. L. Bord.—Comptes Rendus de la Société Linnéenne de Bordeaux.
  (Also Actes.)
- Dan. Selsk. Skr.—K. Danske Videnskabernes Selskabs Skrifter (Copenhagen).
- Denk. Ak. Wien—Denkschriften der k. Akademie der Wissenschaften zu Wien (Vienna). (Also SB.)
- Deutsche s. Z.—Deutsche entomologische Zeitschrift (Kraatz: Berlin).
- Échange—L'Échange, Revue Linnéenne, organe des Naturalistes de la région Lyonnaise.
- E. Mus. Lund.—E museo Lundii En Samling Af Afhandlinger (Lütken : Kjøbnhavn).
- Ent.—The Entomologist (London).
- Ent. Am.—Entomologica Americana (Brooklyn).
- Ent. Gen.—L'Entomologiste Genevois. Journal mensuel d'Entomologie pure et appliquée (Geneva).
- Ent. M. M.—The Entomologist's Monthly Magazine (London).
- Ent. Med.—Entomologiske Meddeldser udgivne af Entomologisk Forening ved Fr. Meinert Copenhagen.
- Ent. Nachr.—Entomologische Nachrichten (Karsch: Berlin).
- Ent. Tidskr.—Entomologisk Tidskrift, på föranstaltande af Entomologiska Föreningen i Stockholm (Spångberg: Stockholm).
- Ért. Term. Kör.—Értekezések a természettudományok köréből, Magyar tudományos Akadémia [Memoirs on Natural Science, Hungarian Academy of Sciences] (Budaposth).
- Ess. Nat.—Essex Naturalist: being the Journal, Transactions, and Proceedings of the Essex Field Club (Buckhurst Hill).
- Études d'Ent.—Études d'Entomologie, Faunes Entomologiques, Descriptions d'Insectes nouveaux ou peu connus (C. Oberthür: Rennes).
- Evkön. Erd. Muz.—Evkönyvek erdélzi muzeumegylét (Kolozsvár = Klausenberg).
- Fauna—Fauna. Comptes rendus des Sciences de la Société de Naturalistes Luxembourgeois.
- Feuill. Nat.—Feuille des Jeunes Naturalistes (Dollfus: Paris).
- Field—The Field (London).

Field Club.—The Field Club, with which is incorporated The Garner (Loudon).

For. & Str.-Forest and Stream.

Forh. Selsk. Chr.—Forhandlinger i Videnskabs-Selskabet i Christiania.

Förh. Sk. Naturf.—Förhandlingar vid de Skandinaviska Naturforskarnes. Frelon.—Le Frelon, journal d'Entomologie.

Garner—The Garner, &c. (now Field Club, q.v.).

Gef. Welt—Die gefiederte Welt: Zeitschrift für Vogelliebhaber, -zuchter und -händler (Russ: Berlin).

Geol. Mag.—Geological Magazine (Woodward: London).

Giorn. Sci. Palerm.—Giornale di Scienze Naturali ed Economiche di Palermo.

Göteborgs Handl.—Göteborgs kongl. Vetenskaps och Vitterhets samhälles Handlingar.

Helios-See Mon. MT. Ver. Naturw. Frankfurt-a-O.

Hist. Berwick Nat. Club—History of the Berwickshire Naturalists' Club (Alnwick).

Hor. Ent. Ross.—Hore Societatis Entomologicae Rossicae (St. Petersburg). Humming Bird.—The Humming Bird (Boucard: London).

Ibis-The Ibis (Sclater: London).

Ind. Mus. Notes-Indian Museum Notes (Calcutta).

Ins. Life—Insect Life (Washington).

Int. J. Micr.—The International Journal of Microscopy & Natural Science (Allen & Spiers: London & New York).

Irish Natural.-The Irish Naturalist.

Isvest, Mosc. Unir.—Isvestiya imperatorskova obshchestva lyubiteleì Estestvoznaniya, Antropologi i Etnografii Sostoyashova, pre Moskovskom Universitet.

- J. Ac. Philad.—Journal of the Academy of Natural Sciences of Philadelphia.
- J. Agric. Soc. India—Journal of the Agricultural and Horticultural Society of India (Calcutta).
- J. Anat. Phys.—Journal of Anatomy and Physiology (London).
- J. A. S. B.—Journal of the Asiatic Society of Bengal (Calcutta).
- J. A. S. (Bombay)—Journal of the Bombay Branch of the Royal Asiatic Society.
- J. A. S. (Ceylon)—Journal of the Ceylon Branch of the Royal Asiatic Society.
- J. A. S. (China)—Journal of the China Branch of the Royal Asiatic Society.
- J. A. S. (Straits)—Journal of the Straits Branch of the Royal Asiatic Society (Singapore).

- JB. Ak. Amst.—Jaarboek van de k. Akademie van Wetenschappen (Amsterdam). (Also Verhandl.)
- JB. Mijnwezen—Jaarboek van het Mijnwezen in Nederlandsch Oost-Indië (Amsterdam).
- JB. geol. Reichsanst.—Jahrbuch der k.-k. geologischen Reichsanstalt (Vienna). (Also Verhandl.)
- JB. Hamb.-Jahrbuch der Hamburgischen wissenschaftlichen Austalten.
- JB. Karpath. Ver.—Jahrbuch des Ungarischen Karpathen-Vereins (Kesmark).
- JB. k. Akad. Erfurt—Jahrbücher der königlichen Akademie gemeinnütziger Wissenschaften zu Erfurt.
- JB. k. preuss. geol. Landesanst.—Jahrbuch der königlich preussischen geologischen Landesanstalt und Bergakademie zu Berlin.
- JB. mal. Ges.—Jahrbuch der deutschen malakozoologischen Gesellschaft (Kobelt: Frankfort).
- JB. Mineral.—Neues Jahrbuch für Mineralogie, Geologie, und Paleontologie (Leonard & Geinitz: Leipzig).
- JB. Mijnwezen Nederl. Indie—Jaarboek van het Mijnwezen van Nederl. Oost Indie (Amsterdam).
- JB. Mus. Kärnt.—Jahrbuch des naturhistorischen Landesmuseums von Kärnthen (Klagenfurt).
- JB. nass. Ver.—Jahrbuch des nassauischen Vereins für Naturkunde (Wiesbaden).
- JB. Sieb. Kurpath. Ver.—Jahrbuch des Siebenbürgischen Karpathen-Vereins (Hermannstadt).
- Jahresber. först.-phän. Stat.—Jahresbericht der förstlich-phänologischen Stationen Deutschlands (Berlin).
- J. Ber. Annab. Ver.—Jahresbericht des Annaberg-Buchholzer Vereius für Naturkunde (Annaberg).
- J. Ber. Ges. Graub.—Jahresbericht der naturforschenden Gesellschaft Graubündens (Chur).
- J. Ber. Ges. Hannov.—Jahresbericht der naturforschenden Gesellschaft in Hannover.
- J. Ber. k. Böhm. Ges. Wiss.—Jahresbericht der königlich Böhmischen Gesellschaft der Wissenschaften.
- J. Ber. Pollichia—Jahresbericht der Pollichia eines naturwissenschaftlichen Vereins der Rheinpfalz (Dürkheim a. d. Hart).
- J. Ber. Schles. Ges.—Jahresbericht der Schlesischen Gesellschaft für vaterländische Cultur (Breslau). (Also Abhandl.)
- J. Ber. Ver. Braunschw.—Jahresbericht des Vereins für Naturwissenschaft zu Braunschweig (Brunswick).
- J. Ber. Ver. Frankfurt—Jahresbericht des physikalischen Vereins zu Frankfurt-am-Main.
- J. Ber. Ver. Magdeburg—Jahresbericht und Abhandlungen des naturwissenschaftlichen Vereins in Magdeburg.

- J. Ber. Ver. Osnabr.—Jahresbericht des naturwissenschaftlichen Vereins zu Osnabrück,
- J. Ber. Ver. Zwickau—Jahresbericht des Vereins für Naturkunde zu Zwickau.
- J. Ber. Westf. Ver.—Jahresbericht der zoologischen Sektion des Westfälischen provinzial-Vereins für Wissenschaft und Kunst (Münster).
- J. Bomb. N. H. Soc.—The Journal of the Bombay Natural History Society (Phipson: Bombay).
- J. Brit. Dent. Ass.—Journal of the British Dental Association (London).
- J. Cincinn. Soc.—Journal of the Cincinnati Society of Natural History.
- J. Coll. Sci. Japan—Journal of the College of Science, Imperial University, Japan (Tökyö).
- J. de Conch.—Journal de Conchyliologie (Crosse & Fischer: Paris).
- J. de l'Anat. Phys.—Journal de l'Anatomie et de la Physiologie (Pouchet : Paris).
- J. Elisha Mitchell Sci. Soc.—Journal of the Elisha Mitchell Scientific Society (Raleigh, N.C.).
- Jen. Z. Nat.—Jenaische Zeitschrift für Naturwissenschaft, herausgegeben von der medicinisch-naturwissenschaftlichen Gesellschaft zu Jena.
- J. f. O.—Journal für Ornithologie (Cabanis: Leipzig).
- JH. Ver. Lüneb.—Jahreshefte des naturwissenschaftlichen Vereins für das Fürstenthum Lüneburg.
- JH. Ver. Württ.—Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg (Stuttgart).
- J. Inst. Jamaica—Journal of the Institute of Jamaica (Kingston).
- J. L. S.—Journal of the Linnean Society; Zoology (London).
- J. London Coll. Soc.—Journal of the City of London College Society, London (Lewes).
- J. Mar. Biol. Ass.—Journal of the Marine Biological Association (London & Plymouth).
- J. Morph,—Journal of Morphology (Whitman & Allis: Boston, U.S.A.).
- J. Microgr.-Journal de Micrographie (Pellétan: Paris).
- J. New Jersey N. H. Soc.—Journal of the New Jersey Natural History Society (Trenton). (Formerly J. Trenton Soc.)
- J. Northampt. Soc.—Journal of the Northamptonshire Natural History Society and Field Club.
- J. N. Y. Micr. Soc.—Journal of the New York Microscopical Society (New York).
- J. of Conch.—Journal [formerly Quarterly Journal] of Conchology (London).
- Johns Hopk, Univ. Circ.—Johns Hopkins University Circulars (Baltimore).
- J. Physiol.—The Journal of Physiology (Foster et alii : Cambridge).
- J. Quek. Club-Journal of the Quekett Microscopical Club (London).

- J. R. Agric. Soc.—Journal of the Royal Agricultural Society (London).
- J. R. Inst. Cornwall—Journal of the Royal Institution of Cornwall (Truro).
- J. R. Micr. Soc.—Journal of the Royal Microscopical Society (London).
- J. B. Geol. Soc. Ireland—Journal of the Royal Geological Society of Ireland (London, Dublin, and Edinburgh).
- J. R. Soc. N. S. W.—Journal and Proceedings of the Royal Society of New South Wales (Sydney).
- J. Sci. Lisb.—Jornal de Sciencias, &c., da Academia de Lisboa (Lisbon).
- J. Tr. Vict. Inst.—Journal of the Transactions of the Victoria Institute, or Philosophical Society of Great Britain (London).
- Kansas quart.—The Kansas University quarterly.
- Kansas Scient.—The Kansas City Scientist (Kansas). (Formerly The Naturalist).
- Kosmos Lemberg-Kosmos: Lemberg.
- L'Ab.—L'Abeille (Paris).
- La Nature-La Nature, Revue des Sciences, &c. (Tissandier: Paris).
- L'Anthrop.—L'Authropologie (Paris).
- Le Nat.-Le Naturaliste (Deyrolle : Paris).
- Leopoldina—Leopoldina: Ämtlichen Organ für der k. Leopold-Carol. deutsch. Acad.
- Lioar Man.—Yn Lioar Manninagh. Published quarterly, for The Isle of Man Natural History and Antiquarian Society (Kermode: Ramsey).
- Lotos-Lotos, Jahrbuch für Naturwissenschaft im Auftrage des Vereines 'Lotos' (Prague).
- Maandbl. Natuurw.—Maandblad voor Naturwetenschappen (Amsterdam).

  Mal. Bl.—Malakozoologische Blätter (Clessin: Cassel).
- Math. Nat. Ber. Ung.— Mathematische und naturwissenschaftliche Berichte aus Ungarn. Mit Unterstützung der Ungarischen Akad. d. Wiss. und der K. Ungar. naturwiss. Ges. herausgegeben von Baron R. Eötvös, &c. (Fröhlich: Buda-Pest).
- Math. term. Értes.—Mathematikai és természettudományi Értesitő (Gyula : Buda-Pesth).
- Matk. term. köz.—Mathematikai és természettudományi közlemények (Buda-Pest).
- Med. Nat.—The Mediterranean Naturalist (Cooke: Malta).
- Med. Soc. Fenn.—Meddelanden af Societas pro Fauna et Flora Fennica (Helsingfors).
- Mél. biol.—Mélanges biologiques tirés du Bulletin de l'Académie impériale des Sciences de St. Petersburg.
- Mem. Ac. Barcel.—Memorias de la real Academia de Ciencias de Barcelona.

- Mém. Ac. Belg.—Mémoires de l'Académie royale des Sciences, des Lettres, et des Beaux Arts de Belgique (Brussels). (Also Bull.)
- Mem. Acc. Bologn.—Memorie della R. Accademia delle Scienze dell' Istituto di Bologna.
- Mém. Ac. Dijon—Mémoires de l'Académie des Sciences, Arts, et Belles-Lettres de Dijon.
- Mém. Ac. Lyon—Mémoires de l'Académie des Sciences, Belles-Lettres et Arts de Lyon (Lyon).
- Mem. Ac. Madrid—Memorias de la real Academia de Ciencias Exactas Fisicas, y Naturales de Madrid (Madrid).
- Mem. Acc. Mod.—Memorie delle R. Accademia di Scienze, Lettere, ed Arti in Modena.
- Mém. Ac. Montp.—Mémoires de la Section des Sciences de l'Académie des Sciences et Lettres de Montpellier.
- Mém. Ac. Pétersb. (7)—Mémoires de l'Académie impériale des Sciences de St. Pétersbourg. 7me série.
- Mém. Ac. Savoie—Mémoires de l'Académie des Sciences, Belles-Lettres, et Arts de Savoie (Chambéry). (Also Comptes Rendus.)
- Mém. Ac. Sci.—Mémoires de l'Académie des Sciences (Paris).
- Mem. Ac. Sci. Lisboa-Memorias da Academia real das Sciencias de Lisboa.
- Mem. Acc. Tor.—Memorie della R. Accademia delle Scienze di Torino (Turin).
- Mém. Ac. Toulouse—Mémoires de l'Académie des Sciences, &c., de Toulouse.
- Mém. Ac. Vaucluse-Mémoires de l'Académie de Vaucluse (Avignon).
- Mem. Am. Ac.—Memoirs of the American Academy of Arts and Sciences (Cambridge). (Also Proceedings.)
- Mem. Boll. Soc. Geogr. Ital. Memorie (Bollettino) della Società Geografica Italiana (Rome).
- Mem. Bost. Soc.—Memoirs of the Boston Society of Natural History.

  (Also Proceedings.)
- Mem. California Acad.—Memoirs of the Californian Academy of Sciences (San Francisco).
- Mém. Cour. Ac. Belg. 4to.—Mémoires Couronnés et Mémoires des Savants Étrangers, publiés par l'Académie Royale des Sciences, des Lettres et des Beaux Arts de Belgique.
- Mém. Cour. Ac. Belg. 8vo.—Ibid. 8vo.
- Mém. Inst. Genév.—Mémoires de l'Institut national Genévois (Geneva).
- Mem. Ist. Lomb.—Memorie del R. Istituto Lombardo di Scienze e Lettere (Milan).
- Mem. Ist. Venet. Memorie del R. Istituto Veneto di Scienze, &c. (Venice).
- Mém. Liège-Mémoires de la Société royale des Sciences de Liège.
- Mem. Mus. C. Z.—Memoirs of the Museum of Comparative Zoology at Harvard College (Cambridge, U.S.A.).

- Mem. Nat. Ac. Sci.—Memoirs of the National Academy of Sciences (Washington).
- Mém.-prés. Ac. Sci.—Mémoires présentés par divers savants a l'Académie des Sciences de l'Institut de France (Paris).
- Mém. Soc. Biol.—(See C.R.)
- Mém. Soc. Bord.—Mémoires de la Société des Sciences Physiques et Naturelles de Bordeaux.
- Mém. Soc. Cannes—Mémoires de la Société des Sciences Naturelles et Historiques des Lettres et des Beaux-Arts de Cannes et de l'Arrondissement de Grasse (Cannes).
- Mém. Soc. Cherb.—Mémoires de la Société nationale des Sciences Naturelles et Mathématiques de Cherbourg.
- Mém. Soc. Géol. Mémoires de la Société Géologique de France (Paris).
- Mém. Soc. Hainault Mémoires et publications de la Société des Sciences, des Arts, et des Lettres du Hainault (Mons).
- Mém. Soc. Lille—Mémoires de la Société de l'Agriculture et des Arts de Lille.
- Mém. Soc. L. N. Fr.—Mémoires de la Société Linnéenne du Nord de la France (Amiens).
- Mém. Soc. Maine et Loire—Mémoires de la Société académique de Maine et Loire (Angers).
- Mem. Soc. Manch.—Memoirs and Proceedings of the Manchester Literary and Philosophical Society (London).
- Mém. Soc. Oise—Mémoires de la Société académique d'Archéolo des Sciences et Arts du département de l'Oise.
- Mém. Soc. Phys. Genèv.—Mémoires de la Société de Physique et d'Histoire Naturelle de Genève.
- Mém. Soc. Saône—Mémoires de la Société des Sciences Naturelles de Saône-et-Loire (Chalon-sur-Saône). (Also Bulletin.)
- Mém. Soc. Seine & Oise—Mémoires de la Société des Sciences Naturelles et Médicales de Seine-et-Oise (Versailles).
- Mém. Soc. Zool.—Mémoires de la Société Zoologique de France (Paris).
- Mid. Nat.—The Midland Naturalist (Badger & Hillhouse: London & Birmingham).
- Monit. Zool. Ital.—Monitore Zoologico Italiano (Chiarugi & Ficalbi: Florence.)
- Month. Int. J. Anat. Hist.—Monthly International Journal of Anatomy and Histology (Paris, Leipsic, London).
- Morph. JB.—Morphologisches Jahrbuch: eine Zeitschrift für Anatomie und Entwickelungsgeschichte (Gegenbaur: Leipzig).
- MT. Aargau. nat. Ges. Mittheilungen der Aargauischen naturforschenden Gesellschaft.

- MT. embr. Inst. Wien (n.s.)—Mittheilungen aus dem embryologischen Institute der k. k. Universität in Wien. New series (Schenck: Vienna).
- MT. Ges. Bern-Mittheilungen der naturforschenden Gesellschaft in Bern.
- MT. min. geol. Mus. Dresden—Mittheilungen aus dem k. mineralogischgeologischen und præhistorischen Museum in Dresden.
- MT. orn. Ver. Wien—Mittheilungen des ornithologischen Vereins in Wien (Vienna).
- MT. Osterlande-Mittheilungen aus dem Osterlande (Altenburg).
- MT. Schw. ent. Ges.—Mittheilungen der Schweizerischen entomologischen Gesellschaft (Schaffhausen). (Also with French title.)
- MT. Ung. geol. Anst.—Mittheilungen aus dem Jahrbuche der k. Ungarischen geologischen Anstalt (Buda-Pest).
- MT. Ver. Steierm. Mittheilungen des naturwissenschaftlichen Vereins für Steiermark (Gratz).
- MT. Vorpomm.—Mittheilungen aus dem naturwissenschaftlichen Vereine von Neu-Vorpommern und Rügen (Griefswald).
- MT. z. Stat. Neap.—Mittheilungen aus der zoologischen Station in Neapel (Leipzig).
- N. Act. Ups.—Nova Acta R. Societatis Scientiarum Upsaliensis (Upsala).
- N. Acta Ac. L.-C. Nat. cur.—Nova Acta Academiæ Css. Leopoldino-Carolins Germanis Naturæ curiosorum (Leipzig). (Also Verhandlungen der k. Leop. Carol. deutschen Acad. d. Naturf.)
- N. Arch. Mus. Nouvelles Archives du Muséum d'Histoire Naturelle (Paris).
- N. Denk. Schwer. Ges.—Neue Denkschriften der allgemeinen Schweizerischen Gesellschaft für die gesammten Naturwissenschaften.
- N. Mém. Soc. Helv.—Nouveaux Mémoires de la Société Helvetique des Sciences Naturelles (Lausanne).
- N. Mém. Soc. imp. Moscou—Nouveaux Mémoires de la Société impériale des Naturalistes de Moscou.
- N. Z. J. Sci. The New Zealand Journal of Science (Dunedin).
- Nuchr. Ges. Götting.—Nachrichten von der k. Gesellschaft der Wissenschaften und der Georg Auguste Universität zu Göttingen.
- Nachr. mal. Ges.—Nachrichtsblatt der deutschen malakozoologischen Gesellschaft (Frankfort).
- Nat. Canad.—Le Naturaliste Canadien (Provancher: Cap Rouge, Quebec).
- Nat.-Hist. Tr. North. Durham.—Natural-History Transactions of Northumberland, Durham, and Newcastle-on-Tyne (London & Newcastle).

- Nat. Mex.—La Naturaleza (Mexico).
- Nat. Sicil.—Il Naturalista Siciliano: Giornale delle Scienze Naturali (Ragusa: Palermo).
- Nat. Tijdschr. Nederl. Ind.—Natuurkundig Tijdschrift voor Nederlandsche Indie (Batavia).
- Nat. Notes—Nature Notes, The Selborne Society's Magazine (Britten: London).
- Naturalist—The Naturalist: Journal of the Yorkshire Naturalists'
  Union, &c. A Monthly Journal of Natural History for North of
  England (Roebuck & Clarke: London & Leeds).
- Natural Sci.—Natural Science: a Monthly Review of Scientific Progress. Nature—Nature (London).
- Nat. Ver. Haarlem—Natuurkundige Verhandelingen van de Hollandsche Maatschappig der Wetenschappen te Haarlem (Haarlem).
- Nat. Ver. Utrecht Natuurkundige Verhandelingen Provinciaal Utregtsch genootschap van Kunsten en Wetenschappen (Utrecht).
- Naut.—The Nautilus (Pilsbry & Averell: Philadelphia).
- Neujahrsbl. Naturf. Ges.—Neujahrsblatt herausgegeben von der Naturforschenden Gesellschaft (Zurich).
- Nor. Selsk. Skr.—K. Norske Videnskabernes Selskabs Skrifter (Thrond-jhem).
- Notes Leyd. Mus.—Notes from the Royal Zoological Museum of the Netherlands at Leyden (Jentink).
- Notizbl. Ver. Erdk. Darmstadt—Notizblatt des Vereins für Erdkunde zu Darmstadt.
- Nunq. ot.—Nunquam otiosus (Schaufuss: Dresden).
- Nyt. Mag. Naturv.—Nyt Magazin for Naturvidenskaberne (Danielssen et alii : Christiania).
- Œfv. Ak. Förh. Œfversigt af k. Vetenskaps Akademiens Förhaudlingar (Stockholm).
- Œfv. Finska Förh.—Œfversigt af Finska Vetenskaps Societetens Förhandlingar (Helsingfors).
- Orn. JB. Ornithologisches Jahrbuch. Organ für die palæarktische Faunengebiet.
- Orn. MB.—Ornithologische Monatsberichte herausgegeben von Dr. A. Reichenow, Berlin.
- Orn. Ool.—Ornithologist & Oologist (Pawtucket, R. I.).
- Ornis—Ornis: Internationale Zeitschrift für die gesammte Ornithologie (Blasius & Hayek: Vienna).
- Orvos-termesz. Értesítő.—Orvos-termeszettudományi Értesítő (Kolozsvar = Klausenberg).
- Ottawa Nat.—The Ottawa Naturalist. The Transactions of the Ottawa Field Naturalists' Club (Ottawa).

- Overs. Dan. Selsk.—Oversigt over det k. Danske Videnskabernes Selskabs Forhandlinger (Copenhagen).
- P. Ac. Philad.—Proceedings of the Academy of Natural Sciences of Philadelphia.
- P. Am. Ac.—Proceedings of the American Academy of Arts and Sciences (Boston). (Also Mem.)
- P. Am. Ass.—Proceedings of the American Association for the Advancement of Science. (Also Msm.)
- P. Am. Micr. Soc.—Proceedings of the American Society of Microscopists (Buffalo).
- P. Am. Phil. Soc.—Proceedings of the American Philosophical Society, &c. (Philadelphia). (Also Trans.)
- Pap. Soc. Brit. Columb.—Papers and communications read before the Natural History Society of British Columbia (Victoria, B. C.).
- P. A. S. B.—Proceedings of the Asiatic Society of Bengal (Calcutta).
- P. Bath. N. H. Soc.—Proceedings of the Bath Natural History and Antiquarian Field Club.
- P. Belf. Soc.—(See Report.)
- P. Biol. Soc. Washington—Proceedings of the Biological Society of Washington.
- P. Birmingh. Phil. Soc.—Proceedings of the Birmingham Philosophical Society.
- P. Birmingh. Soc.—(See Report.)
- P. Bost. Soc.—Proceedings of the Boston Society of Natural History (Boston, U.S.A.) (Also Mem.)
- P. Bristol Soc.—Proceedings of the Bristol Naturalists' Society.
- P. Cal. Ac. Sci.—Proceedings of the Californian Academy of Sciences (San Francisco). (Also Trans. and Occas. Papers.)
- P. Cambr. Phil. Soc.—Proceedings of the Philosophical Society, Cambridge. (Also Trans.)
- P. Colorado Soc. Proceedings of the Colorado Scientific Society (Denver).
- P. Cottesso. Nat. F. C.—Proceedings of the Cotteswold Naturalists' Field Club (Gloucester).
- P. Croydon Club—Proceedings of the Croydon Microscopical and Natural History Club.
- P. Davenport Ac.—Proceedings of the Davenport Academy of Natural Sciences (Davenport, Iowa).
- P. Dorset Field Club—Proceedings of the Dorset Natural History and Antiquarian Field Club (Sherborne).
- P. E. Soc.—Proceedings of the Entomological Society of London.
- P. E. Soc. Wash. Proceedings of the Entomological Society of Washington.

- P. Folkestone Soc.—Proceedings of the Folkestone Natural History Society.
- P. Geol. Ass.—Proceedings of the Geologists' Association (London).
- P. Hampshire Club—Papers and Proceedings of the Hampshire Field Club (Southampton).
- P. Holmesdale Nat. Hist. Club—Proceedings of the Holmesdale Natural History Club (London).
- P. Linn. Soc. N.S.W.—Proceedings of the Linnean Society of New South Wales (Sydney).
- P. Liverp. Biol. Soc. Proceedings of the Liverpool Biological Society.
- P. Liverp. Field Club.—Proceedings of the Liverpool Naturalists' Field Club.
- P. Liverp. Soc.—Proceedings of the Literary and Philosophical Society of Liverpool.
- P. London Amateur Soc.—Proceedings of the London Amateur Scientific Society. Metropolitan Scientific Association and Society of Amateur Geologists (London).
- P. Newport Nat. Hist. Soc.—Proceedings of the Newport Natural History Society.
- P. N. H. Soc. Glasg.—Proceedings of the Natural History Society of Glasgow.
- P. N.-Scot. Inst.—Proceedings and Transactions of the Nova-Scotian Institute of Natural Science (Halifax, N.-S.).
- P. Phil. Soc. Glasg.—Proceedings of the Philosophical Society of Glasgow.
- P. Phys. Soc. Edinb.—Proceedings of the Royal Physical Society of Edinburgh.
- P. Rochester Academy of Science (Rochester, N.Y.)
- P. R. Inst.—Proceedings of the Royal Institution of Great Britain (London).
- P. R. Irish Ac.—Proceedings of the Royal Irish Academy (Dublin).
- P. R. Soc.—Proceedings of the Royal Society (London).
- P. R. Soc. Edinb.—Proceedings of the Royal Society of Edinburgh. (Also Trans.)
- P. R. Soc. Queensl.—Proceedings of the Royal Society of Queensland (Brisbane).
- P. R. Soc. Tasm.—Papers and Proceedings and Reports of the Royal Society of Tasmania (Hobarton).
- P. R. Soc. Vict.—Proceedings of the Royal Society of Victoria (Melbourne). (Also Trans.)
- P. Soc. Manch.—(See Mem.)
- P. Somerset. Soc.—Proceedings of the Somersetshire Archeological and Natural History Society. New series (Taunton).

- P. Tr. Croydon Nat. Hist. Club—Proceedings and Transactions of the Croydon Microscopical and Natural History Club (Croydon).
- P. U. S. Nat. Mus.—Proceedings of the United States National Museum (Washington). (Also Bull.)
- P.-v. Soc. Mal. Belg.—Procès-verbaux des séances de la Société Malacologique de Belgique (Brussels).
- P.-v. Soc. Tosc.—Processi verbali della Società Toscana delle Scienze Naturali (Pisa).
- P. Warwick. Club—Proceedings of the Warwickshire Naturalista' and Archæologists' Field Club (Warwick).
- P. Z. S.—Proceedings of the Zoological Society (London). (Also Trans.)
- Pal. Abh. Palæontologische Abhandlungen (Dames & Kayser: Berlin).
- Pulæontogr.—Palæontographica: Beiträge zur Naturgeschichte der Vorwelt (Cassel).
- Pal. Ind.—Palsontologia Indica. (4to) Memoirs of the Geological Survey of India (Calcutta).
- Pal. Soc.—Monographs of the Palsontological Society.
- Pam. Akad. umiej. wydz. przyr. Krakau Pamietnik Akademii Umiejetności w Krakowie. Wydział matem. przyr (Cracow).
- Pam. Fizjogr.—Pamietnik Fizjograficzny (Warsaw).
- Phil. Tr.—Philosophical Transactions of the Royal Society (London). (Also Proc.)
- Preisschr. Jblonovsk. Gesells. Leipsig Preisschriften gekrönt und herausgegeben von der fürstlich Jablonovski' schen Gesellschaft zu Leipsig.
- Prodr. Zool. Vict.—Prodromus of the Zoology of Victoria (McCoy: Melbourne).
- Protok. obsch. estest. Kazan—Protokolui zasyedanii obshchestva estestvoispuitatelei pri imperatorskom Kazanskom Universitetye.
- Psyche—Psyche, a Journal of Entomology. Published by the Cambridge Entomological Club (Cambridge, Mass., U.S.A.).
- Q. J. Geol. Soc.—Quarterly Journal of the Geological Society (London).
- Q. J. Micr. Sci.—Quarterly Journal of Microscopical Science (Lankester et alii: London).
- Rad jugoslav. akad.—Rad jugoslavenske akademije znanosti i umjetnosti (Zagreb). [Transactions of the South Slav Academy of Science and Art.]
- Rec. Austral. Mus.—Records of the Australian Museum (Ramsay: Sydney).
- Rec. Geol. Surv. Ind.—Records of the Geological Survey of India (Calcutta).
- Rec. Z. Suisse- Recueil Zoologique Suisse (Fol: Geneva & Bâle).

- Rend. Acc. Nap.—Rendiconto dell' Accademia delle Scienze Fisiche e Matematiche (Sezione della Sociétá reale di Napoli).
- Rend. Ist. Lombardo—Rendiconti del R. Istituto Lombardo delle Scienze e Lettere (Milan).
- Rep. Austr. Ass.—Reports of the Australasian Association for the Advancement of Science.
- Rep. Austr. Zool. Soc.—Report of the South Australian Zoological and Acclimatization Society (Adelaide).
- Rep. Belfast Field Club—Annual Report and Proceedings of the Belfast Naturalists' Field Club.
- Rep. Brighton Soc.—Annual Report and Abstract of Proceedings of the Brighton and Sussex Natural History Society (Brighton).
- Rep. Brit. Ass.—Report of the British Association for the Advancement of Science.
- Rep. Cornell Univ. Stat.—Report of the Department of Entomology of the Cornell University Experiment Station (Comstock: Ithaca, N.Y.).
- Rep. Cornwall Polytechn.—Reports of the Royal Polytechnical Society of Cornwall.
- Rep. Dep. Agric. Ent.—Report of the Entomologist. From the Annual Report of the Department of Agriculture (Washington).
- Rep. E. Soc. Ont.—Report of the Entomological Society of the Province of Ontario.
- Rep. Felsted Soc.—Report of the Felsted School Natural History Society (Chelmsford).
- Rep. Fish. Scotl.—Annual Report of the Fishery Board for Scotland (London, Edinburgh, & Dublin).
- Rep. Geol. Surv. Canada—Report of the Geological and Natural History Survey and Museum of Canada (Montreal).
- Rep. Guernsey Soc.—Report and Transactions; Guernsey Society of Natural Science and Local Research.
- Rep. Ins. Illin.—Annual Report of the Noxious and Beneficial Insects of the State of Illinois (Springfield).
- Rep. Ins. N. York—Annual Report of the Injurious and other Insects of New York (Lintner: Albany).
- Rep. Leeds Soc.—Leeds Philosophical and Literary Society. The Annual Report (Leeds).
- Rep. Marlb. Coll. Soc.—Report of the Marlborough College Natural History Society.
- Rep. N. Staffordsh. Club—North Staffordshire Naturalists' Field Club and Archeological Society. Annual Report and Transactions.
- Rep. N. Y. Mus.—Annual Report of the New York State Museum of Natural History (Albany).
- Rep. & P. Belfust N. H. Soc.—Report and Proceedings of the Belfast Natural History and Philosophical Society.

- Rep. Penzance Soc.—Report and Transactions of the Penzance Natural History and Antiquarian Society.
- Rep. Plym. Inst. Annual Report and Transactions of the Plymouth Institution and Devon and Cornwall Natural History Society (Plymouth).
- Rep. Rugby Soc. Report of the Rugby School Natural History Society.
- Rep. Southport Soc.—The . . . Report of the Southport Society of Natural Science.
- Rep. Tr. Devon Ass.—Report and Transactions of the Devonshire Association for the Advancement of Science, &c. (Plymouth).
- Rep. U. S. Ent. Comm.—Report of the United States Entomological Commission (Washington).
- Rep. U. S. Fish. Comm.—Report of the Commissioner, United States Commission of Fish and Fisheries (Washington).
- Rep. U. S. Geol. Surv.—Annual Report of the United States Geological Survey (Washington).
- Rep. U. S. Nat. Mus.—Annual Report of the Board of Regents of the Smithsonian Institution. Report of the National Museum (Washington).
- Rep. Wellington Soc.—Report of the Wellington College Natural Science Society (Wellington Coll.).
- Rep. Yorks. Phil. Soc.—Report of the Council of the Yorkshire Philosophical Society.
- Rev. Arg. Hist. Nat.—Revista Argentina de Historia Natural. (Ameghino: Buenos Ayres.)
- Rev. Biol.—Revue Biologique du Nord de la France (Barrois, Halles, Moniez : Lille).
- Rev. Cien. Madrid—Revista de los progresos de las Ciencias Exactas, Físicas, y Naturales (Madrid).
- Rev. Cient. Univ. Venezuela—Revista Cientifica mensual de la Universidad Central de Venezuela (Caracas).
- Rev. d'Ent.—Revue d'Entomologie, publié par la Société Française d'Entomologie (Fauvel : Caen).
- Rev. Mus. la Plata—Revista del Museo de la Plata (Ameghino: la Plata).
- Rev. Quest. Sci.—Revue des Questions Scientifiques publiée par la Société scientifique de Bruxelles (Brussels).
- Rev. Sci.—Revue Scientifique de la France et de l'Étranger (Paris).
- Rev. Sci. Bourb.—Revue Scientifique du Bourbonnais et du centre de la France (Olivier: Moulins).
- Rev. Sci. Nat. Oporto—Revista de Sciencas Naturaes e Socias (Severo & Peixoto: Porto).
- Rev. Sci. Nat. Ouest-Revue des Sciences Naturelles de l'Ouest (Paris).
- Rev. Sci. Nat. St. Pétersb.—Revue des Sciences naturelles. Publiée par la Société des Naturalistes à St. Pétersburg (St. Petersburg).

- Rev. Soc. Porto-Revista da Societa de Instrucção de Porto (Oporto).
- Rev. Tierkeilkunde-Revue für Tierheilkunde und Tierzucht (Vienna).
- Rev. Tr. Sci.—Revue des Travaux Scientifiques (Paris).
- Rev. Zool.—Revue Zoologique.
- Riv. Ital. Sci. Nat.—Rivista italiana di Scienze Naturali e Bollettino del Naturalista Collettore, Allevatore, Coltivatore (Brogi: Siena).
- Rochester Nat.—The Rochester Naturalist (Rochester).
- S. E. Z.—Stettiner entomologische Zeitung (Dohrn: Stettin).
- Samm. Geol. Mus. Leid.—Sammlungen des Geologischen Reichmuseums in Leiden.
- Samm. naturus. Vorträge—Sammlung naturwissenschaftlicher Vorträge (Huth: Berlin.)
- SB. Ak. Berlin—Sitzungsberichte der königlich Preussischen Akademie der Wissenschaften zu Berlin (formerly Bericht and Monatsbericht). (Also Abhandl.)
- SB. Ak. Wien—Sitzungsberichte der mathematische-naturwissenschaftlichen Classe der k. Akademie der Wissenschaften (Vienna). (Also Denkschriften.)
- SB. Bayer. Ak.—Sitzungsberichte der mathematisch-physikalischen Classe der k. Bayerischen Akademie der Wissenschaften (Munich). (Also Abhandl.)
- SB. Böhm. Ges.—Sitzungsberichte der k. Böhmischen Gesellschaft der Wissenschaften (Prague). (Also Abhandl.)
- SB. Ges. Dorp.—Sitzungsberichte der Naturforscher-Gesellschaft bei der Universität Dorpat (Dragendorff: Dorpat). (Also Schriften.)
- SB. Ges. Isis—Sitzungsberichte und Abhandlungen der naturwissenschaftlichen Gesellschaft 'Isis' (Dresden). (Also Abhandl.)
- SB. Ges. Königsb.—Sitzungsberichte der k. physikalisch-ökonomischen Gesellschaft in Preussen (Königsberg). (See Schr. Ges. Königsb.)
- SB. Ges. Leipzig—Sitzungsberichte der naturforschenden Gesellschaft zu Leipzig.
- SB. Ges. Marb.—Sitzungsberichte der Gesellschaft zur Beförderung der gesammten Naturwissenschaften zu Marburg.
- SB. Ges. Morph.—Sitzungsberichte der Gesellschaft für Morphologie und Physiologie in München (Munich).
- SB. Ges. Würzb.—Sitzuugsberichte des physikalisch-medicinischen Gesellschaft zu Würzburg. (Also Verh.)
- SB. nat. Fr.—Sitzungsberichte der Gesellschaft naturforschender Freunde zu Berlin.
- SB. niederrhein. Ges.—Sitzungsberichte der niederrheinischen Gesellschaft für Natur- und Heilkunde (Bonn). (Published with Verh. & CB. Ver. Rheinl.)
- SB. Soc. Erlangen Sitzungsberichte der physikalisch-medicinischen Societät zu Erlangen.

- SB. z.-b. Wien—Sitzungsberichte der zoologisch-botanischen Gesellschaft in Wien (Vienna). (Also Verh.)
- Schr. Ges. Danz.—Neueste Schriften der naturforschenden Gesellschaft zu Danzig.
- Schr. Ges. Königsb.—Schriften der physikalisch-ökonomischen Gesellschaft zu Königsberg in Preussen.
- Schr. gesammt. Naturvo. Marburg—Schriften der Gesellschaft zur Beförderung der gesammten Naturwissenschaften zu Marburg.
- Schr. Nat. ges. Dorpat.—Schriften herausgegeben von der Naturforscher-Gesellschaft bei der Universität Dorpat. (Also SB.)
- Schr. Nat. Ver. Schleswig-Schriften des naturwissenschaftlichen Vereins für Schleswig-Holstein (Kiel).
- Schr. Ver. Harzes—Schriften des naturwissenschaftlichen Vereins des Harzes in Wernigerode.
- Schr. Univ. Kiel-Schriften der Universität zu Kiel.
- Science-Science (Dall: Cambridge, Mass.).
- Sci. Goss. Hardwick's Science Gossip (Taylor: London).
- Sci. P. R. Dubl. Soc.—Scientific Proceedings of the Royal Dublia Society.
- Sci. Tr. R. Dublin Soc. (2)—The Scientific Transactions of the Royal Dublin Society. Second Series.
- Scot. Nat. The Scottish Naturalist (Clarke: Perth).
- Smiths. Contrib. Knowledge—Smithsonian Contributions to Knowledge (Washington).
- Smiths. Misc. Coll.—Smithsonian Miscellaneous Collections (Washington).
- Smiths. Report—Annual Report of the Board of Regents of the Smithsonian Institution, &c. (Washington).
- Soc. Ent.—Societas Entomologica.
- Soc. Agricole Pyrén.-orient.—Société Agricole Scientifique et Littéraire des Pyrénées-orientales (Perpignan).
- Sprawozd. Kom. fizyjogr.—Sprawozdanie Komisyi fizyjograficznej, &c. (Cracow).
- Stavanger Mus.—Stavanger Museums Aarsberetning.
- Stud. Biol. Lab. J. Hopkins Univ.—Studies at the Biological Laboratory of the Johns Hopkins University (Baltimore).
- Stud. Mus. Dundee—Studies from the Museum of Zoology in University College, Dundee (D'Arcy Thompson: Dundee).
- Sv. Ak. Handl.—K. Svenska Vetenskaps-Akademiens Handlingar (Stockholm).
- Tag. Deut. Nat. Vers.—Tageblatt der Versammlung deutscher Naturforscher und Aertzte.
- Tijdschr. Ent.—Tijdschrift voor Entomologie (The Hague).
- Tijdschr. Ind. Volkenkund.—Tijdschrift voor Indische Taal-, Land- en Volkenkunde, etc. (Brandes & Abendann, Batavia and the Hague).

- Tijdschr. Nederl. Dierk. Ver. (2)—Tijdschrift van de Nederlandsche Dierkundige Vereeniging (Leyden).
- Tijdschr. Nederl. Ind.—Naturkundig Tijdschrift voor Nederlandsch Indië (Batavia).
- Timehri.—Timehri: being the Journal of the Royal Agricultural and Commercial Society of British Guiana (Quelch: Demerara).
- Termes Közlöny Természettudományi Közlöny kiadja a k. magyar termeszett. Társulat (Budapest).
- Term. füzetek—Természetrajzi füzetek: kiadja a magyar remzeti Müseum (Journal of Zoology, &c., edited by Hungarian Museum at Budapesth).
- Tr. Albany Inst.—Transactions of the Albany Institute. (Formerly Proceedings also).
- Tr. Am. Phil. Soc.—Transactions of the American Philosophical Society, &c. (Philadelphia). (Also Proceedings).
- Tr. Am. Ent. Soc.—Transactions of the American Entomological Society (Philadelphia).
- Tr. A. S. Japan—Transactions of the Asiatic Society of Japan (Yokohama).
- Tr. Camb. Phil. Soc.—Transactions of the Cambridge Philosophical Society. (Also Proceedings.)
- Tr. City London Soc.—Transactions of the City of London Entomological and Natural History Society (London).
- Tr. Conn. Ac.—Transactions of the Connecticut Academy of Sciences (New Haven).
- Tr. Croydon Club.—(See Proceedings.)
- Tr. Cumberl. Westmorl. Ass.—Transactions of the Cumberland and Westmoreland Association for the Advancement of Literature and Science (Carlisle).
- Tr. Dumfries Nat. Hist. Soc.—The Transactions and Journal of Proceedings of the Dumfries and Galloway Natural History and Antiquarian Society (Dumfries).
- Tr. Edin. Geol. Soc. Transactions of the Edinburgh Geological Society.
- Tr. Edinb. Nat. Soc.—Transactions of the Edinburgh Field Naturalists' and Microscopical Society.
- Tr. Ess. Club = Ess. Nat.—Essex Naturalist: being the Journal, Transactions, and Proceedings of the Essex Field Club (Buckhurst Hill).
- Tr. E. Kent Nat. Hist. Soc.—Transactions of the East Kent Natural History Society (Canterbury).
- Tr. E. Soc.—Transactions of the Entomological Society of London.
- Tr. Hertf. Soc.—Transactions of the Hertfordshire Natural History Society and Field Club (Watford).
- Tr. Kansas Ac. Transactions of the Kansas Academy of Science (Topeka).

- Tr. Leicester Soc.—Transactions of the Leicester Literary and Philosophical Society (Leicester).
- Tr. L. S.—Transactions of the Linnean Society, London. (Also Journal.)
- Tr. L. S. New York—Transactions of the Linnean Society of New York.
- Tr. Manch. Geol. Soc.—Transactions of the Manchester Geological Society.
- Tr. Manch. Micr. Soc.—Transactions of the Manchester Microscopical Society.
- Tr. Maryland Ac. Sci.—Transactions of the Maryland Academy of Sciences.
- Tr. Norw. Soc.—Transactions of the Norfolk and Norwich Naturalists' Society (Norwich).
- Tr. Notts. Soc.—Transactions of the Nottingham Naturalists' Society.
- Tr. N. York Ac. Sci.—Transactions of the New York Academy of Sciences (New York). (Also Annals.)
- Tr. N. Z. Inst.—Transactions and Proceedings of the New Zealand Institute (Wellington).
- Tr. Odont. Soc.—Transactions of the Odontological Society.
- Tr. Ottawa Nat. Club—Transactions of the Ottawa Field-Naturalists' Club.
  (Also called Ottawa Naturalist.)
- Tr. & P. Pertheh. Soc.—Transactions and Proceedings of the Perthshire Society of Natural Science (Perth).
- Tr. R. Irish Ac.—Transactions of the Royal Irish Academy (Dublin).
- Tr. S. African Phil. Soc.—Transactions of the South African Philosophical Society (Cape Town).
- Tr. R. Soc. Canada—Proceedings and Transactions of the Boyal Society of Canada (Montreal).
- Tr. R. Soc. Edinb,—Transactions of the Royal Society of Edinburgh.
- Tr. R. Soc. S. Austr.—Transactions of the Royal Society of South Australia (Adelaide).
- Tr. R. Soc. Vict.—Transactions of the Royal Society of Victoria. (Also Proceedings.)
- Tr. Shropshire Soc.—Transactions of the Shropshire Archeological and Natural History Society (Shrewsbury and Oswestry).
- Tr. Stirling Soc.—Transactions of the Stirling Natural History and Archeological Society.
- Tr. St. Louis Acad.—Transactions of the Academy of Science of St. Louis.
- Tr. Texas Ac.—Transactions of the Texas Academy of Science.
- Tr. Wagner Inst.—Transactions of the Wagner Free Institute of Science of Philadelphia.
- Tr. Wisconsin Acad. Transactions of the Wisconsin Academy of Sciences, Arts, and Letters (Madison). (Also Bulletin.)

- Tr. Woolhope Nat.—Transactions of the Woolhope Naturalists' Field Club (Hereford).
- Tr. Yorksh. Union—Transactions of the Yorkshire Naturalists' Union (London & Leeds).
- Tr. Z. S.—Transactions of the Zoological Society (London).
- Trav. Inst. Zool. Lille—Travaux le l'Institut Zoologique de Lille et de la Station maritime de Wimereux (Lille).
- Trav. Lab. Histol.—Travaux du Laboratoire d'Histologie du Collège de France, École pratique des Hautes Études (Ranvier : Paris).
- Trav. Soc. Univ. Kharkow—Travaux de la Société des Naturalistes à l'Université impériale de Kharkow.
- Trav. Soc. Varsovie.—Travaux de la Société des Naturalistes de Varsovie. Comptes rendus de la section biologique. (Also Protokolui otdycleniya biologhii [Warsaw].)
- Trencsén term. egy. Trencsén megzei természettudomanyi egysület (Trencsin).
- Tromsö Mus. Aarsh.—Tromsö Museum's Aarshefter.
- Trudui Kazan. Univ. Trudui obshchestva estestvoispuitatelei pri imperatorskom Kazanskom Universitetye.
- Trudui Kharkoff Univ. Tru lui obshchestva ispuitateleĭ prihody pre imp. Kharkovakom Universitetye (Kharkoff).
- Trudui obsch. est.—Trudui obshchestva estestvolspuitatelei pri Imperatorikom Varshavskom Universitetye (Warsaw).
- Trudui Russ. Est.—Trudui sezda russkich estestvolspuitatelei.
- Trudui St. Petersburg Est.—Trudui sanct Petersburghskagho obshchestva estestvoispuitatelei. (Called also Travaux de la Société des Naturalistes de St. Petersburg.)
- Unters. Nat.—Untersuchungen zur Naturlehre des Menschen und der Thiere (Moleschott: Giessen).
- Upsala Univ. Årsskrift—Upsala Universitets Årsskrift (Upsala). (Also Act. Upsala.)
- Verh. Ak. Amst.—Verhande!ingen der koninklijke Akademie van Wetenschappen (Amsterdam). (Also JB.)
- Verh. Anat. Ges. Verhaudlungen der anatomischen Gesellschaft (Jena: published with Anat. Ans.).
- Verh. anthrop. Ges.—Verhandlungen der Berliner authropologischen Gessellschaft (Berlin).
- Verh. deutsch. Naturf. Verhandlungen der Gesellschaft deutscher Naturforscher und Arzte (Leipzig or elsewhere).
- Verk, deutsch. odont. Ges.—Verhandlungen der deutschen odontologischen Gesellschaft.
- Verh. deutsch. Zool. Ges.—Verhandlungen der deutschen zoologischen Gesellschaft.

- Verh. geol. Reichsanst.—Verhandlungen der k.-k. geologischen Reichsanstalt (Vienna). (Also JB.)
- Verh. Ges. Basel—Verhandlungen der naturforschenden Gesellschaft in Basel (Båle).
- Verh. Ges. Würzb. Verhandlungen der physikalisch medicinischen Gesellschaft zu Würzburg. (Also SB.)
- Verh. k, Leopold, Carol. Akad. Naturf.—(See N. Act. Ac. Leop. Carol. Nat. cur.)
- Verh. naturw. Ver. Karlsruhe Verhandlungen des naturwissenschaftlichen Vereins in Karlsruhe.
- Verh. schweiz. Naturf. Ges.—(See Act. Soc. Helv.)
- Verh. Siebenb. Ver.—Verhandlungen und Mittheilungen des Siebenburgischen Vereins für Naturwissenschaften (Hermannstadt).
- Verh. Ver. Brünn-Verhandlungen des naturforschenden Vereins in Brünn.
- Verh. Ver. Hamb.—Verhandlungen des Vereins für naturwissenschaftliche Unterhaltung zu Hamburg.
- Verh. Ver. Heidelb.—Verhandlungen des naturhistorisch-medieinischen Vereins zu Heidelberg.
- Verh. Ver. Presburg.—Verhandlungen des Vereins für Natur- und Heilkunde zu Presburg.
- Verh. Ver. Rheinl.—Verhandlungen des naturhistorischen Vereins der preussischen Rheinlande und Westphalen (Bonn). (Also CB.)
- Verh. z.-b. Wien—Varhandlungen der kaiserlichen-königlichen zoologischbotanischen Gesellschaft in Wien (Vienna). (Also SB.)
- Verh. d. Ver. Santiago—Verhandlungen des Deutschen wissenschaftlichen Vereines zu Santiago (Santiago, Chili).
- Versl. Ak. Amst.—Verslagen en Mededeelingen der k. Akademie van Wetenschappen (Amsterdam).
- Vict. Nat.—Victorian Naturalist, The Journal and Magazine of the Field Naturalists' Club of Victoria (Melbourne).
- Vid. Medd.—Videnskabelige Meddelelser fra den naturhistoriske Forening (Copenhagen).
- Viert. Ges. Zürich—Vierteljahrschrift der naturforschenden Gesellschaft in Zürich (Wolf: Zürich).
- West Am. Scientist—The West American Scientist (San Diego).
- Wien. ent. Z.—Wiener entomologische Zeitung (Vienna).
- Wilt. Mag.—The Wiltshire Archeological and Natural History Magazine (Devizes).
- York. Phil. Soc.—Yorkshire Philosophical Society Annual Report (York).
- Z. Biol.—Zeitschrift für Biologie (München).
- Z. Ent. Bresl.—Zeitschrift für Entomologie (Breslau).

- Ferdinand. Tirol Vorarlberg—Zeitschrift des Ferdinandeums für Tirol und Vorarlberg (Innsbrück).
- Z. f. Ethnol.—Zeitschrift für Ethnologie (Bastian & Hartmann: Berlin).
- Z. f. Thiermed.—Zeitschrift für Thiermedicin und vergleichende Pathologie).
- Z. geol. Ges. Zeitschrift der deutschen geologischen Gesellschaft (Berlin).
- Z. ges. Orn.—Zeitschrift für die gesammte Ornithologie (Von Madarász: Budapest).
- Z. Naturw.—Zeitschrift für Naturwissenschaften. Original Abhandlungen und Berichte herausgegeben im Auftrage der naturwissenschaftlichen Vereins für Sachsen und Thüringen (Brass: Hallea-S.).
- Z. Orn.—Zeitschrift für Ornithologie und praktischen Geflug.
- Z. wiss. Mikr.—Zeitschrift für wissenschaftliche Mikroscopie und für mikroscopische Technik (Berens: Braunschweig).
- Z. wiss. Zool.—Zeitschrift für wissenschaftliche Zoologie (Kölliker & Ehlers: Leipzig).
- Zapiski Kiev. Zapiski Kievskagho Obshchestva Estestvolspuitatelei (Kieff).
- Zupiski Novoross. Obsch. Estestv.—Zapiski Novorossiskagho Obshchestva Estestvolspuitatelei (Odessa).
- Zoe-Zoe: A Biological Journal (San Francisco).
- Zool.—The Zoologist (Harting: London).
- Zool. Anz.—Zoologischer Anzeiger (Carus: Leipzig).
- Zool. Beitr.—Zoologische Beiträge (Breslau).
- Zool. Gart.—Der Zoologische Garten (Noll: Frankfort).
- Zool. Jahrb.—Zoologische Jahrbücher (Spengel: Giessen).
- Zool. JB.—Zoologischer Jahresbericht (Mayer & Giesbrecht: Berlin).
- Zool. Mag.—The Zoological Magazine (Tokyo).
- Zool. Rec.—The Zoological Record (Zool. Soc. : London).
- Zool, Vort.—Zoologische Vorträge herausgegeben von William Marshall (Leipzig).

An asterisk prefixed to a quotation signifies that the Recorder has not seen the journal or work referred to.

# ERRATA.

INSECTA, p. 107, top line, for "Pryacoderus," read "Pyracoderus."

INSECTA, p. 139, line 11 from top, after "Tactoderus" add "n. g."

INSECTA, p. 139, line 12 from top, transfer "Tanophthalmus costifer" to p. 158, it being a Curculionid.

# CONTENTS.

I. GENERAL SUBJECTS.	By J. ARTHUR THOMSON, M.A.
Page         Titles        1         Subject-Index        36         Text Books, Essays, & General       36         Cell and Protoplasm        42	Oogenesis, Spermatogenesis, Fertilisation
II. MAMMALIA. By R.	LYDEKKER, B.A., F.G.S.
Introduction       1         The General Subject       2         Faunas       21         Special Structures, Development, &c.       21         Special Work       22         Primates       22         Chiroptera       25         Insectivora       27	Carnivora       28         Rodentia       32         Ungulata       36         Cetacea       48         Sirenia       51         Edentata       51         Marsupialia       52         Monotremata       54         Multituberoulata       55
III. AVES. By R. Bo	WDLER SHARPE, LL.D.
Titles       1         Faunistic       24         Anatomy and Miscellaneous       24         Systematic       24         Ratitie       24         Struthioniformes       24         Casuariiformes       24         Dinornithiformes       25         Appornithiformes       25         Carinate       25	Carinatæ         32           Lariformes         32           Ægialitornithiformes         32           Charadriformes         33           Gruiformes         34           Pelargiformes         35           Phænicopteriformes         35           Anseriformes         35           Pelecaniformes         36           Cathartidiformes         36
Carinates       25         Crypturiformes       25         Galliformes       25         Columbiformes       28         Opisthocomiformes       29         Ralliformes       29         Podicipedidiformes       31         Colymbiformes       31         Hesperornithiformes       31         Procellariiformes       31         Alciformes       32	Cathartidiformes       36         Accipitriformes       37         Striges       39         Coraciiformes       40         Trogones       46         Coccyges       46         Psittaciformes       47         Scansores       48         Eurylsemi       49         Menures       49         Passeriformes       49

#### IV. REPTILIA AND BATRACHIA. By G. A. BOULENGER. Page . . . . . . . . . . Emydosauria . . . . . . . . . 31 Works and Papers . . . . . . Chelonia . . . . . . . . . . . . . . . . 16 32 Faunæ . . . . . . . . . . . . . . . . 17 33 Ichthyosauria . . . . . . . . . Palseontological . . . . . . . . . 20 Plesiosauria . . . . . . . . . . 33 20 Thecodontia . . . . . . . . . . REPTILIA . . . . . . . . . . . . . . . 34 Squamata . . . . . . . . . . . . 21 Acrosauria . . . . . . . . . . . . . . . 34 Lacertilia....... 21 Rhynchocephalia . . . . . . . 34 Rhiptoglossa . . . . . . . 25 Anomodontia . . . . . . . . . 35 Dolichosauria . . . . . . . 25 35 25 Ecaudata ..... Pythonomorpha . . . . . . 36 26 Caudata . . . . . . . . . . . . . 39 Ornithosauria . . . . . . . . . 30 Apoda . . . . . . . . . . . . . . . 41 Dinosauria . . . . . . . . . . . 31 Stegocephala . . . . . . . . . . . . V. PISCES. By G. A. BOULENGER. Ostracodermi ...... General Subject ..... 15 Antiarcha . . . . . . . . . . . . . Osteostraci . . . . . . . . . . . Faunæ . . . . . . . . . . . . . . . . 17 Palseontology . . . . . . . . . . . . . . . Heterostraci ...... 19 35 Teleostomi....... 20 Holocephali . . . . . . . . . . . . . . . . . . 35 Teleostei . . . . . . . . . . . . . . . . . 20 Elasmobranchii . . . . . . . . . 35 Acanthopterygii . . . . . . . 20 Acanthodii . . . . . . . . . . . . . . 24 Selachii ....... A. Pharyngognathi. . . . . . 35 Anacanthini ...... 24 Asterospondyli..... 36 Physostomi . . . . . . . . . . . Tectospondyli . . . . . . . 25 36 Plectognathi . . . . . . . . . . . . Ichthyotomi....... 31 37 Ganoidei........ Incertæ sedis . . . . . . . . . . 32 37 Crossopterygii . . . . . . . . . 33 Cyclostomi . . . . . . . . . . . . 38 Leptocardii...... 33 VI. TUNICATA. By Prof. W. A. HERDMAN, D.Sc. List of Publications. . . . . . Geographical Distribution . . . 1 Anatomy, &c. . . . . . . . . . . . . . . . VII. MOLLUSCA. By B. B. WOODWARD, F.G.S., F.R.M.S. Gastropoda . . . . . . . . . . . 56 Anatomy, Physiology, &c. . . . Pulmonata . . . . . . . . . 29 56 37 Stylommatophora . . . . . 56 Geographical . . . . . . . . . Basommatophora . . . . . 37 67 Geological . . . . . . . . . . . . . . . Opisthobranchiata . . . . . 48 68 Nudibranchiata . . . . . 53 68 Tectibranchiata . . . . . Cephalopoda . . . . . . . . . . . . 53 69 Dibranchiata . . . . . . . . . Prosobranchiata . . . . . . 53 70 Octopoda...... Nucleobranchiata . . . . 53 70

54

54

55

Decapoda . . . . . . . .

Ammonea . . . . . . . .

Tetrabranchiata . . . . . .

Pectinibranchiata . . . .

Scutibranchiata . . . . .

Polyplacophora.....

71

84

MOLLUSC	A—continued.
Page	Page
Aplacophora 89	Pelecypoda—
Scaphopoda 89	Pseudolamellibranchiata . 94
Pelecypoda 90	Filibranchiata 95
Septibranchiata 90	Protobranchiata 95
Eulamellibranchiata 90	Incerta sedes 96
VIII. BRACHIOPODA. By B	B. WOODWARD, F.G.S., F.R.M.S.
Titles 1	Systematic—
Anatomy, &c 3	Inarticulata 5
Distribution—Geographical 4	Articulata 6
Geological 4	
IX. BRYOZOA (POLYZOA). B	y B. B. Woodward, F.G.S, F.R.M.S.
· · · · · · · · · · · · · · · · · · ·	
Titles	Systematic— Entoprocta4
Distribution—Geographical 3	
Geological 3	Ectoprocta 4
X. CRUSTACEA (including	JIGANTOSTRACA and PANTOPODA).
	NES POCOCK.
Titles 1	Edriophthalma.
Anatomy, Physiology, Embry-	Cumace 25
ology, Distribution 13	Amphipoda 25
Systematic 16	Isopoda 27
Malacostraca.	Entomostraca.
Podophthalma.	Phyllopoda 29
Decapoda.	Cladocera 29
Brachyura 16	Ostracoda 29
Oxystomata 19	Copepoda 30
Anomura 20	Cirripedia 31
Macrura 22	Pantopoda 32
Schizopoda 24	Gigantostraca
XI. ARACHNIDA.	By R. INNES POCOCK.
Titles of Papers 1	Systematic—
Anatomy, Embryology, Physi-	Pedipalpi 14
ology, Habits, &c 11	Araneæ 14
Distribution 13	Pseudoscorpiones 33
Systematic—	Opiliones 34
Scorpiones 14	Acari 35
XII. MYRIOPODA AN	D PROTOTRACHEATA.
	NES POCOCK.
Myriopoda—	Systematic—
Titles of Papers 1	Chilopoda 3
Anatomy, Physiology, Phy-	Diplopoda 4
logeny, &c	Prototracheata 6
Distribution	

# XIII. INSECTA. By D. SHARP.

	Page	Page		
Introduction	1   Economic	83		
Titles 2 Bibliography, Nomenclature,				
Anatomy, Physiology,	Biology 63 Biograp	h <b>y</b> 86		
Faunistic and Palsonto	logy 70 Systematic	88		
_ =====================================	Mg To   Systematic			
	(a.) COLEOPTERA.			
Pago	Page	Page		
Cicindelidæ 88	Lathridiidæ, My-	Melandryidæ,		
Carabidæ 90		1		
Amphizoidæ, Hali-	cetophagidæ 110 Dermestidæ,	Lagriide 141		
	l — • '	Pedilidæ, Xylophi-		
plidæ, Dytis- cidæ98	Byrrhidæ 110	lidæ 141		
	Parnidse, Hetero-	Anthicidæ 141		
	ceridæ 110	Mordellidæ, Rhi-		
Hydrophilidæ 100	Lucanidæ, Passa-	pidophoridæ,		
Staphylinidæ 101	lidæ 111	Stylopids 142		
Pselaphidæ 103	Scarabæidæ 111	Cantharida 142		
Scydmænidæ 103	Buprestides 122	Œdemeridæ 145		
Paussidæ 103	Throscidæ, Monom-	Curculionida 145		
Silphidæ 104	midæ, Eucnemidæ 126	Cerambycidae 159		
Trichopterygidæ . 104	Elateridæ 126	Scolytida 168		
Historidæ 104	Dascillidæ 129	Brenthidse 169		
Phalacride 106	Malacodermata. 129	Anthribidæ 170		
Nitidulidæ 107	Cleridæ 131	Bruchidæ 170		
Trogositide 108	Bostrichidæ,	Chrysomelidse 170		
Colydidæ 108	Ptinidæ, Cioidæ 133	Erotylidæ 182		
Cucujidæ, Rhysso-	Tenebrionidæ 133	Endomychida 182		
didse 109	Cistelidse 140	Coccinellide 183		
Crytophagids 109	Į.	Corylophida 185		
	(b.) HYMENOPTERA.			
Aculeata	Terebrantia !	Terebrantia		
Apidæ 186	Ichneumonidæ. 202			
		Chalcidide 206		
Vespidæ 188	Braconidse 205	Proctotrupidæ. 207		
Fossoria 189	Evaniidæ, Ste-	Cynipidæ207		
Formicide 196	phaniidæ,Mo-	Sessiliventres		
Terebrantia	nomachidæ,	Siricidae 208		
Chrysididæ201	Trigonalidm . 206	Teuthredinidæ. 208		
		•		
	(c.) LEPIDOPTERA.			
	· · ·			
General 211	Rhopalocera	Rhopalocera		
Rhopalocera	Nymphalidæ 215	Hesperiidse 224		
Papilionidæ 211	Morphidæ 218	Heterocera		
Pieridæ 213	Satyridæ 219	General 226		
Danaidæ 214	Elymniidæ, Ery-	Sphingida 226		
Acræidæ, Heli-	cinidæ 221	Ægeriidæ 227		
coniidæ215	Lycænidæ 221	Castniidæ229		

## CONTENTS.

## LEPIDOPTERA—continued.

He erocera Bombyces	Heterocera Pyralidæ, Crambidæ, Phycitidæ, Siculodidæ 273	Heterocera Tortricidæ 279 Tineidæ 280 Agdistidæ, Pterophoridæ 282		
	(d.) DIPTERA.			
General 283 Cecidomylidæ 283 Mycetophilidæ, Simuliidæ, Bibionidæ, Blepharoceridæ 284 Culicidæ, Chironomidæ 285 Tipulidæ 285	Xylophagidæ, Stratiomyiidæ . 285 Tabanidæ 286 Nemestrinidæ, Bombyliidæ 287 Asilidæ 289 Leptidæ, Therevidæ, Empidæ 289 Dolichopodidæ 290 Syrphidæ 290	Catridæ 292 Conopidæ, Pipun- culidæ, Platy- pezidæ 293 Muscidæ calyp- tratæ 293 Muscidæ acalyp- tratæ 301 Phoridæ 302 Pupipara 302		
	( ) Barriaga			
Heteroptera	(e.) RHYNCHOTA.  Heteroptera Reduviidæ 309 Saldidæ, Cimicidæ 310 Capsidæ 311 Hydrocorisæ 312 Homoptera 312 Cicadidæ 312 Fulgoridæ 314	Homoptera Cercopidæ 315 Membracidæ 315 Jassidæ 316 Psyllidæ 316 Aphididæ 316 Coccidæ, Aleurodidæ 317		
(f.) Neuroptera.				
Trichopters 320 Planipennia 320 Thysanura(Aptera)320	Mallophaga, Ter- mitidæ, Psocidæ 321	Ephemeridæ 321 Odonata 322		
(g.) ORTHOPTERA.				
Forficulidæ 325 Blattidæ 325	Mantidæ 326 Phasmatidæ 327 Gryllidæ 327	Locustides 327 Acridiides 331		

XIV. ECHINODERMATA. By F. A. BATHER.				
Titles       Page         1       1         Biology       5         Distribution—       7         Geographical       7         Geological       8         Systematic       9         Holothurioidea       9	Echinoidea         Page           Asteroidea         10           Application         16           Ophiuroidea         17           Crinoidea         18           Cystidea         21           Blastoidea         22			
XV. VERMES. By FLORENCE BUCHANAN, B Sc.				
Titles	Systematic— Trematoda 66 Nemertinea 69 Turbellaria 70 Nematoda, Acanthocephala and Chestognatha— Biological 73			
Faunistic       39         Systematic       43         Gephyrea       43         Chestopoda       44         Hirudinea       51         Platyhelminthes	Faunistic       79         Systematic       80         Nematoda       80         Acanthocephala       81         Chætognatha       82			
Biological       53         Faunistic       61         Systematic       63         Cestoda       63	Biological       82         Faunistic       84         Systematic       85         Mesozoa       88			
XVI. CŒLENTERATA. By List of Papers 1 Morphology 4 Faunistic 9	S. J. Hickson, M.A., D.Sc.  Systematic—  Hydrozoa			
XVII. SPONGIÆ. By	R. HANITSCH, Ph.D.			
List of Publications 1 Anatomy, Embryology, Physiology, Biology	Systematic—       15         Calcarea       15         Hexactinellida       17         Tetractinellida       17         Monaxonida       18         Keratosa       24         Incertæ sedis       24			
XVIII. PROTOZOA. By R. HANITSCH, Ph.D.				
List of Publications       1         Anatomy, Biology       14         Distribution—       16         Geological       17	Systematic— Gymnomyxa 18 Corticata 28 Incertæ sedis 32			

INDEX TO GENERA AND SUBGENERA DESCRIBED AS NEW, &C.

# THE ZOOLOGICAL RECORD

FOR 1892.

# I. GENERAL SUBJECTS.

BY

J. ARTHUR THOMSON, M.A.

### PLAN OF RECORD.

I .- TITLES.

II.—Subject-Index.

- A. Text-Books, Essays, and General.
- B. Cell and Protoplasm.
- c. Oogenesis, Spermatogenesis, Fertilisation.
- D. Sex, Reproduction, and Heredity.
- E. Evolution.

### TITLES.\*

- AAS, E. Sjaeleliv og Intelligens hos Dyr. Del 1. Pattedyr og Fugle. Christiania: 8vo, pp. 1-192, illustr.
- AGASSIZ, A. General sketch of the Expedition of the "Albatroes" from February to May, 1891. Report on the Dredging operations off the West Coast of Central America to the Galapagos, to the West Coast of Mexico, and in the Gulf of California. Bull. Mus. C. Z. xxiii (1892), pp. 1-89, 22 pls.
- Alcock, A. A case of Commensalism between a Gymnoblastic Anthomedusoid (Stylactis minoi, n. sp.) and a Scorpsenoid Fish (Minous inermis). Ann. N. H. x, pp. 207-214, 1 pl.

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

### I. GENERAL SUBJECTS.

4. [Alcock, A.] On the Habits of Gelasimus annulipes, Edw. Ann. N. H. 1892, pp. 415 & 416.

Re Sexual Selection.

- ALLEN, J. A. Definite versus Fortuitous Variation. (Discussion Amer. Soc. Nat.) Am. Nat. xxvi, pp. 85-91.
- ALTMANN, A. Ein Beitrag zur Granulalehre. Verh. Anat. Ges. 1892, pp. 220–223.
- Ueber Kernstructur und Netzstructuren. Inhalt eines Vortrages, gehalten auf der Wiener Anatomen Versammlung 1892.
   Arch. Anat. Entw. 1892, pp. 223–230.
- \*Ammon, O. La selection naturelle chez l'homme. L'Anthropologie, 1892, ii, pp. 720-736.
- Andrews, E. A. Experimental Embryology. Am. Nat. xxvi, pp. 367-382 & 580-592.
- 10. —. Heliotropic Animals. T. c. pp. 400-407.
- 11. —. Bifurcated Annelids. T. c. pp. 725-731.
- Anon. Chronophotography, or Photography as applied to Moving Objects. Nature, xlv, pp. 228-232, 11 figs.
- Discussion über Verbreitung pelagischer Thierformen.
   Verh. deutsch. Zool. Ges. 1892, pp. 119-121.
- 14. APATHY, S. Contractile und leitende Primitivfibrillen. MT. z. Stat. Neap. x, pp. 355-375, 1 pl. General Histology.
- Kritische Bemerkungen über das Frenzel'sche Mesozoon Salinella. Eine biologische Skizze. Biol. Centralbl. xii, pp. 108-123.
- Critical Observations on Frenzel's Mesozoon Salinella: a Biological Sketch. Ann. N. H. ix, pp. 465-481.
- Translation of above.
- 17. APLIN, O. V. Albinism in Birds and Mammals. Zool. xvi, p. 191.
- APSTEIN, C. Quantitative Plankton-Studien im Süsswasser. Biol. Centralbl. xii, pp. 484-512 & 608.
- Das Plankton des Süsswassers und seine quantitative Bestimmung. Schr. Nat. Ver. Schleswig, ix, pp. 267-273.
- Andissone, F. L'Organismo vivente considerato nella sua essenza et nella sua origine. Varese: 8vo, 24 pp.
- 21. ARNDT, B. Biologische Studien. I. Das biologische Grundgesetz. Greifswald: 8vo, ix & 203 pp.
- CARNOLD, A. F. Beiträge zur Kenntnis des Reptilien-Ovariums. Inaug. Diss. Erlangen, 8vo, 39 pp.

- Assileton, R. On the Development of the Optic Nerve of Vertebrates, and the Choroidal Fissure of Embryonic Life. Q. J. Micr. Sci. xxxiv, pp. 85-104, 2 pls.
  - ---. [See also BEARD (37).]
- Aurivillius, C. W. J. Analyse d'un mémoire intitulé du déguisement des Décapodes oxyrhyniques à l'aide d'adaptations singulières du corps. Ann. Sci. Nat. xiii, pp. 343-348.
- AYERS, H. Vertebrate Cephalogenesis. II. A Contribution to the Morphology of the Vertebrate Ear, with a Reconsideration of its Functions. J. Morph. vi, pp. 1-360, 12 pls.
- BALBIANI E. G. Nouvelles recherches expérimentales sur la mérotomie des Infusoires ciliés. Ann. Micrograph. 1892, 79 pp., 3 pls.
- Ballowitz, E. Die Bedeutung der Valentin'schen Querbänder am Spermatozoenkopfe der Säugethiere. Arch. Anat. Phys. 1891, pp. 193-211, 1 pl.
- 28. BANKS, N. Mimicry in Spiders. P. E. Soc. Wash. ii, pp. 174-176.
- von Bardeleben, K. Ueber Spermatogenese bei Säugetieren, besonders beim Menschen. Verh. Anat. Ges. 1892, pp. 202-208.
- 30. BATAILLON, E. Sur le determinisme physiologique de la métamorphose chez le ver à soie. C.R. cxv, pp. 61-64.
- 31. —. Recherches anatomiques et expérimentales sur la métamorphose des Amphibiens anoures. Paris : 127 pp., 6 pls.
- 32. Bates, H. W. The Naturalist on the River Amazon. With a Memoir of the author, by Edward Clodd. Reprint of the unabridged edition, with map and numerous illustrations. London: 8vo, 478 pp.
- 33. Bateson, W. On Numerical Variation in Teeth, with a discussion of the Conception of Homology. P. Z. S. 1892, pp. 102-115, 6 figg.
- 34. —. [Letter to Nature.] The Alleged Aggressive Mimicry of Volucelle. Nature, xlvi, p. 585.
  - Cf. Poulton, op. cit. xlvii, p. 28; Bateson, t. c. p. 77; Poulton, t. c. p. 126.
    - BATHER, F. A. [See BUCKMAN & BATHER (79).]
- 35. Beale, L. S. Protoplasm, Physical Life, and Law; or, Nature as viewed from without. London: 4th Edition, 8vo, 286 pp.
- Beard, J. The Histogenesis of Nerve. Anat. Ana. vii, pp. 290-302.
- [Letter to Nature.] On a supposed Law of Metazoan Development. Nature, xlvii, p. 79, and Anat. Anz. viii, pp. 22.
   Cf. Assileton, R., t. c. p. 176.
- 38. —. Larvæ and their Relations to Adult Forms. Rep. Brit. Ass. 1892, p. 757.

- [Beard, J.] The Transient Ganglion Cells and their Nerves in Ruja batis. Anat. Anz. vii, pp. 191-206, 8 figg.
- Motes on Lampreys and Hags (Myzine). Anat. Anz. viii, pp. 59 & 60.

Re Hermaphroditism.

- 41. BEDDARD, F. E. Animal Colouration: an account of the principal facts and theories relating to the Colours and Markings of Animals. London: 8vo, 258 pp., 4 col. pls., illustrs.
- Some Recent Observations upon Mimicry. Nat. Sci. pp. 10-16.
- BEECHER, C. E. Development of the Brachiopoda. Part. II. Classification of the Stages of Growth and Decline. Am. J. Sci. xliv, pp. 133-155, 1 pl., figg.
- BEHRENS, W. Tabellen zum Gebrauch bei mikroskopischen Arbeiten.
   2nd Ed. Braunschweig: 8vo, vii & 205 pp.
- Benda, C. Ueber die Histiogenese des Sauropeidenspermatoscons.
   Verh. Anat. Ges. vi, 1892, pp. 195–199.
- 6—. Eine Mitteilung zur Samenbildung. Int. Centralbl. Phys. Path. Harn Sexual-organe, iv, pp. 23-25.
- Béraneck, E. Sur le nerf pariétal et la morphologie du troisième œil des Vertébrés. Anat. Anz. vii, pp. 674-689, 6 figg.
- 48. BERGH, R. S. Kritik einer moderner Hypothese von der Uebertragung erblicher Eigenschaften. Zool. Anz. xv, pp. 43-52.
- A Criticism of a Modern Hypothesis of the Transmission of Hereditary Characters. Ann. N. H. x, pp. 467-475.
   BERNARD, H. M. & M. [See Lang (318).]
- 50. Bertacchini, P. La Spermatogénèse chez la Rana temporaria, Arch. Ital. Biol. xvii, pp. 166 & 167.
  - BEZANÇON, P. [See VARIOT & BEZANÇON (500-502).]
- BIEDERMANN, W. Ueber den Farbenwechsel der Frösche. Arch. Ges. Phys. li, pp. 455-508, 1 pl. Re Colouration.
- BIEGELEBEN, A. VON. Betrachtungen über das Verhältnis der menschlichen Seele zur Thierseele. Frankfurt: 8vo, 27 pp.
- Bignell, G. C. Myrmica ruginodes making War on its own Species. Ent. M. M. xxviii, p. 135.
- BLANC, H. Note préliminaire sur la maturation et la fécondation de l'œuf de la Truite. Bull. Soc. Vaud. xxvii, pp. 272-275.
- Sur la maturation et la fécondation de l'œuf de la Truite des lacs. C.R. Soc. helv. Sci. Nat. lxxiv, pp. 55 & 56.

- Blanc, L. Un cas d'ovule à deux noyaux chez un Mammifère (Mus decumanus). C.B. Soc. Biol. iv. pp. 563 & 564.
- Note sur l'influence de la lumière sur l'orientation de l'embryon dans l'œuf de Poule. T. c. pp. 774-776.
- BLANCHARD, R. Deuxième Rapport sur la nomenclature des êtres organisés. Congrès Internat. Zool. ii, pt. 1, pp. 303-314. Cf. Bull. Soc. Z. Fr. xvii, pp. 196-200.
- 59. —. Règles de nomenclature adoptées par le Congrès Zoologique de Moscou. Zool. Ans. xv, pp. 440-442.
- 59A. BLES, E. J. Notes on the Plankton observed at Plymouth during June, July, August, and September, 1892. J. Mar. Biol. Ass. ii, pp. 340-343.
- 60. Bodington, A. Mental Evolution in Man and the Lower Animals (continued). Am. Nat. xxvi, pp. 482-494 & 593-606.

  Boettoer, O. [See Brehm (73).]
- 61. \*Böнм, A. A. Die Befruchtung des Forelleneies. SB. Ges. Morph. vii, pp. 65-73, figg.
- 62. \*Bommeli, R. Die Thierwelt. Eine illustrierte Naturgeschichte der jetzt lebenden Thiere. Stuttgart: 8vo, Heft 1, 64 pp.
- 63. BONNIER, G. Anatomie et Physiologie Animales. 1. Étude spécial de l'homme. 11. Les Organes et leurs fonctions dans la série animale. Paris: 18mo, 310 pp.
- BORDAGE, E. Les Vertébrés descendent-ils des Arthropodes. Rev. Sci. xlix, pp. 333-339, 9 figg.
- 65. Born, G. Die Reifung des Amphibieneies und die Befruchtung unreifer Eier bei Triton tæniatus. Anst. Anz. vii, pp. 772-781 & 803-811, 1 fig.
- 66. BORREL, A. De la division du noyau et de la division cellulaire dans les tumeurs épithéliales. J. de l'Anat. Phys. xxviii, pp. 129– 142, 2 pls.
- 67. Boulenger, G. A. Fecundation in the Tailed Batrachians: a summary Review of Recent Discoveries. Zool. Jahrb. vi, pp. 447-452.
- 68. BOVERI, TH. Ueber die Enstehung des Gegensatzes zwischen den Geschlechtszellen und den somatischen Zellen bei Ascaris megaloce-phala. SB. Ges. Morph. viii, pp. 114-125, 5 figg.
- OBRANDES, G. Ueber Mimicry und andere Anpassungserscheinungen im Thierreiche. CB. Ver. Sachs. Thür. 1891, pp. 168-171.
- BRANDT, A. Essai d'une classification des variations animales selon leurs causes. Congrès Internat. Zool. ii, pt. 1, pp. 66 & 67.
- BRANDT, K. Ueber Anpassungserscheinungen und Art der Verbreitung von Hochseethieren, in Krümmel's Reisebeschreibung d. Plankton Exp. pp. 338-370, 1 map.

- Brauer, A. Ueber das Ei von Branchipus Grubii v. Dyb. Von der Bildung bis zur Ablage. Abh. Ak. Berl. 1892, 66 pp., 3 pls.
- BREHM, A. E. Thierleben. Allgemeine Kunde des Thierreichs.
   gänzlich neubearbeitete Aufl. Hrsg. von Prof. Dr. Pechuel-Loesche. Bd. vi. (Birds, 3rd vol.), 740 pp., 20 pls., 106 figg., 3 maps (Pechuel-Loesche & W. Haacke). Bd. vii. (Reptilia and Amphibia), 825 pp., 16 pls., 167 figg., 1 map (Pechuel-Loesche & O. Bosttger). Bd. viii. (Fishes), 517 pp., 11 pls., 146 figg., 1 map (Pechuel-Loesche & W. Haacke). Bd. ix. (Insects), 764 pp., 21 pls., 287 figg., 1 map (E. L. Taschenberg).
- 74. °—. Thierleben. Kleine Ausgabe für Volk und Schule. 2. Auf. gänzl. neu bearb. von R. Schmidtlein. 1 Bd. Die Saügethiere. Leipzig: 8vo, xvi & 747 pp., 1 pl., 226 figg.
- 75. Bronn, —. Klassen und Ordnungen des Thierreichs.

The following divisions have made progress in 1892, viz.:—Cælenterata, by C. Chun; Echinoderms, by H. Ludwig; Mollusca, by H. Simroth, Vermes, by M. Braun; Crustacea, by A. Gerstaecker; Mammalia, by W. Leche.

See other Records.

- BROOKS, W. K., & HERRICK, F. H. The Embryology and Metamorphosis of the Macroura. Johns Hopk. Univ. Circ. xi, pp. 67 & 68.
   Re Poscilogony.
- 77. Brown-Séquard. Effets produits sur de nombreux états morbides par des injections sous-cutanées d'un extrait liquide retiré des testicules. C.B. cxiv, pp. 1318-1321. Cf. c. pp. 1399-1405 (Brown-Séquard & d'Arsonval), p. 1446 (Grigorescu).
- 78. Effets physiologiques d'un liquide extrait des glandes sexnelles, et surtout des testicules. T. c. pp. 1237-1242.
- 78A. ——. Hérédité d'une affection due à une cause accidentelle. Faits et arguments contre les explications et les critiques de Weismann. Arch. Phys. xxiv, pp. 686-688.
- BUCKMAN, S. S., & BATHER, F. A. The Terms of Auxology. Zool. Ans. xv, pp. 420, 421, & 429-434.
- BUCHNER, O. Die Asymmetrie der Gastropoden in ihren Beziehungen und Wirkungen auf die Lebenaüsserungen der Schalentragenden Schnecken. JH. Ver. Württ. xlviii, pp. 58-92.
- 81. OBÜCHNER, E. Ueber Vergangenheit und Zukunft des Menschengesschlechtes im Sinne der Entwickelungstheorie. Festschr. Ver. Rheinpfalz. 1892, pp. 48-65.
- BÜRGER, O. Was sind die Attraktionsphären und ihre Centralkörper. Anat. Anz. vii, pp. 222-231.

- 83. BÜTSCHLI, O. Untersuchungen über mikroskopische Schäume und das Protoplasma. Versuche und Beobachtungen zur Lösung der Frage nach den physikalischen Bedingungen der Lebenserscheinungen. Leipzig: 4to, iv & 232 pp., 6 pls., 23 figg., and atlas of 19 microphotographs.
- Ueber die sogenannten Centralkörper der Zelle und ihre Bedeutung. Verh. Ver. Heidelb. iv, pp. 535-538.
- 85. Burckhardt, R. Das Centralnervensystem von Protopterus annectens. Eine vergleichend-anatomische Studie. Berlin: 8vo, 64 pp., 5 pls.
  - Chapters on Phyletic History of the Brain, Metamerism, &c.
- 86. Buscalioni, L. Sulla frammentazione nucleare seguita dalla divisione della cellula. Giorn. Acc. Med. Torino, v, pp. 326-328.
- 87. 2—. Contribuzione allo studio della membrana cellulare. Malphighia, 1892, 90 pp., 2 pls.
- 88. CALDERWOOD, W. L. A Contribution to the Knowledge of the Ovary and Intra-ovarian Egg in Teleosteans. J. Mar. Biol. Ass. ii, pp. 298-313, 2 pls.
- 89. Campbell, H. J. Text-book of Elementary Biology. London: 8vo, 286 pp., 136 figg.
  - CANESTRINI. [See Morselli (366).]
- Ocarlet, G. Précis de zoologie médicale. 3 édit. entièrement refondue. Paris: 8vo, 722 pp., 636 figg.
- 91. CARLILE, W. W. Animal Intelligence. Tr. N. Z. Inst. xxiv, pp. 349-354.
- CATTANEO, G. Influenza del letargo sulle forme e i fenomeni delle cellule ameboidi negli Invertebrati. Atti Soc. Ligust. iii, pp. 248-251.
- 93. Chatin, J. La Cellule animale; sa structure et sa vie: étude biologique et pratique. Paris: 8vo, 304 pp., 149 figg.
- 94. Sur la membrane basilaire. C.R. Soc. Biol. iv, pp. 463 & 464.
- CHOLODKOVSKY, N. Contribution à la theorie du mésoderme et de la métamérie. Congr. Internat. Zool. ii, pt. 1, pp. 58-65. Cf. Rev. Sci. Nat. St. Pétersb. iii, pp. 204-209.
- 96. Chun, C. Die Dissogonic, eine neue Form der geschlechtlichen Zeugung. Feschr. R. Leuckart, pp. 77-108, 5 pls., 3 figg. Leipzig: 4to.
- 97. CLARKE, C. B. On Biologic Regions and Tabulation Areas. Phil. Trans. clxxxiii, pp. 371-387, 2 maps; P. R. Soc. l, pp. 472-474. 1892. [VOL. XXIX.]

- 98. CLARK, J. Natural Relations between Temperature and Protoplasmic Movement. Rep. Brit. Ass. 1892, p. 760.
- 99. Experimental Observations on the Function of the Nucleus in the Vegetable Cell. T. c. p. 761.
- COMSTOCK, J. H. The Descent of the Lepidoptera. An application of the Theory of Natural Selection to Taxonomy. (Abstract) P. Am. Ass. xli, pp. 199-201.
- 101. COPE, E. D. Definite versus Fortuitous Variation. (Discussion Amer. Soc. Nat.) Am. Nat. xxvi, pp. 85-91.
- On some points in the Kinetogenesis of the Limbs of Vertebrates.
   P. Am. Phil. Soc. xxx, pp. 282-285.
- 103. Cori, C. J. Ueber Anomalie der Segmentirung bei Anneliden und deren Bedeutung für die Theorie der Metamerie. Z. wiss. Zool. liv, pp. 569-578, 1 pl.
- 104. CORNEVIN, C. Études zootechniques sur la croissance. Arch. Phys. xxiv, pp. 477-487, cf. J. méd. vét. et zootechn. iii, pp. 449-460.
- 105. COSTE, F. H. PERRY. On Insect Colours. Nature, xlv, pp. 513-517, 541, 542 & 605.
- 106. —. Evolution of Insect Colours. Ent. xxv, pp. 28 & 29. —. [See also Hopkins (266).]
- 107. COULON, RAIMOND. Synthèse de transformisme. Description élémentaire de l'évolution universelle. Paris : 8vo, xiv & 215 pp.
- 108. Crato, E. Beitrag zur Kenntnis der Protoplasmastructur. Ber. Deutsche Botan. Ges. x, pp. 451-458.
- 109. CRETY, C. Le noyau vitellin dans les œufs des Trématodes. Arch. Ital. Biol. xvii, pp. 396-399.
- Intorno al nucleo vitellino dei Trematodi. Atti [Rend.] Acc. Rom. 1892, pp. 92-97.
- 111. CREUTZBURG, N. Ueber metaphysische Probleme in der Zoologie. Eine Kritik der Darwiu'schen Theorie. Abh. Ges. Halle, xvii, pp. 461-472.
  - CROTCH, W. DUPPA. [See ROMANES (435).]
- 112. CUÉNOT, L. Les organes phagocytaires chez quelques invertébrés. Arch. Z. expér. x, pp. ix-xi.
- 113. —. Notes sur les Echinodermes. I. Ovogénèse et Spermatogénèse. Zool. Anz. xv, pp. 121-125, figg.
- 114. ——. La valeur respiratoire de l'hemocyanine. C.R. cxv, pp. 127-129.
- CUNNINGHAM, J. T. The Evolution of Flat-Fishes. Nat. Sci. i, pp. 191-199.

- CURTEL, G. [See LANG, A. (318).]
- 116. Danilewsky, B. Ueber die physiologische Wirkung des Cocaïns auf wirbellose Thiere. Arch. ges. Phys. li, pp. 446-454.
- 117. Dannehl, P. Ueber die kadaverösen Veränderungen der Altmann'schen Granula. Arch. pathol. Anat. exxviii, pp. 485-496.
  Re Structure of Protoplasm.
- 118. Dareste, C. Note sur l'évolution de l'embryon de la poule soumis pendant l'incubation à un mouvement de rotation continue. C.R. cxv, pp. 137 & 138.
- 119. —. La tératogénie expérimentale. Rev. Sci. xlix, pp. 33-42, 19 figg.
- 120. DARWIN, CHARLES. Voyage of H.M.S. "Beagle" round the World. London: 8vo, 202 pp., illustr. Sixpenny Edition.
- 121. Darwin, F. Charles Darwin: His Life told in an Autobiographical chapter, and in a selected series of his published letters. London: 8vo, 346 pp., with portrait.
- 122. Dehner, H. Ueber die sogenannte parthenogenetische Furchung des Froscheies. Verh. Ges. Würzb. xxvi. Inaug. Diss. Würzburg: 8vo, 18 pp., 1 pl.
- 123. DETMER, W. Ueber die Natur und Bedeutung der physiologischen Elemente des Protoplasmas. Ber. Deutsche Botan. Ges. x, pp. 433-442.
- 124. DINEUR, E. Note sur la sensibilité des leucocytes à l'electricité. Bull. Soc. Belg. Micr. xviii, pp. 113-118.
- 125. DISTANT, W. L. A Naturalist in the Transvaal. London: 8vo, 16 & 277 pp., 5 pls., 29 figg.
- 126. Dixon, C. The Migration of Birds: an attempt to reduce Avian Season Flight to Law. London: 8vo, 306 pp. Instinct and Evolution.
- DIXON, E. T. [Letter to Nature.] The Limits of Animal Intelligence. Nature, xlvi, p. 392.
- Cf. C. LLOYD MORGAN, t. c. p. 417, and St. George Mivart, t. c. p. 466.
- 128. DREYER, F. Die Principien der Gerüstbildung bei Rhizopoden, Spongien, und Echinodermen. Ein Versuch zur mechanischen Erklärung organischer Gebilde. Jen. Z. Nat. xxvi, pp. 204-468, 14 pls.
- 129. Ziele und Wege biologischer Forschung beleuchtet an der Hand einer Gerüstbildungsmechanik. Jena: 8vo, xii & 103 pp, 6 pls.

- 130. DRIESCH, H. Entwicklungsmechanische Studien:
  - Die Verminderung des Furchungsmaterials und ihre Folgen (Weiteres über Theilbildungen).
  - Experimentelle Veränderungen des Typus der Furchung und ihre Folgen (Wirkungen von Wärmezufuhr und von Druck).
  - v. Von der Furchung doppelt befruchteter Eier.
  - vi. Ueber einige allgemeine Frage der theoretischen Morphologie. Z. wiss. Zool. lv, pp. 1-62, 3 pls.
- 131. ——. Entwicklungsmechanisches. Anat. Auz. vii, pp. 584-586. Am. Nat. xxvi, pp. 1042 & 1043.
- 132. —... Kritische Erörterungen neuerer Beiträge zur theoretischen Morphologie. Biol. Centralbl. xii, pp. 528-540, 545-556.
- 133. ——. Epigenesis or Evolution? (Abstract) Am. Nat. xxvi, pp. 178-180.
- 134. DUBOIS, R. Anatomie et Physiologie comparée de la Pholade dactyle (structure, locomotion, tact, olfaction, gustation, vision, dermatoptique, photogénie) avec une théorie générale des sensations. Paris: 8vo, x & 167 pp., 68 figg., 15 pls.
- 135. DUCLAUX, —. Sur une réaction donnée comme particulière à la spermine. C.R. cxv, pp. 549 & 550.
- 136. °DUTOIT-HALLER, —. Schöpfung und Entwickelung nach Bibel und Naturwissenschaft. Basel: 8vo, 44 pp.
- 137. °EBERTH, S. Referat über den Befruchtungsprocess bei Wirbellosen (Echinodermen) und Wirbeltieren (Selachier, Reptilien). Münch, med. Wochschr. xxxix, p. 371.
- 138. °----. Schlummerzellen und Gewebsbildung. T. c. pp. 755-758.
- Schlummerzellen und Gewebsbildung. Fortschr. Med. x, pp. 990-1000.
- 140. EBNER, V. von. Ueber die Beziehungen der Wirbel zu den Urwirbel. SB. Ak. Wien, ci, pp. 235-260, 1 pl.
- Edinger, L. Untersuchungen über die vergleichende Anatomie des Gehirns.
  - Das Vorderhirn (Neuabdruck, 1892). Frankfurt a-M.: 4to, 28 pp., 4 pls.
  - Das Zwischenhirn. Erster Theil. Das Zwischenhirn der Selachier und der Amphibien. Frankfurt a-M.: 4to, 95 pp., 5 pls.; both reprints from Abh. Senck. Ges.
- 142. —. Zwölf Vorlesungen über den Bau der nervösen Centralorgane. 3 Aufl. Leipzig: 8vo, viii & 196 pp., 139 figg.
- 143. \*EHRMANN, S. Zur Kenntnis von der Entwickelung und Wanderung des Pigmentes bei den Amphibien. Arch. Dermatol. xxiv, pp. 195-222, 1 pl.

- 144. [°EHRMANN, S.] Beitrag zur Physiologie der Pigmentzellen nach Versuchen am Farbenwechsel der Amphibien. Arch. Dermatol. xxiv, pp. 519-539, 1 pl.
  - Re Colouration.
- 145. EIMER, G. H. TH. Die Enstehung und Ausbildung des Muskelgewebes, insbesondere der Querstreifung derselben als Wirkung der Thätigkeit betrachtet. Z. wiss. Zool. liii, Suppl. pp. 67-111, 13 figg.
- Ueber die Enstehung der Arten. Vortrag. JH. Ver Württ. xlviii, pp. lxxxii & lxxxiii.
- 147. <sup>©</sup>EISMOND, J. Beitrag zur Frage über die Theilung des Zell-leibes. Arb. z. Lab. Warschau (1892), 20 pp.
- 148. ELLENBERGER, W. Handbuch der Vergleichende Histologie und Physiologie der Haussäugethiere, Band II. Vergleichende Physiologie der Haussäugetiere, Teil II. Berlin: 8vo, viii & 994 pp., 284 figg., 4 pls.
- 149. \*Elliot, D. G. The Inheritance of Acquired Characters. President's Address Amer. Ornithol. Union, 1891. Auk, ix, pp. 77-104.
- 150. —. Hybridism, and a Description of a Hybrid between Anas boschas and Anas americana. Auk, ix, pp. 160-166.
- 151. EMERY, C. Zur Morphologie der cyklopischen Missbildungen. Eine Hypothese betreffend den Schlundring der Wirbeltiere. Anat. Anz. viii, pp. 52-57.
- 152. Endres, —. Anatomisch-entwickelungsgeschichtliche Studien über die formbildende Bedeutung des Blutgefäss-Apparates unter besonderer Berücksichtigung der damit verbundenen mechanischen Einflüsse. Arch. mikr. Anat. xl, pp. 435-483, 3 pls.
- 153. °ENGEL, R. Nouveaux élements de chimie médicale et de chimie biologique. 4th Ed. Paris: 8vo, 2nd part, pp. 324-694, figg. 37-108.
- 154. \*\*ENGELMANN, T. W. [& OTHERS]. Festschrift. Beiträge zur Psychologie und Physiologie der Sinnesorgane. Hermann von Helmholtz als Festgruss zu seinem 70 Geburtstage dargebracht von Th. W. Engelmann, E. Javal, A. König, J. v. Kries, Th. Lipps, L. Matthiesen, W. Preyer, W. Uhthoff. Gesamm. u. hrsg. von A. König. Hamburg: 8vo, vii & 388 pp., 6 pls.
- 155. ESCHERICH, C. Die biologische Bedeutung der "Genitalanhängo" der Insecten (Ein Beitrag zur Bastardfrage). Verh. z.-b. Wien, zlii, pp. 225-240, 1 pl.
- 156. ESCHERICH, K. Ueber die Gesetzmässigkeit im Abändern der Zeichnung bei Insekten. Beitrag zur Theorie von der Entwicklung aus constitutionellen Ursachen. Deutsche e. Z. 1892, pp. 113-130, 1 pl.

- 156A. EWALD, C. A. The Influence of Light on the Gas Exchange in Animal Tissues. J. Physiol. xiii, pp. 847-859.
- 157. FABANI, C. Nota ad alcuni casi di anomalie nel colore degli uccelli. Riv. Ital. Sci. Nat. xii, pp. 3 & 4.
- Altre osservazioni intorno alle cause dell'albinismo anomalo e periodico. T. c. pp. 70 & 71.
- 159. —. Studi sul mimetismo. T. c. pp. 71 & 72.
- 160. \*FARGES, ALB. La Vie et l'Evolution des espèces, avec une thèse sur l'évolution étendue au corps de l'homme. 3 Ed. rev. et augm. Paris: 8vo, 320 pp.
- 161. FAYOD, V. Réponse aux rémarques de M. le professeur Guignard au sujet de ma communication sur la structure du protoplasme. C.R. Soc. Biol. iv, pp. 60-62.
- 162. FERTON, CH. Notes pour servir à l'histoire de l'instinct des Pompilides. Act. Soc. L. Bord. xliv, pp. 281-294.
- 163. Fick, R. Bemerkungen zu O. Bürger's Erklärungsversuch der Attraktionsphären. Anat. Anz. vii, pp. 464-467.
- 164. . Ueber die Befruchtung des Axolotleies. T. c. pp. 818-821.
- 165. FIELD, G. W. The Problem of Marine Biology. Am. Nat. xxvi, pp. 799-808; Nature, xlvi, pp. 623-625.
- 166. \*FISCHER-SIGWART, H. Das Gebirge, ein Rückzugsgebiet für die Thierwelt. MT. Aargau nat. Ges. 1892, pp. 111-196.
  FLEMING, G. [See NEUMANN, L. G. (374).]
- 167. FLEMMING, W. Zur Nomenclatur der Zellteilung. Anat. Anz. vii, pp. 26-32.
- Ueber Unsichtbarkeit lebendiger Kernstrukturen. T. c. pp. 758-764.
- 169. FLOWER, W. H. The Horse: a Study in Natural History. London: 8vo, 210 pp., figg.
- 170. \*Franceschini, R. Die Biologie als selbständige Wissenschaft. Hamburg: 8vo, 38 pp.
- 171. \*Francke, K. Das Leben der Zelle. Rede. München: 8vo, 9 pp.
- 172. Francken, C. J. Wijnaendts. Sexueele Evolutie. Tijdschr. Nederl. Dierk. Ver. iii, pp. 206-225.
- 173. Frederico, L. Sur l'hémocyanine. C.R. cxv, p. 61.
- Nouvelles recherches sur l'autotomie chez le crabe (Curcinus manas). Arch. Biol. xii, pp. 169-198.
- 175. FRENZEL, J. Ueber den Mitteldarm von Artemia. Ein Beitrag zur Lehre von der Verdauung und Resorption. Zool. Jahrb. (Abth. Anat.) vi, pp. 249-270, 1 pl., 11 figg.

- 176. [FRENZEL, J.] Beiträge zur vergleichenden Physiologie und Histologie der Verdauung. Arch. Anat. Phys. (Physiol. Abth.), 1892, pp. 81-114, 2 pls.
- 177. Die nucleoläre Kernhalbierung. Ein Beitrag zur Kenntnis des Zellkernes und der amitotischen Epithelregeneration. Arch. mikr. Anat. xxxix (1892), pp. 1-32, 1 pl.
- FRORIEP, A. Zur Entwicklungsgeschichte der Kopfnerven.
   Entwicklung der Trochlearis bei Torpedo. II. Kiemenspalten organe der Selachierembryonen. Verh. Anat. Ges. v (1891), pp. 55-65, 6 figg.
- 179. —... Ueber den Gebrauch der Worte proximal und distal. Anat. Anz. vii, pp. 764-767.
- 180. FUSARI, R. Sur les premières phases de développement des Téléostéens. Arch. Ital. Biol. xviii, pp. 204-239.
- 181. Gadow, H. Crop and Sternum of Opisthocomus cristatus: a contribution to the question of the Correlation of Organs and the Inheritance of Acquired Characters. P. R. Irish Ac. 1892, pp. 147-153, 2 pls.
- 182. Gage, S. H. The Comparative Physiology of Respiration. P. Am. Ass. xli, pp. 183-196.
- 183. —... The Comparative Physiology of Respiration. Am. Nat. xxvi, pp. 817-832. Cf. Nature, xlvi, pp. 598-601.
- 184. GARNER, R. L. The Speech of Monkeys. London: 8vo, 260 pp.
- 185. GAUBERT, P. Autotomie chez les Pycnogonides. Bull. Soc. Z. Fr. xvii, pp. 224 & 225.
- 186. °—... Sur l'autotomie chez les Araignées. Bull. Soc. Philom. iv, pp. 78 & 79.
- GAUDRY, A. Similitudes dans la marche de l'évolution sur l'ancien et le nouveau continent. C.R. cxv, pp. 11-13.
- 188. GAUTIER, A., & LANDI, L. Sur la vie résiduelle et les produits du fonctionnement des tissus séparés de l'être vivant. Op. cit. exiv, pp. 1048-1053.
- 189. & Sur les produits du fonctionnement des tissus et particulièrement des muscles, séparés de l'être vivant. Méthodes analytiques. T. c. pp. 1154-1159.
- 190. & . Sur la produits de la vie résiduelle des tissus, en particulier du tissu musculaire séparé de l'être vivant. T. c. pp. 1312-1317.
- 191. & Phénomènes de la vie résiduelle du muscle séparé de l'être vivant. Action physiologiques des bases musculaires. T. c. pp. 1449-1455.

- 192. GEDDES, P., & THOMSON, J. A. L'Evolution du Sexe. Trad. franç. par H. de Varigny. Paris: 18mo, x & 457 pp., illustr. (With some emendations.)
- 193. GEGENBAUR, C. Die Epiglottis. Vergleichend-anatomische Studie. Leipzig: 4to, vii & 69 pp., 2 pls., 15 figg.
- 194. °GERLACH, W. Ein Beitrag zur Lehre von den Elementarorganismen. St. Petersb. med. Wochschr. xvii, pp. 295-297, 1 fig.
- 195. Gehuchten, A. Van. Nouvelles recherches sur les ganglions cérébro-spinaux. Cellule, viii, pp. 235-253, 1 pl.
- Contributions à l'étude des ganglions cérébro-spinaux. Bull.
   Belg. xxiv, pp. 117-154, 11 figg.
- 197. GERMANO, E. Ricerche istologiche sul testicolo dall nascità alla maturità. Month. Int. J. Anat. Hist. ix, pp. 241-255, 1 pl.
- 198. GIACOMINI, E. Matériaux pour l'étude du développement du Seps chalcides. Arch. Ital. Biol. xvi, pp. 332-359, 1 pl.
- 199. GIARD, A. Nouvelles remarques sur la pœcilogonie. C.R. cxiv pp. 1549-1552.
- 200. —... Sur la persistance partielle de la symétrie bilatérale chez un Turbot (Rhombus maximus), et sur l'hérédité des caractères acquis chez les Pleuronectes. C.R. Soc. Biol. iv, pp. 31-34.
- GILSON, G. On the Affinity of Nuclein for Iron and other substances. Rep. Brit. Ass. 1892, pp. 778-780.
- 202. A Method of Staining Chromatin by chemical means. T. c. p. 780.
- 203. Girard, C. Observations sur quelques points de la nomenclature zoologique. Congres Internat. Zool. ii, pt. 1, pp. 74-97.
- 204. °GIROD, P. Manipulations de Zoologie. Guide pour les travaux pratiques de dissection. Animaux vertébrés. Paris : 8vo, vi & 158 pp., 32 pls.
- 205. GOGORZA Y GONZÁLES, J. Influencia del agua dulce en los animales mariuos. Ann Soc. Españ. Hist. Nat. xx, pp. 221-256 & 257-271.
- 206. GRABHAM, —. Heredity. J. Inst. Jamaica, i, No. iv, pp. 121-128.
- 207. Graffenberger, L. Versuche über die Veränderungen, welche der Abschluss des Lichtes in der chemischen Zusammensetzung des thierischen Organismus und dessen N-Umsatz hervorruft. Arch. ges. Phys. liii, pp. 238-280.
- 208. Grassi, B. Conclusioni d'una memoria sulla società dei Termiti. Atti [Rend.] Acc. Rom. 1892, pp. 33-36.
- 209. <sup>o</sup>Grawitz, P. Ueber die Umbildung von Grundsubstanz zu Zellen. Deuts, med. Wochschr. xviii (1892), pp. 712-714.
- Ueber die schlümmernden Zellen des Bindegewebes und ihr Verhalten bei progressiven Ernährungsstörungen. Arch. pathol. Anat. cxxvii, pp. 96-121.

- OGRAWITZ, P.] Ueber Schlummerzellen und Cellularpathologie.
   Deuts. med. Wochschr. xviii, pp. 811-813.
- 212. GRIFFITHS, A. B. The Physiology of the Invertebrata. London: 8vo, ix & 477 pp., 80 figs.
- On the Blood of the Invertebrata. P. R. Soc. Edinb. xix, pp. 116-130.
- 214. —. Sur la composition de l'hémocyanine. Bull. Ac. Belg. xxiii, pp. 842-844.
- 215. —... Recherches sur les couleurs de quelques insectes. ·C.R. cxv, pp. 958 & 959.
- 216. Groom, T. T. Early Development of Cirripedia. P. R. Soc. lii, pp. 158-162. [Oogenesis, Fertilisation.]
- 217. Guignard, L. Remarques sur la communication faite par M. Fayod dans la séance du 26 Decembre, 1891, sur la structure du protoplasme. C.R. Soc. Biol. iv, p. 1.
- 218. —... Remarques au sujet de la deuxième note de M. Fayod sur la structure du protoplasme. T. c. pp. 62-64.
- 219. 2—. Neue Studien über die Befruchtung. Vergleichung der bei Pflanzen und Thieren beobachteten morphologischen Erscheinungen. Nat. Rundschau. vii, pp. 282 & 283, figg.
- 220. GUINARD, L. Précis de tératologie. Anomalies et monstruosités chez l'homme et chez les animaux. Précédé d'une préface par Cam. Dareste. Paris: 8vo, xxiii & 553 pp., 272 figg.
- Guitel, F. Sur l'ovaire et l'œuf du Gobius minutus. C.R. cxiv, pp. 612-616.
- 222. Observations sur les mœurs du Gobius minutus. Arch. Z. espér. x, pp. 499-555, 1 pl. HAACKE, W. [See Brehm (73).]
- 223. HAASE, S. Untersuchungen über die Mimiery auf Grundlage eines natürlichen Systems der Papilioniden. Biblioth. Zool. (Leuckart & Chun). Cassel: 4to, Heft viii, Theil ii, pp. 1-98, 12 pls., 9 figg.
- 224. H.ECKEL, E. Plankton-Composition. Vorlaufige Mittheilung. Jen. Z. Nat. xxvii, pp. 559-566.
- 225. Häcker, V. Die Eibildung bei Cyclops und Canthocamptus. Zool, Jahrb. (Abth. Anat.) vi, pp. 211-248, 1 pl.
- 226. Die heterotypische Kerntheilung im Cyclus der generativen Zellen. Ber. Ges. Freiburg, vi, pp. 160-193, 3 pls.
- 227. Die Kernteilungsvorgänge bei der Mesoderm- und Entodermbildung von Cyclops. Arch. mikr. Anat. xxxix, pp. 556-581 2 pls.
- 228. . Ueber specifische Variation bei Arthropoden, im Besonderen über die Schutzanpassungen bei Krabben. Ber. Ges. Freiburg, vi, pp. 90-100.

- 229. HALLEX, P. Sur l'origine vraisemblablement tératologique de deux espèces de Triclades. C.R. cxiv, pp. 1125-1128; Rev. Sci. xlix, pp. 697 & 698.
- 230. OHALLIBURTON, W. D. Lehrbuch der chemischen Physiologie und Pathologie. Unter Mitwirkung des Verfassers deutsch bearbeitet von K. Kaiser. Heidelberg: 8vo, 12 & 883 pp., 100 figg.
- 231. OHAMANN, O. Entwicklungslehre und Darwinismus, Eine kritische Darstellung der modernen Entwicklungslehre und ihrer Erklärungsversuche mit besonderer Berücksichtigung der Stellung des Menschen in der Natur. Jena: 8vo, xix & 304 pp., 16 figg.
- 232. HANSEMANN, D. Ueber die Anaplasie der Geschwulstzellen und die asymmetrische Mitose. Arch. pathol. Anat. cxxix, pp. 436-449.
- 233. . Ueber Centrosomen und Attractionsphären in ruhenden Zellen. Anat. Anz. viii, pp. 57-59.
- 234. HARTOG, M. M. On Rabl's Doctrine of the Personality of the Segments of the Nucleus, and Weismann's "Idant" Theory of Heredity. Rep. Brit. Ass. 1892, pp. 742-744.
- 235. HATSCHEK, B. Zur Metamerie der Wirbeltiere. Anat. Anz. viii, pp. 89-91.
- 236. OHAUPTMANN, C. Beiträge zu einer dynamischen Theorie der Lebewesen. Bd. 1. Die Metaphysik in der modernen Physiologie. Dresden: 8vo, vii & 388 pp.
- 237. HAYCRAFT, J. B. Development of Vertebrate Muscle. Origin of Vertebrate Fibres. Further Growth. J. Physiol. xiii, pp. 16-18.
- 238. HEIDENHAIN, M. Ueber die Riesenzellen des Knochenmarkes und ihre Centralkörper. SB. Ges. Würzb. 1892, pp. 130-133.
- 239. —. Ueber Kern und Protoplasma. In Festschr.-Kælliker, pp. 109-166, 3 pls.
  - Heider, K. [See Korschelt & Heider (306).]
- 240. Heim, F. La zoologie et la médecine. Rev. Sci. l, pp. 737-743.
- 241. —. Sur la matière colorante bleue du sang des Crustacés. C.R. cxiv, pp. 771-774.
- 242. —. Sur les pigments des œufs des Crustaces. C.R. Soc. Biol. iv, pp. 467-470.
  - Re Colouration.
- 243. OHEINCKE, F. Variabilität und Bastardbildung bei Cyprinoiden. Festschr. Leuckart. pp. 65-73, 3 figg.
- 244. HELLER, K. M. Ein beachtenswerther Fall von Mimicry. Eut. Nachr. (Karsch) xviii, pp. 183-185.
  - HELMHOLZ, H. VON. [FESTSCHRIFT: see ENGELMANN and others (154).]

- 245. Henking, H. Untersuchungen über die ersten Entwicklungsvorgänge in den Eiern der Insecten. 111. Specielles und Allgemeines. Z. wiss. Zool. liv. pp. 1-274, 12 pls., 12 figg.
- 246. —... Künstliche Nachbildung von Kernteilungsfiguren. Arch. mikr. Anat. xli, pp. 28-39, 1 pl.
- 247. \*Henneguy, F. Classification des œufs des animaux basée sur leur constitution. C.R. Soc. Philom. 1891-92, pp. 1 & 2.
- 248. OHENNEGUY, L. F. Essai de classification des œufs des animaux au point de vue embryogénique. Bull. Soc. Philom. iv, pp. 37-44.
- 249. Herbst, C. Experimentelle Untersuchungen über den Einfluss der veränderten chemischen Zusammensetzung des umgebenden Mediums auf die Entwicklung der Thiere. I Theil. Versuche an Seeigeleiern. Z. wiss. Zool. lv, pp. 446-518, 2 pls.
- 250. HERDMAN, W. A. Rules of Nomenclature. Nature, xlvi, pp. 417 & 418.
- HERICOURT, J. La médication de M. Brown-Séquard. Rev. Sci. l, pp. 817-821.
- 252. °HERRICK, C. L. Metamerism of the Vertebrate Head. J. Comp. Neurol. i (1891), pp. 203 & 204.
- 253. °----. Histogenesis and Physiology of the Nervous Elements. Op. cit. ii, pp. 137-149.
  - ---. [See also Brooks & Herrick (78).]
- 254. HERTWIG, O. Text-Book of the Embryology of Man and Mammals. Trans. from 3rd German Ed. by E. L. Mark. London: 8vo, 660 pp., 339 figg., 2 col. pls.
- 255. —... Urmund und Spina bifida, eine vergleichend-morphologische, teratologische Studie an missgebildeten Froscheiern. Arch. mikr. Anat. xxxvii, pp. 353-503, 3 pls.
- 256. —. Die Zelle und die Gewebe. Grundzüge der allgemeinen Anatomie et Physiologie. Erster Theil. Jena: 8vo, xi & 296 pp., 168 figg.
- 257. \*---. Altere und neuere Entwickelungs-Theorien. Berlin: 8vo, 35 pp.
- 258. HERTWIG, R. Lehrbuch der Zoologie. Teil ii. Jena: 8vo, pp. vii & 321-588, figg. 315-568.
- 259. —. Ueber Befruchtung und Conjugation. Verh. deutsch. Zool. Ges. ii, pp. 95-112.
- 260. His, W. Zur allgemeinen Morphologie des Gehirns. Arch. Anat. Phys. (Anat. Abth.) 1892, pp. 346-383, 36 figg.
- Die Entwickelung der menschlichen und tierischen Physiognomien. T. c. pp. 384-424, 51 figg.

- 262. [His, W.] Zur Nomenclatur des Gehirns und Rückeumarks. Arch. Anat. Phys. (Anat. Abth ) 1892, pp. 425-428.
- 263. Hodgkinson, A. Iridescent Colours. Nature, xlvi, pp. 92-94.
- 264. Holborow, A. Evolution and Scripture. London: 8vo, 338 pp.
- 265. HOLT, E. W. L. North Sea Investigations. J. Mar. Biol. Ass. ii (1892), pp. 216-219, 363-393, & 399-404.
  - Re Reproduction of Fishes.
- 266. HOPKINS, F. G. [Letter to Nature.] Pigments of Lepidoptera. Nature, xlv, p. 581.
  - Cf. Coste, t. c. p. 605, and Meldola, ibidem.
- 267. HORNELL, J. [Letter to Nature.] A Strange Commensalism— Sponge and Annelid. Op. cit. xlvii, p. 78.
- 268. Horsley, V. The Structure and Functions of the Brain and Spinal Cord. Being the Fullerian Lectures for 1891. London and Philadelphia: 8vo, 229 pp., illustr.
- 269. HORVATH, —. Sur l'existence des series parallèles dans le cycle biologique des Pemphigiens. C.R. cxiv, pp. 842-844.
- 270. HOUSSAY, F. Sur la théorie des feuillets et le parablaste. T. c. pp. 1128-1130.
- 271. HUDSON, W. H. The Naturalist in La Plata. London: 8vo, 383 pp., illustr.
- 272. Hudson, G. V. Instances of Instinct in Insects. Tr. N. Z. Inst. xxiv, pp. 354-358.
- 273. Hurst, C. H. Biological Theories. I. The Nature of Heredity. Nat. Sci. i, pp. 502-507 & 578-587.
  - —. [See also Marshall, A. M., & Hurst (341).]
- 274. HUTCHINSON, H. N. Extinct Monsters: a Popular Account of some of the Larger Forms of Ancient Animal Life. London: 8vo, 234 pp., numerous illustr.
- 275. HUNLEY, T. H. Essays upon some Controverted Questions. London: 8vo, 616 pp.
- 276. H. W. [Letter to Nature.] On the Attitudes of the Zebra during sleep, and their Influence on the Protective Value of its Stripes. Nature, xlv, p. 248.
- 277. IMHOF, O. E. Programm zu einer monographischen Bearbeitung eines grösseren Sees, enthaltend die verschiedenen Gesichtspunkte, deren eingehendes Studium zur Erkenntnis der Existenzbedingungen des gesammten organischen Lebens der Seen nothwendig ist. Biol. Centralbl. xii, pp. 512-521.
- Die Zusammensetzung der pelagischen Fauna der Süsswasserbecken. Biol. Centralbl. xii, pp. 171-182 & 200-205.

- 279. [IMHOF, O. E.] Vorläufige notiz über die Lebensverhältnisse und Existenz-bedingungen der pelagischen und Tiefsee- Flora und Fauna der Seen. T. c. pp. 521-523.
- 280. —... Les conditions de milieu dans les lacs d'eau douce. Rev. Sci. l, pp. 497-500.
- 281. —... Considérations générales sur la faune des Invertébrés de la Suisse. C.R. Soc. helv. Sci. Nat. lxxiv, pp. 56-66.
- 282. °—. Ueber das Leben und die Lebensverhältnisse zugefrorener Seen. MT. Aargau. nat. Ges. 1892, pp. 42-58. Cf. Beiträge zur Fauna der Schweiz. Thierwelt der Stehenden Gewässer. T. c. pp. 59-110.
- 283. ISCHIKAWA, C. Studies on Reproductive Elements. I. Spermatogenesis, Ovogenesis, and Fertilisation in *Diaptomus*. J. Coll. Sci. Japan, v. pp. 1-34, 1 pl.
- 284. JACQUART, E. Ce qu'il faut penser de l'intelligence des animaux. Lyon: 8vo, 62 pp.
- 285. OJARISCH, —. Ueber die Bildung des Pigmentes in den Oberhautzellen. Arch. Dermatol. xxiv, pp. 223 & 224, 1 pl.

  Re Colouration.
- 286. Jensen, P. Ueber den Geotropismus niederer Organismen. Arch. ges. Physiol. liii, pp. 428-480, 13 figg.
- 287. Methode der Beobachtung und Vivisection von Infusorien in Gelatinelösung. Biol. Centralbl. xii, pp. 556-560.
- 288. Johns, B. G. The Protective Colour in Animals. Nineteenth Century, xxxii, pp. 454-469.
- 289. Johnson, H. P. Amitosis in the Embryonal Envelopes of the Scorpion. Bull. Mus. C. Z. xxii, pp. 127-161, 3 pls.
- 290. Joseph, G. L'influence de l'éclairage sur la disjunction des organes visuels, leur réduction, leur atrophie complète, et leur compensation chez les animaux cavernicoles. Bull. Soc. Z. Fr. xvii, pp. 121-125.
- 291. CKELLER, C. Alpentiere im Wechsel der Zeit. (Marshall's Zool. Vorträge.) Leipzig: 8vo, 48 pp.
- 292. Kennel, J. von. Mimicry bei kleinen Insecten. Ent. Nachr. xviii, pp. 159 & 160.
- 293. —... Questions proposées par M. le Prof. Cosmovici: 1. Sur une division définitive du règne animal en "phyla," division basée sur les recherches morpho-embryologiques. 2. Fondement naturel d'un système du type des vers. 3. Uniformité de la terminologie des organes sécrétoires des vers. Congrès Internat. Zool. ii, pt. 1, pp. 68-73.
- 294. KIPLING, J. L. Beast and Man in India. New Edition. London: 8vo, 356 pp., illustr.

- KNAUTHE, K. Ueber vererbte Verstümmelungen. Zool. Anz. xv,
   p. 5.
- 296. —. Ueber Melanismus bei Fischen. T. c. p. 25.
- 297. Kochs, W. Ueber die Vorgänge beim Einfrieren und Austrocknen von Tieren und Pflanzensamen. Biol. Centralbl. xii, pp. 330-339.
- 298. CKOCKS, J. Ueber die Korrelation des Wachstums der rudimentären organe und ihres Mutterbodens. Verh. Deuts. Ges. Gynäkol, iv, Kongress, pp. 425-432.
- 299. Kenig-Warthausen, Friir. R. Uober die Färbung der Vogeleier und über späte Vogelbruten. JH. Ver. Württ. xlviii, pp. 63-65.
- 300. \*Kohncke, H. Thier und Mensch. Plaudereien aus dem Zoologischen Garten. Part 1,66 pp. Hamburg: 8vo.
- 301. Kölliker, A. von. Nervenzellen und Nervenfasern. Biol. Centralbl. xii, pp. 33-51.
- 302. —. Ueber die Entwickelung der Elemente des Nervensystems, contra Beard und Dohrn. Verh. Anat. Ges. vi (1892), pp. 76-78. —. [Festschrift; see SCHULTZE and others (458).]
- 303. °Köppen, M. Beiträge zur vergleichenden Anatomie des Centralnervensystems der Wirbelthiere. Zur Anatomie des Eidechsengehirns. Morph. Arb. (Schwalbe), i, pp. 496-515, 3 pls.
- 304. Kohl, C. Rudimentäre Wirbeltieraugen. Teil 1. Bibl. Zool. Heft 13, 143 pp., 9 pls.
- 305. °Korschelt, E. Entwicklungsmechanische Studien. Nat. Rundschau, vii, pp. 11, 12, & 89.
- 306. Korschelt, E., & Heider, K. Lehrbuch der vergleichenden Entwicklungsgeschichte der wirbellosen Thiere. Part II. Jena: 8vo, pp. 309-908, 315 figg.
- 307. °KOSTANECKI, K. VON. Ueber die Schicksale der Centralspindel bei karyokinetischer Zellteilung. Anat. Hefte ii, pp. 249-268, 36 figg.
- 308. °KOSTANECKI, V. VON. Ueber Kernteilung bei Riesenzellen nach Beobachtungen an der embryonalen Säugetierleber. Anat. Hefte i, pp. 323-352, 20 figg.
- 309. °KRÄPELIN, K. Die Brutpflege der Thiere (Sammlung Vortr. Virchow und Wattenbach). Hamburg: 8vo, 26 pp.
- 310. Krasser, F. Zur Structur des ruhenden Zellkerns. SB. Ak. Wien, ci, pp. 560-583.
- 311. KROMAYER, E. Die Protoplasmafaserung der Epithelzelle. Arch. mikr. Anat. xxxix (1892), pp. 141-150, 1 pl.
- 312. KRUMMEL, O. Reisebeschreibung der Plankton-Expedition nebst Einleitung von V. Hensen, und Vorberichten von Dahl, Apstein, Lohmann, Borgert, Schütt, und Brandt. Mit 100 figg. in Text, sowie 5 Karten, u. 1 Photogravure, Kiel u. Leipzig: 4to, 370 pp.

- 313. Kunckel d'Herculais. Le Criquet pélerin (Schistocerca peregrina, Oliv.), et ses changements de coloration. Rôle des pigments dans les phénomènes d'histolyse et d'histogènese qui accompagnent la métamorphose. C.R. cxiv, pp. 240-242.
  - Re Colouration.
- 314. \*Kupffer, C. von. The Development of the Cranial Nerves of Vertebrates. J. Comp. Neurol. i (1891), pp. 246-264.
- 315. CLABORDE, J. V. Traité élementaire de Physiologie. Paris: 8vo, 387 pp., 25 pls., 130 figg.
- 316. CLAMEERE, A. Esquisse de la Zoologie. Bruxelles: 12mo, 302 pp. LANDI, L. [See GAUTIER & LANDI (188-191).]
- 317. OLANDOIS, H. Die Ernährung der Larven und Embryonen ist bestimmend für die Eustehung der Geschlechter. Nat. Wochenschr. vii, p. 170.
- 318. LANG, A. Lehrbuch der vergleichenden Anatomie zum Gebrauche bei vergleichend anatomischen und zoologischen Vorlesungen. 3 Abth. Jena: 8vo, pp. 567-870, 219 figg.

Translation of previous parts into English (H. M. & M. Bernard), and into French (G. Curtel).

- 319. Die Asymmetrie der Gastropoden. Viert. Ges. Zurich, xxxvi (1891), pp. 339-371, 22 figg.
- 320. °LATASTE, F. Indications nouvelles et réponse à une réclamation de priorité, à propos de sa publication : Pourquoi, dans le même type de Vertébrés la masse relative de l'encéphale varie en sens inverse de la masse du corps. Act. Soc. Sci. Chili, i, pp. 35-38.
- 321. LA VALETTE, St. GEORGE. Ueber innere Zwitterbildung beim Flusskrebs. Arch. mikr. Anat. xxxix, pp. 504-524, 1 pl.
- 322. Lea, A. Sheridan. The Chemical Basis of the Animal Body: an Appendix to Foster's Text-book of Physiology. London: 8vo, 266 pp.
- 323. LEBRUN, H. Les centrosomes dans l'œuf de l'Ascaris megalocephala. Anat. Anz. vii, pp. 627 & 628.
- 324. LEUCKART, R. Festschrift zum siebenzigsten Geburtstage Rudolf Leuckarts. Dem verehrten Jubilar dargebracht von seinen dankbaren Schülern. Leipzig: 4to, xv & 413 pp., 40 pls., 43 figg., and portrait.
- 325. LILIENVELD, L., & MONTI, ACHILLE. Sulla localizzazione microchimica del fosforo nei tessuti. Atti Acc. Rom. 1892, pp. 354-358.
- 326. LOCARD, A. L'influence du milieu sur la variation. Rev. Sci. xlix, pp. 724-726.

- 327. Looss, A. Schmarotzerthum in der Thierwelt. (Marshall's Vorträge.) Leipzig: 8vo, 180 pp.
  - LORIA, A. [See Morselli (366).]
- 328. LOTHES, R. Præparirmethodik. Eine Anleitung zu den anatomischen Uebungen für die Studierenden der Thiermedicin. Berlin: 8vo, vi & 135 pp., 8 pls.
- 329. °LWOFF, W. Gegenwärtiger Stand der Frage von der Polyspermie (Russian). Rev. Sci. Nat. St. Petersb. iii, pp. 23-30.
- 330. LYDEKKER, R. Phases of Animal Life: Past and Present. London: 8vo, 234 pp.
- 332. McKendrick, J. G. Life in Motion; or Muscle and Nerve. London and Edinburgh; 8vo, xiii & 202 pp., 81 figg.
- 333. McMurricii, J. P. Definite versus Fortuitous Variation. [Discussion.] Am. Nat. xxvi, pp. 85-91.
- 334. \*Macpherson, A. H. Albinism in Birds and Mammals. Zool. xvi, p. 191.
- 335. Mann, G. The Functions, Staining Reactions, and Structure of Nuclei. Rep. Brit. Ass. 1892, pp. 753 & 754.
- 336. —. The Origin of Sex. T. c. pp. 756 & 757.
- 337. MARCHAL, P. Notes sur La Vie et les Mœurs des Insectes: observations sur l'Ammophila affinis, Kirby. Arch. Z. expér. x, pp. 23-36.
  Re Instinct.
- 338. °MARENZELLER, E. VON. Das Heim der Tiefseethiere. Schr. Ver. Verbr. nat. Kenntn. Wien, Bd. xxxii, pp. 149-177.
- 339. MAREY, J. La Photographie du mouvement. Avec chromophotographies originales reproduites en photogravures sans intervention de la main humaine. Paris: 8vo, 90 pp.

  MARK, E. L. [See HERTWIG, O. (254).]
- 340. MARSHALL, A. MILNES. Descriptive Catalogue of the Embryological Models, Manchester Museum. Manchester: 8vo, 160 pp.
- 341. Marshall, A. M., & Hurst, C. H. A Junior Course of Practical Zoology, 3rd Edition. London: 8vo, 502 pp., figg.
- 342. MARSHALL, W. Das Thema der Flughautbildungen von einem allgemeinen Gesichtspunkte. SB. Ges. Leipzig, xvii-xviii Jhg. pp. 11-13.
- 343. °MATTHIESEN, L. Die neueren Fortschritte in unserer Kenntniss von dem optischen Baue des Auges der Wirbelthiere. Hamburg aud Leipzig: 8vo, 63 pp., 2 pls.
  - From Festschrift Helmholtz.

- MATTIEU-WILLIAMS, W. [See ROMANES (435).]
- 344. MEEHAN, T. Definite versus Fortuitous Variation. [Discussion.] Am. Nat. xxvi, pp. 85-91.
  - MELDOLA, R. [See HOPKINS (266).]
- 345. CMELTZER, S. J. On the Importance of Vibration to Cell Life. New York Med. J. v. pp. 708-712.
- 346. MERRIFIELD, F. Double-broodedness: whether influenced by the state of the food-plant? Ent. M. M. xxviii, pp. 20 & 21.
- 347. METSCHNIKOFF, E. Lécons sur la Pathologie comparée de l'inflammation faites à l'Institut Pasteur en Avril et Mai, 1891. Paris : 8vo, 1892, 235 pp., 65 figg., 13 col. pls.
- 348. —... La Lutte pour l'existence entre les diverses parties de l'organisme. Rev. Sci. l, pp. 321-326.
- 349. MIALI, L. C. The Surface-Film of Water, and its relation to the life of Plants and Animals. Nature, xlvi, pp. 7-11.
- 350. MICHAEL, A. D. On the Variations in the Internal Anatomy of the Gamasinæ, especially in that of the genital organs, and on their mode of coition. Tr. L. S. v, pp. 281-324, 4 pls.
- 351. MILES, MANLY. Heredity of Acquired Characters. P. Am. Ass. xli, pp. 202-211.
- 352. ——. Heredity of Acquired Characters. Am. Nat. xxvi, pp. 887-900.
- 353. MINERVINI, R. Contributo alla Morfologia dell' adattamento funzionale degli organi: particolarità di struttura delle arterie della cute. Boll. Soc. Nat. Napoli, vi, pp. 138-152, 1 pl.
- 354. MINGAZZINI, P. L'Oolisi nella Seps chalcides. Atti [Rend.] Acc. Rom. 1892, pp. 41–45. Cf. Mon. Zool. iii, pp. 8 & 9.
- 355. Minot, C. S. Human Embryology. New York: 8vo, xxiii & 815 pp., 463 figg.
  Re Generalities.
- 356. MITCHELL, P. C. Death. Nat. Sci. i, pp. 434-441.

  MIVART, St. GEORGE. [See DIXON (127).]
- 357. °Möbius, K. Die Thiergebiete der Erde, ihre kartographische Abgrenzung und museologische Bezeichnung. Nat. Wochenschr. vii, pp. 107 & 108.
  - Monti, A. [See Lilienfeld & Monti (325).]
- 358. MONTICELLI, F. S. Sul nucleo vitellino del uova dei Trematodi. Boll. Soc. Nat. Napoli, vi, pp. 5-8.
- 359. C. Richerche sulla spermatogenesi nei Trematodi. Month. Int. J. Anat. Hist. ix, pp. 112-118 & 121-149, 2 pls.
  - 1892. [vol. xxix.]

- 360. MOORE, J. E. S. On the Relationships and Rôle of Archoplasm during Mitosis in the Larval Salamander. Q. J. Micr. Sci. xxxiv, 1892, pp. 181-197, 1 pl.
- On the Relationships and Rôle of the Archoplasmic Body during Mitosis in the Larval Salamander. Rep. Brit. Ass. 1892, pp. 755 & 756.
- 362. MORGAN, C. LLOYD. The Method of Comparative Psychology. T. c. pp. 754 & 755.
- 363. —. Factors in the Evolution of Mammals. Nat. Sci. i, pp. 97-101.
  - —. [See also Dixon (127).]
- 364. Morgan, T. H. Spiral Modification of Metamerism. J. Morph. vii, pp. 245-251, 3 figg.
- 365. Morris, C. The Origin of Lungs, a chapter in Evolution. Am. Nat. xxvi, pp. 975-986.
- 366. MORSELLI, E. Carlo Darwin e il Darwinismo nelle Scienze biologiche e sociali. Scritti vari di G. Canestrini, G. Cattaneo, A. Loria, &c. Milano: 8vo, 312 pp.
- 367. Moseley, H. N. Notes by a Naturalist: an Account of Observations made during the Voyage of H.M.S. "Challenger." New and revised edition, with map, portrait, and woodcuts, and a brief memoir of the author. London: 8vo, 562 pp.
- 368. <sup>o</sup>Motto, Coco Λ. Leggendo Darwin: contributo-psicologico. Napoli: 8vo, 16 pp.
- 369. OMULLER, J. Ueber Gamophagie. Ein Versuch zum weiteren Ausbau der Theorie der Befruchtung und Vererbung. Stuttgart: 8vo, 64 pp.
- 370. °NADAILLAC, LE MARQUIS DE. Intelligence et Instinct. Paris: 8vo, 79 pp.
- 371. Nagel, W. Beobachtungen über das Verhalten einiger wirbelloser Thiere gegen galvanische und faradische Reizung. Arch. ges. Phys. lii, pp. 624-631.
- 372. Fortgesetze Beobachtungen über polare galvanische Reizung bei Wasserthieren. Op. cit. liii, pp. 332-347.
- 373. Nathusius, W. von. Die fibrilläre Structur der Hornzellen der Haare. Zool. Anz. xv, pp. 395-400, 9 figg.
- 374. "NEUMANN, L. G. Treatise on the Parasites and Parasitic Diseases of the Domestic Animals. Trans. and ed. by G. Fleming. London: 8vo, 812 pp., 365 figg.
- 375. NEVIANI, A. Riproduzione animale e vegetale. Rev. Ital. Sci. Nat. xii, pp. 86-88, 101, 102, 105-109, 134-136, & 137-141.

- 376. NICOLAS, A. Les sphères attractives et le fuseau achromatique dans le testicule adulte, dans la glande génitale et le rein embryonnaires de la Salamandre. C.R. Soc. Biol. iv (1892), pp. 472-477.
- Les spermatogonies chez la Salamandre d'hiver (noyaux polymorphes; sphère attractive, division directe). C.R. Soc. Biol. iv, pp. 590-595.
- 378. ONORMAN, A. M. Museum Normanicum, or a Catalogue of the Invertebrata of Europe, and the Arctic and North Atlantic Oceans, which are contained in the Collection of the Rev. Canon A. M. Norman. VII. Spongozoa. VIII. Rhizopoda. Printed for private distribution. Durham: 8vo, 21 pp.
- 379. Nussbaum, M. Beiträge zur Lehre von der Fortpflanzung und Vererbung. Arch. mikr. Anat. xli, pp. 129-145.
- 380. NUTTING, C. C. What is an "Acquired character"? Am. Nat. xxvi, pp. 1009-1013.
- 381. OKA, A. Die periodische Regeneration der oberen Körperhälfte bei den Diplosomiden. Biol. Centralbl. xii, pp. 265-268.
- 382. OLIVIER, L. On the Canalisation of Cells, and the Continuity of Living Matter in Plants and Animals. Rep. Brit. Ass. 1892, pp. 792 & 793.
- 383. Oppel, A. Die Befruchtung des Reptilieneies. Arch. mikr. Anat. xxxix, pp. 215-290, 4 pls.
- 384. —.. Unsere Kenntniss von der Enstehung der roten und weissen Blutkörperchen. Zusammenfassendes Referat. Centralbl. Pathol. iii, pp. 193-217 & 241-259.

See for notice of various papers on this subject, not included in this record.

- 385. Osborn, H. F. The Difficulties in Hereditary Theory. Am. Nat. xxvi, pp. 537-567.
- 386. —. Heredity and the Germ Cells. T. c. pp. 642-670.
- 387. OSBORN, H. L. The Protozoa—a Phylum of the Animal Kingdom considered Biologically. Am. Micr. J. xiii, pp. 233-243, 1 pl.
- 388. —. Introduction to Elementary Biology. Op. cit. xii, pp. 241-243; xiii, pp. 9-11 & 26-28.
- 389. Passy, J. La photographie du mouvement. Rev. Sci. xlix, pp. 17-20.
- 390. PEARSON, K. The Grammar of Science. London: 8vo, pp. PECHUEL-LOESCHE. [See BREHM (73).]
- 391. Perényi, J. Die Enstehung des Mesoderms. Math. nat. Ber. Ung. viii, pp. 272-278, 2 pls.
- 392. Perrier, Remy. Éléments d'anatomie comparée. Paris : 8vo, 1200 pp., 600 figg., 8 col. pls.

- 393. \*Perrier, E. Traité de Zoologie. 1 Partie. Zoologie générale, Protozoaires et Phytozoaires. Paris : 8vo, vii & 865 pp., 700 figg.
- 394. Petersen, W. Ueber die Ungleichzeitigkeit in der Erscheinung der Geschlechter bei Schmetterlingen. Zool. Jahrb. (Abth. Syst. Geog. Biol.), vi, pp. 671-679.
- 395. Philippon, G. Effets de la décompression brusque sur les animaux placés dans l'air comprimé. C.R. cxv, pp. 186-188.
- 396. PICTET, C. Spermatogénèse et Fécondation chez les Animaux Invertébrés. Arch. Phys. Nat. xxvii, pp. 175-214 [cf. same in MT. z. Stat. Neap. x (1891), pp. 75-152, 2 pls.].
- 397. PLATEAU, F. La ressemblance protectrice dans le règne animal. Bull. Ac. Belg. xxiii, pp. 89-135.
- 398. Pœhl, A. Action physiologique de la spermiue. Interprétation de ses effets sur l'organisme. C.R. cxv, pp. 129-132.
- 399. —... Du rôle de la spermine dans les oxydations intra-organiques. T. c. pp. 518-521.
- 400. °Posner, C. Weitere Notiz zur Chemie des Samens. CB. med. Wiss. 1892, pp. 225 & 226.
- POUCHET, G. Sur la formation du pigment mélanique. C.R. Soc. Biol. iv, p. 516,
- 402. POULTON, E. B. Further Experiments upon the Colour-relation between certain Lepidopterous larvæ, pupæ, cocoons, and imagines, and their surroundings. Tr. E. Soc. 1892, pp. 293-487, pls. xiv & xv. ——. [See also Bateson (34).]
- 403. PRENANT, A. Sur la signification de la cellule accessoire du testicule et sur la comparaison morphologique des éléments du testicule et de l'ovaire. J. de l'Anat. Phys. xxviii, pp. 292-321 & 529-562.
- 404. —... Contribution à l'étude de la division cellulaire. Le corps intermédiare de Flemming dans les cellules séminales de la Scolopendre et de la Lithobie. Arch. Phys. xxiv, pp. 295-310, 1 pl.
- 405. —. Le corps intermédiare de Flemming dans les cellules séminales de la Scolopendre et de la Lithobie. C.R. Soc. Biol. iv, pp. 172-176.
- 406. —. Le corpuscle central d'E. van Beneden dans les cellules séminales de la Scolopendre. T. c. pp. 221-225.
- L'origine du fuseau achromatique nucléaire dans les cellules séminales de la Scolopendre. T. c. pp. 219-253.
- 409. QUATREFAGES, A. DE. Darwin et ses précurseurs français. Étude sur le transformisme. 2 Ed. rev. et augm. Paris : 8vo, 299 pp.

- 410. Rabl, C. Theorie des Mesoderm (Fortsetzung). Morph. JB. xix, pp. 65-144, 4 pls., 3 figg.
- 411. —... Ueber die Metamerie des Wirbeltierkopfes. Verh. Anat. Ges. vi (1892), pp. 104-135, 1 pl., 4 figg.
- 412. °RAILLET, A. Parasites animaux. Les parasites transmissibles des animaux à l'homme envisagés spécialement au point de vue de la prophylaxie. Paris : 8vo, 48 pp.
- 413. RANDOLPH, H. The Regeneration of the Tail in Lumbriculus. J. Morph. vii, pp. 317-344, 4 pls.
  - General considerations as to regeneration.
- 414. •RANVIER, L. Le système vasculaire. J. Microgr. xv, pp. 295-303, and xvi, pp. 135-142.
- 415. Les éléments et les tissus du système conjonctif (Suite). Op. cit. xv, pp. 321-326.
- 416. RASPAIL, X. Description d'une série de pontes d'Oiseaux au point de vue de la coloration et de la forme des œufs. Mém. Soc. Zool. v, pp. 174-189.
- 417. —... À propos de l'origine de la couleur des œufs des Oiseaux. Bull. Soc. Z. Fr. xvii, pp. 212-218.
- 418. RATH, O. VOM. Kritik einiger Fälle von scheinbarer Vererbung von Verletzungen. Ber. Ges. Freiburg, vi, pp. 101-113.
- 419. —. Zur Kenntnis der Spermatogenese von Gryllotalpa vulgaris, Latr., mit besonderer Berücksichtigung der Frage der Reduktionsteilung. Arch. mikr. Anat. xl, pp. 102-132, 2 pls.
- 420. °RAWITZ, B. Compendium der vergleichenden Anatomic. Leipzig: 8vo, v & 272 pp., 90 figg.
- Beichenow, A. Handwörterbuch der Zoologie, Anthropologie, und Ethnologie. Bd. vi. Breslau: 8vo, 571 pp., figg.
- 422. \*Replachoff, W. Ueber die Gastrulation der Wirbeltiere nebst Bemerkungen über die Homologie der Keimblätter der Metazoen. [Russian.] Mém. Soc. Nat. Nouv. Russ. xvii, 24 pp.
- 423. °RETTERER, E. Anatomie et Physiologie animales. Paris: 16mo, 590 pp., illustr.
- 424. Retzius, G. Biologische Untersuchungen. Neue Folge. Stockholm and Leipzig: 4to, ii, 53 pp., 16 pls.; iii, 68 pp., 23 pls.; iv, 70 pp., 29 pls.
- 425. RILEY, C. V. Fortuitous Variation. Abstract. Am. Nat. xxvi, pp. 861-863.
- 426. ——. The transmission of Acquired Characters through Heredity.

  Abstract. Nature, xlvi, pp. 504 & 505.
- 427. RITZEMA-Bos, J. Zoologie für Landwirthe. Berlin: 8vo, viii & 182 pp., 149 figg.

- 428. Robinson, A. The Nutritive Importance of the Yolk-sac. J. Anat. Phys. xxvi, pp. 308-323, 1 pl.
- 429. ROEMER, F. Die chemische Reizbarkeit tierischer Zellen. Arch. Pathol. cxxviii, pp. 98-131, 1 pl.
- ROHDE, E. Muskel und Nerv. 1. Ascaris. 11. Mermis und Amphioxus. 111. Gordius. Zool. Beitr. iii, pp. 69-106, 6 pls.; pp. 161-182, 4 pls.; pp. 183-192.

General Histology.

431. —... Muskel und Nerve bei Nematoden. SB. Ak. Berlin, 1892, pp. 515-526.

General Histology.

- 432. Muskel und Nerve bei Mermis und Amphioxus. SB. Ak. Berlin, 1892, pp. 659-664.
- 433. ROMANES, G. J. Darwin, and after Darwin: an Exposition of the Darwinian Theory, and a Discussion of Post-Darwinian Questions. Vol. 1. The Darwinian Theory. London: 8vo, 430 pp., illustrated.
- 434. Darwin und nach Darwin. Eine Darstellung der Darwin'schen Theorie, und Erörterung Darwinistischer Streitfragen. Bd. 1. Die Darwin'sche Theorie aus d. Engl. von B. Vetter. Leipzig: 8vo, vii & 542 pp., 1 portrait, 124 figg.
- 435. . [Letter to Nature.] The Migration of the Lemming. Nature, xlv, p. 248. Cf. Crotch, t. c. p. 294, and Mattieu-Williams, t. c. p. 295.
- ROULE, L. L'Embryologie générale. Paris: 16mo, xiv & 510 pp.,
   121 figs.
- Les affinités zoologiques des Vertébrés. Rev. Sci. xlix, pp. 588-594.
- 438. Roux, W. Ueber das entwickelungsmechanische Vermögen jeder der beiden ersten Furchungszellen des Eies. Verh. Anat. Ges. vi, 1892, pp. 22-62.
- 439. Beiträg zur Entwicklungsmechanik des Embryo. Ueber die morphologische Polarisation von Eiern und Embryonen durch den elektrischen Strom, sowie über die Wirkung des elektrischen Strom auf die Richtung des ersten Theilung des Eies. SB. Ak. Wien. ci, pp. 27-234, 3 pls; Ber. Nat. Med. Ver. Innsbruck, xx, pp. 3-57.
- 440. RÜCKERT, J. Zur Entwickelungsgeschichte des Ovarialeies bei Selachiern. Anat. Anz. vii, pp. 107-158, 6 figs.
- Ueber physiologische Polyspermie bei meroblastischer Wirbeltiereiern. Verh. Anat. Ges. vii, pp. 320-333, 2 figs.
- 442. . Ueber die Verdoppelung der Chromosomen im Keimbläschen des Selachiereies. Anat. Anz. viii, pp. 44-52, 2 figg.

- 443. RYDER, J. A. On the mechanical genesis of the Scales of Fishes. P. Ac. Philad. 1892, pp. 219-224.
- 444. The Principle of the Conservation of Energy in Biological Evolution: a Reclamation and Critique. T. c. pp. 455-468.
- 445. A Geometrical Representation of the relative intensity of the Conflict between Organisms. Am. Nat. xxvi, pp. 923-929.
- 446. SABATIER, A. Essai sur la Vie et la Mort. Paris: 8vo, 277 pp.
- 447. La synthèse des êtres vivants. Rev. Sci. l, pp. 16-20.
- 448. SAINT-REMY, G. Sur l'histologie de la glande pituitaire. C.R. cxiv, pp. 770 & 771.
- Scervini, P. Anatomia dei centri nervosi. Napoli: xii & 356 pp., figg.
- 450. SCHAAFFHAUSEN, —. Ueber die Urzeugung. CB. Ver. Rheinl. xlix, pp. 32-40.
- 451. SCHILLEE-TIETZ, —. Folgen, Bedeutung und Wesen der Blutverwandtschaft (Inzucht) im Menschen-, Thier-, und Pflanzenleben.

  2te verm. u. völlig umgearb. Aufl. Berlin u. Neuwied: 8vo, 94 pp.
- 452. SCHIRSCHOFF, D. J. Ueber die Beziehung der Körperform der Fische zur Stromstärke des von ihnen bewohnten Bassins. Trudui St. Petersburg Est. xxiii, pp. 79-99.
- 453. SCHMIDT, H. Schlummernde Zellen im normalen und pathologischveränderten Fettgewebe. Arch. pathol. cxxviii, pp. 58-97, 1 pl.
- 454. SCHMUL, A. Ueber das Schicksal des Eisens im thierischen Organismus. Inaug. Diss. Dorpat: 8vo, 28 pp.
- 455. Schneider, K. C. Zur Frage der Entwicklung des Intellekts. Biol. Centralbl. xii, pp. 30-32.
- 456. SCHNEIDER, R. Verbreitung und Bedeutung des Eiseus im animalischen Körper. Verh. Deutsche Naturf. 64 vers., pp. 111-116.
- 456A. Schofield, A. T. Elementary Physiology for Students. London: 8vo, xii & 372 pp., 227 figg., 2 col. pls.
- 457. SCHOTTLÄNDER, —. Ueber die Enstehung des Graaf'schen Follikels bein Menschen und seinen Untergang bei Mensch und Säugetieren. Z. Geburtshilfe u. Gynäkol. xxiv, pp. 312-314.
- 458. SCHULTZB, O. [and others]. Festschrift Herrn Geheimrath Albert von Kælliker zur Feier seines fünfzigjährigen medicinischen Doctorjubiläums gewidmet vom dem anatomischen Institut der Universität Würzburg. (Schultze, O.; Fick, R.; Voll, A.; Sclavunos, G.; Heidenhain, M.) Leipzig: fol., 166 pp., 11 pls., 2 figg.
- 459. CSCHULZ, P. Ueber die in historischer Zeit ausgestorbenen Thiere. Berlin: 4to, 35 pp., 5 figg.
- 460. Schulze, F. E. Ueber die Bezeichnung der Lage und Richtung im Thierkörper. SB. nat. Fr. 1892, pp. 43-51.

- 461. Schütt, F. Analytische Plankton Studien. Ziele, Methoden, uud Anfangsresultate der quantitative-analytischen Planktonforschung. Kiel and Leipzig: 8vo, 117 pp., 16 tables, 1 map.
- 462. OSCHWEIGER-LERCHENFELD, A. VON. Das Mikroskop. Leitfaden des mikroskopischen Technik nach dem heutigen Stande der theoretischen und praktischen Erfahrungen. Wien & Leipzig: 8vo, 144 pp., 91 figg., 13 pls.
- 463. °SEITZ, AD. Ueber gewisse Eigenthümlichkeiten bei Fortpflanzung einiger Thiere. Ber. Oberhess. Ges. xxviii, pp. 118 & 119.
- 464. SEMON, R. Studien über den Bauplan des Urogenitalsystems der Wirbeltiere. Dargelegt an der Entwickelung dieses Organsystems bei Ichthyophis glutinosus. Jen. Z. Nat. xxvi, pp. 89-203, 14 pls.
- 465. SEMPER, C. Ueber Farbenveräuderung einiger Fische. Arb. Inst. Würzb. x, pp. 15-19, 1 pl.
- 466. SHALER, N. S. Nature and Man in North America. London: 8vo, 290 pp.
- 467. SICARD, H. L'évolution sexuelle dans l'Espèce humaine, Paris : 8vo, 319 pp., 94 figg.
- 468. SIEGFRIED, M. Ueber die chemischen Eigenschaften des reticulierten Gewebes. Ber. Sächs. Ges. 1892, p. 306.
- 469. \*-----. Ueber die chemischen Eigenschaften des reticulierten Gewebes. Habilit. Schrift. Leipzig: 8vo, 21 pp.
- SIMROTH, H. Einige Punkte aus der Oekonomie des Weichthierkörpers, ein Kapitel über Constitution. Leopoldina, xxviii, pp. 100– 102, 121–125, & 141–144.
- 471. —. Ueber die Nahrung der Landthiere. SB. Ges. Leipzig, xvii, xviii, Jhrg., pp. 14-17.
- Weber das Plankton. SB. Ges. Leipzig, Jhrg. 1891-92, pp. 86 & 87.
- 473. SMITH, W. W. Evolution of Colours in the Vanessæ. Ent. xxv, pp. 27 & 28.
- 474. Solger, B. Zelle und Zellkern. (Bd. 111. Thiermedizinische Vorträge herausgegeben von G. Schneidemühl.) Leipzig: 8vo, 61 pp., 1 pl.
- 475. °—. Ueber die Architektur der Stützsubstanzen. Leipzig: 8vo, 36 pp., 6 figg.
- 476. Sorel, G. A propos des lois mathématiques de M. Delbœuf. Rev. Sci. xlix, pp. 461-465.
- 477. SPULER, A. Zur Phylogenie und Ontogenie des Flügelgeäders der Schmetterlinge. Z. wiss. Zool. liii, pp. 597-646, 2 pls. Re Evolution, Eimer's Theory, &c.
- 478. STOLL, O. Zur Zoogeographie der landbewohnenden Wirbellosen. Viert. Ges. Zürich, xxxvii, pp. 233-273.

- 479. STRACHLEY, E. O. Die Karyomitose bei abnormer Entwicklung des Forelleneies. MT. Emb. Inst. Wien, xii, pp. 20-22.
- 480. STRICHT, VAN DER. Nouvelles recherches sur la genèse des globules rouges et des globules blancs du sang. Arch. Biol. xii, pp. 199-344, 6 pls.
- 481. —... Contribution á l'étude de la sphère attractive. Arch. Biol. xii, pp. 741-763, 1 pl.; Bull. Ac. Belg. xxiii, pp. 167-192, 1 pl.
- 482. SUCHETET, A. Les Oiseaux hybrides rencontrés à l'état sauvage. 3 Partie. Mém. Soc. Zool. v, pp. 253-525.
- 483. °Sussdorf, M. Lehrbuch der vergleichenden Anatomie der Haustiere. Stuttgart: 8vo, Lief 2, pp. 161-320, 45 figg.
  TASCHENBERG, E. L. [See Brehm (73).]
- 484. Taschenberg, O. Historische Entwicklung der Lehre von der Parthenogenesis. Abh. Ges. Halle, xvii, pp. 365-454.
- 485. —. Bibliotheca Zoologica. II. Verzeichniss der Schriften über Zoologie, welche in den periodischen Werken enthalten und vom Jahre 1861–1880 selbständig erschienen sind. Bearb. von O. Taschenberg. Lief x, Sign. 361–400. Leipzig: 8vo, pp. 2929–3248.
- 486. TETENS, H. Resultate der anatomischen Untersuchung eines lateralen Zwitters von Smerinthus populi (L.), nebst einiger daran geknüften allgemeinen Betrachtungen. B. E. Z. xxxvi, pp. 457-466, 1 pl.
- 487. Thomson, J. A. Outlines of Zoology. Edinburgh: 8vo, xiv & 640 pp., 32 full-page illustrs.
- 488. . The Study of Animal Life. London: 8vo, xii & 375 pp., figg.
- 489. Thomson, J. A., & Wyld, N. The Facts of Sex in Relation to Metabolism: a Review of some Recent Theories. P. R. Phys. Soc. Edinb. xi, pp. 249-282.
- 490. Tichomirow, A. Signification des recherches embryologiques pour la classification. Congrès. Internat. Zool. ii, part i, pp. 32-48.
- TODARO, F. Sulla struttura, la maturazione, e la fecondazione dell' ovo della Seps chalcides. Atti [Rend.] Acc. Rom. vii, pp. 445-449. Cf. Monit. Zool. Ital. iii, pp. 6-8.
- 492. °TORALBO, L. Contributo alla conoscenza del nucleo cellulare nelle glandole della pelle degli anfibi. Month. Int. J. Anat. Hist. ix, pp. 89-94, 2 pls.
- 493. \*Tschemen, J. A. Darwinismus, eine Untersuchung der Darwin'schen Theorie der Abstammung des Menschen. [Russian.] Odessa: 8vo, 508 pp.

- 494. TYLDEN, H. J. The Bearing of Pathology upon the Doctrine of the Transmission of Acquired Characters. Nature, xlvi, pp. 302– 305.
- 496. Urrein, F. Beobachtungen über die zeitliche Succession des Auftreten der Farbenfelder auf den Puppenflügelchen von Pieris brassicæ. Zool. Anz. xv, pp. 284-290 & 293-299.
- Ueber Eigenschaften der Schuppenpigmente einiger Lepidopteren-species. T. c. pp. 299-306.
- 498. Variony, H. Dr. Experimental Evolution. Lectures delivered in the Summer School of Art and Science, University Hall, Edinburgh. London: 8vo, ix & 271 pp.
- 499. —... Les espèces qui s'en vont [after Lucas]. Rev. Sci. xlix, pp. 560-562.
- 500. Variot, G., & Βεzançon, P. Indépendance de la spermatogénèse et de la sécretion testiculaire proprement dits. Bull. Soc. Anthrop. Paris, iii, pp. 282-288.
- 501. & —. De la spermatogénèse. T. c. pp. 289-294.
- 502. & . Influence de la sécrétion testiculaire sur le développement organique. Iudépendance de cette fonction et de la spermatogénèse dans certains cas. Gaz. med. Paris, lxiii, pp. 229-232.
- 503. Vas, F. Studien über den Bau des Chromatins in der sympathischen Ganglienzelle. Arch. mikr. Anat. xl, pp. 375-389, 1 pl.
- 504. Verhoeff, C. Beiträge zur Biologie der Hymenopteren. Zool. Jahrb. vi (Abth. Syst.), pp. 680-754, 2 pls.
- 505. Neue und wenig bekannte Gesetze aus der Hymenopteren-Biologie. Zool. Anz. xv, pp. 362-370.
- 506. —. Biologische Aphorismen über einige Hymenopteren, Dipteren, und Coleopteren. Verh. Ver. Rheinl. xlviii, pp. 1-80, 3 pls.
- 507. Verworn, M. Die Bewegung der lebenden Substanz. Eine vergleichend physiologische Untersuchung der Contractionserscheinungen. Jena: 8vo, 103 pp., 19 figg.
- 508. ——. Die physiologische Bedeutung des Zellkerns. Arch. Ges. Phys. li, pp. 1-118, 6 pls.
- 509. —. Ueber die Fähigkeit der Zelle, activ ihr specifisches Gewicht zu verändern. Op. cit. liii, pp. 140-155, 1 fig. VETTER, B. [See ROMANES (434).]
- 510. VIALLETON, L. La division indirecte des cellules. Rev. Sci. xlix, pp. 678-687, 12 figg.
- 511. \*—. La spermatogénèse chez les mammifères et chez l'homme. Lyon Médicale, lxix, pp. 383-396.
- 512. VIGUIER, C. L'heliotropisme des Nauplius. C.R. cxiv, pp. 1489-1492.

- 513. Virchow, H. Das Dotter-Organ. Arch. mikr. Anat. xl, pp. 39-101, 2 pls.
- Das Dotterorgan der Wirbelthiere. Z. wiss, Zool. liii, Suppl,
   pp. 161-206, 1 pl.
- 515. —... Dotterzellen und Dotterfurchung bei Wirbeltieren. Verh. Anat. Ges. vi, 1892, pp. 209-219.
- 516. Virchow, R. Transformation and Descent. J. Bacterol. Pathol. i, pp. 1-12.
- 517. VOELTZKOW, A. On the Oviposition and Embryonic Development of the Crocodile. Ann. N. H. ix, pp. 66-72.
- 518. Vogt, C., & Yung, E. Traité d'anatomie comparée pratique. Paris : 8vo, vol. 2, pp. 513-640, illustr.
- 519. VOGT, J. G. Die Menschwerdung. Die Entwicklung des Menschen aus der Hauptreihe der Primaten und die Begründung der weiten Kluft zwischen Thier und Mensch, abschliessend mit der vollständigen Lösung des Willensproblems, &c. Leipzig: 8vo, iv & 393 pp., illustr.
- 520. \*STAATS VON WACQUANT-GEOZELLES. Weitervererbung von Albinismus. Zool. Gart. xxxiii, pp. 356-366.
- 521. WAGNER, F. v. General Observations on Fission and Gemmation in the Animal Kingdom. (Trans.) Ann. N. H. x, pp. 23-54.
- 522. WAGNER, J. Ueberblick über den gegenwärtigen Stand der Frage von dem Wesen und der Bedeutung der Befruchtung (Russian). Rev. Sci. Nat. St. Pétersb. iii, pp. 88-91 & 145-155.
- 523. WAITE, E. R. Albinos and White Varieties. Zool. xvi, p. 25.
- 524. WALLACE, A. RUSSEL. Island Life; or the Phenomena and Causes of Insular Faunas and Floras, including a Revision and Attempted Solution of the problem of Geological Climates. 2nd and Revised Edition. London: 8vo, xx & 563 pp., 26 maps and illustr. Cf. Nature, xlvi, p. 56.
- 525. —. Note on Sexual Selection. Nat. Sci. i, pp. 749 & 750.
- 526. WARNER, F. Co-ordination of Cellular Growth and Action by Physical Forces. Rep. Brit. Ass. 1892, pp. 761 & 762.
- 527. WASMANN, E. Die internationalen Beziehungen von Lomechusa strumosa. Biol. Centralbl. xii, pp. 584-599 & 638-669. Re Instinct.
- 528. Die zusammegesetzen Nester und gemischten Kolonien der Ameisen. Ein Beitrag zur Biologie, Psychologie, und Entwicklungsgeschichte der Ameisengesellschaften. Münster-i.-w.: 8vo, viii & 264 pp., 2 pls., 16 figg.
- 529. WATASE, S. The Origin of the Sertoli's Cell (Spermatoblasts).

  Am. Nat. xxvi, pp. 442-144.

- 530. [WATASE, S.] On the Significance of Spermatogenesis. Am. Nat. xxvi, pp. 624-626.
- On the Phenomena of Sex-differentiation. J. Morph. vi, pp. 481-493.
- 532. WATERS, B. H. Primitive Segmentation of the Vertebrate Brain. Q. J. Micr. Sci. xxxiii, pp. 457-475, 1 pl.
- 533. Webber, H. J. Phenomena and Development of Fecundation. Am. Nat. xxvi, pp. 103-111 & 287-310.
- 534. Weir, J. Jenner. Albinism in Birds and Mammals. Zool. xvi. pp. 141-143.
- 535. Weismann, A. Das Keimplasma. Eine Theorie der Vererbung. Jena: 8vo, xviii & 628 pp., 24 figg.
- 536. Die Continuität des Keimplasmas als Grundlage einer Theorie der Vererbung. 2 Aufl. Jena: 8vo, 112 pp.
- 537. Aufsätze über Vererbung und verwandte biologische Fragen. Jena: 8vo, vii & 848 pp., 19 figg.
- Essays upon Heredity and Kindred Biological Problems.
   Vol. 2. Edited by E. B. Poulton & A. E. Shipley. London: 8vo, 226 pp.
- 539. Weldon, W. F. R. On certain Correlated Varieties in Crangon vulgaris. P. R. Soc. li, pp. 2-21.
- 540. WERNER, F. Untersuchungen über die Zeichnung der Wirbelthiere. Zool. Jahrb. (Abth. Syst.) vi, pp. 155-229, 5 pls.
- Zoologische Miscellen. 1. Konvergenz oder Verwandschaft.
   Noch etwas über die Zeichnung der Tiere. 111. Die epitrichial Skulptur der Schlangenepidermis. Biol. Centralbl. xii, pp. 268-280.
- 542. WILSON, E. B. On Multiple and Partial Development in Amphioxus. Anat. Anz. pp. 732-740, 11 figg.
- 543. WIEDERSHEIM, R. Das Gliedmassenskelet der Wirbelthiere mit besonderer Berücksichtigung des Schulter- und Beckengürtels bei Fischen, Amphibien, und Reptilien. Jena: text 8vo, v & 266 pp., 40 figg., atlas of 17 pls.
- 544. WILDEMAN, E. DE. Les recherches récentes sur la structure cellulaire et la division du noyau. Neptunia, ii, pp. 1400-1410 & 1411-1444.
- 545. Les recherches récentes sur la structure cellulaire. Bull. Soc. Belg. Micr. xviii, pp. 16-24.
- 546. WINKLER, F. Zur Frage nach dem Ursprung des Pigments. MT. embr. Inst. Wien, xii, pp. 64-80.
- 547. °—. Zur Frage nach dem Ursprunge des Pigments. Wien. Med. Wochenschr. xlii, pp. 1153-1155, 1193-1196, 1230-1232, & 1260-1262.

- 548. Wundt, W. Vorlesungen über die Menschen- und Thier-seele. 2 umgearb. Aufl. Hamburg and Leipzig: 8vo, xii & 495 pp.
- 549. WURM, W. Ueber die Farbe der Vogelfedern. JH. Ver. Württ. xlviii, pp. 41-46.
  - WYLD, N. [See Thomson & WYLD (489).]
- 550. Yung, E. De l'influence des lumières colorées sur le développement des animaux. C.R. cxv, pp. 620 & 621.
- Sur l'existence d'un soi-disant sens de direction ou d'orientation chez l'homme et les animaux. C.R. Soc, helv. lxxiv, pp. 49-51.
- 552. ....... Orientirungssinn bei Menschen und Thieren. Nat. Rundschau, vii, pp. 195 & 196.
  - ---. [See also Vogt & Yung (518).]
- 553. CZACHARIAS, O. Katechismus des Darwinismus. Leipzig: 8vo, viii & 176 pp., 30 figg., 1 pl., 1 portrait.
- 554. ZANDER, R. Ueber den gegenwärtigen Stand der Lehre von der Zelltheilung. Biol. Centralbl. xii, pp. 281-309.
- 555. ZIEGLER, H. E. Ueber die embryonale Anlage des Blutes bei den Wirbeltieren. Verh. deuts. Z. Ges. ii, pp. 18-30.
- 556. ... Ueber den Begriff des Instincts. Op. cit. 1892, pp. 122-136.
- 557. ZIEGLER, J. Tierphänologische Beobachtungen zu Frankfurt am Main. Ber. Senck. Ges. 1892, pp. 47-69.
- 558. ZIMMERMANN, K. W. Pigmentzellen mit Attraktionsphären. Verh. Anat. Ges. vi (1892), pp. 274 & 275.
- 559. ZITTEL, K. A. VON. Handbuch der Paläontologie. 1 Abt. Paläozoologie, Bd. iv, Lief. 1, Mammalia. München and Leipzig: 8vo, 1-304 pp., 245 figg.
- 560. Zoja, L. & R. Intorno al plastiduli fuesinophili (bioplasti dell' Altmann). Mem. Ist. Lomb. vii, pp. 237-270, 2 pls.
- 561. —. Sur les plastidules fuchsinophiles. Bioplastes d'Altmann (conclusion). J. Microgr. xvi, pp. 13-15.
- 562. ZOPF, W. Beiträge zur Physiologie und Morphologie Niederer-Organismen. 1 Heft. Leipzig: 8vo, iv & 97 pp., 3 pls.
- 563. °ZSCHOKKE, E. Weitere Untersuchungen über das Verhältnis der Knochenbildung zur Statik und Mechanik des Vertebraten-Skelettes. Zürich: 4to, 102 pp., 11 col. pls., illustrs.
  - Re Evolution.

## II.—SUBJECT-INDEX.

The object of this part of the RECORD is to bring together those contributions which have a direct bearing on the larger biological problems. Such of these papers as are based on special examples are also included in the Special Records; these latter, moreover, include many observations on variation, mimicry, habits, &c., that cannot be noticed in this General Record.

- A. Text-Books, Essays, and General, p. 36.
- B. Cell and Protoplasm, p. 42.
- c. Oogenesis, Spermatogenesis, Fertilisation, p. 44.
- D. Sex, Reproduction, and Heredity, p. 46.
- E. Evolution, p. 49.

## A. TEXT-BOOKS, ESSAYS, AND GENERAL.

## (1) TEXT-BOOKS, &c.

Text-books: Bonnier (63), Campbell (89), Carlet (90), Ellenberger (148), Engel (153), Girod (204), Griffiths (212), Hertwig, O. (254), Hertwig, R. (258), Hurst (341), Laborde (315), Lameere (316), Lang (318), Marshall (341), Minot (355), Perrier, E. (393), Perrier, R. (392), Rawitz (420), Retterer (423), Ritzema-Bos (327), Roule (436), Schofield (456a), Sussdorf (483), Thomson (487), Vogt & Yung (518), Zittel (559).

Other books of a general nature: BATES (32), BREHM (73), BRONN (75), DISTANT (125), HUDSON (271), HUTCHINSON (274), KIPLING (294), MOSELEY (367), NEUMANN (374), THOMSON (488).

# (2) Nomenclature.

Orientation of the body: SCHULZE (460) proposes a complete series of unambiguous terms. All not quite irregular bodies may be grouped according as their middle is defined by a point, a line, or a plane,—synstigma, syngramma, sympeda. Around the centre is central or proximal, the direction leading towards this is centrad or proximad, the position is centran or proximan. So distal, distad, distan; axial, axiad, axian. The two ends of the principal axis are rostral and caudal. The two ends of the dorso-ventral axis are dorsal and ventral. Related terms end in n and in d as above. The median plane separates dextral and sinistral, the frontal plane separates ventral and dorsal, the transversal plane separates caudal and rostral.

The terms proximal and distal; FRORIEP (179).

Terms of auxology: Buckman & Bather (79) suggest the following terminology:—

Ontogeny.		PHYLOGENY.	
Anaplasis .	. Embryonic. Brephic. Neanic.	Same terms,	Epacme.
Metaplasis .	. {Ephebic.	with prefix	}Acme.
Cataplasis .	.{Gerontic. Catabatic. Hypostrophe.	"phyl."	Paracme,

Zoological nomenclature: BLANCHARD (58, 59), GIRARD (203), HERD-MAN (250), TICHOMIROW (490).

# (3) Taxonomic.

Kennel (293) sees seven classes or cycles of animals, viz. :-

- I. Protozoa.
- II. Metazoa.

## Сногоркомзку (95) gives the following arrangement:-

| Enterocala: incl. Brachiopoda, Echinoderma, Chatognatha, Chordata, Enteropneusta.
| Genitocala: incl. all Calomata not Enterocala, in which the coelome is said to arise in the compact mass of mesoderm, which corresponds to the sexual tissue of Platodes.
| Acala: incl. Calentera, Porifera, Platodes, Nemertini, Orthonectida, and Dicyemida.

Embryology and classification: TICHOMIROW (490).

Affinities of Vertebrata; ROULE (437).—Affinities between Arthropoda and Vertebrata; BORDAGE (64).—Affinities of Vertebrata; ROULE (437).

## (4) DISTRIBUTION.

AGASSIZ (2), AFSTEIN (18, 19), BLES (69A), BRANDT (71), CLARKE (97), FIELD (165), FISCHER-SÍGWART (166), IMHOF (277-282), KELLER (291), KRÜMMEL (312), MARENZELLER (338), MÖBIUS (357), SCHÜTT (461), SIMBOTH (472), STOLL (478), WALLACE (524).

## (5) Morphological.

General questions of theoretical morphology; DRIESCH (130-132).—Suggested correction of the prevalent conception of homology; BATESON (33).

Principles of skeleton-making in Rhizopods, Sponges, and Echinoderms; DREYER (128, 129).—Architecture of skeletal substances; Solger (475).

Mesoderm and Metamerism; PERENYI (391), RABL (410, 411).—Metamerism of Enterocala due to ramifications of the gut; of Genitocala due to linear budding; Cholodkowsky (95).—Metamerism of Vertebrata; Burckhardt (85): of Vertebrate head; Herrick (252).—Primitive segmentation of the Vertebrate brain; Waters (532).—Spiral modification of metamerism; Morgan (364).

Relation of size of brain to size of body; LATASTE (320).

Vertebrate nervous system: Brain, EDINGER, Selachii, Amphibia (141); KÖPPEN, Lizard (303); SCERVINI, General (449); RETZIUS, Cerebellum, sympathetic ganglia (424); BURCKHARDT, Phyletic history of the brain (85).—Cranial nerves; FRORIEP (178), KUPFFER (314).—Parietal eye; BERANECK (47).

Pituitary body; SAINT-REMY (448).

Eye; Matthiesen (343).

Vertebrate ear; AYERS (25).

Epiglottis: GEGENBAUER (193).

Pectoral and pelvic girdles, especially of Fishes, Amphibians, and Reptiles; WIEDERSHEIM (543).

Urino-genital system; Semon (464).

Morphology of Cyclopian monsters; EMERY (151).

# (6) PHYSIOLOGICAL.

Physiology of the *Invertebrata*; GRIFFITHS (212).—Comparative physiology of domestic animals; ELLENBERGER (148).—Chemical basis of animal body; Lea (322).

Comparative pathology: "Tracing the significance of amœboid cells, or phagocytes, from the *Protozoa* upwards, through various groups of animals to the higher Vertebrates"; Metschnikoff (347).—Struggle for existence among the parts of the organism; Metschnikoff (348).

Theory of vital movement: In the expansion of a pseudopodium there must be local lessenings of the surface tension; this diminution of the surface tension depends on the chemical affinity of the protoplasm for oxygen; the plasmic particles are drawn within the operative sphere of the oxygen molecules in the medium, and, being satisfied, remain indifferent, or are shoved aside by their successors. Contraction is an expression of increased surface tension; this increase occurs when the complex plasmic particles break up on being stimulated; the disrupted particles are drawn to the centre of the unit mass; the reason of this is in the nuclear substance, for which the exhausted protoplasm now exhibits chemotropism. Application of these ideas to Vorticella, and to muscle; Verworn (507).

Origin of striped muscle; EIMER (145), HAYCRAFT (237).

Muscle and nerve; McKendrick (332).

Movement of animals studied by photographs; MAREY (339), PASSY (389).

General physiology of the cell; Hertwig (256).—Comparison of "microscopic foams" with protoplasm; Bütschli (83).

Studies on growth; Cornevin (104).

Life and death; SABATIER (446).

Death; MITCHELL (356).—La synthèse des êtres vivants; Sabatier (447).

Residual life; GAUTIER & LANDI (188-191).

Nervous system; Horsley (268).

Digestion and absorption (Artemia, &c.); FRENZEL (175, 176).

Comparative physiology of respiration; GAGE (182, 183).

Blood of the *Invertebrata*; GRIFFITHS (213, 214).—Hæmocyanin; FRÉDERICQ (173), CUÉNOT (114), HEIM (241).

Iron in the animal body; SCHMUL (454), SCHNEIDER (456) [cf. SIM-ROTH, Die Enstehung der Landthiere, 1891].

Autotomy of shore-crab; FRÉDERICQ (174).—Autotomy of Spiders and Pycnogonids; GAUBERT (185, 186).

# (7) Animal Intelligence and Instinct.

AAS (1).—Limits of animal intelligence; DIXON (127), MIVART (127), MORGAN (127, 362).—Relation of human and animal "Seele"; BIEGE-LEBEN (52).—Mental evolution in man and animals; BODINGTON (60).—Theory of sensation; DUBOIS (134).—History of instinct of Pompilidæ; FERTON (162).—Societies of Termites; GRASSI (208).—Numerous observations; HUDSON, W. H. (271), and HUDSON, G. V. (272).—Estimate of animal intelligence; JACQUART (284).—Parental care; KRÄPELIN (309).—Analysis of instinct of Anmophila; MARCHAL (337).—Intelligence and instinct; NADAILLAC (370).—Development of intellect; Schneider (455).—Psychology of societies of Ants; WASMANN (528).

-International relations between Lomechusa and Ants; WASMANN (527).—Sense of direction; Yung (551).—Conception of instinct; ZIEGLER (556).—Revised edition of lectures; WUNDT (548).


## (8) Embryological.

Introduction to study of embryology; ROULE (436).

Text-Books: Herrwig (254), Korschelt & Heider (306), Minot (355).

Old and new conceptions of development; Herrwig (257).—Das biologische Grundgesetz; ARNDT (21).—On a supposed law of Metazoan development: Alternation of generations seems to underlie Metazoan development. The birth of the Nemertine on the Pilidium, that of the Echinoderm on the Pluteus, may be rather normal than exceptional. A larval stage lurks in all cases, but it may be represented merely by a blastoderm upon a yolk-sac, or by less than that. But the larva never disappears, however much it may undergo degeneration. In Mammals, where it is thoroughly parasitic, it is most reduced; BEARD (37, 38). See also Assheton (37).

Theory of germinal layers; Houssay (270):-



Theory of mesoderm and metamerism; Cholodkovsky (95). Theory of mesoderm; RABL (410).

Homology of germinal layers: gastrulation in Vertebrata; REPIACHOFF (422).

Development of optic nerve of Vertebrates: The optic stalk has no share in forming the nervous parts of the eye; the nerve-fibres grow inwards from the retina towards and into the brain; Assheton (23).

Histogenesis of nerve; Beard (36), and Herrick (253): contra, Beard and Dohrn; Kölliker (302).

Nerve-cells and nerve-fibres: All nerve-fibres are direct processes of nerve cells; they are not formed by the concrescence of rows of cells; Kölliker (301).

Transient ganglion cells; BEARD (39).—Cerebro-spinal ganglia: at a certain stage in the development of Amniota all the nerve cells of the spinal ganglia are opposito-bipolar (a form permanent in fishes), but they become unipolar. The spinal ganglia of Vertebrates are the origins of the sensory part of all the spinal nerves. The ganglia of 5th, 9th, and 10th cranial nerves are comparable to spinal ganglia; Genuchten (195, 196).

Wirbel and Urwirbel; EBNER (140).

Blood corpuscles; STRICHT (480).—Report on state of knowledge as to the history of blood corpuscles; Oppel (384).—Origin of the blood in Vertebrates; Ziegler (555).

Yolk-organ of Vertebrates; VIRCHOW (513-515).

Physiological factors determining metamorphoses: "Theory of metamorphosis as due to asphyxia"; BATAILLON (30).

Experimental embryology: General accounts of recent work; Andrews (9), Korschelt (305), Roux (438).—Experiments on Frog ova: a loss of even one-sixth of the substance of the ovum produced only circumscribed local defects or disturbances. Puncturing one of the two first segmentation cells resulted in a typical half-morula, half-gastrula, half-embryo. Right and left half-embryos, and also anterior and posterior half-embryos were produced. By "post-generation" the half-embryo may form a whole by vitalising the injured half-egg, which has been lying passive, while the uninjured half has been developing; but Roux has also succeeded in rearing an entire frog-embryo from half an egg, without any co-operation on the part of the other injured half; Roux (305).

Influence of electric current on ova, and their segmentation; Roux (306).

Experiments on ova of Echinoids: With increased temperature the ova of sea-prchius [Spherechinus] formed double blastulæ, gastrulæ, and even Plutei. The mode of segmentation is also modified. If one cell of the 4-celled stage be shaken off or destroyed, the remainder may form a normal Pluteus; and an isolated cell may form a blastula, or even a Pluteus. Pressure modifies very considerably the course of segmentation, but yet after removal of the disturbing influence normal larvæ result; DRIESCH (130).—The addition of various salts (of potassium, lithium, &c.) to sea-water delayed the formation of the larval skeleton, inhibited the characteristic Pluteus outgrowths, or resulted in the production of altogether abnormal larvæ. It seems that the influence on the developing ova is not directly chemical, but is due to the altered osmotic pressure of the sea-water; Herbst (249).

Artificial stimulus resulting in the production of an egg-membrane: Unfertilised ova of sea-urchins placed in sea-water and shaken up with chloroform, benzol, toluol, &c., form a protective membrane similar to that which appears after the entrance of a spermatozoon. Even on fertilised ova, robbed of their normal membrane, a second membrane was formed in response to artificial stimulus; Herbst (249).

Multiple and partial development of Amphioxus ova: From the cells of the 2-celled stage shaken apart, two embryos of half-size result; similarly from the 4-celled stage four embryos of quarter-size were produced. Incomplete separation of the cells resulted in various forms of connected dwarf-embryos. A unit from the 8-celled stage does not, however, form an embryo. According to Wilson this inability is due to

qualitative limitations. The segmentation of isolated blastomere of the 2- or 4-celled stage agrees exactly with that of the entire ovum; WILSON (542).

Incubation during continuous rotation; Dareste (118).—Influence of light on orientation of chick embryo; Blanc (57).—Influence of coloured light on development; Yung (550).

Urmund and Spina bifida: Study of abnormalities which occur in the development of Frog ova, in consequence of over-ripeness and polyspermy. The segmentation may be strangely irregular, large parts of the yolk may remain undivided, and the formation of the gastrula may be so disturbed that the Urmund remains partially or wholly unclosed. Morphological interpretations of resulting abnormalities; Herrwig (255).

The terms of auxology; BUCKMAN & BATHER (79).

## B. CELL AND PROTOPLASM.

General treatises and essays on the Cell: HERTWIG (256), CHATIN (93), HEIDENHAIN (239), SOLGER (474), FRANCKE (171).

Physiology of the Cell and Protoplasm: Theory of protoplasmic movements; Verworn (507).—The power cells have of altering their specific gravity; Verworn (509).—Chemical irritability of protoplasm; ROEMER (429).—Physiological elements of protoplasm; Detmer (123).—Cell physiology; Lilienfeld & Monti (325).

Protoplasm: General considerations; Beale (35).—Activities of protoplasm compared with the phenomena seen in observing fine foams; BUTSCHLI (83). The author's general conclusion is that protoplasm has a frothy or foam-like structure, and that the physical conditions observable in the drops of emulsion do, mutatis mutandis, help us towards an understanding of the streaming movements and changes of shape in protoplasmic units. Cytoplasm and nucleoplasm have a foam-like structure, and consist of a honeycomb-like arrangement of lamellæ, with fluid or enchylema in the interspaces. The structure may be mimicked by means of fine emulsions or foams. Appearances of fibrillation, reticulation, nodal thickenings, &c., are readily produced artificially. In various conditions amoeboid streaming movements occur.

Structure of Cytoplasm: Altmann (6, 7), Bütschli (83), Crato (108), Dannehl (117), Fayod (161), Flemming (168), Guignard (217, 218), Kromayer (311), Prenant (405-408).

Nucleus: Nucleus and Protoplasm; Heidenhain (239).—Physiological import of nucleus; Verworn (508); only through its metabolism does it influence the life of the cell.

Structure and Functions of Nucleus; MANN (335). The nuclear chromatin segments are organs for the transformation of simple into more complex compounds. The nucleolus is a storehouse of elaborated material. The achromatin is the essential plasma. Two special seats of the achromatin substance are the archoplasma and the endo-nucleolar substance.—

Experiments on function of nucleus; CLARK (99).—Structure of nucleus; Altmann (7).—Structure of resting nucleus; Krasser (310).

Rabl's doctrine of the nuclear segments; HARTOG (234).

Affinity of nucleus for iron-staining chromatin; GILSON (201).

Nuclei in glands of Amphibian skin; TORALBO (492).

Attractive spheres and centrosomata; Bürger (82), Bütschli (84), Fick (163), Hansemann (233), Heidenhain (238), Lebrun (323), Moore (360), Nicolas (376), Stricht (481), Zimmermann (558).

Cell-division: Present state of knowledge; Zander (554).—Rôle of archoplasm; Moore (360).—Indirect division, general account; Vialleton (510).—Direct division; Nicolas (377).—Division of nucleus and of cell in epithelial tumours; Borrel (66).—Nuclear fragmentation followed by the division of the cell; Buscalioni (86).—Amitotic cell-division; Frenzel (177).—Heterotypical nuclear division in origin of reproductive elements; Hæcker (226).—Nuclear division in formation of mesoderm and endoderm in Cyclops; Hæcker (227).—Asymmetrical mitosis; Hansemann (232).—Amitosis in embryonal envelopes of Scorpion; Johnson (289).—Eismond (147).

The fate of the central spindle in Karyokinesis; Kostanecki (307).

Nuclear division in giant cells; KOSTANECKI (308).

Terminology, for Aster, Dyaster, &c., it seems desirable to substitute Astroid, Dyastroid, &c.; FLEMMING (167).

Flemming's "intermediate body" in the seminal cells of Scolopendra and Lithobius; PRENANT (404).—Origin of achromatic spindle; PRENANT (408).

Primitive fibrils; APATHY (14). The primitive fibrils of smooth muscle and of nerve have been hitherto overlooked; the interfibrillar spaces have been seen, not the fibrils. Contractile substance and conducting substance are alike intracellular plasmic products, and consist of primitive fibrils and of intrafibrillar substance.—Origin of muscle fibres; EIMER (145), HAYCRAFT (237).— Muscle and nerve; ROHDE (430, 431).

Cell-membrane; Buscalioni (87).—The basilar membrane; Chatin (94).

Continuity between cells; OLIVIER (382).

Biology of Protozoa; OSBORN (387).

Histology of nervous system; RETZIUS (424), ROHDE (430, 431), VAS (503).

Chemical properties of reticular tissue; SIEGFRIED (468).

Schlummerzellen und Gewebsbildung; EBERTH (138), GRAWITZ (210), SCHMIDT (453).

Miscellaneous; Meltzer (345).

c. OOGENESIS, SPERMATOGENESIS, FERTILISATION. General biological considerations; Weismann (535), Henking (245).

## (1) OGENESIS, MATURATION, &c.

Embryological classification of ova; Henneguy (247).—Comparison of male and female reproductive cells; Prenant (403).—Heterotypical nuclear division in oogenesis and spermatogenesis; Hæcker (226).—Graafian follicle, history of; Schottländer (457).

Yolk-cells and yolk-segmentation; Virchow (515).—Nutritive importance of the yolk-sac; Robinson (428).

Yolk-organ of Vertebrates; VIRCHOW (513-515).—Binucleate ovum of rat; Blanc (56).

Ovum of reptiles; Arnold (32). — Oogenesis in Seps chalcides; Giacomini (198), Mingazzini (354), Todaro (491).

Maturation in ovum of Trout; Blanc (54).—Ovary and ovum of Gobius minutus; Guitel (221).—Ova of Turbot; Holt (265).—Oogenesis, &c., in Teleosteans; Fusari (180).—Oogenesis in Teleostei: Ovarian ova of Dab, Hake, &c.; in all ripening ovaries, ova for three consecutive spawning periods are present; Calderwood (88).—Abnormal karyomitosis in Trout ovum; Strachley (479).—Ovarian ovum of Selachii: Complex alterations of mass and form exhibited by the chromosomata. It can hardly be a "Keimplasma" which undergoes such alterations of mass; it must be an involved somatic plasma. Proposed theory of mutual fertilisation, or amphimixis of chromosomata; Rückert (440, 442).

History of the germ-cells in *Insecta*: An important treatise on cogenesis, maturation, and fertilisation in *Insecta*. Discussion of eggenvelopes, the formation of polar bodies, the "thelyid" (an achromatic polar body), &c. The first and second divisions of the spermatocytes are comparable to the divisions which form polar bodies. Polyspermy common. The arrhenoid of the spermatozoon comparable to the thelyid. Comparison of the processes of fertilisation in Insects, *Protozoa*, and Angiosperms; Henking (245).—Oogenesis in *Cyclops* and *Canthocamptus*; H.ECKER (225).—History of ovum in *Branchipus grubii*; Brauer (72).

Early development of Cirripedia, maturation, fertilisation, &c.; Groom (216).—Oogenesis in Diaptomus; ISCHIKAWA (283).—Centrosomata in ovum of Ascaris megalocephala; LEBRUN (323).—Vitelline nucleus of ova of Trematoda; Monticelli (358), and Crety (109).—In Echinoderms; Cuénot (113).

# (2) FERTILISATION.

Fertilisation of ova in Echinoderms, Selachii, Reptilia; EBERTH (137): in Reptiles; Oppel (383): in Axolotl; Fick (164). The middle portion of the spermatozoon gives rise to an attractive sphere, which seems to divide; Fick (164).

Immature ova of *Triton teniatus*; BORN (65): Trout; BÖHM (61). Trout: In fertilisation, the two pronuclei fuse, and so do their solar centres; BLANC (54, 55).

Invertebrates; PICTET (396).—Insects; HENKING (245).—Diaptomus: The primary sperm-cells and the primitive ova have eight chromatin elements; the pronuclei have four; there is a reducing division; in the first two segmentation-spheres there are likewise eight chromosomes; ISCHIKAWA (283).

Parthenogenesis (?) in Frog: of about 1500 unfertilised ova (of Rana fusca) about 100 exhibited irregular segmentation furrows; Dehner (122).

General problems of fertilisation; Hertwig (259), Lwoff (329), Müller (369), Wagner (522), Webber (533).

Comparison of the process of fertilisation in plants and in animals; GCIGNARD (219).

The protective membrane formed after fertilisation; Herbst (249) [see Embryology].

Polyspermy: In Seluchii: Most, if not all, of the merocyte nuclei (vitelline or parablast nuclei) in the ova of the Selachians are modified male pronuclei; polyspermy appears to be normal; RUCKERT (441).—In Reptiles: Accessory sperm-nuclei in Anguis, Tropidonotus, Lucerta; polyspermy appears to be frequent; Oppel (383).—In Axolotl: Sometimes as many as nine spermatozoa in ovum; Fick (164).—In Insects; Henking (245).

Results of over-fertilisation: In the over-fertilised ovum opposing factors conflict; on the one hand there are forces tending to development and incited by the fertilisation, on the other hand there are forces of an inhibiting and disturbing nature, due to the injury the ovum has sustained by over maturation, or by other abnormal conditions operative before fertilisation; HERTWIG (255) [see also Embryology].

# (3) SPERMATOGENESIS.

General significance of spermatogenesis; WATASE (530).—Comparison of spermatogenesis and oogenesis; HENKING (245).—Morphological comparison of ovary and testis; PRENANT (403).—Heterotypical nuclear division in oogenesis and spermatogenesis; H.ECKER (226).—Especial details in regard to reduction-divisions in Gryllotalpa vulgaris; RATH (419).

Histology of testis; GERMANO (197).—Distinction between spermatogenesis and testicular secretion; VARIOT & BEZANÇON (500-502).—Histology of spermatozoon; BALLOWITZ (27).

Spermatogenesis: in various Invertebrates; PICTET (396): in Trematoda; Monticelli (359): in Echinoderms; Cuénot (113): in Diaptomus; Ischikawa (283): in Scolopendra, Lithobius; PRENANT (403-408): in Frog; Bertacchini (50): in Salamander; Nicolas (376): in Sauropsida; Benda (45): in Mammals and Man; Bardeleben (29), Vialleton (511).

Origin of Sertoli's cells; WATASE (529).

Spermine, &c.; Brown-Sequard (77), Duclaux (135), Héricourt (251), Piehl (398, 399).

## p. SEX, REPRODUCTION, HEREDITY.

## (1) SEX AND REPRODUCTION.

General considerations; Weismann (535), Henking (245).—Summary of recent observations and theories; Webber (533).—Reproduction among plants and animals; Neviani (375).—Life and death; Sabatier (446).—Origin of contrast between somatic and reproductive cells in Ascaris; Boyeri (68).

Evolution of sex: The male organism inclines to greater activity, the female to greater passivity, a conclusion similar to that maintained by Geddes and Thomson in "The Evolution of Sex;" FRANCKEN (172).

Sex-differentiation: Important to distinguish (a) the phenomena of heredity, or the mixture of the parental characteristics in the offspring; and (b) the phenomena of sexual differentiation in the organism, in which the parental characters have already been mixed. The micro-chemical reaction of the sperm-nucleus, which is entirely different from that of the ovum-nucleus during the earlier stages of its existence, becomes essentially like that of the female pronucleus at the moment of fecundation. The phenomenon of sex-differentiation is one with that of irritability, and is the most pronounced of all the phenomena of protoplasmic irritability, because the stimulus acts at an early stage of ontogeny, and thus at the outset profoundly deflects the course of behaviour of the whole organism; Watase (531).

Nature of sex: Critical review of some recent theories in regard to sex and reproduction. In contrasting the sexes, their reproductive elements, &c., the contrast must be expressed in contrasted ratios of anabolism (A) and Katabolism (K). Thus, in the female the ratio  $\frac{A}{K}$  is greater than the corresponding vital ratio in the male. This is designed as a correction of the general thesis of the "Evolution of Sex;" Thomson & Wyld (489).

Origin of sex: Conditions unfavourable to ready assimilation are believed to tend to produce male offspring, reverse for female. In conjugation the starving cell is fed on surplus nourishment of the over-fed cell; Mann (336).

Parthenogenesis: History of our knowledge; TASCHENBERG (484).—Apparent parthenogenesis in ova of Frog; DEHNER (122).

Dissogonie: a new form of sexual reproduction; CHUN (96).

Fertilisation and conjugation; HERTWIG (259).

"Gamophagie:" According to the author, one germinal substance assimilates the other in fertilisation; Müller (369).—Fission and germation: Wagner (521).

Hermaphroditism: Traces of hermaphroditism in the Lamprey, not without interest in reference to speculations as to the possible hermaphroditism of ancestral Vertebrates; Beard (40).—Hermaphroditism of Astacus; La Valette, St. George (321).—Certain spermatogonia, "untrue to their original destination," may, instead of dividing into spermatocytes, grow into ova.—Lateral hermaphroditism; Tetens (486).

Hybridism; Elliot (150).—Hybrid birds in nature; Suchett (482).

—Fertile sexual union can take place only between those insects whose genital appendages precisely correspond. Hybridism between different species seems mechanically impossible; Escherich (156).

Variability and hybridism of Cyprinoids; Heincke (243).

Breeding in and in; SCHILLER-TIETZ (451).

Biological conclusions illustrated in reference to certain Hymenoptera, Dîptera, and Coleoptera; Verhoeff (506).—Sex and reproduction in Hymenoptera. (1) Proterothesia: in nests of Fossoria, Vesparia, and their parasites, the tenants of the anterior cells are male of the posterior female. (2) Proterocraty: the individuals which appear first in a brood are the strongest. (3) Proterandry, correlated with Proterothesia. (4) Polyandry; Verhoeff (504, 505).—Dichogamy in Lepidoptera. The advantage of dichogamy, or "dichogenesis," lies in the prevention of in-breeding; its occurrence may be readily accounted for by natural selection; Petersen (394).

Peculiarities of reproduction; Seitz (463).

Fecundation in tailed Batrachians; BOULENGER (67).—Detailed study of reproduction in Gobius minutus; GUITEL (222).—Variations in genital organs of Gamasina, and mode of coition; MICHAEL (350).

Determination of sex by nutrition; Landois (317).

Double-broodedness of *Lepidoptera*: whether influenced by the state of the food-plant; MERRIFIELD (346).

Parental care among animals; KRÄPELIN (309).

Conclusions as to the societies of Termites; GRASSI (208).

Regeneration: Regeneration of the tail in Lumbriculus; general considerations as to regenerative processes; RANDOLPH (413).—Regeneration in Annelids: Bifurcated Annelids. Collection of cases. Some general considerations. "The bifurcated monstrosities exhibit what may be a universally present but latent ability of parts of a highly-organized animal to form a complete individual like that to which they belong;" Andrews (11).

Regeneration; WEISMANN (535).—Periodic regeneration of the upper half of the body in *Diplosomidæ*; OKA (381).—Post-generation; Roux (438).

Influence of nutrition on sex; LANDOIS (317).—Influence of nutrition

on reproduction: It seems likely that the nutritive supply has much to do with Proterothesia and Proterocraty in Hymenoptera; Verhoeff (504-506).

## (2) HEREDITY.

Historical and critical survey; sums up in favour of Lamarckian conceptions; Osborn (385, 366).—Nature of heredity; Hurst (273).

Das Keimplasma: This statement of Weismann's theory of heredity centres around his conception of the germ-plasm. This nuclear substance is believed to contain reserve vital units, or biophors, which are combined into units of a higher order—the determinants. Each determinant is capable of controlling one kind of cell, and the germ-cell contains as many determinants as there are different cells, or groups of cells, in the fully-formed organism, which are capable of being individually determined from the germ onwards. The determinants have a definite arrangement in the germ-plasm, forming aggregates of a higher order—the It is suggested that parts of the chromosomes are the ids, and that the nuclear rods are aggregates of ids, spoken of as idants. Apart from the elaborate working out of the conception of a germ-plasm, there are important discussions of regeneration, multiplication by fission and gemmation, alternation of generations, the formation of the germ-cells, amphimixis, reversion, dimorphism, and polymorphism, the non-transmission of acquired characters, variation, &c.; WEISMANN (535).

Translation of Weismann's essays; POULTON & SHIPLEY (538).—Weismann's "Idant" theory of heredity; HARTOG (234).

Transmission or non-transmission of acquired characters; Elliot (149), Knauthe (295), Miles (351), l'ath (418), Riley (426).

Answer to Weismann; Brown-Séquard (78A).

Pathology, so far from offering any support to the hypothesis of the transmission of acquired characters, pronounces against it; Tylden (494).

It is assumed, on insufficient evidence, that the nucleus is the sole bearer of hereditary qualities. The cell-plasma, and especially the centrosomata, must be taken account of; Bergii (48).

In reference to the Hoazin, GADOW (181) states that the crop, by its downward growth, gradually encroaches upon, and modifies, the conformation of neighbouring organs; some of these modifications of the sternal apparatus are already pre-formed to a considerable extent, before they are actually necessitated. They are inherited.

Transmission of acquired characters in Pleurowetidee; GIARD (200).

HENKING (245) does not depreciate the importance of the nucleus as a bearer of hereditary qualities, but he also believes in the importance of the plasma, and in the modifiability of the chromosomata by influences from the protoplasm.

What is an "acquired character"? "Acquired characters, however evident they may be to all but the nec-Darwinians, have the convenient

property of becoming blastogenic, or not acquired, whenever they are proved to be transmitted"; Nutting (380).

Inheritance of albinism; STAATS VON WACQUANT-GEOZELLES (520).

Heredity; GRABHAM (206), MÜLLER (369).

#### E. EVOLUTION.

#### GENERAL QUESTIONS.

Das Keimplasma; Weismann (535) [cf. Heredity, suprà].

Darwin and after Darwin; ROMANES (433); German translation (434).

Catechism of Darwinism; Zacharias (553).

Critical account of modern theories; HAMANN (231).

French predecessors of Darwin: historical and critical; QUATREFAGES (409).

Darwin and Darwinism; Morselli, Canestrini, Cattaneo, & Loria (366).

Present state of the Evolution problem; some arguments for Lamarckian position; Osborn (385,486).

Criticism of Darwinian theory from philosophical standpoint; CREUTZ-BURG (111).

Controverted questions re Evolution, &c.; HUXLEY (275).

Experimental evolution; VARIGNY (498).

Origin of species; EIMER (146).

Elementary description of universal evolution; RAIMOND COULON (107).

On Delboeuf's laws; SOREL (476).

Dynamical theory of organisms; HAUPTMANN (236).

Protest against a materialistic conception of the organism; ARDISSONE (20).

Transformation and descent; VIRCHOW (516).

Convergence or relationship? WERNER (541).

Das biologische Grundgesetz; ARNDT (21).

Urzeugung; Schaaffhausen (450).

Theories of nutrition and heredity, in which energy is not recognised as one of the prime factors in every vital process, should be received with caution; MILES (351).

The importance of recognising the principle of the conservation of energy when considering the evolution of organisms [cf. Herbert Spencer's "Principles of Biology"]; RYDER (444).

General works: see also Coulon (107), Dutoit-Haller (136), FARGES (160), MOTTO (368), TSCHEMEN (493).

#### NATURE AND ORIGIN OF VARIATIONS.

Classification of animal variations; BRANDT (70):-

- II. Acquired . 

  d. Due to external influences.
  b. Due to function.
  c. Due to pathological processes.
  d. Correlated with the preceding.

Variations may be inherited (a) directly from parents, (b) indirectly by influence, (c) by atavism from ancestors.

Definite versus Fortuitous Variation: Discussion; Meehan (344), McMurrich (333), Allen (5), Cope (101).

Fortuitous variation; RILEY (425).

On the "constitution" of Molluses; SIMROTH (470).—Constitutional variation of markings in insects; Escherich (156).—Correlated variations in Crangon rulgaris; Weldon (539).

Struggle for existence among the parts of an organism; Metschnikoff (348).

Translation of essays on Amphimixis, &c.; Weismann (538).

Factors in evolution of Mammals; Morcan (363).

Précis of teratology—human and animal; Guinard (220).—Probable teratogenic origin of two species of Tricladidæ (Dendrocælum nausicaæ and Phagocata gracilis); Hallez (229).—Teratological variations: The malformations which occur in Frog embryos in consequence of the imperfect closure of the "Urmund" are analogous to the Terata mesodidyma and Katadidyma in Teleostean fishes, and to Spina bifida in higher Vertebrates; Herrwig (255).

Variation in internal structures of Gamasina; MICHAEL (350). Variability and Hybridism of Cyprinoids; Heincke (243).

#### INFLUENCE OF ENVIRONMENT.

Results of experimental Evolution; Varigny (498).

Specific variation in Arthropods: A certain kind of variation occurs with peculiar frequency in certain groups, thus sexual dimorphism in Copepods, protective adaptation in *Decapoda*. The habit of masking; HÆCKER (228).

Influence of artificial environmental conditions on ova; DRIESCH (130), HERBST (249), DARESTE (119) and others [see Embryology].

Influence of surroundings on variation; Locard (326).

Colour-relation between Lepidopterous larvæ, pupæ, cocoons, and imagines, and their surroundings; POULTON (402).—Influence of surroundings on colour; BEDDARD (41).—Influence of coloured light on

development: Green Planarians and Hydræ do not grow better with violet than green light, are exceptions to the rule that green light is prejudicial to development. The Hydra grows more quickly in red than white light; Yung (550).—Influence of light on orientation of chick embryo; Blanc (57).

Atrophy, &c., of eyes in cave-inhabiting animals; Joseph (290).

Heliotropic animals; Andrews (10), after Loeb. — Heliotropic Nauplii; Viguier (512).

Influence of light on metabolism: Exclusion of light is associated with increase of weight, more fat, less hæmoglobin, &c.; GRAFFENBERGER (207), EWALD (156A).

Influence of electrical stimulus on ova; ROUX (439).—Influence of electrical stimulus on leucocytes; DINEUR (124).—Galvanic stimulation of aquatic animals; NAGEL (371, 372).

Relations between temperature and protoplasmic movement; CLARK (98).—Influence of refrigeration and desiccation; Kochs (297).

Influence of fresh water on marine animals; GOGORZA Y GONZÁLES (205).

Influence of cocain on Coelenterates, worms, arthropods, &c.; Dani-Lewsky (116).

General discussion of parasitism; Looss (327).

Functional adaptation; MINERVINI (353) and ENDRES (152).

On the mechanical origin of fish scales: The scales have a segmental arrangement; their imbrication is explained by reference to the musculature which, in contraction, throws the integument into rhombic areolæ; RYDER (443).

Evolution of skeletal structures: "An essay towards a mechanical explanation of organic structures." The occurrence of the tetraxial type of skeletal element in Sponges, Echinoderms, and Polycistina, in different materials, suggests independence of vital forces. Vesicular bodies tend to exhibit a tetraxial skeletal structure quite apart from metabolism, for merely physical reasons; DREYER (128).

Relation of ossification to the statics and mechanics of the skeleton; ZSCHOKKE (563).

Peccilogony: when members of the same species have a different outogeny; Giard (199). Cf. Brooks & Herrick (76).

On the relation between the shapes of fishes and the strength of the currents in which they live; SCHERCHOFF (452).

The surface film of water, its relation to the life of plants and animals; MIALL (349).

Effect of sudden pressure; PHILIPPON (395).

Geotropism of lower organisms; Jensen (286).

Experiments on metamorphosis of Amphibians; BATAILLON (31).

### SELECTION AND ELIMINATION.

Struggle for existence: Geometrical representation of relative intensity of the conflict between organisms. For organisms swimming in water

or flying in air, we may say that their accessibility, inter se, and liability to attack is from 60,000 directions; for sessile organisms, or those lying on a plane, from 30,000 directions; for heavy terrestrial forms, from 400 directions; for burrowing or tubicolous, from 314 to 0; for testaceous, operculate, mimetic, or parasitic forms, accessibility almost or quite nil; Ryder (445)

Interspecific competition: Myrmica ruginodis making war on its own species; BIGNELL (53).

Extermination of animals; SCHULZ (459), VARIGNY (499). [Cf. LUCAS, F. A. Animals recently extinct, or threatened with extinction, as represented in the collections of the U. S. National Museum. Washington: 8vo, 1891, 40 pp., 11 pls.]

Natural selection among mankind; Ammon (8): in Lepidoptera; Comstock (100).

Struggle between males: Not only is there a struggle between males, but the large number of males allows a selection of the fittest amongst the fit, and the strongest are first in the field (proterocraty); Verhoeff (505).

Sexual selection; WALLACE (525).

Case of Gelasimus annulipes: The brightly-coloured chola of the male is (1) a club in contests with rival males, and (2) a signal to allure the females, and may be also a stopper for the burrow and a nuptial support; ALCOCK (4).

#### PARTICULAR PROBLEMS.

#### (a) COLOURATION.

Ætiology of animal colouration; Beddard (41).—Physiology of colour-change; Biedermann (51).—Colour-changes in Fishes; Semper (465).

Physiological chemistry of colouration; URECH (496).—Analysis of green pigment in wings of *Lepidoptera*: Regarded as probably excretory; derived from uric acid; GRIFFITHS (215).—Rôle of pigments in histolysis and histogenesis; KUNCKEL D'HERCULAIS (313).

The origin of pigment: Pigment is present in the ova of Bufo almost from the first, and seems to arise from a modification of yolk-plates; WINKLER (546).—Melanic pigment; POUCHET (401).—Melanism and melanochroism in British Lepidoptera; TUTT (495).

Albinism: APLIN (17), FABANI (157), MACPHERSON (334), WAITE (523), WEIR (534).

Colouration in birds; Wurm (549): in birds' eggs; Koenig-Warthausen, Friir. (299), and Raspail (416, 417).

Origin and migration of pigment in Amphibia; EHRMANN (143, 144). Melanism in Fishes; KNAUTHE (296).

Colours of insects: Coste (105, 106), GRIFFITHS (215), HOPKINS (266),

MELDOLA (266), URECH (496, 497), and others [see INSECTA].—Evolution of colours in the *Vanessa*; SMITH (473).

Pigments of ova of Crustacea; Heim (242).

Iridescent colours; Hodgkinson (263).

Protective colouring; JOHNS (288).—Markings of Vertebrates; WERNER (540).

Hints from pathology: CASPARY, EHRMANN, JARISCH, KAPOSI—Die Pathogenese der Pigmentierungen und Entfärbungen der Haut: Verh. 10 Internat. Med. Congress Berlin (1890); Abth. Dermatol. 1892, pp. 91-114.

## (b) MIMICRY.

General discussion of mimicry; special study of  $Papilionid\alpha$ ; HAASE (223).

Recent observations, with criticism; BEDDARD (42).

Mimicry and other adaptations; BRANDES (69).

Alleged "aggressive mimicry" of Volucellæ; Bateson (34).

Aggressive mimicry of Volucellæ; corrections on latter author's book on "Colours of Animals"; POULTON (34).—A notable case: Peridexia (Coleoptera) and Pogonius (Hymenoptera); Heller (244).—Mimicry in small insects; Kennel (292).—Mimicry in Spiders; Banks (28).—Mimicry; Fabani (159).

Protective resemblance in the animal kingdom; PLATEAU (397).

#### (c) MISCELLANEOUS.

Evolution of Flat-fishes; CUNNINGHAM (115) and GIARD (200).

Origin of muscle: The cross-striation is the expression of alternately strong and weak waves in the muscle substance. It is incomplete in sluggish animals, and more marked as activity increases. The striation is, in short, the result of increased function—the mechanical result of contraction; EIMER (145).

The origin of lungs: Air-breathing in fishes supposed to have been originally performed by the unchanged walls of the cesophagus; then a pouch, . . . an air-bladder, . . . a lung. [But how?]; MORRIS (365).

Origin of *Metazoa*: Discussion of *Salinella*; the higher organisation of the multicellular individual is to be regarded as the consequence of the gradual degeneration of the separate cell-individuals which compose it; APATHY (15).

Adaptations facilitating "masking" in Crabs; Aurivillius (24), and H.ECKER (228).

Migration of birds; DIXON (126).—Speech of Monkeys; GARNER (184). Segmentation: Occurrence of intercalated segments perhaps due to very favourable developmental conditions, to rapid growth causing "slips" in segmentation. Perhaps such cases help to bridge the gulf between Nemerteans and Annelids; Cori (103).

#### I. GENERAL SUBJECTS.

Rudimentary organs; Kocks (298).—Study of rudimentary eyes in Vertebrata; Kohl (304).—Flughaut Bildung; Marshall (342).

Genital appendages of Insects: The intricacies of the genital appendages are not readily interpreted as adaptive. Their final cause is das Princip der Reinerhaltung der Arten; Their efficient cause ein unbekanntes Kraft; ESCHERICH (155).

Venation of wings in Lepidoptera; SPULER (477).

Parallel series in life-histories of Pemphigiida; HORVATH (269).

Asymmetry of Gastropods; Buchner (80), Lang (319).

Parallelism of evolution in Old and New World; GAUDRY (187).

Stages of growth and decline [in development of Brachiopoda]; BEECHER (43).

Evolution of ant-societies; WASMANN (528).

Man's past and future considered according to the doctrine of evolution; BÜCHNER (81).

# II. MAMMALIA.

BY

R. LYDEKKER, B.A., F.G.S.

A FEATURE of the year is the large amount of work which has been devoted to the elucidation of the morphology of the dentition, Ballowitz, Bateson, Freund, Kükenthal, Leche, Osborn, Röse, Scott, Schlosser, Thomas, and Topinard having all contributed memoirs on this subject. The view that Mammals were originally diphyodont is gaining universal acceptation. Attention may be drawn to Scott's investigations as to the homology of the cusps of the premolars, which, if confirmed, show a remarkable difference between those teeth and the molars. Another subject which has received especial attention is that of the brain, and its relations in the higher and lower Mammals; Symington having reverted to the old view that the brain of the non-Placentals has no corpus-callosum. Beddard, Beevor, Chapman, Hill, and Turner have also contributed to this branch.

In distribution, HART MERRIAM'S important memoir on the life-zones of North America is likely to become classical.

Important advances have been made in our knowledge of the Mammals of Africa, several interesting new Antelopes having been described during the year—notably those respectively named by SCLATER and THOMAS. The latter's revision of the genus Cephalophus has cleared up the synonymy of a very difficult group; while TRUE's memoir on the Mammals of East Africa is an important addition to our knowledge of the soology of that region. The revision of the Hyracoidea, by THOMAS, is likewise of great value. The same writer has also contributed papers on the Mammals of the East, recording a new species of Hemigale, as well as several of Tupaia and Sciurus. The large number of new species of Monkeys described is also a feature of the year's work; while the observations of THOMAS on the dentition of Centetes claim special notice. Of high value are Winge's memoir on the Bats of Lagoa Santa, and CAMERON'S observations as to the relations of the antiers of the American Deer to those of the Old World.

Among palsontological work, reference may be made to Earle's memoirs on Coryphodon and Palsosyops, and also to the remarkable horned Artiodactyle—Protoceras, now fully described by Osborn and Wortman. In faunistic palsontology, Boule and Gauder, as well as Depart, have written on the Mammals of France; Rütimeyer has continued his researches on those of Switzerland, and Hofmann on those of Styria; while Cope, Osborn, and Scott have devoted themselves to the North American forms. The first part of the Mammalian portion of Zittel's Palsontologie indicates that when complete the work will be of the highest value to students of both fossil and recent Mammals.

### I.—THE GENERAL SUBJECT.\*

ABBOTT, W. J. L. [See p. 30, Pinnipedia.]

ALLEN, H. On the Foramen Magnum of the Common Porpoise, and on a Human Lower Jaw of unusual size. P. Ac. Philad. 1892, p. 289.

In two specimens of the skull of the Porpoise the basioccipital was seen to be entirely excluded from the condyles by the junction of the exoccipitals below the foramen magnum.

- ---. [See also Chiroptera, p. 25, and Carnivora, p. 28.]
- ALLEN, J. A. On a small Collection of Mammals from the Galapagos Islands, collected by Dr. G. Baur. Bull. Am. Mus. Nat. Hist. iv, pp. 47-50.

Includes a new Atalapha, a new Oryzomys, and the common and black Rats. [See Vespertilionidæ and Muridæ.]

---. The Geographical Distribution of North American Mammals. T. c. pp. 199-243.

Adopts zone-divisions considerably differing from those taken by Merriam in the memoir cited on p. 11; giving the characteristic fauna of each.

Ameghino, F. Répliques aux Critiques du Dr. Burmeister sur quelques Genres de Mammifères Fossiles de la République Argentine. Bol. Ac. Arg. xii, pp. 437-469.

A reply to the criticisms of Burmeister quoted in Zool. Rec. xxviii, Mamm. pp. 5 & 6, in which the author maintains the correctness of his own views. The distinctness of the genera Acrotherium, Adinotherium, and Notohippus, from Nesodon, is insisted upon, and it is urged that the critic has confounded milk with permanent teeth, and that his restoration of the skull of the latter is quite incorrect. The substitution of Pachym-

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

odon in place of Trigodon, &c., is shown to be quite unjustifiable. That the early Megathericids had a thin layer of enamel between the cement and dentine of their teeth, is held to be fully proved; and it is mentioned that this feature harmonizes with the rudiments of an enamel-organ recently discovered among living Edentates. It is further stated that the alleged existence of a plastron among the Glyptodonts is incorrect, that which has been mistaken for such being in reality the head-shield. It is also pointed out that Burmeister has made certain errors in regard to the proper tails belonging to the various genera of this group. Finally, the author mentions that his own name, Pontoplanodes (Zool. Rec. xxviii, Mamm. p. 48), has priority over his critic's Saurodelphis (loc. cit.).

Anderson, J. On a small Collection of Mammals, Reptiles, and Batrachians from Barbary. P. Z. S. 1892, pp. 3-24.

11 species of Mammals are mentioned, among which *Plecotus auritus* is, for the first time, recorded from Tunisia, while its alleged occurrence in Algeria is authenticated.

---. [See also p. 35, Spalacidæ.]

BALLOWITZ, E. [See p. 51, Edentata.]

Bateson, W. On Numerical Variation in Teeth, with a discussion of the Conception of Homology. P. Z. S. 1892, pp. 102-115, figs.

Numerous instances are given of individual variation in the number of teeth of Mammals; especial interest attaching to a skull of Ommatophoca, in which an incipient division of the first premolar on one side of the upper jaw is accompanied on the opposite side with the development of two complete teeth. It is considered that the Owenian theory of the homologies of Mammalian teeth, based on the hypothesis that the typical series is composed of a definite number of teeth, some of which may be absent in particular cases, is untenable; and a new view is propounded as to the homology of multiple parts.

BAUR, G. [See p. 54, Monotremata.]

BEAUREGARD, H. [See p. 25, Chiroptera, and p. 48, Cetacea.]

BEDDARD, F. E. [See p. 22, Primates, p. 32, Rodentia, and p. 36, Octodontidæ.]

BEEVOR, C. E. [See p. 22, Primates.]

Beneden, P. J. v. [See p. 48, Cetacea.]

BLANC, L. Sur un Cas Remarquable de Dédoublement de la Région Tarso-Métatarsienne (Schistomelie). Ann. Soc. Linn. Lyon (2) xxix, pp. 40-51, figs.

Describes the anatomy of the hind foot of a Calf exhibiting a duplication of the metatarsus and pes, with the development of a third digit on one side of the duplicated pes. BLANFORD, W. T. The Age of the Himalayas. Geol. Mag. (3) ix, pp. 161-168.

A list is given of the Mammalian fauna of Tibet, with notes on its evolution and insulation. [See p. 45, Cervidæ, and p. 46, Camelidæ.]

BOCAGE, J. V. B. DU. [See p. 25, Chiroptera.]

BOLAM, G. [See p. 26, Vespertilionidæ.]

Bonnet, B. Grundriss der Entwickelungsgeschichte der Haussäugethiere.
 Berlin: 1891, 282 pp., figs.

BOULE, M. [See p. 6, GAUDRY, and p. 37, Proboscidea.]

BOUVIER, E. L. [See p. 30, Pinnipedia, and p. 48, Cetacea.]

BÜCHNER, E. [See p. 33, Sciurida, and p. 35, Dipodida.]

BUCKLEY, J. E. [See p. 7, HARVIE-BROWN & BUCKLEY.]

Buxton, E. N. Short Stalks; or Hunting Camps, North, South, East, and West. London: 1892, 8vo, 405 pp., illustrated.

A fully illustrated sporting account of many of the species of large game from various parts of the world.

Cacciamali, G. B. Sulla Classificazione dei Mammiferi. Riv. Ital. Sci. Nat. xii, pp. 49-51.

CAMERON, A. G. [See p. 44, Cervidæ.]

CANNIEU, A. [See p. 32, Rodentia.]

CAPELLINI, G. [See p. 48, Cetacea.]

CHAPMAN, F. M. Notes on Birds and Mammals observed near Trinidad, Cuba, with Remarks on the Origin of West Indian Bird Life. Bull. Am. Mus. Nat. Hist. iv, pp. 279-350.

Contains a few notes on the Mammals of S. Cuba, with description of a new subfossil species of *Hutia*. [See p. 36, Octodontidas.]

CHAPMAN, H. [See p. 22, Primates, and p. 36, Octodontides.]

CHIEVITZ, J. H. Sur l'Existence de l'Area Centralis Retinse dans les Quatres Premières Classes des Vertébrés. Overs. Dan. Selsk. 1891, pp. 239-253.

CHRISTY, M. [See p. 36, Leporidæ.]

CIACCIO, G. V. [See p. 40, Equidæ.]

COLLETT, R. [See p. 24, Cercopitheoidæ.]

COOK, J. H. [See p. 29, Ursidæ.]

COPE, E. D. A Contribution to a Knowledge of the Fauna of the Blanco Beds of Texas. P. Ac. Philad. 1892, pp. 226-229.

Several Mammals are recorded from these beds, which are of late Tertiary age; among them being a new Mastodon. [See p. 37, Proboscidea.]

- [Cope, E. D.] On False Elbow-Joints. P. Am. Phil. Soc. xxx, pp. 285-291, pls. ix & x.
- —. Points in the Kinetogenesis of the Limbs of Vertebrates. T. c. pp. 282-285.
- ---. [See also p. 40, Equidæ, and p. 52, Marsupialia.]

CRAGIN, F. W. [See p. 46, Camelida.]

DEPÉRET, C. La Faune de Mammifères Miocénes de la Grive-St-Alban (Isère), et de quelques autres Localités du Bassin du Rhone. Arch. Mus. Lyon, v, art. 2, 96 pp., 4 pls.

An important revision of all the Mammals hitherto described from these formations, with notices of a number of recently discovered specimens and descriptions of new forms. The author recognizes a total of 47 named species from Grive-St-Alban, of which 4 are new, while there are also several new varieties. Especial attention is directed to the *Chalicotheriida*, and it is concluded that the older forms from Sansan and Grive-St-Alban belong to *Macrotherium*, while the Eppelsheim species pertains to *Chalicotherium*. The name *M. grands* is adopted for the species described. It is concluded, on account of the structure of both the skull and the limbs, that the group should be placed among the *Perissodactyla*. The fauna of Mt. Ceindre includes 18 named species, 3 of which are new. [See Simiida, Rhinolophida, Vespertilionida, Soricida, Felida, Viverrida, Mustelida, Chalicotherida, and Suida.]

Les Animaux Pliocènes du Roussillon. Mém. Soc. Géol.-Pal. iii, pp. 117-127, pls. i & ii.

The completion of the Mammalian portion of the memoir quoted in Zool. Rec. xxviii, Mamm. p. 7. The forms described include Felis, Ursus, Lagomys, Hipparion, Cervus, Capreolus, and Dolichopithecus. [See, p. 35, Muridæ and, p. 46, Cervidæ.]

Dollo, L. Sur la Morphologie des Côtes. Bull. Sci. Fr. Belg. xxiv, pp. 1-19.

It is contended, in the first place, that ribs have not an intervertebral origin; and, in the second, that has mapophyses are homologous with one another throughout the length of the vertebral column. The former view is illustrated by the conditions obtaining in the Sirenia.

Dun, W. S. [See p. 52, Marsupialia.]

DUVAL, M. [See p. 32, Rodentia.]

EARLE, C. [See p. 38, Amblypoda, and p. 42, Titanotheriidæ.]

EGLINGTON, W. [See p. 12, NICOLLS.]

FABRINI, W. [See p. 28, Felida.]

FAUVEL, A. A. La Faune du Chan-toung (Chine). Rev. Quest. Sci. (2) i, pp. 455-492.

Contains a notice of the prehistoric and living Mammals.

FILHOL, H. [See p. 27, Macroscelididæ, and p. 30, Creodonta.]

FISERIUS, E. [See p. 32, Rodentia.]

Forbes, H. O. [See p. 30, Pinnipedia.]

FRENKEL, M. Du Tissu Conjonctif dans le Lobule Hépatique de certains Mammifères. C.R. Soc. Biol. 1892, pp. 38 & 39.

FREUND, P. [See p. 33, Rodentia.]

GADOW, H. [See p. 52, Marsupialia.]

GAUDRY, A. Excursion dans l'Amérique du Nord. Bull. Soc. Géol. (3) xix, pp. 936-1024.

Contains notes on American fossil Mammals and their phylogeny.

- ---. [See also p. 34, Cercopithecidæ.]
- ---- & Boule, M. Materiaux pour l'Histoire des Temps Quaternaires, pt. 4, pp. 105-130, pls. xx-xxiv. Paris: 1892, sm. 4to.

Treats of Hyana, Ursus, and Canis lupus.

- Geberg, A. Ueber die Endigung des Gehörnerven in der Schnecke der Saugethiere. Anst. Ans. viii, 20-22.
- GEHUCHTEN, A. v. Les Cellules Nervenser du Sympathique chez quelques Mammifères et chez l'Homme. Cellule, viii, pp. 83-85, pl.

It is concluded that the nervous elements in the sympathetic of Mammals are strictly comparable to those of the cerebro-spinal system. They are provided with two kinds of prolongations—one long and the other short; and always end freely.

GERMAIN, H. [See p. 24, Cebidæ.]

GILIS, P. Anatomie des Scalènes (Costotrachélians) chez les Ruminants, les Solipèdes, et les Carnassiers. C.R. Soc. Biol. 1892, pp. 464–467.

---. [See also p. 33, Rodentia.]

GORJANOVIC-KRAMBERGER, D. O. [See p. 48, Cetacea.]

GREVÉ, C. [See p. 28, Carnivora,]

Hamilton, D. Records of Sport in Southern India. London: 1892, 4to, 284 pp., illustrated,

Contains sporting accounts of many Indian Mammals.

HARTING, J. E. Vermin of the Farm, J. R. Agric. Soc. (3) iii, pp., 205-231 & 463-478, figs.

A popular illustrated account of the various Mammals supposed to be injurious to the English farmer. Contains an account of the present plague of *Microtus* (= Arvicola) agrestis in the south of Scotland.

HARVIE-BROWN, J. A., & BUCKLEY, E. A Vertebrate Fauna of Argyll and the Inner Hebrides. Edinburgh: 1892, 8vo, 252 pp., illustrated. Includes an account of the Mammals.

Henneguy, L. F. Sur la Constitution de l'Endoderm des Mammifères. C.R. Soc. Biol. 1892, pp. 277-279.

HEPBURN, D. [See p. 23, Primates.]

HICKS, H. [See p. 37, Proboscidea.]

HILL, A. The Hippocampus. P. R. Soc. lii, p. 5 (Abstract).

The subject of this paper was to test the deviation of the Hippocampus from the normal type in such Mammals as are devoid of smell, or possess this sense in a very limited degree. It was discovered that the fascia dentata was absent in those forms which possess no olfactory bulb or tract.

----. [See also p. 54, Monotremata.]

Hoernes, R. [See p. 47, Charopotamida.]

HOFMANN, A. Beiträge zur Miocsenen Säugethier-fauna der Steiermark. JB. geol. Reichsanst, xlii, pp. 63-76, pls. ii & iii.

A continuation of the author's previous researches on the Miocene fauna of Styria. The species described are Chalicomys jægeri, Hyænarctus brevirhinus, Trochictis taxodon, Hyomoschus peneckei, and Sorex styriacus. [See, p. 27, Soricidæ, p. 29, Ursidæ, and, p. 46, Tragulidæ.]

Howes, G. B. Notes upon the Shoulder-Girdle of certain Dicynodont Reptiles. J. Anat. Phys. xxvi, pp. 403-405.

It is incidentally mentioned that the coracoid process of the *Eutheria* and *Metatheria* represents the epicoracoid of the Monotremes, and also that the term precoracoid is inapplicable to the latter element.

HOWORTH, Sir H. H. [See p. 37, Proboscidea.]

HUDSON, W. H. The Naturalist in La Plata. London: 1892, 8vo, 388 pp., illustrated.

Contains accounts of the habits of many of the Argentine Mammals, more especially the Huanaco and the Viscacha.

HUTCHINSON, H. N. Extinct Monsters. London: 1892, 8vo, 254 pp., illustrated.

Contains restored figures of several species of extinct Mammals.

JENTINE, F. A. [See p. 34, Murida.]

KOLMANN, -. [See p. 24, Cercopithecidæ.]

Kříž, M. Die Hohlen in den Mährischen Devonkalken und ihre Vorzeit. JB. geol. Reichsanst, xli, pp. 442-570, pls. viii & ix.

An account of explorations in the caverns of northern Moravia, with details of their fauna.

KÜKENTHAL, W. Ueber den Ursprung und die Entwickelung der Säugetierzahne. Jen. Z. Nat. xxvi, pp. 469-489. Translated in Ann. N. H. (6) ix, pp. 279-285.

A recapitulation and further development of the author's views as to the evolution of Mammalian teeth. The theory that the Whales are connected by means of Zeuglodon with the Seals is disputed.

—. Ueber die Enstehung und Entwicklung der Säugetierstammes. Biol. Centralbl. xiii, pp. 400-413. Translated in Ann. N. H. (6) x, pp. 365-380.

The author summarizes his views as to the relationship between Mammals and the lower Vertebrates. He concludes that in the Theromorous or Anomodont Reptiles the original state of the dentition was that of polyphyodontism, while in the marsupial and placental Mammals it was diphyodont. In consequence, however, of the specialization of the individual teeth in the Auomodonts, all dentitions except the first have been suppressed; while in the Marsupials at least one tooth of the second dentition became functional. On the other hand, in spite of the specialization, both series of teeth appear in Placentals. Later on, the author considers that his view as to the formation of complex Mammalian teeth by the fusion of simple denticules is supported by the cheek-teeth of the Multituberculata. In the Vertebrata, as a whole, it appears, however, that the development of the dentition is primarily traceable to the fission of individual teeth. In regard to the relations of Mammals to other groups, it is urged that their ancestors were not Anomodont Reptiles, but primeval forms (from which the Anomodonts themselves may have diverged), living during the Palæozoic period. and having a generalized dentition of simple conical teeth. From these were developed the Multituberculate Mammals; of which the existing Monotremes are considered to be a specialized side-branch. The Marsupials are likewise considered to be another early branch from this same ancient stem, their dentition being traceable to a modification of the Multituberculate type. Finally, it is concluded that Placentals are not descended from Marsupials; the one having the second dentition as the permanent series, while in the other the first set persists.

---. [See also p. 43, Cetacea.]

KULAGIN, N. [See p. 29, Canida.]

LANGKAVEL, B. [See p. 29, Canida.]

LATASTE, F. [See p. 25, Chiroptera, and p. 36, Leporida.]

LECHE, W. Studien über die Entwicklung des Zahnsystems bei den Säugethieren. Morph. JB. xix, pp. 502-547, figs.

An important memoir, still further elaborating the subject of the succession and homologies of the Mammalian dentition. Kükenthal's conclusions are discussed in detail; and the author's own observations on dental development fully illustrated. Taking Erinaceus as a starting-

point, the author has studied the development of the teeth and the relations of the milk and permanent series in the following genera, viz.: Talpa, Sorex, Didelphys, Myrmecobius, Perameles, Trichosurus, Phascolarctus, Tatusia, Dasypus, Balænoptera, Procavia, Felis, and Canis. In discussing the relations between the milk and permanent dentitions, it is shown that the teeth of the latter series are developed on the inner side of those of the former; while, contrary to the view of Kükenthal, reasons are given for considering the milk-dentition to be both autogenetically and phylogenetically the elder of the twain. As the result of all the forms examined, it is further concluded that the molars belong morphologically to the first dentition, i.e., they are milk-teeth without vertical successors. The author takes the opportunity of mentioning that his previous views as to the teeth of Bats are incorrect; it now appearing that the almost completely tritubercular nature of their molars is the result of a kind of degeneration, and is in nowise primitive. It is also urged that the term, second dentition, is not the equivalent of persistent dentition, seeing that in the latter the molars belong to the first series; while, as in Erinaceus, some of the anterior teeth may likewise pertain to the first series. Hence the following stages of development of Mammalian dentition may be distinguished—(1) When the permanent dentition consists solely of teeth of the first series, e.g., Odontoceti. (2) When the persistent dentition is composed of teeth of the first series, with the exception of the last premolar, e.g., Marsupialia. (3) When the permanent dentition comprises molars of the first dentition, with incisors, canines, and premolars belonging partly to the first and partly to the second series, e.g., Erinaceus. (4) When the permanent dentition is formed exclusively, with the exception of the molars, of teeth of the second series, as in all the higher Mammals. Later on it is argued that the monophyodontism of the inferior Mammals is the result of the non-appearance of the second dentition, whilst in the higher types it is produced by the suppression of the first series, that is to say by retrogressive evolution. In opposition to the view that the milkdentition is a recent addition, the author is inclined to admit that it is, on the contrary, the second dentition, which has been but gradually sequired. With regard to the so-called "Zahmoall" and "Zahnfurche," the author arrives at the same results in Erinaceus, as Röse in Man; Zahnwall and Zahnfurche having no relation to the origin or development of teeth. On the contrary, they have merely a relation to the configuration of the buccal cavity during that period of life in which teeth are wanting. To the foregoing ontogenetical researches are added observations on the formation of the dental system in general. Amongst these may be noted the important statement that in all Mammalia where homodontism is found (Odontoceti, Edentata), it is not a primitive character, but the result of retrogressive evolution. Additional highly interesting remarks are given on progressive and retrogressive evolution of the Mammalian dentitions.

LECHE, W. [See also p. 52, Marsupialia.]

- LENHOSSEK, M. v. Die Nervenursprünge und Endigungen im Jacobson'schen Organ des Kaninchens. Anst. Ans. vii, pp. 628-635.
- LYDEKKER, R. Discovery of Australian-like Mammals in South America. Nature, xlvi, pp. 11 & 12.

Notices the discovery in the Tertiary of South America of Marsupials allied to the Thylacine.

- ——. [See also p. 28, Viverridæ, p. 41, Rhinocerotidæ, p. 43, Bovidæ, p. 45, Cervidæ, p. 47, Anoplotheriidæ, p. 49, Cetacea, and p. 51, Sirenia.]
- MACPHERSON, H. A. A Vertebrate Fauna of Lakeland. Edinburgh: 1892, 8vo, 650 pp., illustrated.

Contains notices of the Mammals of Cumberland, Westmoreland, and North Lancashire, and two special articles on the Red Deer and Pine Marten.

MAGGI, D. [See p. 23, Primates.]

Malaquin, A. Sur la Presence de Vertebres dans l'Eocene Inférieur du Nord de la France. Ann. Soc. Géol. Nord. xix, pp. 315-319.

Records a large Coryphodon.

Major, C. J. F. Le Gisement Ossifère de Mitylini. In "Samos," by C. Stefani. Lausanne: 1892, 4to.

The author recapitulates his views as to the Upper Miocene age of the deposits of Samos, Pikermi, and Maragha. An important discussion is given on the affinities of *Chalicotherium*, in which it is suggested that its resemblances to *Palæosyops* do not indicate genetic relationship. It is concluded that the genus is quite distinct from *Macrotherium*, of the Middle Miocene of Sansan, and that the two together represent a distinct order—the *Ancylopoda*. A list is given of the Mammals of Samos.

- MARES, F. Expériences sur l'Hibernation des Mammifères. Mém. Soc. Biol. 1892, pp. 313-320.
- MARSH, O. C. Discovery of Cretaceous Mammals: Part 3. Am. J. Sci. (3) xliii, pp. 249-262, pls. v-xi.

Describes a number of teeth from the Laramie Cretaceous of Wyoming, many of which are referred to new genera and species of *Multituberculata* and *Marsupialia* [q.v.].

MARTIN, H. T. [See p. 34, Castoridæ.]

MARTIN, K. [See p. 37, Proboscidea.]

MAŚKA, K. J. Die Diluviale Fauna und Spuren des Menschen in der Schoschunker Höhle in Mähren. JB. geol. Reichsanst. xli, pp. 415-422, pl. vii.

Records the occurrence of human remains with those of other Vertebrates in a Moravian cavern. The milk-dentition of *Hyana crocuta* (spelaa) is figured.

- MATSCHIE, P. Ueber einige Afrikanische Säugethiere. SB. nat. Fr. 1892, pp. 110-115 & 130-140. [See Felide, Procaviide, Equide, and Bovide.]
- —. Ueber eine kleine Sammlung von Säugethieren und Reptilien, welche Herr L. Conradt aus Usambaru (Deutsch-Ost-Afrika), heimgebracht hat. T. c. pp. 101-110.

A few Micro-Mammalia are mentioned.

—. Ueber die Verbreitung einiger Säugethiere in Afrika. T. c. pp. 223-235.

Chiefly devoted to a criticism of True's paper (infra, p. 19), but also contains a description of a new Cynocephalus [see Cercopithecidæ, p. 24].

——. Ueber einige Neuerwerbungen des Berliner Zool. Gartens. T. c. pp. 220-223.

Mainly deals with certain Felines.

MELI, M. Sopra alcuni Resti di Mammiferi Fossili nei Terreni Quaternari della Provincia di Roma. Boll. Soc. Geol. Ital. x, 5 pp.

Records remains of a few recent and extinct species from the Pleistocene deposits in question.

MERRIAM, C. H. The Geographic Distribution of Life in North America, with special reference to the *Mammalia*. P. Biol. Soc. Washington, vii, pp. 1-64, map.

North America is divided, from north to south, into the Boreal, Transition, Sonoran, and Tropical zones, severally distinguished by a more or less characteristic Mammalian fauna. These zones are proved to coincide very closely with those founded by different writers respectively on the eastern and western sides of the continent, and are now shown to be transcontinental. The Boreal region is divided into an Arctic and a Boreal zone, the latter comprising the area south of the treeless Arctic districts, sweeping across the continent from Newfoundland and the southern shore of Hudson Bay to Alaska; and on its southern border giving off three prolongations running into the United States, down the Alleghanies, the Rocky Mountains, and the Cascade and Sierra Nevada ranges. The Transitional zone, as its name implies, is a kind of neutral area between the Boreal and the Sonoran. It is strongly urged that the Boreal zone and the northern portion of the Palmarctic region really form one soological region, but that from the latter, as ordinarily understood, should be removed an area corresponding with that of the Sonoran zone. Full lists are given of the Mammalian fauna of the several zones, among which are some new subgeneric names, together with certain amendments on some generic terms generally in use. It is considered that the Glacial period has played a very important part in the present distribution of the North American fauna; it being even urged that the S. American Llamas were originally northern forms driven as far south as the latitude of the Andes by the cold. The paper concludes with a criticism of some of Wallace's views, in which it is considered that he has not allowed sufficient importance to the effects of climate as a factor in distribution. [See, p. 33, Sciuridæ, p. 34, Muridæ, and, p. 35, Geomyidæ.]

[MERRIAM, C. H.] Descriptions of Nine New Mammals collected by E. W. Nelson, in the States of Colima and Jalisco, Mexico. T. c. pp. 164-174.

---. [See p. 27, Soricidæ, p. 34, Muridæ, and p. 35, Geomyidæ.]

MEYER, A. B. [See p. 25, Cercopithecidæ.]

MÖBIUS, K. [See p. 37, Proboscidea.]

MOEWES, F. [See p. 44, Giraffida.]

MORGAN, C. L. Factors in the Evolution of the Mammalia. Nat. Sci. i, pp. 97-101.

Criticizes some of Scott's views on this subject.

Moreno, F. P. [See p. 49, Cetacea.]

MUMMERY, J. H. Some Points in the Structure and Development of Dentine. Phil. Tr. clxxxii B, pp. 527-545, pls. xxxvi-xxxix.

The full text of the memoir quoted in Zool. Rec. xxviii, Mamm. p. 13.

NACHTRIEB, H. F. [See p. 24, Lemuroidea.]

Nehring, A., Verh. Anthrop. Ges. 1892, p. 86, records 14 species of fossil Mammals from a cave near Schaffhausen; all but one being small forms.

---. [See also p. 45, Cervidæ, and p. 48, Suidæ.]

NEWTON, E. T. [See p. 34, Castoridæ.]

NICOLLS, J. A., & EGLINGTON, W. The Sportsman in South Africa. London: 1892, 8vo, pp. vi & 147, pls.

An illustrated descriptive work on all the large game of the districts lying to the south of the Zambesi [see, p. 44, Bovidæ].

OGILBY, J. D. Catalogue of Australian Mammals, with Introductory Notes on General Mammology. Sydney: 1892, 8vo, 142 pp., figs.

The completion of the work, of which an advanced publication of the first portion is noticed in Zool. Rec. xxviii, Mamm. p. 14. The list includes both terrestrial and aquatic forms; and the author adopts the view of the indigenous origin of the Dingo. [See p. 35, Muridæ.]

Osborn, H. F. The History and Homologies of the Mammalian Molar Cusps. Anat. Ans. vii, pp. 740-747.

In the course of this paper the author emphasizes his views that the primitive type of Mammalian molar was a simple cone; and contests the theory that the molars of the *Multituberculata* are due to the fusion of originally separate teeth.

---.. [See also p. 31, Creodonta, p. 37, Amblypoda, p. 39, Condylarthra and Perissodactyla, p. 40, Lophiodontidæ, and p. 46, Protoceratidæ.]

OSBORN, H. F., & WORTMAN, J. L. Fossil Mammals of the Wahsatch and Wind River Beds. Bull. Am. Mus. Nat. Hist. iv, pp. 81-148, figs., pl. iv.

A very important paper, recording the results of working out a collection obtained in 1891. The Palscontological portion is divided into the four following sections, of which the third is by Wortman, while the others are by Osborn:—viz., (1) Homologies and Nomenclature of the Mammalian Molar Cusps; (2) the Classification of the Perissodactyla; (3) the Ancestry of the Felidæ; (4) Taxonomy and Morphology of the Primates, Creodonts, and Ungulates. In the first part of the paper attention is principally directed to the subsidiary cusps of the Mammalian molar, for which the general term style is suggested. [See Creodonta, Perissodactyla, and Amblypoda.]

PALMER, T. S. [See p. 30, Otariidæ.]

PAVLOW, M. [See p. 41, Rhinocerotidæ.]

Petrini, —. [See p. 28, Carnivora.]

POHLIG, H. [See p. 37, Proboscidea, and p. 45, Cervidæ.]

POLAND, H. Fur-bearing Animals in Nature and in Commerce. London: 1892, 8vo, pp. lvi & 392, illustrated.

Contains tables of the number of pelts annually obtained from the various species of fur-bearing animals, with notices of the animals themselves. Reviewed in Nat. Science, i, pp. 624-626.

POMEL, A. [See p. 33, Sciuridæ, p. 35, Muridæ, and p. 44, Giraffidæ.]

POMMEROL, F. [See p. 40, Equidæ.]

POUCHET, G. [See p. 49, Cetacea.]

Pousargues, E. De. [See p. 33, Rodentia, and p. 45, Cervide.]

QUELCH, J. J. [See p. 25, Chiroptera.]

BEGNAULT, F. (1) Fouilles dans le Terrain Miocène Moyen de St. Gaudens (Haute-Garonne)—Le Dryopitheque, (2) Fouilles dans les Grotees de Gargas et de Malarnaud. C.R. Ass. Fr. Sci. xix, pt. 2, pp. 408-411, figs.

Figures a mandible of *Dryopithecus*, and a human lower jaw very similar to the one from Naulette.

RETTERER, E. DE. Sur la Morphologie et l'Evolution de l'Epithelium du Vagin des Mammifères. Mém. Soc. Biol. 1892, pp. 101-107.

ROBINSON, A. Development of Segmentation-Cavity, Archenteron, Germinal Layers, and Amnion of Mammals. Q. J. Micr. Sci. xxxiii, pp. 369-455, 5 pls.

---. [See also p. 33, Rodentia.]

Röse, C. Ueber die Enstehung und Formabänderungen der Menschlichen Molaren. Anat. Anz. vii, pp. 392–421, figs.

The author gives a detailed description of the cheek-teeth of human embryoes, of various races, and indicates their mutual resemblances and differences, and their general bearing on the phylogeny of molars. It is stated that although, in some respects, the immature molars of Europeans resemble those of non-European nations, the reduction in the number of the tubercles found in the former is peculiar to them. It is further shown that this reduction-process tends to reproduce the original tritubercular type; the tubercles which disappear being those which have been superadded to the primitive triangle. It is added that Cope is correct in regarding the human first upper milk-molar as being formed strictly on the tritubercular type. It is shown that in all the upper cheek-teeth, the first cusp to appear is the antero-external, hitherto regarded as the paracone, but which is identified by the author with the protocone.

- —. Ueber die erste Anlage der Zahnleiste beim Menschen. Op. cit. viii, pp. 29-32.
- —. Ueber die schmelzlosen Zahnrudimente des Menschen. Verh. deutsch. odont. Ges. iv, pp. 1-28, figs.

Concludes that the toothlike bodies devoid of enamel frequently found in the front of human jaws are due to caries or other disease, and do not represent missing teeth.

—. Zur Phylogenie des Säugetiergebiss. Biol. Centralbl. xii, pp. 624-638.

A discussion as to the relationship of the teeth of Mammals to those of Reptiles. It is concluded that Placentals, Marsupials, and Menotremes are correlated, but not genetically related groups.

- ---. [See also p. 51, Edentata, and p. 52, Marsupialia.]
- ROTHSCHILD, W. Descriptions of Two New Mammals from New Guinea. P. Z. S. 1892, pp. 545 & 546. [See p. 53, Phalangeridæ and p. 54, Echidnidæ.]
- RÜTIMEYER, L. Die Eocäne Säugethier-welt von Egerkingen. Abh. Schw. pal. Ges. xviii, pp. 1-153, pls. i-viii.

The full memoir, of which a preliminary resumé (Zool. Rec. xxvii, Mamm. p. 13) had been previously published, descriptive of recent additions to our knowledge of the Mammals of the Swiss Siderolites. Many of the new species were mentioned in the preliminary notice, but as they were unaccompanied by any diagnosis could not be recognized. The title-page is dated 1891, but the description of the plates 1892; and from internal evidence (p. 62), it is evident that the memoir was not published until the latter year.

The first of the descriptive portion of the memoir treats of the Ungulates, which are classified as follows: viz.—

- A. Ungulata Trigonodontia (? = Condylarthra).
- B. Ungulata Zygodontia.
  - 1. Imparidigitata.
    - a. Palæotherida.
    - b. Lophiodontida.
  - II. Paridigitata.
    - a. Suina.
    - b. Selenodontia (Hyopotamus, Bachitherium, etc.).

In the Trigonodonts it is argued that less importance should be assigned to the structure of the carpus and tarsus than to the trituberculism of the teeth. In the Palæotheridæ the genus Paloplotherium is maintained, and a new species described; while in the Lophiodontidæ there is likewise a new species of the type genus. Pachynolophus is divided into several subgenera, of which there is one new species; the name Hyracotherium siderolithicum being still maintained for a member of this group, although its marked distinction from the typical Hyracotherium is mentioned.

Among the Artiodactyles, Tapinodon gresslyi is identified with Hyopotamus; the skull from Hordwell referred by the Recorder to the former under the name of Anthracotherium gresslyi being regarded as indicating a new form. Hyopotamus picteti, Lyd., is also considered to be identical with H. gresslyi, although identified by its founder with Dacrytherium [see p. 47]. Several new species of Dichobunus and Xiphodontherium are named; and Tetraselenodon and Bachitherium are recorded from these beds, the former being placed near Dichodon, of which a new species is described.

Among the Rodents a new *Plesiarctomys* is described, and certain Insectivores are identified with some from Quercy. In the Creodonts a lower molar, described as *Ailuravus*, is referred to the *Arctocyonida*, with a suggestion of affinity to *Elurus*. New species of *Hyanodon* and *Pterodon* are described, as well as a new genus of *Proviverrida*, approximating to the living African *Suricata* (*Rhyzana*).

In the Lemuroids, Adapis and Cænopithecus are compared, and new species of Pelycodus, Hyopsodus, and Necrolemur described.

The memoir concludes with a discussion of the homologies and nomenclature of the elements of the Mammalian molar, and a general survey of the fauna of the Siderolites and its relationships.

[See p. 24, Lemuroidea, p. 31, Proviverridæ and Arctocyonidæ, p. 32, Hyænodontidæ, p. 33, Sciuridæ, p. 40, Lophiodontidæ, p. 41, Palæotheriidæ, p. 47, Dichodontidæ, Anoplotheriidæ, Cænotheriidæ, and Anthracotheriidæ.]

RÜTIMEYER, L. Die Eocanen Saugethiere von Egerkingen. Verh. Ges. Basel. x (Festschrift), pp. 101-129.

The first part of this paper deals with the author's earlier works in homologizing the cusps of Mammalian molars; his own nomenclature

being contrasted with that subsequently published in the United States. The second portion has some general observations on the fauna of Egerkingen.

RUFFINI, H. [See p. 28, Carnivora.]

Ruge, G. [See p. 23, Primates.]

SAFFORD, J. M. The Pelvis of a *Megalonyx* and other Bones from Big-Bone Cave. Bull. Geol. Soc. Amer. iii, pp. 121-123.

SCHACHT, H. [See p. 28, Carnivora.]

SCHÄFF, E. [See p. 29, Canidæ.]

SCHELER, G. [See p. 45, Cervidæ.]

Schlosser, M. Ueber die Deutung des Milchgebisses der Säugethiere. Verh. deutsch. odont. Ges. iv, 10 pp.

A recapitulation and discussion of recent views as to the homology and development of the milk dentition. In the course of the paper it is stated that living Elephants have no deciduous incisors.

—. Ueber die systematische Stellung der Gattungen Plesiadapis, Protoadapis, Pleuraspidotherium, and Orthaspidotherium. JB. Mineral. 1892, ii, pp. 238-240.

It is concluded that the first two of the genera named, which had been previously assigned to the Lemuroids, are more probably Rodents, although having a fuller dentition than any existing member of the order. On the other hand the genera *Pleuraspidotherium* and *Orthaspidotherium* are considered to represent Condylarthrous Ungulates, probably allied to the *Perissodactyla*. They had been previously referred to the *Insectivora*.

SCHÖNLAND, S. [See p. 41, Equidæ.]

Sclater, P. L. On a small Collection of Mammals brought by Mr. A. Sharpe from Nyassaland. P. Z. S. 1892, pp. 97 & 98.

8 species are noticed, and among these the range of Tragelaphus angasi is extended more to the north than was previously known.

- [See also p. 24, Cercopithecidæ, and p. 43, Bovidæ.]

Scott, O. B. The Evolution of the Premolar Teeth in the Mammals. P. Ac. Philad. 1892, pp. 405-440.

In the first place the author refers to Fleischmann's objections to the view of the development of the more complex types of Mammalian molars from a simple cone, and to the homology between the cusps of those of the upper and lower jaws. It is considered that these objections cannot be sustained, more especially as the author believes that the Marsupials are not the ancestors of the Placentals. The author then proceeds to trace, in detail, the origin of the more complex types of premolars. Even when these teeth have assumed a completely molariform structure, their history is quite different from that of the molars, the elements which correspond in function and position to those of the latter not being

homologous therewith. The key to the homologies is afforded by the position of the protocone. As the premolars do not present quite the same degree of developmental constancy as the molars, the last of the series is adapted as the standard. The various cusps of the premolars are then contrasted with those of the molars, and new names proposed where necessary. It is concluded that while the inferior premolars may contain three elements which are homologous with molar cusps, the upper premolars contain but one cusp which can be homologized with a molar cusp; and, furthermore, that while in the molar crown the protocone is on the inner, and the protoconid on the outer side, in both sets of premolars, upper as well as lower, it retains its primitive position on the outer side. The following table exhibits the correspondence of position (not of homology) between the molar and premolar elements, when the whole are developed:—

### UPPER JAW.

011211 01211		
MOLARS.		PREMOLARS.
Protocone	=	Deuterocone.
Paracone	=	Protocone.
Metacone	=	Tritocone.
Hypocone	=	Teturtocone.

#### LOWER JAW.

Protoconid	=	Protoconid.
Paraconid		Paraconid.
Metaconid	=	Deuteroconid.
Hypoconid	=	Metaconid.
Entoconid	_	Tetartoconid.

The author next proceeds to point out the peculiarities of the premolar cusps in several of the orders. In an addendum it is mentioned that if Röse's views as to the position of the protocone and paracone in the molars (supra, p. 14) prove to be well-founded, then the cusps of the premolars and molars will be homologous, and the new terms proposed unnecessary.

### ---. [See also p. 31, Creodonta.]

SNELL, O. Das Gewicht des Gehirnes und des Hirnmantels der Säugethiere in Beziehung zu deren geistigen Fahigkeiten. SB. Ges. Morph. vii, pp. 90-94.

Southwell, T. [See p. 50, Cetacea.]

SPENCER, W. B. The Fauna and Zoological Relationships of Tasmania. Rep. Austr. Assoc. 1892; Address to Sect. D, 44 pp.

STEFANESCU, G. [See p. 37, Proboscidea.]

STEPHENS, F. [See p. 33, Sciurida.]

1892. [VOL. XXIX.]

STRUTHERS, J. [See p. 23, Primates.]

Sussdorf, J. [See p. 48, Suidæ.]

SWAYNE, H. G. C. [See p. 43, Bovidæ.]

TAEKER, J. [See p. 36, Ungulata.]

THOMAS, O. Notes on Dr. W. Kükenthal's Discoveries in Mammalian Dentition. Ann. N. H. (6) ix, pp. 308-311.

In consequence of Kükenthal's recent investigations, the author is now disposed to admit that Mammals may have been originally diphyodont; although, at the same time, this view renders it very difficult to account for the condition obtaining in the Mesozoic Triconodon. It is contended that Kükenthal is incorrect in homologizing the changing premolar of the Marsupials with the third of the typical series, and that it really corresponds with the fourth of the same. After some discussion as to the question whether the premolars or milk-molars are homologous with the true molars, it is argued that Kükenthal's theory of the fusion of the teeth of the permanent and milk sets to form the molars cannot be sustained; but his view of the fission of multicuspid teeth to form the unicuspid ones of the Cetacea is supported.

- ---. On Mammals from Nyassaland. P. Z. S. 1892, pp. 546-554.
- Describes 27 species of Mammals sent home by Mr. H. H. Johnston, which include 3 new varieties, among them the Lesser Pala. *Isomys* is regarded as distinct from *Mus*. [See p. 34, *Muridæ*, and p. 43, *Bovidæ*.]
- —. On some Mammals from Mount Dulit in North Borneo. T. c. pp. 221-227, pls. xviii & xix.
- 21 species are recognized, among which are the new Hemigale hosei, Tupaia montana, T. melanura, and Sciurus brookei. [See p. 27, Tupaiidæ, p. 29, Viverridæ, and p. 33, Sciuridæ.]
- —. On some New Mammalia from the East Indian Archipelago. Ann. N. H. (6) ix, pp. 250-254.

A preliminary notice of the species described in the preceding paper, together with that of a new *Tragulus*. [See p. 46, *Tragulida*.]

—. On the Mammalia collected by Signor Leonardo Fea in Burma and Tenasserim. Ann. Mus. Genov. (2) x, pp. 913-949, pls. x & xi.

The list includes 114 species, all of which have been previously described. The name Linsang is adopted in lieu of Prionodon, and Cervulus few is figured. Orcella fluminalis is identified with O. brevirostris.

- On the probable Identity of certain Specimens, formerly in the Lidth de Jeude Collection, and now in the British Museum, with those figured by Albert Seba in his "Thesaurus," in 1734. P. Z. S. 1892, pp. 309-318.

TOPINARD, B. [See p. 23, Primates.]

TROUESSART, E. L. [See p. 41, Equida.]

TRUE, F. W. An Annotated Catalogue of the Mammals collected by Dr. W. L. Abbott in the Kilima-njaro Region, East Africa. P. U. S. Nat. Mus. xv, pp. 445–480, pls. lxxv-lxxx.

Contains a complete list of all the Mammals from the district, with descriptions of three new species, and figures of others recently named. [See p. 33, Sciuridæ, and p. 34, Muridæ.]

Techerski, J. D. Wissenschaftliche Resultate der v. d. k. Ak. Wiss. zur Eforschung des Janalandes und der Neusibirischen Inseln in den Jahren 1885 und 1886 ausgesandten Expedition. Pt. 4. Beschreibung der Sammlung Posttertiärer Säugethiere. Mém. Ac. Petersb. xl, pp. 1-511, pls. i & vi.

The species described include Felis tigris, Canis lupus, C. familiaris, Vulpes lagopus, Gulo luscus, Ursus maritimus, U. arctos, Phoca fatida, Trichechus rosmarus, Spermophilus eversmanni, Lemmus obensis, Cuniculus torquatus, Lepus variabilis, Bison priscus, Ovibos moschatus, Ovis nivicola, Colus saiga [= Saiga tartarica], Alces palmatus, Rangifer tarandus, Cervus canadensis, v. maral, Equus caballus, and Rhinoceros tichorhinus. A skeleton of the recent Equus przewalskii is also described; and there are notices of the distribution of certain groups, of which the fossilized remains occur in the new Siberian islands. The occurrence of remains of the tiger at such a distance within the Arctic Circle is noteworthy.

TUCKERMANN, F. [See p. 23, Primates.]

TURNER, SIR W. [See p. 50, Cetacea, and p. 54, Monotremata.]

VAUGHAN, J. [See p. 41, Equidæ.]

WALDEYER, W. [See p. 51, Sirenia.]

WARD, B. Horn-Measurements and Weights of the Great Game of the World. London: 1892, 4to, 264 pp., illustrated.

—. [See also p. 43, Bovidæ.]

WHITE, T. [See p. 29, Canida.]

WINGE, H. [See p. 25, Chiroptera.]

WITCHELL, C. A., & STRUGNELL, O. B. The Fauna and Flora of Gloucestershire. Stroud: 1892, 8vo, 301 pp., illustrated.

Contains an account of the Mammals of the district.

WOODWARD, M. F. [See p. 38, Hyracoidea.]

WORTMAN, J. L. [See Osborn, and p. 46, Proteroceratide.]

ZIETZ, A. [See p. 53, Marsupialia.]

#### II. MAMMALIA.

ZITTEL, K. A. Handbuch der Palæontologie. 1st Abhtheil.—Palæozoologie, band iv, lief. 1. Munich: 1892, 8vo, 304 pp., illustrations.

This part of the work commences the Mammals, and includes the orders Monotremata, Marsupialia, Edentata, Cetacea, Sirenia, and the Perissodactyle Ungulates. The class is subdivided as follows:—

Subclass A. EPLACENTALIA.

Order 1. Monotremata.

- 2. Allotheria (Multituberculata).
- 3. Marsupialia.

Suborder a. Polyprotodontia.

b. Diprotodontia.

Subclass B. PLACENTALIA.

Order 4. Edentata.

5. Cetacea.

Suborder a. Archæoceti.

- b. Odontoceti.
- c. Mystacoceti.

Order 6. Sirenia.

7. Ungulata.

Suborder a. Condylarthra.

- b. Perissodactyla.
- c. Artiodactyla.
- d. Amblypoda.
- e. Proboscidea.
- f. Toxodontia.
- g. Typotheria.
- h. Hyracoidea.

Order 8. Tillodontia.

- 9. Rodentia.
- 10. Insectivora.
- 11. Chiroptera.
- 12. Carnivora.

Suborder a. Creodonta.

- b. Fissipedia.
- c. Pinnipedia.

Order 13. Primates.

Suborder a. Prosimiæ.

- b. Simiæ.
- c. Bimana.

The Polyprotodont Secondary Mammals are classed among the Marsupials; while the existing Myrmecobius is made the type of a separate family.

### II.—FAUNAS.

#### A .- RECENT.

Britain. See HARVIE-BROWN & BUCKLEY, p. 7; MACPHERSON, p. 10; and WITCHELL & STRUGNELL, p. 19.

Asia. See Blanford (Tibet), p. 4; FAUVEL (China), p. 6; THOMAS (Malayana and Burma), p. 18.

Africa. See Anderson, p. 3; Matschie, p. 11; Nicolis & Eglington, p. 12; Sclater, p. 43; Thomas, p. 18; True, p. 19.

America. See Allen (Galapagos), p. 2; Chapman (Cuba), p. 4; LATASTE (Chilian Bats), p. 25; MERRIAN, p. 11; QUELCH (Guiana Bats), p. 25; WINGE (Brazilian Bats), p. 25.

Australia. See OGILBY, p. 12; SPENCER, p. 17; ZIETZ, p. 19.

#### B.—Fossil.

Europe. See Depéret (France), p. 5; Gaudry & Boule (France), p. 6; Hofmann (Styria), p. 7; Maška (Moravia), p. 10; Major (Pikermi and Samos), p. 10; Meli (Italy), p. 11; Nehring (Germany), p. 12; Rütimeyer (Switzerland), p. 14.

Asia. See TSCHERSKI (Siberia), p. 19.

N. America. See Cope, p. 4; MARSH, p. 10; OSBORN & WORTMAN, p. 13; Scott, p. 16.

# III.—SPECIAL STRUCTURES, DEVELOPMENT, Erc.

CIRCULATORY SYSTEM .- See BEAUREGARD, Chiroptera, p. 25.

DENTITION.—See Allen, Chiroptera, p. 25; Ballowitz, Edentata, p. 51; Bateson, p. 3; Freund, Rodentia, p. 33; Kükenthal, p. 49; Leche, p. 8; Lydekker, Sirenia, p. 31; Osborn, p. 12; Röse, Edentata, p. 51; and Marsupialia, p. 52; Scott, p. 16; Schlosser, p. 17; Taeker, Ungulata, p. 36; Thomas, p. 18, and Centetidæ, p. 27; Topinard, Primates, p. 23; Woodward, Hyracoidea, p. 38.

DEVELOPMENT, EMBRYOLOGY, AND REPRODUCTION.—See CANNIEU, Rodentia, p. 32 (penis); DUVAL, Rodentia, p. 32 (placenta); FISERIUS, Rodentia, p. 32; FLEISCHMANN, Rodentia, p. 33 (placenta); POUSARGUES, Rodentia, p. 33 (penis); ROBINSON, p. 13 (germinal layers), and Rodentia, p. 33 (volk-sac and placenta).

DIGESTIVE SYSTEM.—See CARLIER, Erinaceidæ, p. 27 (histology); TURNER, Cetacea, p. 50; WALDEYER, Sirenia, p. 51.

EAR.—See GEBERG, p. 6.

EYE .- See CHIEVITZ, p. 4.

GLANDS.—See PETRINI, Carnivora, p. 28; TUCKERMANN, Primates, p. 23. HISTOLOGY. — See CARLIER, Erinaceidæ, p. 27 (digestive system) FRENKEL, p. 6 (hepatic connective tissue); HENNEGUY, p. 7 (endoderm); MUMMERY, p. 12 (dentine); RETTERER, p. 13 (epithelium).

MORPHOLOGY.—See CAMERON, p. 44 (antlers); DOLLO, p. 5 (ribs); HEPBURN, *Primates*, p. 23 (grooves in hand and foot); Howes, p. 7 (shoulder-girdle); KÜKENTHAL, p. 8 (teeth); OSBORN, p. 12 (teeth); RÖSE, p. 14 (teeth); SCOTT, p. 16 (teeth); SCHLOSSER, p. 16 (teeth); THOMAS, p. 18 (teeth); TOPINARD, *Primates*, p. 23 (teeth).

Muscles.—See Allen, Carnivora, p. 28; Beddard, Primates, infrå, and Octodontidæ, p. 36; Gilis, p. 6, and Rodentia, p. 33; Hepburn, Primates, p. 23; Ruge, Primates, p. 23.

NERVES AND BRAIN.—See BEDDARD, Rodentia, p. 32, and Octodontidæ, p. 36; BEEVOR, Primates, infrà; CHAPMAN, Primates, infrà; GEBERG, p. 6; GEHUCHTEN, p. 6; HEPBURN, Primates, p. 23; HILL, p. 7, and Monotremata, p. 54; LENHOSSÉK, p. 10; RUFFINI, Carnivora, p. 28; SYMINGTON, Marsupialia, p. 53; TURNER, Monotremata, p. 54.

Nose.—See Lenhossék, p. 10 (Jacobson's organ); Symington, Marsupialia, p. 53 (Jacobson's organ).

OSTEOLOGY.—See ALLEN, p. 2, and Carnivora, p. 28; BAUR, Monotremata, p. 54; COPE, p. 5; HOWES, p. 7; MAGGI, Primates, p. 23; STRUTHERS, Primates, p. 23.

### IV.—SPECIAL WORK.

(Extinct groups and species are indicated by a †.)

### 1. PRIMATES.

BEDDARD, F. E. Contributions to the Anatomy of the Anthropoid Apes. P. Z. S. 1892, pp. 118-120 (Abstract).

Compares the myology of Anthropopithecus [Troglodytes] calvus with A. niger; and also shows that an Ape recently exhibited in the Zoological Society's Gardens, and believed to be an adult example of Simia morio, was immature.

Beevor, C. E. On the Course of the Fibres of the Cingulum and the Posterior Parts of the Corpus Callosum and Fornix in the Marmoset Monkey. Phil. Tr. clxxxii B, pp. 135-199, pls. xx-xxiv.

The full text of the memoir quoted in Zool. Rec. xxviii, Mamm. p. 22.

CHAPMAN, H. Observations upon the Brain of the Gorilla. P. Ac. Philad. 1892, pp. 203-212, pls. xi & xii.

The author points out the features in which the Gorilla's brain differs from those of other Anthropoids, and suggests that the Gorilla may be descended from *Cynocephalus*; the other forms having totally distinct origins.

HEPBURN, D. Comparative Anatomy of the Muscles and Nerves of the Superior and Inferior Extremities of the Anthropoid Apes. J. Anat. Phys. (2) vi, pp. 149-186 & 324-356, pls. iii & ix.

An elaborate comparison of these structures in a large number of the Primates.

—. The Integumentary Grooves on the Palm of the Hand and Sole of the Foot of Man and the Anthropoid Apes. Op. cit. xxvii, pp. 112– 130, pls. vii-ix.

It is found that both hand and foot bear external evidence of the muscular homologies existing between them; and as these homologies are almost identical in man and apes, so are the lines produced in the skin by muscles. Modifications in the size of muscles have produced some variation in the direction of the lines, but not sufficient to obscure the general adherence to one plan. While the human hand has developed into a better grasping organ than that of an ape, and the foot into a more secure organ of support, both retain evidence of their fundamental unity of structure, and both are capable of performing the same kinds of movements, subject to certain modifications suited to the necessities of each case.

- MAGGI, L. La Sutura Endomesognatica alla Superficie Facciale degli Intermascellari nel Semnopithecus entellus. Rend. Ist. Lombardo (2) xxv, pp. 89 & 90, pl. i.
- Ruge, G. Zeugnisse für die metamere Verkurzung des Rumpfes bei Säugethieren; Der Musculus Rectus Thoraco-abdominalis der Primaten. Morph. JB. xix, pp. 376-427, pls. xii & xiv.

An elaborate study of the relations of this muscle in a number of Primates, with an account of its morphological bearings.

- STRUTHERS, J. The Articular Processes of the Vertebræ of the Gorilla compared with those in Man, and on Costo-Vertebral Variation in the Gorilla. J. Anat. Phys. xxvii, pp. 131-138.
- TOPINARD, P. De l'Évolution des Molaires et des Prémolaires chez les Primates, et en particalier chez l'Homme. L'Anthrop. iii, pp. 641-710, figs.

An elaborate dissertation on the formation and homologies of the cusps and roots of the cheek-teeth of the *Primates*, with observations on the relation of these teeth to the tritubercular type. The author is of opinion that the theory of the direct descent of Man from Lemuroids cannot be sustained; and he inclines to the view that the *Hominida* should be separated from the other Anthropoids to constitute a distinct suborder.

Tuckerman, F. The Guetatory Organs of Ateles ater. J. Anat. Phys. xxvi, pp. 391-393.

#### A.—ANTHROPOIDEA.

#### a. Similda.

- †Dryopithecus: note on its femur; Pohlig, SB. niederrhein. Ges. 1892, pp. 41-43.
- Hylobates hainanus, n. sp., Thomas, Ann. N. H. (6) ix, pp. 85 & 86, Hainan. Distinguished by absence of light frontal band.
- † Pliopithecus antiquus, n. var. chantrei, DEPÉRET, Arch. Mus. Lyon, v, p. 9, pl. ii, fig. 14, Mt. Ceindre.

# b. CERCOPITHECIDE.

COLLETT, R. On a New Monkey from North-East Sumatra. P. Z. S. 1892, pp. 613-617, pl. xlii.

GAUDRY, C.R. cxiv, pp. 1236 & 1237, records the occurrence of fragmentary remains of a Monkey from a cavern in the Haute-Garonne.

Macacus cynomolgus: embryo described by Kolmann, Anat. Auz. vii, pp. 335-340.

† Macacus trarensis, n. sp., Pomel, C.R. cxiv, pp. 157-160, Pleistocene, Algeria.

Semnopithecus everetti, n. sp., Thomas, P. Z. S. 1892, p. 582, N. Borneo.

Semnopithecus cruciger, n. sp., Thomas, Ann. N. H. (6) x, pp. 475-477, Borneo.

Semnopithecus pyrrhus, its distinctness from S. maurus; JENTINK, Notes Leyd. Mus. xiv, pp. 119-121, pls. iii & iv.

Semnopithecus thomasi, n. sp., Collett, op. cit., Sumatra.

Semnopithecus nemæus, recorded by MEYER, P. Z. S. 1892, p. 665, from Hainan.

Cercopithecus schmidti, n. sp., MATSCHIE, Zool. Anz. xv, p. 161, Uganda. Cercopithecus stairsi, n. sp., Sclater, P. Z. S. 1892, p. 580, pl. xl, Lower Zambeei

Cynocephalus langheldi, n. sp., MATSCHIE, SB. nat. Fr. 1892, p. 232, Ungaru, E. Africa.

### c. CEBIDÆ.

GERMAIN, H. Les Sapajous. Act. Soc. Chili, i, pp. 105-107 Notes on the habits of these Monkeys.

#### B.—LEMUROIDEA.

NACHTRIEB, H. F. A New Lemur. Zool. Anz. xv, pp. 147 & 148.

A preliminary notice of a Lemur from the Philippines regarded as indicating a new species.

Lemur menagensis, n. sp., NACHTRIEB, loc. cit.

†Necrolemur cartieri, n. sp., L. RÜTIMEYER, Abh. Schw. pal. Ges. xviii, p. 114, Up. Eccene, Switzerland.

†Pelycodus helveticus, n. sp., RÜTIMEYER, t. c. p. 117, ibid.

†Hyopsodus jurensis, n. sp., RÜTIMEYER, t. c. p. 121, ibid.

### 2. CHIROPTERA.

ALLEN, H. On a New Subfamily of Phyllostome Bats. P. U. S. Nat. Mus. xv, pp. 437-439.

It is concluded that Natalus should be removed from the Vespertilionida to form a subfamily of Phyllostomatida.

—. On the Molars of the Pteropine Bats. P. Ac. Philad. 1892, pp. 172 & 173.

The author believes that he is able to identify the constituent cusps of these teeth with those of the quadritubercular type, in which case they cease to be aberrant.

Beauregard, H. Le Canal Carotidien des Roussettes. C.R. Soc. Biol. 1892, pp. 914-916.

BOCAGE, J. V. B. DU. Observations sur les Espèces du Genre Cynonycteris rencontrées en Angola par M. d'Anchieta. J. Sci. Lisb. (2) vii, pp. 173-178, figs.

It is concluded that the species represented are probably C. straminea, C. agyptiaca, and C. colluris.

LATASTE, F. Études sur la Faune Chilienne—Note sur les Chauves-Souris. Act. Soc. Chili, i, pp. 70-91.

The list includes 1 Desmodus, 1 Sturnira, 1 Nyctinomus, 3 Vespertilio, 2 Atalapha, and 4 Vesperugo.

Quelch, J. J. The Bats of British Guiana. Timehri (2) vi, pp. 90-109.

Notes on the habits of the various species. It is stated that there are blood-sucking Bats in districts where Desmodus is unknown.

WINGE, H. Jordfundne og nulevende Flagermus (Chiroptera) fra Lagoa Santa, Minas Geraes, Brasilien; nach Udsigt over Flagermusenes indtyrdes Scægtskab. E. Mus. Lund. vol. iii, pp. 1-65, pls. i & ii.

An important memoir on the Bats of Lagoa Santa, uniform with the one on the Rodents noticed in Zool. Rec. xxiv, Mamm. p. 32, but, unfortunately, in Danish. As in that case, the recent and fossil forms are taken together, and figures of the skulls and dentition of the majority given. No species are new; but a full account is given of the forgotten Phyllostoma humerale, Lund, now referred to Stenoderma. Numerous species, described more or less imperfectly by the same author, are assigned to their proper genera. It is proposed to alter the names of the Bats termed by Dobson Lonchoglossa wiedi and Molossus temmincki into

L. ecaudata and L. histipes respectively; but both these changes are inadmissible, according to the rules of nomenclature. Two appendices follow the systematic portion, the first containing an account of the author's investigations on the myology of Bats and his proposed classification, while the second deals with the work of other writers. This part appears to contain a number of valuable observations, and also comprises a complete series of dental formulæ, given in such a manner as to indicate what the author considers to be the homologies of the individual teeth throughout the order.

### a. PTEROPODIDÆ.

Cynopterus blanfordi, n. sp., THOMAS, Ann. Mus. Genov. (2) x, p. 884 (1891), Karin Hills, Burma. See also t. c. p. 921, pl. xi.

#### b. RHINOLOPHIDE.

- Rhinolophus ferrum-equinum: DE KERVILLE, Le Nat. 1891, p. 239, describes and photographs a colony hibernating in a cave in the department of Eure.
- †Rhinolophus lugdunensis et collongengis, n. spp., DEPÉRET, Arch. Mus. Lyon, v, pp. 13 & 16, pl. ii, figs. 4-9, Miocene, Mt. Ceindre.

#### c. VESPERTILIONIDÆ.

- BOLAM, G. Daubenton's Bat in Northumberland, with remarks upon other Species recorded from the district. Naturalist, 1892, pp. 269-272.
- †Vespertilio grivensis, n. sp., DEPÉRET, Arch. Mus. Lyon, v, p. 11, pl. ii, figs. 2 & 3, Miocene, Grive-St-Alban.
- Rhogeëssa alleni, n. sp., Thomas, Ann. N. H. (6) x, pp. 477 & 478, Mexico.
- Scotophilus temmincki, regarded by H. Allen, P. U. S. Nat. Mus. xv, pp. 442 & 443, as nearly allied to Atalapha and Dasypterus.
- Atalapha brachyotis, n. sp., J. A. Allen, Bull. Am. Mus. Nat. Hist. iv, pp. 47 & 48, Galapagos I.
- Nyctophilus walkeri, n. sp., Thomas, Ann. N. H. (6) x, pp. 405 & 406, Adelaide R., Australia.
- Harpiocephalus fex, n. sp., Thomas, Ann. Mus. Genov. (2) x, p. 884 (1891), Karin Hills, Burma. See also t. c. p. 926.
- Natalus, Gray, amended by Winge (suprà, p. 25), p. 38, to Natalis; transferred by Allen, suprà, to Phyllostomatidæ.

### d. Phyllostomatidæ.

Chilonycteris davyi, n. var. fulvus, THOMAS, Ann. N. H. (6) p. 410, Mexico.

Ectophylla alba, n. g. & sp., H. Allen, P. U. S. Nat. Mus. xv, pp. 441 & 442, Honduras. Founded on a mutilated skin without skull.

Artibeus, Leach, amended by Winge (suprà, p. 25), p. 38, to Artobius.

Artibeus harti, n. sp., Thomas, Ann. N. H. (6) ix, pp. 408-410, Trinidad.

### 3. INSECTIVORA.

### a. TUPAHDA.

Tupaia everetti, n. sp., Thomas, Ann. N. H. (6) ix, pp. 250 & 251, Mindanao, Philippines.

Tupaia picta, n. sp., Thomas, t. c. pp. 251 & 252, N. Borneo.

Tupaia montana, n. sp., Thomas, t. c. p. 252, Mt. Dulit, N. Borneo. Also P. Z. S. 1892, p. 223.

Tupaia melanura, n. sp., Thomas, t. c. pp. 252 & 253, ibid. Also P. Z. S. 1892, p. 224.

#### b. MACROSCELIDIDAS.

Filhol, H. Note sur un Insectivore Nouveau. Bull. Soc. Philom. (8) iv. p. 134.

Describes and figures the hinder end of a mandibular ramus considered to be allied to the existing Rhynchocyon.

†Pseudorhynchocyon cayluxi, n. g. & sp., Filhol, l. c., Phosphorites, Caylux.

### c. Soricidæ.

†Sorex styriacus, n. sp., HOFMANN, JB. geol. Reichsanst. xlii, p. 74, pl. iii, fig. 4, Miocene, Styria.

Sorex oreopolus et saussurei, n. spp., MERRIAM, P. Biol. Soc. Washington, vii, pp. 173 & 174, Mexico.

### d. Erinaceidæ.

CARLIER, E. W. Contributions to the Histology of the Hedgehog (Erinaceus europæus). J. Anat. Phys. xxvii, pp. 85-111 & 169-178, pls. iv-vi & xi.

The first part deals with the alimentary canal, the second with the liver, and the third with the skin. The spines are shown to be the morphological equivalents of hairs.

#### 6. CENTETIDE.

THOMAS, O. On the Insectivorous Genus Echinops, Martin, with Notes on the Dentition of the Allied Genera. P. Z. S. 1892, pp. 500-505.
After noticing in detail the leading dental and other characteristics of Echinops, the author shows that the adult dentition of Centetes exhibits the Marsupial feature of having four molars. It is suggested that the presumed affinity between Solenodon and Centetes is merely due to the feature of trituberculism in the molar dentition.

Echinops telfuiri, n. var. palescens, THOMAS, t. c. p. 501, S. Madagascar.

### 4. CAUNIVORA.

#### A.—CARNIVORA VERA.

ALLEN, H. On the Cephalo humeral Muscle, and the so-called rudimentary Clavicle of Curnivora. P. Ac. Philad. 1892, p. 217.

Suggests that the ossification met with in the muscles of the shoulder of Carnivores is not really a representative of the clavicle.

GREVÉ, C. Die Geographische Verbreitung der Barenartigen. Zool. Jahrb. Abth. f. Syst. vi, pp. 588-616, pl. xxvii.

The geographical distribution of the various species of Ursidee, Procyonidee, Arcticis, Cerculeptes, &c.

- Petrini, —. Note sur la Présence de Corpuscules de Pacini et de Gauglions Nerveux dans le Pancreas du Chat. CR. Soc. Biol. 1892, pp. 275 & 276.
- Schacht, H. Die Raubsäugethiere des Teutoburger Walder. VIII, Der Fischotter (*Lutra vulgaris*); 1x, Der Dachs (*Meles taxus*). Zool. Gart. xxxiii, pp. 41-48.
- RUFFINI, A. Sur un Réticule Nerveux Spécial, et sur quelques Corpuscules de Pacini qui se trouvent en connexion avec les Organes Musculo-tendineux du Chat. Arch. Ital. Biol. xvii, pp. 101-106.

### a. FELIDE.

FABRINI, E. Su alcuni Felini del Pliocene Italiano. Atti [Mem. Rend.] Acc. Rom. (5) i, pp. 257-263.

Preliminary notes on Pliocene Feline remains in the Museum at Florence.

- † Felis hillianus [a], n. sp., Cope, Am. Nat. xxvi, p. 1029, Pliocene, Texas. Caracal berberorum, n. sp., MATSCHIE, SB. nat. Fr. 1892, p. 114, N. Africa.
- †Pseudælurus transitorius, n. sp., Déperet, Arch. Mus. Lyon, v, p. 21, pl. i, figs. 5 & 6, Miocene, Grive-St-Alban.

#### b. VIVERRIDE.

LYDEKKER, R. On the Occurrence of the so-called *Viverra hastingsia* of Hordwell in the French Phosphorites. Q. J. Geol. Soc. xlviii, pp. 373 & 374, fig.

It is shown that Viverra hastingsia, Davies, is common to the Oligocene

of France and Hordwell, and since there is no character by which the lower jaw of the type of the latter can be satisfactorily distinguished from the type of V. angustidens, Filhol, it is considered that V. hastingsiae is specifically inseparable from V. angustidens. Hence the cranium, which is the subject of the communication, is figured under the latter and earlier name. There is added a list of 7 Mammals known to be common to the Headon Beds of Hordwell and the Isle of Wight, and the French phosphorites.

Genetta vulgaris, recorded by DE KERVILLE, Bull. Soc. Rouen, 1890, pp. 79 & 80, from the Department of the Eure.

+Progenetta, n. g., Déperet, Arch. Mus. Lyon, v, p. 34, pl. i, fig. 3; type, Mustela incerta, Lartet.

Linsang, adopted by Thomas, Ann. Mus. Genov. (2) x, p. 9, in place of *Prionodon*; the latter being preoccupied by *Priodon*.

Hemigale hosei, n. sp., Thomas, Ann. N. H. (6) ix, p. 250, and P. Z. S. 1892, p. 222, pl. xviii, Mt. Dulit, Borneo.

#### c. HYÆNIDÆ.

†Borophagus diversidens, n. g. & sp., Cope, Am. Nat. xxvi, p. 1029, Pliocene, Texas.

#### d. Canidæ.

- Kulagin, N. Mittheilung über die Hunderasse Laika (Eskimo-hunde) in Russland. Zool. Jahrb. Abth. f. Syst. vi, pp. 435-441, pl. xx.
- LANGKAVEL, B. Ueber Dingos, Pariah- und Neuseeländische Hunde. Zool. Gart. xxxiii, pp. 33-38.
- —. Der Polarfuchs (Canis lagopus). T. c. pp. 79-88 & 111-119.
- Schäff, E. Ueber den Schädel von Canis adustus, Sund. Zool. Jahrb. Abth. f. Syst. vi, pp. 522-531, pl. xxv.
  - A comparison of the skull of this Jackal with those of other species.
- WHITE, T. On the Native Dog of New Zealand. Tr. N. Z. Inst. xxiv, pp. 540-557.

### e. Ursidæ.

- COOKE, J. H. On the occurrence of *Ursus arctos* in the Malta Pleistocene. Med. Nat. ii, pp. 267-269.
- Hyanarctus brevirhinus, Hofmann, JB. geol. Reichsanst. xlii, p. 64, pl. ii, figs. 1-3, = Cephalogale brevirhinus, Hofmann [Zool. Rec. xxiv, Mamm. p. 25], = Hyanarctus minutus, Koken [Zool. Rec. xxv, Mamm. p. 31].

# f. MUSTELIDÆ.

HARTING, J. E. The British Marten (Martes sylvatica, Nilsson). Zool. (3) xv, pp. 450-459 (1891), and xvi, pp. 131-138.

An account of its distribution and habits in the British Islands.

- † Martes delphinensis, n. sp., DEPÈRET, Arch. Mus. Lyon, v, p. 26, pl. i, fig. 10, Miocene, Grive-St-Alban.
- † Mustela incerta, Lartet, referred by Depèret to Viverridæ (vide suprà).
- †Canimartes cumminsi, n. g. & sp., Cope, Am. Nat. xxvi, p. 1029, Pliocene, Texas.

#### B.—PINNIPEDIA.

- Abbott, W. J. L. On the occurrence of Walrus in the Thames Valley. P. Geol. Assoc. xii, p. 357.
- Bouvier, E. L. Plexus formes par les Artères Intercostales du *Phoca vitulina*. Bull. Soc. Philom. (8) v, pp. 81-84.
- FORBES, H. O. On some points in the Anatomy of a Sea Bear caught off Sumner, Canterbury, New Zealand; with notes on the New Zealand Eared Seals. Tr. N. Z. Inst. xxiv, pp. 198-200.

A preliminary paper dealing with some points in the anatomy of Arctocephalus forsteri. The author objects to the division of the Otariidæ into Otaria and Arctocephalus; and he would subdivide the latter into Eumetopias (Hair-seals, or Sea-lions), and Arctocephalus proper (Fur-seals, or Sea-bears). In the New Zealand region he recognizes Eu. hookeri (Auckland Island Hair-seal), and A. forsteri, of which Otaria cinerea is the female. The skins described by J. W. Clarke as Otaria cinerea, Péron, are referred to A. forsteri; being those of Fur-seals. Péron's name must be dropped. Eu. hookeri appears to be confined to the Auckland Isles; while A. forsteri is common to New Zealand, Chatham Island, and the Seal-Rocks, near Port Stephens, New South Wales.

## g. OTARIIDÆ.

+Callotaria, n. nom., PALMER, P. Biol. Soc. Washington, vii, p. 156; to replace Callorhinus, Gray (preoccupied).

### †C.—CREODONTA.

FILHOL, H. Note sur le Quercitherium tenebrosum. Bull. Soc. Philom. (8) iv, pp. 135-137.

Describes and figures the upper and lower dentition of this remarkable Creedont from the Phosphorites.

OSBORN, Bull. Am. Mus. Nat. Hist. iv, pp. 103-118, gives a synopsis of the Wahsatch forms, which are divided into *Palæonictidæ*, *Oxyænidæ*, *Proviverridæ*, *Miacidæ*, *Mesonychidæ*, and *Arctocyonidæ*. *Oxyæna* is shown to have  $\frac{3}{4}$  incisors.

Scorr, W. B. A Bevision of the North American *Creedonta*, with notes on some Genera which have been referred to that Group. P. Ac. Philad. 1892, pp. 291–323.

The group is divided into the families Oxyclænidæ, Arctocyonidæ, Triisodontidæ, Mesonychidæ, Proviverridæ (= Leptictidæ, in part), Palæonyctidæ (= Ambloctynidæ), Hyænodontidæ (= Oxyænidæ), and Miacidæ, each of which is fuller defined. Then follows a classified synopsis of the genera, among which several are new (vide infrå).

#### h. Oxyclænidæ.

- †Protochriacus, n. g., Scott, P. Ac. Philad. 1892, p. 296; type, Chriacus priscus, Cope.
- † Epichriacus, n. g., Scott, l. c.; type, Chriacus schlosserianus, Cope.
- † Pentacodon, n. g., Scott, l. c.; type, Chriacus inversus, Cope.
- † Ellipsodon, n. g., Scott, t. c. p. 298; type, Tricentes inæquidens, Cope.

### i. †ARCTOCYONIDÆ.

- †Ailurarus picteti, n. g. & sp., L. RÜTIMEYER, Abh. Schw. pal. Ges. xviii, p. 97, Up. Eocene, Switzerland.
- †Clanodon, n. g., Scott, op. cit. p. 298; type, Mioclanus ferox, Cope.
- †Tetraclænodon, n. g., Scott, t. c. p. 299; type, Mioclænus floverianus, Cope.

# j. †Triisodontidæ.

† Microclanodon, n. g., Scott, t. c. p. 302; type, Triisodon assurgens, Cope.

# k. + MESONYCHIDÆ.

- † Dissacus leptognathus, n. sp., Osborn, Bull. Am. Mus. Nat. Hist. iv, p. 112, Eocene, Wahsatch Beds.
- † Pachyana gigantea, n. sp., Osborn, t. c., pp. 113 & 114, ibid.

### l. +Phoviverridae.

†Prorhyzana egerkingio, n. sp., RÜTIMEYER, Abh. Schw. pal. Ges. xviii, p. 105, Up. Eocene, Switzerland.

### m. +PALEONYCTIDE.

WORTMAN, Bull. Am. Mus. Nat. Hist. iv, pp. 95-103, points out how the dentition of *Palæonictis* and *Patriofelis* shows a transition towards that of the *Felidæ*, the upper p. 4 and the lower m. 1 developing into carnassials. *Ambloctonus* is included in the family, of which a diagnosis is given by OSBORN, t. c. p. 104.

† Patriofelis leidyanus [a] n. sp., Wortman, t.c. p. 98, Eocene, Wahsatch Beds.

†Palæonictis occidentalis, n. sp., Osborn, t. c. p. 104, ibid.

### n. †HYÆNODONTIDÆ.

†Hyænodon schlosseri, n. sp., RÜTIMEYER, Abh. Schw. pal. Ges. xviii, p. 101, Up. Eocene, Switzerland.

†Pterodon magnus, n. sp., Rütimeyer, t. c., p. 99, ibid.

#### INCERTA SEDIS.

† Protogonodon, n.g., Scott, op.cit. p. 322: type, Mioclanus pentacus, Cope. † Paradoxodon, n.g., Scott, l. c.; type, Chriacus ruetimeyeranus, Cope.

†Carcinodon, n. g., Scott, t. c., p. 323; type, Mioclænus filholianus, Cope.

### 5.—RODENTIA.

BEDDARD, F. E. On the Convolutions of the Cerebral Hemispheres of certain Rodents. P. Z. S. 1892, pp. 596-613, figs.

This paper refers chiefly to Dasyprocta, Calogenys, Lagostomus, Hydrocharus, and Dolichotis, being the genera of Rodents in which the brains show the greatest development of convolutions. The author illustrates the form of the convolutions in these and other genera, and discusses the bearing of his observations upon classification.

- Cannieu, A. Recherches sur l'Appareil Reproducteur Male du Cavia. Rev. Sci. Nat. Ouest. ii, pp. 55-63, and figs. [incomplete].
- Duval, M. Le Placenta des Rongeurs et l'Inversion des Feuillets Blastodermiques. T. c. pp. 917-919.
- —... Le Placenta des Rongeurs. J. de l'Anat. Phys. xxviii, pp. 58-98 & 333-453, pls. iii-v, xvi, & xvii.

The completion of the memoir, of which the earlier portions have already been recorded.

FISERIUS, E. Beiträge zur Entwicklungsgeschichte von Sciurus vulyaris. Verh. Ges. Wurzb. xxvi, pp. 103-122, pl. ii.

An epitome of the developmental history of this species.

FLEISCHMANN, A. Die einheitliche Plan der Placentarbildung bei Nagethieren. SB. Ak. Berlin, 1892, pp. 445-457, pl. ii.

It is concluded that the view of Byder that the discoidal placenta of the Rodents is derived from the diminution of a zonary one, is well supported.

FREUND, P. Beiträge zur Entwicklungsgeschiete der Zahnanlagen bei Nagethieren; Arch, mikr. Anat. xxix, pp. 525-555, 2 pls.

Confirms previous observations as to the presence in Lepus of a rudimentary tooth in each jaw in front of the large incisor, which is regarded as a much-reduced deciduous predecessor of the latter. In the upper diastema of Lepus there is a distinct dental ridge, without enamel-germs; but the latter occur in Sciurus. It is concluded that the Duplicidentata present a very old type of the order, next to which comes the Sciurus morpha. The view that Rodents are directly descended from Marsupials is supported. The memoir is translated in Ann. N. H. (6) xi, pp. 301-327 (1893).

- GILIS, P. Note sur un Muscle Costo-Basilaire chez le Cochon d'Inde. CR. Soc. Biol. 1892, p. 1018.
- POUSARGUES, E. DE. Note sur l'Appareil Génital Mâle du Cochon d'Inde (Cavia cobaya). Bull. Soc. Philom. (8) v, pp. 45-48.
- ROBINSON, A. Some Points in the Early Development of Mus musculus and Mus decumanus; the Relation of the Yolk-sac to the Decidua and Placenta. Rep. Brit. Ass. 1891, pp. 690 & 691.
- —... Observations upon the Development of the Spinal Cord in Mus musculus and Mus decumanus; the Formation of the Septa and Fissures. T.c. pp. 691 & 692.

### a. SCIURIDE.

STEPHENS, F. Notes on Sciurus fossor, Peale. Zoe, iii, pp. 118 & 119.

The author thinks it probable that the name S. leporinus will eventually have to be adopted for this Carolina Squirrel.

Sciurus brookei, n. sp., Thomas, Ann. N. H. (6) ix, p. 253, and P. Z. S. 1892, p. 225, Mt. Dulit, N. Borneo.

Sciurus lowi, n. sp., THOMAS, t. c. p. 253, Lumbidan.

Sciurus pryeri and hosei, n. spp, Thomas, op. cit. x, pp. 214 & 215, N. Borneo.

Sciurus undulatus, n. sp., TRUE, P. U. S. Nat. Mus. xv, pp. 465-467, E. Africa.

Sciurus getulus: POMEL, C.R. cxiv, pp. 53 & 54, has a note on this almost unknown African species.

Xerospermophilus, n. subg., MERRIAM, P. Biol. Soc. Washington, vii, p. 27; type, Spermophilus mohavensis.

1892. [vol. xxix.]

Ammospermophilus, n. subg., MERRIAM, l. c.; type, S. leucurus.

Arctomys caudatus: notes on its distribution by BÜCHNER, Mél. biol. xiii, pp. 309-324, where it is shown that this species has a wide range in Central Asia, extending to regions north of the Altai.

Cynomys mexicanus, n. sp., MERRIAM, op. cit. p. 157, Mexico.

†Plesiarctomys schlosseri, n. sp., RUTIMEYER, Abh. Schw. pal. Ges. xviii, p. 89, Up. Eocene, Switzerland.

### b. Castoridæ.

MARTIN, H. T. Castorologia: the History and Traditions of the Canadian Beaver. London & Montreal: 1892, 8vo, 238 pp., illus.

Describes the Beaver from the historical, zoological, distributional, and commercial aspects; with notices of other American Rodents, recent and fossil.

NEWTON, E. T. On a Skull of *Trogontherium cuvieri* from the Forestbed of East Runton, near Cromer. T. Z. S. xiii, pp. 165-175, pl. xix.

The complete memoir, of which the abstract is quoted in Zool. Rec. xxviii, Mamm. p. 30.

#### c. MURIDÆ.

HARTING, J. E. Plague of Field Voles in the South of Scotland. Zool. (3) xvi, pp. 161-173.

JENTINK, F. A. On a New Species of Rat from the Island of Flores. In Weber's Zool. Ergebnisse einer Reise in Niederländish Ost-Indien, ii, pp. 78-83, pl. iii.

The species in question is the largest Rat yet known. The Indian Mus setifer is referred to Nesocia, which is regarded as a subgenus of Mus (vide infrà).

Gerbillus calurus, hab. unknown, G. gracilis, Gambia. and G. emini, Wadelai, n. spp., Thomas, Ann. N. H. (6) ix, pp. 76-79.

Meriones tristrami, n. sp., THOMAS, t. c. p. 148, Palestine.

Dendromys nigrifrons, n. sp., TRUE, P. U. S. Nat. Mus. xv, pp. 462-464, E. Africa.

Steatomys bocagei, n. sp., Thomas, Ann. N. H. (6) x, p. 264, Angola. It is also noted that the proper name of S. edulis is S. pratensis.

Rhithrodontomys, Giglioli, 1873, should replace Ochetodon, Coues, 1874; MERRIAM, P. Biol. Soc. Washington, vii, p. 26.

Sitomys, Fitzinger, should replace the untenable Hesperomys; MERRIAM, t. c. p. 27.

Sitomys musculus, n. sp., MERRIAM, t. c. p. 170, Mexico.

Oryzomys bauri, n. sp., J. A. Allen, Bull. Am. Mus. Nat. Hist. iv, pp. 48-50, Galapagos I.

- Neotoma alleni and tenuicauda, n. spp, MERRIAM, P. Biol. Soc. Washington, vii, pp. 168-170, Mexico.
- Xenomys nelsoni, n. g. & sp., MERRIAM, t. c. pp. 159-163, Mexico.
- Arcicola phaus, n. sp., MERRIAM, t. c. pp. 171 & 172, Mexico.
- †Bramus barbarus, n. g. & sp., POMEL, C.R. exiv, pp. 1159-1163, Pleistocene, Tunis. Allied to Microtus.
- Isomys pumilio, n. varr. diminutus and bechuana, Thomas, P. Z. S. 1892, pp. 551 & 552, Masailand and Bechuanaland.
- Mus chiropus, n. sp, Thomas, Ann. Mus. Genov. (2) x, p. 889 (1891), Karin Hills, Burma. See also t. c. p. 935, pl. xi.
- Mus (Dasyonys) bentleyæ, p. 179, Lower Congo, M. daltoni, p. 181, W. Africa, M. burtoni, pp. 182 & 183, Ashanti, n. spp., Thomas, Ann. N. H. (6) x.
- Mus aquilus, n. sp., TRUE, P. U. S. Nat. Mus. xv, pp. 460-462.
- Mus armandvillei, n. sp., JENTINK, op. cit., Flores.
- Acomys wilsoni, n. sp, THOMAS, Ann. N. H. (6) x, p. 22, Mombass.
- Coniturus, W. Ogilby, 1838, adopted by J. D. Ogilby, Cat. Australian Mammals (supra, p. 12), p. 113, for Hapalotis, Licht., preoccupied.
- Pithechir melanurus, JENTINK, Notes Leyd. Mus. xiv, pp. 122-126, pls. iii & iv, gives a description of this rare red-coloured Rat from Java.
- †Trilophomys, n. g., Depéret, Mém. Soc. Géol.-Pal. iii, p. 121; to replace Lophiomys, Dep. [Zool. Rec. xxvii, Mamm. p. 37], preoccupied.

#### d. SPALACIDÆ.

Spalaz typhlus, recorded by Anderson, P. Z. S. 1892, pp. 472-476, from Egypt, with notes on its habits.

### e. GROMVIDÆ.

- Geomys nelsoni, formosus, and gymnurus, n. spp., MERRIAM, B. Biol. Soc. Washington, vii, pp. 164-167, Mexico. (The first of these is apparently identical with the undermentioned species, which has the priority.—R. L.)
- Geomys bulleri, n. sp., THOMAS, Ann. N. H. (6) x, p. 196, Jalisco, Mexico. Perodipus, FITZINGER, SB. Ak. Wien. 1867, p. 126, is stated by MERRIAM, P. Biol. Soc. Washington, vii. p. 26, to be the proper name of the genus termed Dipodops by Merriam, 1890 [see Zool. Rec. xxvii, Mamm. p. 37].
- Perognathus merriami, n. sp., J. A. Allen, Bull. Am. Mus. Nat. Hist. iv, pp. 45 & 46, pl. iii, S.E. Texas.

### f. Dipodida.

Sminthus concolor, n. sp., BÜCHNER, Mél. biol. xiii, pp. 268-271, Kansu, China.

# g. Octodontidæ.

BEDDARD, F. E. On the Brain and Muscular Anatomy of Aulacodus-P. Z. S. 1892, pp. 520-527, figs.

Describes the anatomy of the muscles and brain in detail, and that of the viscera more briefly; with figures of the hard palate and the brain.

†Capromys columbianus, n. sp., Chapman, Bull. Am. Mus. Nat. Hist. iv, pp. 314 & 315, Subfossil, Cuba.

#### h. LEPORIDE.

CHRISTY, M. On the Extermination of the Rabbit in Australasia. Zool. (3) xvi, pp. 377-388.

In the course of this article the author alludes to the remarkable variation in the numbers of *Lepus americanus* in the north-west of Canada; the animals being apparently attacked by au epidemic.

LATASTE, F. À propos de Lapins Domestiques Vivant en Liberté dans l'Îlot de l'Étang de Cauquenes (Colchagua). Act. Soc. Chili, ii, pp. 210-222.

The author discusses the bearings of this case in regard to feral Rabbits in general. He concludes that domestic Rabbits run wild, never revert to the original wild race, nor do they ever form a new species. Generally such feral Rabbits tend to lose individual variations, and to form a more or less uniform race.

Lepus cuniculus: a wild Albino recorded by DE KERVILLE, Bull. Soc. Rouen, 1891, pp. 61 & 62. For formula of milk-dentition, see Woodward, infrå, p.

Lepus sinensis, n. var. coreanus, THOMAS, Ann. N. H. (6) ix, pp. 146 & 147, Corea.

### 6.—UNGULATA.

TAEKER, J. Zur Kenntniss der Ontogenese bei Ungulaten. Dorpat: 1892, 8vo, 27 pp., 4 pls.

An elaborate treatise on the mode of formation and development of the cheek-teeth of the Ungulates. The forms described are the Horse, the Pig, Dorcatherium, the Elk, the Roe, the Sheep, and the Ox. It is concluded that all selenodont types show an initial bunodont condition. In all forms cones are first developed on the crowns of the molars, which in the bunodont types tend to assume a pyramidal form, and unite with one another at their bases; while in the selenodonts they take a crescent shape. The accessory column and cingulum in the latter are of secondary origin. The more complicated types of selenodont molars result from

additions of new elements to the original cones, such additions being added in the upper molars to the paracone, and not, as has been supposed, to the protocone. In selenodonts it is the antero-external cone which first makes its appearance, and it is accordingly urged that this, instead of the inner cone, ought to be called the protocone.

### A.—PROBOSCIDEA.

HICKS, H. On the Discovery of Mammoth and other Remains in Endsleigh Street, &c. Q. J. Geol. Soc. xlviii, pp. 453-467.

It is concluded that the Mammoth lived very early in the Glacial period.

Howorth, Sir H. H. The Mammoth and the Glacial Drift. Geol. Mag. (3) ix, pp. 250-258 & 396-405.

The author endeavours to prove that the Mammoth is of pre-rather than post-glacial age.

- MARTIN, K. Mammuthreste aus Niederland. JB. Mineral. 1892, i, pp. 45-48.
- Möbius, K. Die Behaarung des Mammuths und der Lebenden Elephanten, vergleichend untersucht. SB. Ak. Berlin, 1892, pp. 527-538. pl. iii.

Records the existence in living Elephants of a remnant of the woolly coat of the Mammoth.

Pohlig, H. Monographie des *Elephas antiquus*; Nachträge zur Dentition des *Elephas antiquus*. N. Acta Ac. L.-C. Nat. cur. lvii, pp. 260-466, pls. and text, figs.

The completion of the monograph quoted in Zool. Rec. xxv, Mamm. p. 43.

- STEFANESCU, G. On the Existence of the *Dinotherium* in Roumania. Bull. Geol. Soc. Am. iii, pp. 81-83.
- † Mastodon longirostris from Lahr, Eck, JB. Mineral, 1892, i, p. 151.
- †Mastodon successor, n. sp., Cope, P. Ac. Philad. 1892, p. 227, Up. Tertiary, Texas.
- †Mastodon giganteus, restored by Marsh, Am. J. Sci. (3) xlv, p. 350, pl. viii.

#### B.—†AMBLYPODA.

# a. †Coryphodontidæ.

Osborn, Bull. Am. Mus. Nat. Hist. iv, pp. 120-122, shows that while the fore-foot was of the semi-digitigrade type of the Elephant, the pes was truly plantigrade.

EARLE, C. Revision of the Species of Coryphodon. Bull. Am. Mus. Nat. Hist. iv, pp. 149-166, figs.

It is concluded that Bathmodon and Metalophodon are synonyms of Coryphodon, and that there are 8 American species of the latter, together with 1 each of Manteodon and Ectacodon. The cusps of the molars are homologized with those of other Ungulates.

#### C.—HYRACOIDEA.

### b. Procaviida.

THOMAS, O. On the Species of the Hyracoidea. P. Z. S. 1892, pp. 50-76, pl. iii.

After a careful revision of all the known species, it is concluded that it appears impossible to refer the Hyraces to more than a single genus, for which the name *Procavia* is adopted. 14 species (one of which is new), and 4 varieties are recognized, so far as available material admits of forming a final conclusion. These species are defined, and their distribution described, with the aid of a map. The skull of *P. pallida* is figured.

WOODWARD, M. F. On the Milk-Dentition of *Procavia* (*Hyrax*) capensis and of the Rabbit, with remarks on the relation of the Milk and Permanent Dentitions of the *Mammalia*. T. c. pp. 38-49, pl. ii.

It is concluded that in the Hyrax the milk-dentition is  $i\frac{2}{3}$ ,  $c\frac{1}{1}$ ,  $mm\frac{4}{4}$ , and the permanent  $i\frac{1}{3}$ ,  $c\frac{2}{0}$ ,  $p\frac{4}{4}$ ,  $m\frac{2}{3}$ . The upper milk-canine may, however, sometimes be retained in the permanent series. In the Rabbit the formula of the milk-teeth is found to be  $i\frac{2}{1}$ ,  $mm\frac{3}{4}$ . The common Hare and the Rabbit are the only known Rodents with deciduous incisors, while the Rabbit is peculiar in having milk-predecessors to both incisors.

Procavia latastei, n. sp., THOMAS, t. c. p. 69, Senegal.

Procavia stuhlmanni, n. sp., MATSCHIE, SB. nat. Fr. 1892, pp. 110-112, Victoria Nyanza.

Dendrohyrax validus: skull figured by TRUE, P. U. S. Nat. Mus. xv, pl. lxxv.

#### D.-+CONDYLARTHRA.

MARSH, O. C. A New Order of Extinct Eccene Mammals. Am. J. Sci. (3) xliii, pp. 445-449, figs.

Regards Meniscotherium and a form described as Hyracops as indicating a new order, allied to the Hyracoidea, but presenting very generalized features in the dentition and feet; the digits terminating neither in claws nor in hoofs. From this character the name Mesodactyla is proposed. The paper is criticized by COPE, Am. Nat. xxvi, pp. 411 &

412, who doubts whether Hyracops is distinct from Meniscotherium, and confirms his previous reference of the latter to the Condylarthra.

OSBORN, H. F. Is Meniscotherium a Member of the Chalicotheroidea? Am. Nat. xxvi, pp. 507-509, figs.

The question is provisionally answered in the affirmative; Meniscotherium being perhaps the ancestor of Chalicotherium.

#### c. †PHENACODONTIDAL.

† Meniscolon picteti, n. sp., RÜTIMEYER, Abh. Schw. pal. Ges. xviii, p. 10, Up. Eocene, Switzerland. The Genus Meniscolon, Rüt., dates from 1888, but is quoted in Zool. Rec. xxv, Mamm. p. 45, as being a misprint for Meniscotherium.

### d. †Meniscotheriidæ.

†Hyracops socialis, n. g. & sp., Marsh, Am. J. Sci. (3) xliii, pp. 446 & 447, Low. Eocene, New Mexico.

#### E.—PERISSODACTYLA.

The following classification is proposed by Osborn, Bull. Am. Mus. Nat. Hist. iv, p. 92:—

Buno-selenodont. . Titanotheriida . . Palaosyopina . . Palaosyops.

Titanotheriina. . Titanotherium.

Lopho-selenodont . Equidæ . . . . . . Hyracotheriuæ . Hyracotherium.

Anchitheriinæ . . Anchitherium.

Equina . . . . . Equus.

Sub-lophodont . . . Tapiridæ . . . . . Systemodontinæ . Systemodon.

Tapirinæ . . . . . Tapirus.

Transitional . . . . Helaletite . . . . Helaletes.

Lophiodontidæ. . Lophiodon.

Lophodont . . . . . Hyracodontidæ . Hyrachyinæ . . . Hyrachyus.

Hyracodontina. . Hyracodon.

Triplopodinæ. . . Triplopus.

Amynodontidæ.. Amynodon. Rhinocerotidæ.. Aceratheriinæ.. Aceratherium.

Diceratheriina. . Diceratherium.

Rhinocerotinæ . . Rhinoceros.

Elasmotheriinæ . Elasmotherium.

Systemodon is regarded as an ancestral Tapir (t. c. p. 124), and is distinguished from Hyracotherium and the equine line by the approximation of the first lower premolar to the canine.

### e. †LOPHIODONTIDE.

OSBORN, Am. Nat. xxiv, pp. 763-765, considers that the European forms included under the genus Lophiodon are really referable to three distinct families. The more typical forms, which might be retained in Lophiodon, have no representatives in America, but the others seem allied to transatlantic types, as indicated in the following table:—

- $\left. \begin{array}{c} L. \ isselensis, \ parisiensis, \ tapiroides. \end{array} \right\} \ = \ Lophiodon \ \ Lophiodontidæ.$
- L. annectans = Isectolophus Tapiridæ.
- L. cartieri = Hyrachyus Hyracodontidæ.
- L. rhinoceroides = Amynodon Rhinocerotidæ.
- †Lophiodon annectans, n. sp., RÜTIMEYER, Abh. Schw. pal. Ges. xviii, p. 34, Up. Eocene, Switzerland.
- †Propalæotherium minutum, n. sp., Rütimeyer, t. c. p. 35, ibid.

## f. Equide.

- CIACCIO, G. V. Sur une Étrange et Remarquable Particularité de Structure observée dans le Cornée d'un Cheval. Arch. Ital. Biol. xvii, pp. 295-297.
- COPE, E. D. On the Permanent and Temporary Dentitions of certain Three-toed Horses. Am. Nat. xxvi, pp. 942-944, pls. xxv & xxvi.

The changes in the upper molars of *Protohippus placidus* are pointed out; in the early state the dentition is that of *Hippotherium*, and it is not till middle life that the protocone fuses with the paraconule to form the typical *Protohippus* molar. It is added that in all the group the milk-dentition is much simpler than the permanent one; the milk-molars approximating to the permanent molars of *Anchitherium*.

Marsh, O. C. Recent Polydactyle Horses. Am. J. Sci. (3) xliii, pp. 339-354, figs.

Describes and figures feet of several existing Horses having supernumerary digits, which are regarded as indicating reversions to primitive polydactyle forms. A general survey of the extinct North American Perissodactyles allied to the Horse is appended, with illustrations of the foot-structure. In the course of this the new name Hippops is suggested for the undiscovered pentadactyle ancestor of the Horse; and it is also stated that Phenacodus, Cope, should be replaced by the earlier Helohyus, Marsh. Criticized by Cope, Am. Nat. xxvi, pp. 410 & 411, who states that Helohyus was never defined.

POMMEROL, F. Sur un Petit Cheval Quaternaire trouve dans la Limagne. C.R. Ass. Fr. Sci. xix, pt. 2, pp. 567-573, figs.

A replica of the paper quoted in Zool. Rec. xxvii, Mamm. p. 41.

SCHÖNLAND, S., Nature, xlvi, pp. 6 & 7, confirms the statement as to the coloration of the Zebra being indistinguishable in moonlight from surrounding objects, and adds that it is likewise protective in daylight.

TROUESSART, E. L. The Fiction of the American Horse, and the Truth on this Disputed Point. Science, xx, pp. 128 & 129.

It is concluded that the statement as to the occurrence of Horses in La Plata at the time of the discovery of the New World is unfounded. Some observations are added on fossil species.

- VAUGHAN, T. Strangeway's Veterinary Anatomy. 4th edition. Edinburgh: 1892, 8vo, 601 pp., illustrated.
- † Equus simplicidens, n. sp., COPE, P. Am. Phil: Soc. 1892, p. 124, Pliocene (?), Texas.
- Equus boehmi, n. sp., MATSCHIE, SB. nat. Fr. 1892, p. 131, German East Africa.

## q. †PALEOTHERIIDÆ.

† Paloplotherium magnum, n. sp., RÜTIMEYER, Abh. Schw. pal. Ges. xviii, p. 19, Up. Eocene, Switzerland.

#### h. RHINOCEROTIDÆ.

- LYDERKER, R. Notes on Rhinoceroses, Ancient and Modern. Field, lxxix, p. 903, & lxxx, p. 38.
- Rhinoceros simus, note on by H. A. BRYDEN, ibid. lxxx, p. 84.
- PAVLOW, (MADAME) M. Les Rhinoceridæ de la Russie, et le Developpement des Rhinoceridæ en general. Bull. Mosc. 1892, pp. 137-221, pls. ii-v.

After recording the occurrence of the Russian representatives of the family, the author enters into a long discussion as to the mutual relationships of the various species of Rhinoceros, Aceratherium, etc.; her views as to the phylogeny of the group being expressed in tabular form. It is considered that R. antiquitatis has no affinity with R. simus and R. platyrhinus, but is allied to R. leptorhinus; while both R. indicus and R. sondaicus (javanicus) are derived from R. sivalensis, the latter through the bicorn R. karnuliensis. Aceratherium croizeti, of the European Oligocene, is referred to the American genus Amynodon, although the characteristic third upper molar is at present unknown. The Phosphorites of France also yield a species nearly allied to the American Aceratherium occidentale. The resemblance of Lophiodon rhinoceroides to the Rhinocerotides is referred to (cf. Obborn, supra, p. 40).

## i. †TITANOTHERIIDÆ.

EARLE, C. A Memoir upon the Genus *Palæosyops*, Leidy, and its Allies. J. Ac. Philad. ix, pp. 267-388, pls. x-xiv.

The author concludes that the family Lambdotheriidæ should be abolished, and that Palæosyops and its allies should be included in the Titanotheridæ, of which they constitute a subfamily, the Palæosyopinæ. This subfamily is characterized by all the premolars being unlike the molars, whereas in the Titanotherinæ some of the former are molariform. The genera of the Palæosyopinæ are distinguished as follows:—

- I. All three incisors present in each jaw.
  - A. External lobes of upper molars very oblique and shallow . . . 1. Lambdotherium.
  - B. External lobes of upper molars straight and deep.
    - a. Premaxillary symphysis short
       and round, last upper molar
       with two inner cones . . . 2. Limnohyops.
       Last upper molar with one inner
      - cone . . . . . . . . 3. Palæosyops.
    - b. Premaxillary symphysis elongate
      - and narrow . . . . . . 4. Telmatotherium.
- II. Only two incisors in each jaw . . . . 5. Haplacodon.

The author has some doubts as to the propriety of separating Limno-hyops and Telmatotherium from Palxosyops, and concludes that the relationship between the first and third is closer than that between the second and third. The abortion of one of the inner cones in the upper molars of Limnohyops is considered, as in the analogous instance of the Rhinocerotide, to indicate the greater specialization. As regards the succession in time of the genera of the whole family, the horned Titano-therium is Miocene, the hornless Diplacodon Uinta Eocene, Telmato-therium, Palxosyops, and Limnohyops are from the upper Bridger Eocene, while Lambdotherium is from the lower Bridger, and perhaps the underlying Wasatch. Lambdotherium is closely allied to the ancestral stock of the family, but the absence of p may indicate that it is not the actual ancestor.

#### INCERTÆ SEDIS.

## j. CHALICOTHERIIDÆ.

Macrotherium grande, n. var. rhodanicum, Deperet, Arch. Mus. Lyon, v, p. 63, pls. ii-iv, Miocene, Rhone Valley (skull figured).

#### F.—ARTIODACTYLA.

#### k. BOVIDE.

LYDEKKER, R. Wild Oxen. Land and Water, lii, pp. 738-740 (1891), and liii, pp. 25, 53, 54, 79, & 80, woodcuts.

An illustrated popular account of all the living and some of the fossil species.

—. Wild Goats. Field, lxxix, pp. 312, 313, 355, 356, & 393.
A popular illustrated account of all the species.

NEHRING, SB. nat. Fr. 1892, pp. 129 & 130, has notes on the atlas and axis vertebres of Bos primigenius.

Sclater, P. L. On a New Antelope from Somaliland, and on some other Specimens of Antelopes from the same country. P. Z. S. 1892, pp. 98-102, pl. v, figs.

The new Antelope is a species of *Bubalis*, that generic name being adopted in place of *Alcelaphus*. The other species referred to are those mentioned in the following paper.

—. The Antelopes of Somaliland. Nat. Science, i, pp. 255-265, figs.

An account of all the Antelopes known to inhabit this region. The full list of the species from North Somaliland is now as follows, viz.:—

- Swayne, H. G. C. Field-Notes on the Antelopes of Northern Somaliland, P. Z. S. 1892, pp. 300-308.
- THOMAS, O. On the Antelopes of the Genus Cephalophus. T. c. pp. 413-430.

A revision of all the species, which are considered to be 18 in number.

WARD, R. New African Antelopes. Field, lxxx, p. 84.

Notices and figures heads of Oryx callotis, Bubalis jacksoni, and Connochastes albo-jubatus.

- Bubalis swaynei, n. sp., Sclater, P. Z. S. 1892, p. 98, pl. v, Somaliland; see also t. c. p. 257.
- Bubalis jucksoni, n. sp., Thomas, Ann. N. H. (6) ix, pp. 386 & 387, country between the Victoria Nyanza and the Naivasha.
- Bubalis leucoprymmus, n. sp., MATSCHIE, SB. nat. Fr. 1892, p. 137, German East Africa.
- Damalis jimela, n. sp., MATSCHIE, t. c. p. 135, ibid.
- Connochates taurinus, n. var. albojubatus, Thomas, Ann. N. H. (6) ix, p. 387, Uganda.
- Oryx callotis, n. sp., Thomas, P. Z. S. 1892, pp. 195 & 196, pl. xiv, East Africa. Figured by TRUE, P. U. S. Nat. Mus. xv, pl. lxxvi.
- †Antilope (? Nagor) maupasi, n. sp., Pomel, C.R. cxv, pp. 213-216, Pleistocene, Algeria.
- Æpyceros melampus, n. var. johnstoni, Thomas, P. Z. S. 1892, pp. 553 & 554, Nyassaland.
- Cephalophus jentinki, n. sp., THOMAS, t. c. p. 417, Liberia.
- Cephalophus dorsalis, n. var. castaneus, THOMAS, t. c. p. 421, Cameroons.
- Cephalophus abyssinicus, n. n., Thomas, t. c. p. 427, to replace the inadmissible C. madoqua.
- Cephalophus æquinoctialis, n. sp., MATSCHIE, SB. nat. Fr. 1892, p. 112, Victoria Nyanza.
- Cephalophus spadix, figured by TRUE, P. U. S. Nat. Mus. xv, pl. lxxviii.
- Strepsiceros suara, n. sp., MATSCHIE, op. cit. p. 135, German E. Africa.
- Antilope triangularis, Günther (= Doratoceros, Lydekker), provisionally considered by NICOLLS and EGLINGTON (supra, p. 12) to be founded upon abnormal horns of the Kudu.

#### l. GIRAFFIDÆ.

- Moewes, F. Ueber die gegenwärtige Verbreitung der Giraffe im Süden des Sambesi. Zool. Gart. xxxiii, pp. 51-53.
- †Libytherium macrusium, n. g. & sp., Pomel, C.R. cxv, pp. 100-102, Pliocene, Algeria.

## m. CERVIDE.

CAMERON, A. G. The Value of the Antlers in Classification of Deer. Field, lxxix, pp. 625, 703, 741, 860, & 861, figs.

The author urges that antiers are of primary importance in the classification of the Deer, but that they must be studied from an evolutionary and palæontological point of view; and that those of the first and second year are, in some respects, of more value than those of adult animals. It is considered that Rangifer, Alces, Cervus, and Cariacus all indicate primary branches of the family, which have had no connection with one another since antiers first originated, and hence that the component parts

of these antiers cannot be homologized. The brow-tine of Cervus is considered to be quite unrepresented in Cariacus, the vertically directed tine found in some species of the latter being regarded as distinct from a brow-tine. Hence the Deer are divided into brow-antiered and fork-antiered types. And it is further considered that Père David's Deer belongs to the second group, and should therefore be regarded as generically distinct from Cervus, under the title of Elaphurus.

- LYDEKKER, R. The Deer of Asia. Land and Water, liii, suppl., Jan. 23, pp. x & xi, & pp. 130, 131, 164, 165, 192, 193, 221, 247, & 248.
- Pohlig, H. Die Cerviden des Thüringischen Diluvial-Travertines, mit Beiträgen über undere diluviale und über recente Hirschformen. Palsontographica, xxxix, pp. 215-261, pls. xxiv-xxvii.

This forms the second fasciculus of the fauna and flora of the formation in question. The species include Cerous giganteus, C. dama, C. elaphus, Alces machlis, Rangifer tarandus, and Cupreolus capræa. [Vide infrà.]

Pousargues, E. de. Sur un Anomalie des Pattes Anterieures d'une Biche (Cervus xanthopygus). Bull. Soc. Philom. (8) iv, pp. 95-97.

Describes and figures a horny growth near the carpus in this stag.

Scheler, G. v. Ueber die Ursachen abnormer Geweihbildung bei den Hirscharten, insbesoudere die Bildung von mehr als zwei Geweihstangen. JH. Ver. Württ. xlviii, pp. 135-178, pls. iii-vi.

Describes and figures a number of abnormal antiers of Deer, with a discussion of the mode of formation.

Cercus elaphus, n. var. antiqui, Pohlig, op. cit. p. 246, Pleistocene. Cercus cashmirianus, n. var. yarkandensis, Blanford, P. Z. S. 1892, pp. 116 & 117, Yarkand.

tCerrus giganteus: A. NEHRING. SB. nat. Fr. 1892, pp. 3-8, describes a head of the variety ruff, dredged from the Rhone near Worms. This specimen (figured in Deutsche Jäger-Zeitung, xviii, pp. 571-573) shows that the antiers, instead of extending horizontally, are directed upwards and outwards, like those of the Fallow Der. In Verh. anthrop. Ges. 1892, pp. 125-127, the same author gives reasons for believing that the word Schelch, occurring in the "Niebelungen Lied," is not, as has been thought, applicable to this species, but probably refers either to the Elk or to a wild Stallion.

†Cercus giganteus (= euryceros), n. varr. germaniæ and italiæ, Pohlig, op. cit. pp. 220 & 228, Pleistocene.

Cerrus dama, n. var. gastaldii, Pohlig, op. cit. p. 241, Pleistocene.

†Cervus pachygenys, n. sp., POMEL, C.R. cxv, pp. 213-216, Pleistocene, Algeria.

Cervulus feæ, figured by THOMAS, Ann. Mus. Genov. (2) x, pl. x.

Alces machlis (= Cerous alces), n. varr. diluvii and europæ, Pohlig, op. cit. pp. 236 & 237, Pleistocene.

†Capreolus ruscinensis, n. sp., DEPÉRET, Pliocene, Roussillon, France.

## n. TRAGULIDÆ.

Tragulus nigricans, n. sp., THOMAS, Ann. N. H. (6) ix, p. 254, Balabac, Philippines.

† Hyomoschus peneckei, n. sp., Hofmann, JB. geol. Reichsanst. zlii, p. 72, pl. ii, figs. 4, 5, Miocene, Styria.

## o. PROTOCERATIDÆ (INCERTÆ SEDIS).

OSBORN, H. F., & WORTMAN, J. L. Characters of *Protoceras* (Marsh), the New Artiodactyl from the Lower Miocene. Bull. Am. Mus. Nat. Hist. iv, pp. 351-372, figs.; see also Nature, xlvii, pp. 321 & 322 (1893).

The authors describe a skull remarkable for the presence of a number of paired irregular bony processes, comparable to those of *Uintatherium*. Hornless (female) skulls of the same type were also obtained; and it was found by comparison that both belonged to *Protoceras*, Marsh (Zool. Rec. xxviii, *Mamm*. p. 44). That genus must accordingly indicate a distinct family group of Artiodactyles, approximating, in some degree, to the *Tragulina* through *Leptomeryx*. The feet, however (of which specimens were discovered with the skulls) are of the inadaptive modification, and there are important differences in the structure of the cranium. The resemblance of the male skull to that of *Uintatherium* is further increased by the presence of large sabre-like upper tusks.

## p. Camelidæ.

CRAGIN, F. W. Observations on Llama Remains from Colorado and Kansas. Amer. Geol. ix, pp. 257-260.

Restricting the term Auchenia to those forms in which the premolars are  $\frac{2}{1}$ , the author considers it doubtful whether any of the West American Llamas belong to that genus. The remainder of the paper is devoted to the description of specimens from the localities mentioned. Auchenia hesterna, Leidy, is provisionally referred to Holomeniscus.

- †Auchenia huerfanensis, n. sp., CRAGIN, l. c. (?), Pliocene, Huerfano River, Colorado.
- Camelus bactrianus: Blanford, P. Z. S. 1892, pp. 370 & 371, notices the skull and skin of a specimen shot by Major Cumberland in the neighbourhood of Yarkand; and concludes that all the examples found in that district are feral.

## q. †Dichodontida.

- +Dichodon cartieri, n. sp., RÜTIMEYER, Abh. Schw. pal. Ges. xviii, p. 79, Up. Eocene, Switzerland.
- †Tetraselenodon kowalevskii, recorded by Rütimeyer, t. c. p. 82, from same.

## r. Anoplotheriidæ.

- †Xiphodontherium pygmæum and obliquum, n. spp., RÜTIMEYER, Abh. Schw. pal. Gea. xviii, p. 74, Up. Eocene, Switzerland.
- †Xiphodontherium schlosseri, n. sp., RÜTIMEYER, t. c. p. 75, ibid.
- † Mixtotherium gresslyi, n. sp., RÜTIMEYER, t. c. p. 78, ibid.
- LYDEKKER, R. On Dacrytherium ovinum from the Isle of Wight and Quercy Q. J. G. S. xlviii, pp. 1-4, pl. i.

Describes a cranium and mandible of Dacrytherium cayluxi from the Quercy Phosphorites, which proves the identity of this form with the Dichobune ovina of Owen, from the Oligocene of the Isle of Wight. The species should thus be known as Dacrytherium ovinum. It is shown that the mandible referred by Filhol to D. cayluxi belongs to another animal. Hyopotamus picteti, Lyd., is identified with Dacrytherium (see also RÖTIMEYER, supra, p. 15).

## s. †Cænotheriidæ.

†Dichobune muelleri, langii, p. 70, and pygmæa, p. 72, n. spp., RÜTI-MEYER, Abh. Schw. pal. Ges. xviii, Up. Eocene, Switzerland.

#### t +Anthracotheride.

†Hypotamus gresslyi, n. var. minor, Rütimeyer, t. c. p. 61, ibid.

#### u. CHŒROPOTAMIDÆ.

HOERNES, B. Zur Kenntniss der Milchbezahnung der Gattung Entelodon, Aym. SB. Ak. Wien, 1892, Abth. ii, pp. 17-24, 1 pl.

Describes part of a mandibular ramus with the milk dentition, showing that the last milk-molar has the usual trilobed form.

## v. SUIDÆ.

- Nehring, A. Zwei Javanische Wildschweine des Berliner Zoologischen Gartens (wahrscheinlich Sus longirostris, Nehring). Zool. Gart. xxxiii, pp. 7-11.
- —. Einige neue Notizen über die Langrüsselschweine (Sus longirostris) im Berliner Zoologischer Gartens. T. c. pp. 240-242.

Notices two pigs from Java in the Berlin Gardens, which may belong to S. longirostris, that form being intermediate between S. barbatus and S. verrucosus.

Sussporf, J. Der Hauer eines Suiden, ein interressanter Bodenseefund. JH. Ver. Württ. xlviii, pp. 238-247.

Describes a lower canine of Sus scrofu exhibiting a pathological peculiarity in structure.

†Chæromorus pygmæus, n. sp., Depéret, Arch. Mus. Lyon, v, p. 87, pl. i, Miocene, Grive-St-Alban.

#### 7.—CETACEA.

- BEAUREGARD, H. Note sur Deux Échouements Récents de Balænoptera musculus. CR. Soc. Biol. 1892, pp. 202 & 203.
- Beneden, P. J. van. Un Cétacé Fluviatile d'Afrique. Mém. Ac. Belg. (3) xxiii, pp. 350-355, 1 pl.

Note on Sotalia teuszii, with figure of skull.

- BOUVIER, E. L. Sur un Échouement d'Hyperoodon à l'entree de la Baie de Carentan. CR. Soc. Biol. 1892, pp. 835-838.
- ——. Observations Anatomiques sur l'Hypercodon rostratus. Ann. Sci. Nat. (7) xiii, pp. 259-320, pls. vii & viii.

Describes, in detail, the external characters and visceral anatomy of a recently stranded specimen.

CAPELLINI, G. Un Delfinide Miocenico, Ossia il Supposto Uomo Fossile di Acquabona presso Arcevia nelle Marche. Atti (Mem. Rend.) Acc. Rom. (5) i, pp. 325-330.

It is considered that the specimen referred to belongs probably either to a Schizodelphis or a Squalodon.

Gorjanovic-Kramberger, D. O. Fosilnih Cetaceih Hrvutske i Kranjske. (De Fossilibus Cetaceis Croatiæ et Carneoliæ.) Rad jugoslav. akad. cxi, pp. 1-21, pls. i-iii.

A memoir in Croatian on fossil Cetaceans from Croatia and Carniola. The specimens are referred to Champsodelphis, Platanista, Delphinapterus, and Mesocetus; the two new species being mentioned below.

KÜKENTHAL, W. Mittheilungen über den Carpus des Weisswals (Die Bildung des Hamatums und das Vorkommen von Zwei und Drei Centralien). Morph. JB. xix, pp. 56-64, pl. iii.

The occurrence of two centralia is noticed in both *Delphinapterus* and *Monoceros*; and the morphological bearing of this feature, as well as the presence of five distal carpalia, fully discussed.

- Ichthyosaurier und Wale. JB. Mineral. 1892, i, pp. 161-166.

The analogy between Ichthyosaurs and Whales is traced; and as it is probable that the former have descended from armoured reptiles, so the latter (from the discovery of traces of scutes with a fossil Dolphin and Zeuglodon) may have taken origin from Mammals furnished with a dermal armour.

- ---. Sotalia teuszii, n. sp., ein Pflanzenfressender (?) Delphin aus Kamerun. Zool. Jahrb. Abth. f. Syst. vi, pp. 442-446, pl. xxi.
- LYDEKKER, R. On Zeuglodont and other Cetacean Remains from the Tertiaries of the Caucasus. P. Z. S. 1892, pp. 558-564, pls. xxxvi-xxxviii.

The specimens are from Eocene beds, and are referred to a new species of Zeuglodon, and to two representatives of the Platanistide, one of which is regarded as indicating a new genus, while the other is not named. It is considered that the specimens help to confirm the view as to the Cetacean nature of Zeuglodon (vide infrů).

Moreno, F. P. Noticias sobre algunos Cetáceos Fósiles y Actuales de la República Argentina. Rev. Mus. la Plata, iii, pp. 381-398, pls. viii-xi.

Describes a recent Cetacean referred to a new species of Lageno-rhynchus, and two fossil crania provisionally assigned to the Delphinida (q. r.).

POUCHET, G. Note sur la Baleine Observée par Nearque. CR. Soc. Biol. 1892, pp. 422 & 423.

Concludes that the Whale seen by Nearchus in the Persian Gulf was a Megaptera.

- & BEAUREGARD, H. Note sur l'Ambre Gris. T. c. pp. 588-590.
- —. Recherches sur le Cachalot. N. Arch. Mus. (3) iv, pp. 1-90, pls. i-xii.

A continuation of the elaborate monograph quoted in Zool. Rec. xxvi, Mamm. p. 48; the present part dealing with the anatomy of the head and brain.

1892. [vol. xxix.]

- Southwell, T. Notes on the Seal and Whale Fishery. Zool. (3) xii, pp. 100-105.
- TURNER, SIR W. The Lesser Rorqual (Balanoptera rostrata) in the Scotch Seas, with observations on its Anatomy. P. R. Soc. Edinb. xix, pp. 36-75, 2 pls.

Describes a female stranded at Granton in 1888, and tabulates all the examples hitherto recorded from the Scotch coasts.

—... Notes on some of the Viscera of Risso's Dolphin (Grampus griseus).
J. Anat. Phys. (2) vi, pp. 258-270.

## a. PHYSETERIDÆ.

- Mesoplodon bidens (sowerbiensis). FISCHER, C.R. cxiv, pp. 1283-1286, describes the osteology of a specimen stranded at Cap Breton, August, 1888. TEGETMEIER, Field, lxxx, p. 1003, records the stranding of another specimen on the Norfolk coast, Dec. 18th, 1892.
- Hypercodon rostratus: BOUVIER'S notes, quoted in Zool. Rec. xxviii, Mamm. p. 48, translated in Ann. N. H. (6) ix, pp. 484-486.

#### b. Platanistida.

- † Iniopsis caucasica, n. g. & sp., Lydekker, P. Z. S. 1892, p. 562, Eocene, Caucasus.
- †Platanista croatica, n. sp., Gorjanovic-Kramberger, Rad jugoslav. akad. cxi, p. 12, Tertiary, Croatia.
- † Champsodelphis carniolicus, n. sp., Gorjanovic-Kramberger, t. c. p. 6, Tertiary, Carniola.

#### c. DELPHINIDE.

- Grampus grissus: Sir W. Turner, P. Phys. Soc. Edinb. xi, pp. 1-6, describes one of six examples caught at Hillswick, Shetland, September, 1889. Another captured on the Solway, September 30th, 1892, recorded in Land and Water, liv, p. 405.
- Delphinus dussumieri, n. n., Blanford, Mammals of British India, p. 588 (1891), for D. longirostris, Dussumier, nec Gray. [Omitted from Zool. Rec. xxviii.]
- Sotalia teuszii, n. sp., KÜKENTHAL, Zool. JB. vi, pp. 442-446, pl. xxi, Cameroon district. Believed to be herbivorous.
- Lagenorhynchus floweri, n. sp., Moreno, Rev. Mus. La Plata, iii, p. 383, pla. viii & ix.

- † Mesocetus poucheti, n. g. & sp., Moreno, t. c. p. 393, pl. x, Tertiary, Argentina.
- † Notocetus vanbenedeni, n. g. & sp., Moreno, t. c. p. 395, ibid.
- †Orcella fluminalis, identified with O. brevirostris; Thomas, Ann. Mus. Genov. (2) x, p. 39.

## d. †Zeuglodontidæ.

†Zeuglodon caucasicus, n. sp., Lydekker, P. Z. S. 1892, p. 559, Eoceue, Caucasus.

## 8. SIRENIA.

LYDERKER, R. On a remarkable Sirenian Jaw from the Oligocene of Italy, and its bearing on the Evolution of the Sirenia. P. Z. S. 1892, pp. 77-83, woodcut.

Describes part of an upper jaw of a Sirenian from the Tertiaries of Northern Italy, containing milk-teeth. As these teeth showed a masked selenodont structure, it is urged that the specimen indicates the descent of the Sirenia from Selenodont Artiodactyle-like Ungulates. It was incidentally shown that Halitherium veronense, Zigno, from the same deposits, belongs to Prorastoma, Owen.

WALDEYER, W. Ueber den feineren Bau des Magens und Darmkanales von Manatus americanus. SB. Ak. Berlin, 1892, pp. 79-85.

#### 9. EDENTATA.

Ballowitz, E. Das Schmelzorgan der Edentaten, seine Ausbildung im Embryo, und die Persistenz seines Keimrandes bei dem erwachsenen Thier. Arch. mikr. Anat. zl, pp. 133–156, pls. vi & vii.

The existence of an enamel-organ in the embryo of *Dasypus* is confirmed; and it is now shown that a portion of this persists as an epithelial ring at the base of the pulp-cavity.

Rose, C. Ueber rudimentare Zahnanlagen der Gattung Mania. Anat. Anat. Ana. vii, pp. 618-622.

Records rudimentary dental structures in embryoes of Manis tricuspis and M. jevanica.

## +MEGATHERLIDE.

For presence of enamel in teeth of earlier forms, see Ameghino, supra, pp. 2 & 3.

#### +GLYPTODONTIDÆ.

The alleged existence of a plastron is stated by AMEGHINO (supra, p. 3) to be incorrect.

## 10.—MARSUPIALIA.

For list of Australian forms, see OGILBY, suprà, p. 12.

COPE, E. D. On the Habits and Affinities of the New Australian Mammal, Notoryctes typhlops. Am. Nat. xxvi, pp. 121-128, pl. ix.

Concludes that the affinities of this animal are rather with Insectivores than Marsupials, and that its nearest ally is probably *Chrysochloris*.

DUN, W. S. Notes on the Teeth known as Sceparnodon ramsayi, Owen (Phascolonus gigas, Lydekker). Rec. Geol. Surv. N.S.W. iii, pp. 25–28, pl. vi.

New specimens from Bingera serve to disprove some of the views of De Vis as to the lower incisors of this type; and thus leave it still probable that Sceparnodon may be identical with Phascolonus.

GADOW, H. On the Systematic Position of Notoryctes typhlops. P. Z. S. 1892, pp. 361-370, figs.

After an exhaustive consideration of the hard and soft parts, the author concludes that *Notoryctes* is an undoubted polyprotodont Marsupial, and that its relationship is nearer to the *Dasyurida* than to the *Peramelida*.

LECHE, Biol. Fören. iv, pp. 1 & 2, likewise has a note on the affinities of the same genus.

Röse, C. Ueber die Zahnentwickelung der Beuteltiere. Anat. Ans. vii, pp. 639-650 & 693-707, figs.

From an examination of a series of feetal specimens, it is concluded that the teeth of Marsupials are developed in the same manner as in other Mammals; the anterior ones being diphyodont, while the true molars are produced as lateral outgrowths, with the exception of the last premolar. The germs of the replacing teeth are, however, aborted, and the permanent teeth consequently belong to the milk-series. In some cases, as in Didelphys, the last premolar takes a place among the milk-molars, without displacing one of the latter; although it generally succeeds a milk-tooth in the ordinary manner. It is also considered that the outermost upper incisor of Perameles, Macropus, and Phalangista belongs to the second series, which takes its place alongside those of the first series without any displacement. Further evidence is also obtained in support of the view that the premolars and molars are formed by the coalescence of three or more simple conical teeth.

SYMINGTON, J. On the Organ of Jacobson in the Kangaroo and Rock-Wallaby (Macropus giganteus and Petrogale penicillata). J. Anat. Phys. xxvi, pp. 371-374, pl. x.

It is shown that in *Ornithorhynchus* the cartilage of the organ in question, in the greater part of its extent, forms a complete tube, from which passes, internally, a turbinative process; no indications of such a process occurring in the *Metatheria* and *Eutheria*. Further, in *Ornithorhynchus* the organ, which is very large, extends forwards as well as backwards beyond its opening into the naso-palatine foramen, while in other Mammals the opening, which may be either into that foramen or into the nasal cavity, represents the anterior end of the organ.

—... The Cerebral Commissures in the Marsupialia and Monotremata.

T. c. pp. 22-84, pls. i-iii, figs.

It is concluded that the cerebral commissures of Monotremes and Marsupials may be distinguished from those of Eutherians by the following characters:—(1) The anterior commissure is as large as, and generally much larger than, any other transverse commissure of the cerebrum, and it unites the whole of the cortex of the two hemispheres, except the gyri dentati and hippocampi majores. (2) They have no true corpus callosum. (3) The superior transverse commissure is simply one for the gyri dentati and hippocampi majores. In the Eutherians, on the other hand, the following conditions obtain:—(1) The anterior commissure is much smaller than the other transverse commissures, and it never extends to the cortex on the upper surface of the cerebral hemispheres, or to that on the upper part of the inner surface of the same. (2) A true corpus callosum, and likewise (3) a hippocampal commissure are present.

ZIETZ, A. List of South Australian Species of Kangaroos and Wallabies. Tr. R. Soc. S. Austr. xv, pp. 18 & 19.

## a. PHALANGERIDÆ.

Acrobates pulchellus, n. sp., ROTHSCHILD, P. Z. S. 1892, p. 546, Island in Dutch New Guinea.

#### b. +AMPHITHERIIDE.

† Telacodon lavis, n. g. & sp., MARSH, Am. J. Sci. (3) xliii, p. 258, Cretaceous, Wyoming.

† Batodon tenuis, n. g. & sp., MARSH, l. c. ibid.

## c. INCERTÆ SEDIS.

†Stagodon validus, n. sp., MARSH, Am. J. Sci. (3) xliii, p. 256, Cretaceous, Wyoming.

† Thisodon pasianicus, n. g. & sp., Cope, Am. Nat. xxvi, pp. 758-760, Cretaceous, Wyoming. Founded upon an imperfect mandible, in which the angle is wanting, considered to belong either to a Marsupial or a Monotreme.

## 11. MONOTREMATA.

BAUR, Am. Nat. xxvi, pp. 72 & 435, records the absence of sygapophyses in the cervical vertebræ as a new fact. CARUS, t. c. p. 965, points out that Owen had long since noticed the fact of their absence, except in the atlas of *Echidna*.

HILL, A. The Cerebrum of Ornithorhynchus paradoxus. P. R. Soc. lii, pp. 163 & 164. [Abstract.]

Although the brain is distinctly Mammalian, it presents certain peculiar features connected with the rhinencephalon, hippocampus, and cerebral commissures, which have not hitherto been noticed. The olfactory bulb has a thin stalk, and is quite free from the frontal portion of the hemisphere. The hippocampus is very large, and lies entirely dorsal of the velum interpositum, while it is continued forwards to the extreme front of the brain, thus forming with the rhinencephalon a loop open in front. No commissural fibres cross the incisura pallii longitudinalis dorsally of the hippocampus; and it is concluded such fibres as do cross from one hemisphere to another do not represent a true corpus callosum. Exclusive of the incisura rhinalis and dentary fossa, which scarcely come under this category, the cortex is completely destitute of convolutions.

Turner, Sir W. The Cerebral Hemispheres of Ornithorhynchus paradozus. J. Anat. Phys. (2) vi, pp. 358-361.

The author finds that the rhinencephalon is more highly organized and larger in *Echidna* than in *Ornithorhynchus*; while the pallial part is on a larger and more complicated plan. Moreover, the cerebellum is likewise a larger organ, with its surface more minutely fissured in *Echidna*. Hence it is concluded that the brain of the latter is decidedly more highly organized than that of *Ornithorhynchus*.


#### a. Echidnida.

Echidna aculeata: remarks on young specimens; W. N. PARKER, Rep. Brit. Ass. 1891, p. 693.

Procehidna nigroaculeata, n. sp., ROTHSCHILD, P. Z. S. 1892, p. 545, Dutch New Guinea.

## 12. +MULTITUBERCULATA.

- †Cimolodon parvus, n. sp., O. C. Marsh, Am. J. Sci. (3) xliii, p. 254, Cretaceous, Wyoming.
- † Cimolodon agilis, n. sp., MABSH, t. c. p. 255, ibid.
- † Allacodon fortis, n. sp., MARSH, l. c. ibid.
- † Allacodon rarus, n. sp., MARSH, t. c. p. 256, ibid.
- †Oracodon conulus, n. sp., MARSH, l. c. ibid.



# III. AVES.

BY

## R. BOWDLER SHARPE, LL.D.

The year 1892 has been rendered remarkable by the publication of Dr. Gadow's Classification of Birds, by Professor Mivart's popular Introduction to Ornithology, and by the excellent work by Captain Charles Bendire on the Life-histories of American Birds. The zoo-geographical work of Dr. C. Hart Merriam and Mr. J. A. Allen, though based upon studies of the Mammalia, possesses the deepest interest for Ornithologists. The year will also probably be notable for the foundation in this country of the 'British Ornithologists' Club,' whose monthly 'Bulletin' contains a record of most important work, while the 'Ibis' is more flourishing than ever.

#### ARRANGEMENT OF RECORD.

I.—TITLES OF SEPARATE WORKS, AND OF THE MORE IM-PORTANT PAPERS PUBLISHED IN PROCEEDINGS OF SOCIETIES, &c.

II .- FAUNISTIC.

III.—Anatomy and Miscellaneous.

IV.—Systematic.

#### I.—TITLES.\*

Albarda, N. Ueber das Vorkommen seltener Vögel in den Niederlanden. J. f. O. 1892, pp. 417-431.

ALESSI, S. Briefliches aus Tunisien. T. c. pp. 313-317.

ALFARO, A. Resena de las principales Aves que habitan la parte superior del Volcán de Poás. An. Mus. Costa Rica, iii, pp. 160 & 161.

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

<sup>1892. [</sup>vol. xxix.]

- ALLEN, C. S. [See Pandiones, p. 39.]
- ALLEN, J. A. Notice on some Venezuelan Birds collected by Mrs. H. H. Smith. Bull. Am. Mus. iv, pp. 51-56.
  - [Tanagridæ, p. 52; Tyrannidæ, p. 61; Piciformes, p. 49.]
- On a Collection of Birds from Chapada, Matto Grosso, Brazil, made by Mrs. H. H. Smith. Part II. Tyrosnido. T. c. pp. 330-350. [Part I, cf. Zool. Rec. xxviii, Aves, p. 2.]
- ---. The Geographical Distribution of North American Mammals. T. c. pp. 199-243, pls. v-viii [maps].
- ---. [See also Ralliformes, p. 30; Piciformes, p. 49.]
- Anonymous. Fourth Supplement to the A. O. U. Check-list of North-American Birds. Auk, ix, pp. 105-108.
- Anthony, A. W. Birds of South-western New Mexico. Auk, ix, pp. 357-369.
- APLIN, O. V. Notes on Birds seen in Switzerland. Zool. 1892, pp. 8-14 & 65-73.
- Supplementary Notes on the Birds of Oxfordshire. T. c. pp. 241-258.
- —. Notes from a Winter Journal, 1890 & 1891. Mid. Nat. xv, pp. 40-43 & 58-62.
  - [Emberizinæ, p. 53; Laniidæ, p. 55; Cinclidæ, p. 58.]
- ATTWATER, H. P. List of Birds observed in the vicinity of San Antonio, Bexar County, Texas. Auk, ix, pp. 337-345.
- Austen, E. E. On Bird-life at Heidelberg. Rep. Rugby Soc. 1891, pp. 36-43.
- BAKER, E. C. S. [See Pycnonotida, p. 59; Troglodytida, p. 58.]
- Baldamus, E. Das Leben der europaischen Kuckucke, nebst Beiträgen zur Lebenskunde der übrigen parasitischen Kuckucke und Stärlinge. Berlin: 1892, 224 pp., viii pls.
- BARBOZA DU BOCAGE, J. V. Additions et Corrections à l'Ornithologie d'Angola. I. J. Sci. Lisb. (2) viii, pp. 248-264.
  - [Irrisoride, p. 43; Nectariniida, p. 54.]
- —. Aves do Sertão de Benguella. J. Sci. Lisb. (2) vii, pp. 157-172. [Turdidæ, p. 57.]
- ...... Subsidios para a Fauna da Guiné Portugueza. T. c. pp. 179-184.
- BAUR, G. Besuch der Galapagos-Inseln. Biol. Centralbl. xii, pp. 221-250.
- BECKWITH, W. Shropshire Birds. Tr. Shropshire Soc. 1892, pp. 183-198.
- Beddard, F. E. Animal Coloration: an account of the principal facts and theories relating to the colours and markings of animals. London: 8vo, 1892, pp.
  - [See also Plotide, p. 36.]

- BENDIRE, C. Life Histories of North American Birds, with special reference to their breeding, habits, and eggs. U. S. Nat. Mus. Special Bulletin, No. i. Washington: 1892, 4to, pp. viii & 416, pls. i-xii.
- [Galliformes, p. 25; Columbiformes, p. 28; Accipitriformes, p. 37; Striges, p. 39.]
- Berlepsch, H. von. Die Vernichtung unserer Vögel im Süden und der daraus resultirende Schaden. C.R. ii, Congr. Orn. Intern. ii, pp. 179-189.
- ---. Die Vögel der Insel Curação, nach einer von Herrn Ernst Peters daselbst angelegten Sammlung. J. f. O. 1892, pp. 61-104.
  - [Carebida, p. 52; Falconida, p. 39; Odontophorina, p. 28.]
- ---. [See also BOUCARD & BERLEPSCH.]
- ---- & STOLZMANN, J. Résultats des recherches ornithologiques faites au Pérou par M. Jean Kalinowski. P. Z. S. 1892, pp. 371-411.
- [Tanagridæ, p. 52; Icteridæ, p. 51; Tyrannidæ, p. 62; Dendrocolaptidæ, p. 62.]
- & —. [See also Crypturiformes, p. 25; Trochili, p. 62; Furnariinæ, p. 62.]
- Besse, C., & St. Maurice, Abraye de. Liste des Oiseaux palmés du Valais. Bull. Soc. Murith. livr. xvi-xviii, pp. 36-42 (1890).
- Bird, M. Bird Life of the Norfolk Broads. Field Club, iii, pp. 1-4, 20-24, 36-60, 52-55, 65-69, 86-88, 104-108, 116-120, 133-138, 151-154, 164-167, & 177-180.
- BLAAUW, F. E. [See Platalea, p. 34.]
- BLAKISTON, T. W. [THE LATE]. Obitnary Notice. Ibis, 1892, pp. 190 & 191; Auk, ix, pp. 75 & 76.
- BLASIUS, R. Zur Oologie. C.R. ii, Congr. Orn. Intern. ii, pp. 148-152.
- BLOMEFIELD, L. Remarks on the Distribution and Movements of British Animals and Plants, in Past and Present Times, as instanced in the Nightingale and some other cases. P. Bath Nat. Hist. Club, 1892, pp. 185-199.
- BOCAGE. [See BARBOZA DU BOCAGE.]
- BOLAM, G. Ornithological Notes. Hist. Berwick Nat. Club, 1890-91, pp. 345-368.
- Bolles, F. [See Piciformes, p. 49.]
- BONVALOT, G. De Paris au Tonkin a travers Le Tibet-Inconnu. Paris: 8vo, 1892, Oiseaux, pp. 488 & 489.
- BOUCARD, A. Aves colectadas en Costarica. An. Mus. Costa Rica, iii, pp. 141–159.

- [Boucard, A.] [See also Phasiani, p. 26; Trochili, p. 46.]
- ---. H. W. Bates [Obituary Notice]. Humming Bird, ii, pp. 26-31, with portrait.
- BOUCARD, A., & BERLEPSCH, H. von. List of Birds collected by M. Hardy, at Porto Real, Brazil, with description of one supposed new species. Humming Bird, ii, pp. 41-45.

[Formicariidæ, p. 63.]

- BREWSTER, W. [See Mniotiltidæ, p. 53.]
- Brooks, W. E. A few remarks on Mr. Oates's "Birds of British India." Ibis, 1892, pp. 59-62.
- [Corvidæ, p. 47; Regulidæ, p. 55; Sylviidæ, p. 57; Muscicapidæ, p. 61; Turdidæ, p. 57; Motacillidæ, p. 53; Alaudidæ, p. 53.]
- BRUSINA, S. Vögel Croato-Serbische mit Berücksich des übrig slav. Südens. Verbreitung für eine croato-serbische Ornithologie. Fortzetzung. Denkschr. Serb. K. Akad. xii, 168 pp.
- —. K. Ornitologyi Kotora i Crne Gore. Glasnik Naravosl. Druž. vi, pp. 59-81.
- Ornitoložke bilježke za hrvatsku faunu. Op. cit. iii, pp. 129-150 (1888).
- ---. Nore ornitološke bilješke. Op. cit. iv, pp. 205-221 (1889).
- BRUTTINI, A. Appunti storici sulla Falconeria. Riv. Ital. Sci. Nat., Ann. xii (Boll.), pp. 89, 90, 100-102, 107-109, 121, & 122.
- BRYANT, W. E. A Check list of the Water Birds of California. Zoe, iii, pp. 135-140.
- BÜCHNER, E. Rapport sur le deuxième Congrès Ornithologique International de Budapest. 8vo, pp. 1-50.
- Buckley, T. E. Contributions to the Vertebrate Fauns of Sutherland and Caithness. Ann. Scott. Nat. Hist. i, pp. 156-168.

[See also Harvie-Brown & Buckley.]

- Buda, A. von. Unsere seltenere Gäste im Comitate Hunyad. C.R. ii, Congr. Orn. Intern. ii, pp. 145-148.
- Buller, W. L. Notes and Observations on New Zealand Birds. Tr. N. Z. Inst. xxiv, pp. 64-74, pl. xiv.
- —. Further Notes and Observations on certain species of New Zealand Birds. T. c. pp. 75-92.

[Apterygiformes, p. 25.]

- BURMEISTER H. [THE LATE]. Obituary Notice. Ibis, 1892, pp. 581 & 582.
- BUTLER, A. W. [See Psittaciformes, p. 47.]
- BÜTTIKOFER, J. On the Collections of Birds sent by the late R. T. Demery, from the Sulyman River (W. Africa). Notes Leyd. Mus. xiv, pp. 19-30.

[Ploceida, p. 51.]

- [BÜTTIKOFER, J.] On a Collection of Birds from the Islands of Flores, Sumba, and Rotti. T. c. pp. 193-206.
- [Sylviidæ, p. 58; Diceidæ, p. 54; Meliphagidæ, p. 54; Ploceidæ, p. 51; Muscicapidæ, p. 61.]
- ----. Europäische Zug-vögel als Gäste in Liberia. C.R. ii, Congr. Orn. Intern. ii, pp. 172-176.
- —. J. P. van Wickevoort Crommelin [Obituary Notice]. Orn. Jahrb. iii, pp. 76-79.
- —... [See also Plocsidæ, p. 51; Sylviidæ, p. 57; Turdidæ, p. 57; Muscicapidæ, p. 61.]
- CABANIS, J. [See Pterocletes, p. 28.]
- CAMPBELL, A. G. Notizen ueber Australische Oologie. C.R. ii, Congr. Orn. Intern. ii, pp. 155-158.
  - [Menuræ, p. 49; Sterninæ, p. 32.]
- ---. [See also Paradiseida, p. 50.]
- CAMPBELL, C. W. A List of the Birds collected in Corea. Ibis, 1892, pp. 230-248, pl. v.
  - [Picida, p. 49; Timeliida, p. 59; Sylviida, p. 57; Muscicapida, p. 62]
- Cannaviello, E. Casi d'Albinismo nella Provincia di Napoli. Boll. Ital. Sci. Nat. xii, p. 72.
- CAPEK, V. [See Cuculidæ, p. 46.]
- CHAPMAN, F. M. Notes on Birds and Mammals observed near Trinidad, Cuba, with remarks on the origin of West Indian Bird Life. Bull. Am. Mus. iv, pp. 279-330.
- [Ralliformes, p. 30; Columbiformes, p. 29; Buteoninæ, p. 38; Tyrannidæ, p. 62; Icteridæ, p. 51; Mniotiltidæ, p. 53.]
- —. A Preliminary Study of the Grackles of the Subgenus Quiscalus. Bull. Am. Mus. iv, pp. 1-20, map.
  - [Icteridæ, p. 51.]
- \*\*CHENEY, S. P. Wood Notes Wild, Notations of Bird Music. Boston, U.S.A.: 1892, 12mo, i-xvi & 1-261 pp.
- CHERNEL, S. VON. [See Scolopacida, p. 34.]
- CHERRIE, G. K. A preliminary list of the Birds of San José, Costa Rica. [Zool. Rec. xxviii, Aves, p. 7.] Auk. ix, pp. 21-27, 247-251, 322-329.
  - [Tanagridæ, p. 52; Icteridæ, p. 51.]
- ——. Descripcion de tres Especies nuevas para la Avifauna Costaricense. An. Mus. Costa Rica, iii, pp. 135-137.
  - [Tanagridæ, p. 52; Formicariidæ, p. 63.]
- \_\_\_\_. [See also Tyrannida, p. 62.]
- CHIEVITZ, J. H. Sur l'existence de l'area centralis retine dans les vertébrès. Overs. Dan. Selsk. 1891, Aves, pp. 246-252.

- Church, A. H. Researches on Turacin, an animal pigment containing copper. Phil. Tr. clxxxviii, pp. 511-536. [Musophagida, p. 47.]
- CLARKE, W. EAGLE. [See Gruiformes, p. 84; Lariformes, p. 32; Muscicapida, p. 60.]
- COCKERELL, T. The West Indian fauna in South Florida. Nature, xlvi, pp. 458-459.
- COLLETT, R. On a collection of Birds from Tongoa, New Hebrides. Forh. Selsk. Chr. 1892, pp.
- —. Das Leben der Vögel im Arctischen Norwegen. C.R. ii, Congr. Orn. Intern. ii, pp. 21-39.
- COPE, E. D. Occurrence of Bird remains in Pueres Series, New Mexico. Tr. Am. Phil. Soc. xvi, p. 302.
- CORDEAUX, J. On the Migration of Birds as observed on the east coast of Great Britain. C.B. ii, Congr. Orn. Intern. ii, pp. 165-167.
- —. A recent visit to the Farne Islands. Zool. 1892, pp. 293-303.
- —. Migration in the Humber District in the Autumn of 1892. T. c. pp. 417-420.
- ---. [See also Sylviidæ, p. 57.]
- CORY, C. B. Catalogue of West Indian Birds, containing a list of all species known to occur in the Bahama Islands, the Greater Antilles, the Caymans, and the Lesser Antilles, excepting the Islands of Tobago and Trinidad. Boston: 1892, roy. 8vo, 163 pp., with map. [Columbiformes, p. 29; Striges, p. 40; Fringillidæ, p. 52; Vireonidæ, p. 56; Carebidæ, p. 52.]
- —. A list of Birds taken on Maraguana, Watling's Island, and Inagua, Bahamas, during July, August, September, and October, 1891. Auk. ix, pp. 48 & 49.
- —. Remarks on a Collection of Birds made by W. W. Brown, jun., on Mona and Porto Rico during February and a part of March, 1892. T. c. pp. 228 & 229.
- ---. [See also Picidæ, p. 49.]
- COUVREUR, E. Influence du Pneumogastrique sur les fonctions circulatoires (cœur, pression sanguine, effets vasomoteurs), chez les Oiseaux, et comparaison avec les Mammiféres. Ann. Soc. L. Lyon (n.s.) xxxix, pp. 11-24.
- —. Influence du Pneumogastrique sur la sécrétion urinaire chez les Oiseaux et comparaison avec les Mammifères. T. c. pp. 25-40.
- —. Sur la Pneumogastrique des Oiseaux. Ann. Univ. Lyon, pp. 1-104, pls. i-iii.
- CRAIG, A. Stray notes on the Birds of Anglesea. Tr. Edinb. Nat. Soc. ii, p. 361, 1889-90.

- [CRAIG, A.] Birds of the Great Glen, Inverness-ahire. Tr. Edinb. Nat. Soc. pp. 439-457, 1890-91.
- CREAGHE-HAWARD, L. Notes on some East Anglian Birds. Tr. Leicester Soc. ii, pp. 538-554.
- CROMMELIN, J. P. VAN WICKEVOORT [THE LATE]. Obituary Notice. Ibis, 1892, p. 352.

[See also BÜTTIKOFER, J.]

- CROWLEY, P. [See Paradiseidæ, p. 50; Ptilonorhynchidæ, p. 50.]
- CZATO, J. VON. [See Accipitriformes, p. 37.]
- CZÝNK, E. VOS. Der Herbstaug, 1891, im Fogaras-cher Comität. Orn. Jahrb. iii, pp. 191-195.
- DALGLEISH, J. J. [See Procellariida, p. 31.]
- Dareste, C. Note sur l'evolution de l'embryon de la poule soumis pendant l'incubation à un mouvement de rotation continu. C.R. cxv, pp. 137 & 138.
- DAVIES, S. A. A Naturalist's Ramble in Swedish Lapland. Zool. 1892, pp. 81–87.
- Davison, W. R. Descriptions of some new species of Birds from the Eastern Coast of the Malayan Peninsula. Ibis, 1892, pp. 99-108.
- [Campophagidæ, p. 60; Muscicapidæ, p. 60; Timeliidæ, p. 59; Eulabetidæ, p. 50.]
- DE LA TOUCHE, J. D. On Birds collected or observed in the vicinity of Foochow and Swatow in South-eastern China. Part I [with notes by H. H. SLATER]. Ibis, 1892, pp. 400-430.
- —. Part II. Id. t. c. pp. 477-503, pl. xii.

## [Ralliformes, p. 80.]

- DÉPERET, C. Sur la faune d'Oiseaux pliocènes du Roussillon. C.R. cxiv, pp. 690-692. [Corvidæ, p. 48; Perdicidæ, p. 27; Anatidæ, p. 35.]
- DE Vis, C. W. Report on the Zoological Gleanings of the Administration during the year 1890-91 [Annual (Parliamentary) Report on British New Guinea]. Brisbane: 1892, App. cc, pp. 93-97, cum tab. [Muscicapidæ, p. 56; Meliphagidæ, p. 54; Timeliidæ, p. 59; Ptilosorhynchidæ, p. 50; Laniidæ, p. 56.]
- —. Zoology of British New Guinea. Ann. Queensland Mus. ii, pp. 4-11.
- [Muscicapida, p. 58; Timeliida, p. 59; Meliphagida, p. 54; Paramythiida, p. 50; Sylviida, p. 58; Laniida, p. 56.]
- —... Residue of the Extinct Birds of Queensland as yet detected. P. Linn. Soc. N.S.W. (2) vi, pp. 437–456, pls. xxiii & xxiv,
- [Casuariiformes, p. 24; Apterygiformes, p. 25; Aquilinæ, p. 38; Ralliformes, p. 31; Pelargiformes, p. 34; Plataleæ, p. 34; Pelecani, p. 36.]

DISTANT, W. L. A Naturalist in the Transvaal. London: 8vo, xvi & 277 pp., 5 pls.

[Aves, pp. 163-169.]

- DIXON, L. The Migration of Birds. An attempt to reduce Avian Flight to Law. London: 1892, 8vo, viii & 300 pp.
  [See Nature, xlvi, pp. 169-172.]
- DRESSER, H. E. [See Laniida, p. 55; Parida, p. 55; Cinclida, p. 58.]
- DUBALEU, P. E. Monographie raisonnée des Oiseaux, observés dans les départements des Laudes, de la Gironde, du Gers, des Basses-Pyrènées, des Hautes-Pyrènées, et sur le Golfe de Gascogne. Bull. Soc. Borda-Dax, xvii, pp. 173-191.
- DUCHAUSSOY, H. [See Hirundinidæ, p. 61.]
- DUBOIS, A. Établissement d'une classification Internationale. C.R. ii, Congr. Orn. Intern. ii, pp. 92-94.
- Dugès, A. Instrucciones para colectores de Aves. Nat. Mex. (2) ii, pp. 175-179, pl. ix.
- D'Urban, W. S. M., & Mathew, M. A. The Birds of Devon. London: 1892, 8vo, i-lxxxvii & 1-460 pp.
- DWIGHT, I. Summer Birds of the Crest of the Pennsylvania Alleghanies. Auk, ix, pp. 129-142.
- Dybowski, J., & Oustalet, E. L'extreme sud Algérien. Arch. Miss. Sci. i, 1891, pp. 354-359.
- EGLINGTON, W. [See NICOLLS & EGLINGTON.]
- ELLIOT, D. G. The inheritance of acquired characters. [Presidential address to the A. O. U. Nov. 18, 1891.] Auk, ix, pp. 77-104.
- ----. Hybridism, and a description of a Hybrid between Anas boschas and A. americana. T. c. pp. 160-166.

  [Anseriformes, p. 35.]
- EVANS, A. H. [See WILSON & EVANS.]
- Evans, W. Some further notes on the periods occupied by Birds in the Incubation of their Eggs. Ibis, 1892, pp. 55-58. [Cf. Zool. Rec. xxviii, Aves, p. 10.]
- Fabani, C. Nota intorno ad alcuni casi di anomalie nel colore degli uccelli. Riv. Ital. Sci. Nat., Ann. xii (Boll.), pp. 2 & 3.
- —. Spiegazione intorne alle cause dell' isabellismo, melanismo ed albinismo negli uccelli. T. c. pp. 3 & 4.
- —. Epoche, mutazioni et varieta nel Canto degli uccelli. T. c. pp. 35-39.
- Teoria delle somiglianze nel Canto degli uccelli. T. c. pp. 90 & 91.

- [FABANI, C.] Studio sul mimetismo nel colorito degli uccelli. Riv. Ital. Sci. Nat., Ann. xii (Boll.), pp. 105, 106, & 122-124.
- FATIO, V. [See STUDER & FATIO.]
- FAUVEL, A. La Faune du Chan-toung. Rev. Quest. Sci. ii (i) pp. 455-492.
  - Oiseaux, pp. 466-476.
- FERGUSON, R. S. [See MACPHERSON, H. A.]
- FICALBI, E. Sulla architettura istologica di alcuni peli degli uccelli con considerazioni sulla filogenia dei peli e delle penne. Atti [Mem.] Soc. Tosc. xi, 1890, pp. 227-266, pl. ix.
- FIELD, H. C. [See Dinornithiformes, p. 25.]
- FINN, F. An Evolutionist in East Africa. Nat. Sci. i, pp. 744-748.
- FISCHER-SIEGWART, H. L'albinisme chez les Oiseaux. Arch. Sci. Nat. (3) xxviii, pp. 555-559.
- FLOERICKE, C. Winterbeobachtungen 1891-92. Orn. Jahrb. iii, pp. 182-189.
- —. Zur charakteristik der schlesischen Vogelwelt. J. f. O. 1892, pp. 151-170.
- Versuch einer avifauna der Provinz Schlesien. Lief i, 8vo, Marburg, 1892.
- XII Jahresbericht (1887) des Ausschusses für Beobachtungsstationen der Vögel Deutschlands. J. f. O. 1892, pp. 237-253.
- FORBES, H. O. Preliminary notice of additions to the extinct Avifauna of New Zealand. Tr. N. Z. Inst. xxiv, pp. 185-189.
  - [Accipitrina, p. 38; Ralliformes, p. 30; Anseriformes, p. 35; Phalacrocoraces, p. 36; Dinornithiformes, p. 25.]
- ——. Aphanapteryx and other remains in the Chatham Is. Nature, xlvi, pp. 252 & 253.
- ---. [See also Ralliformes, p. 30; Corvidæ, p. 49.]
- FOSTER, L. S. Bibliographies of American Naturalists: IV. The Published Writings of George Newbold Lawrence, 1844-1891. Bull. U. S. Nat. Mus. No. 40, pp. xi-124.
- FRITSCH, A. Die Vogelwelt am Gatterschlager Teiche bei Neuhaus in Böhmen. Orn. Jahrb. iii, pp. 19-23.
- —. Die Vogelwelt am unter-Porernitzer Teiche bei Prag. T. c. pp. 30-32.
- FROHAWK, F. W. [See Ralliformes, p. 30.]
- FÜRBRINGER, M. L. Stejneger's Vogelsystem und Th. Studer's Untersuchungen über die Embryonalentwickelung der antarktischen Vögel. J. f. O. 1892, pp. 137-151.

- [FÜRBRINGER, M.] Untersuchungen sur Morphologie, &c., der Vögel. [Review by F. Helm.] Biol. Centralbl. xii, pp. 146-157.
- Gadow, H. On the Classification of Birds. P. Z. S. 1892, pp. 229-256. [See also Opisthocomi, p. 29.]
- GITKE, H. Ornithologisches aus Helgoland. Orn. Jahrb. iii, pp. 201 & 202.

[Platalece, p. 34; Fringillidee, p. 53; Alaudida, p. 53.]

- ——. [See also Seeвони, H.]
- GIBBS, M. Acorn-eating Birds in Michigan. Nature, xlvi, p. 495.
- GIGLIOLI, H. H. L'Avigeographia. C.R. ii, Congr. Orn. Intern. ii, pp. 159-164.
- & Manzella, A. Iconographia dell' Avifauna Italica, ovvero Tavole illustranti le specie di Uccelli che trovansi in Italia con brevi descrizione e note. Prato (Toscano): 1892, fol., pts. xlvii-l.
- [Accipitriformes, p. 39; Pandiones, p. 39; Striges, p. 40; Cuprinulgi, p. 44; Cypseli, p. 44; Laniidæ, p. 56; Certhiidæ, p. 54; Sittidæ, p. 54; Turdidæ, p. 57; Muscicapidæ, p. 60.]
- Godman, F. D., & Salvin, O. Biologia Centrali-americana; or Contributions to the Knowledge of the Fauna and Flora of Mexico and Central America. [Zool. Rec. xxviii, Aves, p. 12.] London: 1892, 4to, Zoology, pts. xcix, c, cii-cv, pp. 193-368, pls. 49-57.

[Formicariidæ, p. 62; Trochili, p. 45.]

- ----. [See also Salvin & Godman.]
- GOERING, A. Zur kenntnis des Pflauzen- und Thier-lebens der Paramos. MT. Osterlande (n.s.) v, pp. 249-256.
- GRANT, W. R. OGILVIE. [See OGILVIE-GRANT.]
- Gurney, J. H. Ornithological Notes from Norfolk. Zool. 1892, pp. 398-401.
- HAIGH, G. H. CATON. The Motions of Divers on Land. T. c. p. 226.
- HALL, H. G. Odd-coloured eggs. C.R. ii, Congr. Orn. Intern. ii, pp. 153 & 154.
- Hänisch, H. Aus Dalmatien. Orn. Jahrb. iii, pp. 27-30. [Scolopacidæ, p. 33.]
- —. Ornithologische Mittheilungen aus dem unteren Narentathale. T. c. pp. 55-60.

[Ardeidæ, p. 34.]

- [See also Otides, p. 34.]
- Hamilton, A. [See Ralliformes, p. 30; Dinornithiformes, p. 25.]
- HARTERT, E. Notes on the Caprimulgiida. Ibis, 1892, pp. 274-288, pl. vi.

[Caprimulgi, p. 44.]

- [HARTERT, E.] Catalogue of the *Picaria*: in the Collection of the British Museum. *Caprimulgi* [q.v.], *Cypeeli* [q.v.]. [Vol. xvi of the Catalogue of Birds.] London: 8vo, pp. 434-654, pls. x-xiv.
  - [Caprimulgi, p. 44; Cypseli, p. 44; Podargi, p. 40; Steatornithes, p. 40.]
- ---... On the Birds of Eastern Prussia. Part 1. Ibis, 1892, pp. 353-372. [Sylviida, p. 57; Sittida, p. 54; Fringillida, p. 53; Corvida, p. 49.]
- ---. Part II. T. c. pp. 504-522.
- —. Trivial-namen und Notizen aus Schlesien. Orn. Jahrb. iii, pp. 32 & 33.
- —. [See also Striges, p. 40; Pioninæ, p. 47; Conurinæ, p. 47; Alaudidæ, p. 53; Tyrannidæ, p. 62.]
- HARTING, J. E. Hawks and Hawking. P. R. Inst. xiii, pp. 357-361.
- HARTLAUB, G. Ein Beitrag zur Ornithologie Chinas. Abh. Ver. Brem. xii, pp. 295-335.
- ---. [See also Ralliformes, p. 30; Muscicapida, p. 61.]
- HARVIE-BROWN, J. A., & BUCKLEY, T. E. A Vertebrate Fauna of Argyll and the Inner Hebridea. Edinburgh: 1892, Roy. 8vo, xxiv & 314 pp., map.

[See also Piciformes, p. 48.]

- HELM, F. [See FÜRBRINGER, M., & MEYER, A. B.]
- HATCH, P. L. Notes on the Birds of Minnesota, with specific characters. Minneapolis: 1892, 8vo, ix & 487 pp.
- HENKE, R. G. Auch einiges über Rackelwild und Hahnenfedrigkeit. J. f. O. 1892, pp. 170-177.
- HERBANO, A. L. [See Psittaciformes, p. 47.]
- Hewerson, H. B. Causes of Variation in Shape and Colour of Eggs. Tr. Leeds Nat. Club, ii, pp. 64-66 (1890).
- HILL, J. P. [See Howes & HILL.]
- Holland, A. H. Short Notes on the Birds of the Estancia Espartilla, Argentine Republic. Ibis, 1892, pp. 193-214.
- Homeyer, A. von. Ueber das Leben der Vögel in Central West Afrika.

  Denkwürdigkeiten meiner Reise nach und durch Angola, 1875.

  C.B. ii, Cong. Orn. Intern. ii, pp. 1-17.
- Howes, G. B., & Hill, J. P. Pedal Skeleton of the Dorking Fowl, with remarks on hexadactylism and phalangeal variation in the *Amniota*. J. Anat. Phys. xxvi, pp. 395-403.
- Hubbard, S. Additions to the Birds of the Gray's Harbour Region, Washington. Zoe, iii, pp. 140-144.
- Hudson, W. H. The Naturalist in La Plata. London: 1892, 8vo, viii & 388 pp.

HUTCHINSON, H. N. Extinct Monsters. A Popular Account of some of the larger forms of Animal Life. London: 1892, 8vo, i-xx & 1-254 pp.

[Hesperornithiformes, p. 31.]

- HUTTON, F. W. [See Dinornithiformes, p. 25.]
- IJIMA, J. Notes on a collection of Birds from Tsushima. J. Coll. Sci. Japan, v, pp. 105-128, pl. xii.
  [Piciformes, p. 49.]
- IRBY, L. H. British Birds, Key List. Second Edition. London: 1892, 8vo, viii & 70 pp.
- JACKSON, F. J. [See SHARPE, R. B.]
- James, H. Berkeley [THE LATE]. A New List of Chilian Birds, with a preface by P. L. Sclater. London: 8vo, 1892, viii & 16 pp.
- JOHNSON, F. O. Bird Notes from Alameda County. Zoe, iii, pp. 115-117.
- JOHNSON, F. P. Birds observed on a Fishing Expedition to South Trondhjem. Zool. 1892, pp. 267-269.
- KAISER, A. Zur Ornis der Sinaihalbinsel. Orn. Jahrb. iii, pp. 207-248.
- KEELER, C. A. Evolution of the Colours of North American Land Birds. Occas. papers Calif. Acad. iii, 362 pp., xix pls.. [Cf. Auk, xi, pp. 189-195.]
- A supposed new feather structure (Pseudopilum). Zoe, iii, p. 257.
- —. On the Natural History of the Farallon Islands. Zoology. T. c. pp. 144-165, pls. xviii-xxi.
- KERR, J. GRAHAM. On the Avifauna of the Lower Pilcomayo. Ibis, 1892, pp. 120-152, pl. iii.

[Piciformes, p. 48.]

- KERVILLE, H. G. DE. Faune de la Normandie. Fasc. iii. Columbidæ, Gallinaceæ, Grallæ, Palmipedes. Bull. Soc. Rouen, 1892, pp. 201-582, pl. i.
- KHLYEBNIKOV, V. Spisok Ptitz Astrakhanskoi ghubernii. Protok. obsch. estest. Kazan, 1890-91 (1891), No. 121, 10 pp.
- OKLEIN, E. The Etiology and Pathology of Grouse Disease, Fowl Enteritis, and some other Diseases affecting Birds. London: 8vo.
- KLEINSCHMIDT, O. Einige bemerkenswerte Notizen aus der Umgebung von Marburg a. d. Lahn (Sommer 1891 und winter 1891-92). Orn. Jahrb. iii, pp. 166-168.
- ——. Vögel des Grossherzogthums Hessen, insbesondere der Rheinebene bei Nierstein. T. c. pp. 195-212.

- Klug, K. Beiträge zur Kenntniss der Verdauung der Vögel insbesondere der Gänse. C.R. ii, Congr. Orn. Intern. ii, pp. 43-59, taf. i.
- KNUBLEY, E. P. Is Legislative Protection required for Wild Birds' Eggs? Ann. Scott. Nat. Hist. i, pp. 213-218.
- KCENIG, A. Zweiter Beitrag zur Avifauna von Tunis. J. f. O. 1892, pp. 266-312 & 330-416, taf. iii.
- KCENIG-WARTHAUSEN, BARON. Ueber späte Vogelbruten. JH. Ver. Württ, xlviii, pp. lxiv & lxv.
- ---. Färbung der Vogeleier. T. c. pp. lxiii & lxiv.
- KOEPERT, O. [See Sturnidæ, p. 50.]
- KOLOMBATOVIC, G. [See Anseriformes, p. 35.]
- KOLLIBAY, —. Ornithologisches aus Schlesien. J. f. O. 1892, pp. 317-328.
- Kosic, V. Gragja za dubrovačku nomenklaturu i faunu ptica. Glasnik. Naravosl. Druž. iii, pp. 118-128 (1888).
- LAWRENCE, G. N. [See FOSTER, L. S.]
- LAWRENCE, R. H. A Preliminary List of the Birds of the Gray's Harbour Region, Washington. Auk, ix, pp. 39-47.
- —... Further Notes on Birds of the Gray's Harbour Region, Washington. T. c. pp. 352-357.
- LEGGE, W. V. Protection of Tasmanian Owls. P. R. Soc. Tasm. 1889, p. 40.
- ----. Protection of the Cape Barren Goose. T. c. pp. 41 & 42.
- ----. [See also Limicola, p. 33.]
- LEGTARD, J. Sur la disparition ou l'extension de diverses espèces animales. CR. Ass. Fr. Sci. xx, Birds, pp. 563-565.
- LEEBODY, J. R. Notes on Birds of Lough Swilly. Irish Nat. i, pp. 173-177.
- LEVERKÜHN, P. Bericht über eine Reise nach Ungarn im Früjahre, 1891. C.R. ii, Congr. Orn. Intern. ii, pp. 210-238.
- Lewis, H. Notes on Birds observed in Hertfordshire during the Year 1891. Tr. Hertf. Soc. vii, pp. 62-68.
- LILFORD [LORD]. Coloured Figures of the Birds of the British Islands. Pts. xxi & xxii [cf. Zool. Rec. xxviii, Aves, p. 16].
- The species figured are mentioned under the headings of their respective families.
- ---. Notes on the Ornithology of Northamptonshire. Zool. 1892, pp. 201-210.
- —. Notes on the Birds of Northamptonshire. J. Northampt. Soc. v. pp. 41-53, 85-93, 131-141, 161-171, 178-1 6, 207-222, 293-302.
- \_\_\_\_. Op. cit. vi, pp. 1-7, 41-49, 69-76, 116-125, 162-171, 196-207, 262-266, & 269-277.

- [LILFORD (LORD).] Notes on the Birds of Northamptonshire. J. Northampt. Soc. vii, pp. 3-12, 58-63, & 140-146.
- ---. [See also Accipitriformes, p. 37; Hemipodii, p. 28.]
- Linstow, Dr. v. Beobachtungen an Vogeltänien. CB. Bakt. Parasit. xii, pp. 501-504.
- LOOMIS, L. M. A further Review of the Avian Fauna of Chester County, South Carolina [Zool. Rec. xxviii, Aves, p. 17]. Auk, ix, pp. 28-39.
- LORENZ, T. Die Vögel des Moskauer Gouvernements. Bull. Soc. Mosc. 1892, pp. 263-321.
- LORENZ VON LIBURNAU, L. Die Ornis von Oesterreich-Ungarn und den Occupationslandern im k. k. Naturhistorischen Hofmuseum zu Wien. Ann. Hofmuseum Wien, vii, pp. 306-372.
- LUCAS, F. A. [See Alciformes, p. 32.]
- ---. [See also Trochili, p. 45.]
- LUZECKI, O. J. Zur Ornis der sädöstlichen Bukowina. Orn. Jahrb. iii, pp. 199-201.
- LYDEKKER, R. Remarks on some recently described Extinct Birds of Queensland. Ibis, 1892, pp. 530-533.
- --- Recent Researches in Fossil Birds. Nat. Sci. i, pp. 266-271.
- [See also Dinornithiformes, p. 25.]
- MACKAY, G. H. [See Charadriida, p. 33; Scolopacida, p. 33.]
- MACPHERSON, H. A. A Vertebrate Fauna of Lakeland; including Cumberland and Westmoreland, with Lancashire north of the Sanda. With a Preface by R. S. Ferguson. Edinburgh: 1892, 8vo, i-civ & 1-552 pp.
- —. Winter Notes from Switzerland and North Italy. Zool. 1832, pp. 258-263.
- Manzella, A. [See Giglioli, H. H.]
- MARTORELLI, G. Le Mute Regressive degli Uccelli Migranti, e il loro scambio tra gli Emisferi Nord e Sud. Atti Soc. Ital. xxxiv, pp. 36-97.
- MATHEW, M. A. [See D'URBAN & MATHEW.]
- McLeon, J. C. Ornithological Notes from New Zealand. Ibis, 1892, pp. 250-254.
- MEARNS, E. A. [See Falconida, p. 39.]
- MERRIAM, C. HART. The Geographic Distribution of Life in North America, with special reference to the *Mammalia*. P. Biol. Soc. Wash. vii, pp. 1-64, pl. i.

- [Merriam, C. Hart.] The West Indian Fauna in South Florida. Nature, xlvi, pp. 458 & 459.
- ---. [See also Striges, p. 40.]
- MEVES, F. W. [THE LATE]. Obituary Notice. Ibis, 1892, pp. 191 & 192.
- MEYER, A. B. Abbildungen von Vogel-akeletten herausgegeben mit Unterstützung der Generaldirektion der Königl. Sammlungen fur Kunst und Wissenschaft in Dresden. Lief. xvi, xvii, xviii, taf. clickxx. [Zool. Rec. xxvii, Aves, p. 127.]
  - The species figured are mentioned under their respective families.
- —. Beitrag zur Kenntniss der Vogelfauna von Kaiser Wilhelms-Land. J. f. O. 1892, pp. 254–266.
  - [Timeliidæ, p. 59; Paradiseidæ, p. 50; Columbiformes, p. 28.]
- ---. The Birds of Sumba. Notes Leyd. Mus. xiv, pp. 265 & 266.

  [Paleornithing, p. 48.]
- ---. [See also Aquilina, p. 38.]
- [See also Pycnonotides, p. 59.]
- & Helm, F. vi. Jahresbericht (1890) der Ornithologischen Beobachtungstationen im Konigreiche Sachsen. Dresden: 1892, i-iv & 1-135 pp.
- MIDDENDORF, E. v. Vogelleben an den Russischen Leuchthürmen des Schwarzen, Kaspischen und Weissen Meeres. Ornis, 1893, pp. 1– 124.
- MILLAIS, J. G. [See Tetraonida, p. 26.]
- MILNE-EDWARDS, A. Sur les oiseaux fossiles des depots Eccénes de phosphate de Chaux du sud de la France. C.R. ii, Congr. Orn. Intern. ii, pp. 60-80.
- [Accipitriformes, p. 38; Striges, p. 40; Cuculidæ, p. 46; Trogones, p. 46; Coraciæ, p. 41; Cypseli, p. 44; Pseudopisthocomi, p. 29; Pterocletes, p. 28; Galliformes, p. 27; Grues, p. 34; Arden, p. 34; Ralliformes, p. 30; Tupinopiformes, p. 35.]
- MIVART, St. GEORGE. Birds: The Elements of Ornithology. London: 1892, 8vo, 330 pp.
- Mochler-Ferryman, A. F. Up the Niger: Narrative of Major Claude Macdonald's Mission to the Niger and Benue Rivers, W. Africa. London: 8vo, 1892, Birds, Appendix, p. 310.
- MORE, A. G. Recent Additions to the List of Irish Birds. Irish Nat. i, pp. 3 & 4.
- NACHTRIEB, H. F. First Report of the State Zoologist, accompanied with Notes on the Birds of Minnesota, by Dr. P. L. Hatch [q.r.]. Minnespolis: 1892, 8ve, 487 pp.

- NEWTON, A. Errors concerning the Sanderling. [Add. to Ibis for July, 1892], 8vo, 5 pp. [Scolopacidæ, p. 33.]
- NICHOLLS, H. Notes on the Habits of some of our British Birds. Tr. Eistb. Nat. Hist. Soc. 1892, pp. 253-258.
- NICOLLS, J. A., & EGLINGTON, W. The Sportsman in South Africa. The haunts, habits, description, and the pursuit of all game, fur and feather, found south of the Zambesi (including the Cape Colony, Transvaal, Bechuanaland, Natal, and Damaraland), at the present day, with brief notices of the best-known fresh and saltwater fish. 1892, 8vo, vi & 148 pp., pls. i-xii.
- North, A. J. Supplement to the Descriptive Catalogue of Nests and Eggs of Birds found Breeding in Australia and Tasmania. Rec. Austral. Mus. ii, pp. 11-22.
- [Striges, p. 40; Cacatuida, p. 47; Coccyges, p. 47; Paradiseida, p. 50; Timeliida, p. 59; Campophagida, p. 60; Meliphagida, p. 54.]
- Additions to the Avifaunas of Tasmania, and Norfolk and Lord Howe's Islands. T. c. pp. 36-38.
- [Anseriformes, p. 35; Pelargiformes, p. 34; Phalacrocoraces, p. 36; Scolopacida, p. 33.]
- OATES, E. W. [See BROOKS, W. E.]
- OGILVIE, F. M. [See Larida, p. 32.]
- OGILVIE-GRANT, W. R. Catalogue of the Picariæ in the Collection of the British Museum. Bucerotes [q.v., p. 42] and Trogones [q.v., p. 46]. [Vol. xvii of the "Catalogue of Birds."] 1892, 8vo, pp. 347-502, pls. xiii-xvii.
- -, [See also Perdicidæ, p. 27.]
- ---. [See also Hemipodii, p. 28.]
- OLPHE-GALLIARD, L. Christian Ludwig Brehm's Ornithologische Briefe (with Portrait). Orn. Jahrb. iii, pp. 127-162, taf. i.
- Onslow [Earl of]. Native New Zealand Birds. Nature, xlvi, pp. 502 **&** 503.
- Oustalet, M. E. Catalogue des oiseaux rapportés de l'Ile Nias, par M. J. Claine en 1891. Bull. Soc. Philom. viii, iv, pp. 107-122.
- -. Contributions à la Faune de la Chine et du Thibet. Description des espéces et de races nouvelles d'oiseaux données recemment au Musée d'Histoire Naturelle, par le Prince Henri d'Orleans. Ann. Sci. Nat. (7) xii, pp. 271-318.
  - [See Perdicidæ, p. 27; Cinclidæ, p. 58; Timeliidæ, p. 58; Fringillidæ, p. 52; Paridæ, p. 55; Zosteropidæ, p. 55.]

- [Oustalet, M. E.] Notice sur quelques espéces nouvelles ou peu connues de la Collection ornithologique du Muséum d'Histoire Naturelle. N. Arch. Mus. Paris (2) iv, pp. 211-220, pls. xiv & xv. [Halcyones, p. 42; Paradiseidæ, p. 50; Ampelidæ, p. 56; Muscicapidæ, p. 61; Tyrannidæ, p. 62.]
- Arts sur le Congrès Ornithologique International de Budapest en 1891. Arch. Miss. Sci. 1892, 34 pp.; t. c. pp. 441-473 (1891).
- ----. Rapport sur la Biologie des Oiseaux. C.R. ii, Congr. Orn. Intern. ii, pp. 95-117.
- ----. [See Dybowski, J.]
- PACKARD, A. S. The Labrador Coast. A Journal of Two Summer Cruises to that Region. With Notes on its Early Discovery, on the Eskimo, on its Physical Geography, Geology, and Natural History. New York: 1891, 8vo, 1-513 pp., with maps and illustrations.
- PALACKY, J. Ueber die Nordostasiatische Ornis. Verh. z.-b. Wien, xlii, pp. 43-46.
- PALMER, J. E. Birds of Lambay Island. Irish Nat. i, pp. 114-117.
- PARKER, T. JEFFERY. [See Apterygiformes, p. 25.]
- PATTERSON, R. L. Birds of Rathlin Island, Antrim. Irish Nat. i, pp. 50-53, 70-73, 85, & 86.
- ---. [See Procellariiformes, p. 32; Coraciidæ, p. 41.]
- PELZELN, A. VON [THE LATE]. Obituary Notice. Ibis, 1892, pp. 188-190; Auk, ix, pp. 74 & 75.
- Perzina, E. Wiener Trivialnamen. Orn. Jahrb. iii, pp. 66-71.
- Peters, E. Die Vögel Curaçaos nach meinem Journalnotizen vom 21 August bis 5 Sept. 1890. J. f. O. 1892, pp. 104-122.
- PIGOTT, D. London Birds and London Insects. Roy. 8vo, 1892, pp.
- PLESKE, T. Die ornithologische Ausbeute der Expedition der Gebrüder G. und M. Grum-Grzimailo nach Central Asien (1889-90). Mél. biol. xiii, livr. 2, pp. 273-301.
  - [Phasianidæ, p. 26; Turdidæ, p. 57.]
- —. Ornithographia Rossica. Die Vogelfauna des Russischen Reichs. Bd. ii, Lief. v (Locustella, Cettia, Urosphena), pp. xli-liv & 561-666, taf. iv.

[Sylviida, p. 57.]

- Prain, D. Fauna of Narcondam and Barren Island. J. A. S. B. 1892, pp. 109-120. Birds, pp. 111-114.
- PRATT, A. E. The Snows of Tibet. London: 1892, 8vo. Birds, pp. 235-237.
  - 1892. [vol. xxix.]

- Quelch, J. J. Avifauna of Georgetown (British Guiana). Journ. Agr. & Comm. Soc. Brit. Guian. (2) vi, pp. 182 & 183.
- —. Additions to the Avifauna of British Guiana. T. c. p. 186.
- [See also Accipitriformes, p. 37; Cotingidæ, p. 62.]
- RABÉ, DR. Observations sur les passages d'oiseaux dans le département de l'Yonne. Bull. Soc. Yonne, xlvi, pp. 1-44.
- RAEBURN, H. Some Further Notes on the Summer Birds of Shetland. P. Phys. Soc. Edinb. xi, pt. 1, pp. 67-73.
- ---. [See also WINGE, H.]
- RALFE, P. Birds of the Isle of Man. Zool. 1892, pp. 93-100.
- REICHENOW, A. Zur Vogelfauna von Kamerun. Erster Nachtrag. J. f. O. 1892, pp. 176-195, tab. ii.
- [Ralliformes, p. 30; Columbiformes, p. 28; Coliidæ, p. 46; Capitones, p. 48; Muscicapidæ, p. 60; Campophagidæ, p. 60; Laniidæ, p. 56; Eulabetidæ, p. 50; Ploceidæ, p. 51; Pycnonotidæ, p. 59; Nectariniidæ, p. 54; Zosteropidæ, p. 55; Sylviidæ, p. 58; Timeliidæ, p. 59; Turdidæ, p. 57.]
- —. Zur Vogelfauna von Togolaud. Nachtrag. T. c. pp. 233-236.
- ——. [See also Columbiformes, p. 28; Perdicidæ, p. 27; Phalacocoraces, p. 36; Capitones, p. 48; Colii, p. 46; Indicatores, p. 48; Caprimulgi, p. 44; Ploceidæ, p. 46; Fringillidæ, p. 53; Nectariniidæ, p. 54; Zosteropidæ, p. 55; Laniidæ, p. 56; Turdidæ, p. 56; Sylviidæ, p. 58; Pycnonotidæ, p. 59; Muscicapidæ, p. 60; Hirundinidæ, p. 61.]
- RENDALL, P. Notes on the Ornithology of the Gambia. Ibis, 1892, pp. 215-230.
- RHOADS, S. N. The Breeding Habits of the Florida Burrowing Owl (Spectyto cunicularia floridana). Auk, ix, pp. 1-8.
- —. The Birds of South-eastern Texas and Southern Arizona observed during May, June, and July, 1891. P. Ac. Philad. 1892, pp. 98-126.
- RIDGWAY, R. [See Trochili, p. 45.]
- RODICZKY, E. VON. Volkswirthschaftliche Bedeutung der Geflügelzucht in Ungarn. C.R. ii, Congr. Orn. Intern. ii, pp. 190-197.
- ROTHSCHILD, W. Descriptions of Seven New Species of Birds from the Sandwich Islands. Ann. N. H. (6) x, pp. 108-112.
- [See Anseriformes, p. 35; Drepanidæ, p. 52; Fringillidæ, p. 52; Meliphagidæ, p. 54; Sylviidæ, p. 57.]
- Russ, K. Ueber die Nest- und Jugendkleider in der Gefangenschaft gezüchteter fremländischer Vögel nebst angeben über Nestbau,
   Gelege, Verfärbung, und Geschlects-Verschiedenheiten. C.R. ii,
   Congr. Orn. Intern. ii, pp. 129-136.

- Ruzskii, M. Kratkii otchet ob ornitologhicheskikh izslyedovaniyakh Kazanskoi ghubernii. Protok. obech. estest. Kazan, 1890-91 (1891), No. 126, 13 pp.
- RZEHAK, E. C. F. Oesterreichisch-schlesisches Vogel-Idiotikon. Orn. Jahrb. iii, pp. 114-118.
- St. Maurice, Abbaye de. [See Besse & St. Maurice.]
- Salvadori, T. Uccelli di Engano raccolti dal dott. Modigliani. Ann. Mus. Genov. (2) xii, pp. 123-142.
- [See Campophagidæ, p. 60; Zosteropidæ, p. 55; Turdidæ, p. 56; Eulabetidæ, p. 50; Columbiformes, p. 29.]
- —. Viaggio di Lamberto Loria nella Papuasia Orientale. I. Uccelli della Nuova Guinea Meridionale-Orientale e delle Isole Heath E Killerton. II. Uccelli delle Isole D'Entrecasteaux. III. Collezioni Ornitologiche. T. c. pp. 797–834.
- ---. [See also Columbiformes, p. 29; Conurinæ, p. 47; Pioninæ, p. 47; Paridæ, p. 55.]
- Salvin, O. Catalogue of the *Picariæ* in the Collection of the British Museum. Upupæ, Trochili. [Vol. xvi of the Catalogue of Birds.] London: 8vo, 1892, pp. 1-433, pls. i-ix.

[Upupæ, p. 43; Trockili, p. 45.]

- —. [See also Godman & Salvin.]
- & GODMAN, F. D. Biologia Centrali-americana; or, Contributions to the Knowledge of the Fauna and Flora of Mexico and Central America. London: 1892, 4to, Aves, pts. xcix, c, & cii-cv, pp. 193-368, pls. xlix-lvii.
- [Formicariidæ, p. 62; Trockili, p. 45.]
- & —. On a Collection of Birds from Central Nicaragua. Ibis, 1892, pp. 324-328.

[Trochili, p. 46; Certhiidæ, p. 54.]

- Sandman, J. A. Fågelfaunan på Karlö och Kringliggande Skäv. Med. Soc. Fenn. xvii, pp. 187-272.
- SAUNDERS, H. [See Procellariidæ, p. 31; Stercorariidæ, p. 32.]
- Schaeck, F. de. Les variétés d'oiseaux du Muséum de Paris. C.R. ii, Congr. Orn. Intern. ii, pp. 81-91.
- —. Drei Wocher in der N\u00e4he von Fontainebleau. Orn. Jahrb. iii, pp. 24-27.
- Schaeff, E. Die Excursion nach dem Hanság und dem Neusiedler-See. C.R. ii, Congr. Orn. Intern. ii, pp. 198-201.
- Schollmayer, --. Beiträge zur Ornis Krains. [Zool. Rec. xxviii, Aves, p. 25.]

- Schöpwinkel, A. Die Vogelwelt der Grafschaft Weruigerode. Schr. Ver. Harzes. vi, pp. 1-62.
- Schrader, G. Ornithologischer Beobachtungen auf meinen Sammelreisen. Orn. Jahrb. iii, pp. 11-19 & 43-54. [Zool. Rec. xxviii, Avcs, p. 25.]
- SCHULZ, F. [See STEMPLEMANN, H.]
- Sclater, P. L. On a Collection of Birds from the Island of Anguilla, West Indies. P. Z. S. 1892, pp. 498-500.
- ---. [See also JAMES, H. B.]
- [See also Columbiformes, p. 29; Fringillida, p. 52.]
- SCLATER, W. L. On the Indian Museum and its Collection of Birds. Ibis, 1892, pp. 65-87.
- A valuable memoir, giving a list of the types still extant in the Indian Museum, Calcutta.
- Scott, W. E. D. Observations on the Birds of Jamaica. I. A list of the Birds recorded from the island, with annotations. [Zool. Rec. xxviii, Aves, p. 25.] Auk, ix, pp. 120-129, 273-277, & 369-375.
- —. Notes on the Birds of the Caloosahatchie Region of Florida. T. c. pp. 209-218.
- —. [See Ardeidæ, p. 34.]
- Sebisanovic, G. Ornitolozko-mamaložke viesti iz varaždinske okohee. Glasnik Naravosl. Druž. iii, pp. 236–248 (1888).
- —. Ornitoloska viest iz varazdinstrog Kraja za. god. 1889. Op. cit. iv, pp. 312-315, 1889.
- Seebohm, H. List of the Birds of Heligoland as recorded by Herr Gätke. Ibis, 1892, pp. 1-32.
- —. Further additions to the Avifauna of Tsu-sima, in the Japanese Empire. T. c. pp. 399 & 400.
  - [Piciformes, p. 49.]
- —... [See also Scolopacida, p. 33; Phasianida, p. 26; Piciformes, p. 49; Sylviida, p. 57.]
- Sennett, G. B. [See Meleagrida, p. 28.]
- SHARPE, R. B. Catalogue of the Picariæ in the Collection of the British Museum. Coraciæ and Halcyones. Leptosomatidæ [q.v. p. 41], Coraciidæ [q.v. p. 41], Meropidæ [q.v. p. 43], Alcedinidæ [q.v. p. 41], Momotidæ [q.v. p. 43], Todidæ [q.v. p. 43], Coliidæ [q.v. p. 46.] 1892, pp. 1-345, pls. i-xii. [Vol. xvii of The Catalogue of Birds.]

- [Sharpe, R. B.] On the Birds collected by Mr. F. J. Jackson, F.Z.S., during his recent expedition to Uganda, through the territory of the Imperial British East African Company. [Zool. Rec. xxviii, Area, p. 26.] With notes by the Collector. Part III. Ibis, 1892, pp. 152-164, pl. iv.
  - [Muscicapida, p. 61; Sylciida, p. 58.]
- —. Part IV. T. c. pp. 299-322, pl. vii.
  [Muscicapida, p. 61; Capitones, p. 48; Meropes, p. 43.]
- —. Part v. T. c. pp. 534-555, pl. xiv. [Perdicida, p. 27.]
- —. Descriptions of New Species of Birds discovered by Mr. C. Hose on Mount Dulit in N. W. Borneo. T. c. pp. 322-324.
  - [Podargi, p. 40; Zosteropidæ, p. 55; Turdidæ, p. 56; Capitones, p. 48.]
- ---. On a Collection of Birds from Mount Dulit, in North Western Borneo. T. c. pp. 430-442, pls. x & xi.
  - [Campophagida, p. 60; Eurylami, p. 49; Capitonida, p. 48.]
- ——. [See also Rallidæ, p. 30; Striges, p. 40; Falconidæ, p. 39; Corvidæ, p. 50; Oriolidæ, p. 51; Eulabetidæ, p. 50; Timeliidæ, p. 59; Fringillidæ, p. 53; Sylviidæ, p. 58; Muscicapidæ, p. 60; Cumpophagidæ, p. 60; Pycnonotidæ, p. 59; Eurylæmi, p. 49.]
- SHARPE, R. B., & WYATT, C. W. A Monograph of the Hirundinida, or Family of Swallows. Part xv. 1892, 4to.
  - [*Hirundinida*, p. 61.]
- SHELLEY, G. E. [See Parida, p. 55; Zosteropida, p. 55; Nectarinida, p. 54.]
- Shufeldt, R. W. A Study of the Fossil Avifauna of the Equus Beds of the Oregon Desert. J. Ac. Philad. ix, pp. 383-425, pls. xv-xvii. [Lariformes, p. 32; Ralliformes, p. 31: Tetraonida, p. 27: Pelargiformes, p. 34; Phanicopteriformes, p. 35; Accipitriformes, p. 38; Icterida, p. 51.]
- —. Concerning the Taxonomy of the North American Pygopodes, based upon their osteology. J. Anat. Phys. xxvi, pp. 199-203.
  [See also Trockili, p. 46]
- Sibree, J. On the Birds of Madagascar, and their connection with native Folklore, Proverbs, and Superstitions. [Zool. Rec. xxviii, Aces, p. 27.] Part IV. Ibis, 1892, pp. 103-119.
- —. The same. Part v. T. c. pp. 261-274.
- SJÖSTEDT, Y. Neue Vogelarten von Kamerun. J. f. O. 1892, pp. 313 & 314.
  - [Cuculi, p. 47; Pici, p. 49; Ploceida, p. 51.]
- SLATER, H. H. [See DE LA TOUCHE, J. D.]

- STEJNEGER, L. Two additions to the Japanese Avifauna, including description of a new species. P. U. S. Nat. Mus. xv, pp. 371-373. [Scolopacidæ, p. 33; Sylviidæ, p. 57.]
- STEMPLEMANN, H., & SCHULZ, F. Enumeración de las Aves de la provincia de Córdoba República Argentina. Bol. Ac. Arg. pp. 393-408, x, 4a, 1890.
- STONE, W. Birds collected by the West Greenland Expedition. P. Ac. Philad. 1892, pp. 145-152.
- STOSSICH, M. I distomi degli uccelli. Boll. Soc. Adr. xiii, pp. 143-194.
- STRUGNELL, W. B. [See WITCHELL & STRUGNELL.]
- STUDER, TH., & FATIO, V. Katolog der schweizerischen Vögel und ihrer Verbreitungsgebiete. 8vo, 70 pp., with map.
- STYAN, F. W. [See Perdicidæ, p. 27; Corvidæ, p. 50; Sylviidæ, p. 58; Muscicapidæ, p. 61.]
- Suchetet, A. Oiseaux hybrides rencontrés à l'etat sauvage. 111e Partie. Les Passereaux [Zool. Rec. xxviii, Aves, p. 29]. Bull. Soc. Z. Fr. (2) v, pp. 253-352, 352-524.
- TINKLER, JOHN E. The Avifauna of Arkengarthdale, Swaledale, and the New Forest. Nat. 1892, pp. 309-325.
- TIPPENHAUER, L. Die Insel Haiti. Leipzig: 4to, 1893 (1892), xviii & 693 pp., 32 pls., Aves, pp. 317-323.
- TRISTRAM, H. B. On two small collections of Birds from Bugotu and Florida, two of the smaller Solomon Islands. Ibis, 1892, pp. 293-299.
  - [Campophagidæ, p. 60.]
- [See also Nestoridæ, p. 47.]
- TRUMBULL, G. [See Anseriformes, p. 36.]
- TSCHUSI-SCHMIDHOFFEN, V. RITTER VON. Am Velenezeer und Kleinen Balatonsee. C.R. ii, Congr. Orn. Intern. ii, pp. 202-209.
- ---. August von Pelzeln. Obituary Notice. Orn. Jahrb. iii, pp. 1-11.
- —. Ueber die Anlegung von Privat-Sammlungen. T. c. pp. 71-73.
- ---. Pfarrer Carl (P. Blasius), Hanf. Obituary Notice. T. c. pp. 87-97.
- ---. Ornithologische Miscellen. T. c. pp. 254-256.
- [See also Pterocletes, p. 28.]
- VALENTINITSCH, F. [See Tetraonida, p. 26.]
- Vallon, G. Seltenheiten der italianische Ornis. Orn. Jahrb. iii, pp. 97-108.
- ---. Escursioni Ornitologiche nel Trentino. Boll. Soc. Adr. xiii, pp. 117-14!.

- VAN DYCK, W. T. [See Nectariniida, p. 54.]
- VORDERMAN, A. G. Java Vögels, I. Nat. Tijdschr. Nederl. Ind. li, Afl. 4, pp. 373-416.
- ---. Bijdrage tot de kennis der Avifauna van het Eiland Bawean.
  T. c. pp. 417-422.
- WAITE, E. R. Sterna and Shoulder Girdle of Birds. Tr. Leeds Nat. Club, ii, p. 67.
- WALKER, J. J. The Bird-life of Adele Island, North-West Australia. Ibis, 1892, pp. 254-261.
  - [Sulæ, p. 36; Fregati, p. 36.]
- WALLACE, A. R. Island Life. Second Edition. London: 8vo, xx & 563 pp.
- •Walters, Alax. Palms and Pearls; or, Scenes in Ceylon. London: 8vo, 1892, pp.
- WESTERN, R. W. The tactics adopted by certain Birds when flying in the wind. P. R. Soc. Edinb. pp. 76-80, pl. xviii.
- WHYMPER, E. Travels amongst the Great Andes of the Equator, with Supplementary Appendix. London: 1892, 2 vols., 8vo.
- WILSON, S. B., & EVANS, A. H. Aves Hawaienses: The Birds of the Sandwich Islands. Part III. 1892, 4to.
  - [Drepanida, p. 52; Scolopacida, p. 33.]
- WIEGE, H. Fuglene ved de danske Fyr i 1890, Vid. Medd. 1891, pp. 61-132.
- On the Breeding of the Sand Grouse in Captivity in Denmark. Op. cit. 1892, pp. 126-130. [Translated by H. Raeburn, Zool. 1892, pp. 341-345.]
- WITCHELL, C. A. Bird-song and its Scientific Value. P. Cottesw. Nat. F. C. x, pp. 238-257.
- WITCHELL, C. A., & STRUGNELL, W. B. The Fauna and Flora of Gloucestershire. Strond: 1892, roy. 8vo, i-xxiv & 1-300 pp.
- Wolf, T. Geografía y Geología del Ecuador. 8vo, xii & 671 pp. [Notes on the Birds, pp. 456-460.]
- WOLLEY-DOD, C. [See Cuculida, p. 47.]
- Wurm, W. Über die Farben der Vogelfedern. JH. Ver. Württ. xlviii, pp. xli-xlvi.
- WYATT, C. W. [See SHARPE & WYATT.]
- Young, C. J. Notes on the Natural History of the Blue Mountain, Co. of Leeds, Ontario. Ottawa Nat. p. 45.
- ZIEGLER, J. Tierphänologische Beobachtungen zu Frankfurt a. Main. Ber. Senck. Ges. 1892, pp. 47-69.

## SPECIAL SUBJECTS.

### II.—FAUNISTIC.

PALEARCTIC REGION. See Alessi, Aplin, Bonvalot, Brusina, Campbell, Collett, Dresser, Dybowski, Fatio, Gätke, Giglioli, Hartert, Hartlaub, Ijima, Kaiser, Kerville, Koenig, Lilford, Lorenz, Macpherson, Oustalet, Pleske, Pratt, Schrader, Seebohm, Stejneger, Studer, Tschusi, Winge.

ETHIOPIAN REGION. See Bocage, Büttikofer, Eglington, Homeyer, Nicholls, Ogilvie-Grant, Reichenow, Rendall, Sharpe, Shelley.

INDIAN REGION. See Brooks, Büttikofer, Davison, De la Touche, Oustalet, Salvadori, Sharpe, Vorderman.

Australian Region. See Buller, Büttikofer, Campbell, Collett, De Vis, Forbes, Meyer, North, Rothschild, Salvadori, Tristram, Wilson.

NEARCTIC REGION. See Allen, "Anon.", Anthony, Bendire, Bryant, Cory, Keeler, Loomis, Merriam, Packard.

NEOTROPICAL REGION. See Alfaro, Allen, Berlepsch, Boucard, Chapman, Cherrie, Cory, Godman, Hartert, Holland, Hudson, Kerr, Peters, Quelch, Salvin, Sclater, Scott, Tippenhauer, Wolf,

## III.-ANATOMY AND MISCELLANEOUS.

ANATOMY. See Beddard, Couvreur, Dareste, Fabani, Fürbringer, Helm, Klug, Parker.

OSTEOLOGY. See Beddard, Meyer.

OOLOGY. See Bendire, Blasius, Crowley, Hall, Hewetson, North.

NEO-OOLOGY. See Evans.

PTERYLOGRAPHY. See Church, Ficalbi, Keeler, Martorelli.

FOSSIL BIRDS. See Dèperet, Forbes, Lydekker, Milne-Edwards, Shufeldt.

BIOGRAPHY. See Boucard, Burmeister, Büttikofer, Pelzeln, Tschusi-Schmidhoffen.

LITERATURE, See Persina, Sclater (W. L.).

CLASSIFICATION. See Fürbringer, Gadow, Mivart.

GEOGRAPHICAL DISTRIBUTION. See Giglioli, Palacky,

MIGRATION. See Cordeaux, Czýnk, Dixon.

#### IV.—SYSTEMATIC.

## Subclass RATITÆ.

#### Order STRUTHIONIFORMES.

See Nicolls, J. A., & Eglington, W.

## Order CASUARIIFORMES.

Dromaius gracilipes, n. sp. (fossil); C. W. DE VIS, P. Linn. Soc. N.S.W. (2) vi, pp. 445-448, pl. xxiii, figg. 7a & 7b.

### Order DINORNITHIFORMES.

- FIELD, H. C. Discoveries of Mos Bones. Tr. N. Z. Inst. xxiv, pp. 558-561.
- HUTTON, F. W. On the Moas of New Zealand. T. c. pp. 93-172, pls. xv-xvii.
- Dinornis potens, n. sp. (= D. ingens, var. robustus, Owen), p. 115;
  Tylopteryx, n. subg. D. (T.) torosus, n. sp. (= D. gracilis, Haast),
  p. 116; Palapteryx plenus, n. sp., p. 122 (= P. dromioides, Owen).
  Anomalopteryx antiquus, p. 124, n. sp.; Mesopteryx, n. g., p. 129,
  Euryapteryx ponderosus, p. 137 (= Dinornis gravis, Hutton), E. pygmaus, p. 139, n. spp.
- D. potens, D. torosus; P. dromioides; Anomalopteryx didiformis; Cela curtus; Syornis crassus; Mesopteryx didinus; Euryapteryx ponderosus: bones figured. Id. ut suprà.
- LYDEKKER, R. The History of the Moss, or Extinct Flightless Birds of New Zealand. Nat. Sci. i, pp. 588-595.
- Palaocasuarius, n. g. (fossil). Type, P. elegans, n. sp., New Zealand; H. O. Forbes, Tr. N. Z. Inst. xxiv, p. 189.

#### Order ÆPYORNITHIFORMES.

See Newton, R. B., Geol. Mag. (3) x, pp. 197 & 198.

\*Epyornis medius?, egg of; P. L. Sclater, P. Z. S. 1892, p. 299.

### Order APTERYGIFORMES.

- PARKER, T. J. Observations on the Anatomy and Development of the *Apteryx*. P. R. Soc. p. 340; *id*. Phil. Trans. clxxxii B, pp. 25-134, pls. viii-xix.
- Apteryx maxima, notes on; W. L. Buller, Tr. N. Z. Inst. xxiv, p. 91.

  Metapteryx, n. g. Type, M. bifrons, n. sp. (fossil), Queensland; C. W. de
  Vis, P. Linn. Soc. N.S.W. (2) vi, pp. 448-454, pl. xxiii, figg. 8a & 8b.

## Subclass CARINATÆ.

#### Order CRYPTURIFORMES.

Nothoprocta moebiusi, n. sp., Bolivia, H. von Berlepsch, Ber. Allg. Deutschl. Orn. Ges., Dec. 1892, p. 13.

Crypturus garleppi, n. sp., Bolivia, id. t. c. p. 12.

#### Order GALLIFORMES.

BENDIRE, C. Life Histories of North American Birds, with special reference to their Breeding Habits and Eggs. U.S. Nat. Mus., Special Bulletin, No. 1, Washington, 1892, 4to, figs.; eggs of Colinus virginianus (pl. i, fig. 1); Oreortyx pictus plumiferus (pl. i, figs. 2

& 3); Callipepla squamata (pl. i, figs. 4 & 5), C. squamata castanogastris (pl. i, figs. 6 & 7), C. californica (pl. i, figs. 8-10), C. gambeli (pl. i, figs. 11-14); Cyrtonyx montesumes (pl. i, fig. 15); Dendragapus obscurus fuliginosus (pl. i, figs. 16-19), D. canadensis (pl. i, figs. 20-23); Bonasa umbellus (pl. ii, fig. 1), B. umbellus togata (pl. ii, fig. 2), B. umbellus umbelloides (pl. ii, fig. 3), B. umbellus sabini (pl. ii, fig. 4); Lagopus lagopus (pl. ii, figs. 5-10), L. rupestris (pl. ii, figs. 11-15), L. lueurus (pl. ii, figs. 16 & 17); Tympanuchus americamus (pl. ii, figs. 18-20), T. cupido (pl. iii, figs. 2), T. pallidicinctus (pl. iii, fig. 1); Pediocates phasianellus (pl. iii, figs. 3-5), P. phasianellus columbianus (pl. iii, figs. 6-8), P. phasianellus campestris (pl. iii, figs. 9 & 10); Centrocercus urophasianus (pl. iii, figs. 11-13).

#### Suborder CRACES,

Ortalis retula maccalli: egg figured; C. Bendire, t. c. pl. iii, fig. 16.

Penelope purpurusceus: skeleton figured; A. B. Meyer, Abbild. Vögel.

Skel., Taf. clxxviii.

#### Suborder PHASIANI.

## Family Phasianidz.

- Crossoptilon leucurum, n. sp., E. Thibet; H. Seebohm, Bull. B. O. C., No. 4, p. xvii.
- Rheinhardius phænix, nom. emend. [?], pro R. ocellata (Verr.); A. Bou-CARD, Humming Bird, ii, pp. 10-12.
- Phasianus satcheuensis, n.sp., North of Nian Schan Mountains; Th. Pleske, Mèl. Biol. xiii, livr. 2, pp. 296 & 297.
- P. tarnovskii, n. sp., Zarafshan, Central Asia; H. SEEBOHN, P. Z. S. 1892, p. 271.

## Family TETRAONIDE.

- See BENDIRE, C., ut suprà, p. 25.
- MILLAIS, J. G. Game Birds and Shooting-sketches; illustrating the habits, modes of capture, stages of plumage, and the hybrids and varieties which occur amongst them. London: 1892, roy. 4to, 72 pp., xv pls.
- Valentinitsch, F. Das Haselhuhn (*Tetrao bonasia*). Dessen Naturgeschichte und Jagd. Eine ornithologische und jägdliche Monographie. 8vo, xii & 288 pp., pls. 1-4. [Orn. Jahrb. iii, pp. 82 & 83.]
- Lagopus lagopus, in Maine; H. MERRILL, Auk, p. 300: in the Faroe Is.;
  H. W. Feilden, Zool. 1892, p. 413.
- Bonasia bonasia, skeleton figured; A. B. MEYER, Abbild. Vögel. Skel.,
  Taf. clxxx.
- Pediocetes nanus, n. sp. (fossil), Pliocene of Oregon; R. W. Shufeldt, J. Ac. Philad. ix, p. 414.

- Palæotetrix, n. g. Type, P. gilli, n. sp. (fossil), Equus Beds, Oregon; B. W. Shufeldt, J. Ac. Philad. ix, p. 416.
- Dendragapus obscurus fuliginosus, its food; L. Belding, Zoe, iii, pp. 232 & 233.

## Family PERDICIDE.

## Subfamily Perdicinæ.

- NICOLLS, J. A., & EGLINGTON, W. The Sportsman in South Africa. 8vo, 1892.
- The Francolins and Quails described, and many of them figured (pp. 100-106, pls. i & xi).
- Tetraogallus henrici, n. sp., Tà-tsién-lou; E. Oustalet, Ann. Sci. Nat. (7) xii, p. 295.
- Palæortyx ocyptera, n. sp. (fossil), Eocene of S. France; A. MILNE-EDWARDS, C.R. ii, Congr. Orn. Intern. ii, pp. 71 & 72. P. caylux-ensis, n. sp. (fossil), id. t. c. p. 72.
- Palæocryptonyx donnezani, n. sp. (fossil), Pliocene of France; C. Déperet, C.B. cvi, p. 690.
- Perdiz hispaniolensis, n. sp., Spain; A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges. xi, p. 5.
- Francolinus and Pternistss: notes on these genera; W. R. OGILVIE-GRANT, Ibis, 1892, pp. 32-55, pl. i.
- F. vulgaris: its range; F. DE SCHAECK, Orn. Jahrb. iii, pp. 108-112.

  F. subtorquatus stuhlmanni, n. subsp., Victoria Nyanza, A. REICHENOW, J. f. O. 1892, pp. 17 & 18. F. uluensis, n. sp., Ukambani, W. R.
  OGILVIE-GRANT, Ibis, 1892, p. 44. F. sharpii, n. sp., Abyssinia, id.
  t. c. p. 47. F. jacksoni [Zool. Rec. xxviii, Aves, p. 35], figured, id. t. c. pl. i. F. gedgii, figured, R. B. Sharpe, Ibis, 1892, pl. xiv.
- Caloperdix borneensis, n. sp., Mt. Dulit, Borneo, W. R. OGILVIE-GRANT, Bull. B. O. C. ii, p. v. C. sumatrana, n. sp., Sumatra, id. ibid.
- Bambusicola: notes on the genus; id. Ibis, 1892, pp. 387-389.
- Arboricola: notes on the genus; id. t. c. pp. 390-399. A. gingica, figured; id. t. c. p. ix. A. ardens, n. sp., Hainan; F. W. STYAN, Bull. B. O. C. ii, p. vi.
- Coturnix: notes on the genus; W. R. OGILVIE-GRANT, Ann. N. H. (6) x, pp. 166-173.
- Coturniz communis, in Wigtownshire; P. Adair, Ann. Scott. Nat. Hist.i, pp. 168-171: nesting in N. Devon; F. H. C. Gould, Zool. 1892, p. 427. C. emini [Zool. Rec. xxviii, Ares, p. 36], figured; A. Reichenow, J. f. O. 1892, tab. i, fig. 3. [Is C. adansoni, W. R. Ogilvie-Grant, ut suprà.]

#### Subfamily ODONTOPHORINÆ.

Oreortyx pictus plumiferus: its food; R. Belding, Zoe, iii, pp. 233 & 234.

- Odontophorus guianensis: akeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxxix.
- Eupsychortyz gouldi, n. sp., Curação; H. von Berlepsch, J. f. O. 1892, pp. 98-101.

## Family MELEAGRIDE.

Meleagris gallopavo ellioti, n. subsp., Rio Grande, Texas; G. B. SENNETT, Auk, ix, pp. 167-169, pl. iii. M. gallopavo (fig. 14) and M. gallopavo mexicana (fig. 15): eggs figured; C. BENDIRE, Life Histories N. A. Birds, pl. iii.

#### Suborder HEMIPODII.

See Nicolls, J. A., & Eglington, W.

Turniz nigricollis: sexual differences; W. R. OGILVIE-GRANT, Ibis, 1892, pp. 346 & 347: nesting in captivity; LORD LILFORD, t. c. pp. 466 & 467.

#### Suborder PTEROCLETES.

See Nicolls, J. A., & Eglington, W.

- Syrrhaptes paradoxus: notes on the incursion of 1891; V. RITTER VON TSCHUSI ZU SCHMIDHOFFEN, Orn. Jahrb. iii, pp. 118-122: in the North of France; M. DUBOIS, Bull. Soc. L. Nord France, ix, pp. 11 & 12.
- Pterocles coronata, in Tunis; J. CABANIS, Ber. Allg. Deutschl. Orn. Ges., Sept. 1892, p. 5. P. varius and P. larvatus, n. spp. (fossil), Eocene of S. France, A. MILNE-EDWARDS, C.R. ii, Congr. Orn. Intern. ii, pp. 70 & 71.

### Order COLUMBIFORMES.

- BENDIRE, C. Life Histories of North American Birds, with special reference to their Breeding Habits, and Eggs. U. S. Nat. Mus. Special Bulletin, No. 1, Washington, 1892, 4to.
- Eggs of the following species are figured:—Columba fasciata (pl. iii, fig. 17). C. fasciata viosca (pl. iii, fig. 18), C. favirostris (pl. iv, fig. 2), C. leucocephula (pl. iv, fig. 4); Ectopistes migratorius (pl. iv, fig. 6); Zenaidura macrura (pl. iv, figs. 8 & 9); Zenaida zenaida (pl. ii, fig. 21); Engyptila albifrons (pl. ii, fig. 22); Melopelia leucoptera (pl. ii, fig. 23); Columbigallina passerina pallescens (pl. ii, fig. 24); Scardafella inca (pl. ii, fig. 25); Geotrygon montana (pl. ii, fig. 26).
- Columba domestica × Turtur risorius; H. GADBAU DE KERVILLE, Bull. Soc. Rouen (3) xxvii, pp. 195-199. C. domestica: its parasites; C.R. Soc. Biol. (9) iv, pp. 49-53.
- Haplopelia inornala. n. sp., Cameroons, A. REICHENOW, Ber. Allg. Deutschl. Orn. Ges., Feb. 1892, p. 6; id., J. f. O. 1892, p. 179,

- Turtur: notes on the African species of the genus; A. REICHENOW, J. f. O. 1892, pp. 12-15.
- Leucosarcia picata: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxxvii.
- Carpophaga ænothorax, n. sp., I. of Engano, T. Salvadori, Ann. Mus. Genov. (2) xii, pp. 130 & 140.
- Myristicivora luctuosa: skeleton figured; A. B. MEYER, Abbild. Võgel. Skel., Taf. clxxvi.
- Ptilopus coronulatus huonensis, n. subsp., Huon Gulf, E. New Guines; id. J. f. O. 1892, pp. 263 & 264. P. salvadorii, n. sp., Jobi I, W. Rothschild, Bull. B. O. C. iii, p. x. P. tristrami, n. sp., Marquesas L., T. Salvadori, Boll. Mus. Zool. Tor. vii, pp.
- Interior melanospila: akeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxxv, fig. 1.
- Phlogenas bimaculata, n. sp., Makassar, T. Salvadori, Bull. B. O. C., No. 3, p. 8. P. albicullis, n. sp., Box L, id. t. c. p. x.
- Macropygia albicapilla: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxxv, fig. 2.
- Macropygia cinnumomea, n. sp., I. of Engano, T. Salvadori, Ann. Mus. Genov. (2) xii, pp. 140 & 141.
- Zenaida richardsoni, = Z. zenaida; C. B. Corv, Cat. W. Ind. Birds, p. 138.
- Columbigallina passerina terrestris, n. subsp., E. United States, F. M. Chapman, Bull. Am. Mus. iv, p. 293.
- Geophaps scripta: notes on nestlings; P. L. Sclater, P. Z. S. 1892, pp. 76 & 77.
- Omotreron griseicauda: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxxiv.

#### Order OPISTHOCOMIFORMES.

### Suborder OPISTHOCOMI.

Opisthocomus cristatus, crop and sternum of: a contribution to the question of the correlation of organs and the inheritance of acquired characters; H. Gadow. P. R. Irish Ac. (3) ii, pp. 147-154, pls. vii & viii.

### Suborder PSEUDOPISTHOCOMI.

Filholornis, n. g. Type, F. paradoza, n. sp. (fossil), Eocene of S. France;
A. MILNE-EDWARDS, C.R. ii, Congr. Orn. Intern. ii, pp. 67 & 68.
F. gravis and F. debilis, n. spp., id. t. c. p. 69.

### Order RALLIFORMES.

Elaphrocnemus, n. g. Type, E. phasianus, n. sp., Eocene of S. France;
A. MILNE-EDWARDS, C.R. ii, Congr. Orn. Intern. ii, pp. 77 & 78. E. gracilis and E. crez, n. sp. 78 & 79.

- Orthocnemus, n. g. Type, O. gallicus, n. sp. (fossil), Eocene of S. France; id. t. c. pp. 74 & 75. O. cursor, n. sp., id. t. c. p. 76. O. major, n. sp., id. t. c. pp. 76 & 77.
- Rallus dasypus, n. sp. (fossil), id. t. c. pp. 73 & 74. R. arenarius, n. sp., id. t. c. p. 74. R. longirostris cubanus, n. subsp., Cuba, F. M. CHAPMAN, Bull. Am. Mus. iv, pp. 288 & 289.
- Cabalus modestus, Hutton, is the young of C. dieffenbachii, Gray; H. O. Forbes, Bull. B. O. C., No. 4, p. xx. [Cf. H. O. Forbes, Bull. B. O. C. ix, p. xxxix.]
- Aramides plumbeicollis, n. sp., Costa Rica, J. C. Zeledon, Ann. Mus. Costa Rica, iii, p. 134.
- Crecoides, n. n. for Crecioides, Shuf.; SHUFELDT, J. Ac. Philad. ix, pp. 411 & 412, notes.
- Porzana bailloni, in Herefordshire; W. H. M. Ayres, Zool. 1892, p. 27: in Yorkshire; J. BACKHOUSE, Nat. 1892, p. 308: in Somersetshire; H. St. B. Goldsmith, Zool. 1892, p. 412.
- Porzanula, n. g. Type, P. palmeri, n. sp., Laysan I., North Pacific; F. W. Frohawk, Ann. N. H. (6) ix, pp. 247-249.
- Rallus ecaudatus, King, and R. sandvicensis, Gm., belong to the genus Pennula; G. HARTLAUB, Abh. Ver. Brem. xii, heft. 3, pp. 395-397.
- Porzanula (ut suprà) = Pennula; id. t. c. p. 399.
- Rallus sandwichensis, is a Pennula, and probably = P. ecaudata, King; R. B. SHARPE, Bull. B. O. C., No. 4, p. xx. [Cf. HARTLAUB, op. cit. No. 5, p. xxiv, and SHARPE, op. cit. No. 5, p. xxiv.]
- Kittlitsia, n. g. [but cf. Hartert, Zool. Rec. xxviii, Aves, p. 57]. Type, Rallus monasa, Kittl., from Ualan; G. Hartlaub, Abh. Ver. Brem. xii, heft. 3, p. 391.
- Aphanolimnas, n. n. pro Kittlitzia, Hartl. (nec Hartert); R. B. SHARPE, Bull. B. O. C., No. 4, p. xx (Dec., 1892).
- Corethrura elegans, from the Cameroons; A. REICHENOW, J. f. O. 1892, pp. 178 & 179.
- Ocydromus insignis, n. sp. (subfossil), Chatham Is., H. O. FORBES, Tr. N. Z. Inst. xxiv, p. 188.
- Aphanapteryx hawkinsii, n. sp. (subfossil), Chatham Is., id. t. c. p. 189.
- Diaphorapteryx, n. g. for the preceding; id., Bull. B. O. C., No. 4, p. xxi. [Genus cancelled, id., Bull. B. O. C., No. 9, p. l.]
- Porphyrio mackintoshi, n. sp. (fossil), Queensland, C. W. DE VIS, P. Linn. Soc. N.S.W. (2) vi, p. 440, pl. xxiv, fig. 2a.
- Notornis parkeri, n. sp. (fossil), New Zealand, H. O. Forbes, Tr. N. Z. Inst. xxiv, p. 187.
- Aptornis: notes on the genus; A. Hamilton, t. c. pp. 175-184.
- Gallinula coccineipes [Zool. Rec. xxviii, Aves, p. 37], figured; J. D. DE LA TOUCHE, Ibis, 1892, pl. xii [= Amaurornis akool]. G. peralata, n. sp. (fossil), Queensland, C. W. DE VIS, P. Linn. Soc. N.S.W. (2) vi, pp. 440 & 441, pl. xxiv, figs. 3a & 3b. G. chloropus × Fulica atra; H. Kreye, Orn. Jahrb. iii, p. 172.

- Porphyriornis, n. g. Type, P. comeri, n. sp., Gough I.; J. A. Allen, Bull, Am. Mus. iv, pp. 57 & 58.
- Tribonyx effluxus, n. sp. (fossil), Queensland, C. W. DE VIS, P. Linn. Soc. N.S.W. (2) vi, pp. 439 & 440.
- Fulica minor, n. sp. (fossil), Pliocene of Oregon, R. W. Shufeldt, J. Ac. Philad. ix, p. 412.

## Order PODICIPEDIDIFORMES.

- Shuffeldt, R. W. Concerning the Taxonomy of the North American Pygopodes, based upon their osteology. J. Anat. Phys. xxvi, pp. 198-203.
- Podicipes nigricollis, in Merionethshire; G. H. CATON HAIGH, Zool. 1892, p. 311: in Anglesea; W. Banks, t. c. p. 411: in Norfolk; T. Bidwell, t. c. p. 358. P. rubricollis, in Devon; E. A. S. Elliot, t. c. p. 331.
- Podicipes auritus, supposed nesting in Ross-shire, = P. fluviatilis; A. W. Evans, Ann. Scott. Nat. Hist. i, pp. 171 & 172.

### Order COLYMBIFORMES.

Motions on land; HAIGH, Zool. 1892, p; 226.

Taxonomy; see Shufeldt, R. W., ut suprà.

Urinator imber: migration in Massachusetts; G. H. MACKAY, Auk, ix, p. 292.

### Order HESPERORNITHIFORMES.

Hesperornis regalis: restoration; H. N. HUTCHINSON, Extinct Monsters, p. 109, pl. ii.

#### Order PROCELLARIIFORMES.

- Dalgleish, J. J. Notes on the Petrels of Madeira and adjoining Seas. P. Phys. Soc. Edinb. xi, pt. 1, pp. 27-30.
- Puffinus major, in Tiree; J. A. HARVIE-BROWN, Ann. Scott. N. H. i, p. 74. P. major and P. cinereus, in France; L. Bureau, Bull. Soc. Ouest France, ii, pp. 39 & 40, pl. iii, figs. 1 & 2.
- Cymochorea leucorrhoa, off Scottish Coasts; W. Evans, Ann. Scott. N. Hist. i, pp. 74-76.
- Oceanitis oceanicus, in Ireland; R. L. PATTERSON, Ibis, 1892, pp. 180 & 181: in Scotland; H. SAUNDERS, t. c. p. 183: in Jura; H. EVANS, Ann. Scott. N. H. i, p. 18.
- Procellaria pelagica used as a candle in the Orkneys and S. Kilda; E. LOVETT, P. Croydon Club, pp. 331 & 332; id., J. Northampt. Soc. 1892, pp. 113 & 114.

### Order ALCIFORMES.

- KEELER, C. A. Notes on the Auks breeding in the Farallone Islands. Zoe, iii, pp. 144-165, pls. xviii-xxi.
- Alca impennis: an unrecorded egg; F. A. Lucas, Auk, ix, p. 198: figured; H. Gadeau de Kerville, Faun. Norm. iii, pl. i.
- Uria troile, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxi.

#### Order LARIFORMES.

## Family STERCORARIIDÆ.

Stercorarius catarrhactes, in Shetland; W. E. CLARKE, Ann. Scott. N. H. i, pp. 87-92. S. parasiticus and S. crepidatus: notes on; H. SAUNDERS, Ibis, 1892, pp. 183-185.

Lestris pomarinus, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxi.

## Family LABIDE.

## Subfamily LARINÆ.

- Larus robustus, n. sp. (fossil), and L. oregonus, n. sp. (fossil), Equus Beds, Oregon, R. Shufeldt, J. Ac. Philad. ix, p. 398. L. argentatus smithsonianus, Habits in New England; G. H. Mackay, Auk, ix, pp. 221-228. L. leucopterus, and L. glaucus, in Ireland; R. Warren, Irish Nat. i, pp. 129-134 & 154-156. L. leucopterus, near Aldeburgh; F. Menteith Ogilvie, Zool. 1892, p. 114: In West and North of Scotland; Duke of Argyll, Ann. Scott. N. H. i, pp. 140-141. L. eburneus, in co. Dublin; E. G. Waddilove, Zool. 1892, p. 228. L. ridibundus, figured; Lord Lilford, Col. Fig. Br. Birds, part xxi. L. canus, a monstrosity; G. de Kerville, J. de l'Anat. Phys. xxviii, pp. 564-566, pl. xviii.
- Rissa tridactyla, its rudimentary hallux; W. E. CLARKE, Ibis, 1892, pp. 442-447.
- Xema sabinii, near Bournemouth; F. Coburn, Zool. 1892, p. 22: in Cornwall; T. H. Cornish, t. c. p. 22: in Norfolk; H. W. Feilden, t. c. p. 423: in Pembrokeshire; C. Jeffreys, t. c. p. 423: in San Francisco Bay; W. E. Bryant, Zoe, iii, p. 165.

### Subfamily STERNINÆ.

Anous stolidus, nests photographed; A. J. CAMPBELL, C.R. ii, Congr. Orn. Intern. ii, p. 157.

### Order ÆGIALORNITHIFORMES.

[INCERTÆ SEDIS. Cf. Zool. Rec. xxviii, Aves, p. 39.]

Ægialornis gallicus, Lydekker, = Tachyornis [Cypselidæ, q.v.]; A. MILNE-EDWARDS, C.R. ii, Congr. Orn. Intern. ii, pp. 66 & 67.

### Order CHARADRIIFORMES.

Legge, W. V. On the Geographical Distribution of the Australian Limicola. Pr. Austr. Ass. 1892, pp. [1-50].

#### Suborder CHARADRIL

# Family Charadelida.

- Charadrius squatarola, habits in Massachusetts; G. H. MACKAY, Auk, ix, pp. 143-152.
- Vanellus vulgaris, notes on nesting; R. Howse, Nat. 1892, pp. 137-139. Ægialius asiatica, figured; Lord Lilford, Col. Fig. Br. Birds, part xxi.

## Family Scolopacida.

- Scolopax rusticula, in Dalmatia; H. Hänisch, Orn. Jahrb. iii, pp. 27-30 & 189-191: nesting in North Uist; J. C. Orde, Ann. Scott. N. H.i. p. 73.
- Numenius borealis, habits and migration in New England; G. H. Mackay, Auk, ix, pp. 16-21: figured; Lord Lilford, Col. Fig. Br. Birds, part xxi: in California; W. E. Bryant, Zoe, iii, p. 165, [= N. hudsonicus; L. Belding, t. c. p. 257]. N. arquata, in Worcestershire in July; F. Coburn, Zool. 1892, p. 23: on Long Island; W. Dutcher, Auk, ix, p. 392. N. hudsonicus, in Massachusetts; G. H. Mackay, Auk, ix, pp. 345-352. N. cyanopus and allies; W. Legge, P. R. Soc. Tasm. 1888, pp. 133-136.
- Pavoncella pugnax, in North Carolina; H. H. BRIMLEY, Auk, ix, pp. 299 & 300: in Orkney; T. Peace, Ann. Scott. N. H. i, p. 73: in the Outer Hebrides, id., t. c. p. 139.
- Totamus fuscus, in Ireland; E. Williams, Zool. 1892, p. 35. T. incanus, figured; S. B. Wilson & A. H. Evans, Av. Haw. part iii.
- Eurinorhynchus pygmæus, Hainan; G. HARTLAUB, Abh. Ver. Brem. xii, p. 330.
- Tringa temminckii, in Japan; L. Stejneger, P. U. S. Nat. Mus. xv, p. 371. T. minutilla, in Devonshire; H. A. Evans, Zool. 1892,
  - p. 141. T. acuminata, in Norfolk; T. Southwell, Zool. 1892,
  - pp. 356-358: Н. Seebohm, Bull. B. O. C. No. 3, p. ix.
- Ereunetes occidentalis, in Connecticut; W. E. TREAT, Auk. ix, p. 389.
- Calidris arenaria, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxi: notes on its distribution; A. Newton, Add. to Ibis for July, 1892, p. 5; id. Nature, xlvi, p. 222; id. Rec. Austral. Mus. ii, p. 22: in the Malay Archipelago; H. Seebohm, Ibis, 1892, pp. 469-470.
- Tringoides hypoleucus, akeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxii, fig. 2.
- Limosa lapponica, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxii. Himantopus leucocephalus, in Norfolk Island, Lord Howe Island, and the New Hebrides; A. J. NORTH, Rec. Austral. Mus. ii, p. 38.
- Crymophilus fulicarius, in Nantucket Island, Massachusetts; G. H. MACKAY, Auk, ix, pp. 294-298: in Princetown Harbour, Mass.; G. S. MILLER, t. c. pp. 288 & 289: in Leicestershire; A. MATTHEWS, Zool. 1892, p. 26: in Warwickshire; F. COBURN, t. c. p. 26: in 1892. [VOL. XXIX.]

Ireland; R. Patterson, Zool. 1892, pp. 26, 32, & 33: in Islay, R. S. Skirving, Ann. Scott. N. H. i, p. 72: in Jura; H. Evans, t. c. p. 72: in S.W. Scotland; R. Service, t. c. p. 73: in Devon; W. E. H. Pidsley, t. c. p. 28. *P. hyperboreus*: notes on nesting and migration; S. von Chernel, C.R. ii, Congr. Orn. Intern. ii, pp. 137-144, pl. 2: in Ireland; E. Williams, Zool. 1892, p. 28.

#### Suborder PARRÆ.

Jacana spinosa, in Jamaica; T. D. A. COCKERELL, P. Z. S. 1892, p. 472.

#### Suborder ŒDICNEMI.

Œdicnemus crepitans, in Kent; W. VERNER, Zool. 1892, p. 24.

### Suborder OTIDES.

See Nicolls, J. A., and Eglington, W.

Otis tarda, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxi: in Orkney, T. PEACE, Ann. Scott. N. H. i, p. 138. O. tetrax, in Silesia; E. Urban, Orn. Jahrb. iii, p. 34: in Moravia; F. Schaller, ibid.: in Steiermark; A. J. Mojsisovics, ibid.: in Northern Dalmatia; H. Hänisch, t. c. pp. 112 & 113.

## Order GRUIFORMES.

Geranopsis elatus, n. sp. (fossil), A. MILNE-EDWARDS, C.B. ii, Congr. Orn. Intern. ii, pp. 72 & 73.

Grus leucogeranus, in the Hebrides [Zool. Rec. xxviii, Aves, p. 41], was an escaped specimen; W. E. CLARKE, Ibis, 1892, p. 181.

### Order PELARGIFORMES.

### Suborder ARDEÆ.

Hänisch, H. Ornithologische Mittheilungen aus dem unteren Narentathale. Reiher (Ardeidæ). Orn. Jahrb. iii, pp. 55-60.

Palæopelagus, n. g. Type, P. nobilis, n. sp. (fossil), Queensland; C. W. DE VIS, P. Linn. Soc. N.S.W. (2) vi, pp. 441-443, pl. xxiv, figs. 4a, 4a<sup>1</sup>.

Ardea amissa, n. sp. (fossil), Éocene of S. France; A. MILNE-EDWARDS, C.R. ii, Congr. Orn. Intern. ii, p. 73. A. paloccidentalis, n. sp. (fossil), Pliocene of Oregon; R. W. SHUFELDT, J. Acad. Philad. ix, p. 411.
A. garzetta, figured; Lord Lilford, Col. Fig. Br. Birds, part xxi.

Herodias egretta, in Norfolk I.; A. J. North, Rec. Austral. Mus. ii, p. 37.

Botaurus neoxenus: male described; W. E. D. Scott, Auk, ix, pp. 141 & 142.

### Suborder PLATALEÆ.

Platalea leucerodia, in Heligoland; H. GÄTKE, Orn. Jahrb. iii, p. 201: its breeding places in Holland; F. E. BLAAUW, Ibis, 1892, p. 468. P. subtenuis, n. sp. (fossil), Queensland, C. W. DE Vis, P. Linn. Soc. N.S.W. (2) vi, pp. 443 & 444, figs. 5a & 5b.

### Order TAPINOPIFORMES.

Tapinopus, n. g. Type, T. ellieti, n. sp. (fossil), Eocene of S. France;
A. MILNE-EDWARDS, C.B. ii, Congr. Orn. Intern. ii, pp. 79 & 80.

## Order PHÆNICOPTERIFORMES.

Phanicopterus copei, n. sp. (fossil), Equus Beds, Oregon; R. W. Shu-Feldt, J. Ac. Philad. ix, pp. 410 & 411.

### Order ANSERIFORMES.

- See Nicolle, J. A., and Eglington, W.
- Olor paloregonus (feesil): notes on; R. W. Shufeldt, J. Ac. Philad. ix, p. 409.
- Chemiornis gracilis and C. minor, n. spp. (fossil); H. O. FORBES, Tr. N. Z. Inst. xxiv, pp. 187 & 188.
- Bernicla rusicollis, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxi. B. munroii, n. sp., Kauai, Sandwich, Is., W. ROTHSCHILD, Ann. & Mag. N. H. (6) x, p. 108. B. leucopsis, in Venezia, E. ARRIGONI DEGLI ODDI, Atti Soc. Ital. xxxiv, pp. 117-121. B. propinqua, n. sp. (fossil), Equus Beds, Oregon, Shuffldt, J. Ac. Philad. ix, pp. 408 & 409.
- Anser anatoides, n. sp. (fossil), Pliocene of France; C. Déperer, C.R. cxiv, p. 690. A. cinereus, A. segetum, and A. albifrons, in Dalmatia; G. Kolambatovic, Glasnik. Naravosl. Druž. vi, pp. 168-171 (1891).
- Biziura lautouri, n. sp. (fossil), from New Zealand; H. O. FORBES, Tr. N. Z. Inst. xxiv, p. 188.
- Assertante melanoleuca, in Tasmania; W. F. Petterd, P. R. Soc. Tasm. 1888, pp. 91 & 92.
- Tadorna casarca: on the recent occurrence in the British Is.; F. M. OGILVIE, Zool. 1892, pp. 392-398: in Ireland; R. J. USSHER, t. c. pp. 334 & 335. T. rutila, in co. Donegal; J. S. Elliott, t. c. p. 311.
- Anas condoni, n. sp. (fossil), Equus Beds, Oregon, R. W. SHUFELDT, J. Ac. Philad. ix, pp. 406 & 407. A. laysanensis, n. sp., Laysan I., W. ROTHSCHILD, Bull. B. O. C., No. 4, p. xvii. A. angustirostris and Fuligula marila, in Bohemia; RIHA, Orn. Jahrb. iii, pp. 260 & 261. A. boschas × A. americana; D. G. Elliot, Auk, ix, pp. 160-166.
- Spatula clypeata nesting in Orkney; W. Harvey, Ann. Scott. N. H. i, p. 138: in co. Westmeath, F. J. Battersby, Zool. 1892, p. 311.
- 8. variegata in Tasmania; A. J. North, Rec. Austral. Mus. ii, p. 37.
- Harelda glacialis on the Suffolk Coast in Summer; F. M. OGILVIE, Zool. 1892, p. 109.
- Somateria mollissima, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxii.
- Aythya marila nearctica, n. sp. (fossil), Equus Beds, Oregon; R. W. Shupellot, J. Ac. Philad. ix, p. 405.

- Edemia: notes on the American species; G. TRUMBULL, Auk, ix, pp. 153-157. E. nigra, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxii: migration in spring; G. H. MACKAY, Auk, ix, pp. 292-294: nesting in Sussex; C. Fowler, Zool. 1892, p. 151. E. fusca, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxii.
- Erismatura australis in Tasmania; A. J. North, Rec. Austral. Mus. ii, p. 37.
- Clangula hyemalis, in New England; G. H. MACKAY, Auk, ii, pp. 330-337.
- Mergus serrator, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxi.

## Order PELECANIFORMES.

#### Suborder SULÆ.

Sula cyanops and S. fiber, nesting on Adéle I.; J. WALKER, Ibis, 1892, pp. 257 & 258.

#### Suborder PHALACROCORACES.

## Family Phalacrocoracidm.

Phalacrocorax carbo and P. graculus, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxii. P. gutturalis, n. sp., Victoria Nyansa, A. REICHENOW, J. f. O. 1892, pp. 133 & 134. P. novæ scalandiæ, n. var., major (fossil); H. O. FORBES, Tr. N. Z. Inst. xxiv, p. 189. P. sulcirostris, in Norfolk I.; A. J. NORTH, Rec. Austral. Mus. ii, p. 38.

## Family PLOTIDE.

Plotus melanogaster: notes on anatomy and osteology; F. E. Beddard, P. Z. S. 1892, pp. 291–296.

### Suborder PELECANI.

Pelecanus proavus, n. sp. (fossil), Queensland, C. W. DE VIS, P. Linn. Soc. N.S.W. (2) vi, pp. 444 & 445, figs. 6a & 6b. P. erythrorhynchus, in Maine; H. MERRILL, Auk, ix, p. 292.

### Suborder FREGATI.

Tachypetes minor nesting on Adéle I.; J. WALKER, Ibis, 1892, pp. 258-260.

### Order CATHARTIDIFORMES.

- Bendire, C. Life Histories of North American Birds, with special reference to their breeding habits and eggs. U.S. Nat. Mus., Special Bulletin No. 1, Washington, 1892, 4to.
- Eggs of the following species are figured:—Pseudogryphus californianus (pl. iv, fig. 5); Cathartes aura (pl. iv, figs. 1 & 3); Catharista utrata (pl. iv, figs. 7 & 10).

### Order ACCIPITRIFORMES.

- Bender, C. Life Histories of North American Birds, with special reference to their breeding habits and eggs. U. S. Nat. Mus., Special Bulletin No. i, Washington, 1892, 4to.
- Eggs of the following species are figured:—Elanoides forficatus (pl. v. figs. 1 & 2), Elanus leucurus (pl. v, figs. 3 & 4); Ictinia mississipiensis (pl. 5, fig. 5); Rosthramus sociabilis (pl. v, figs. 6 & 7); Circus hudsonius (pl. v, figs. 8-10); Accipiter velox (pl. v, figs. 11-17), A. cooperi (pl. v, figs. 18-20), A. atricapillus (pl. vi, fig. 1), A. atricapillus striatulus (pl. vi, fig. 2); Parabuteo unicinctus harrisi (pl. vi, figs. 3 & 3), Buteo borealis (pl. vi, figs. 5 & 6), B. borealis calurus (pl. vi, figs. 7 & 8), B. lineatus (pl. vii, figs. 1-5), B. lineatus elegans (pl. vi, fig. 9), B. abbreviatus (pl. vii, fig. 6), B. albicaudatus (pl. vii, figs. 8 & 9), B. swainsoni (pl. viii, figs. 1-6), B. latissimus (pl. vii, figs. 10-13), B. brachyurus (pl. viii, fig. 7); Urubitinga anthracina (pl. viii, figs. 8 & 9), Asturina plagiata (pl. vii, fig. 7), Archibuteo lagopus sanctijohannis (pl. viii, figs. 10-12), A. ferrugineus (pl. ix, figs. 1, 2, & 4); Aquila chrysuetos (pl. ix, figs. 3 & 5); Haliætus leucocephalus (pl. ix, fig. 7); Falco rusticolus gyrfalco (pl. ix, figs. 6, 8, & 9), F. rusticolus obsoletus (pl. x, fig. 1), F. mexicanus (pl. x, figs. 2 & 3), F. peregrinus anatum (pl. x, figs. 5-7), F. columbarius (pl. x, figs. 4 & 8), F. fusco-carulescens (pl. x, figs. 9 & 10), F. sparverius (pl. x, figs. 11-16); Polyborus cheriway (pl. xi, figs. 1-4); Pandion halidetus carolinensis (pls. x, fig. 17, & xi, figs. 5-9).
- CZATÓ, J. VON. Die Verbreitung und Lebendsweise der Tagraubvögel in Siebenbürgen. C.R. ii, Congr. Orn. Intern. ii, pp. 118-128.
- LILFORD, LORD. Notes on Birds in the Lilford Aviaries. J. Northampt. Soc. v, pp. 258-269.
- [Haliaetus albicilla, Aquila mogilnik, A. adulberti, A. ropax, A. nepalensis, A. pennata, Pseudaetus bonelli, Circaetus gallicus, Vultur monachus, Gypäetus barbatus, Neophron perenopterus.]
- MEARNS, E. A. A Study of the Sparrow Hawks (Subgenus Tunnunculus) of America, with especial reference to the Continental species (Falco sparrerius, Linn.). Auk, ix, pp. 252-270.
  - [See Falconida, p. 39.]
- Quelch, J. J. Our [British Guiana] Birds of Prey. Journ. R. Agr. & Comm. Soc. Brit. Guian. (2) vi, pp. 117-162.

# Family Vulturida.

Neophron pileatus: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. cli.

## Family FALCONIDE.

### Subfamily POLYBORINAS.

- Polyborus cherinosy: egg and chick; A. Dugès, Nat. Mex. (2) ii, pp. 173 & 174, pl. ix.
- Ibycter chimango: skeleton figured; A. B. Meyer, Abbild. Vögel. Skel., Taf. cliii.

### Subfamily ACCIPITRINA

- Circus cyaneus, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxi.
  O. hamiltoni and C. teautensis, n. spp. (subfossil), New Zealand;
  H. O. FORBES, Tr. N. Z. Inst. xxiv, p. 186.
- Astur brevipes, notes on; N. v. Ssomow, Orn. Jahrb. iii, pp. 179-182.
- Melierax polyzonus: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clix.

## Subfamily BUTEONINÆ.

- Busarellus nigricollis: skeleton figured; A. B. MEYER, Abbild. Vögel, Skel., Taf. cliv.
- Buteo borealis calurus in Cuba; F. M. CHAPMAN, Bull. Am. Mus. iv, p. 294. B. albicaudatus colonus, n. subsp., Curação; H. von Berlepsch, J. f. O. 1892, pp. 89-91.

### Subfamily AQUILINA

- Necrastur, n. g. Type N. alacer (fossil), Queensland; C. W. DE VIS, P. Linn. Soc. N.S.W. (2) vi, pp. 437-439, pl. xxiv, figs. 1a, 1b.
- Aquila hypogaa, n. sp. (fossil); A. MILNE-EDWAEDS, C.R. ii, Congr. Orn. Intern. ii, pp. 60 & 61. A. pliogryps, n. sp. (fossil), and A. sodalis, n. sp. (fossil), Equus Beds, Oregon, R. W. SHUFELDT, J. Ac. Philad. ix, pp. 416-418. A. rapax, in Astrachan; A. B. MEYER, Abh. Ges. Isis, 1892, pp. 67-71: notes on A. bocki, id.: t. c. A. navia, in Essex; MILLER CHRISTY, Zool. 1892, p. 76.
- Spizaetus coronatus, skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clvi.
- Nisaetus fasciatus, skeleton figured, id. t. c. Taf. clv.
- Spilornis rufipectus, skeleton figured, id. t. c. Taf. clvii.
- Circaetus gallicus, in Bohemia, F. v. Fiskal, Orn. Jahrb. iii, pp. 173 & 174: in the Forest of Andaine (Orne); Abbé Letace, Bull. Soc. L. Norm. vi, pp. 180-183.
- Haliaetus albicilla: notes on; V. RITTER VON TSCHUSI ZU SCHMID-HOFFEN, Orn. Jahrb. iii, pp. 74-76: figured, E. H. GIGLIOLI & A. Manzella, Iconogr. Avif. Ital. part xlix.
- Gypohierax angolensis, skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clii.
- Milvus migrans, figured; E. H. GIGLIOLI & A. MANZELLA, Iconogr. Avif. Ital. part xlvii.

## Family FALCONIDE.

- Pernis apivorus, in co. Wexford; E. WILLIAMS, Zool. 1892, p. 428.
- Baza lophotes: notes on its nest and eggs; T. A. HAUXWELL, J. Bomb. N. H. Soc. viii, pp. 403 & 404.
- Falco peregrinus, from Pappan I., N. W. Borneo, R. B. Sharpe, Bull. B. O. C. ii, p. iv. F. peregrinus anatum, skeleton figured; A. B. Meyer, Abbild. Vögel. Skel., Taf. clviii. F. barbarus and F. punicus, figured; E. H. Giglioli & A. Manzella, Iconogr. Avif. Ital., parts. xlvii & xlviii. F. rusticolus obsoletus, in Rhode I.; A. O'D. Taylor, Auk, ix, pp. 300 & 301.
- Gennaia saker and G. feldeggii, figured; E. H. GIGLIOLI & A. MANZELLA, Iconogr. Avif. Ital., part xlviii.
- Hierofalco candicans, in Achill; R. M. BARRINGTON, Zool. 1892, p. 77.
- Falco cenchrie, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxii.
- Falco tinnunculus × F. æsalon; George Bolam, Naturalist, 1892, pp. 369 & 370. F. sparverioides: notes on; F. M. Chapman, Bull. Am. Mus. iv, pp. 294-296. F. sparverius deserticolus, n. subsp., Arizona; E. A. Mearns, Auk, ix, p. 263. F. sparverius peninsularis, n. subsp., Lower California, id. t. c. p. 267. F. sparverius australis, n. subsp., S. America, id. t. c. p. 269. F. sparverius æquatorialis, n. subsp., Guayaquil, Ecuador, id. t. c. p. 269. F. sparverius brevipensis, n. subsp., Curação; H. von Berlepsch, J. f. O. 1892, pp. 91-95.

#### Suborder PANDIONES.

Pandion haliactus, figured; E. H. GIGLIOLI & A. MANZELLA, Icongr. Avif. Ital., part xlvii: formerly nesting in the Lake District of England; A. G. More, Zool. 1892, pp. 1-3: breeding on Plum I., N. Y.; C. S. Allen, Auk, ix, pp. 313-321, pls. iv & v.

### Order STRIGES.

- BENDIRE, C. Life Histories of North American Birds, with special reference to thier breeding habits and eggs. Washington: 1892, 4to, viii & 414 pp., pls. i-xii.
- The eggs of the following species are figured:—Strix pratincola (pl. xii, fig. 1); Asio wilsonianus (pl. xii, fig. 2), A. accipitrinus (pl. xii, fig. 3); Syrnium nebulosum (pl. xii, fig. 4); Scotiaptex cinerea (pl. xii, fig. 5); Nyctala tengmalmi richardsoni (pl. xii, fig. 6), N. acadica (pl. xii, fig. 7); Megascops asio (pl. xii, fig. 8), M. asio maxwelliæ (pl. xii, fig. 10), M. asio trichopsis (pl. xii, fig. 11), M. asio macfarlanei (pl. xii, fig. 9), M. flammeolus (pl. xii, fig. 15); Bubo virginianus (pl. xii, fig. 12); Nyctea nyctea (pl. xii, fig. 19); Surnia ulula caparoch (pl. xii, fig. 18); Spectyto cunicularia hypogea (pl. xii, fig. 14), S. cunicularia floridana (pl. xii, fig. 16); Glaucidium phalænoides (pl. xii, fig. 17); Micropallas whitneyi (pl. xii, fig. 13).

- Bubo maximus, figured; E. H. GIGLIOLI & A. MANZELLA, Iconogr. Avif. Ital, part xlvii. B. incertus, n. sp. (fossil), Eccene of S. France; A. MILNE-EDWARDS, C. B. ii, Congr. Orn. Intern. ii, pp. 63 & 64.
- Necrobyas harpax, n. sp. (fossil), and N. rossignoli, n. sp. (fossil), Eccene of France, id. t. c. pp. 61-63.
- Megascops flummeolus idahoensis, figured: C. H. MERRIAM, Auk, ix, pp. 169-171, pl. ii.
- Scops menadensis, skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxi, fig. 2. S. mantananensis, n. sp., Mantanani I., N. W. Borneo; R. B. SHARPE, Bull. B. O. C., No. 2, p. iv. S. brookii, n. sp., Mt. Dulit, Borneo, id. ibid.
- Nyctea ulula, in Austrian Silesia; E. Urban, Orn. Jahrb. iii, p. 122.
- Asio accipitrinus, increase of nests in vole-stricken districts of Scotland; P. Adair, Ann. Scott. N. H. i, pp. 219-231.
- Otus (potius Asio) henrici, n. sp. (fossil), Eocene of S. France, A. MILNE-EDWARDS, C.R. ii, Congr. Orn. Intern. ii, p. 63.
- Carine noctua, in Oxfordshire; O. V. APLIN, Zool, 1892, p. 332.
- Athene noctua plumipes, skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clx, fig. 2.
- Ninox connivens, egg figured; A. J. NORTH, Rec. Austral. Mus. ii, pl. ii, fig. 1.
- Spectyto cunicularia bahamensis, = S. cunicularia floridanus; C. B. CORY, Cat. W. Ind. Birds, p. 140. S. cunicularia floridana, its breeding habits; S. N. RHOADS, Auk, lx, pp. 1-8.
- Syrnium occidentale: notes on; F. STEPHENS, Auk, ix, pp. 392 & 393.

  S. aluco, figured; A. H. GIGLIOLI & A. MANZELLA, Iconogr. Avif.

  Ital., part xlvii.
- Strix flammea bargei, n. subsp., Curação I.; E. HARTERT, Bull. B. O. C., No. 3, p. xiii.

#### Order CORACIIFORMES.

#### Suborder STEATORNITHES.

HARTERT, E. Catalogue of the *Picaria* in the Collection of the British Museum. Coracia, Fam. *Steatornithida*. [Vol. xvi of The Catalogue of Birds.] London: 1892, 8vo, pp. 653 & 654.

### Suborder PODARGI.

- HARTERT, E., ut suprà: Fam. Podargidæ, pp. 629-636 & 652, pl. xiv. Podargus strigoides, remarks on; W. SAVILLE KENT, P. Z. S. 1892, p. 477.
- Batrachostomus harterti, n. sp., Mt. Dulit, N. W. Borneo; R. B. SHARPE, Ibis, 1892, p. 323: figured, E. HARTERT, Cat. Birds, xvi, pl. xiv. B. poliolophus, n. sp., W. Sumatra; E. HARTERT, Notes Leyd. Mus. xiv, p. 63. B. mixtus, n. sp., Mt. Dulit, N. W. Borneo; R. B. SHARPE, Bull. B. O. C., No. 2, p. iv.
- Egotheles salvadorii, n. sp., S. E. New Guinea; E. HARTERT, Cat. Birds, xvi, p. 649.

#### Suborder CORACIÆ.

## Family Coraciida.

- SHARPE, R. B. Catalogue of the *Picariæ* in the Collection of the British Museum. Coraciæ and Halcyones. *Coraciidæ*. [Vol. xvii of The Catalogue of Birds.] 1892, pp. 4–40, pls. ii & iii.
- Geranopterus, n. g. Type, G. elutus, n. sp. (fossil), Eocene of S. France;
  A. MILNE-EDWARDS, C.R. ii, Orn. Congr. Intern. ii, p. 66.
- Coracias olivaceiceps, n. sp. [nom. emend. pro C. mosambicus, Dresser];
  R. B. Sharpe, t. c. p. 25. C. garrula, in Ireland; R. L. Patterson,
  Ibis, 1892, p. 181: in Caithness; L. Dunbar, Ann. Scott. N. II.
  i, p. 136.
- Eurystomus orientalis (pl. ii, fig. 1), E. calonyx (pl. ii, fig. 2), E. solomonensis [Zool. Rec. xxvii, Aves, p. 38] (pl. iii, fig. 1), E. azureus (pl. iii, fig. 2), figured; R. B. SHARPE, ut suprà. E. afer, n. var. rufobuccalis, Uganda; A. Reichenow, J. f. O. 1892, p. 27.

#### Suborder LEPTOSOMATI.

## Family LEPTOSOMATIDE.

SHARPE, R. B. Catalogue of the *Picariæ* in the Collection of the British Museum. Coraciæ and Halcyones. *Leptosomatidæ*. [Vol. xvii of The Catalogue of Birds.] 1892, pp. 1-3.

#### Suborder HALCYONES.

## Family ALCEDINIDE.

SHARPE, R. B. Catalogue of the *Picariæ* in the Collection of the British Museum. Coraciæ and Halcyones. *Alcedinidæ*. [Vol. xvii of The Catalogue of Birds.] 1892, pp. 93-312.

### Subfamily ALCEDININÆ.

- Ceryle septentrionalis, n. subsp., nom. emend. pro C. cabanisi (auct., nec Tschudi), S. United States to Panama; id. t. c. pp. 134-136. C. aquatorialis, n. sp., Ecuador, id. t. c. p. 140, pl. iv, fig. 1. C. stictoptera, figured; id. t. c. pl. iv, fig. 2.
- Alcedo floresiana, n. subsp., Flores; id. t. c. p. 151. A. guentheri, n. sp., W. Africa, id. t. c. pl. iv, fig. 2. Distinct from A. quadribrachys, (pl. iv, fig. 1), id. t. c. pp. 156 & 157. A. ispida: nesting habits; R. Scobie, P. N. H. Soc. Glasg. (n.s.) iii, p. 255.

### Subfamily DACELONINÆ.

Ceyx everythra, n. sp., Malaisia; R. B. Sharpe, Cat. B. Brit. Mus. xvii, p. 179. C. sumarensis [Zool. Rec. xxvii, Aves, p. 38] = C. melanura; id. t. c. p. 180. C. sulaensis [Zool. Rec. xxvii, Aves, p. 38] = C.

malamaui [Zool. Rec. xxvii, Aves, p. 38], id. t. c. p. 184. C. margarethæ [Zool. Rec. xxvii, Aves, p. 38] = C. bournsii [Zool. Rec. xxvii, Aves, p. 38]; id. t. c. p. 185. C. steerii, n. sp., Mindoro, id. t. c. p. 187.

Halcyon torquatus (pl. vi, fig. 1), H. forbesi, n. sp., Sierra Leone (pl. vi, fig. 2), H. malimbicus (pl. vi, fig. 3): heads figured; R. B. SHARPE, t. c. pp. 246 & 247, pl. vi. H. admiralitatis, n. sp., Admiralty Is., id. t. c. p. 251. H. rufigularis, n. subsp., hab. ignot., id. t. c. p. 260. H. mediocris, n. subsp., Ponapé, id. ibid. H. tutuilæ, n. sp., Tutuila, id. t. c. p. 266. H. tannensis, n. sp., Tanna I., id. ibid. H. chloris: head figured; id. t. c. pl. vii, fig. 3. H. armstrongi, n. subsp., Burmah, id. t. c. p. 277, pl. vii, fig. i. H. vidali, n. subsp., S. Konkan, id. t. c. p. 278. H. solomonis, head figured; id. t. c. pl. vii, fig. 2. H. suvensis, n. sp., Suva I., id. t. c. p. 281. H. humii, n. sp., Malacca, id. t. c. p. 281, pl. viii. H. meyeri, n. subsp., Togian Is., id. t. c. p. 282. H. davisoni, n. subsp., Andamans, id. t. c. p. 282. H. (Cyanalcyon) quadricolor, figured; E. Oustalet, N. Arch. Mus. (2) (3) iv, pl. xiv.

Todirhamphus youngi, n. sp., Morea I., p. 289; R. B. Sharpe, ut supra, p. 289.

#### Suborder BUCEROTES.

## Family BUCEROTIDE.

OGILVIE-GRANT, W. R. Catalogue of the *Picariæ* in the Collection of the British Museum. Coraciæ and Halcyones. *Bucerotidæ*. [Vol. xvii of The Catalogue of Birds.] 1891, pp. 347-428, pls. xiii & xiv.

Buceros exaratus: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel, Taf. clxix.

Gymnolæmus, n. g. Type, G. marchii, W. R. OGILVIE-GRANT, t. c. p. 370. Ptilolæmus, n. g. Type, P. tickelli; id. t. c. p. 392.

Lophoceros damarensis, figured; id. t. c. p. 411, pl. xiv. L. flavirostris, id. t. c. p. 412, L. leucomelas, p. 414, and L. elegans, p. 415: tails figured; id. t. c. p. 413. L. jacksoni [Zool. Rec. xxviii, Aves, p. 49], figured; id. t. c. p. 416.

Ortholophus, n. g. Type, O. albocristatus; id. t. c. p. 424.

Tockus gingalensis: nest and eggs; F. Lewis, J. A. S. (Ceylon) xii, pp. 63-65.

#### Suborder UPUPÆ.

Salvin, O. Catalogue of the *Picaria* in the Collection of the British Museum. Upupe. [Vol. xvi of The Catalogue of Birds.] 1892, pp. 1-26, pls. i-iii.

## Family UPUPIDE.

5 species recognized; O. Salvin, ut suprà. U. somalensis, n. sp., Somaliland, id. t. c. p. 13, pl. i.

## Family IRRISORIDE.

- 3 genera recognized: Irrisor, Scoptelus, and Rhinopomastus; O. Salvin, t. c. pp. 16-26.
- Irrisor bollii (pl. ii), and I. jacksoni [Zool. Rec. xxvii, Aves, p. 39], pl. iii, fig. 1, figured; id. t. c.
- Scoptelus castansiceps, figured; id. t. c. pl. iii, fig. 2. S. anchietæ, n. sp., Benguela, J. v. Barboza du Bocage, J. Sci. Lisb. (2) No. viii, p. 254.

### Suborder MEROPES.

## Family MEROPIDE.

- SHARPE, R. B. Catalogue of the *Picaria* in the Collection of the British Museum. Coracia and Halcyones. *Meropida*. [Vol. xvii of The Catalogue of Birds.] 1892, pp. 41-92, pl. i.
- Dicrocercus furcatus (pl. i, fig. 2), distinct from D. hirundineus (pl. i, fig. 1): both species (heads) figured; id. t. c. pp. 42-44.
- Melittophagus meridionalis, n. sp. (pl. i, fig. 4), distinct from M. pusillus (pl. i, fig. 5), and M. cyanostictus (pl. i, fig. 3): heads of all 3 species figured; id. t. c. pp. 45-48. M. oreobates, n. sp., Mt. Elgon, E. C. Africa; id. Ibis, 1892, pp. 320 & 321.
- Merops apiaster, in Moravia; J. Talský, Orn. Jahrb. iii, p. 195: in co. Wicklow; E. Williams, Zool. 1892, p. 428. M. philippinus, skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxii, fig. 1.

#### Suborder MOMOTI.

## Family Momotida.

- SHARPE, R. B. Catalogue of the *Picaria* in the Collection of the British Museum. Coracise and Halcyones. *Momotida*. [Vol. xvii of The Catalogue of Birds.] 1892, pp. 313-332.
- Momotus parensis, n. sp., Para; id. t. e. p. 320. M. bartletti, n. sp., Colombia to Upper Amazonia; id. t. c. pp. 320 & 321, pl. ix. M. subrufescens, figured; id. t. c. pl. x, fig. i. M. microstephanus figured; id. t. c. pl. x, fig. 2. M. squatorialis, figured; id. t. c. pl. xi. M. venezuela, n. sp., Venezuela; id. t. c. p. 321.
- Aspatha, n. g. Type, A. gularis (Lafresn.); id. t. c. p. 331.

#### Suborder TODI.

## Family Todida.

SHARPE, R. B. Catalogue of the *Picariæ* in the Collection of the British Museum. Coraciæ and Halcyones. *Todidæ*. [Vol. xvii of The Catalogue of Birds.] 1892, pp. 333-337.

#### Suborder CAPRIMULGI.

HARTERT, E. Notes on the Caprimulgidæ. Ibis, 1892, pp. 274 288, pl. vi. Caprimulgus europæus and its allies; id. t. c. pp. 277-279. C. tristigma, Rüpp., = C. lentiginosus, Smith; id. t. c. p. 279. C. eximius, figured; id. t. c. pp. 279 & 280, pl. vi. C. asiaticus: notes on; id. t. c. p. 280. C. arundinaceus, Jacq. & Pucher, and C. faberi, Meyer, = C. affinis, id. t. c. pp. 280 & 281. C. salvadorii, Sharpe, = C. macrurus, of which C. albonotatus is only a race; id. t. c. pp. 281-283. C. keluarti is a race of C. jotaka; id. t. c. pp. 283-285. C. melanopogon, Salvad., = C. jotaka; id. t. c. p. 285. C. concretus: notes on; id. t. c. p. 285.

Antrostomus, = Caprimulgus; id. t. c. p. 286. A. vociferus, arizonæ, vocifer, = C. macromystax, Wagl.; id. t. c. p. 286. C. salvini, n. sp. (=C. macromystax, auct., nec Wagl.) id. t. c. p. 287.

Scotornis nigricans, = S. climacurus; id. t. c. pp. 287 & 288.

HARTERT, E. Catalogue of the Picariæ in the Collection of the British Museum. Coraciæ. Family Caprimulgidæ. [Vol. xvi of The Catalogue of Birds.] 8vo, pp. 519-652, pls. xi-xiii.

Caprimulgus griseatus, figured; id. t. c. pl. xi. C. whiteleyi, n. sp., Roraima; id. t. c. pp. 573 & 574, pl. xii. C. clarus, n. sp., Uganda; A. REICHENOW, Ber. Allg. Deutschl. Orn. Ges., Jan. 1892, p. 1; id. J. f. O. p. 29. C. europæus, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxii; E. H. GIGLIOLI & A. MANZELLA, Iconogr. Avif. Ital. pt. l.

C. ruficollis and C. asiaticus, figured; iid. t. c. pt. 1.

Lyncornis mindanensis, figured; E. HARTERT, t. c. pl. ziii.

#### Suborder CYPSELI.

HARTERT, E. Catalogue of the *Picariæ* in the Collection of the British Museum. Coraciæ. Family *Cypselidæ*. [Vol. xvi of The Catalogue of Birds.] London: 1892, pp. 434-518, pl. x.

Micropus africanus, distinct from M. apus; id. t. c. pp. 440 & 441.

Aeronautes, n. g. Type, A. melanoleucus; id. t. c. pp. 459 & 460.

Claudia, n. g. Type, C. squamata; id. t. c. pp. 469 & 470.

Chetura guianensis, n. subsp., Guiana; id. t. c. p. 486. O. ussheri, figured; id. t. c. pl. x.

Collocalia merguiensis, n. subsp., Tenasserim; id. t. c. p. 506.

[See RIDGWAY, R., Am. Nat. xxvi, pp. 1040 & 1041; SHUFELDT, R. W., t. c. pp. 869 & 870.]

Cypselus affinis, figured; E. H. GIGLIOLI & MANZELLA, Iconogr. Avif. Ital. pt. xlviii.

Collocalia in the Andaman Is.; [Anon.] Ibis, 1892, pp. 578 & 579.

Tachyornis, n. g. Type, T. hirundo, n. sp. (fossil), Eocene of S. France;
A. MILNE-EDWARDS, C.R. ii, Congr. Orn. Intern. ii, pp. 66 & 67.

[Ægiulornis (Zool. Rec. xxviii, Aves, p. 39) is identical; id. t. c.]

#### Suborder TROCHILL.

Food: Bolles, F., Science, xx, p. 318, and Gibbs, M., t. c. p. 329.

BOUCARD, A. Genera of Humming Birds, pp. 1-56, 8vo [published in the 'Humming Bird,' ii, Jan.-July, 1892].

Lophornis hauxwelli, n. sp., Nauta. id. t. c. p. 37.

- —. A complete list up to date of the Humming Birds found in Columbia, with descriptions of several supposed new species. T. c. pp. 73-87.
- The following species are supposed to be new:—Heliodoxa berlepschi, n. sp., Heliotrypha simoni, n. sp., Heliangelus rothschildi, n. sp., Eriocnemis albigularis, n. sp., Chlorostillon speciosa, n. sp., Saucerottia nunesi, n. sp., Uranomitra colombiana, n. sp., Phaethornis panamensis, n. sp., Homophania lawrencii, n. sp.
- RIDGWAY, R. The Humming Birds. Rep. U.S. Nat. Mus. 1890, pp. 253-383, pl. xlv, and 47 cuts in the text. A review of the family of *Trochilida*, with special reference to the species of the United States. T. c. pp. 255-292.
- Notes on the anatomy, by F. A. Lucas.

  [See also Am. Nat. xxvi, pp. 1040 & 1041.]
- SALVIN. O., & GODMAN, F. D. Biologia Centrali-americana., Aves, Vol. ii, pp. 249-366.
- The following species are figured:—Oreopyra calolama (pl. liv, figs. 1 & 2); Delattria hemileuca (pl. liv, figs. 3 & 4), D. margaretha and D. sybilla (pl. livA); Hemistephania veraguensis (pl. lv, fig. 1); Hypuroptila (s. n. Chalybura) melanorrhoa (pl. lv, figs. 2 & 3); Callipharus (s. n. Eupherusa) nigriventris (pl. lvii, figs. 3 & 4); Selasphorus torridus and S. ardens (pl. lvi); Lophornis adorabilis (pl. lvii, figs. 1 & 2).
- Salvin, O. Catalogue of the *Picariæ* in the Collection of the British Museum. [Vol. xvi of The Catalogue of Birds.] *Trochili*, pp. 27-433, pls. iv-ix.

Panychlora micans, n. sp., hab.?, id. t. c. p. 71, pl. iv, fig. 1.

Chlorostilbon peruanus, figured; id. t. c. pl. iv, fig. 2.

Neolesbia, n. g. Type, N. nehrkornæ; id. t. c. p. 145.

Heliangelus laticlavius, Ecuador, id. t. c. p. 160, and H. violicollis, n. spp., p. 162, pl. v, figs. 1 & 2.

Heliotrypha barrali and H. speciosa, n. sp., Colombia?, figured, id. t. c. pl. vi, figs. 1 & 2.

Iolama luminoso, figured; id. t. c. pl. viii, fig. 1.

Phaolama cerrinigularis, n. sp., Ecuador?, id. t. c. p. 325, pl. viii, fig. 2.

Eriocnemis ventralis, n. sp., Colombia, id. t. c. p. 364, pl. ix, fig. 2. E. dyselius figured; id. t. c. pl. ix, fig. 1.

Agyrtria cæruleiceps, figured; id. t. c. pl. vii, fig. 1.

Amazilia sumichrasti, n. sp., S. Mexico, id. t. c. p. 213, pl. vii, fig. 2.

- Polycrata decora, n. sp., Chiriqui, id. t. c. p. 238.
- Oreopyra pectoralis, n. sp., Costa Rica, id. t. c. p. 308.
- See Shuffeldt, R. W., Am. Nat. xxvi, pp. 869 & 870; Nature, xlvi, p. 465.
- Cyanolesbia emmæ, n. sp., Colombia, H. von Berlepsch, Ber. Allg. Deutsch. Orn. Ges., Dec. 1892, p. 10. C. caudata, n. sp., Venezuela, id. t. c. p. 12.
- Delattria sybillæ, n. sp., Nicaragua, O. Salvin & F. D. Godman, Ibis, 1892, p. 327.
- Cyanomyia guerrerensis, n. sp., Guerrero, iid. t. c. p. 290; iid., Biol. Centr.-Am. Aves, ii.
- Phaethornis rupurumii, n. sp., British Guiana, A. BOUCARD, Humming Bird, ii, p. 1.

### Suborder COLII.

## Family Colidz.

- SHARPE, R. B. Catalogue of the *Picaria* in the Collection of the British Museum. *Collidæ*. [Vol. xvii of The Catalogue of Birds.] 1891, pp. 338-346, p. xii.
- C. leucotis (fig. 1), and C. affinis (fig. 2), figured; id. t. c. pl. xii.
- Colius nigriscapalis, n. sp., Cameroons, A. REICHENOW, Ber. Allg. Deutsch. Ges., Feb. 1892, p. 3; id. J. f. O. 1892, p. 180.

### Order TROGONES.

# Family Trogonidz.

- OGILVIE-GRANT, W. R. Catalogue of the *Picaria* in the Collection of the British Museum. Coracise and Halcyones. *Trogonida*. [Vol. xvii of The Catalogue of Birds.] 1891, pp. 429-497, pls. xiii-xvii.
- Archæotrogon, n. g. Type, A. venustus, n. sp. (fossil), Eccene of S. France; A. MILNE-EDWARDS, C.R. ii, Congr. Orn. Intern. ii, pp. 64-66.
- Trogon bolivianus, n. sp., Amazonia and Ecuador, W. R. OGILVIE GRANT, t. c. p. 473, pl. xv.
- Hapaloderma vittatum, figured; id. t. c. pl. xvi.
- Harpactes vidua, n. sp., Mt. Dulit, Borneo, id. t. c. p. 501. H. dulitensis, n. sp., Mt. Dulit, id. t. c. p. 502, pl. xvii.

## Order COCCYGES.

# Family Cuculidae.

- Dynamopterus, n. g. Type, D. veloz, n. sp. (fossil); A. MILNE-EDWARDS, C.R. ii, Congr. Orn. Intern. ii, p. 64.
- Cuculus canorus: two eggs in a nest; V. CAPER, Orn. Jahrb. iii, pp. 248-

- 254. C. canorus and Hirundo rustica reared in the same nest; C. Wolley-Dod, Ibis, 1892, pp. 524-530. C. aurivillii, n. sp., Cameroons, Y. Sjöstedt, J. f. O. 1892, p. 313.
- Cacomantis virescens: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxxiii, fig. 2.
- Coccysus americanus occidentalis, in Washington; R. H. LAWRENCE, Auk, ix, pp. 393 & 394.
- Centropus phasianus: egg described and figured; A. J. North, Rec. Austral. Mus. ii, p. 17, pl. ii, fig. 4.
- Pyrrhocentor celebensis: akeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxxii.
- Guira guira: skeleton figured; id. t. c. Taf. clxxi, fig. 2.

## Family MUSOPHAGIDA.

CHURCH, A. H. Researches on Turacin, an animal pigment containing copper. Phil. Trans. clxxxiii, pp. 511-530.

#### Order PSITTACIFORMES.

A. L. H[ERRANO]. Sinopsis de los Psitacidos mexicanos. Nat. Mex. (2) ii, pp. 213-218.

## Family NESTORIDE.

Nestor norfolcensis: notes on a specimen; H. B. Tristram, Ibis, 1892, pp. 557 & 558.

# Family CACATUIDE.

Calyptorhynchus funereus and C. solandri: eggs described and figured;
A. J. North, Rec. Austral. Mus. ii, p. 18, pl. ii, figs. 2 & 3,

# Family PSITTACIDE.

## Subfamily Conurinz.

Conurus carolinensis: its range and habits; A. W. BUTLER, Auk, ix, pp. 49-56: in Missouri; C. H. MERRIAM, t. c. p. 301. C. rubritorques, Scl., distinct from C. holochlorus; T. SALVADORI, Bull. B. O. C., No. 3, p. xi. C. arubensis, n. sp., Aruba I., E. HARTERT, op. cit. No. 4, p. xvi. C. gundlachi, from Mona = C. chloropterus of San Domingo; C. CORY, Auk, ix, pp. 228 & 229.

#### Subfamily PIONINÆ.

Chrysotis kecki = C. lilacina (Lees.); T. Salvadori, Ibis, 1892, pp. 467 & 468. C. rothschildi, n. sp., Bonaire I., E. Hartert, Bull. B. O. C., No. 3, p. xiii.

### Subfamily PALEORNITHINE.

- Geoffroyius tjindanæ, u. sp., Sumba I., A. B. MEYER, Notes Leyd. Mus. xiv, p. 267.
- Polytelis alexandra: egg described and figured; A. J. NORTH, Rec. Austral. Mus. ii, p. 19, pl. 2, fig. 5.
- Loriculus stigmatus: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxx, fig. 1.

## Subfamily PLATYCERCINA.

- Platycercus erythrotis, from Antipodes I.; H. O. FORBES, Tr. N. Z. Inst. xxiv, pp. 190-192.
- Neophema elegans: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxx, fig. 2.

### Order SCANSORES.

#### Suborder CAPITONES.

- Pogonorhynchus rubrifacies, n. sp., Kimoani, Equat. Africa; A. REICH-ENOW, Ber. Allg. Deutsch. Orn. Ges., Jan. 1892, p. 1: id. J. f. O. 1892, p. 25.
- Mesobucco eximius, n. sp., Mt. Dulit, N. W. Borneo, R. B. SHARPE, Ibis, 1892, p. 324: figured, id. t. c. pl. xi.
- Trachyphonus erythrocephalus, T. boehmi and T. arnaudi: notes on, id. t. c. pp. 310 & 311.
- Barbatula ugandæ, n. sp., Uganda; A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., Jan. 1892, p. 1: id. J. f. O. 1892, p. 25. B. coryphæa, n. sp., Cameroons, id. Ber. Allg. Deutsch. Orn. Ges., Feb. 1892, p. 3, and J. f. O. 1892, p. 181, Tab. ii, fig. 2. B. leucomystax, n. sp., Sotik, E. C. Africa, R. B. SHARPE, Ibis, 1892, p. 311.

#### Suborder INDICATORES.

Indicator pygmæus, n. sp., Victoria Nyanza, A. Reichenow, J. f. O. 1892, p. 132.

#### Suborder PICI.

## Family PICIDE.

#### Subfamily PICINAL

- Celeus kerri [Zool. Rec. xxviii, Aves, p. 55], figured; J. Graham Kerr, Ibis, 1892, pl. iii.
- Brachypternus ceylonus: eggs and nesting; F. Lewis, J. A. S. (Ceylon) xii, pp. 63-65.
- Picus Isuconotus, in Carinthia; F. SCHULZ, Orn. Jahrb. iii, p. 203. P. major, in Scotland; J. A. HARVIE BROWN, Ann. Scott. N. H. i, pp. 1-17.

- Sphyrapicus varius: young in captivity; F. Bolles, Auk, ix, pp. 109-119. Thriponax kalinouskii, figured; C. W. Campbell, Ibis, 1892, pl. v; not distinct from T. richardsi, Tristr.; H. Seebohm, t. c. pp. 248-250: P. Z. S. 1892, p. 195: from Tsu-sima, id. Ibis, 1892, p. 399: figured; J. Ijima, J. Sci. Coll. Tokio, v, pl. xii.
- Conturus bahamensis, n. sp., Great Bahama I., Bahamas, C. Cory, Auk, ix, p. 270.
- Campothera tullbergi, n. sp., Cameroons, Y. Sjöstedt, J. f. O. 1892, pp. 313 & 314.
- Colaptes: remarks on the North American species; J. A. Allen, Bull. Am. Mus. iv, Art. ii, pp. 21-44, pl. ii.

### Subfamily PICUMNINÆ.

Picumnus obsoletus, n. sp., Venezuela, J. A. Allen, Bull. Am. Mus. iv, Art. v, p. 55.

### Order EURYLÆMI.

Calyptomena hosii, n. sp., Mt. Dulit, Sarawak, R. B. SHARPE, Ann. & Mag N. H. (6) ix, p. 249: figured, Ibis, 1892, pl. xi.

### Order MENURÆ.

Menura victoriæ: nest figured; A. J. CAMPBELL, C.R. ii, Congr. Orn. Intern. ii, p. 156.

#### Order PASSERIFORMES.

#### Section A. OSCINES.

# Family Corvide.

- Polæocorax, n. g. Type, Corvus moriorum, Forbes; H. O. Forbes, Bull. B. O. C., No. 4, p. xxi.
- Corvus procorax, n. sp. (fossil), Pliocene of France, see C. Déperet, C.R. cxiv, p. 690. C. annectens, n. sp. (fossil), Pliocene of Oregon, R. W. Shufeldt, J. Ac. Philad. ix, p. 419. C. intermedius, remarks on; W. E. Brooks, Ibis, 1892, p. 59.
- Trypancorax frugilegus: on the feathered bill; R. SERVICE, Ann. Scott. N. H. i, pp. 218 & 219.
- Nucifraga caryocatastes: breeding in East Prussia; E. HARTERT, Ibis, 1892, p. 370.
- Garrulus glandarius: migration in Belgium; E. DE SÉLYS-LONGCHAMPS, C.B. ii, Congr. Orn. Intern. ii, p. 178.
- Aphelocoma californica, in Washington; R. H. LAWRENCE, Auk, ix, p. 301.
- Cyanocorax eyanopogon: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clx, fig. 1.

1892. [vol. xxix.]

Perisoreus canadensis, in Massachusetts; R. O. Morris, Auk, ix, p. 395. Crypsirhina nigra, n. sp., Hainan, F. W. Styan, Bull. B. O. C., No. 2, p. vi [? = Temnurus truncatus, R. B. Sharpe, op. cit. No. iv, p. xi.] Pyrrhocorax primigenius (fossil), in France; E. Rivière, C.R. Ass. Fr. Sci. xx, p. 400.

## Family PARADISEIDÆ.

- Ptilorhis victoriæ: nest and egg; A. J. CAMPBELL, Vict. Nat. viii, pp. 134 & 135.
- Paradisea raggiana: nest and egg; P. Crowley, Bull. B. O. C., No. 4, p. xvi (3), v, pp. 218-220, pl. xv.
- Craspedophora mantoui, figured; E. OUSTALET, N. Arch. Mus.
- Diphyllodes chrysoptera septentrionalis, n. subsp., N. E. New Guinea; A. B. MEYER, J. f. O. 1892, pp. 260-262.
- Manucodia comrii: nest and egg described; A. J. NORTH, Bec. Austral. Mus. ii, p. 32.

## Family PTILONOBHYNCHIDE.

- Amblyornis subalaris: plan of its bower; C W. DE VIS, Ann. Report Brit. New Guinea, 1890-91, App. cc, p. 95, cum fig.
- Chlamydodera cerviniventris and C. maculata: notes on eggs; P. CROWLEY, Bull. B. O. C., No. 4, p. xvi.

## Family STURNIDE.

- KOEPERT, O. Der Star (Sturnus vulgaris) in volkswirtschaftlicher und biologische Beziehung. Altenburg: 1892, 115 pp.
- Sturnus vulgaris, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxii.

# Family EULABETIDE.

- Acridotheres torquatus, n. sp., Pahang, E. Malay Archipelago; W. R. DAVISON, Ibis, 1892, pp. 102 & 103: is an Æthiopear; R. B. SHARPE, Bull. B. O. C., No. 2, p. vii.
- Calornis enganensis, n. sp., I. of Engano; T. SALVADORI, Ann. Mus. Genov. (2) xii, p. 137.
- Onychognathus preussi, n. sp., Cameroons; A. REICHENOW, Ber. Allg. Deutsch. Ges., Feb. 1892, p. 4; id. J. f. O. 1892, p. 184.
- Gracula enganensis, n. sp., I. of Eugano; T. Salvadori, Ann. Mus. Genov. (2) xii, pp. 137 & 138.

# Family PARAMYTHIIDE.

Paramythia, n. g. Type, P. montium, n. sp., Mt. Suckling, S. E. New Guinea; C. W. De Vis, Ann. Rep. Brit. New Guinea, App. cc, p. 95.

## Family DICEURIDE.

Chibia bracteata, in Tasmania; W. V. LEGGE, P. R. Soc. Tasm. 1888, pp. xix, xx & pp. 93-94 (1889).

## Family ORIOLIDE.

- Oriolus hosii, n. sp., Mt. Dulit, Borneo; R. B. SHARPE, Bull. B. O. C., No. 2, p. iv.
- Oriolus celebensis: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxvi, fig. 1.

## Family ICTERIDE.

- Quiscalus: notes on the species; F. M. CHAPMAN, Bull. Am. Mus. iv, pp. 1-16, pl. i.
- Ptiloxena, n. g. Type, P. atroviolacea, of Cuba; id. t. c. p. 307.
- Scolecophagus affinis, n. sp. (fossil), Equus Beds, Oregon; R. W. SHUFELDT, J. Ac. Philad. ix, pp. 418 & 419.
- Dives kalinowskii, n. sp., Ica, Peru; H. von Berlepsch & J. Stolzmann, P. Z. S. 1892, p. 378.
- Molothrus occidentalis, n. sp., Lima, Peru; iid. t. c. p. 378. M. ater, in Mexico; A. Dugès, Nat. Mex. (2) ii, pp. 103-106. M. robustus, parasitic on Buarremon gutturalis; G. K. Chebrik, Auk, ix, pp. 25-27.

# Family PLOCEIDA.

## Subfamily VIDUINÆ.

- Cryptospiza reichenowi: & described; A. REICHENOW, J. f. O. 1892, p. 187. C. salvadorii, n. sp., Shoa; id. Ber. Allg. Deutsch. Ges., Feb. 1892, p. 6.
- Hypochera: notes on the species; id. J. f. O. 1892, p. 49.
- Munia sinensis: notes on; J. BÜTTIKOFER, Notes Leyd. Mus. xiv, p. 132.

  M. nisoria and its allies; id. t. c. pp. 202-204.
- Estrelda roseicrissa, n. sp., Bukoba, Victoria Nyanza; A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., Feb. 1892, p. 7; id. J. f. O. 1892, p. 47.

#### Subfamily PLOCEINÆ,

- Symplectes croconotus and S. tephronotus, n. spp., Cameroons; A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., Feb. 1892, p. 4; id. J. f. O. 1892, p. 186. S. melanogaster, & described; id. J. f. O. 1892, p. 185. S. preussi, n. sp., Cameroons; id. t. c., Sept. 1892, p. 6.
- Malimbus: notes on species in Liberia; J. BÜTTIKOFER, Notes Leyd.

  Mus. xiv, pp. 24-28. M. rachiliæ: female described; Y. SJÖSTEDT,
  J. f. O. 1892, p. 313.

## Family TANAGRIDE.

- Euphonia: notes on the Costa Rican species, with 'Key'; G. K. CHERRIE, Auk, ix, pp. 23-25.
- Rhamphocelus costaricensis, n. sp., Costa Rica; id. An. Mus. Costa Rica, iii, p. 135. R. atrosericeus capitalis, \*n. subsp., Venezuela; J. A. Allen, Bull. Am. Mus. iv, Art. v, p. 51.
- Saltator immaculatus, n. sp., Lima, Peru; H. von Berlepsch & J. Stolz-Mann, P. Z. S. 1892, p. 375.
- Pyranga testacea techudii, n. subsp., Lima, Peru; iid. t. c. p. 375.

## Family CEREBIDE.

Cæreba bartolemica and allies: notes on; C. B. Cory, Cat. W. Ind. Birds, pp. 154 & 155. C. uropiggialis, n. sp., Curação; H. von Berlepsch, J. f. O. 1892, pp. 77-81.

## Family DREPANIDE.

- Hemignathus obscurus, H. olivaceus, H. procerus, H. hanapepe, figured; S. B. Wilson & A. H. Evans, Av. Haw. part iii.
- Vestiaria coccinea: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxiii, fig. 2.
- Himatione fraithii, n. sp., Laysan I., Sandwich Group, W. ROTHSCHILD, Ann. & Mag. N. H. (6) x, p. 109. H. parva, H. montana, and H. stejnegeri, figured; S. B. WILSON & A. H. EVANS, Av. Haw., part iii. H. sanguinea: skeleton figured; A. B. MEYER, Abbild. Vögel, Skel., Taf. clxiii, fig. 1.

Loxioides baillieni: akeleton figured; id. op. cit. Taf. clxv, fig. 2. Psittacirostra psittacea: akeleton figured; id. t, c. fig. 1.

# Family FRINGILLIDA.

## Subfamily Coccothraustina.

- Coccothraustes vespertina, in Ontario; Trans. Can. Inst. iii, pp. 111-124: notes on; A. W. Butler, Auk, ix, pp. 238-274.
- Pyrrhulagra for Loxigilla: L. STEJNEGER, in Cory, Cat. W. Ind. Birds, p. 148. P. noctis grenadensis, n. subsp., Grenada, and P. n. ridgwayi, Antigua; C. B. Cory, t. c. p. 150.

### Subfamily FRINGILLINÆ.

Acanthis, Bechst., = Carduelis; P. L. SCLATER, Ibis, 1892, pp. 555-557.
Serinus serinus, in Devonshire; W. E. H. PIDSLEY, Zool. 1892, p. 114.
Rhodacanthis, n. g. Type, R. palmeri, Kona, Hawaii; W. ROTHSCHILD,
Ann. & Mag. N. H. (6) x, pp. 110 & 111. R. flaviceps, n. sp., Kona,

Hawaii, id, t. c. p. 111.

- Telespysa flavissima, n. sp., Laysan I., Sandwich Group, id. t. c. p. 110. Carpodacus crythrinus, breeding in East Prussia; E. Hartert, Ibis, 1892, p. 368.
- Uragus henricii, n. sp., Tibet; E. OUSTALET, Ann. Sci. Nat. (7) xii, p. 291, pl. xi, fig. 2.
- Eurhinospisa, n. g. Type, E. henrici, n. sp.; E. Oustalet, Ann. Sci. Nat. (7) xii, p. 293, pl. xi.
- Hyphantospiza, n. g. Type, H. olivacea (Fraser); A. REICHENOW, Ber. Allg. Dentsch. Orn. Ges. zi, p. 6.
- Loxia curvirostra, in Ireland; R. J. USSHER, Irish Nat. i, pp. 6-9 & 28-31. L. bifasciata, in Belgium; A. DE SELYS-LONGCHAMPS, C.R. ii, Congr. Orn. Intern. ii, p. 177.
- Pinicola enucleator, in Heligoland; H. GATKE, Orn. Jahrb. iii, p. 202.

### Subfamily EMBERIZINAL,

- Emberisa cirlus: its distribution in Great Britain; O. V. Aplin, Zool. 1892, pp. 121-128 & 174-181. E. hortulana, in E. Bohemia; K. Kněžowbek, Orn. Jahrb. iii, pp. 64-66. E. cia, near Graz; A. v. Mojsisovics, t. c. p. 79. E. pusilla, from Borneo; R. B. Sharpe, Bull. B. O. C. ii, p. iv.
- Zonotrichia albicollis, in California; R. RIDGWAY, Auk, ix, p. 302.

## Family ALAUDIDE.

- Alauda gulgula and allies: remarks on; W. E. BROOKS, Ibis, 1892, pp. 61 & 62. A. brachydactyla, near Vichy, France; Rev. Sci. Bourb. v, p. 53. A. yeltoniensis, in Heligoland; H. Gätke, Orn. Jahrb. iii, p. 202.
- Alamon margarita: eggs described; S. Alessi, J. f. O. 1892, pp. 314 & 315.
- Otocorys berlepschi, figured; E. HARTERT, Ibis, 1892, pl. xiii.

# Family MOTACILLIDE.

- Motacilla raii, on the migration of; J. CORDEAUX, Zool. 1892, pp. 389-391.
- Anthus blakistoni, remarks on ; W. E. BROOKS, Ibis, 1892, p. 61. A. gustavi: akeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxxi, fig. 1.

# Family MNIOTILTIDE.

- Sciurus noveboracensis notabilis, in New Jersey; W. C. Southwick, Auk, ix, p. 303.
- Dendræcu petechia flaviceps, n. subsp., Bahama Is., M. CHAPMAN, Bull. Am. Mus. iv, pp. 310 & 311. D. nigrifrons, & Q figured; W. Brewster, Auk, ix, pl. i. D. dugesi, figured; A. Duges, Nat.

Mex. (2) ii, pp. 97 & 98, pl. v, fig. 1. D. pennsylvanica nesting in Missouri; B. F. GAULT, Auk, ix, p. 396.

Helminthophila pinus × H. leucobronchialis; F. M. CHAPMAN, t. c. p. 302. H. leucobronchialis, in Queen's Co., Long I.; A. H. HOWELL, t. c. p. 306.

## Family CERTHIDE.

Certhia familiaris, figured; E. H. GIGLIOLI & A. MANZELLA, Iconogr. Avif. Ital. pt. l. C. mexicana from Nicaragua; O. Salvin & F. D. Godman, Ibis, 1892, p. 325.

## Family SITTIDE.

Sitta casia homeyeri, in East Prussia; E. HARTERT, Ibis, 1892, pp. 364 & 365. S. neumayeri, figured; E. H. GIGLIOLI & A. MANZELLA, Iconogr. Avif. Ital. pt. xl.

## Family MELIPHAGIDE.

- Viridonia, n. g. Type, V. sagittirostris, n. sp., Sandwich Is.; W. ROTHSCHILD, Ann. N. H. (6) x, p. 112.
- Melipotes maculata, n. sp., Mt. Suckling, S. E. New Guinea; C. W. DE VIS, Ann. Rep. Brit. New Guinea, 1890-91, App. cc, p. 94; id. Ann. Queensl. Mus. No. 2, p. 5.
- Ptilotis frenata: nest and eggs described and figured; A. J. North, Rec. Austral. Mus. ii, p. 15, pl. iii.
- Stigmatops ocularis: remarks on ; J. BÜTTIKOFER, Notes Leyd. Mus. xiv, pp. 199 & 200.
- Acrulocercus braccatus: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxiv, fig. 1.

## Family NECTARINIDE.

- Nectarinia gadowi, n. sp., Benguela; J. V. BARBOZA DU BOCAGE, Jorn. Lisb. (2) No. viii, p. 256. N. osea, near Beyrout; W. T. VAN DYCH, Ibis, 1892, p. 469.
- Cinnyris nesophilus, n. sp., Anjuan I., G. E. SHELLEY, Bull. B. O. C., No. 2, p. iv. C. oritis and C. preussi, n. spp., Cameroons, A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., March, 1892, p. 4: id. J. f. O. 1892, p. 190.

# Family DICEIDE.

Diccum wilhelmina, n. sp., Sumba I., J. BÜTTIKOFER, Notes Leyd. Mus. xiv, p. 199.

## Family ZOSTEBOPIDE.

Zosterops incerta, n. sp., I. of Engano, Sumatra, T. Salvadori, Ann. Mus. Genov. (2) xii, p. 133. Z. squamifrons, n. sp., Mt. Dulit, N. W. Borneo, R. B. Sharpe, Ibis, 1892, p. 323. Z. anderssoni, n. sp., Damaraland, G. E. Shelley, Bull. B. O. C., No. 2, p. v. Z. stuhlmanni, n. sp., Victoria Nyansa, A. Reichenow, Ber. Allg. Deutsch. Orn. Ges., March, 1892, p. 7: id. J. f. O. 1892, p. 54. Z. stenocricota, n. sp., Cameroons, id. Ber. Allg. Deutsch. Ges., March, 1892, p. 7: id. J. f. O. 1892, p. 191. Z. flavilateralis, n. sp., East Africa, id. t. c. p. 192. Z. superciliosa, n. sp., Equatorial Africa, id. ibid. Z. atrifrons: skeleton figured; A. B. Meyer, Abbild. Vögel. Skel., Taf. clxxiii, fig. 1. Z. mussoti, n. sp., Szechuen, E. Oustalet, Ann. Sci. Nat. (7) xii, p. 289.

## Family REGULIDE.

Regulus: notes on the genus, with key to species; T. PLESKE, Mél. biol., Taf. xiii, livr. 2, pp. 303-337. B. tristis, n. sp., id. t. c. p. 306, Turkestan. R. himalayensis, remarks on; W. E. Brooks, Ibis, 1892, p. 59.

## Family PARIDE.

Parus cristatus, in West of France; L. Bureau, Bull. Soc. Sci. Nat. Ouest France, ii, p. 143. P. major, double nest; E. Lovett, P. Tr. Croydon Nat. Hist. Club, 1892, pp. 330 & 331. P. pleskei, in Belgium; E. DE SÉLYS-LONGCHAMP, C.R. ii, Congr. Orn. Intern. ii, p. 178. Notes on synomymy of European species; v. Tschusi zu Schmidhoffen, Orn. Jahrb. iii, pp. 175-178. P. xanthostomus, n. sp., R. Zambesi, G. E. Shelley, Bull. B. O. C., No. 2, p. vi. P. rovumæ, n. sp., Rovuma River, id. ibid.

Leptopocile henrici, n. sp., Tibet; E. OUSTALET, Ann. Sci. Nat. (7) xii, p. 287, pl. x, fig. 1.

Acredula irbii, in South Tirol; V. RITTER VON TSCHUSI ZUR SCHMID-HOFFEN, Orn. Jahrb. iii, p. 122. A. macedonica, n. sp., Macedonia, T. Salvadori & H. E. Dresser, Bull. B. O. C. No. 4, p. xvi. A. bonvaloti, n. sp., Tà-tsién-lou; E. Oustalet, Ann. Sci. Nat. (7) xii, pp. 284-286, pl. ix, fig. 1.

# Family LANIIDE.

- APLIN, O. V. Further remarks upon Grey Shrikes, Zool. 1892, p. 112.
  DRESSER, N. E. Remarks on Lanius lahlorn and its allies. Ibis, 1892, pp. 288-293.
- —. Notes on the Palsearctic species of the genus Lanius, t. c. pp. 374-380.

Lanius excubitor, in Gloucestershire; A. L. CLARKE, Zool. 1892, p. 151.

L. mackinnoni [Zool. Rec. xxviii, Aves, p. 62], in Cameroons;
A. Reichenow, J. f. O. 1892, p. 184. L. minor, figured; E. H. Giglioli & A. Manzella, Icongr. Avif. Ital. part xlix. L. rufus, in Great Britain; O. V. Aplin, Zool. 1892, pp. 345-352; in Sussex;
T. Parkin, t. c. p. 230: in South Devon; E. A. Elliot, t. c. p. 412.

Laniarius gubernator, n. sp., Cameroons, A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., Sept. 1892, p. 5.

Nilaus nigritemporalis, n. sp., Ngoma, Equatorial Africa, id. t. c. p. 3: id. J. f. O. 1892, pp. 36 & 37.

Hypocolius ampelinus, & Q figured; P. L. Sclater, P. Z. S. p. 470.

Eopsaltria sudestiensis, n. sp., Sud-est I., Louisiade Archipelago, C. W. De Vis, Ann. Rep. Brit. New Guinea, 1890-91, p. 96: Ann. Queensl. Mus. No. 2, p. 8.

## Family AMPELIDE.

Ampelis macsi, n. sp., Japan, E. Oustalet, N. Arch. Mus. Paris (3) iv, pp. 213-216.

## Family VIREONIDE.

Vireo: notes on West Indian species; C. B. Cory, Cat. W. Ind. Birds, pp. 152-154. V. olivaceus, in British Colombia and Washington; C. F. BATCHELDER, Auk, ix, pp. 395 & 396.

# Family TURDIDE.

- Geocichla everetti, n. sp., Mt. Dulit, Sarawak, Borneo, R. B. Sharpe, Ibis, 1892, pp. 323 & 324. G. leucolæma, n. sp., Isle of Engano, T. Salvadori, Ann. Mus. Genov. (2) xii, pp. 135 & 136.
- Turdus nigrilorum, n. sp., Cameroons, A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., March, 1892, p. 4: id. J. f. O. 1892, p. 194.
- Merula celebensis, n. sp., Macassar, Celebes, J. BÜTTIKOFER, Notes Leyd.

  Mus. xv, p. 109. Notes on M. javanica, M. seebohmi and M. schlegeli,
  id. t. c. pp. 107-110. M. elgonensis [Zool. Rec. xxviii, Aves, p. 63],
  fully described; R. B. Sharpe, Ibis, 1892, pp. 160 & 161.
- Phæornis obscura: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxiv, fig. 2.
- Callene hypoleuca, n. sp., Cameroons, A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., Feb. 1892, p. 6: id. J. f. O. 1892, p. 194, tab. ii, fig. 3.

Cossypha polioptera, n. sp., Victoria Nyanza, id. t. c. p. 133.

- Erythacus sibilans, from Hainan; G. HARTLAUB, Abh. Ver. Brem. xii, pp. 304 & 305. E. rubecula, figured; LORD LILFORD, Col. Fig. Br. Birds, part xxii.
- Cyanecula succica, in Orkney; T. E. BUCKLEY, Ann. Scott. N. H. 1892, p. 70.

- Ruticilla tithys, in Lancashire; F. Brownswood, Zool. 1892, p. 115. R. moussieri, figured; E. H. GIGLIOLI & A. MANZELLA, Icongr. Avif. Ital. part xlviii.
- Pratincola pallidigula, n. sp., Cameroons, A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., April, 1892, p. 4: id. J. f. O. 1892, p. 194. P. indica, remarks on; W. E. BROOKS, Ibis, 1892, p. 60. P. rubicola, in N. E. Germany; H. SCHALOW, SB. nat. Fr. 1892, pp. 141-145. P. maura presualskii; TH. PLESKE, Bull. Pétersb. (3) xxxv, p. 129.
- Cercomela (Myrmecocichla) melanura, note on the habits; H. E. BARNES, J. Bomb. N. H. Soc. vii, pp. 252 & 253.
- Thamsolas nigra and its allies; J. BÜTTIKOFER, Notes Leyd. Mus. xiv, pp. 17 & 18. T. shelleyi, notes on, J. V. BARBOZA DU BOCAGE, J. Sci. Lisb. (2) vii, pp. 167-170.
- Saxicola leucomela, figured; E. H. GIGLIOLI & A. MANZELLA, Iconogr. Avif. Ital. p. xlix.

## Family Sylvinde.

- Sylvia misoria in Yorkshire; J. CORDEAUX, Zool. 1892, p. 424; id. P. Z. S. 1892, p. 542: records of capture in Great Britain; H. SEEBOHM, Bull. B. O. C., No. 3, p. xi.
- Hypolais caligata and H. roma: remarks on; W. E. Brooks, Ibis, 1892, pp. 59 & 60. H. rama and Acrocephalus streperus: hybrid described; Th. Pleske, Orn. Ross. pp. 561-563.
- Phylloscopus neglectus and P. sindianus: remarks on; W. E. BROOKS, Ibis, 1892, p. 60. P. rufus, wintering in Somerset; R. H. READ, Zool. 1892, p. 111: unusual nesting; id. ibid.: notes on nesting; J. BACKHOUSE, t. c. p. 150; C. F. ARCHIBALD, t. c. p. 226. P. superciliosus, in Lincolnshire; G. H. CATON HAIGH, Zool. 1892, p. 413.
- Acanthopneuste ijima, n. sp., Seven Is., Japan; L. Stejneger, P. U. S. Nat. Mus. xv, p. 372.
- Lusciniola melanopogon, in Hungary; A. Grunack, J. f. O. 1892, pp. 213 & 214.
- Locustella navia, nesting in S. W. Scotland; R. SERVICE, Zool. 1892, p. 333. L. ochotensis, ad. et juv., figured; Th. Pleske, Orn. Ross., tab. 4, figs. 1 & 2.
- Cettia canturiens, & ad. figured; id. t. c., tab. iv, fig. 3. C. minuta; id. t. c., tab. iv, fig. 4. C. canturiens septentrionalis, n. subsp., Russian Manchuria; C.W. Campbell, Ibis, 1892, p. 235. O. canturiens borealis, n. subsp., Corea; id. t. c. p. 235.
- Urosphena squamiceps, ad. figured; T. PLESKE, Orn. Ross., tab. iv, fig. 5.
  Acrocephalus fruticola (Naum.), in E. Prussia; E. HARTERT, Ibis, 1892, pp. 361 & 362. A. palustris, in Oxfordshire; W. WARDE FOWLER, Zool. 1892, pp. 303-309.
- Tatars: notes on the species; J. BÜTTIKOFER, Notes Leyd. Mus. xiv, pp. 13-16. T. familiaris, n. sp., Laysan I., Sandwich Group; W. ROTHSCHILD, Ann. N. H. (6) x, p. 109.

- Burnesia epichlora, n. sp., Camaroons; A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., Feb. 1892, p. 6; figured, id. J. f. O. 1892, p. 193, tab. ii, fig. 1.
- Drymaca sahara, figured; A. Koenig, J. f. O. 1892, tab. iii.
- Apalis mystacalis [Zool. Rec. xxviii, Aves, p. 65], figured; A. REICHENOW, t. c. tab. i, fig. 2. A. jacksoni [Zool. Rec. xxviii, Aves, p. 65], figured; R. B. SHARPE, Ibis, 1892, pl. iv, fig. 2. A. pulchra [Zool. Rec. xxviii, Aves, p. 65], figured; id. t. c. pl. iv, fig. 1.
- Camaroptera griseigula, n. sp., Teita, E. Africa; id. t. c. p. 158.
- Malurus mortoni, n. sp., Sudest I., Louisiade Archipelago; C. W. DE VIS, Ann. Rep. Brit. New Guinea, 1890-91, p. 97; id. Ann. Queensland Mus. No. 2, p. 9.
- Acanthiza tenkatei, n. sp., Flores; J. Büttikofer, Notes Leyd. Mus. xiv, p. 195.
- Cisticola emini, n. sp., Victoria Nyanza; A. REICHENOW, J. f. O. 1892, p. 56. C. chubbi, n. sp., Mt. Elgon, E. C. Africa; R. B. SHARPE, Ibis, 1892, pp. 157 & 158.
- Graminicola striata, n. sp., Hainan; F. W. STYAN, Bull. B. O. C., No. 2, p. vi.

## Family TrogLodyTIDE.

Elachura haplonota, n. sp. [E. immaculata on plate], from N. Cachar, Assam; E. C. S. BAKER, J. Bomb. N. H. Soc. vii, pp. 319-322; Ibis, 1892, pp. 62-64, pl. i, fig. 1. E. punctata, figured; id. t. c. pl. i, fig. 2.

# Family CINCLIDE.

Cinclus: notes on some Palmarctic species; H. E. DRESSER, Ibis, 1892, pp. 380-387. C. melanogaster pyrenaicus, range of; O. V. APLIN, Zool. 1892, p. 359. C. pallasi, var. soullisi, n. subsp., Tibet; E. Oustalet, Ann. Sci. Nat. (7) xii, p. 299.

# Family Minida.

- Minus polyglottus, formerly in New Jersey; S. LOCKWOOD, Am. Nat. xxvi, pp. 635-641.
- Harporhynchus curvirostris palmeri: its nesting habits; H. Brown, Zoe, iii, pp. 243-248.

# Family TIMELIIDE.

- Bubax lanceolatus, var. bonvaloti, n. subsp., So, Tibet; E. Oustalet, Ann. Sci. Nat. (7) xii, p. 273.
- Pomatorhinus maclellandi, var. dedekensi, n. subsp., Tioungeu; P. m., var. armandi, n. subsp., Aio, Tibet; E. Oustalet, t. c. pp. 276 & 277.

- Trochalopteron henricii, n. sp., So, Tibet; E. Oustalet, Ann. Sci. Nat. (7) xii, p. 274. T. bonvuloti, n. sp., Tioungeu, id. t. c. p. 275.
- Suthora fulvicauda and S. longicauda, n. spp., Corea; C. W. CAMPBELL, Ibis, 1892, p. 237.
- Amalocichla, n. g. Type, A. sclateriana, n. sp., Mt. Suckling, S. E. New Guinea; C. W. De Vis, Ann. Rep. Brit. New Guinea, App. cc, 1890-91, p. 95; id. Ann. Queensland Mus., No. 2, pp. 5 & 6.
- Orthonyx spinicauda: nest described and egg figured; A. J. NORTH, Rec. Austral. Mus. ii, pp. 15 & 16, pl. ii, fig. 5.
- Turdinus monachus, n. sp., Cameroons; A. REICHENOW, Ber. Allg.
   Deutsch. Orn. Ges., Feb. 1892, p. 4; id. J. f. O. 1892, p. 193. T.
   moloneyanus, n. sp., Gold Coast; R. B. SHARPE, P. Z. S. 1892, p. 228, pl. xx, fig. 2.
- Amaurocichla, n. g. Type, A. bocagii, n. sp., St. Thomas, West Africa; id. t. c. p. 228, pl. xx, fig. 1.
- Ptilocichla leucogastra, n. sp., Pahang, E. Malay Peninsula, W. R. DAVIson, Ibis, 1892, pp. 100 & 101.
- Malacopteron melanocephalum, n. sp., Pahang, E. Malay Peninsula, id. t. c. pp. 101 & 102.
- Eupetes geislerorum, n. sp., Butaueng, N. E. New Guinea; A. B. MEYER, J. f. O. 1892, pp. 259 & 260.
- Stachyris davisoni, n. sp., Pahang, E. Malay Peninsula, R. B. Sharpe, Bull. B. O. C., No. 2, p. vii.
- Alcippe bieti, n. sp., E. Oustalet, Ta-tsién-lou, Ann. Sci. Nat. (7) xii, p. 304.

# Family Pycnonotidæ.

- Hypsipetes psaroides and H. concolor: account of nesting habits; E. C. S. Baker, J. Bomb. N. H. Soc. vii, pp. 129 & 265.
- Hemixus flavala: account of nesting habits; id. t. c. p. 7.
- Criniger aureus, note on; A. B. MEYER, Ibis, 1892, pp. 178 & 180. C. flaveolus, figured with nest; E. C. S. BAKER, J. Bomb. N. H. Soc. vii, pl. ii.
- Otocompsa emeria and O. flaviventris, figured with nests; id. t. c. pls. viii & xv.
- Xanthixus flavescens, figured with nest; id. t. c. pp. 1-4, pl. i.
- Pinarocichla schmackeri, n. sp., Hainan; F. W. STYAN, Bull. B. O. C., No. 2, p. vi [= Criniger pallidus, Swinhoe; R. B. SHARPE, op. cit. No. 4, p. xix].
- Xenocichla poliocephala, n. sp., Cameroons; A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., Feb. 1892, p. 5; id. J. f. O. 1892, p. 189.
- Spisizus canifrons: account of nesting habits; E. C. S. BAKER, J. Bomb. N. H. Soc, vii, p. 10.
- Andropadus montanus, n. sp., Cameroons; A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., Feb. 1892, p. 5; id. J. f. O. 1892, p. 188. A. cameronensis, n. sp., Cameroons, id. J. f. O. 1892, pp. 126 & 127. A. sugenius, n. sp., Victoria Nyanza, id. t. c. p. 133.

# Family CAMPOPHAGIDE.

- Graucalus preussi, n. sp., Cameroons, A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges. xi, p. 4. G. enganensis, n. sp., I. of Engano, T. Salvadori, Ann. Mus. Genov. (2) xii, pp. 129 & 130. G. nigrifrons, n. sp., Bugotu, Solomon Is., H. B. Tristram, Ibis, 1892, p. 294. G. welchmani, n. sp., Bugotu, id. t. c. pp. 294 & 295. G. leucopygius: skeleton figured; A. B. Meyer, Abbild. Vögel. Skel., Taf. clxvi, fig. 2.
- Edoliisoma tenuirostre: nest and eggs described; A. J. NORTH, Rec. Austral. Mus. ii, pp. 13 & 14. E. morio: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxvii, fig. 2.
- Campophaga minor, n. sp., Pahang, E. Malay Peninsula, W. R. DAVISON, Ibis, 1892, p. 99 [= Lalage culminata; R. B. SHARPE, Bull. B. O. C., No. 2, p. vii].
- Pericrocotus modiglianii, n. sp., I. of Engano, T. SALVADORI, Ann. Mus. Genov. (2) xii, pp. 130-132. P. wrayi = P. montanus: P. cinereigula also probably identical; R. B. SHARPE, Ibis, 1892, p. 435.
- Lalage leucopygialis: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxvii, fig. 1.

# Family MUSCICAPIDE.

- Alseonax pumila, n. sp., Bukoba, Victoria Nyanza, A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., Jan. 1892, xi, p. 3; id. J. f. O. 1892, pp. 32 & 33.
- Platystira jacksoni [Zool. Rec. xxviii, Aves, p. 67], fully described and figured; R. B. Sharpe, Ibis, 1892, p. 301, pl. vii, fig. 2.
- Musicapa grisola: supposed occurrence in N. Zealand; F. W. HUTTON, Tr. N. Z. Inst. xxiv, p. 710 [Determination erroneous; teste H. O. Forbes]. M. collaris, figured; E. H. Giglioli & A. Manzella, Iconogr. Avif. Ital. pt. xlix.
- Erythrosterna parva, figured; iid. t. c. pt. xlix.
- Muscicapa johnstoni = Dioptrornis fischeri; R. B. SHARPE, Ibis, 1892, pp. 300 & 301.
- Rubecula tytleri, Jameson = Muscicapa parva; W. E. CLARKE, Ibis, 1892, pp. 558 & 559.
- Pedilorhynchus stuhlmanni [Zool. Rec. xxviii, Aves, p. 67], figured; A. REICHENOW, J. f. O. 1892, tab. i, fig. 1: in Cameroons; id. t. c. p. 183.
- Gerygone insperata, n. sp., Mt. Suckling, S. E. New Guinea, C. W. De Vis, Ann. Rep. Brit. New Guinea, 1890-91, App. cc, p. 94; id. Ann. Queensland Mus. No. 2, p. 4. G. pectoralis, n. sp., Pahang, E. Malay Peninsula; W. R. Davison, Ibis, 1892, pp. 99 & 100 [= G. modiglianii, Salvad. (Zool. Rec. xxviii, Aves, p. 67); R. B. Sharpe, Bull. B. O. C., No. 2, p. vii.
- Pacilodryas vicaria, n. sp., Mt. Suckling, S. E. New Guinea, C. W. DE VIS, Ann. Rep. Brit. New Guinea, 1890-91, App. cc, p. 94: Ann. Queensland Mus. No. 2, p. 4.

- Hyliota nehrkorni, n. sp., Accra, G. HARTLAUB, Ibis, 1892, p. 373, pl. viii. Review of the species of Hyliota; id. pp. 373 & 374.
- Parisoma lugens, fully described; R. B. Sharpe, Ibis, 1892, pp. 302 & 303.
- Trochocercus albonotatus [Zool. Rec. xxviii, Aves, p. 67], figured; R. B. Sharpe, Ibis, 1892, pl. vii, fig. 1.
- Rhipidura: review of the genus; J. BÜTTIKOFER, Notes Leyd. Mus. xv, pp. 65-98 & 113-115. R. celebensis, n. sp., Macassar, p. 79. R. teyesmanni, n. sp., Macassar, p. 80. R. meyeri, n. sp. (R. braehyrhyncha, auct., nec Schl.), pp. 82 & 113. R. sumbawensis, n. sp., Sumbawa, p. 85. R. rosenbergi, n. sp., Aru I., p. 88. R. hoeedti, n. sp., Lettie I., p. 93. R. tenkatei, n. sp., Rotti I.; id. Notes Leyd. Mus. xiv, p. 205. R. concinna, n. sp., Mt. Suckling, S. E. New Guinea; C. W. DE Vis, Ann. Rep. Brit. New Guinea, 1890-91, App. cc, p. 94. R. büttikoferi, Dammar I., Banda Sea; R. B. Sharpe, Bull. B. O. C., No. 4, p. xviii.
- Elminia echwebischi, n. sp., Congo; E. Oustalet, N. Arch. Mus. Paris (3) iv, pp. 216 & 217.
- Cryptolopha bicolor, n. sp., Hainan; F. W. STYAN, Bull. B. O. C., No. 2, p. vi. [= Herpornis tyrannulus, Swinh.; R. B. SHARPE, Bull. B. O. C., No. 4, p. xix.] C. mackensiana, n. sp., Kikuyu, E. C. Africa; R. B. SHARPE, Ibis, 1892, p. 153. C. xanthoschistus: remarks on; W. E. BROOKS, Ibis, 1892, p. 60.
- Siphia luteola: female described; C. W. CAMPBELL, Ibis, 1892, pp. 233 & 234.

# Family HIRUNDINIDE.

- Duchassov, H. De l'arrivée et du départ des Hirondelles. Bull. Soc. L. Nord France, i, pp. 85-92 (1891).
- SHARPE, R. B., & WYATT, C. W. A Monograph of the *Hirundinida*, or Family of Swallows. Part xv. 1892, 4to.
- The following species are described and figured:—Atticora cinerea; Hirundo javanica, H. tahitica, H. namiyei (no figure), H. arcticincta; Cotile rupestris; Tachycineta thalassinus.
- Hirundo emini, n. sp., Victoria Nyanza; A. REICHENOW, Ber. Allg. Deutsch. Orn. Ges., Jan. 1892, p. 2; id. J. f. O. 1892, pp. 30 & 31.
- Psalidoprocne chalybea, n. sp., Cameroons; id. op. cit., Sept. 1892, p. 6.

#### Section B. OLIGOMYODÆ.

## Family TYRANNIDE.

- ALLEN, J. A. On a Collection of Birds from Chapada, Matto Grosso, Brazil, made by Mrs. H. H. Smith [Zool. Rec. xxviii, Aves, p. 2]. Part·II. Tyrannidæ. Bull. Am. Mus. iv, pp. 331-350.
- Lophotriccus subcristatus, n. sp., Venezuela; id. t. c. p. 53.

- Anæretes sclateri, n. sp., Chili; E. Oustalet, N. Arch. Mus. Paris (3) iv, pp. 217 & 218.
- Mnionectes semischistaceus, n. sp., Costa Rica; G. K. CHERRIE, P. U. S. Nat. Mus. xv, p. 27.
- Ornithion pusillum subflavum, n. subsp., Costa Rica; id. t. c. p. 28.
- Pitangus jamaicensis, n. sp., Jamaica; F. M. Chapman, Bull. Am. Mus. iv, p. 303.
- Pyrocephalus rubineus heterurus, n. subsp., Lima, Peru; H. von Berlepsch & J. Stolzmann, P. Z. S. 1892, p. 381.
- Myiarchus nuttingi, in Arizona; A. K. FISHER, Auk, ix, p. 394. M. brevipennis, n. sp., I. of Aruba, E. HARTERT, Bull. B. O. C., No. 3, p. xii.

## Family Cotingida.

Chasmorhynchus niveus: position of the Caruncle and method of song; J. J. QUELCH, J. R. Agr. & Comm. Soc. Brit. Guiana (2) vi, pp. 164-166. C. tricarunculatus: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clviii.

# Family PITTIDE.

- Pitta granatina borneensis, n. subsp., Borneo, D. G. Elliot, Auk, ix, pp. 218-221.
- Erythropitta celebensis: skeleton figured; A. B. MEYER, Abbild. Vögel. Skel., Taf. clxi, fig. 1.

## Division II. TRACHEOPHONÆ.

# Family DENDROCOLAPTIDE.

### Subfamily FURNARIINA.

- Upucerthia harterti, n. sp., Bolivia, H. von Berlepsch, Ber. Allg. Deutsch. Orn. Ges., Dec. 1892, p. 10.
- Cinclodes taczanowskii, n. sp., Chorillos, Peru, id. & J. Stolzmann, P. Z. S. 1892, p. 381.

# Family FORMICABIIDE.

- Thamnophilus punctatus and T. bridgesi, figured; O. Salvin & F. D. Godman, Biologia Centrali-americana, Aves, ii, pl. xlix, figs. 1 & 2. T. atrinucha, n. sp., Honduras to Ecuador, iid. t. c. pp. 200 & 201. T. immaculatus, Lafr., is a Myrmelastes; iid. t. c. p. 225.
- Myrmelastes lawrencii, n. sp. [= M. corvinus, Lawr., nec Gould], Panama, iid. t. c. pp. 226 & 227. M. intermedius, figured [n. s. Myrmeciza immaculata], iid. t. c. pl. li, figs. 2 & 3.
- Thamnestes anabatinus, & figured; iid. t. c. pl. l, fig. 1.

- Dysithamnus puncticeps, & Q figured; iid. t. c. pl. 1, figs. 2 & 3.
- Formicivora stictocorypha, n. sp., Porto Real, Brazil, A. BOUCARD & H. VON BERLEPSCH, Humming Bird, ii, p. 44.
- Myrmeciza leucosticta, figured; iid. t. c. pl. li, fig. 1. M. nigricauda, n. sp., Ecuador, iid. t. c. p. 230. M. occidentalis, n. sp., Costa Rica, G. K. CHERRIE, An. Mus. Costa Rica, iii, p. 135.
- Pithys leucaspis is a Gymnopithys; O. Salvin & F. D. Godman, t. c. p. 221.

  G. rusicops, n. sp., Colombia, iid. t. c. p. 222, note.
- Grallaria princeps (pl. lii), G. dives (pl. liii, fig. 1), and G. perspicillata (pl. liii, fig. 2), figured; iid. t. c. G. lizanoi, n. sp., Costa Rica, G. K. Cherrie, An. Mus. Costa Rica, iii, p. 137.



# IV. REPTILIA AND BATRACHIA.

#### BY

#### G. A. BOULENGER.

This subject is treated as follows, viz:-

- I. TITLES OF ALL WORKS AND PAPERS ARRANGED ALPHABETICALLY ACCORDING TO AUTHORS' NAMES.
- II. Works and Papers dealing with both Reptiles and Batrachians.
- III. FAUNISTIC: A, RECENT; B, FOSSIL.
- IV. REFERENCES, ARRANGED SYSTEMATICALLY.

#### I. TITLES.\*

- Adolphi, H. Ueber Variationen der Spinalnerven und der Wirbelsäule unserer Amphibien. 1. Bufo variabilis, Pall. Morph. JB. xix, 1892, pp. 313-375, figs., pl. xii.
- Ambrosetti, J. B. Observaciones sobre los Reptiles fósiles oligocenos de los terrenos terciarios antiguos del Paraná. Bol. Ac. Cordoba, x. 1890, pp. 409-426.
- ANDERSON, J. On a small Collection of Mammals, Reptiles, and Batrachians from Barbary. P. Z. S. 1892, pp. 3-24, pl. i.
- Andersson, O. A. Zur Kenntniss des sympathischen Nervensystems der urodelen Amphibien. Morph. JB. (Anat.), v, 1892, pp. 184-210, pls. xv-xviii.
- AYERS, H. Vertebrate Cephalogenesis. II. A Contribution to the Morphology of the Vertebrate Ear, with a Reconsideration of its Functions. J. Morph. vi, 1892, pp. 1-360, pls. i-xii.
- BATAILLON, E. Contribution à l'étude de la circulation artérielle des Chéloniens. Ann. Soc. L. Lyon (2) xxxix, 1892, pp. 81-86, pl.

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

<sup>1892. [</sup>vol. xxix.]

- BAUR, G. Das Variiren der Eidechsen-Gattung Tropidurus auf den Galapagos-Inseln. Leuckart Festschrift, 1892, pp. 259-277, figs.
- —... On the Morphology of the Skull in the Mosasaurida. J. Morph. vii, 1892, pp. 1-22, pls. i & ii.
- ——. Bemerkungen über verschiedene Arten von Schildkröten. Zool. Anz. xv, 1892, pp. 155-159.
- —. Der Carpus der Schildkröten. Erwiderung an Herrn Prof. Dr. Emil Rosenberg. Anat. Anz. vii, 1892, pp. 206-211, figs.
- —... On the Taxonomy of the genus *Emys*, C. Duméril. P. Am. Phil. Soc. xxx, 1892, pp. 40-44. Additional note, t. c. p. 245.
- Bedriaga, J. De. A propos de l'importation et du croisement des Reptiles et des Amphibiens. Lettre à M. le professeur Anatole Bogdanow. C.R. Congrès Internat. Zool. i, 1892, pp. 244 & 245.
- —. Chalcides simonyi, Steind., et Molge luschani, Steind. T. c. pp. 242 & 243.
- Tableaux synoptiques pour servir à la détermination des larves des Batraciens Urodèles d'Europe. C.R. Ass. Fr. Sci. 1891, ii, pp. 540-546.
- ——. Ueber die Begattung bei einigen geschwänzten Amphibien (Fortsetzung). Zool. Anz. xvi, 1893, pp. 102-104.
- —. Notice sur les Amphibiens et Reptiles recueillis par M. A. F. Moller aux îles de la Guinée. O Istituto (Coimbra), ser. 2, Nos. 6 & 7, 1892 [quoted from a Reprint].
- Benda, C. Ueber die Histiogenese des Sauropsidenspermatozoons. Verh. Anat. Ges. 1892, pp. 195-199.
- BÉRANEK, E. Sur le nerf pariétal et la morphologie du troisième œil des Vertébrés. Anat. Anz. vii, 1892, pp. 674-689, figs.
- Bertelli, D. Sulla membrana timpanica della Rana esculenta. Mon. Zool. Ital. iii, 1892, pp. 203-207. French translation: Arch. Ital. Biol. xviii, 1893, pp. 458-462.
- Bertoldo, J. M. Sur le Développement des Axolotis. Le Nat. xiv, 1892, p. 15.
- Biedermann, W. Ueber den Farbenwechsel der Frösche. Arch. ges. Phys. li, 1892, pp. 455-508, pl. xi.
- Bocage, J. V. B. Du. Subsidios para a Fauna da Guiné Portugueza. J. Sci. Lisb. (2) ii, 1892, pp. 179–184.
- ----. Sur le Hemidactylus mabouiu, var. molleri, Bedriaga, de St. Thomé. T. c. p. 221.
- —. Notice sur les Amphibiens et Reptiles recueillis par M. A. F. Moller aux Iles de la Guinée. T. c. pp. 229-232.

- [BOCAGE, J. V. B. DU.] Note sur le *Dendraspis* de l'île St. Thomé. J. Sci. Lisb. (2) ii, 1892, pp. 265-267, fig.
- BOCOURT, F. Note sur la variabilité dans le nombre des plaques céphaliques chez certains Ophidiens. Bull. Soc. Z. Fr. xvii, 1892, pp. 40 & 41.
- —. Note sur un petit Ophidien appartenant au genre Tropidoclonium. Le Nat. xiv, 1892, p. 132.
- —. Note sur un Ophidien appartenant au genre Eutenia. T. c. p. 278.
- Bœttger, O. Katalog der Batrachier-Sammlung im Museum der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt am Main. Frankfort a/M.: 1892, 8vo, 73 pp.
- —. Kriechtiere der Kaukasusländer, gesammelt durch die Radde-Valentin'sche Expedition nach dem Karabagh und durch die Herren Dr. J. Valentin und P. Reibisch. Ber. Senck. Ges. 1892, pp. 131-150.
- —. Mitteilung über eine Reptiliensendung von Aden in Süd-Arabien. Ber. Offenb. Ver. xxix-xxxii, 1892, pp. 61-63.
- —. Liste von Kriechtieren und Lurchen aus dem tropischen Asien u. aus Papuasien. T. c. pp. 65-157.
- —. Drei neue colubriforme Schlangen. Zool. Anz. xv, 1892, pp. 417–420.
- BORN, G. Die Reifung des Amphibieneies und die Befruchtung unreifer Eier bei Triton tæniatus. Anat. Anz. vii, 1892, pp. 772-781 & 803-811, figs.
- BOULENGER, G. A. Catalogue of the Snakes in the British Museum (Natural History). Vol. 1, containing the families Typhlopidæ, Glauconiidæ, Boidæ, Ilysiidæ, Uropeltidæ, Xenopeltidæ, and Colubridæ aglyphæ, part. London: 1893, 8vo, xiii & 448 pp., 28 pls.
- —... Reptilia and Batrachia. Appendix to Distant's "Naturalist in the Transvaal." London: 1892, 8vo, pp. 174-176, fig.
- —. Remarks on an Iguana with reproduced tail. P. Z. S. 1891, pp. 466 & 467.
- ——. A Synopsis of the Tadpoles of the European Batrachians. T. c. pp. 593-627, pls. xlv-xlvii.
- —. Notes on Transcaspian Reptiles. T. c. pp. 628-633.
- —... An Account of the Reptiles and Batrachians collected by Mr. C. Hose on Mt. Dulit, Borneo. Op. cit. 1892, pp. 505-508, pls. xxix & xxx.
- On newly-discovered East African Chameleons, with Remarks on some other Reptiles described by Dr. Steindachner. Ann. N. H. (6) ix, 1892, pp. 72-74.
- —. Description of a new Snake from Nubia. T. c. pp. 74-76.

- [BOULENGER, G. A.] Description of a new Frog from Burma. Anu. N. H. (6) ix, 1892, pp. 141 & 142, pl. ix.
- --- On Strauch's Triton longipes. T. c. pp. 144 & 145.
- —... Description of new Reptiles and Batrachians from the Loo Choo Islands. Op. cit. x, 1892, pp. 302-304.
- On the Larva of Molye montandoni. T. c. pp. 304 & 305, fig.
- ---. On some newly-described Jurassic and Cretaceous Lizards and Rhynchocephalians. Op. cit. xi, 1893, pp. 204-210.
- ---. Description of a new Toad from Travancore. J. Bomb. Soc. vii, 1892, pp. 317 & 318.
- ---. Description of a new Earth-Snake from Travancore. 7. c. p. 318.
- Description of a new Tree-Frog from New South Wales. P. Linn. Soc. N.S.W. (2) vii, 1892, p. 403.
- ---. An Investigation into the Variations of the Viper in Great Britain. Zool. (3) xvi, 1892, pp. 87-93.
- --... Fecundation in the Tailed Batrachians. A summary Review of recent Discoveries. Morph. JB., Syst. vi, 1892, pp. 447-452.
- ---. The Poisonous Secretion of Batrachians. T. c. pp. 185-190, figs.
- BOUTAN, L. Excursion zoologique à la Montagne de Hummoun ul Faroun (Montagne des bains de Pharaon). Arch. Z. expér. (2) x, 1892, pp. 1-22.
- BREHM, A. C. Tierlebeu, Dritte Auflage. VII. Die Kriechthiere und Lurche. Neubearbeitet von O. Beettger and Pechuel-Loesche. Leipzig & Vienna: 1892, 8vo, xiv, 825 pp., figs., 16 pls., 1 map.
- Burckhardt, R. Das Gehirn von Triceratops flabellatus, Marsh. JB. Mineral. 1892, ii, pp. 71 & 72, fig.
- BUTLER, G. W. On the Subdivision of the Body-cavity in Snakes. P. Z. S. 1892, pp. 477-493, pl. xxviii.
- CAMERANO, L. Nota di Biologia alpina. 111. Dell'azione dell'acqua corrente e della luce sullo sviluppo degli Anfibi Anuri. Boll. Mus. Zool. Tor. viii, 1893, No. 140.
- Caruccio, A. Su' Serpenti non velenosi della Provincia di Roma. Boll. Soc. Rom. i, 1892, pp. 39-57.
- CHIARUGI, G. Sullo sviluppo del nervo olfattivo nella Lacerta muralis. Nota preliminare. Monit. Zool. Ital. iii, 1892, pp. 211 & 212. French translation, Arch. Ital. Biol. xviii, 1893, pp. 363 & 364.
- COPE, E. D. On the Homologies of the Posterior Cranial Arches in the Reptilia. Tr. Am. Phil. Soc. xvii, 1892, pp. 11-26, pls. i-v. Abstract in Am. Nat. xxvi, 1892, pp. 407 & 408, pl. xvii.

- [COPE, E. D.] A Synopsis of the Species of the Teid Genus Cnemidophorus. Tr. Am. Phil. Soc. xxvii, 1892, pp. 27-52, pls. vi-xiii [recorded from separate copy].
- ---.. A Contribution to the Vertebrate Palsontology of Texas. P. Am. Phil. Soc. xxx, 1892, pp. 123-131.
- —. On Tiaporus, a new Genus of Teiidæ. T. c. pp. 132 & 133, pl. —.
- —. The Osteology of the Lacertilia. T. c. pp. 185-221, pls. ii-vi.
- —. On the Skull of the Dinosaurian Lælaps incrassatus, Cope. T. c. pp. 240-245.
- ---. On the Phylogeny of the Vertebrata. T. c. pp. 278-281, figs.
- —... The Batrachia and Reptilia of North Western Texas. P. Ac. Philad. 1892, pp. 331-337.
- A Critical Review of the Characters and Variations of the Snakes of North America. P. U. S. Nat. Mus. xiv, 1892, pp. 589-694.
- —... On Degenerative Types of Scapular and Pelvic Arches in the Lacertilia. J. Morph. vii, 1892, pp. 223-244, pl. xiii.
- —. On a New Spade-Foot from Texas. Am. Nat. xxvii, 1893, pp-155 & 156, fig.
- —. A New Species of Eutania from Western Pennsylvania. Op. cit. xxvi, 1892, pp. 964 & 965, fig.
- —... The Vertebrata of the Tertiary and Cretaceous Rocks of the North-West Territory. Contr. Canad. Pal. iii, 1891, 25 pp., 14 pls.
- ---... Remarks on the communication "Ein fossiler Giftzahn," by Dr. F. Kinkelin. Zool. Anz. xv, 1892, p. 224.
- Dehner, H. Ueber die sogenannte parthenogenetische Furchung des Frosch-Eies. Verh. Ges. Würzb. (2) xxvi, 1892, pp. 1-18, pl. i.
- DEKHUYSEN, M. C. Ueber das Blut der Amphibien. Verh. Anat. Ges. 1892, pp. 90-103, pl. i.
- DEL PRATO, D. A. Le Raccolte Zoologiche fatte nel Congo dal Cav. Giuseppe Corona. Parma: 1893, 8vo, 14 pp.
- DE Vis, C. W. Zoology of British New Guinea. Part 1. Vertebrata. Ann. Queensland Mus. i, 1892, No. 2, pp. 3-12.
- Dollo, L. Nouvelle Note sur le Champsosaure, Rhynchocéphalien adapté à la vie fluviatile. Bull. Soc. Belg. Géol. v, Mém. pp. 151-199, pls. vi-viii (1892).
- —. Sur l'origine de la nageoire caudale des Ichthyosaurus. Op. cit. vi, Proc. Verb. pp. 167-174, figs. (1892).
- —. Sur la morphologie des Côtes. Bull. Sci. Fr. Belg. xxiv, 1892, pp. 113-129.
- —. Sur la morphologie de la Colonne Vertébrale. T. c. 19 pp. [separate copy].

- Douglas-Ogilby, J. Descriptions of three new Australian Lizards. Rec. Austral, Mus. ii, 1892, pp. 6-11.
- —. On some undescribed Reptiles and Fishes from Australia. T. c. pp. 23-26.
- Douglass, G. N. On the Herpetology of the Grand Duchy of Baden. Zool. (3) xvi, 1892, pp. 15-19 & 211-222.
- DRASCH, —. Ueber die Giftdrüsen des Salamanders. Verh. Anat. Gez. 1892, pp. 244-253.
- DRECKMANN, F. A rare Snake (Psammophis longifrons). J. Bomb. N. H. Soc. vii, 1892, pp. 406 & 407.
- Dugės, A. Descripcion del esqueleto del Rhinophrynus dorsalis, D. B. Nat. Mex. (2) ii, 1892, pp. 98 & 99, pl. v.
- El Dendrophidium dendrophie, Schl. T. c. pp. 100 & 101, pl. v.
- Edinger, L. Untersuchungen über die vergleichende Anatomie des Gehirns: 2. Das Zwischenhirn. Erster Teil: Das Zwischenhirn der Selachier und der Amphibien. Abh. Senck. Ges. xviii, 1892, 55 pp., 5 pls. Abstract in An. Anz. vii, 1892, pp. 472-476.
- EMERY, C. Ulteriori studi sullo scheletro della mano degli Anfibi Anuri. Atti Acc. Pontif. Lincei (5) i, 1892, Rend. pp. 203-206.
- ETHERIDGE, R., JUN. On Further Traces of *Meiolania* in New S. Wales. Rec. Austral. Mus. ii, 1893, pp. 39-41.
- EYCLESHYMER, A. C. The Development of the Optic Vesicles in Amphibia. J. Morph. viii, 1893, pp. 189-193, figs.
- —. Paraphysis and Epiphysis in Amblystoma. Anat. Anz. vii, 1892, pp. 215-217.
- FARRINGTON, O. C. The Nephrostomes of *Bana*. Tr. Conn. Ac. viii, 1892, pp. 309-314, pl. xxiv.
- FEORTISTOW, A. E. On the Physiology of the Rattle of Crotulus durissus. Translation. Ann. N. H. (6) xi, 1893, pp. 54-58.
- FERREIRA, J. B. Sobre o "Acanthodaetylus" de Portugal. J. Sci. Lisb. (2) ii, 1892, pp. 188-194.
- —. Sur l'existence du "Triton palmatus" (Schn.) en Portugal. T. c. pp. 195-198.
- —. Revisão dos Reptis e Batrachios de Portugal. T. c. pp. 268-290.
- FICK, R. Ueber die Befruchtung des Axolotleies. Vorläufige Mitteilung. Anat. Anz. vii, 1892, pp. 818–821.
- Field, H. H. Ueber streng metamere Anlage der Niere bei Amphibien. Verh. deutsch. Zool. Ges. ii, 1892, pp. 113-117.
- FLETCHER, J. J. Contributions to a more exact knowledge of the Geographical Distribution of the Australian Batrachia. No. III. P. Linn. Soc. N.S.W. (2) vii, 1892, pp. 7-19.

- FRAAS, E. Ueber einen neuen Fund von Ichthyosaurus in Württemberg. JB. Mineral. 1892, ii, pp. 87-90, figs.
- ----. Ichthyosaurus numismalis, E. Fraas. JH. Ver. Württ. xlviii, 1892, pp. 22-31, pl. i.
- GAGE, S. P. A preliminary account of the brain of *Diemyctylus virides-cens*, based upon sections made through the entire head. Abstract. P. Am. Ass. xli, 1892, p. 197.
- GARMAN, H. A Synopsis of the Reptiles and Amphibians of Illinois. Bull. Illin. Lab. N. H. iii, 1892, pp. 215-389, pls. ix-xv.
- GARMAN, S. The Reptiles of the Galapagos Islands. From the collections of Dr. Geo. Baur. Bull. Ess. Inst. xxiv, 1892, pp. 73-87.
- —... On Reptiles collected by Dr. Geo. Baur near Guayaquil, Ecuador. T. c. pp. 88-95.
- ---. On Cophias and Bachia. T. c. pp. 96 & 97.
- —. On Texas Reptiles collected by Mr. F. W. Wamsley for Professor J. W. P. Jenks. T. c. pp. 98-109.
- The Reptilian Rattle. Science, xx. 1892, pp. 16 & 17, figs.
- GAUDRY, A. Les Pythonomorphes de France. C.R. cxv, 1892, pp. 303 & 304, and Mém. Soc. Géol. iii, 1892, No. 3, 13 pp., pls. i & ii.
- —. L'Ichthyosaurus burgundiæ. Bull. Soc. Autun, v, 1892, pp. 365-372, pl. xv.
- GAUPP, E. Grundzüge der Bildung und Umbildung des Primordialcraniums von Rana fusca. Verh. Anat. Ges. 1892, pp. 183-190.
- GIACOMINI, E. Contributo alla migliore conoscenza degli annessi fetali nei Rettili. Nota preventiva. Monit. Zool. Ital. iii, 1892, pp. 126–128, 157–164, & 185–196.
- GIRTANNER, A. Die Griechische Landschildkröte im Garten. Zool. Gart. xxxiii, 1892, pp. 349 & 350.
- GORJANOVIĆ-KRAMBERGER, C. Aigialosaurus, eine neue Eidechse aus den Kreideschiefern der Insel Lesina, mit Rücksicht auf die bereits beschriebenen Lacertiden von Comen und Lesina. Glasnik Nadar. Drustva, vii, 1892, pp. 74–106, pls. iii–iv. The same paper in Croatian, Rad jugoslav. akad. cix, 1892, pp. 96–123, pls. i & ii.
- GREENOUGH, H. S. Sur les homologies des premiers stades suivant la segmentation chez les Batraciens. Bull. Soc. Z. Fr. xvii, 1892, pp. 57-59.
- GREGORIO, A. DE. Sulla Trionyx ragusensis, De Greg., et la T. melitensis, Lydek. Nat. Sicil. xi, 1892, pp. 182 & 183.
- GRIESBACH, H. Ueber Plasmastructuren der Blutkörperchen im kreisenden Blute der Amphibien. Leuckart Festschrift, 1892, pp. 215–227, pl. xxiii.

- GÜNTHER, A. List of the Species of Reptiles and Fishes collected by Mr. A. E. Pratt on the Upper Yang-tze-Kiang and in the province Sze-chuen, with Description of the New Species. Appendix 11 to PRATT'S "Snows of Tibet" (London: 1892, 8vo), pp. 238-250, pls. i-iv.
- —. Report on a Collection of Reptiles and Batrachians transmitted by Mr. H. H. Johnston, C.B., from Nyassaland. P. Z. S. 1892, pp. 555– 558, pl. xxxiii–xxxv (1893).
- HAHN, G. Les Axolotls et leur Métamorphose. Rev. Quest. Sci. (2) i, 1892, pp. 178-205.
- HALY, A. Essay on the Construction of Zoological Tables, with a Tabular Diagnosis of the Snakes of Ceylon. J. A. S. (Ceylon), xi, 1892, pp. 22-32, 183-232, figs.
- HARVIE-BROWN, J. A., & BUCKLEY, T. E. A Vertebrate Fauna of Argyll and the Inner Hebrides. Edinburgh: 1892, 8vo, 252 pp.
- HASSE, C. Die Entwicklung der Wirbelsäule von Triton tæniatus. Erste Abhandlung über die Entwicklung der Wirbelsäule. Z. wiss. Zool. liii (Suppl.) 1892, pp. 1-20, pls. i-iii.
- —. Die Entwicklung der Wirbelsäule der ungeschwänzten Amphibien. Zweite Abhandlung über die Entwicklung der Wirbelsäule. Z. wiss. Zool. lv, 1892, pp. 252-264, pl. xii.
- —... Allgemeine Bemerkungen über die Entwickelung und die Stammesgeschichte der Wirbelsäule. Anat. Anz. viii, 1893, pp. 288-289.
- HAY, O. P. On the Ejection of Blood from the Eyes of Horned Toads. P. U. S. Nat. Mus. xv, 1892, pp. 375-378.
- —. On the Breeding Habits, Eggs, and Young of certain Snakes. T. c. pp. 385-397.
- ---. Description of a supposed new Species of Storeria from Florida, Storeria victa. Science, xix, 1892, p. 199.
- Heidenhein, M. Notiz betreffend eine rudimentäre Drüse bei den Weibehen der einheimischen Tritonen. Anat. Anz. vii, 1892, pp. 432-435, figs.
- HERRERA, A. L. Sur le pleur de sang chez le Batrachosoma asio, Cope. Le Nat. xiv, 1892, p. 114.
- HERTWIG, O. Urmund und Spina bifida, eine vergleichend-morphologische, teratologische Studie an missgebildeten Froscheiern. Arch. mikr. Anat. xxxvii, 1892, pp. 353-503, pls. xvi-xx.
- Hochstetter, F. Beiträge zur Entwicklungsgeschichte des Venensystems der Amnioten. II. Reptilien (*Lacerta*, *Tropidonotus*). Morph. JB. xix, 1892, pp. 428-501, pl. xv-xvii.
- HOFFMANN, C. K. Bijdrage tot de Kennis der Ontwikkelingsgeschiedenis van het Aderlijke Bloedvatenstelsel bij de Reptiliën, Verh. Ak. Amst. xxix, 1892, pp. 1-19, pls. i & ii,

- HOLLA: D, W. J. A Nest Bu'lding Frog. Am. N.t. xxvii, 1893, pp. 67 & 68.
- HONNORAT-BASTIDE, E. F. Reptiles et Batraciens des Basses-Alpes. C.R. Ass. Fr. Sci. 1891, ii, pp. 583-586.
- Houssay, F. Sur la circulation embryonnaire dans la tête chez l'Axolotl. C.R. exv, 1892, pp. 132-135.
- Howes, G. B. Notes upon the Shoulder Girdle of certain Dicynodontoid Reptiles. J. Anat. Phys. xxvi, 1892, pp. 403-405.
- HULKE, J. W. On the Shoulder Girdle in Ichthyosauria and Sauropterygia. P. R. Soc. lii, 1892, pp. 233-255, figs.
- JACK, R. L., & ETHERIDGE, R., JUN. The Geology and Paleontology of Queensland and New Guinea. Brisbane and London: 1892, 8vo, 768 pp.
- JORDAN, E. O., & EYCLESHYMER, A. C. The Cleavage of the Amphibian Ovum. Anat. Anz. vii, 1892, pp. 622-624.
- JUNGERSEN. H. F. E. Om Udviklingen af den Müllerske Gang (Æggelederen) hos Padderna. Vid. Medd. 1892 (1893), pp. 32-72, pls. ii & iii.
- KERMODE, P. M. C. Contributions to a Vertebrate Fauna of the Isle of Man. Zool. (3) xvii, 1893, pp. 64 & 65.
- KINGSLEY, J. S. The Head of an Embryo Amphiuma. Am. Nat. xxvi, 1892, pp. 671-680.
- KINKELIN, F. Ein fossiler Giftzahn. Zool. Anz. xv, 1892, pp. 93 & 94.
- KLAATSCH, H. Zur Morphologie der Mesenterialbildungen am Darmkanal der Wirbelthiere. I. Theil. Amphibien und Reptilien. Morph. JB. xviii, 1892, pp. 385-450, figs., pl. xii.
- KLINCKOWSTRÖM, A. DE. Le premier développement de l'oeil pinéal, l'epiphyse et le nerf parietal chez *Iguana tuberculata*. Anat. Anz. viii, 1893, pp. 289-299, figs.
- KNAUTHE, K. Zur Biologie der Amphibien. Zool. Anz. xv, 1892, pp. 20-23.
- König, A. Die Kriechthierfauna Tunesiens. SB. niederrhein. Ges. 1892, pp. 3-26.
- KÜKENTHAL, W. Zur Entwicklung des Handskelettes des Krokodils. Morph. JB. xix, 1892, pp. 42-55, pl. ii.
- Ichthyosaurier und Wale. JB. Mineral. 1892, i, pp. 161-166.
- LATASTE, F. Etudes sur la faune Chilienne. 1. Note sur les Lézards (Ordre des Sauriens). Act. Soc. Sci. Chili, i, 1892, pp. 7-40.
- —... Le péritoine du Python de Séba, réponse aux critiques de M. Gérard W. Butler. Op. cit. ii, 1893, pp. clxvii-clxx.

- LEYDIG, F. Receptacula seminis der Urodelen. Zur Frage der Deutung. Zool. Anz. xv, 1892, pp. 309-312, 356.
- —. Integument brünstiger Fische und Amphibien. Biol. Centralbl. xii, 1892, pp. 205-221.
- —. Zum Integument niederer Wirbeltiere abermals. T. c. pp. 444-467.
- —. Blaufarbiger Wasserfrosch; Leuchtflecken der Ellritze. Zool. Gart. xxxiii, 1892, pp. 1-7.
- LORTET, L. Les Reptiles fossiles du Bassin du Rhône. Arch. Mus. Lyon, v, 1892, pp. 1-139, figs., pls. i-xii.
- LOVISATO, D. Nuovi resti di coccodrilliano fossile nel miocene di Nurri. Atti Acc. Lincei (5) i, 1892, Rend., pp. 436-439.
- Lucas, A. H. S. Notes on the Distribution of Victorian Batrachians, with Descriptions of two new Species. P. R. Soc. Vict. (2) iv, 1892, pp. 59-64.
- LYDEKKER, R. On Part of the Pelvis of Polacanthus. Q. J. Geol. Soc. xlviii, 1892, pp. 148 & 149, figs.
- —. Note on Two Dinosaurian Foot-bones from the Wealden. T. c. pp. 375 & 376, fig.
- —. Recent Advances in Knowledge of the Ichthyosaurian Reptiles. Nat. Sci. i, 1892, pp. 514-521, figs.
- MACALLUM, A. B. Studies on the Blood of Amphibia. Tr. Can. Inst. ii, 1892, pp. 221-260, pl. i.
- MARSH, O. C. The Skull of *Torosaurus*. Am. J. Sci. (3) xliii, 1892, pp. 81-84, pls. ii & iii.
- —. Notice of New Reptiles from the Laramie Formation. T. c. pp. 449-453, figs.
- Notes on Triassic Dinosauria. T. c. pp. 543-546, pls. xv-xvii.
- —. Notes on Mesozoic Vertebrate Fossils. Op. cit. xliv, 1892, pp. 171-176, pls. ii-v.
- —. Restorations of Claosaurus and Ceratosaurus. T. c. pp. 343-349, pls, vi & vii.
- ---. The Skull and Brain of Claosaurus. Op. cit. xlv, 1893, pp. 83-86, pls. iv & v.
- --- Restoration of Anchisaurus. T. c. pp. 169-170, pl. vi.
- MARSHALL, W. B. Necturus maculatus in the Hudson River. Am. Nat. xxvi, 1892, pp. 779 & 780.
- MARTIN, C. J. Observations on the Poisonous Constituents of the Venom of the Australian Black Snake (*Pseudechis porphyriacus*). P. Linn. Soc. N.S.W. (2) vii, 1892, pp. 319-322.

- MARTIN, C. J.] The Venom of the Australian Black Snake (Pseudechis porphyriacus). J. R. Soc. N.S.W. xxvi, 1892, pp. 240-264.
- ARTIN, R. Sur l'habitat de la Couleuvre verte et jaune. Bull. Soc. Ouest France, ii, 1892, pp. 149 & 150.
- ARTIN, R., & ROLLINAT, R. Catalogue des Reptiles, Batracieus et Poissons du Département de l'Iudre. Mém. Soc. Zool. v, 1892, pp. 30-45.
- ATSCHIE, P. Ueber eine kleine Sammlung von Säugethieren und Reptilien, welche Herr L. Conradt aus Usambara (Deutsch-Ostafrika) heimgebracht hat. SB. nat. Fr. 1892, pp. 101-110.
- AURER, F. Haut-Sinnesorgane, Feder- und Haaranlagen, und deren gegenseitige Beziehungen, ein Beitrag zur Phylogenie der Säugethierhaare. Morph. JB. xviii, 1892, pp. 717-804, figs., pls. xxiv-xxvi.
- AZZA, F. Caso di dicefalia derodimica in un Anguis fragilis. Atti Soc. Ligust. iii, 1892, pp. 256-261, pl. ii.
- ÉHELŸ, L. v. Magyarország Barna Békál (Ranæ fuscæ Hungariæ).

  Math. term. köz. xxv, 1892, pp. 1-63, pls. i-viii.
- —. Beiträge zur Kenntniss der Bombinator-Arten sowie deren Standorte und Verbreitung in Ungarn. Math. Nat. Ber. Ung. x, 1892, pp. 55-79, pls. iv & v.
- —. Die Herpetologischen Verhältnisse des siebenbürgischen Burzenlandes. Beiträge zu einer Monographie der k. freien Stadt Kronstadt. Kronstadt: 1892, 8vo, 91 pp.
- [EYER, A. Ueber das Vorderhirn einiger Reptilien. Z. wiss. Zool. lv, 1892, pp. 63-133, pls. iv & v.
- [EYER, A. B., & HELW, F. vi. Jahresbericht (1890) der Ornithologischen Beobachtungstationen im Königreich Sachsen. Berlin: 1892, 4to, 135 pp.
- Іїма-Рацияво, F. Rettili ed Anfibi Nebrodensi (Continua). Nat. Sicil. xi, 1892, pp. 114-120, 239-252, and xii, 1892-93, pp. 52-56, 75-80, & 148-150.
- IINGAZZINI, P. L'Oolisi nella Seps chalcides. Atti Acc. Lincei (5) i, 1892, Rend., pp. 41-45.
- IITROPHANOPH, P. Researches on the development of Vertebrate Animals [Russian]. Trav. Soc. Varsovie, ii, 1892, 251 pp., 7 pls.
- ITTSUKURI, K. On the Foetal Membranes of Chelonia. J. Coll. Sci. Japan, iv, 1891, pp. 1-53, pl. i-x.
- —. Further Studies on the Formation of the Germinal Layers in Chelonia. Op. cit. v, 1892, pp. 35-52, pls. ii-iv.
- IOCQUARD, F. Voyage de M. Chaper à Bornéo. Nouvelle contribution à la faune herpétologique de Bornéo. Mém. Soc. Zool. v, 1892, pp. 190-206, pl. vii.

- [Mocquard, F.] Description de deux Ophidiens et d'un Batracien d'espèces nouvelles. Le Nat. xiv, 1892, p. 35.
- McLLER, F. Siebenter Nachtrag zum Katalog der herpetologischen Sammlung des Basler Museums. Festschr. 75. Best. Nat. Ges. Basel, 1892, pp. 195-215, pl. iii.
- NEGRI, A. Trionici eocenici ed oligocenici del Veneto. Mem. Soc. Ital. (3) viii, 1892, No. 7, 53 pp., 5 pls.
- NEWTON, E. T. On some new Reptiles from the Elgin Sandstone. (Abstract). P. R. Soc. lii, 1892, pp. 389-391.
- —. Note on an Iguanodont Tooth. Geol. Mag. (3) ix, 1892, pp. 49 & 50, fig.
- NIEMANN, F. Beiträge zur Morphologie und Physiologie der Oberlippendrüsen einiger Ophidier. Arch. f. Nat. lviii, 1, 1892, pp. 262-286, pl. xiv.
- Norris, H. W. Studies on the Development of the Ear of Amblystoma.

  Part I. Development of the Auditory vesicle. J. Morph. vii, 1892,
  pp. 23-34, pls. iii & iv.
- Oppel, A. Die Befruchtung des Reptilieneies. Arch. mikr. Anat. xxxix, 1892, pp. 215-290, pls. ix-xii.
- OSTROUMOFF, A. Zoologische Excursion nach Mangyschlak und Busatchi (Russian text). Protok. obsch. estest. Kazan, 1889-90, Append. No. 113 (1890), 16 pp.
- PARKER, W. N. Experiments on Respiration in Tadpoles of the Common Frog (Rana temporaria). Rep. Brit. Ass. 1891, p. 694.
- Peracca, M. G. Descrizione di nuove specie di Rettili e Anfibi di Madagascar. Boll. Mus. Zool. Tor. vii, 1892, No. 112.
- Perrin, A. Remarques sur le pied des Batraciens et des Sauriens. C.R. cxv, 1892, pp. 885-887.
- —. Contribution à l'étude de la myologie comparée : Membre postérieur chez un certain nombre de Batraciens et de Sauriens. Bull. Sci. Fr. Belg. (4) iii, 1892, pp. 373-552, pls. xvi-xxiii.
- PIEL DE CHURCHEVILLE, H. & T. Note sur la présence du Tropidonote à collier, variété à deux raies, *Tropidonotus natrix*, var. bilineata, dans les environs de Nantes. Bull. Soc. Ouest France, ii, 1892, pp. 35-38, pl. ii.
- POHLIG, H. Altpermische Saurierfährten, Fische und Medusen der Gegend von Friedrichroda i. Thür. Leuckart Festschrift, 1892, pp. 59-64, figs., pl. vii.
- Positano-Spada, D. Sulla identitá specifica della Lacerta muralis e della Lacerta serpa. Boll. Soc. Rom. i, 1892, pp. 89-94.
- —. Sopre une nuova varietà di Lacerta muralis. T. c. pp. 154-157, figs.

- Quelch, J. J. Venom in Harmless Snakes. Zool. (3) xvii, 1893, pp. 30 & 31.
- REMY SAINT-LOUP, —. Sur une disposition intermédiaire à celles qui ont fait etablir un caractère anatomique différentiel des Plagiotrèmes et des Hydrosauriens. CR. Soc. Biol. (9) iv, 1892, pp. 176-179, figs.
- RETZIUS, G. Die Nervenendigungen in dem Geschmacksorgan der Sängethiere und Amphibien. Biol. Unters. (2) iv, 1892, pp. 19-32, pls. vii-x.
- ---. Die Nervenendigungen in den Endknospen resp. Nervenhügeln der Fische und Amphibien. T. c. pp. 33-36.
- ----. Ueber die sensiblen Nervenendigungen in den Epithelien bei den Wirbelthieren. T. c. pp. 37-44, pls. xi-xv.
- Rex, H. Beiträge zur Morphologie der Hirnvenen der Amphibien. Morph. JB. xix, 1892, pp. 295-311, pl. xi.
- Röse, C. Ueber die Zahnleiste und die Eischwiele der Sauropsiden. Anat. Anz. vii, 1892, pp. 748-758, figs.
- —. Ueber die Zahnentwickelung der Krokodile. Verh. Anat. Ges. 1892, pp. 225-227.
- Roux, W. Ueber das entwickelungsmechanische Vermögen jeder der beiden ersten Furchungszellen des Eies. T. c. pp. 22-62.
- Rumpf, G. Aus dem Leben der griechischen Landschildkröte, Testudo græca. Zool. Gart. xxxiii, 1892, pp. 260-264.
- Sarasin, P. & F. Ueber das Gehörorgan der Cæciliiden. Anat. Anz. vii, 1892, pp. 812-815.
- Scharff, R. F. Is the Frog a Native of Ireland? Irish Nat. i, 1893, pp. 1-6.
- SCHLAMPP, K. W. Das Auge des Grottenolmes (Proteus anguineus). Z. wiss. Zool. liii, 1892, pp. 537-557, pl. xxi.
- SCHULZE, E., & BORCHEDING, F. Fauna Saxonica. Amphibia et Reptilia. Verzeichniss der Lurche und Kriechtiere des Nordwestlichen Deutschlands. Jena: 1893, 8vo, 94 pp., figs.
- Schulze, F. E. Ueber die inneren Kiemen der Batrachierlarven.

  II. Mittheilung. Skelet, Musculatur, Blutgefässe, Filterapparat,
  Respiratorische Anhänge und Athmungsbewegungen erwachsener
  Larven von Pelobates fuscus. Abb. Ak. Berl. 1892, 66 pp., 6 pls.
- Sclater, P. L. Remarks upon the habits of a South-African Snake (Dasypeltis scabra). P. Z. S. 1892, p. 476.
- Sclater, W. L. List of the Batrachia in the Indian Museum. London: 1892, 8vo, 43 pp.
- —. On some Specimens of Frogs in the Indian Museum, Calcutta, with Descriptions of several new Species. P. Z. S. 1892, pp. 341-348, pl. xxiv.

- SEELLEY, H. G. Researches on the Structure, Organization and Classification of the Fossil Reptilia. VII. Further observations on Pareiasuurus. Phil. Tr. claxxiii (B), 1892, pp. 311-370, figs., pls. xvii-xxiii.
- —. The Nature of the Shoulder Girdle and Clavicular Arch in Sauropterygia. P. R. Soc. li, 1892, pp. 119-151, figs.
- —. On the Os pubis of Polacanthus foxii. Q. J. Geol. Soc. xlviii, 1892, pp. 81-85, figs.
- —. Contribution to a Knowledge of the Saurischia of Europe and Africa. T. c., Proc., pp. 188-191.
- ---. On a New Reptile from Welte Vreden (Beaufort West), Eunoto-saurus africanus (Soeley). T. c. pp. 583-585, figs.
- ---. The Mesosauria of South Africa. T. c. pp. 586-604, pl. xviii.
- SEILLER, R. v. Die Zungendrüsen von Lacerta. Leuckart Festschrift, 1892, pp. 250-258, pls. xxv & xxvi.
- SIEBENROCK, F. Zur Kenntniss des Kopfskelettes der Scincoiden, Anguiden, und Gerrhosauriden. Ann. Hofmuseum Wien, vii, 1892, pp. 163-196, pls. i & ii.
- —. Ueber Wirbelassimilation bei den Sauriern. T. c. pp. 373-378, figs.
- SKUPHOS, T. G. Vorläufige Mittheilung über Parthanosaurus zitteli, cinen neuen Saurier aus der Trias. Zool. Anz. xvi, 1893, pp. 67-69.
- SLUITER, C. P. Das Jacobson'sche Organ von Crocodilus porosus (Schn.). Anat. Anz. vii, 1892, pp. 540-545, figs.
- SMIRNOW, A. E. Ueber die Nerven der Sehnen bei Rana temporaria, R. esculenta, und Bufo vulgaris (Russian text). Protok. obsch. estest. Kazan, 1889-90, Append. No. 112 (1890), 6 pp.
- STEINDACHNER, F. Ueber die typischen Exemplare von Lacerta mosorensis, Kolomb., 1886 (= Lacerta koritana, Tom., 1889). Ann. Hofmuseum Wien, vii, 1892, pp. 235-240, pl. xvi.
- —. Ueber zwei noch unbeschriebene Nototrema-Arten aus Ecuador und Bolivia. Anz. Ak. Wien, 1892, pp. 175 & 176, and SB. Ak. Wien, ci, i, 1892, pp. 837-842, 2 pls.
- STEJNEGER, L. Preliminary Description of a new Genus and Species of Blind Cave Salamander from North America. P. U. S. Nat. Mus. xv, 1892, pp. 115-117, pl. ix.
- Strahl, —. Die Rückbildung reifer Eierstockseier am Ovarium von Lacerta agilis. Verh. Anat. Ges. 1892, pp. 190-195.
- STRONG, O. S. The Structure and Homologies of the Cranial Nerves of the Amphibia, as determined by their Peripheral Distribution and Internal Origin. Part 2. Anat. Anz. vii, 1892, pp. 467-471.
- TAYLOR, W. E. Ophidia of Nebraska. Report of the Nebraska State Board of Agriculture, 1891.

- [Taylor, W. E.] Catalogue of the Snakes of Nebraska, with Notes on their Habits and Distribution. Am. Nat. xxvi, 1892, pp. 742-752.
- Test, F. C. Fish-Cultural Investigations in Montana and Wyoming. Annotated List of Reptiles and Batrachians collected. Bull. U. S. Fish Comm. f. 1891, pp. 57-59 (1892).
- VAILLANT, L. Recherches biologiques faites à la Ménagerie des Reptiles (premier Article). Contribution à l'étude de l'alimentation chez les Ophidiens. N. Arch. Mus. (3) iv, 1892, pp. 221-234.
- ---. Remarques sur l'alimentation chez les Ophidiens. C.R. cxv, 1892, pp. 277-279.
- VERRILL, G. E. Notes on the Fauna of the Island of Dominica, British West Indies, with Lists of the Species obtained and observed by G. E. and A. H. Verrill. Tr. Conn. Ac. viii, 1892, pp. 315-355.
- VINCIGUERRA, D. Rettili e Batraci di Engano raccolti dal Dott. Elio Modigliani. Ann. Mus. Genov. (2) xii, 1892, pp. 517-526.
- Virchow, H. Das Dotterorgan der Wirbelthiere. Z. wiss. Zool. liii (Suppl.), 1892, pp. 161-206, pl. x.
- Viel. Tzkow, A. On the Oviposition and Embryonic Development of the Crocodile. Translation. Ann. N. H. (6) ix, 1892, pp. 66-72.
- WERNER, F. Untersuchungen über die Zeichnung der Wirbelthiere. Zool. JB., Syst. vi, 1892, pp. 155-229, pls. vi-x.
- Zoologische Miscellen. 11. Noch etwas über die Zeichnung der Tiere. 111. Die Epitrichialskulptur der Schlangenepidermis. Biol. Centralbl. xii, 1892, pp. 272-280.
- Herpetologische Nova. Zool. Anz. xvi, 1893, pp. 81-84.
- ----. Ausbeute einer herpetologischen Excursion nach Ost-Algerien. Verh. z.-b. Wien, xlii, 1892, pp. 350-355.
- —. Tierleben in Ost-Algerien. Zool. Gart. xxxiii, 1892, pp. 264-273.
- ---. Biologische Notizen aus der Reptilienwelt. T. c. pp. 367-372.
- WERNER, H. Bemerkungen über den Scheltopusik und die Treppennatter. T. c. pp. 38-41.
- WIEDERSHEIM, R. Das Gliedmassenskelet der Wirbelthiere, mit besonderer Berücksichtigung des Schulter- und Beckengürtels bei Fischen, Amphibien und Reptilien. Jena: 1892, 8vo, 266 pp., figs., 17 pls.
- Die Phylogenie der Beutelknochen. Eine entwicklungsgeschichtlich-vergleichend anatomische Studie. Z. wiss. Zool. liii (Suppl.) 1892, pp. 43-66, pls. vi & vii.

- WILDER, H. H. Die Nasengegend von Menopoma alleghaniense und Amphiuma tridactylum, nebst Bemerkungen über die Morphologie des Ramus ophthalmicus profundus trigemini. Morph. JB. v, 1892, pp. 155-176, pls. xii & xiii.
- WILL, L. Beiträge zur Entwicklungsgeschichte der Reptilien. 1. Die Anlage der Keimblätter beim Gecko (*Platydactylus facetanus*, Schreib.). Op. cit. vi, 1892, pp. 1-160, figs., pls. i-xi.
- —. Zur Kenntniss der Schildkröten-Gastrula. Biol. Centralbl. xii, 1892, pp. 182-192, figs.
- WILLISTON, S. W. Kansas Pterodactyls. Kansas Quart. i, 1892, pp. 1-13, pl. i.
- --- & Case, E. C. Kansas Mosasaurs. T. c. pp. 15-32, pls. ii-vi.
- WITCHELL, C. A., & STRUGNELL, W. B. The Fauna and Flora of Gloucestershire. Stroud: 1892, 8vo, 301 pp.
- ZIEGLER, F. Zur Kenntniss der Oberflächenbilder bei Rana-Embryonen.
  Anat. Anz. vii, 1892, pp. 211-215, figs.

# II.—WORKS AND PAPERS DEALING WITH BOTH REPTILES AND BATRACHIANS.

O. Bettger & Pechuel-Loesche have prepared the seventh volume, dealing with the Reptiles and Batrachians, of the third revised edition of Brehm's Tierleben (Leipzig & Vienna: 1892).

Remarks on phylogeny of the Reptiles and Batrachians; E. D. COPE, P. Am. Phil. Soc. xxx, 1892, pp. 278-280.

Development and phylogeny of the vertebral column; C. Hasse, Anat. Anz. viii, 1893, pp. 288 & 289, summarises his views on this. He traces two phyla: one embracing the Ganoids, Teleostei, Tailless Batrachians, and Amniota; the other the Selachians, Dipnoi, and Tailed Batrachians.

The morphology of the vertebral column and ribs, by L. Dollo, Bull. Sci. Fr. Belg. xxiv, 1892 [quoted from separate copies].

R. WIEDERSHEIM'S book, "Das Gliedmasenskelet der Wirbelthiere" (see p. 15), deals chiefly with the pectoral and pelvic arches and limbs of Reptiles and Batrachians.

Philogeny of the marsupial bones of Mammals. R. WIEDERSHEIM, à propos of this, enters into details on the morphology of the pelvis in Reptiles and Batrachians; Z. wiss. Zool. liii (Suppl.), 1892, pp. 1-20, pls. i-iii.

Myology of the hind limb in Lizards and Batrachians; A. Perrin, C.R. cxv, 1892, pp. 885-887, and Bull. Sci. Fr. Belg. (4) iii, 1892, pp. 373-552, has a contribution on.

On the nerve-terminations in the skin of Reptiles and Batrachians. G. Retzius, Biol. Unters. (2) iv, 1892, pp. 19-44, pls. vii-xv.

Copulatory organs in the various groups of Reptiles and Batrachians; REMY SAINT-LOUP, C.R. Soc. Biol. (9) iv, 1892, pp. 176-179, figs., remarks on the difference in these.

Morphology of the vertebrate ear; H. AYERS, J. Morph. vi, 1892, pp. 1-360, pls. i-xii.

Morphology of the peritoneum in Reptiles and Batrachians; H. KLAATSCH, Morph. JB. xviii, 1892, pp. 385-450, figs., pl. xii.

Markings of Reptiles and Batrachians; F. WERNER, Zool. JB., Syst. vi, 1892, pp. 155-229, pls. vi-x, deals with their signification.

Seventh supplement to his Catalogue of the Herpetological Collection in the Basle Museum; F. Müller, Festschr. Nat. Ges. Basel, 1892, pp. 195-215, pl. iii.

#### III.—FAUNÆ.

#### A.—RECENT.

#### EUROPE.

European Batrachians; G. A. BOULENGER, P. Z. S. 1892, pp. 593-627, pls. xlv-xlvii, describes and figures the Tadpoles.

Isle of Man; KERMODE, Zool. (3) xvii, 1893, pp. 64 & 65.

"Vertebrate Fauna of Argyll and Inner Hebrides" (Edinburgh: 1892, 8vo); J. A. HARVIE-BROWN & T. E. BUCKLEY devote pp. 216–218 to the Reptiles and Batrachians.

"Vertebrate Fauna of Lakeland" (Edinburgh: 1892, 8vo); H. A. MAC-PHERSON, pp. 461-465, deals with the Reptiles and Batrachians.

"Fauna and Flora of Gloucestershire" (Stroud: 1892, 8vo); C. A. WITCHELL & W. B. STRUGNELL, Reptilia and Amphibia, pp. 124-148, figs. Department Indre, France: R. Martin & R. Rollinat, Mém. Soc. Zool. v, 1892, pp. 30-38, enumerate the Reptiles and Batrachians of.

Department Basses Alpes: E. F. HONNORAT-BASTIDE, CR. Ass. Fr. Sci. 1891, ii, pp. 583-586, remarks on the herpetological fauna of the.

France and Germany: J. DE BEDRIAGA, CR. Congrès. Internat. Zool. i, 1892, p. 244, draws attention to some recent introductions of foreign Reptiles and Batrachians.

N. W. Germany: ERWIN SCHULZE & F. BORCHERDING publish a catalogue of the Reptiles and Batrachians of; Jena: 1893, 8vo, 94 pp., figs.

Grand Duchy of Baden: G. N. Douglass concludes his Herpetology of; Zool. (3) xvi, 1892, pp. 15-19 & 211-222.

Kronstadt, Transylvania: L. VON MÉHELŸ, in Beiträge zu einer Monographie der k. freien Stadt Kronstadt (Kronstadt: 1892, 8vo), contributes an article (91 pp.) on the Herpetology of the environs of.

Portugal: J. BETTENCOURT FEREIRA, J. Sci. Lisb. (2) ii, 1892, pp. 268-290, list of the Reptiles and Batrachians.

Province of Rome: A. CARUCCIO, Boll. Soc. Rom. i, 1892, pp. 39-57, describes the harmless Snakes.

Sicily: F. MINA-PALUMBO, Nat. Sicil. xi, 1892, pp. 114-120, 239-252, and xii, 1892-3, pp. 52-56, 75-80, 148-150, articles on the Herpetology of.

#### Asia.

Caucasus District: O. BŒTTGER, Ber. Senck. Ges. 1892, pp. 131-150, has notes on Reptiles and Batrachians.

Mangyechlak and Busatschi: A. OSTROUMOFF, Protok. obsch. estest. Kazan. 1889-90, Append. No. 113 (1890), gives an account, in Russian, of a zoological excursion.

Transcaspian Reptiles: G. A. BOULENGER, P. Z. S. 1891, pp. 622-633, has notes on.

Aden: O. BŒTTGER, Ber. Offenb. Ver. xxix-xxxii, 1892, pp. 61-63, records some Reptiles from.

Upper Yang-tse-kiang and Sze-Chuen: A. GÜNTHER gives a list of the Reptiles and Batrachians collected by Mr. A. E. Pratt, Appendix ii to Pratt's "Snows of Tibet" (London: 1892, 8vo), pp. 238-242, pl. i.

Great Loo Choo I.: G. A. BOULENGER, Ann. N. H. (6) x, 1892, pp. 302-304, remarks on a collection of Reptiles and Batrachians made by Mr. Holst.

India and Ceylon, Salanga I. in Siam, Camboja, Sumatra, and Java: O. Bettger, Ber. Offenb. Ver. xxix-xxxii, 1892, pp. 65-145, reports on collections from.

India and Burma: W. L. SCLATER, P. Z. S. 1892, pp. 341-348, pl xxiv, has notes on the Frogs preserved in the Calcutta Museum.

Ceylon: A. HALY, J. A. S. (Ceylon) xi, 1892, pp. 22-32 & 183-232, table for the identification of the Snakes.

Engano I., S. E. of Sumatra: D. VINCIGUERRA, Ann. Mus. Genov. (2) xii, 1892, pp. 517-526, reported on the Reptiles and Batrachians collected by Modigliani.

Borneo: F. Mocquard, Mém. Soc. Zool. v, 1892, pp. 190-206, pl. vii, reports on a collection of Reptiles and Batrachians from W. Borneo, and also remarks on some of the specimens previously described by himself. G. A. Boulenger, P. Z. S. 1892, pp. 505-508, pls. xxix & xxx, reports on the Reptiles and Batrachians collected by Mr. C. Hose on Mt. Dulit.

#### AFRICA.

Algeria and Tunisia: J. Anderson, P. Z. S. 1892, pp. 11-24, list of Reptiles and Batrachians, with notes.

Algeria: F. WERNER, Verh. z.-b. Wien. xlii, 1892, pp. 350-355, and Zool. Gart. xxxiii, 1892, pp. 264-273, has notes on the Reptiles and Batrachians collected by himself.

Tunisia: A. König, SB. niederrhein. Ges. 1892, pp. 3-26, enumerates, with notes, the Reptiles and Batrachians collected by himself.

Portuguese Guinea: J. V. BARBOZA DU BOCAGE, J. Sci. Lisb. (2) ii, 1892, pp. 183 & 184, enumerates the Reptiles.

Is. of the Gulf of Guinea: J. DE BEDRIAGA, O Istituto, ser. 2, Nos. 6 & 7, 1892, describes the Reptiles and Batrachians obtained by A. F. Moller in the; remarks by J. V. BARBOZA DU BOCAGE, J. Sci. Lisb. (2) ii, 1892, pp. 229-232.

Congo: D. A. DEL PRATO, in a separately published pamphlet (Parma: 1893, 8vo, 14 pp.), gives a list of Reptiles.

Usambara, German East Africa: P. MATSCHIE, SB. nat. Fr. 1892, pp. 102-110, reports on a small collection of Reptiles and Batrachians.

Nyassaland: A. GÜNTHER, P. Z. S. 1892, pp. 555-558, pls. xxxiii-xxxv (1893), reports on a collection of Reptiles and Batrachians.

Transvaal: G. A. BOULENGER gives a list of the Reptiles and Batrachians obtained by W. L. Distant, Appendix to Distant's "Naturalist in the Transvaal" (London: 1892, 8vo), pp. 174-176, fig.

Madagascar: M. G. Peracca, Boll. Mus. Zool. Tor. vii, 1892, No. 112, describes two new Snakes and a new Frog.

#### AMERICA.

North America: E. D. COPE, P. U. S. Nat. Mus. xiv, 1892, pp. 589-694, critical review of the characters and variations of the Snakes.

Nebraska: W. E. TAYLOR, Catalogues of the Snakes, with notes on their habits and distribution; Report of the Nebraska State Board of Agriculture, 1891, and Am. Nat. xxvi, 1892, pp. 742-752.

Montana and Wyoming: F. C. Test, Bull. U. S. Fish. Comm. 1891, pp. 57-59 (1892), remarks on some Reptiles and Batrachians.

Illinois: H. GARMAN, Bull. Illin. Lab. N. H. iii, 1892, pp. 215-389, pls. ix-xv, synopsis of the Reptiles and Batrachians.

Texas: S. GARMAN, Bull. Ess. Inst. xxiv, 1892, pp. 98-109, reports on a collection of Reptiles.

North Western Texas: E. D. Cope, P. Ac. Philad. 1892, pp. 331-337, list of Reptiles and Batrachians.

Dominica, West Indies: G. E. VERRILL, Tr. Conn. Ac. viii, 1892, pp. 350-352 & 355.

Galapagos and Guayaquil: S. GARMAN, Bull. Ess. Inst. xxiv, 1892, pp. 73-95, reports on the Reptiles obtained by G. Baur.

Chili: F. LATASTE, Actes Soc. Sci. Chili, i, 1892, pp. 7-40, publishes a revision of the *Lacertilia*, minus the genus *Liolæmus*.

#### PAPUASIA AND AUSTRALIA.

Ceram, Amboyna, Barda, and New Guinea: O. BETTGER, Ber. Offenb. Ver. xxix-xxxii, 1892, annotated list of Reptiles and Batrachians.

British New Guinea: C. W. DE VIS, Ann. Queensland Mus. i, 1892, No. 2, pp. 11 & 12, enumerates some Reptiles.

Australia: J. J. FLETCHER, P. Linn. Soc. N.S.W. (2) vii, 1892, pp. 7-19, publishes a third contribution to a more exact knowledge of the geographical distribution of the Batrachia.

Victoria: A. H. S. Lucas, P. R. Soc. Vict. (2) iv, 1892, pp. 59-64, has notes on the distribution of Victorian Batrachians.

#### B.-FOSSIL.

Rhone Basin: L. LORTET, Arch. Mus. Lyon, v, 1892, pp. 1-139, figs., pls. i-xii, monographs the fossil Reptiles and Batrachians. Critical remarks on some of the Lizards and Rhynchocephalians are made by G. A. BOULENGER, Ann. N. H. (6) xi, 1893, pp. 204-210.

United States: O. C. Marsh, Am. J. Sci. (3) xliii, 1892, pp. 449-453, figs., and xliv, 1892, pp. 171-176, pls. ii-v, describes various Reptilian remains from the Laramie formation.

Texas: E. D. Cope, P. Am. Phil. Soc. xx, 1892, pp. 123-131, describes fossil Reptiles.

Parana: J. B. Ambrosetti, Bol. Ac. Cordoba, x, 1890, pp. 409-426, describes fossil Oligocene Reptiles.

Queensland: R. L. JACK & R. ETHERIDGE, JUN., Geol. and Pal. Queensland and New Guinea (Brisbane and London: 1892, 8vo), pp. 648-653, enumerate the fossil Reptiles.

## IV.—REFERENCES, ARRANGED SYSTEMATICALLY.

# REPTILIA.

- H. G. SEELEY, dealing with the systematic position of Pariosaurus, expresses his views on the relations of the various groups of Reptilia. The Labyrinthodonts are referred to this class. Phil. Tr. clxxxiii (B), 1892, pp. 367 & 368.
- E. D. COPE, Tr. Am. Phil. Soc. xvii, 1892, pp. 11-26, pls. i-v, discusses the homologies of the posterior cranial arches in the Reptilia. The author admits as probable Baur's hypothesis, which supposes the bars to have been derived from a roofed skull by perforation. The elements usually termed quadratojugal, squamoral, and supratemporal, are respectively named zygomatic, supratemporal, and supramastoid. The paper concludes with systematic considerations which do not differ materially from the views expressed by the same author in 1885.
- J. W. HULKE, P. R. Soc. lii, 1892, pp. 233-255, figs., discusses the homology of the various elements in the Reptilian shoulder-girdle.
- A. MEYER, Z. wiss. Zool. lv, 1892, pp. 63-133, pls. iv & v, writes on the brain of Reptiles.
- E. BERANEK, Anat. Anz. vii, 1892, pp. 674-689, figs., writes on the pineal eye and its nerve.
- A. DE KLINCKOWSTRÖM, Anat. Anz. viii, 1893, pp. 289-299, figs., studies the early development of the pineal eye, the epiphysis, and the parietal nerve in *Iguana*.

- F. MAURER, Morph. JB. xviii, 1892, pp. 739-747, remarks on the formation and structure of scales in the Reptiles.
- C. Röse, Anat. Anz. vii, 1892, pp. 748-758, figs., writes on the development of teeth and egg-tooth in Reptiles.
- C. K. HOFFMANN, Verh. Ak. Amst. xxix, 1892, pp. 1-19, pls. i & ii, writes on the development of the venous system in Reptiles.
- F. HOCHSTETTER, Morph. JB. xix, 1892, pp. 428-501, pls. xv-xvii, contributes to the knowledge of the development of the venous system in Reptiles, his researches being chiefly based on Lucerta and Tropidonotus.
- E. GIACOMINI has preliminary notes on the fætal membranes of Reptiles; Mon. Zool. Ital. iii, 1892, pp. 126-128, 157-164, & 185-196.
- H. Virchow, Z. wiss. Zool. liii (Suppl.), 1892, pp. 170-206, pl. x, deals specially with the vitelline sac of Reptiles.
- A. OPPEL, Arch. mikr. Anat. xxxix, 1892, pp. 215-290, pls. ix-xii, deals with the fecundation of the egg in Anguis fragilis and Tropidonotus natrix.
- C. Benda, Verh. Anat. Ges. 1892, pp. 195-199, writes on the histiogenesis of the Sauropidan spermatozoon.
- F. WERNER, Biol. Centralbl. xii, 1892, pp. 272-277, offers further remarks on the markings of Reptiles.
- F. WERNER, Zool. Gart. xxxiii, 1892, pp. 367-372, has biological notes on various Reptiles.

# SQUAMATA.

#### LACERTILIA.

- E. D. COPE, P. Am. Phil. Soc. xxx, 1892, pp. 185-221, pls. ii-vi, writes on the osteology of the Lacertilia, and J. Morph. vii, 1892, pp. 223-244, pl. xiii, on degenerative types of scapular and pelvic arches in the Lacertilia.
- F. SIEBENROCK, Ann. Hofmuseum Wien, vii, 1892, pp. 163-196, pls. i & ii, makes a comparative study of the skulls in the principal genera of Scincidæ, Anguidæ, and Gerrhosauridæ. The same author, t. c. pp. 373-378, writes on the sacrum of Lacertilia.
- R. v. Seiller, Leuckart Festschrift, 1892, pp. 250-258, pls. xxv & xxvi, writes on the lingual glands of Lacerta.
- G. CHIARUGI, Monit. Zool. Ital. iii, 1892, pp. 211 & 212, and Arch. Ital. Biol. xviii, 1893, pp. 363 & 364, has a preliminary note on the development of the olfactory nerve in Lacerta muralis.
- L. Will, Morph. JB., Anat. vi, 1892, pp. 1-160, figs., pls. i-xi, deals with the embryology of Tarentola mauritanica, L.
- P. MINGAZZINI, Atti Acc. Lincei (5) i, 1892, Rend. pp. 41-45, has a note on the gestation in Chalcides tridactylus.
- STRAHL, Verh. Anat. Ges. 1892, pp. 190-195, remarks on retrogression of ripe ovarian eggs in Lacerta agilis.

#### GECKONIDÆ.

- Teratoscincus scincus, Schleg.: note on Transcaspian specimens; Bou-LENGER, P. Z. S. 1891, p. 629.
- Gymnodactylus caspius, Eichw., and G. fedtschenkoi, Strauch: notes by Boulenger, t. c. pp. 629 & 630. G. louisiadensis, n. sp., De Vis, Ann. Queensland Mus. i, 1892, No. 2, p. 11, Sudest I., British New Guinea. G. sphyrurus, p. 6, New South Wales, and cornutus, p. 8, N. E. Queensland, n. spp., Douglas-Ogilby, Rec. Austral. Mus. ii, 1892.
- Gonatodes collaris, n. sp., GARMAN, Bull. Ess. Inst. xxiv, 1892, p. 83, Chatham I., Galapagos.
- Diplodactylus intermedius, n. sp., Douglas-Ogilby, t. c. p. 10, Interior of New South Wales.
- Phyllodactylus galapagensis, Ptrs.: notes by GARMAN, Bull. Ess. Inst. xxiv, 1892, p. 81. P. baurii, n. sp., id. t. c. p. 82, Charles I., Galapagos.
- Ptyodactylus lacazii, n. sp., Boutan, Arch. Z. expér. (2) x, 1892, p. 17, N. W. Arabia [= P. lobatus, L.—Rec.].
- Hemidactylus mabouia, Mor.: on variations in the femoral pores; BOCAGE, J. Sci. Lisb. (2) ii, 1892, p. 221: n. var molleri; BEDRIAGA, O Istituto (Coimbra), xxxix, 1892 (p. 19 of reprint), S. Thomé, Gulf of Guinea. H. greeffii, Bocage: remarks, id. ibid. p. 16.
- Lygodactylus conradti, n. sp., MATSCHIE, SB. nat. Fr. 1892, p. 109, Usambara, German E. Africa. L. angularis, n. sp., Günther, P. Z. S. 1892, p. 555, pl. xxxiii, fig. 1 (1893), Nyassaland.
- Ptychozoon horsfieldii, Gray, described and figured by F. Müller, Festschr. Nat. Ges. Basel, 1892, p. 209, pl. iv.

#### AGAMIDÆ.

- Draco modiglianii, n. sp., VINCIGUERRA, Ann. Mus. Genov. (2) xii, 1892, p. 523, Engano I.
- Gonyocephalus beyschlagi, n. sp., BETTGER, Ber. Offenb. Ver. xxix-xxxii, 1892, p. 104, N. E. Sumatra.

#### IGUANIDÆ.

- Iguana tuberculata, L.: on a specimen with reproduced, striped tail; BOULENGER, P. Z. S. 1891, p. 466.
- Conolophus subcristatus, Gray, and Amblyrhynchus cristatus, Bell: notes by Garman, Bull. Ess. Inst. xxiv, 1892, pp. 77 & 79. The same author, Science, xx, 1892, p. 16, fig., remarks on the dorsal crest of Conolophus subcristatus, each of the spines of which resembles externally the rattle of a small rattlesnake.

- Tropidurus, Wgm.: G. BAUR, Leuckart Festschrift, 1892, pp. 259-277, figs., describes the variations of the Lizards of this genus in the Galapagos Is. T. barringtonensis, p. 268, and jacobii, p. 269, n. spp., id. ibid. T. bocourtii, Blgr., = T. occipitalis, Ptrs.; GARMAN, Bull. Ess. Inst. xxiv, 1892, p. 94.
- Phrynosoma, Wgm.: on the ejection of blood from the eyes; O. P. HAY, P. U. S. Nat. Mus. xv, 1892, p. 375; A. L. HERRERA, Le Nat. xiv, 1892, p. 114.
- Iguanavus teres, n. sp. (foss.), MARSH, Am. J. Sci. (3) xliii, 1892, p. 451, Laramie formation, Wyoming.

#### ANGUIDÆ.

- Ophisaurus apus, Pall.: notes on habits; H. WERNER, Zool. Gart. xxxiii, 1892, p. 38.
- Anguis fragilis, L.: on a dicephalous monster; Mazza, Atti Soc. Ligust. iii, 1892, p. 256, pl. ii.

#### VARANIDÆ.

Varanus griseus, Daud.: on its habits in confinement; H. WERNER, Zool. Gart. xxxiii, 1892, pp. 304 & 374. V. heteropholis, n. sp., Boulenger, P. Z. S. 1892, p. 506, pl. xxix, Mt. Dulit, Borneo.

#### TEHDÆ.

- Ameira pleii, n. var. brachiosquamatus, Cope, in Verrill, Tr. Conn. Ac. viii, 1892, p. 352, Dominica, W. Indies [=? A. fuscata, Garm.—Rec.]
- Tiaporus, n. g., like Ameiva, but without femoral pores. T. fuliginosus, n. sp., Cope, P. Am. Phil. Soc. xx, 1892, p. 132, pl. —, Swan I., off N. E. Coast of Honduras.
- Cnemidophorus, Wagl.: synopsis of the species, by Cope, Tr. Am. Phil. Soc. xvii, 1892, p. 27, pls. vi-xiii. C. tessellatus, rubidus, and multiscutatus, pp. 36 & 37, Lower California, C. gularis scalaris, p. 47, Chihuahua, N. Mexico, sericeus, p. 48, Texas, and semifasciatus, p. 49, Chihuahua, n. subspp., id. ibid. C. variolosus, p. 39, Coahuila, and septemvittatus, p. 40, California, n. spp., id. ibid. C. lentiginosus, n. sp., Garman, Bull. Ess. Inst. xxiv, 1892, p. 92, Guayaquil, Ecuador.
- Propodinema, n. g. for Podinema paranensis, Scalabrini (foss.), Ambrosetti, Bol. Ac. Cordoba, x, 1890, p. 410. P. oligocena, n. sp. (foss.), id. t.c. p. 413, Oligocene, Parana.
- Chamops, n. g. for a Lacertilian maxillary. C. segnis, n. sp. (foss.), MARSH, Am. J. Sci. (3) xliii, 1892, p. 450, fig., Laramie formation of Wyoming [apparently nearest to the Teiidæ.—Rec.].
- Bachia, Gray, 1845 [not 1849, as stated] should supersede Cophias, Fitz., 1826, nec Merr., 1820; GARMAN, Bull. Ess. Inst. xxiv, 1892, p. 96.

#### LACERTIDE.

- Lacerta muralis, Laur.: notes on Italian specimens; Positano-Spada, Boll. Soc. Rom. i, 1892, p. 89: n. var. multifasciata, id. t. c. p. 154, fig., Italy. L. muralis, vars. depressa, Camer., and difilippii, Camer.: notes by Bœttger, Ber. Senck. Ges. 1892, pp. 140 & 144: n. vars. raddei, p. 142, and ralentini, p. 145, Caucasus. L. mosorensis, Kolomb. (= koritana, Tom.), redescribed and figured by Steindachener, Ann. Hofmuseum Wien, vii, 1892, p. 235, pl. xvi.
- Acanthodactylus vulgaris, D. & B.: remarks on Portuguese specimens; FERREIRA, J. Sci. Lisb. (2) ii, 1892, p. 188.
- Holaspis guentheri, Gray, recorded from East Africa by MATSCHIE, SB. nat. Fr. 1892, p. 110.

#### SCINCIDE.

- E. D. Cope, J. Morph. vii, 1892, p. 233, gives a synopsis of the genera of Scincida, in which the following new generic names are introduced:— Monophyaspis, Oncopus, Ollochirus, Tridentulus, Furcillus, Haploscincus, Dimeropus, Dicloniscus, Mesomycterus.
- Lygosoma africanum, Gray, redescribed by Bedriaga, O Istituto (Coimbra), xxxix, 1892 (p. 23 of reprint). L. travancoricum, Bedd., n. var., palnica; Bettger, Ber. Offenb. Ver. xxix-xxxii, 1892, p. 72, Pulney Hills, S. India. L. (Siaphos) relictum, n. sp., Vinciguerra, Ann. Mus. Genov. (2) xii, 1892, p. 524, Engano I.
- Mocoa caruleocauda, n. sp., DE Vis, Ann. Queensland Mus. i, 1892, No. 2, p. 12, Sudest I., British New Guinea.
- Eumeces scutatus, Theob.: note on Transcaspian specimens; Boulenger, P. Z. S. 1891, p. 630.
- Ophiomorus brevipes, Blanf.: note on Transcaspian specimens; Bou-LENGER, t. c. p. 631.
- Chalcides simonyi, Stdr., is regarded as a variety of C. viridanus, Gravh., by Boulenger, Ann. N. H. (6) ix, 1892, p. 74: note by Bedriaga, C.R. Congrès Internat. Zool. i, 1892, p. 242. C. boulengeri, n. sp., Anderson, P. Z. S. 1892, p. 17, pl. i, fig. 1, Duirat, Tunisia.

#### INCERTA SEDIS.

Euposaurus, n. g. (Jourdan MS.). for small fossil Lizards, which are referred to the Rhynchocephalians; LORTET, Arch. Mus. Lyon, v, 1892, p. 73. These fossils are regarded by the Recorder as Lacertilian; Ann. N. H. (6) xi, 1893, p. 208. E. thiollieri, p. 73, pl. vi, figs. 7-10, and cerinensis, p. 75, figs., n. spp., LORTET, t. c., Jurassic, Cerin, France.

#### RHIPTCGLOSSA.

- G. A. BOULENGER, Ann. N. H. (6) ix, 1892, p. 72, remarks on some newly-discovered E. African Chameleons, regarding Chameleon höhnelii, Stdr., and C. leikipiensis, Stdr., as identical with C. bitæniatus, Fisch., and C. abbotti, Stejn., as identical with C. tuvetensis, Stdr. The female of the latter species is described.
- Chamaleon fischeri, Reich.: note by MATSCHIE, SB. nat. Fr. 1892, p. 102.

  C. deremensis, p. 103, spinosus, p. 105, and tenuis, p. 106, n. spp., id. ibid., Usambara, German E. Africa. C. isabellinus, n. sp., GÜNTHER, P. Z. S. 1892, p. 556, pl. xxxiii, fig. 2 (1893), Nyassaland [ = C. dilepis, Leach.—Rec.] C. (Brookesia) brevicaudatus, p. 107, and temporalis, p. 108, n. spp., MATSCHIE, t. c., Usambars, German E. Africa.
- Rhampholeon platyceps, p. 556, pl. xxxiv, fig. 1, and brachyurus, p. 557, pl. xxxiv, fig. 2, n. spp., Günther, t. c. Nyassaland.

#### DOLICHOSAURIA.

- C. GORJANOVIĆ-KRAMBERGER, Glasnik Nadar. Drustva, vii, 1892, pp. 74-106, pls. iii & iv, and Rad jugoslav. akad. cix, 1892, pp. 96-123, pls. i & ii, discusses the relations of this group of Reptiles, for which he proposes the name Ophiosauria.
- Aigialosaurus, n. g. for A. dalmaticus, n. sp. (foss.), C. Gorjanović-Kramberger, ibid., Chalk of Lesina I., Dalmatia.
- Pontosaurus, n. g. for Hydrosaurus lesinensis, Kornh. (foss.) C. GORJANOVIĆ-KRAMBERGER, ibid.
- Remarks by Boulenger, Ann. N. H. (6) xi, 1893, pp. 204-207.

#### PYTHONOMORPHA.

- G. BAUR, J. Morph. vii, 1892, pp. 1-22, reviews the history of our know-ledge of the Mosasauroid skull, and discusses the systematic position of the Mosasauridæ.
- A. GAUDRY, Mém. Soc. Géol. iii, No. 3, 1892, 13 pp., 2 pls., describes the remains of Mosasaurs hitherto found in France, which are referred to Liodon anceps, Ow., L. compressidens, Gaudry, L. mosasauroides, Gaudry, and Mosasaurus, sp.
- Liodon mosasauroides and compressidens, n. spp., GAUDRY, CR. cxv, 1892, p. 304, Chalk, Pau and Sens, France.
- Clidastes velox, Marsh: description of a nearly complete specimen, with figures of the pectoral and pelvic arches; Williston & Case, Kansas Quart. i, 1892, p. 17, pls. ii & iii. C. westii, n. sp., iiid., t. c. p. 29, pls. iv & v, Cretaceous of Kansas.
- Platecarpus coryphæus, Cope: on the skull; BAUR, J. Morph. vii, 1892, p. 1, pls. i & ii.

#### OPHIDIA.

BOULENGER, G. A. Catalogue of the Snakes in the British Museum (Natural History). Vol. i, containing the families Typhlopidæ, Glauconiidæ, Boidæ, Ilysiidæ, Uropeltidæ, Xenopeltidæ, and Colubridæ Aglyphæ, part. London: 1893, 8vo, xiii, 448 pp., 28 pls.

The classification adopted is the same as proposed by the Recorder in 1890. This volume contains descriptions of 523 species, and the enumeration of 3324 specimens.

- G. B. BUTLER, P. Z. S. 1892, pp. 477-498, pl. xxviii, deals with the subdivision of the body-cavity in Snakes.
- F. LATASTE, Act. Soc. Sci. Chili, ii, 1893, pp. clxvii-clxx, replies to some criticisms of his own work made by the author of the above paper.
- F. WERNER, Biol. Centralbl. xii, 1892, pp. 277-280, remarks on the sculpture of the epidermis in Snakes.
- S. GARMAN, Science, xx, 1892, p. 16, figs., has further remarks on the origin and evolution of the Ophidian "Rattle."
- F. NIEMANN, Arch. f. Nat. lviii, i, 1892, pp. 262-286, pl. xiv, deals with the morphology and physiology of the labial and poison glands of Snakes.
- J. J. QUELCH writes on "Venom in Harmless Snakes," and refers in particular to the effect on himself of a bite from the opisthoglyphous Erythrolamprus venustissimus. Zool. (3) xvii, 1893, p. 30.
- C. J. Martin, P. Linn. Soc. N.S.W. (2) vii, 1892, pp. 319-322, and J. R. Soc. N.S.W. xxvi, 1892, pp. 240-264, makes observations on the poisonous constituents of the venom of Pseudechis porphyriacus.
- L. VAILLANT records observations made in the Garden of Plants on the feeding of Snakes. N. Arch. Mus. (3) iv, 1892, pp. 221-234. Abstract in CR. cxv, 1892, pp. 277-279.
- O. P. Hay, P. U. S. Nat. Mus. xv, 1892, pp. 385-397, has notes on the breeding habits, eggs, and young of various North American Snakes.
- G. A. BOULENGER, Nat. Sci. i, 1892, pp. 44-49, figs., writes on Marine Snakes.

#### TYPHLOPIDÆ.

Typhlops curtus, n. sp., DOUGLAS-OGILBY, Rec. Austral. Mus. ii, 1892, p. 23, Gulf of Carpentaria [= T. ligatus, Ptrs.—Rec.] T. boettgeri, p. 39, pl. ii, fig. 6, Madagascar, and crossii, p. 52, pl. iii, fig. 5, Niger, n. spp., BOULENGER, Cat. Sn. i, 1893.

#### GLAUCONIIDÆ.

Glauconia distanti, n. sp., Boulenger, in Distant's Naturalist in the Transvaal, p. 175, fig., Transvaal.

#### Boidæ.

Eryx, Daud., and Gongylophis, Wagl.: A synopsis of the species is given by Boulenger, Ann. N. H. (6) ix, 1892, p. 75. Gongylophis muelleri, n. sp., id., ibid. p. 74, Sennar. E. jaculus, L.: Note on Transcaspian specimens; Boulenger, P. Z. S. 1891, p. 631.

#### UROPELTIDÆ.

Rhinophis travancoricus, n. sp., BOULENGER, Cat. Sn. i, p. 143, pl. ix, fig. 3, 1893, and J. Bomb. N. H. Soc. vii, 1892, p. 318, Travancore.

#### COLUBRIDÆ.

# Aglyphæ.

- Dromicodryas, n. g. for Herpetodryas bernieri, D. & B., BOULENGER, Cat. Sn. i, p. 189, 1893.
- Eutania elegans brunnea and lineolata, n. subspp., Cope, P. U. S. Nat. Mus. xiv, 1892, pp. 654 & 655, California. E. infernalis vidua, n. subsp., id., ibid. p. 658, California. E. sirtalis semifasciata, p. 662, Illinois and Wisconsin, and trilineata, p. 664, Oregon and Montana, n. subspp., id., ibid. E. brachystoma, n. sp., Cope, Am. Nat. xxvi, 1892, p. 964, fig., W. Pennsylvania [= E. leptocephala, B. & G.— Rec.] E. aurata, n. sp., Cope, P. U. S. Nat. Mus. xiv, 1892, p. 659, New Mexico. E. pracocularis, n. sp., Bocourt, Le Nat. xiv, 1892, p. 278, Belize, Honduras.
- Tropidonotus infernalis, Blainv.: On variations in the labial shields; Bocourt, Bull. Soc. Z. Fr. xvii, 1892, p. 40. T. natrix, L.: On the occurrence of the striped variety near Nantes, Brittany; H. & T. PIEL DE CHURCHEVILLE, Bull. Soc. Ouest France, ii, 1892, p. 35, pl. ii. T. conspicillatus, Gthr.: redescribed, from Sumatra, by Bœttger, Ber. Offenb. Ver. xxix-xxxii, 1892, p. 109. T. tessellatus, var. flavescens, Werner, an albino; Werner, Verh. z.-b. Wien, xlii, 1892, p. 355. T. obliquus, Hallow., and transversus, Hallow.: notes by Garman, Bull. Ess. Inst. xxiv, 1892, p. 103. T. petersii, n. sp., Boulenger, Cat. Sn. i, p. 225, pl. xiv, fig. i, 1893, Borneo.
- Natrix fasciata pleuralis, n. subsp.; Cope, P. U. S. Nat. Mus. xiv, 1892, p. 672, Mississippi and Georgia.
- Macropophis, n. subg. for Tropidonotus hypometas, Gthr., and T. dendrophiops, Gthr.; BOULENGER, Cat. Sn. i, p. 264, 1893.
- Macropisthodon, n. g. for Tropidonotus flaviceps, D. & B., and allies;
  BOULENGER, t. c. p. 265.
- Storeria victa, n. sp.; O. P. HAY, Science, xix, 1892, p. 199, Florida.
- Rhabdops, n. g., for Ablabes olivaceus, Bedd., and A. bicolor, Gthr.; BOULENGER, Cat. Sn. i, p. 300, 1893.
- Plagiopholis, n. g., for P. blakewayi, n. sp.; Boulenger, t. c. p. 301, pl. xix, fig. 3, Shan States.

- Trirhinopholis, n. g., for T. nuchalis, n. sp.; BOULENGER, Cat. Sn. i, p. 419, pl. xxviii, fig. 1, Shan States.
- Tetralepis, n. g., allied to Ablabes, D. & B., but with the dentition of Homalosoma, Wagl., and only four upper labial shields; BETTGER, Ber. Offenb. Ver. xxix-xxxii, 1892, p. 124. T. fruhstorferi, n. sp.; id. t. c., p. 125, Tengger Mts., Java.
- Oxyrhabdium, n. n. for Stenognathus, D. & B., nec Chaud.; BOULENGER, Cat. Sn. i, p. 302.
- Lycognathophis, n. g., for Psammophis seychellensis, Schleg.; Boulenger, t. c. p. 317.
- Ablabophis, n. g., for Coronella rufula, Licht.; BOULENGER, t. c. p. 318.
- Lamprophis fuscus, n. sp., BOULENGER, t. c. p. 322, pl. xx, fig. 4, Cape of Good Hope.
- Gonionotophis, n. n. for Gonionotus, Mocq. nec Gray, Boulenger, t. c. p. 323.
- Gonionotus vossii, n. sp., BŒTTGER, Zool. Anz. xv, 1892, p. 418, Cameroons.
- Lycophidium meleagris, p. 337, pl. xxi, fig. 2, Angola, jacksoni, p. 340, pl. xxi, fig. 3, E. Africa, and abyssinicum, p. 342, pl. xxii, fig. 2, Abyssinia, n. spp., Boulenger, t. c.
- Lycodon striatus, Shaw, recorded from Transcaspia by BOULENGER, P.Z. S. 1891, p. 631. L. stormi, n. sp., BETTGER, Zool. Anz. xv, 1892, p. 418, Celebes.
- Stegonotus heterurus, n. sp., BOULENGER, Cat. Sn. i, p. 367, pl. xxv, fig. 1, Duke of York I. and New Britain. S. dumerilii, n. n. for Lycodon muelleri, D. & B.; id. t. c. p. 368.
- Zamenis, Wagl.: a synopsis of the species recorded from S. W. Asia;
  BOULENGER, P. Z. S. 1891, p. 632. Z. gemonensis, Laur.: on its distribution in France; R. MARTIN, Bull. Soc. Ouest France, ii, 1892, p. 149.
- Bascanium semilineatum, n. sp., Cope, P. U. S. Nat. Mus. xiv, 1892, p. 626, Arizona.
- Philophyllophis, Pantherophis, and Potamophis. These nomina nuda of Fitzinger are revived by GARMAN, Bull. Ess. Inst. xxiv, 1892, pp. 105 & 108, to replace Haldea, B. & G., Cyclophis, Gthr., and Scotophis, B. & G.
- Coluber enganensis, n. sp., VINCIGUERRA, Ann. Mus. Genov. (2) xii, 1892, p. 524, Engano I.
- Rhinechis scalaris, Mich.: on its habits in confinement; H. WERNER, Zool. Gart. xxxiii, 1892, p. 40.
- Dromicus dolicocercus, n. sp., Peracca, Boll. Mus. Zool. Tor. vii, 1892, No. 112, pl. i, fig. 1, Madagascar.
- Herpetodryas biserialis, Gthr., brunneus, Gthr., and reticulatus, Ptrs.: notes by Garman, Bull. Ess. Inst. xxiv, 1892, pp. 85, 90, & 91.
- Dendrophidium dendrophis, Schl., described and figured by Dugks, Nat. Mex. (2) ii, 1892, p. 100, pl. v.
- Dendrophis helena, n. sp., WERNER, Zool. Anz. xvi, 1893, p. 81, Ceylon.

- Simotes meyerinkii, Stdr., = S. octolineatus, Schn.; Boulenger, Ann. N. H. (6) ix, 1892, p. 74.
- Conopsis lineatus, Kenn.: on variations in the head-shields; Boccourt, Bull. Soc. Z. Fr. xvii, 1892, p. 41.
- Cyclophis modestus, Mart., n. var. punctutolineatu; BETTGER, Ber. Senck. Ges. 1892, p. 147, Armenia.
- Pseudocyclophis walteri, Bttgr.: note by Boulenger, P. Z. S. 1891, p. 632.
- Tropidocionium annulatum, n. sp., Bocourt, Le Nat. xiv, 1892, p. 132, Godines, Guatemala.
- Chilomeniscus stramineus fasciatus, n. subsp., Cope, P. U. S. Nat. Mus. xiv, 1892, p. 595, California.
- Idiopholis, n. g. Body cylindrical; head not distinct from neck; tail very short; three internassls and two præfrontals; no loreal, no præocular, no supraocular, no anterior temporal; nostril between two shields; eye small, with round pupil; two pairs of chin-shields, anterior in contact with symphysial; scales smooth; anal single; subcaudals double; maxillary teeth numerous, equal, very small. I. collaris, n. sp., Mocquard, Le Nat. xiv, 1892, p. 35, and Mém. Soc. Zool. v, 1892, p. 191, pl. vii, fig. 1, W. Borneo.
- Dasypeltis scalra, L.: on the manner in which it feeds on eggs; Sclater, P. Z. S. 1892, p. 476.

# Opisthoglyphæ.

- Coniophanes signatus, n. sp., GARMAN, Bull. Ess. Inst. xxiv, 1892, p. 91, Guayaquil, Ecuador.
- Tachymenis boulengeri, n. sp., PERACCA, Boll. Mus. Zool. Tor. vii, 1892, No. 112, pl. i, fig. 2, Madagascar.
- Psammophylax variabilis, n. sp., GÜNTHER, P. Z. S. 1892, p. 557, pl. xxxv (1893), Nyassaland.
- Psammophis longifrons, Blgr.: note by F. DRECKMANN, J. Bomb. Soc. vii, 1892, p. 406.
- Dipsas trigonata, Schn., recorded from Transcaspia by Boulenger, P. Z. S. 1891, p. 633.
- Leptodira annulata, L.: notes by GARMAN, Bull. Ess. Inst. xxiv, 1892, p. 90.
- Tantilla annulata, n. sp., BETTGER, Zool. Anz. xv, 1892, p. 419, Nicaragua.
- Gerardia prevostiana, Eyd. & Soul., recorded from Ceylon by HALY, J. A. S. (Ceylon), xi, 1892, p. 197.

# Proteroglyphæ.

Hoplocephalus suboccipitalis, n. sp., DOUGLAS-OGILBY, Rec. Austral. Mus. ii, 1892, p. 23, Moree, New South Wales.

- Dendraspis angusticeps, Smith, described from S. Thome by Bedriaga, O Istituto, No. 6, 1892. The same Snake is referred to D. jamesonii, Trail, by Bocage, J. Sci. Lisb. (2) ii, 1892, p. 265.
- Hydrus platurus, L.: notes by GARMAN, Bull. Ess. Inst. xxiv, 1892, p. 88.

### VIPERIDÆ.

- Causus rostratus, Gthr., redescribed as C. rostratus, n. sp., by MOCQUARD, Le Nat. xvi, 1892, p. 35; the error corrected, p. 64.
- Vipera, Laur.: J. DE BEDRIAGA, C.R. Congrès Internat. Zool. i, 1892, pp. 236-241, diagnoses the species of the Mediterranean district. V. berus, L.: on variations in British specimens; BOULENGER, Zool. (3) xvi, 1892, p. 87. A. B. MEYER & F. HELM, J. Ber. ornith. Beobacht. Stat. Sachsen, vi, pp. 55-63, have compiled interesting information on its occurrence and habits in Saxony. V. lebetina, L., u. var. deserti; Anderson, P. Z. S. 1892, p. 20, pl. i, fig. 3, Duirat, Tunisia.
- A. E. FECKTISTOW'S paper on the physiology of the rattle of *Crotalus durissus* is translated in Ann. N. H. (6) xi, 1893, pp. 54-58.
- Crotalus adamanteus ruber, n. subsp., Cope, P. U. S. Nat. Mus. xiv, 1892, p. 690, N. America.
- Trimeresurus xanthomelus, Gthr. (= jerdoni, Gthr.), figured by GÜNTHER, in Pratt's Snows of Tibet, pl. i, fig. A. T. okinavensis, n. sp., Boulenger, Ann. N. H. (6) x, 1892, p. 302, Great Loo Choo I.
- Provipera bættgeri, n. g. & sp., for a fossil poison-tooth from the Lower Miocene of Morbach-Biebrich, Germany; Kinkelin, Zool. Anz. xv, 1892, p. 93. Remarks by Cope, t. c. p. 224.

### INCERTÆ SEDIS.

Coniophis, n. g., for Ophidian vertebræ, the affinities of which are still unsettled. C. precedens, n. sp. (foss.), Marsh, Am. J. Sci. (3) xliii, 1892, p. 450, fig., Ceratops beds (Cretaceous) of Wyoming.

# ORNITHOSAURIA.

- S. W. WILLISTON, Kansas Quart. i, 1892, pp. 1-13, pl. i, describes remains of Pteranodon, Marsh, and Nyctodactylus, Marsh, from Kansas, and proposes to divide the Order into three families, viz.: Pterodactylidæ (with subfamilies Pteranodontinæ and Pterodactylinæ), Rhamphorhynchidæ, and Ornithochiridæ.
- Pterodactylus cerinensis, H. v. Mey., described and figured by LORTET, Arch. Mus. Lyon, v, 1892, p. 129, fig., pl. v, figs. 3 & 4.

# DINOSAURIA.

- H. G. SEELEY, Q. J. Geol. Soc. xlviii, 1892, Proc. pp. 108-191, discusses the relations of the Saurischian Dinosaurs.
- C. Marsh, Am. J. Sci. (3) xliii, 1892, pp. 81-83, pls. ii & iii, has remarks on the skull of the Ceratopsidα, and a restoration of the skull of Torosaurus latus, Marsh.
- B. LYDEKKER, Q.J. Geol. Soc. xlviii, 1892, p. 375, fig., has a note on some Sauropodous Dinosaurian foot-bones from the Wealden.
- R. Burckhardt, JB. Mineral. 1892, ii, pp. 71 & 72, fig., criticizes Marsh's interpretation of the brain of Triceratops flabellatus.
- Iguanodon hillii, n. sp., for a tooth from the Lower Chalk near Hitchin, England; E. T. Newton, Geol. Mag. (3) ix, 1892, p. 49.
- Claosaurus annectens, n. sp., MARSH, Am. J. Sci. (3) xliii, 1892, p. 453, & xliv, 1892, p. 171, pl. ii, and restoration, p. 343, pl. vi, Laramie formation, Wyoming. On its skull and brain; id. op. cit. xlv, 1893, p. 83, pls. iv & v.
- Polacanthus, Hulke: on the pelvis; Seeley, Q. J. Geol. Soc. xlviii, 1892, p. 81, figs.; Lydekker, t. c. p. 148, figs.
- Palæoscincus costatus, Leidy, is regarded as a Stegosaurian tooth by Marsh, Am. J. Sci. (3) xliv, 1892, p. 173. P. latus, n. sp., id. ibid. pl. iii, fig. 2, Ceratops beds, Wyoming.
- Stegosaurus ungulatus, Marsh: tooth figured; MARSH, ibid. pl. iv, fig. 1.
  Lalaps incrassatus, Cope: description of the skull; Cope, P. Am. Phil.
  Soc. xxx, 1892, p. 240.
- Ceratosaurus nusicornis, Marsh: restoration by MARSH, Am. J. Sci. (3) xliv, 1892, p. 347, pl. vii.
- Amhisaurus colurus, Marsh: further notes and figures by MARSH, Am. J. Sci. (3) xliii, 1892, p. 543, pls. xv & xvi, & xlv, 1893, p. 169, pl. vi. A. solus, n. sp., id. op. cit. xliii, p. 545, Trias, Connecticut River.
- Ornithonimus, Marsh, should probably be referred to the Theropoda;
  MARSH, Am. J. Sci. (3) xliii, 1892, p. 451. O. sedens, p. 451, and minutus, p. 452, n. spp., id. ibid., Ceratops beds, Wyoming.
- Aublysodon, Leidy: remarks by MARSH, Am. J. Sci. (3) xliv, 1892, p. 174.
  A. amplus, p. 174, and cristatus, p. 175, n. spp., MARSH, t. c., Laramie formation, North America

# EMYDOSAURIA.

- W. KÜKENTHAL, Morph. JB. xix, 1892, pp. 42-55, pl. ii, describes the skeleton of the manus in young Crocodiles, and draws special attention to the hyperphalangy in the fourth and fifth fingers, which shows that the existing Crocodilians are derived from forms more thoroughly adapted to aquatic life.
- C. ROESE, Verh. Anat. Ges. 1892, pp. 225-227, remarks on the development of the teeth in Crocodiles.

- C. P. SLUITER, Anat. Anz. vii, 1892, pp. 540-545, figs., writes on Jacobson's organ in Crocodilus porosus.
- Tomistoma, S. Müll.: on fossil remains from the Miocene of Nurri, Italy; LOVISATO, Atti Acc. Pontif. Lincei (5) i, 1892, Rend. p. 436.
- Rhamphostoma neogea, Burm. (foss.): note by Ambrosetti, Bol. Ac. Cordobs, x, 1890, p. 414.
- Leptorrhamphus, n. g., near Gavialis, for L. entrerrianus, n. sp. (foss.);
  Ambrosetti, t. c. p. 417, Oligocene, Parana.
- Oxysdonsaurus, n. g., near Gavialis, for O. striatus, n. sp. (foss.); AMBRO-SETTI, t. c. p. 419, Oligocene, Parana.
- Proalligator, n. g. for Crocodilus australis, Bravard (foss.); Ambrosetti, t. c. p. 420.
- Atoposaurus jourdani, H. v. Mey. (foss.), described and figured by LORTET, Arch. Mus. Lyon, v, 1892, p. 92, figs., pl. xi, figs. 9-16.
- Alligatorellus beaumonti, Gerv. (foss.), described and figured; LORTET, t. c. p. 99, pls. ix bis. & xi, figs. 1-8.
- Alligatorium meyeri, Gerv. (foss.), described and figured; LORTET, t. c. p. 108, pl. x.
- Gavialinum, n.g. (Gavialidæ?) for G. rhodani, n. sp. (foss.); LORTET, t. c. p. 122, pl. viii, Bathonian, Villebois, France.
- Crocodileimus robustus, Gerv. (foss.), described and figured by LORTET, t. c. p. 114, pl. ix.

# CHELONIA.

- G. BAUR, Anat. Anz. vii, 1892, pp. 206-211, figs., has further remarks on the carpus of Chelonians.
- E. BATAILLON, Ann. Soc. L. Lyon (2) xxxix, 1892, pp. 81-86, pl. —, has a contribution to the study of the arterial circulation in the land Tortoise.
- L. WILL, Biol. Centralbl. xii, 1892, pp. 182-192, figs., writes on gastrulation in the Chelonians.
- K. MITSUKURI'S second and third contributions to the embryology of Reptilia, J. Coll. Sci. Japan, iv, 1891, pp. 1-53, pls. i-x, & v, 1892, pp. 35-52, pls. ii-iv, deals with the foetal membranes of Chelonia, the researches being based on Clemmys japonica and Trionyx japonicus.
- Trionyx melitensis, Lyd. (foss.), == ? T. ragusensis, De Greg. 1883; De Gregorio, Nat. Sicil. xi, 1892, p. 182. T. leucopotamicus, n. sp. (foss.), Cope, Contr. Canad. Pal. iii, 1891, p. 5, pl. i, figs. 8 & 9, Lower Miocene, Canada. T. gemmellaroi, p. 11, pls. i & v, figs. 2-5, capellinii, p. 26, pls. ii & iii, and affinis, p. 47, pl. v, fig. 1, n. spp. (foss.); Negri, Mem. Soc. Ital. Sci. Nat. (3) viii, 1892, No. 7, Eocene of Monte Bolca.
- Miolania owenii, A. S. Woodw. (foss.): on some remains from New S. Wales; R. ETHERIDGE, Jun, Rec. Austral. Mus. ii, 1893, p. 39.

- L. LORTET, Arch. Mus. Lyon, v, 1892, p. 5, proposes a new family, "Halmyrachelides," for Idiochelys, H. v. Mey., and Hydropelta, H. v. Mey.
- Idiochelys fitzingeri, H. v. Mey. (foss.), described and figured by LORTET, t. c. p. 7, pl. i, & pl. ii, figs. 1 & 2.
- Hydropelta meyeri, H. v. Mey. (foss.), described and figured by LORTET, t. c. p. 18, pl. iii, figs. 3-5.
- Eurysternum crassipes, Wagn. (foss.), described and figured; LORTET, t. c. p. 24, pl. ii, figs. 4-7, and pl. ii bis.
- G. BAUR, Zool. Anz. xv, 1892, pp. 155-159, remarks on various genera of *Testudinidæ*, and insists on the distinctive characters of the families *Emydidæ* and *Testudinidæ*.
- Testudo graca, L.: on its habits; G. Rumpf, Zool. Gart. xxxiii, 1892, p. 260; A. GIRTANNER, t. c. p. 349. T. turgida, n. sp. (foss.), Cope, P. Am. Phil. Soc. xx, 1892, p. 27, Upper Cenozoic of Texas.
- Emys, Dum.: G. BAUR, op. cit. xxx, 1892, pp. 40 & 245, discusses which species should be regarded as the type of this genus.
- Cistudo cinosternoides, Gray: notes by GARMAN, Bull. Ess. Inst. xxiv, 1892, p. 99.
- Cinosternum flavescens, Ag.: note by Cope, P. Ac. Philad. 1892, p. 333.

## ICHTHYOSAURIA.

- E. Fraas, JB. Mineral. 1892, ii, pp. 87-90, figs., describes and figures a specimen of *Ichthyosaurus quadriscissus*, Qu., showing the contours of the soft parts, including dorsal and caudal fins.
- A propos of Fraas's discovery, L. Dollo, P.-v. Soc. Belg. Géol. vi, 1892, pp. 167-174, figs., discusses the origin of the dorsal fin in these Reptiles.
- R. LYDEKKER, Nat. Sci. i, 1892, pp. 514-521, reviews the various accessions to our knowledge of the *Ichthyosauria*.
- J. W. HULKE, P. R. Soc. lii, 1892, pp. 233-255, has remarks on the Ichthyosaurian shoulder-girdle.
- W. KÜKENTHAL, JB. Mineral. 1892, i, pp. 161-166, remarks on the hyperphalangy of Ichthyosaurs as compared to that of Cetaceans.
- Ichthyosaurus numismalis, n. sp., E. FRAAS, JH. Ver. Württ. xlviii, 1892, p. 22, pl. i, Lias, Württemberg. I. burgundiæ, Gaudry, described and figured by GAUDRY, Bull. Soc. Autun, v, 1892, p. 365, pl. xv.

# PLESIOSAURIA.

H. G. SEELEY, P. R. Soc. li, 1892, pp. 119-151, figs., deals with the nature of the shoulder-girdle and clavicular arch in the *Plesiosauria*, which he divides into two groups, termed *Dicranopleura* and *Cercidopleura*.

1892. [vol. xxix.]

- The subject further discussed by J. W. Hulke, P. R. Soc. lii, 1892, pp. 233-255, figs.
- H. G. SEELEY, Q. J. Geol. Soc. xlviii, 1892, pp. 586-604, discusses the characters and systematic position of the Mesosauria, which suborder he divides into two groups: Proganosauria and Neusticosauria.
- Mesosaurus tenuidens, Gerv., described and figured; SEELEY, t. c. p. 591, pl. xviii, fig. 5. M. pleurogaster, n. sp., id. t. c. p. 587, pl. xviii, figs. 1-4, Kimberley, S. Africa.
- Eunotosaurus, n. g., probably referable to the Mesosauria. E. africanus, n. sp., Seeley, t. c. p. 583, figs., Beaufort West, S. Africa.
- Parthanosaurus, n.g., allied to Nothosaurus, for P. zittelii, n. sp., Skuphos, Zool. Anz. xvi, 1893, p. 67, Trias of Vorarlberg.

# THECODONTIA.

Belodon, H. v. Mey., and Aëtosaurus, Fraas, are regarded by SEELEY, Q. J. Geol. Soc. xlviii, 1892, Proc., p. 189, as primitive Cetiosaurians. Episcoposaurus haplocerus, n. sp., Cope, P. Am. Phil. Soc. xx, 1892, p. 129, Trias of Texas.

# ACROSAURIA.

Pleurosaurus goldfussi, H. v. Mey. (foss.), described and figured by LORTET, Arch. Mus. Lyon, v, 1892, p. 81, figs., pl. vii. The Recorder points out, Ann. N. H. (6) xi, 1893, p. 208, that this Lizard should be regarded as the type of a distinct order of Reptiles, combining characters of the Proterosaurian Rhynchocephalia and Squamata, for which the name Acrosauria, H. v. Meyer, 1860, may be used.

# RHYNCHOCEPHALIA.

- L. Dollo, Bull. Soc. Belg. Géol. v, Mém. 1892, pp. 151-199, pls. vi-viii, completes his description of Champsosaurus, and discusses its position among the Rhynchocephalians, which he pronounces to be near the Proterosauridæ.
- G. A. BOULENGER gives a revised scheme of the classification proposed by him in 1891; the *Champsosauridæ* are transferred to the suborder *Proterosauria*, and the *Sauranodontidæ* are added to the suborder *Rhynchocephalia vera*; Ann. N. H. (6) xi, 1893, p. 209.
- Diopeus, Cope (foss.), is regarded as a Rhynchocephalian, somewhat allied to Palæohatteria, Credn.; Cope, Tr. Am. Phil. Soc. xvii, 1892, p. 15.
- Homoeosaurus jourdani, p. 69, pl. vi, figs. 1-5, and rhodani, p. 72, pl. iv, fig. 6, n. spp. (foss.), Lortet, Arch. Mus. Lyon, v, 1892, Jurassic, Corin, France.

Sauranodon incisivus, Jourdan (foss.), [= Saphæosaurus thollierii, H. v. Mey.], described and figured by Lortet, Arch. Mus. Lyon, v, p. 30, figs., pls. iiter, iii, iv, & v. A new family, Sauranodontidæ, is established by Lortet for its reception.

## ANOMODONTIA.

- H. G. SEELEY, Phil. Tr. clxxxiii (B), 1892, pp. 311-370, pls. xvii-xxiii, describes remains of Pariosaurus and Procolophon, at the same time discussing the classification of the Anomodontia, which are now divided into six groups.
- G. B. Howes remarks on the shoulder-girdle of Keirognathus, Seeley, and other Dicynodonts; J. Anat. Phys. xxvi, 1892, pp. 403-405.
- Gordonia, n. g., allied to Dicynodon, Ow., for G. traquairi, huxleyana, dufiana, and juddiana, n. spp.; E. T. Newton, P. R. Soc. lii, 1892, p. 389, Trias of Elgin, Scotland.
- Geikia, n. g., allied to Ptychognathus, Ow., for G. elginensis, n. sp., E. T. NEWTON, t. c. p. 390, Elgin.
- Elginia, n. g., nearest allied to Pariosaurus, Ow., for E. mirabilis, n. sp.; E. T. Newton, t. c. p. 391, Elgin.
- Parsiasaurus baini, n. sp., Seeley, t. c. p. 322, Trias, Karroo Group, S. Africa.

# BATRACHIA.

- O. BOTTGER issues a catalogue of the collection of Batrachians in the Senckenberg Museum. Frankfort a/M.: 1892, 8vo, 73 pp.
- W. L. SCLATER has compiled a list of the Batrachians in the Indian Museum, Calcutta. London: 1892, 8vo, 43 pp.
- O. S. STRONG, Anat. Anz. vii, 1892, pp. 467-471, writes on the structure and homologies of the cranial nerves of the Batrachians, as determined by their peripheral distribution and internal origin.
- H. Rex, Morph. JB. xix, 1892, pp. 295-311, pl. xi, contributes to the morphology of the cerebral veins in Batrachians.
- L. EDIEGER, Abh. Senck. Ges. xviii, 1892, 55 pp., 5 pls., has a memoir on the brain of Batrachians. Abstract in Anat. Anz. vii, 1892, pp. 472-476.
- A. B. Macallum, Tr. Can. Inst. ii, 1892, pp. 221-260, pl. i, has studies on the blood of Batrachians.
- M. C. DEKHUYSEN, Verh. Anat. Ges. 1892, pp. 90-103, pl. i, writes on the same subject.
- H. GRIESBACH, Leuckart Fetschrift, 1892, pp. 215-227, pl. xxiii, further contributes to the knowledge of the blood-corpuscles of Batrachians.
- O. C. FARRINGTON, Tr. Conn. Ac. viii, 1892, pp. 309-314, pl. xxiv, describes the kidney, with nephrostomes in Batrachians.

- H. F. E. Jungersen, Vid. Medd. 1892 (1893), pp. 32-72, pls. ii & iii, writes on the development of the Müllerian duct in Batrachians.
- A. C. EYCLESHYMER, J. Morph. viii, 1893, pp. 189-193, figs., has a note on the development of the optic vesicles in Batrachians.
- H. H. FIELD remarks on the metameric development of the kidney in Batrachians; Verh. Deutsch. Zool. Ges. ii, 1892, pp. 113-117.
- W. Roux, Verh. Anat. Ges. 1892, pp. 22-62, reviews recent researches on the segmentation of the Batrachian ovum.
- G. BORN, Anat. Anz. vii, 1892, pp. 772-781 & 803-811, figs., writes on the first phases of the development of the Batrachian ovum.
- E. O. JORDAN & A. C. EYCLESHYMER, Anat. Anz. vii, 1892, pp. 622-624, have a note on the cleavage of the Batrachian ovum.
- H. S. GREENOUGH, Bull. Soc. Z. Fr. xvii, 1892, pp. 57-59, writes on the same subject.
- F. LEYDIG, Biol. Centralbl. xii, 1892, pp. 205-221, writes on the dermal excrescences of breeding Batrachians, and pp. 444-467, on the structure of the skin.
- G. A. BOULENGER, Nat. Sci. i, 1892, pp. 185-190, figs., reviews our knowledge of the poisonous secretion of Batrachians.
- K. KNAUTHE, Zool. Anz. xv, 1892, pp. 20-23, records his experiments on the influence of temperature on Batrachians.

# ECAUDATA.

- C. EMERY, Atti Acc. Pontif. Lincei (5) i, 1892, Rend., pp. 203-206, publishes the results of further studies of the skeleton of the manus in tailless Batrachians.
- H. Adolphi, Morph. JB. xix, 1892, pp. 315-375, figs., pl. xii, writes on variations in the spinal nerves and vertebral column of Bufo viridis.
- C. HASSE, Z. wiss. Zool. lv, 1892, pp. 252-264, pl. xii, writes on the development of the vertebral column in tailless Batrachians.
- E. GAUPP, Verh. Anat. Ges. 1892, pp. 183-190, writes on the endocranium of Runa temporaria.
- D. BERTELLI, Monit. Zool. Ital. iii, 1892, pp. 203-207, remarks on the tympanic membrane of *Rana esculenta*. French translation in Arch. Ital. Biol. xviii, 1893, pp. 458-462.
- A. E. Smirnow has a paper, in Russian, on the nerves of Rana and Bufo; Protok. obsch. estest. Kazan, 1889-90, Append. No. 112 (1890), 6 pp.
- W. BIEDERMANN, Arch. Ges. Phys. li, 1892, pp. 455-508, pl. xi, investigates the movements of the chromatophores in tailless Batrachians.
- G. A. BOULENGER, P. Z. S. 1891, pp. 593-627, pls. xlv-xlvii, gives descriptions and figures of the Tadpoles of the European Batrachians, with a key to their identification.
- F. E. Schulze, Abh. Ak. Berl. 1892, 66 pp., 6 pls., continues his studies of the structure of the larva of *Pelobates fuscus*.

- L. CAMERANO, Boll. Mus. Zool. Tor. vii, 1892, No. 140, writes on the action of running water and of light on the development of tailless Batrachians.
- W. N. PARKER reports on Experiments on the Respiration in Tadpoles of Rana temporaria, Rep. Brit. Ass. 1891, p. 694.
- F. ZIEGLER, Anat. Anz. vii, 1892, pp. 211-215, figs., describes the external characters of the frog-embryo.
- H. Dehner, Verh. Ges. Würzb. (2) xxvi, 1892, pp. 1-18, pl. i, has a paper on the segmentation of the frog-ovum.
- HERTWIG, Arch. Mikr. Anat. xxxvii, 1892, pp. 353-503, pls. xvi-xx, has a paper on malformed frog-eggs.

### RANIDÆ.

Rana, L.: L. v. Méhely, Math. term. köz. xxv, 1892, pp. 1-63, pls. i-viii, has an illustrated memoir (in Hungarian) on the brown Frogs of Hungary (R. temporaria, L., arvalis, Nilss., and agilis, Thom.). R. temporaria, L.: on its occurrence in Ireland; SCHARFF, Irish Nat. i, 1893, p. 1. R. pretiosa, B. & G.: Notes by Test, Bull. U. S. Fish. Comm. 1891, p. 58 (1892). R. agilis, Thom.; on its occurrence in the Main district; LEYDIG, Zool. Gart. xxxiii, 1892, p. 321. R. camerani, Blgr., and R. macrocnemis, Blgr.: notes by BETTGER, Ber. Senck. Ges. 1892, pp. 134 & 136. R. esculenta, L.: on blue specimens; LEYDIG, Zool. Gart. xxxiii, 1892, p. 1. R. newtoni, Bocage: redescribed by BEDRIAGA, O Istituto (Coimbra), xxxix, 1892, p. 2 of reprint. R. masoni, Blgr. : redescribed by BETTGER, Ber. Offenb. Ver. xxix-xxxii, 1892, p. 138. R. vicina, Stol.: p. 342, pl. xxiv, fig. 1, and granulosa, And., p. 346, redescribed by W. L. SCLATER, P. Z. S. 1892, who also has notes on R. hascheana, Stol., p. 344, R. pulla, Stol., p. 344, R. gracilis, Gravh., p. 345, R. nigrovittata, Blyth, p. 345, and R. erythræa, Schleg., p. 345. R. assamensis, p. 343, pl. xxiv, fig. 2, Khasi Hills, limborgi, p. 344, fig. 3, Tenasserim, and tenasserimensis, p. 345, fig. 4, n. spp., W. L. SCLATER, t. c. R. holsti, n. sp., BOULENGER, Ann. N. H. (6) x, 1892, p. 302, Great Loo Choo I. R. octoplicata, p. 83, United States [= areolata, B. & G.—Rec.], and schlueteri, p. 84, Borneo, n. spp., WERNER, Zool. Anz. xvi, 1893. R. microdisca, n. sp., BETTGER, Ber. Offenb. Ver. xxix-xxxii, 1892, p. 137, Tengger Mts., Java. R. nyassæ, n. sp., Günther, P. Z. S. 1892, p. 558 (1893), Nyassaland. R. schmackeri, n. sp., BETTGER, Katal. Senckenb. p. 11, 1892, Ichang, China. R. oatesii, n. sp., Boulenger, Ann. N. H. (6) ix, 1892, p. 141, pl. ix, Toungoo, Burma. R. albofrenata, n. sp., F. MÜLLER, Festschr. Nat. Ges. Basel, 1892, p. 197, pl. iii, fig. i, Madagascar.

Cornufer boulengeri, n. sp., BŒTTGER, Katal. Senckenb. p. 18, 1892, New Britain.

- Rhacophorus difficilis, n. sp., Bettger, Katal. Senckenb. p. 14, 1892, Madagascar. R. boulengeri, n. sp., Peracca, Boll. Mus. Zool. Tor. vii, 1892, No. 112, pl. i, fig. 3, Madagascar. P. dulitensis, n. sp., Boulenger, P. Z. S. 1892, p. 507, pl. xxx, fig. 1, Mt. Dulit, Borneo.
- Chiromantis rufescens, Gthr.: on the 'nest' of a West African Tree Frog, probably of this species; W. J. Holland, Am. Nat. xxvii, 1893, p. 67.
- Ixalus cinerascens, Stol., and I. punctatus, And., (= glandulosus, Jerd.):
  note on the type specimens; W. L. Sclater, P. Z. S. 1892, p. 347.
  I. aurifasciatus, Schleg.: redescribed by Bættger, Ber. Offenb.
  Ver. xxix-xxxii, 1892, p. 142. I. schmackeri, n. sp., id. Katal.
  Senckenb. p. 17, 1892, Mindoro, Philippines.
- Rappia thomensis, Bocage: redescribed by BEDRIAGA, O Istituto (Coimbra), xxxix, 1892 (p. 5 of reprint). R. molleri, n. sp., id. t. c. p. 13, S. Thomé, Gulf of Guinea.

### ENGYSTOMATIDE.

- Phryniscus lævis, Gthr., n. var. exigua, Better, Katal. Senekenb. p. 22, 1892, Andes of Ecuador.
- Cophizalus, n. g., allied to Callula, for C. geislerorum, n. sp., BŒTTGER, Katal. Senckenb. p. 24, 1892, New Guines.
- Chaperina, n. g.: Tongue elliptic, free behind; no vomerine teeth: a transverse dermal fold on the palate in front of the esophagous; tympanum distinct; digits free, dilated into small disks; distal phalanx T-shaped; outer metatarsals bound together; pupil horizontal; transverse processes of sacrum rather strongly dilated; præcoracoids present, very slender; sternum cartilaginous; no omosternum. C. fasca, n. sp., MOCQUARD, Le Nat. xiv, 1892, p. 35, and Mém. Soc. Zool. v, 1892, p. 194, pl. vii, fig. 2, Sintang, Borneo.

### DYSCOPHIDÆ.

- Anodonthylu, n. g.: no teeth on palate; toes free; pupil horizontal.
  A. boulengeri, n. sp., F. MÜLLER, Festschr. Nat. Ges. Basel, 1892, p. 198, pl. iii, fig. 2, Madagascar.
- Fanchonia, n. g., referred to the Dyscophidæ, for F. elegans, n. sp., WERNER, Zool. Anz. xvi, 1893, p. 82, Tropical Africa?

### CYSTIGNATHIDE.

Calyptocephalus gayi, D. & B. Lucerta caudiverbera, L., is based on the larva of this frog; LATASTE, Act. Soc. Sci. Chili, i, 1892, p. 7. Hylodes lehmanni, p. 26, Colombia, fleischmanni, p. 27, Costa Rica, and W-nigrum, p. 28, Andes of Ecuador, n. spp., Bættger, Katal. Senckenb, 1892.

### BUFONIIDE.

- Pseudophryns dendyi, p. 62, North Gippsland, Victoria, and semimar-morata, p. 63, S. E. Victoria, n. spp., Lucas, P. R. Soc. Vict. (2) iv, 1892.
- Nectophryne hosii, n. sp., BOULENGER, P. Z. S. 1892, p. 508, pl. xxx, fig. 2, Mt. Dulit, Borneo.
- Bufo asper, Gravh.: on a remarkable specimen from Borneo; BOULENGER, P. Z. S. 1892, p. 508. B. fergusonii, n. sp., BOULENGER, J. Bomb. N. H. Soc. vii, 1892, p. 317, Travancore.
- Rhinophrynus dorsalis, D. & B.: description and figure of the skeleton, by A. Dugès, Nat. Mex. (2) ii, 1892, p. 98, pl. v.

### HYLIDÆ.

- Hyla amboinensis, 'Horst: notes by BŒTTGER, Ber. Offenb. Ver. xxix-xxxii, 1892, p. 156. H. albolimbata, p. 40, S. America, columbiana, p. 41, Colombia, and prosoblepon, p. 45, Costa Rica, n. spp., BŒTTGER, Katal. Senckenb. 1892. H. chloris, n. sp., BOULENGER, P. Linn. Soc. N.S.W. (2) vii, 1892, p. 403, New South Wales. H. (Trachycephalus) angustifrons, n. sp., WERNER, Zool. Anz. xvi, 1893, p. 83, Brazil [= ? H. nigromaculata, Tsch.—Rec.].
- Nototrema weinlandii, pp. 175 & 837, pl. i, Ecuador, and bolivianum, pp. 176 & 840, pl. ii, Bolivia, n. spp., Steindachner, Anz. Ak. Wien, 1892, and SB. Ak. Wien, ci, i, 1892.

### PELOBATIDÆ.

Spea laticeps, n. sp., Cope, Am. Nat. xxvii, 1893, p. 155, fig., Texas.

Leptobrachium carinense, Blgr.: on a specimen with vomerine teeth;

W. L. Sclater, P. Z. S. 1892, p. 347.

### DISCOGLOSSIDÆ.

L. v. Méhely publishes a translation in German of his paper on Bombinator; Math. Nat. Ber. Ung. x, 1892, pp. 55-79, pls. iv & v.

# CAUDATA.

- J. DE BEDRIAGA gives a French translation of his synopsis of the larvæ of European tailed Batrachians. C.R. Ass. Fr. Sci. 1891, ii, pp. 540-546.
- S. P. GAGE, P. Am. Ass. xli, 1892, p. 197, investigates the brain of *Molge viridescens*, Raf.
- A. C. EYCLESHYMER, Anat. Anz. vii, 1892, pp. 215-217, writes on paraphysis and epiphysis in the brain of Amblystoma.
- O. A. Andersson, Morph. JB., Anat. v, 1892, pp. 184-210, pls. xv-xviii, writes on the sympathetic nervous system of tailed Batrachians.

- K. W. SCHLAMPP, Z. wiss. Zool. iii, 1892, pp. 537-557, pl. xxi, investigates the structure of the eye of *Proteus anguineus*.
- H. H. WILDER, Morph. JB., Anat. v, 1892, pp. 155-176, pls. xii & xiii, describes the anatomy of the nasal region of Cryptobranchus and Amphiuma.
- C. HASSE, Z. wiss. Zool. liii (Suppl.), 1892, pp. 1-20, pls. i-iii, describes the development of the vertebral column of Molge vulgaris.
- J. S. KINGSLEY, Am. Nat. xxvi, 1892, pp. 671-680, describes the head of an embryo Amphiuma means.
- F. Houssay, C.R. cxv, 1892, pp. 132-135, writes on the circulation in the head of the Axolotl-embryo.
- H. W. Norris, J. Morph. vii, 1892, pp. 23-34, pls. iii & iv, investigates the development of the auditory vesicle in Amblystoma.
- R. Fick, Anat. Auz. vii, 1892, pp. 818-821, has a preliminary note on the fecundation of the Axolotl-egg.
- F. LEYDIG, Zool. Anz. xv, 1892, pp. 309-312 & 356, remarks on the receptaculum seminis of tailed Batrachians.
- M. HEIDENHEIN, Anat. Anz. vii, 1892, pp. 432-435, figs., records the presence of rudimentary anal glands in female Newts.
- G. A. BOULENGER, Morph. JB., Syst. vi, 1892, pp. 447-452, reviews recent discoveries on the fecundation in the tailed Batrachians.
- Drasch, Verb. Anat. Ges. 1892, pp. 244-253, makes observations on the poison-glands of the Salamander.
- Salamandra, Laur.: Molge luschani, Stdr., is referred to this genus by Boulenger, Ann. N. H. (6) ix, 1892, p. 74, and to the genus Spelerpes by Bedriaga, C.R. Congrès. Internat. Zool. i, 1892, p. 243.
- Triton longipes, Strauch., = Molge cristata, Laur.; BOULENGER, Ann. N. H. (6) ix, 1892, p. 144. Molge montandoni, Blgr.: larva described and figured by BOULENGER, op. cit. (6) x, 1892, p. 304, fig.
- Molge palmata, Schn.: Portuguese specimens are referred to this species by Ferreira, J. Sci. Lisb. (2) ii, 1892, p. 195 [belong, no with, to M. bosca, Lat.—Rec.]. M. rusconii, Gené: on the pairing; Bedriaga, Zool. Anz. xvi, 1893, p. 102.
- Tylototriton andersoni, n. sp., BOULENGER, Ann. N. H. (6) x, 1892, p. 304, Great Loo Choo I.
- Amblystoma tigrinum, Green: G. Haiin, Rev. Quest. Sci. (2) i, 1892, pp. 178-205, reviews our knowledge of the Axolotl and its metamorphoses; Spanish translation in Nat. Mex. (2) ii, 1892, pp. 218-230.
  J. M. Bertoldo, Le Nat. xiv, 1892, p. 15, records notes on the growth of Axolotls in confinement.
- Typhlotriton, n. g., of the group Desmognathinæ, but with the eyes concealed under the continuous skin of the head. T. spelæus, n. sp., Stejneger, P. U. S. Nat. Mus. xv, 1892, p. 115, pl. ix, Rock House Cave, Missouri.
- Necturus maculatus, Raf.: on its introduction in the Hudson River; W. B. Marshall, Am. Nat. xxvi, 1892, p. 779.

# APODA.

- P. & F. Sarasin, Anat. Anz. vii, 1892, pp. 812-815, have further remarks on the organ of hearing in Caecilians.
- Dermophis thomensis, Bocage, redescribed by BEDRIAGA, O Istituto (Coimbra), xxxix, 1892 (p. 13 of reprint).
- Siphonops paulensis, n. sp., BŒTTGER, Katal. Senckenb. p. 62, 1892, S. Paulo, Brazil.

# STEGOCEPHALA.

- This Order is proposed to be transferred to the Class Reptilia; H. G. SEELEY, Phil. Tr. claxxiii (B), 1892, p. 367.
- Ichniotherium cottæ, n. n. for foot-prints from the Permian of Friedrichroda, Thuringia; Pohlig, Leuckart Festschrift, 1892, p. 60, fig., pl. vii, fig. 1.



# V. PISCES.

BY

### G. A. BOULENGER.

This subject is treated as follows, viz. :-

- I. TITLES OF WORKS AND PAPERS ARRANGED
  ALPHABETICALLY ACCORDING TO AUTHORS' NAMES.
- II. GENERAL SUBJECT.
- III. FAUNÆ.
- IV. PALÆONTOLOGY.
  - V. REFERENCES, ARRANGED SYSTEMATICALLY.

## I. TITLES.\*

- AGASSIZ, A. Preliminary Note on some Modifications of the Chromatophores of Fishes and Crustaceans. Bull. Mus. C. Z. xxiii, 1892, pp. 189-193, 1 pl.
- ALCOCK, A. On Utero-gestation in Trygon bleekeri. Ann. N. H. (6) ix, 1892, pp. 417-427, pl. xix.
- ... Some Observations on the Embryonic History of Pteroplatea micrura. Op. cit. (6) x, 1892, pp. 1-8, pl. iv.
- A Case of Commensalism between a Gymnoblastic Anthomedusoid (Stylactis minoi), and a Scorpænoid Fish (Minous inermis). T. c. pp. 207-214, fig.
- Natural History Notes from H. M. Indian Marine Survey Steamer 'Investigator,' Lieut. G. S. Gunn, R.N., commanding. Series 11, No 5: On the Bathybial Fishes collected during the Season of 1891-92. T. c. pp. 345-365, pl. xviii.
- ALCOCK, R. The Digestive Processes of Ammocates. P. Cambr. Phil. Soc. vii, 1892, pp. 252-255.

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

- Andreae, A. Vorläufige Mittheilung über die Ganoiden (Lepidosteus und Amia) des Mainzer Beckens. Verh. Ver. Heidelb. (2) v, 1893, pp. 7-15, figs.
- Antipa, G. Ueber die Beziehungen der Thymus zu den sog. Kiemenspaltorganen bei Selachiern. Anat. Anz. vii, 1892, pp. 690-692, fig.
- AYERS, H. Vertebrate Cephalogenesis. II. A Contribution to the Morphology of the Vertebrate Ear, with a Reconsideration of its Functions. J. Morph. vi, 1892, pp. 1-360, pls. i-xii.
- Bassani, F. Gl' Ittioliti delle Marne di Salcedo e di Novale nel Vicentino. Atti Ist. Venet. (7) iii, 1892, pp. 1031-1045.
- Bean, T. H. Notes on the Fishes collected in Mexico by Professor Dugès, with Descriptions of new species. P. U. S. Nat. Mus. xv, 1892, pp. 283-287, pl. xliv.
- —. Una nuevo especio de Lampera. Nat. Mex. (2) ii, 1892, pp. 171 & 172, pl. viii.
- Beard, J. The Transient Ganglion Cells and their Nerves in Raia batis. Anat. Anz. vii, 1892, pp. 191-206, figs.
- —. Notes on Lampreys and Hags (My.cine). Op. cit. viii, 1892, pp. 59 & 60.
- Bellotti, C. Note al Manuale d'Ittiologia francese del Dott Emilio Moreau. Atti Soc. Ital. xxxiv, 1892, pp. 19-35.
- —. Un nuovo Siluroide giapponese. T. c. pp. 99-101, fig.
- BLANC, H. Note préliminaire sur la maturation et la fécondation de l'œuf de la truite. Bull. Soc. Vaud. (3) xxvii, 1892, pp. 272-275.
- Bohr, C. Sur la sécrétion de l'oxygène dans la vessie natatoire des Poissons. C.R. cxiv, 1892, pp. 1560-1562.
- BOLLMAN, C. H. A Review of the *Centrarchidæ*, or Freshwater Sunfishes, of North America. Rep. U. S. Fish Comm. f. 1888, pp. 557-579, pls. lxviii-lxxii (1891).
- BOULENGER, G. A. Third Account of the Fishes obtained by Surgeon-Major A. S. G. Jayakar at Muscat, East Coast of Arabia. P. Z. S. 1892, pp. 134-136.
- ---. On Lucioperca marina, C. & V. T. c. pp. 411-413, pl. xxv.
- —. Description of a new Blennioid Fish from Kamtschatka. T. c. pp. 583-585, fig. (1893).
  - ---. Note on Toxotes microlepis, Gthr., and Toxotes microlepis, Blyth. Ann. N. H. (6) ix, 1892, pp. 143 & 144.
  - —. Description of a new Siluroid Fish from China. T. c. p. 247.

- [BOULENGER, G. A.] On some new or little-known Fishes obtained by Dr. J. W. Evans and Mr. Spencer Moore, during their recent Expedition to the Province of Matto Grosso, Brazil. Ann. N. H. (6) x, 1892, pp. 9-12, pls. i & ii.
- —. Note on the Variations of the Lateral Shields in the Three-spined Stickleback (Gastrosteus aculeatus). Op. cit. xi, 1893, pp. 228 & 229.
- BOVERI, T. Die Nierencanälchen des Amphioxus. Ein Beitrag zur Phylogenie des Urogenitalsystems der Wirbelthiere. Morph. JB., Anat. v. 1892, pp. 429-510, figs., pls. xxxi-xxxiv.
- Ueber die Bildungsstätte der Geschlechtsdrüsen und die Entstehung der Genitalkammern bei Amphiocus. Anat. Anz. vii, 1892, pp. 170-181, figs.
- BOYER, E. R. The Mesoderm in Teleosts; especially its Share in the Formation of the Pectoral Fin. Bull. Mus. C. Z. xxiii, 1892, pp. 91-133, 8 pls.
- \*BEAUS, H. Ueber die Rami ventrales der vorderen Spinalnerven einiger Selachier. Inaugural Dissertation. Jena: 1892, 8vo, 35 pp.
- Breum, A. C. Tierleben. Die Fische. Dritte Auflage, unter Mitwirkung von W. HAACKE neubearbeitet von Pechuel-Loesche. Leipzic and Vienna: 1892, 8vo, 517 pp., 146 figs., 11 pls., 1 map.
- BRINGE, T. W., & HADDON, A. C. Contributions to the Anatomy of Fishes.

  11. The Air-Bladder and Weberian Ossicles in the Siluroid Fishes. Abstract, P. R. Soc. lii, 1892, pp. 139-157.
  - Browne, M. Notes upon Colobodus, a genus of Mesozoic Fossil Fishes. Rep. Brit. Ass. 1891, pp. 644 & 645 (1892).
  - BURCKHARDT, P. Das Centralnervensystem von Protopterus annectens.

    Eine vergleichend-anatomische Studie. Berlin: 1892, 8vo, 64 pp.,
    4to atlas, 5 pls,
  - —. Ueber das Centralnervensystem der Dipnoër. Verh. deutsch. Zool. Ges. ii, 1892, pp. 92-95.
  - Das Centralnervensystem von Protopterus annectens. SB. nat.
     Fr. 1892, pp. 23-25.
  - Burne, R. H. On the Presence of a Branchial Basket in Myxine glutinosa. P. Z. S. 1892, pp. 706-708, pl. xlvii (1893).
  - BUXBAUM, L. Ein Zug Aale auf der Wanderschaft im Main. Zool. Gart. xxxiii, 1892, pp. 20 & 21.
  - —. Der Wanderzug der Mainfische im Frühling 1892. T. c. pp. 184– 188.
  - CALDERWOOD, W. L. A Contribution to our Knowledge of the Ovary and Intra-ovarian Egg in Teleosteans. J. Mar. Biol. Ass. (2) ii, 1892, pp. 298-312, pls. xi & xii.
  - . Notes on Centrina salviani. T. c. pp. 322-324, pl. xiii.

- CAVAZZANI, E. L'ichthyotoxique chez les Petromyzon marinus. Arch. Ital. Biol. xviii, 1892, pp. 182-186.
- CLAYPOLE, E. W. On the Structure of the American Pteraspidian, *Palæaspis* (Claypole); with Remarks on the Family. Q. J. Geol. Soc. xlviii, 1892, pp. 542-561, figs.
- —... A new Gigantic Placoderm from Ohio. Am. Geol. x, 1892, pp. 1-4, fig.
- —. The Head of Dinichthys. T. c. pp. 199-207.
- —. A new Coccostean—Coccosteus cuyahagæ. Op. cit. xi, 1893, pp. 167-171, figs.
- COCKERELL, T. D. A. A provisional List of the Fishes of Jamaica. Institute of Jamaica, Bulletin No. 1, 1892, 16 pp.
- COLLINGE, W. E. On the Preservation of Teleostean Ova. Ann. N. H. (6) x, 1892, pp. 228-230.
- COPE, E. D. The Vertebrata of the Tertiary and Cretaceous Rocks of the North-West Territory. Contr. Canad. Pal. iii, 1891, 25 pp., 14 pls.
- —. On some New and Little-known Palæozoic Vertebrates. T. c. pp. 221-229, pls. vii & viii.
- —. On the Phylogeny of the Vertebrata. T. c. pp. 278-281, figs.
- —. On a new Horizon of Fossil Fishes. Abstract, P. Am. Ass. 1891 (1892), p. 285.
- CROOK, A. R. Ueber einige fossile Knochenfische aus der mittleren Kreide von Kansas. Palæontogr. xxxix, 1892, pp. 107-124, pls. xvxviii.
- CUENOT, L. Les Poissons commensaux et parasites. Le Nat. xiv, 1892, pp. 53-57, figs.
- CUNNINGHAM, J. T. Recent Additions to our Knowledge of the Natural History of some Devonshire Sea Fish. Rep. Tr. Devon Ass. xxiv, 1892, pp. 470-475.
- On the Rate of Growth of some Sea Fishes, and the Age and Size at which they begin to breed. J. Mar. Biol. Ass. (2) ii, 1892, pp. 222-264.
- . Ichthyological Contributions. T. c. pp. 325-332, pl. xiv.
- —... Report on the Probable Ages of Young Fish Collected by Mr. Holt in the North Sea. T. c. pp. 344-362.
- ---.. On the Growth of Food-Fishes and their Distribution at different Ages. Rep. Brit. Ass. 1891, pp. 685 & 686 (1892).
- —... The Reproduction of the Pilchard. T. c. p. 686.

- [CUNNINGHAM, J. T.] Distribution of Crystallogobius nilssonii. Rep. Brit. Ass. 1891, p. 687.
- ---. The Growth of the Pilchard or Sardine. Nature, xlv, 1892, pp. 255 & 256.
- —. Pilchards. T. c. p. 558.
- —. The Evolution of Flat-Fishes. Nat. Sci. i, 1892, pp. 191-199 & 635-638.
- DAGUIN, A., & BARDIES, C. Faune de la Haute Marne. Les Poissons. Paris: 1892, 16mo, 89 pp.
- Dalil., F. Zur Frage der Bewegung fliegender Fische. Zool. Anz. xv, 1892, pp. 106-108.
- Daimeries, A. Notes Ichthyologiques. Bull. Soc. Mal. Belg. (4) vi, 1891, pp. lxxiii-lxxvii.
- Danilewsky, B. Zur Physiologie des Centralnervensystems vom Amphioxus. Arch. ges. Phys. li, 1892, pp. 393-400.
- DAVIS, J. W. On the Fossil Fish-Remains of the Coal-Measures of the British Islands. Part I. Pleuracanthidæ. Sci. Tr. R. Dublin Soc. (2) iv, 1892, pp. 703-748, pls. lxv-lxxiii.
- DEAN, B. Notes on the Common Catfish. Rep. U. S. Fish. Comm. xix, 1892, pp. 301-305, pl. —. [Quoted from Separate Copy.]
- —... Pineal Fontanelle of Placoderm and Catfish. T. c. pp. 308-333, 14 pls. [Quoted from Separate Copy.]
- Dohrn, A. Die Schwann'schen Kerne der Selachierembryonen. Anat. Anz. vii, 1892, pp. 348-351.
- Dollo, L. Sur la Morphologie des Côtes. Bull. Sci. Fr. Belg. xxiv, 1892, pp. 113-129.
- —. Sur la Morphologie de la Colonne vertébrale. T. c. 19 pp. [separate copy].
- —. Sur le Lepidosteus suessoniensis. T. c. 5 pp. [separate copy].
- DOUGLAS-OGILBY, J. On some undescribed Reptiles and Fishes from Australia. Rec. Austral. Mus. ii, 1892, pp. 23-26.
- Dubois, R. Contribution à l'étude du mécanisme respiratoire des Dipnoïques et de leur passage de la torpeur estivale à la vie active. Ann. Soc. L. Lyon (2) xxxix, 1892, pp. 65-72, figs.
- DUNCKER, G. Ein muthmasslicher Bastard zwischen Pleuronectes platessa, L., und P. limanda, L. Zool. Anz. xv, 1892, pp. 451-453.
- Dunn, M. Pilchards. Nature, xlv, 1892, pp. 511 & 512.
- EDINGER, L. Untersuchungen über die vergleichende Anatomie des Gehirns: 2. Das Zwischenhirn. Erster Teil: Das Zwischenhirn der Selachier und der Amphibien. Abh. Senck. Ges. xviii, 1892, 55 pp., 5 pls. Abstract in Anat. Anz. vii, 1892, pp. 472-476.

- EIGENMANN, C. H. The Fishes of San Diego, California. P. U. S. Nat. Mus. xv, 1892, pp. 123-178, pls. x-xviii.
- ---. Branchiostoma elongatum, Sundevall, at San Diego, California. Am. Nat. xxvi, 1892, p. 70.
- —. On the Presence of an Operculum in the Aspredinidæ. T. c. p. 71.
- ---. The Percopsida of the Pacific Slope. Science, xx, 1892, pp. 233 & 234.
- EIGENMANN, C. H. & R. S. New Fishes from Western Canada. Am. Nat. xxvi, 1892, pp. 961-964.
- —. Preliminary Descriptions of New Fishes from the North-West. Op. cit. xxvii, 1893, pp. 151-154.
- ETHERIDGE, R., JUN., & WOODWARD, A. S. On the Occurrence of the genus *Belonostomus* in the Rolling Downs Formation (Cretaceous) of Central Queensland. Tr. R. Soc. Vict. ii, 1892, pp. 1-7, pl. i.
- EVERMANN, B. W. Report on the Establishment of Fish-Cultural Stations in the Rocky Mountain Region and Gulf States. Bull. U. S. Fish Comm. f. 1891, pp. 1-90, pls. i-xxxvi (1892).
- ---. Description of a new Sucker, Pantosteus jordani. Op. cit. 1892, pp. 51-56, fig. (1893).
- EWART, J. C. The Electric Organ of the Skate.—Observations on the Structure, Relations, Progressive Development, and Growth of the Electric Organ of the Skate. Phil. Tr. clxxxiii (B), 1892, pp. 389-420, pls. xxvi-xxx. Abstract, P. R. Soc. l, 1892, pp. 474-476.
- The lateral sense organs of Elasmobranchs. 1. The sensory canals of Læmargus. 11. The sensory canals of the common Skate, Raia batis. Zool. Anz. xv, 1892, pp. 116-120.
- FACCIOLÀ, L. Sull'esistenza di passaggio da alcune specie di Leptocefalidi agli adulti corrispondenti. Nat. Sicil. xi, 1893, pp. 144-148.
- FESTA, E. I Pesci del Piemonte. Boll. Mus. Zool. Tor. vii, 1892, No. 129, 125 pp.
- Forbes, H. O. On a Species of Regalecus or Great Oar-fish, caught in Okain's Bay. Tr. N. Z. Inst. xxiv, 1892, pp. 192-198.
- FUSARI, R. Sur les premières phases de développement des Téléostéens. Arch. Ital, Biol. xviii, 1892, pp. 204-239.
- GADEAU DE KERVILLE, H. Description d'un poisson et d'un oiseau monstreux. J. de l'Anat. Phys. xxviii, 1892, pp. 563-566, pl. xviii,
- GARDINER, E. J. L. Note on the Fish Remains from the Old Red Sandstone of Portishead. P. Bristol Soc. (2) vii, 1892, pp. 34 & 35.
- Garman, S. The Discoboli. Cyclopteridæ, Liparopsidæ, and Liparididæ. Mem. Mus. C. Z. xiv, No. 2, 1892, 96 pp., 13 pls.
- —. The Distribution of Fishes. Science, xix, 1892, p. 187.

- GIARD, A. The Evolution of Flat-Fish. Nat. Sci. i, 1892, pp. 356-359.
- GILI., T. Notes on the Tetraodontoidea. P. U. S. Nat. Mus. xiv, 1892, pp. 705-720, pl. xxxiv.
- GRASSI, G. B., & CALANDRUCCIO, S. Le Leptocefalide e la loro transformazione in Murenide. Atti Acc. Pontif. Lincei (5) i, 1892, Rend. pp. 375-379.
- GRIMM, G. Fischerei und Jagd in den russischen Gewässern. Uebersetzt von G. Josephy. Arch. f. Nat. lviii, i, 1892, pp. 191-208.
- GUITEL, F. Sur l'ovaire et l'œuf du Gobius minutus. C.R. exiv, 1892, pp. 612-616, figs.
- ---... Sur les mœurs du Clinus argenteus. Op. cit. cxlv, 1892, pp. 295-297.
- Observations sur les mœurs du Gobius minutus. Arch. Z. expér. (2) x, 1892, pp. 499-555, pl. xxii.
- GÜNTHER, A. List of the Species of Reptiles and Fishes collected by Mr. A. E. Pratt on the Upper Yang-tze-kiang and in the province Sze-Chuen, with Description of the New Species. Appendix 11 to PRATT'S "Snows of Tibet" (London: 1892, 8vo), pp. 238-250, pls. i-iv.
- ---. Description of a remarkable Fish from Mauritius, belonging to the Genus Scorpana. P. Z. S. 1891, pp. 482 & 483, pl. xxxix.
- —. Additional Note on the Occurrence of Lichia vadigo on the Cornish Coast. Ann. N. H. (6) x, 1892, pp. 335 & 336.
- HAASE, H. Beobachtungen über fliegende Fische. Zool. Anz. xv, 1892, p. 26.
- HASSE, C. Die Entwicklung der Wirbelsäule der Elasmobranchier. Dritte Abhandlung über die Entwicklung der Wirbelsäule. Z. wiss. Zool. lv, 1892, pp. 519-531, pl. xxi.
- Die Entwicklung der Wirbelsäule der Dipnoi. Vierte Abhandlung über die Entwicklung der Wirbelsäule. T. c. 1893, pp. 533-542, pl. xxii.
- ----. Allgemeine Bemerkungen über die Entwickelung und die Stammesgeschichte der Wirbelsäule. Anat. Anz. viii, 1893, pp. 288 & 289.
- HATSCHER, B. Zur Metamerie der Wirbeltiere. Nachtrag und Berichtigung. Op. cit. 1892, pp. 89-91.
- —. Die Metamerie des Amphioxus und des Ammocætes. Verh. Anat. Ges. 1892, pp. 136-161, figs.
- HEINCRE, F. Variabilität und Bastardbildung bei Cyprinoiden. Leuckart Festschrift, 1892, pp. 64-73, figs., pl. viii.
- HERRICK, C. L. Notes upon the Anatomy and Histology of the Prosencephalon in Teleosts. Am. Nat. xxvi, 1892, pp. 112-120, pls. vii & vii.

1892. [vol. xxix.]

- [HERRICK, C. L.] Additional Notes on the Teleost Brain. Anat. Anz. vii, 1892, pp. 422-431, figs.
- HERZENSTEIN, S. Ichthyologische Bemerkungen aus dem Zoologischen Museum der Kaiserlichen Akademie Wissenschaften. Mél. Biol. xiii, 1892, pp. 219-235, fig., and Bull. Pétersb. (2) iii, 1892, pp. 49-65, fig.
- HILGENDORF, F. Neue Stör-Art von Nord-Japan, Acipenser mikadoi. SB. nat. Fr. 1892, pp. 98-100.
- HOFFMAN, H. A., & JORDAN, D. S. A Catalogue of the Fishes of Greece, with Notes on the Names now in use, and those employed by classical authors. P. Ac. Philad. 1892, pp. 230-285.
- HOFFMANN, C. K. Zur Entwicklungsgeschichte des Herzens und der Blutgefässe bei den Selachiern. Ein Beitrag zur Kenntniss des unteren Keimblattes. Morph. JB. xix, 1893, pp. 592-648, figs., pls. xxii-xxv.
- —. Ueber die Entstehung der endothelialen Anlage des Herzens und der Gefässe bei Hai-Embryonen (Acanthias vulgaris). Anat. Anz. vii, 1892, pp. 270-273, figs.
- HOLMBERG, E. L. Sobre algunos Peces nuevos ó poco conocidos de la República Argentina. Rev. Argent. i, 1891, pp. 180-193.
- Hollt, E. W. L. On some Young Specimens of Centrolophus pompilus (Art.) from the Coast of Cornwall. J. Mar. Biol. Ass. (2) ii, 1892, pp. 265-267.
- —... North Sea Investigations. T. c. pp. 216-219 & 363-393.
- HOPKINS, G. S. On the digestive tract of some North American Ganoids.

  Abstract. P. Am. Ass. xli, 1892, p. 197.
- Howes, G. B. On the Arrangement of the Living Fishes, as based upon the Study of their Reproductive System. Rep. Brit. Ass. 1891, pp. 694 & 695 (1892).
- —. On the Customary Methods of Describing the Gills of Fishes. T. c. p. 702.
- JAEKEL, O. Ueber Dichelodus, Gieb., und einige Ichthyodoruliten, eine Entgegnung an Herrn A. Smith Woodward. JB. Mineral. 1892, i, pp. 145-151, figs.
- ---. Ueber Cladodus und seine Bedeutung für die Phylogenie der Extremitäten. SB. nat. Fr. 1892, pp. 80-92, fig.
- —. Ueber Chalcodus permianus. T. c. pp. 156-158.
- JORDAN, D. S., & DAVIS, B. M. A Preliminary Review of the Apodal Fishes or Eels inhabiting the waters of America and Europe. Rep. U. S. Fish. Comm. f. 1888, pp. 581-677, pls. lxxiii-lxxx (1892).
- KERMODE, P. M. C. Contributions to a Vertebrate Fauna of the Isle of Man. Fish. Zool. (3) xvii, 1893, pp. 65-70.

- KIRSCH, P. H. Notes on the Streams and Fishes of Clinton County, Kentucky, with a Description of a new Darter. Bull. U. S. Fish. Comm. f. 1890, pp. 289-292, fig. (1892).
- —. Notes on a Collection of Fishes from the Southern Tributaries of the Cumberland River in Kentucky and Tennessee. Op. cit. 1891, pp. 259-268 (1893).
- KLAATSCH, H. Beiträge zur vergleichenden Anatomie der Wirbelsäule.

  1. Ueber den Urzustand der Fischwirbelsäule. Morph. JB. xix, 1893, pp. 649-680, fig., pl. xxvi.
  - \*Klunzinger, C. B. Bodenseefische, deren Pflege und Fang. Stuttgart: 1892, 8vo, xii, 232 pp., figs.
  - KNAUTHE, K. Ueber Melanismus bei Fischen. Zool. Anz. xv, 1892, p. 25.
  - —. Ichthyologische Notiz. Op. cit. xvi, 1893, pp. 109 & 110.
  - KOLOMBATOVIC, G. Blennius zvonimiri, n. sp., Nova vrsta babice dalmatinskoga mora. Glasnik. Naravosl. Druztva, vii, 1892, pp. 107-112, pl. v.
  - KUNSTLER, M. Sur la fécondation artificielle du Cantharus griseus. Act. Soc. L. Bord. (5) v, 1893, pp. 241-243.
  - LATASTE, F. Sur les poissons vomis par les puits artésiens. Actes Soc. Sci. Chili, i, 1892, pp. xxxviii & xxxxix.
  - I.ENHOSSEK, M. v. Die Nervenendigungen in den Eudknospen der Mundschleimhaut der Fische. Vorläufige Mittheilung. Festschr, Nat. Ges. Basel, 1892, pp. 92-100.
  - —. Beobachtungen an den Spinalganglien und dem Rückenmark von Pristiurusembryonen. Anat. Anz. vii, 1892, pp. 519-539, figs.
  - LEYDIG, F. Integument brünstiger Fische und Amphibien. Biol. Centrabl. xii, 1892, pp. 205-221.
  - —. Blaufarbiger Wasserfrosch; Leuchtflecken der Ellritze. Zool. Gart. xxxiii, 1892, pp. 1-7.
  - —. Zur Kenntniss der Legeröhre des Bitterlings. T. c. pp. 129-133.
  - —. Springfrosch, Rana agilis; Ellritze, Phoxinus lavis. T. c. pp. 321-326.
  - LIDTH DE JEUDE, T. W. On Orthagoriscus nasus, Ranzani. Notes Leyd. Mus. xiv, 1892, pp. 127-128, pl. v.
  - LILLJEBORG, W. Sveriges och Norges Fiskar. Upsala: 3 vols. 8vo, 1884-91.
    - Fascicle 8, concluding the work, issued 1891.
  - Lucas, A. H. S. A new species of Freshwater Fish from Lake Nigothoruk, Mount Wellington, Victoria. P. R. Soc. Vict. (2) iv, 1892, pp. 27 & 28.

- LUCAS, F. A. On Carcharodon mortoni, Gibbes. P. Biol. Soc. Washington, vii, 1892, pp. 151 & 152.
- LÜTKEN, C. F. Spolia Atlantica. Scopelini Musei Zoologici Universitatis Hauniensis. Bidrag til Kundskab om det aabne Havs Laxesild eller Scopeliner. Med et Tillæg om en anden pelagisk Fiskeslægt. Vid. Selsk. Skr. (6) vii, 1892, pp. 223–297, 3 pls.
- LWOFF, B. Ueber die Keimblätterbildung bei den Wirbeltieren. Biol. Centralbl. xiii, 1893, pp. 40-50, 76-81.
- MACPHERSON, H. A. A Vertebrate Fauna of Lakeland, including Cumberland and Westmoreland with Lancashire North of the Sands. Edinburgh: 1892, 8vo, 552 pp.
- MARCY, —. Des Mouvements de natation de la Raie. CR. cxvi, 1893, pp. 77-81, figs.
- MARION, F. Reproduction et croissance de la Sardine dans le golfe de Marseille durant la campagne 1891. C.R. Ass. Fr. Sci. 1891, i, pp. 238 & 239 (1892).
- MARTIN, R., & ROLLINAT, R. Catalogue des Reptiles, Batraciens, et Poissons du Département de l'Indre. Mém. Soc. Zool. v, 1892, pp. 30-45.
- MAXWELL, H. Notes on the Freshwater Fishes of the Solway Area. Ann. Scott. N. H. i, 1892, pp. 103 & 104.
- McCormick, L. M. Descriptive List of the Fishes of Lorain County, Ohio. Bull. Oberlin Coll. Labor. No. 2, 1892, 34 pp., 14 pls.
- McIntosh, W. C. Notes from the St. Audrews Marine Laboratory. No. xiii. Ann. N. H. (6) x, 1892, pp. 97-108, pl. viii.
- Mojsisovics, A. v. Ueber eine auffälige (neue) Varietät des Acipenser ruthenus, L. SB. Ak. Wien, ci, i, 1892, pp. 328-330, pl. —.
- Mollier, S. Zur Entwickelung der Selachierextremitäten. Vorläufige Mitteilung. Anat. Anz. vii, 1892, pp. 351-365.
- Moreau, E. Manuel d'Ichthyologie française. Paris: 1892, 8vo, 650 pp., 3 pls.
- Morris, C. The Origin of Lungs, a Chapter in Evolution. Anat. Aux. xxvi, 1892, pp. 975-986.
- Newton, E. T. Note on a new species of *Onychodus* from the Lower Old Red Sandstone of Forfar. Geol. Mag. (3) ix, 1892, pp. 51 & 52, fig.
- NITSCHE, P. Die Zucht des Schleierschwanzes und des Teleskopfisches in Zimmer und Garten. Zool. Gart. xxxiii, 1892, pp. 289-300, 336-348.
- OLT, A. Lebensweise und Entwicklung des Bitterlings. Z. wiss. Zool. lv, 1893, pp. 543-575, pl. xxiii.

- OSTROUMOFF, A. Zoologische Excursion nach Mangyschlack und Basatschi (Russian text). Protok. obsch. estest. Kazan, 1889-90, Append. No. 113 (1890), 16 pp.
- OWSJANNIKOW, P. On the Embryology of the River-Lamprey. Translation. Ann. N. H. (6) xi, 1893, pp. 30-43.
- Perugia, A. Intorno ad alcuni pesci raccolti al Congo dal Capitano/Giacomo Bove. Ann. Mus. Genov. (2) x, 1892, pp. 967-977, figs.
- ——. Descrizione di una nuova specie di Myzus raccolta nel Rio Senegal dal Cap. G. Capurro. Op. cit. xii, 1892, pp. 1007 & 1008.
- —... Descrizione di due nuove specie di pesci raccolti in Sarawak dai Sig. G. Doria ed O. Beccari, T. c. pp. 1009 & 1010.
- PLATT, J. B. Fibres connecting the Central Nervous System and Chorda in Amphioxus. Anat. Anz. vii, 1892, pp. 282-284, figs.
- Poillig, H. Altpermische Saurierfährten, Fische und Medusen der Gegend von Friedrichroda i. Thür. Leuckart Festschrift, 1892, pp. 59-64, figs., pl. vii.
- Pollard, H. B. On the Anatomy and Phylogenetic Position of *Polypterus*. Morph. JB., Anat. v, 1892, pp. 387-428, figs., pls. xxvii-xxx.
- —. The Lateral Line System in Siluroids. T. c. pp. 525-551, pls. xxxv & xxxvi.
- POUCHET, G. Remarque sur deux Turbots à face nadirale pigmentée. C.R. Soc. Biol. (9) iv, 1892, pp. 200-202.
- Purvis, G. C. On the Pineal Eye of Lamna cornubica, or Porbeagle Shark. P. Phys. Soc. Edinb. 1890-91, pp. 62-67, pl. — (1892).
- RABL, C. Ueber die Entwicklung des Venensystems der Selachier. Leuckart Festschrift, 1892, pp. 228-235, figs.
- —. Ueber die Metamerie des Wirbeltierkopfes. Verh. Anat. Ges. 1892, pp. 104-135, figs., pl. ii.
- RAFFAELE, F. Ricerche sullo sviluppo del sistema vascolare nei Selacei. MT. z. Stat. Neap. x, 1892, pp. 441-479, pls. xxix-xxxi.
- RAMON Y CAJAL, S. La Retina de los Teleósteos y algunas observaciones sobre la de los vertebrados superiores. Ann. Soc. Esp. (2) i, 1892, pp. 281-305, figs.
- Reis, O. Zur Osteologie der Coelacanthinen. I Teil. Rumpfskelet, Knochen des Schädels und der Wangen, Kiemenbogenskelet, Schultergürtel, Becken, Integument und innere Organe. Inaugural Dissertation. Munich: 1892, 8vo, 39 pp., 2 pls.
- —. Zur Osteologie und Systematik der Belonorhynchiden und Tetragonolepiden. Geogn. JH. iv, 1892, pp. 143-166.
- Anhang über Dorypterus. T. c. pp. 167-170.

١

- [Reis, O.] Ueber die Zurechnung der Acanthodier zu den Selachiern. SB. nat. Fr. 1892, pp. 153-156. With remarks by O. JAEKEL, t. c. pp. 156-158.
- RETZIUS, G. Die sensiblen Nervenendigungen in der Haut des Petromyzon. Biol. Untersuch. (2) iii, 1892, pp. 37-40, pl. xiii.
- —. Die Nervenendigungen in den Endknospen resp. Nervenhügeln der Fische und Amphibien. Op. cit. iv, pp. 33-36.
- —. Ueber die sensiblen Nervenendigungen in den Epithelien bei den Wirbelthieren. T. c. pp. 37-44, pls. xi-xv.
- —. Das Caudalskelet der Myzine glutinosa. Biol. Fören. iii, 1892, pp. 81-84, pl. iii.
- —. Das hintere Ende des Rückenmarks und der Chorda dorsalis bei Petromyzon fluviatilis. Op. cit. iv, 1892, pp. 36-41, figs.
- ROHDE, E. Muskel und Nerv bei Mermis und Amphioxus. SB. Ak. Berlin, 1892, pp. 659-664.
- Rohon, J. V. Die Obersilurischen Fische von Oesel. 1. Theil. Thyestidæ und Tremataspiidæ. Mém. Ac. Pétersb. (7) xxxviii, 1892, No. 13, 88 pp., 2 pls.
- Röse, C. Ueber Zahnbau und Zahnwechsel der Dipnoer. Anat. Anz. vii, 1892, pp. 821-839, figs.
- RÜCKERT, J. Zur Entwickelungsgeschichte des Ovarialeies bei Selachiern. T. c. pp. 107-158, figs.
- —. Ueber die Verdoppelung der Chromosomen im Keimbläschen des Selachiereies. Op. cit. viii, 1892, pp. 44-52, figs.
- RYDER, J. A. The Sturgeons and Sturgeon Industries of the Eastern Coast of the United States, with an Account of Experiments bearing upon Sturgeon Culture. Bull. U. S. Fish. Comm. viii, 1890, pp. 231-281, pls. xxxvii-lix.
- —. On the Mechanical Genesis of the Scales of Fishes. P. Ac. Philad. 1892, pp. 219-224, figs., and Ann. N. H. (6) xi, 1893, pp. 243-248, figs.
- Sauvage, H. E. La faune ichthyologique du terrain permien français. C.R. cxvi, 1893, pp. 656 & 657.
- —... Recherches sur les Poissons du Lias Supérieur de l'Yonne. Zone à Ciment de Vassy. Bull. Soc. Autun, iv, 1891, pp. 59-81, pls. vi-ix, and v, 1892, pp. 386-399, pls. xvi & xvii.
- V. Schmidt, F. Ueber neue silurische Fischfunde auf Oesel. JB. Mineral. 1893, i, pp. 99-101.
  - SCHMIDT, L. Untersuchungen zur Kenntniss des Wirbelbaues von Amia calva. Z. wiss. Zool. liv, 1892, pp. 748-764, figs., pl. xxxiv.
  - Schulze, E. Fauna piscium Germaniæ. 11. Auflage. Königsberg: 1892, 8vo, 94 pp., figs.

- SCHULZE, F. E. Freie Nervenenden in der Epidermis der Knochenfische. SB. Ak. Berlin, 1892, pp. 87 & 88, fig.
- SEDGWICK, A. Notes on Elasmobranch Development. Q. J. Micr. Sci. xxxiii, 1892, pp. 559-586, pl. xxxv.
- SERVICE, R. Freshwater Fishes of the Solway Area. Ann. Scott. N. H. i, 1892, pp. 18-25.
- Sieglin, —. Einbürgerung fremder Nutzfische in Württemberg. JH. Ver. Württ. xlviii, 1892, pp. xxvii-xli.
- SIM, G. Occurrence of Triglops murrayi, Günther, on the East Coast of Scotland. Ann. Scott. N. H. i, 1892, p. 76.
- ---. Occurrence of Sebastes viviparus, Kroyer, off the East Coast of Scotland. Op. cit. ii, 1893, p. 47.
- SPENCER, W. B. Note on the Habits of Ceratodus forsteri. P. R. Soc. Vict. (2) iv, 1892, pp. 81-84.
- STEINDACHNER, F. Ueber einige neue und seltene Fischarten in den Sammlungen des k. k. naturhistorischen Hofmuseums. Anz. Ak. Wien, 1892, pp. 130-134, and Denk. Ak. Wien, lix, 1892, pp. 357-384, 6 pls.
- STÖHR, P. Die Entwickelung von Leber und Pankreas der Forelle. Anat. Anz. viii, 1893, pp. 205-208.
- STORMS, R. Note sur l'Ichthyologie fossile. Bull Soc. Belg. Géol. v, 1891, pp. 114-128 & 142-157.
- ----. Sur le Cybium (Enchodus) bleekeri, du Terrain Bruxellien. Op. cit. vi, Mém. pp. 3-14, pl. i (1892).
- STRACHLY, O. Die Karyomitose bei abnormer Entwicklung des Forelleneies. MT. embr. Inst. Wien (2) v, 1892, pp. 20-22.
- THOMSON, G. M. Notes on Sea-Fishes. Tr. N. Z. Inst. xxiv, 1892, pp. 202-215.
- TRAQUAIR, R. H. On the British Species of Asterolepidæ. P. Phys. Soc. Edinb. 1891-92, pp. 283-286 (1893).
- —. On a New Palsoniscid Fish, Myriolepis hibernicus, from the Coal-Measures, co. Kilkenny, Ireland. Geol. Mag. (3) x, 1893, pp. 54-56, pl. iii.
- ——. Notes on the Devonian Fishes of Campbelltown and Scaumenac Bay in Canada. No. 2. T. c. pp. 145-149, figs.
- —. Note on an abnormally developed Thornback (Raia clavata, L.).
  Ann. Scott. N. H. i, 1892, pp. 29 & 30, fig.
- —. List of the type and figured Specimens in the "Powrie Collection" of Fossils. T. c. pp. 31-39.
- —. On Malformed Trout from Scottish Waters. No. 1. T. c. pp. 92-103, pls. iii-v.
- ---. A new Fossil Fish from Dura Den. T. c. pp. 233-235.

- [Traquair, R. H.] An unusually Coloured Example of the Thornback (Raia clavata, Linn.). Ann. Scott. N. H. ii, 1893, p. 25, pl. i.
- TRIMEN, R. On the Occurrence of a rare Fish (Lophotes cepedianus) at the Cape of Good Hope. P. Z. S. 1891, pp. 483 & 484.
- VAILLANT, L. Remarques sur quelques Poissons du haut Tonkin. C.R. cxiv, 1892, pp. 1028 & 1029.
- ---. Sur le genre Megapleuron. T. c. pp. 1083 & 1084.
- —... Sur les affinités du genre Oreosoma, Cuvier. Op. cit. cxvi, 1893, pp. 598-600.
- —. Sur quelques Poissons rapportés du haut Tonkin par M. Pavie. Bull. Soc. Philom. (8) iv, 1892, pp. 125-127.
- —. Sur la possibilité du transport des galets dans l'appareil digestif des Poissons. Bull. Soc. Géol. (3) xx, 1892, pp. 111-113.
- Virchow, H. Das Dotterorgan der Wirbelthiere. Z. wiss. Zool. liii, Suppl., 1892, pp. 161-206, pl. x.
- WERNER, F. Untersuchungen über die Zeichnung der Wirbelthiere. Morph. JB., Syst. vi, 1892, pp. 155-229, pls. vi-x.
- WHITEAVES, J. F. Contributions to Canadian Palseontology. Vol. 1. Part IV. The Fossils of the Devonian Rocks of the Islands, Shores, or immediate vicinity of Lakes Manitoba and Winnipegosis. Geological Survey of Canada. Ottawa: 1892, 359 pp., 47 pls.
- WIEDERSHEIM, R. Das Gliedmassenskelet der Wirbelthiere, mit besonderer Berücksichtigung des Schulter- und Beckengürtels bei Fischen, Amphibien, und Reptilien. Jena: 1892, 8vo, 266 pp., figs., 17 pls.
  - WIJHE, J. W. VAN. Ueber Amphioxus. Anat. Anz. viii, 1893, pp. 152–172.
  - WILSON, E. B. On Multiple and Partial Development in Amphicaus. Op. cit. vii, 1892, pp. 732-740, figs.
  - WITCHELL, C. A., & STRUGNELL, W. B. The Fauna and Flora of Gloucestershire. Stroud: 1892, 8vo, 301 pp.
  - WOOD-MASON, J., & ALCOCK, A. Further Observations on the Gestation of Indian Rays; being Natural History Notes from H.M. Indian Marine Survey Steamer "Investigator." Series II, No. 2. P. R. Soc. 1, 1892, pp. 202-209.
  - WOODWARD, A. S. On the Skeleton of a Chimæroid Fish (*Ischyodus*) from the Oxford Clay of Christian Malford, Wiltshire. Ann. N. H. (6) ix, 1892, pp. 94-96.
  - ---. On some Teeth of new Chimeroid Fishes from the Oxford and Kimmeridge Clays of England. Op. cit. x, 1892, pp. 13-16, pl. iii.
  - —. Doubly-armoured Herrings. T. c. pp. 412 & 413.

- [Woodward, A. S.] Note on a Case of Subdivision of the Median Fin in a Dipnoan Fish. Ann. N. H. (6) xi, 1893, pp. 241 & 242, fig.
- ——. On the Lower Devonian Fish-Fauna of Campbelltown, New Brunswick. Geol. Mag. (3) ix, 1892, pp. 1-6, pl. i.
  - Further Contributions to the Devonian Fish-Fauna of Canada. T. c. pp. 481-485, figs., pl. xiii.
  - ---. Description of the Cretaceous Saw-Fish, Sclerorhynchus atavus. T. c. pp. 529-534, fig., pl. xiv.
  - —. Supplementary observations on some Fossil Fishes of the English Lower Oolites. P. Geol. Ass. xii, 1892, pp. 238-241, pl. iv.
  - —. Remarks on the Lower Tertiary Fish-Fauna of Sardinia. Rep. Brit. Ass. 1891, pp. 634 & 635 (1892).
  - ---. The Evolution of Fins. Nat. Sci. i, 1892, pp. 28-35, figs.

  - The Evolution of Sharks' Teeth. T. c. pp. 671-675, figs.
  - WOOLMAN, A. J. New Fishes from Chihuahua, Mexico. Am. Nat. xxvi, 1892, pp. 259-261.
  - Young, M., & Eagle Clarke, W. On Anarrhichas minor, Olafsen, and its occurrence on the Aberdeenshire Coast. Ann. Scott. N. H. i, 1892, pp. 26-29, pl. i.
  - ZENK, F. Ueber die Laichzeit des Aales. Mitth. Oesterr. Fischereiver. 1891, No. 39, and Zool. Gart. xxxiii, 1892, pp. 61 & 62.
- ZIEGLEB, H. E., & F. Beiträge zur Entwickelungsgeschichte von Torpedo. Arch. mikr. Anat. xxxix, 1892, pp. 56-102, figs., pls. iii & iv.

### II.—GENERAL SUBJECT.

- W. HAACKE & PECHUEL-LOESCHE have revised the volume dealing with Fishes in the new (third) edition of Brehm's Tierleben; Leipzic & Vienna: 1892, 8vo, 517 pp., 146 figs., 11 pls., 1 map.
- G. B. Howes, Rep. Brit. Ass. 1891, p. 694 (1892), expresses his views on the arrangement of Fishes (cf. Zool. Rec. xxviii, Pisces, p. 2).
- E. D. COPE, P. Am. Phil. Soc. xxx, pp. 280 & 281, figs., remarks on the phylogeny of the Fishes.
- R. WIEDERSHEIM, "Das Gliedmassenskelet der Wirbelthiere" (see p. 14); pp. 24-84 & 143-183 are devoted to the Fishes.
- A. S. WOODWARD, Nat. Sci. i, 1892, pp. 28-35, figs., gives a summary of the state of our knowledge of the evolution of fins.
- O. JAEKEL, SB. nat. Fr. 1892, pp. 80-92, also discusses the evolution of the limbs of Fishes.
- C. Hasse, Anat. Anz. viii, 1893, pp. 288 & 289, summarises his views on the development and phylogeny of the vertebral column.

- H. KLAATSCH, Morph. JB. xix, 1893, pp. 649-680, fig., pl. xxvi, contributes to the morphology and embryology of the vertebral column in Fishes.
- L. Dollo, Bull. Sci. Fr. Belg. xxiv, 1892 [quoted from separate copies], discusses the morphology of the vertebral column and ribs.
- C. Rabl, Verh. Anat. Ges. 1892, pp. 104-135, figs., pl. ii, discusses the metamery of the head in Vertebrates.
- H. AYERS, J. Morph. vi, 1892, pp. 1-360, pls. i-xii, has a contribution to the morphology of the Vertebrate ear.
- G. B. Howes, Rep. Brit. Ass. 1891, p. 702, remarks on the customary methods of describing the gills of Fishes. Gills of the Marsipobranch-Plagiostome type are, for anatomical purposes, termed *Cystobranchiæ*, and those of the higher Teleosteans *Pectinobranchiæ*.
- J. A. RYDER, P. Ac. Philad. 1892, pp. 219-224, figs., and Ann. N. H. (6) xi, 1893, pp. 243-248, figs., writes on the mechanical genesis of the scales of Fishes, and their segmental relation to the hard and soft parts.
- C. MORRIS, Am. Nat. xxvi, 1892, pp. 975-986, writes on the air-bladder and the origin of lungs.
- P. MITROPHANOPH, Trav. Soc. Varsovie. ii, 1892, 251 pp., 7 pls., has a paper, in Russian, dealing with the embryology of Fishes.

On the nerve-terminations in the epithelium of Fishes; G. Retzius, Biol. Untersuch. N. F. (2) iii, 1892, pp. 37-40, pls. xiii, & iv, 1892, pp. 33-44, pls. xi-xv.

On the nerve-terminations in the buccal epithelium of Fishes; Lenhossek, Festschr. Nat. Ges. Basel, 1892, pp. 92-100.

- A. AGASSIZ, Bull. Mus. C. Z. xxiii, 1892, pp. 189-193, 1 pl., has a preliminary note on some modifications of the chromatophores of Fishes.
- F. LEYDIG, Biol. Centralbl. xii, 1892, pp. 205-221, writes on the dermal excrescences of breeding Fishes, and, pp. 444-449, on the structure of the skin.
- J. T. CUNNINGHAM, Rep. Tr. Devon Ass. xxiv, 1892, pp. 470-475, reviews recent additions to our knowledge of the life-history of some Devonshire Sea Fishes.
- C. Bohr, C.R. cxiv, 1892, pp. 1560-1562, records his experiments on the secretion of oxygen in the air-bladder of Fishes.
- H. HAASE, Zool. Anz. xv, 1892, p. 26, and F. Dahl, t. c. pp. 106-108, further discuss the movements of flying Fishes.
- F. LATASTE, Actes Soc. Sci. Chili, i, 1892, pp. xxxviii & xxxix, remarks on Fishes expelled from artesian wells in France and the Algerian Sahara.
- L. Vaillant, Bull. Soc. Géol. (3) xx, 1892, pp. 111-113, remarks on stones transported in the digestive tract of Fishes.
- L. Curnot, Le Nat. xiv, 1892, pp. 53-57, figs., publishes a popular article on commensalism and parasitism among Fishes.

### III.—FAUNÆ.

### EUROPE.

"Vertebrate Fauna of Argyll and the Iuner Hebrides" (Edinburgh: 1892, 8vo); J. A. HARVIE-BROWN & T. E. BUCKLEY devote pp. 219-234 to Fishes.

Solway: R. Service, Ann. Scott. N. H. i, 1892, pp. 18-25, gives a list of the Freshwater Fishes of the Solway area. A few additions to the list by H. Maxwell, t. c. pp. 103 & 104.

Isle of Man: P. M. C. KERNODE, Zool. (3) xvii, 1892, pp. 65-70, enumerates the Fishes.

"Vertebrate Fauna of Lakeland;" H. A. MACPHERSON deals with the Fishes on pp. 466-526.

"The Fauna and Flora of Gloucestershire" (Stroud: 1892, 8vo); C. A. WITCHELL & W. B. STRUGNELL, Pisces, pp. 149-166.

Scandinavia: "Sveriges och Norges Firkar" (Upsala: 8vo, 3 vols., 1884-91), by W. LILLJEBORG; fascicle 8, concluding the work, was issued in 1891.

Russia in Europe: revised list of the Fishes; O. GRIMM, Arch. f. Nat. lviii, 1, 1892, pp. 195-207.

France: Manual of French Ichthyology, E. Moreau (Paris: 1892, 8vo, 650 pp., 3 pls); criticised by C. Bellotti, Atti Soc. Ital. xxxiv, 1892, pp. 19-35.

Dep. Haute Marne: Faune de la Haute Marne, Poissons; A. DAGUIN & C. BARDIES (Paris: 1892, 16mo, 89 pp.).

Dep. Indre: R. MARTIN & R. ROLLINAT, Mém. Soc. Zool. v, 1892, pp. 38-45, enumerate the Fishes.

Germany: ERWIN SCHULZE brings out a new, illustrated edition of his synopsis of the Fishes of Germany (Könisberg: 1892, 8vo, 94 pp., 49 figs.).

River Main: L. BRUXBAUM, Zool. Gart. xxxiii, 1892, pp. 184-188, reports on the migrations of Fishes in this river during the spring of 1892.

Bodensee: C. B. Klunzinger, Bodenseefische, deren Pflege und Fang (Stuttgart: 1892, 8vo, xii & 232 pp., figs.).

Württemberg: on the introduction of foreign Fish to; Sieglin, JH. Ver. Württ. xlviii, 1892, pp. xxvii-xli.

Piedmont: E. Festa, Bol. Mus. Zool. Tor. vii, 1892, No. 129, describes the Fishes.

Greece: H. A. HOFFMAN & D. S. JORDAN, P. Ac. Philad. 1892, pp. 230-285, publish a catalogue of the Fishes, with notes on the names now in use and those employed by classical authors.

### ARIA.

Mangyschlak and Busatchi: A. OSTROUMOFF, Protok. obsch. estest. Kazan, 1889-90, Append. No. 113 (1890), gives an account, in Russian, of a zoological excursion to.

Upper Yang-tse-Kiang and Province Sze-Chuen: A. GÜNTHER gives a list of the Fishes collected by Mr. A. E. Pratt; Appendix ii to PRATT'S "Snows of Tibet" (London: 1892, 8vo), pp. 243-250, pls. i-iv.

Upper Tonkin: L. Vaillant, C.R. cxiv, 1892, pp. 1028 & 1029, and Bull. Soc. Philom. (8) iv, 1892, pp. 125 & 127, records some Fishes.

Muscat, East Coast of Arabia: G. A. BOULENGER, P. Z. S. 1892, pp. 133-136, has a third report on the Fishes obtained by Surgeon-Major Jayakar.

Laccadive Sea, Gulf of Manaar, and Coast of Coromandel: A. Alcock, Ann. N. H. (6) x, pp. 345-365, pl. xviii, reports on the bathybial Fishes obtained by the "Investigator," during the season of 1891-92.

### AFRICA.

Congo: A. Perugia, Ann. Mus. Genov. (2) x, 1892, pp. 967-977, figs., reports on a collection of Fishes.

### AMERICA.

Montana and Wyoming: B. H. EVERMANN, Bull. U. S. Fish. Comm. f. 1891, pp. 41-55 (1892), gives an annotated list of Fishes.

Ohio: L. M. McCormick, Bull. Oberlin Coll. Labor., No. 2, 1892, 34 pp., 14 pls., gives a descriptive list of the Fishes of Lorain County.

Kentucky: P. H. Kirsch, Bull. U. S. Fish Comm. f. 1890, pp. 291 & 292, gives a list of the Fishes of Clinton County.

Kentucky and Tennessee: P. H. Kirsch, op. cit. 1891, pp. 259-268 (1893), reports on a collection from the Southern tributaries of the Cumberland River.

Texas: B. H. EVERMANN, t. c. pp. 75-88, has notes on Fishes.

California: C. H. EIGENMANN, P. U. S. Nat. Mus. xv, 1892, pp. 123-178, pls. x-xviii, gives an account of his observations on the Fishes of San Diego, with special reference to the spawning habits, embryology and migrations.

Jamaica: T. D. A. COCKERELL has compiled a provisional list of the Fishes, published as Bulletin No. 1, by the Institute of Jamaica, 1892.

Matto-Grosso, Brazil: G. A. BOULENGER, Ann. N. H. (6) x, 1892, pp. 9-12, pls. i & ii, describes some new or little-known Fishes.

Argentina: E. L. Holmberg, Rev. Argent. i, 1891, pp. 180-193, describes new or little-known Fishes.

### NEW ZEALAND.

G. M. THOMSON, Tr. N. Z. Inst. xxiv, 1892, pp. 202-215, has notes on the occurrence, habits, weight, &c., of various Marine Fishes in New Zealand.

### IV.—PALÆONTOLOGY.

R. STORMS, Bull. Soc. Belg. Géol. v, 1891, pp. 114-128 & 142-157, reviews recent progress in fossil Ichthyology.

R. H. TRAQUAIR, Ann. Scott. N. H. i, 1892, pp. 31-39, gives a list of the type and figured specimens of fossil Fishes in the "Powrie Collection," now in the Edinburgh Museum.

British Islands: J. W. DAVIS, Sci. Tr. R. Dublin Soc. (2) iv, 1892, pp. 703-748, pls. lxv-lxxiii, describes and figures the *Pleuracanthidæ* from the Coal Measures.

England: A. S. WOODWARD, P. Geol. Ass. xii, 1892, pp. 238-241, pl. iv, makes supplementary observations on some fossil Fishes of the Lower Oolites. E. J. L. GARDINER, P. Bristol Soc. (2) vii, 1892, pp. 34 & 35, has a note on Fish-remains from the Old Red Sandstone of Portishead.

France: H. E. SAUVAGE, C.R. cxvi, 1893, p. 656, remarks on the fossil Fish of the Permian.

Dep. Yonne: H. E. SAUVAGE, Bull. Soc. Autun, iv, 1891, pp. 59-81, pls. vi-ix, and op. cit. v, 1892, pp. 386-399, pls. xvi & xvii, publishes researches on the Fish of the Upper Lias.

Belgium: A. Daimeries, Bull. Soc. Mal. Belg. (4) vi, 1891, pp. lxxiii-lxxvii, describes fossil Fish-remains.

Isle of Oesel, Baltic: J. V. ROHON monographs the *Thyestidæ* and *Tremataspidæ* of the Upper Silurian; Mém. Ac. Pétersb. (7) xxxviii, 1892, No. 13, 88 pp., 2 pls. F. SCHMIDT, JB. Mineral. 1893, pp. 99-101, reports on the discovery of New Silurian Fish-remains on the Isle of Oesel.

Italy: F. BASSANI, Atti Ist. Venet. (7) iii, 1892, pp. 1031-1045, has a note on Tertiary Fish-remains from Salcedo and Novale.

Sardinia: A. S. WOODWARD, Rep. Brit. Ass. 1891, p. 634 (1892), makes short remarks on the Lower Tertiary Fish fauna.

Canada: A. S. WOODWARD, Geol. Mag. (3) ix, 1892, pp. 1-6, pl. i, and pp. 481-485, pl. xiii, and R. H. Traquair, op. cit. x, 1893, pp. 145-149, figs., have notes on some Devonian Fish-remains. J. F. Whiteaves, Contr. Can. Pal. i, pt. iv, pp. 353-356, has notes on the Fishes of the Devonian of Lakes Manitoba and Winnipeg. E. D. Cope describes some Fish-remains from the Oligocene or Lower Miocene beds of the Cypress Hills; op. cit. iii, 1891, pp. 1-4.

Pennsylvania: E. D. COPE, P. Am. Phil. Soc. xxx, 1892, pp. 221-229, pls. vii & viii, describes various new and little-known palæozoic Fishremains, mostly from the Chemung beds.

Kansas: A. R. Crook, Paleontogr. xxxix, 1892, pp. 107-124, pls. xv-xviii, describes some Teleosteans from the Chalk.

South Dakota: E. D. Cope, P. Am. Ass. 1891 (1892), p. 285, remarks on the discovery of Oligocene (?) Fish-remains in the Ree Hills.

## V.—REFERENCES, ARRANGED SYSTEMATICALLY.

# TELEOSTOMI.

## TELEOSTEI.

- C. L. HERRICK, Am. Nat. xxvi, 1892, pp. 112-120, pls. vii & viii, has notes on the anatomy and histology of the prosencephalon of Teleosteans. The same author, Anat. Anz. vii, 1892, pp. 422-431, figs., has additional notes on the Teleost brain.
- S. RAMON Y CAJAL, An. Soc. Esp. (2) i, 1892, pp. 281-305, figs., remarks on the retina of Teleosteans.
- F. E. SCHULZE, SB. Ak. Berlin, 1892, pp. 87 & 88, fig., has a note on nerve-terminations in the epidermis of Teleosteans.
- R. Fusari, Arch. Ital. Biol. xviii, 1892, pp. 204-239, gives a French abstract of his researches on the early stages of development in Teleosteans.
- P. STÖHR, Anat. Anz. viii, 1893, pp. 205-238, writes on the development of the liver and pancreas in the Trout.
- W. L. CALDERWOOD, J. Mar. Biol. Ass. (2) ii, 1892, pp. 298-312, pls. xi & xii, contributes to our knowledge of the ovary and intra-ovarian egg in Teleosteans.
- E. R. BOYER, Bull. Mus. C. Z. xxiii, 1892, pp. 91-133, 8 pls., writes on the mesoderm in Teleosteans, and its share in the formation of the pectoral fin.
- J. T. CUNNINGHAM, J. Mar. Biol. Ass. (2) ii, 1892, pp. 222-264 & 344-362, writes on the rate of growth of some sea Fishes (Gadoids, Pleuronectoids, and Clupeoids), and the age and size at which they begin to breed; also, t. c. pp. 325-332, on a few larval forms of Sole and Mackerel.
- W. E. COLLINGE, Ann. N. H. (6) x, 1892, pp. 228-230, reports on the results of his method of preserving Teleostean ova.

### ACANTHOPTERYGII.

### PERCIDE.

- Lucioperca, Cuv.: BOULENGER, P. Z. S. 1892, p. 411, gives a synopsis of the species, and describes and figures, pl. xxv, a specimen of L. marina, C. & V.
- Etheostoma cinereum, Storer, redescribed by Kirsch, Bull. U. S. Fish Comm. f. 1891, p. 264 (1893). E. guentheri, p. 962, Winnipeg and Iowa, and quappella, p. 963, Qu'Appelle, W. Canada, n. spp., Eigenmann, Am. Nat. xxvi, 1892, p. 963. E. scovellii, n. sp., Woolman, t. c. p. 260, Chihuahua, N. Mexico. E. wrightii, n. sp., McCormick, Bull. Oberlin Coll. Labor., No. 2, 1892, p. 30, Lorain County, Ohio. E. obeyense, n. sp., Kirsch, Bull. U. S. Fish Comm. f. 1890, p. 292, fig. (1892), Kentucky.

- Siniperca scherzeri, n. sp., Steindachner, Anz. Ak. Wien, 1892, p. 130, and Denk. Ak. Wien, lix, 1892, p. 357, pl. i, fig. 1, Shanghai.
- Percalabraz japonicus, C. & V.: on a large specimen; STEINDACHNER, Denk. Ak. Wien, lix, 1892, p. 359.
- Serranus dermopterus, Schleg., redescribed and figured; id. t. c. p. 359, pl. ii, fig. 1.
- C. H. BOLLMAN, Rep. U. S. Fish. Comm. f. 1888, pp. 557-579, pls. lxviii-lxxii (1891), reviews the N. American Centrarchide.

### SQUAMIPINNES.

Toxotes microlepis, Gthr., is distinct from T. microlepis, Blyth, which is renamed T. blythii; BOULENGER, Ann. N. H. (6) ix, 1892, p. 143.

### SPARIDE.

Box lineatus, n. sp., BOULENGER, P. Z. S. 1892, p. 134, Muscat, Arabia. Cantharus griseus, C. & V.: on artificial fecundation; Kunstler, Act. Soc. L. Bord. (5) v, 1893, p. 241.

### SCORPÆNIDÆ.

- Sebastes riviparus, Kroy.: on its occurrence off the East Coast of Scotland; G. Sim, Ann. Scott. N. H. ii, 1893, p. 47. [The fish described as S. norvegicus by Sim is Scorpæna dactylopterus.—Rec.]
- Scorpæna frondosa, n. sp., GÜNTHER, P. Z. S. 1891, p. 482, pl. xxxix, Mauritius.
- Minous inermis, Alcock: description of an Anthomedusoid commensal on this fish; Αιcock, Ann. N. H. (6) x, 1892, p. 207.

### NANDIDE.

Acharnes niederleinii, p. 181, Missiones, Argentina, and chacoensis, p. 182, Chaco, Argentina, n. spp., Holmberg, Rev. Argent, i, 1891.

### BERYCIDÆ.

Oreosoma, Cuv., is referred to the Berycidæ; VAILLANT, C.R. cxvi, 1893, p. 598.

### SCIÆNIDÆ.

Sciena albiflora, Richards., (= Pagrus macrophthalmus, Basil.), p. 361, amoyensis, Blkr., p. 362, and amblyceps, Blkr., p. 363, redescribed by STEINDACHNER, Denk. Ak. Wien, lix, 1892.

### CARANGIDE.

Lichia vadigo, Risso: on specimens from the Cornish Coast; GÜNTHER, Ann. N. H. (6) x, 1892, p. 335.

# STROMATEIDE.

Centrolophus pompilus, Art.: on some young specimens from the Coast of Cornwall; Holt, J. Mar. Biol. Ass. (2) ii, 1892, p. 265.

# SCOMBRIDÆ.

- Thynnus macropterus, Schleg., = T. albacora, Lowe; Boulenger, P. Z. S. 1892, p. 135.
- Cybium (Enchodus) bleekeri, Winkler (foss.), described and figured by STORMS, Mém. Soc. Belg. Géol. vi, 1892, p. 3, pl. i.

#### COTTIDÆ.

- Triglops murrayi, Gthr., recorded from the E. Coast of Scotland by G. Sim, Ann. Scott. N. H. i, 1892, p. 76.
- Cottus philonips and onychus, n. spp., EIGENMANN, Am. Nat. xxvi, 1892, p. 963, W. Canada.
- Argyrocottus, n. g., resembling Melletes, Bean, in the very long ventrals, but distinguished by the absence of cleft behind the last gill, of teeth on the palate, and of scales. A. zanderi, n. sp., HERZENSTEIN, Mél. biol. xiii, 1892, p. 219, Sachalien I.

#### DISCOBOLI.

- S. GARMAN brings out a monograph of these Fishes, which he divides into three families: Cyclopteridæ, Liparopsidæ, Liparididæ; Mem. Mus. C. Z. xiv, No. 2, 1892, 96 pp., 13 pls.
- Cyclopteroides, n. g., closely allied to Cyclopterus, for C. gyrinops, n. sp.; GARMAN, t. c. p. 37, pl. xi, figs. 4-9, Alaska.
- /Liparops, n. g., for Cyclopterus stelleri, Pall.; GARMAN, t. c. p. 42.
  - Liparis, Act.: the following species are figured by GARMAN, t. c.:—L. agassizii, Putn., pls. i-iii, pulchellus, Ayres, pl. iv, mucosus, Ayres, pl. v, calliodon, Pall., pl. vi, figs. 1-5, antarctica, Putn., pl. vi, figs. 6-10, liparis, L., pl. vii, figs. 1-5, and montagui, Don., pl. vii, figs. 6-20.
  - Remarks on the distribution of the Cyclopteri; GARMAN, Science, xix, 1892, p. 187.

#### GOBIIDÆ.

- Gobius minutus, Gm.: extensive notes on variations of fin-rays, scaling and colour, sexual differences, and habits of this Fish, which is figured; Guitel, Arch. Z. expér. (2) x, 1892, p. 499, pl. xxii: on the ovary and eggs; Guitel, C.R. cxiv, 1892, p. 612. G. beccarii, n. sp., Perugia, Ann. Mus. Genov. (2) xii, 1892, p. 1010, Sarawak.
- Crystallogobius nilssonii, Gill: on its abundance at Plymouth; J. T. Cunningham, Rep. Brit. Ass. 1891, p. 687 (1892).

# BLENNIIDE.

- Blennius zvonimiri, n. sp., Kolombatovic, Glasnik Naravoal. Brustza, vii, 1892, p. 107, pl. v, Coast of Dalmatia.
- Clinus argenteus, C. & V.: on nesting and oviposition; F. Guitel, C.R. cxv, 1892, p. 295.
- Anarrhichas minor, Olafs., figured, with notes on its occurrence on the Scottish Coast; M. Young & W. Eagle Clarke, Ann. Scott. N. H. i, 1892, p. 26, pl. i.
- √ Blenniophidium, n. g., allied to Cebedichthys; BOULENGER, P. Z. S. 1892, p. 583 (1893). B. petropauli, n. sp., id. t. c. p. 584, fig., Kamtschatka.

# MASTACEMBELIDÆ.

Mastacembelus marmoratus, n. sp., Perugia, Ann. Mus. Genov. (2) x, 1892, p. 968, Congo.

#### MUGILIDA.

- Mugil alatus, n. sp., STEINDACHNER, Anz. Ak. Wien, 1892, p. 133, and Denk. Ak. Wien, lix, 1892, p. 364, Madagascar.
- Myzus capurrii, n. sp., Perugia, Ann. Mus. Genov. (2) xii, 1892, p. 1007, Senegal.

## GASTROSTEIDÆ.

Gastrosteus aculeatus, L.: Remarks on the variations of the lateral shields in specimens from Ostend Harbour; BOULENGER, Ann. N. H. (6) zi, 1893, p. 228.

#### TRACHYPTERIDÆ.

Regalecus grillii, Lindr., redescribed by H. O. FORBES, Tr. N. Z. Inst. xxiv, 1892, p. 192.

#### LOPHOTIDE.

Lophotes cepedianus, C. & V., recorded from the Cape of Good Hope by TRIMEN, P. Z. S. 1891, p. 483.

#### INCERTA SEDIS.

V Pseudoscopelus, n. g. (Trachinidarum?). Body naked; spinous dorsal ahort, soft dorsal and anal long; a ventral lateral line; mouth deeply cleft. P. scriptus, n. sp., LÜTKEN, Vid. Selsk. Skr. (6) vii, 1892, p. 285, pl. i, figs. 3-5, Bahamas.

1892. [vol. xxix.]

# ACANTHOPTERYGII PHARYNGOGNATHI.

#### CHEOMIDES.

- Chromis fusciatus, n. sp., Perugia, Ann. Mus. Genov. (2) x, 1892, p. 970, Congo.
- Acara bartoni, n. sp., BEAN, P. U. S. Nat. Mus. xv, 1892, p. 286, pl. xliv, fig. 3, San Luis Potosi, Mexico.
- Heros centralis, n. sp., Holmberg, Rev. Argent. i, 1891, p. 183, Santiago del Estero, Argentina.
- Crenicichla sazutilis, L., n. var., semicincta, STEINDACHNER, Denk. Ak. Wien, lix, 1892, p. 376, Bolivia.

# ANACANTHINI.

# GADIDÆ.

J. T. CUNNINGHAM, Rep. Brit. Ass. 1891, p. 685, remarks on the growth of various British Gadoids, and their Distribution at different ages Also J. Mar. Biol. Ass. (2) ii, 1892, p. 222-264.

#### OPHIDIDE.

- Ammodytes lanceolatus and A. tobianus: on the eggs and young stages; McIntosh, Ann. N. H. (6) x, 1892, p. 97.
- Lamprogrammus fragilis, n. sp. (?), ALCOCK, t. c. p. 348, Bay of Bengal, 678 faths.
- V Hephthocara, n. g.: head large, with thin, smooth, uncrested bones, scaleless; no armature, but a weak opercular spine; body compressed, tapering, covered with deciduous cycloid scales; eye moderate; snout not overhanging the jaws; mouth with obliquely ascending cleft, and with the lower jaw prominent; villiform teeth in the jaws, palatines, and vo.ner; no barbel or hyoid filaments; gill-openings wide; gill-membranes separate; four gills; no pseudobranchiæ; eight branchiostegals; lateral line indistinguishable; vertical fins confluent; pectoral fins entire; no ventral fins. H. simum, n. sp., Alcock, t. c. p. 349, pl. xviii, fig. 1, Gulf of Manaar, 902 faths.

#### MACRURIDE.

Macrurus macrolophus, Alcock, figured by Alcock, Ann. N. H. (6) x, 1892, p. 351. M. hextii, Alc., and wood-masoni, Alc.: further notes; id. t. c. pp. 351 & 353. M. calorhynchus, Risso: description of an anomalous specimen; Herzenstein, Mél. biol. xiii, 1892, p. 223.

# PLEURONECTIDÆ.

W. C. McIntosh, Ann. N. H. (6) x, 1892, p. 102, remarks on the ova and larve of certain British Peuronectoids.

- E. W. L. HOLT, J. Mar. Biol. Ass. (2) ii, 1892, pp. 363-379, writes on the relation of size to sexual maturity in Pleuronectoids.
- J. T. CUNNINGHAM, Nat. Sci. i, 1892, pp. 191-199 & 635-638, and A. GIARD, t. c. pp. 356-359, discuss the evolution of Flat-Fishes.
- G. POUCHET, CR. Soc. Biol. (9) iv, 1892, pp. 200-202, remarks on Turbots with partially pigmented blind side.
- Pleuronectes, L.: on an apparent hybrid between P. platessa, L., and P. limanda, L.; Duncker, Zool. Anz. xv, 1892, p. 451.
- Zeugopterus norvegicus, Gthr.: note by J. T. Cunningham, J. Mar. Biol. Ass. (2) ii, 1892, p. 325.

## PHYSOSTOMI.

# SILUBIDE.

- B. Dean, Rep. U. S. Fish Comm. xix, 1892, pp. 308-333, 14 pls., has notes on the anatomy of the Siluroids, with special reference to the pineal fontanelle; a list is given of the characters in which the Fish of this family differ from the other Teleosteans.
- H. B. POLLARD, Morph. JB., Anat. v, 1892, pp. 525-551, pls. xxxv & xxxvi, describes the lateral line system in a few Siluroids of the genera Clarias, Callichthys, Auchenaspis, Trichomycterus, and Chatostomus.
- T. W. BRIDGE & A. C. HADDON, P. R. Soc. lii, 1892, pp. 139-157, publish an abstract of their paper on the air-bladder and Weberian ossicles in the Siluroids.
- C. H. EIGENMANN, Am. Nat. xxvi, 1892, p. 71, corrects a previous statement that the operculum is absent in Aspredo.
- Diastatomycter, Vaill., = Hemisilurus, Blkr.; VAILLANT, Bull. Soc. Philom. (8) v. 1893, p. 15.
- Macrones pluriradiatus, n. sp., Valllant, op. cit. iv, 1892, p. 126, Upper Tonkin. M. pratti, n. sp., Günther, in Pratt's "Snows of Tibet," p. 245, pl. i, fig. B, Province of Sze-Chuen, China.
- Pseudobagrus eupogon, n. sp., Boulenger, Ann. N. H. (6) ix, 1892, p. 247, Shanghai.
- √ Neobagrus, n. g., distinguished from Pseudobagrus by the small number
  (13) of anal rays, the rounded caudal, the absence of teeth on the
  palate, and the attachment of the gill-membranes to the isthmus.
  Bellotti, Atti Soc. Ital. xxxiv, 1892, p. 100. N. fuscus, n. sp.,
  id. t. c. p. 101, fig., Yokohama, Japan.
  - Rhineaster rheas, n. sp. (foss.), COPE, Contr. Canad. Pal. iii, 1891, p. 3, pl. i, fig. 3, Lower Miocene, Canada.
  - Amiurus catus, L., note and figure; B. DEAN, Rep. U. S. Fish Comm. xix, 1892, p. 302, pl. —. A. cancellatus and maconnellii, n. spp. (foss.), Cope, Contr. Canad. Pal. iii, 1891, pp. 3 & 4, pl. i, figs. 4-7, Lower Miocene, Canada.
  - Amblyceps marginatus, n. sp., GÜNTHER, in Pratt's "Snows of Tibet," p. 245, pl. ii, fig. A, Province of Sze-Chuen, China.

- Achilognathus guichenoti, Blkr., redescribed by Steindachner, Denk. Ak. Wien, lix, 1892, p. 367. A. coreanus, n. sp., Steindachner, Anz. Ak. Wien, 1892, p. 133, and t. c. p. 366, Corea.
- Opsariichthys bidens, Gthr., redescribed by STEINDACHNER, t. c. p. 368.
- Chanodichthys affinis, n. sp., VAILLANT, Bull. Soc. Philom. (8) iv, 1892, p. 127, Upper Tonkin.
- Leucaspius delineatus, Sieb.: on malformed specimens; KNAUTHE, Zool. Anz. xvi, 1893, p. 109.
- Homaloptera fimbriata, Gthr., figured by GÜNTHER, in Pratt's "Snows of Tibet," pl. iii, fig. A. H. abbreviata, n. sp., id. t. c. p. 248, pl. iii, fig. B, Province of Sze-Chuen, Chins.
- Nemachilus sturanyi, n. sp., STEINDACHNER, Anz. Ak. Wien, 1892, p. 131, and Denk. Ak. Wien, lix, 1892, p. 378, pl. ii, fig. 3, Macedonia.
- Botia pratti and superciliaris, n. spp., GUNTHER, in Pratt's "Snows of Tibet," p. 250, pl. iv, Province of Sze-Chuen, China.
- √ Eucirrichthys, n. g., allied to Acanthophthalmus and Apua. Body cylindrical, elongate, slender; mouth with pendant lip; eight barbels at the end of the snout, six of which are maxillary; a minute spine below the eye; dorsal far back, with the last ray just above the first anal ray. E. doriæ, n. sp., Perugia, Ann. Mus. Genov. (2) xii, 1892, p. 1009, Sarawak.

#### CHABACINIDÆ.

- Curimatus nitens, p. 184, and conspersus, p. 185, n. spp., Holmberg, Rev. Argent. i, 1891, Rio Paraguay.
- Prochilodus platensis, n. sp., Holmberg, t. c. p. 186, Rio de la Plata, Buenos Ayres.
- Leporinus solarii, Holmb., redescribed by Holmberg, t. c. p. 187.
- Alestes brevipinnis, n. sp., Perugia, Ann. Mus. Genov. (2) x, 1892, p. 975, Congo.
- Nannæthiops unitæniatus, Gthr., has ctenoid scales; VAILLANT, Bull, Soc. Philom. (8) v. 1893, p. 13, fig.
- Brycon (Chalcinopsis) alburnus, Gthr., and dentex, Gthr., redescribed by STEINDACHNER, Denk. Ak. Wien, lix, 1892, pp. 374 & 375.
- Tetragonopterus (Hemibrycon) polyodon, Gthr., redescribed, id. t. c. p. 375.

  T. lineatus, Stdr., = T. lineatus, Perugia; T. anomalus, Stdr., = T. nigripinnis, Perugia, id. t. c. p. 376.
- T. correntinus, p. 188, Rio Parana, Corrientes, and erythropterus, p. 189, Rio de la Plata, n. spp., Holmberg, Rev. Argent. i, 1891. T. moorii, n. sp., Boulenger, Ann. N. H. (6) x, 1892, p. 11, pl. ii, fig. 1, Matto Grosso, Brazil.
- √ Brachychalcinus, n. g., intermediate between Tetragonopterus, Cuv., and Luctkenia, Stdr., but with a movable spine, directed forward, in front of the dorsal fin; Boulenger, t. c. p. 11. B. retrospina, n. sp., id. t. c. p. 12, pl. ii, fig. 2, Matto Grosso, Brazil.
  - Piabuca melanostoma, n. sp., Holmberg, t. c. p. 192, Argentina.

- Myletes mesopotamicus, Holmb., redescribed; Holmberg, Rev. Argent. i, 1891, p. 193.
- Bergia altipinnis, Stdr., = Pseudocorynopoma doriæ, Perugia; Boulen-Ger, Ann. N. H. (6) x, 1892, p. 12; Steindachner, Denk. Ak. Wien, lix, 1892, p. 376.
- ✓ Chalcinopelecus, n. g. [= Pseudocorynopoma, Perugia.—Rec.], for C. argentinus, n. sp., Holmberg, t. c. p. 190, Argentina.

## CYPRINODONTIDE.

- Characadon variatus, Bean: note by BEAN, P. U. S. Nat. Mus. xv, 1892, p. 286.
- Haplochilus dayi, n. sp., STEINDACHNER, Anz. Ak. Wien, 1892, p. 131, and Denk. Ak. Wien, lix, 1892, p. 376, pl. i, fig. 2, Ceylon.
- Fundulus pallidus, n. sp., EVERMANN, Bull. U. S. Fish Comm. f. 1891, p. 84, pl. xxxv (1892), Texas. F. robustus, n. sp., BEAN, P. U. S. Nat. Mus. xv, 1892, p. 285, pl. xliv, fig. 2, Mexico.
- Zygonectes funduloides, p. 85, pl. xxxv, fig. 3, pulvereus, p. 85, pl. xxxvi, fig. 1, and jenkinsi, p. 86, pl. xxxvi, fig. 2, n. spp., EVERMANN, t. c., Texas.

# GALAXIIDÆ.

Galaxias nigothoruk, n. sp., Lucas, P. R. Soc. Vict. (2) iv, 1892, p. 28, Lake Nigothoruk, Gippaland, Victoria.

#### STERNOPTYCHIDÆ.

- Diplophos tania, Gthr., redescribed and figured by LUTKEN, Vid. Selsk. Skr. (6) vii, 1892, p. 278, pl. ii, figs. 1-3.
- Gonostoma microdon, Gthr. (?): note on, and figure of a specimen referred to this species; LÜTKEN, t. c. p. 280, pl. iii, figs. 4 & 5. G. elongatum, Gthr.: note by ALCOCK, Ann. N. H. (6) x, 1892, p. 354.
- Chauliodus pammelas, n. sp., ALCOCK, t. c. p. 355, Laccadive Sea, 1370 faths.

## STOMIATIDÆ.

Astronesthes, Rich.: remarks on the three species of this genus; Lütken, Vid. Selsk. Skr. (6) vii, 1892, p. 273.

Photostomias guernei, Coll.: note by Lütken, t. c. p. 280, fig.

#### SALMONIDÆ.

- H. Blanc, Bull. Soc. Vaud. (3) xxvii, 1892, pp. 272-275, has a preliminary note on the maturation and fecundation of Trout ova.
- O. STRACHLY, MT. Embr. Inst. Wien (2) v, 1892, pp. 20-22, has a note on abnormal development of the Trout ovum.

Salmo fario, L.: R. TRAQUAIR, Ann. Scott. N. H. i, 1892, p. 92, pls. iii-v, describes and figures malformed specimens from Scottish waters. S. ohridanus, n. sp., Steindachner, Anz. Ak. Wien, 1892, p. 132, and Denk. Ak. Wien, lix, 1892, p. 23, pl. iii, figs. 1-3, Macedonia.

Coregonus coulterii, n. sp., Eigenmann, Am. Nat. xxvi, 1892, p. 961, Rocky Mountains.

## PERCOPSIDE.

Remarks by Eigenmann, Science, xx, 1892, p. 233.

V Columbia, n. gen., allied to Percopsis, Ag., but dorsal and anal each with two very strong spines, scales most strongly ctenoid on anterior part of body, and posterior margin of preopercle with a few short but strong spines. C. transmontana, n. sp., id. t. c. p. 234, Umatilla, Columbia River.

#### CLUPEIDE.

Clupea pilchardus, L.: on its growth; J. T. Cunningham, Nature, xlv, 1892, pp. 255-558; M. Dunn, t. c. p. 511: on its growth in the Mediterranean; Marion, CR. Ass. Fr. Sci. 1891, p. 228: on its reproduction; J. T. Cunningham, Rep. Brit. Ass. 1891, p. 686 (1892).

V Hyperlophus, n. g., distinguished by the dorsal serrature, for Clupea spratelloides, n. sp., DOUGLAS-OGILBY, Rec. Austral. Mus. ii, 1892, p. 24, Coast of New South Wales. A. S. WOODWARD, Ann. N. H. (6) x, 1892, p. 412, points out that this doubly-armed herring falls in the genus Diplomystus, Cope, established for Cretaceous and early Tertiary species.

Pellonula vorax, Gthr.: note by Perugia, Ann. Mus. Genov. (2) x, 1892, p. 977.

#### HALOSAURIDÆ.

Halosaurus parvipennis, n. sp., Alcock, Ann. N. H. (6) x, 1892, p. 362, Laccadive Sea, 865-880 faths.

#### SAUBODONTIDÆ.

Ichthyodectes anaides, Cope (foss.): notes by Crook, Palseontogr. xxxix, 1892, p. 111, pl. xv. I. polymicrodus, n. sp. (foss.), id. t. c. p. 112, pl. xvi, Chalk of Kansas.

Portheus molossus, Cope: notes by CROOK, t. c. p. 114, pl, xviii.

#### MURÆNIDÆ.

D. S. Jordan & B. M. Davis, Rep. U. S. Fish. Comm. f. 1888, pp. 581-677, pls. lxxiii-lxxx (1892), review the Apodal Fishes inhabiting the waters of America and Europe,

- G. B. Grassi & S. Calandruccio, Atti Acc. Pontif. Lincei (5) i, 1892, Rend. pp. 375-379, write on the transformation of Leptocephali into Mursenoids.
- Anguilla rulgaris, Cuv.: on an extraordinary passage of Eels in the Main; Buxbaum, Zool. Gart. xxxiii, 1892, p. 20: on its breeding season; Zenk, t. c. p. 61.
- Uroconger vicinus, Vaill.: note on the stomach; Alcock, Ann. N. H. (6) x, 1892, p. 363.
- ✓ Rabula, n. subg. for the species of Gymnothorax which have the dorsal fin inserted behind the head; JORDAN & DAVIS, t. c. p. 590.
- V Quassiremus, n. g. for Ophichthys evionthias, Jord. & Bollm.; JORDAN & DAVIS, t. c. p. 622.
- No Scytalichthys, n. subg. for Ophichthys miurus, Jord. & Gilb.; JORDAN & DAYIS, t. c. p. 635.
  - Nettastoma melanurum, Raf.: note on its larval (Leptocephulus longirostris, Kaup; Hyoprorus messanensis, Küll.) and adult characters; Facciolà, Nat. Sicil. xi, 1893, p. 144.
  - Gavialiceps microps, Alcock (? = Nemichthys infans, Vaill. nec Gthr.): further notes by Alcock, t. c. p. 361.
- √ Venefica, n. g. for Nettastoma procerum, Goode & Bean; JORDAN & DAVIS, t. c. p. 651.
- V Avocettina, n. g. for Nemichthys infans, Gthr.; JORDAN & DAVIS, t. c. p. 655.
- "Ilyophis, n. g., type of a new family, Ilyophididæ, intermediate between Simenchelys and Synaphobranchus; JORDAN & DAVIS, t. c. p. 670.

  I. brunneus, n. sp., iid. t. c. p. 671, Galapagos Is.
- Ahlia, n. g. for Myrophis egmontis, Jord.; Jordan & Davis, t. c. p. 639.
   Myrophis frio, n. sp., Jordan & Davis, t. c. p. 640, Coast of Brazil.
- Gordiichthys, n. g., allied to Muranesox, JORDAN & DAVIS, t. c. p. 644;
  G. irreritus, n. sp., iid. t. c., Gulf of Mexico.
- Xenomystax, n. g., allied to Muranesox; JORDAN & DAVIS, t. c. p. 648.
  X. atrarius, n. sp., iid. t. c. p. 649, off Coast of Ecuador, 401 faths.
  - Chlopsis equatorialis, n. sp., JORDAN & DAVIS, t. c. p. 650, off Coast of Ecuador, 401 faths.

# PLECTOGNATHI.

- Balistes aureolus, Richards., redescribed by Herzenstein, Mél. biol. xiii, 1892, p. 233.
- Ostracion meretrix, n. sp. (foss.), Daimeries, Bull. Soc. Mal. Belg. (4) vi, 1891, p. lxxv, Eocene, Brussels.
- T. GILL, P. U. S. Nat. Mus. xiv, 1892, pp. 705-720, pl. xxxiv, has notes on the Tetrodons, which group he raises to the rank of a superfamily, Tetraodontoidea, divided into three families, viz.: Chonerhinidæ, Canthigasteridæ, and Tetraodontidæ; the latter is again divided into two subfamilies, viz., Tetraodontinæ and Colomesinæ.
- Orthagoriscus nasus, Ranz., described and figured by VAN LIDIH DE JEUDE, Notes Leyd. Mus. xiv, 1892, p. 127, pl. v.

#### GANOIDEI.

- G. S. HOPKINS gives an abstract of a paper on the digestive tract of some N. American Ganoids; P. Am. Ass. xli, 1892, p. 197.
- L. SCHMIDT, Z. wiss. Zool. liv, 1892, pp. 748-764, figs., pl. xxxiv, deals with the morphology of the vertebral column of Amia calca.
- Amia whiteavesiana and macrospondyla, n. spp. (foss.), Cope, Contr. Canad.
  Pal. iii, 1891, p. 2, pl. i, figs. 1 & 2, Lower Miocene, Canada. A.
  kehreri, n. sp. (foss.), Andreae, Verh. Ver. Heidelb. (2) v, 1893, p. 9,
  Lower Miocene, Messel, near Darmstadt.
- Caturus, Ag. (foss.): notes by SAUVAGE, Bull. Soc. Autun, iv, 1891, pp. 63-81, pls. vi-ix. C. gigas, p. 74, pl. vii, and retrodorsalis, p. 79, pl. ix, n. spp., id. ibid., Upper Lias, Yonne, France. C. chaperi, Sauvg., and stenoura, Sauvg., figured, pls. vi & viii.
- Leptolepis affinis, p. 389, pl. xvi, and autissiodorensis, p. 393, pl. xvi, n. spp., SAUVAGE, Bull. Soc. Autun, v, 1892, Upper Lias, Yonne, France.
- VLepidopterus, n. g., allied to Lepidotus, for L. crassus, n. sp. (foss.);
  POILIG, Leuckart Festschrift, 1892, p. 63, pl. vii, fig. 3, Permian of Friedrichroda, Thuringia.
  - Belonostomus sweeti, n. sp. (foss.), R. ETHERIDGE, JUN., & A. S. WOODWARD, Tr. R. Soc. Vict. ii, 1892, p. 1, pl. i, Cretaceous of Central Queensland.
  - Pholidophorus gaudryi, n. sp., SAUVAGE, Bull. Soc. Autun, v, 1892, p. 398, pl. xvii, Upper Lias, Yonne, France.
  - Otenolepis cyclus, Ag. (foss.): figures of typical specimens; A. S. WOODWARD, P. Geol. Ass. xii, 1892, pl. iv, figs. 5-8.
  - Lepidosteus suessoniensis, Gerv. (foss.), is a Fish, not a Reptile as stated by Lemoine; Dollo, Bull. Sci. Fr. Belg. xxiv, 1892 [quoted from separate copy].
  - Tetragonolepis, Bronn (foss.), is regarded as the type of a new family, Tetragonolepidæ, by REIS, Geogn. JH. iv, 1892, p. 160.
- Colobodus, Ag. (foss.): remarks by Montagu Brown, Rep. Brit. Ass. 1891, p. 644 (1892).
- Microdus dumbelii, n. sp. (foss.), COPE, P. Am. Phil. Soc. xx, 1892, p. 128, Lower Cretaceous (?) of Texas.
- Mesodon rugulosus, Ag. (foss.): note by A. S. WOODWARD, P. Geol. Soc. xii, 1892, p. 239, pl. iv, figs. 2-4.
- Dorypterus, Germ. (foss.); notes by Reis, Geogn. JH. iv, 1892, p. 167.
- Acipenser, L: on the Sturgeons of the East Coast of the United States; RYDER, Bull. U. S. Fish Comm. viii, 1890, p. 231, pls. xxxvii-lix. A. ruthenus, L.: on a remarkable new variety: A. v. Mojsisovics, SB. Ak. Wien, ci, i, 1892, p. 328, pl.—. A. mikadoi, n. sp., HILGENDORF, SB. nat. Fr. 1892, p. 98, N. Japan.
- Myriolepis hibernicus, n. sp. (foss.), TRAQUAIR, Geol. Mag. (3) x, 1893, p. 54, pl. iii, Coal Measures, co. Kilkenny, Ireland.
- Amblypterus traquairi, n. sp. (foss.), Pohlig, Leuckart Festschrift, 1892, p. 63, pl. vii, fig. 4, Permian of Friedrichroda, Thuringia.

- O. M. Reis, Geogn. JH. iv, 1892, p. 157, discusses the systematic position of the Belonorhynchida.
- Saurichthys striolatus, Bronn, apicalis, Ag., and mougeoti, Ag. (foss.); notes, id. t. c. p. 148, figs. S. gypsophilus, n. sp., id. t. c. p. 153, fig., Keuper of Franconia.
- √ Saurorhynchus, n. g. for Belonostomus acutus, Ag. (foss.), Reis, t. c. p. 145.

# CROSSOPTERYGII.

- H. B. POLLARD, Morph. JB., Anat. v, 1892, pp. 387-428, figs., pls. xxvii-xxx, has a paper on the anatomy and phylogenetic position of *Polypterus*, which is regarded as representing the ancestral form of the Stegocephalous Batrachians.
- M. Reis, Inaugural Dissertation, Munich, 1892 (cf. p. 11), describes the osteology of the Calacanthidae.
- Onychodus scoticus, n. sp., E. T. NEWTON, Geol. Mag. (3) ix, 1892, p. 51, Old Red Sandstone of Forfar, Scotland.
- Gyroptychius heddlei, n. sp., TRAQUAIR, Ann. Scott. N. H. i, 1892, p. 233, Lower Old Red Sandstone, Dura Den, Scotland.
- Megalichthys macropomus, n. sp. (foss.), Cope, P. Am. Phil. Soc. xxx, 1892, p. 226, pl. viii, Carboniferous, Kansas.
- Holoptychius? filosus, n. sp. (foss.), COPE, t. c. p. 228, pl. vii, fig. 3, Devonian, Pennsylvania.

# DIPNOI.

- C. Röse, Anat. Anz. vii, 1892, pp. 821-839, figs., writes on the teeth of the Dipnoi.
- C. Hasse, Z. wiss. Zool. lv, 1893, pp. 533-542, pl. xxii, deals with the development of the vertebral column in the Dipnoi.
- B. Dean, Rep. U. S. Fish. Comm. xix, 1892, pp. 309-314, writes on the pineal fontanelle of *Dinichthys*, and compares the Placoderms and Siluroids.
- P. Burckhard brings out a work on the Central Nervous System of the *Dipnoi* (Berlin: 1892, 8vo, 64 pp., 4to atlas, 5 pls.). Abstracts of his researches in Verh. deutsch. Zool. Ges. ii, 1892, pp. 92-95, and SB. nat. Fr. 1892, pp. 23-25.
- R. Dubois, Ann. Soc. L. Lyon (2) xxxix, 1892, pp. 65-72, figs., writes on the mechanism of the respiration in the *Dipnoi* and the passage from the torpid to the active condition.
- Ceratodus forsteri, Krefft: notes on the habits; W. B. Spencer, Pr. R. Soc. Vict. (2) iv, 1892, p. 81. C. (Meglapleuron) rochei, Gaudry (foss.): note on its scales; Vaillant, C.R. cxiv, 1892, p. 1083.
- Coccosteus macromus, n. sp. (foss.), Cope, P. Am. Phil. Soc. xxx, 1892, p. 225, Devonian, Pennsylvania. C. canadensis, n. sp. (foss.), A. S. Woodward, Geol. Mag. (3) ix, 1892, p. 483, pl. xiii, fig. 2, Upper Devonian, Scaumenac Bay, Canada. C. cuyahoga, n. sp. (foss.), CLAYPOLE, Am. Geol. xi, 1893, p. 167, figs., Devonian of Ohio.

- Aspidichthys (?) nobilis, n. sp. (foss.), WHITEAVES, Contr. Can. Pal. i, pt. iv, p. 354, pl. xlvii, fig. i, Devonian, South Manitou L, Lake Winnipeg.
- Phlyctenaspis acadica, Whiteaves: notes and figures by A. S. Woodward, Geol. Mag. (3) ix, 1892, p. 5, pl. i, figs. 7 & 8, and p. 481, fig., and by Traquair, op. cit. x, 1893, p. 147.
- Dinichthys, Newb. (foss.): description of the skull, by CLAYPOLE, Am. Geol. x, 1892, p. 199, figs. D. canadensis, n. sp., WHITEAVES, Contr. Canad. Pal. i, pt. iv, 1892, p. 353, pl. xlvi, fig. 12, Devonian, Snake I., Lake Winnipeg.
- √ Gorgonichthys, n. g., allied to Dinichthys. G. clarki, n. sp. (foss.), CLAYPOLE, Am. Geol. x, 1892, p. 2, fig., Devonian of Ohio.
  - Ganorhynchus oblongus, n. sp. (foss.), COPE, P. Am. Phil. Soc. xxx, 1892, p. 225, Devonian, Pennsylvania.
  - A. S. WOODWARD, Ann. N. H. (6) xi, 1893, pp. 241 & 242, fig., describes a case of subdivision of the median fin in the fossil Dipnoan *Phaneropleuron curtum*.

# OSTRACODERMI.

A. S. WOODWARD, Nat. Sci. i, 1892, pp. 596-602, figs., has an article on the Earliest Fishes (*Palæospondylus* and *Ostracodermi*).

# ANTIARCHA.

- B. H. TRAQUAIR, P. Phys. Soc. Edinb. 1891-92, pp. 283-286 (1893), offers remarks on the British species of Asterolepidæ.
- Cephalaspis campbelltownensis, Whiteaves: note and figure by A. S. Woodward, Geol. Mag. (3) ix, 1892, p. 4, pl. i, fig. 6: note by Traquair, op. cit. x, 1893, p. 146. C. jexti, n. sp., id. t. c. p. 147, Lower Devonian, Campbelltown, New Brunswick.
- Holonema rugosa, Clayp.: notes by Cope, P. Am. Phil. Soc. xxx, 1892, p. 223. H. horrida, n. sp., id. t. c. p. 222, pl. vii, fig. 2, Devonian, Pennsylvania.
- Bothriolepis minor, Newb.: note by COPE, t. c. p. 224. B. canadensis, Whiteaves: note on the supposed jaws; A. S. WOODWARD, Geol. Mag. (3) ix, 1892, p. 484, fig. B. leptochirus, n. sp., TRAQUAIR, P. Phys. Soc. Edinb. 1891-92, p. 286 (1893), Upper Old Red Sandstone of Heads of Ayr, Scotland.

## OSTEOSTRACI.

- J. V. Rohon discusses the systematic position of the families Thyestidas (n. n.) and Tremataspidae, and recharacterises the genera Thyestes, Eichw., and Tremataspis, F. Schmidt.; Mém. Ac. Pétersb. (7) xxxviii, 1892, No. 13, p. 84.
- Tremataspis schmidti, p. 39, mickwitsi, p. 56, and simonsoni, p. 63, n. spp., Rohon, t. c., Upper Silurian, Isle of Oesel, Baltic.
- Thyestes verrucosus, Eichw.: desord. & figrd. by Bohon, t. c. p. 15, pls. i & ii. Chalcodus permianus, Zitt., = Menaspis armata, Jackel; JAEKEL, SB. nat. Fr. 1892, p. 157.

#### HETEROSTRACI.

Palæaspis, Claypole: notes and restoration by CLAYPOLE, Q. J. Geol. Soc. xlviii, 1892, p. 542, figs.

∨ Tolypaspis, n. n. for Tolypelepis, Pander; F. SCHMIDT, JB. Mineral.
1893, i, p. 100.

# HOLOCEPHALL

- Ischyodus emarginatus, Egert. (foss.): note by A. S. WOODWARD, P. Geol. Ass. xii, 1892, p. 238, pl. iv, fig. 1.
- Packymylus, n. g., for P. leedsii, n. sp. (foss.), founded on teeth from the Oxford Clay of Peterborough; A. S. WOODWARD, Ann. N. H. (6) x, 1892, p. 14, pl. iii, figs. 1 & 2.
  - v Brackymylus, n. g., A. S. WOODWARD, t. c. p. 14. B. altidens and minor, n. spp. (foss.), Oxford and Kimmeridge Clays, Peterborough and Weymouth.
    - Elasmodectes secans, n. sp. (foss.), A. S. Woodward, t. c. p. 16, pl. iii, fig. 3, Kimmeridge Clay, Weymouth.

# ELASMOBRANCHII.

# ACANTHODII.

On the systematic position of the Acanthodii; REIS, SB. nat. Fr. 1892, pp. 153-156, and JAEKEL, t. c. pp. 156-158.

# SELACHII.

- A. S. WOODWARD, Nat. Sci. i, pp. 671-675, figs., discusses the evolution of Sharks' teeth.
- L. EDINGER, Abh. Senck. Ges. xviii, 1892, 55 pp., 5 pls., has a memoir on the brain of Selachians. Abstract in Anat. Anz. vii, 1892, pp. 472– 476.
- H. Braus publishes researches on the rami ventrales of the anterior spinal nerves of Selachians. Jena: 1892, 8vo, 35 pp.
- A. SEDGWICK, Q. J. Micr. Sci. xxxiii, 1892, pp. 559-586, pl. xxxv, has notes on Elasmobranch development.
- S. Mollier, Anat. Anz. vii, 1892, pp. 351-365, has a preliminary note on the development of the limbs in Selachians.
- C. HASSE, Z. wiss. Zool. lv, 1892, pp. 519-531, pl. xxi, publishes a contribution to the development of the vertebral column of the Selachians.
- G. Antipa, Anat. Anz. vii, 1892, pp. 690-692, figs., remarks on the relations of the Thymus to the so-called branchial cleft organs in Selachians.
- C. Rabl, Leuckart Festschrift, 1892, pp. 228-235, figs., writes on the development of the venous system in Selachians.

- F. RAFFAELE, MT. z. Stat. Neap. x, 1892, pp. 441-479, pls. xxix-xxxi, investigates the development of the vascular system in Selachians.
- A. Dohrn, Anat. Anz. vii, 1892, pp. 348-351, remarks on the nervecells of Selachian embryos.
- C. K. HOFFMANN, Morph. JB. xix, 1893, pp. 592-648, figs., pls. xxii-xxv, deals with the embryology of the heart and blood-vessels in the Selachians.
- J. RÜCKERT, Anat. Anz. viii, 1892, pp. 44-52, has a further contribution to the segmentation of the Selachian ovum.
- J. Bückert, op. cit. vii, 1892, pp. 107-158, figs., has a paper on the development of the ovarian egg in Selachians.
- F. WERNER, Morph. JB., Syst. vi, 1892, pp. 220-224, pl. x, discusses the signification of the markings of Selachians.

#### ASTEROSPONDYLI.

- G. C. Purvis, P. Phys. Soc. Edinb. 1890-91, pp. 62-67, pl. (1892), writes on the pineal gland of Lamna cornubica.
- M. V. Lenhossék, Anat. Anz. vii, 1892, pp. 519-539, figs., has observations on the spinal ganglia and spinal marrow in embryos of Pristiurus.
- Curcharodon mortoni, Gibbes, = C. megalodon, Ag. (foss.); F. A. Lucas, P. Biol. Soc. Washington, vii, 1892, p. 151.
- Galeus lefevrei, n. sp. (foss.), DAIMERIES, Bull. Soc. Mal. Belg. (4) vi, 1891, p. lxxiv, Eocene, Brussels.

# TECTOSPONDYLI.

- J. C. EWART has a further paper on the electric organ of the Skate; Phil. Tr. clxxxiii (B), 1892, pp. 389-420, pls. xxvi-xxx. Abstract, P. R. S. l, 1892, pp. 474-476.
- J. C. EWART, Zool. Anz. xv, 1892, pp. 116-120, writes on the sensory canals of Lamargus and Raia batis.
- J. BEARD, Anat. Anz. vii, 1892, pp. 191-206, figs., has a paper on the transient ganglion cells and their nerves in Raia batis.
- A. Alcock, Ann. N. H. (6) ix, 1892, pp. 417-427, pl. xix, and x, 1892, pp. 1-8, pl. iv, describes the gravid uterus and feetus of *Trygon bleekeri* and *Pteroplatea micrura*.
- J. WOOD-MASON & A. ALCOCK, P. R. Soc. l, 1892, pp. 202-209, record further observations on the gestation of Indian Rays.
- C. K. HOFFMANN, Anat. Anz. vii, 1892, pp. 270-273, figs., has a note on the early development of the heart and blood-vessels in Acanthias vulgaris.
- H. E. & F. Ziegler, Arch. mikr. Anat. xxxix, 1892, pp. 56-102, figs., pls. iii & iv, publish contributions to the embryology of Torpedo ocellata.
- MARCY, C.R. cxvi, 1893, pp. 77-81, figs., has observations on the natation of the Skate.

- Acanthias vulgaris, Risso: on a dicephalous embryo; GADEAU DE KERVILLE, J. de l'Anat. Phys. xxviii, 1892, p. 563, pl. xviii, figs. 1 & 2.
- Centrina salviani, Risso: note by W. L. CALDERWOOD, J. Mar. Biol. Ass. (2) ii, 1892, p. 322, pl. xiii.
- Sclerorhynchus atarus, A. S. Woodw. (foss.), redescribed and figured by A. S. Woodward, Geol. Mag. (3) ix, 1892, p. 529, pl. xiv.
- Dichelodus acutus, Gieb. (foss.): remarks by JAEKEL, JB. Mineral. 1892, i, p. 148.
- Raia clavata, L.: on an abnormally coloured example; TRAQUAIR, Ann. Scott. N. H. ii, 1893, p. 25, pl. i.
- Trygon pastinaca, L., var. marmorata, Stdr., figured by STEINDACHNER, Denk. Ak. Wien, lix, 1892, pl. iii, fig. 1. T. navarræ and simensis, n. spp., STEINDACHNER, Anz. Ak. Wien, 1892, pp. 132 & 133, and t. c. pp. 381 & 382, pls. v & vi, Shanghai.

#### ICHTHYOTOMI.

- Claulodus, Ag.: remarks by JAEKEL, SB. nat. Fr. 1892, p. 80.
- Pleuracanthidæ: on the remains from the British Coal Measures; J. W. DAVIS, Sci. Tr. R. Dublin Soc. (2) iv, 1892, pp. 703-748, pls. lxv-lxxiii.
- Pleuracanthus undulatus, p. 733, pl. lxxii, fig. 16, Burnley, howsei, p. 735, pl. lxxii, fig. 21, Newcastle-on-Tyne, thomsoni, p. 742, pl. lxxiii, fig. 17, Kilmarnock, obtusus, p. 742, pl. lxxiii, fig. 18, Clifton, Yorkshire, serratus, p. 744, pl. lxxiii, figs. 19 & 20, Newcastle-on-Tyne, and woodwardi, p. 744, pl. lxxiii, fig. 21, Dalkeith, n. spp., id. ibid.
- Diplodus problematicus, n. sp. (foss.), A. S. WOODWARD, Geol. Mag. (3) ix, 1892, p. 2, pl. i, fig. 2, Lower Devonian, Campbelltown, New Brunswick.
- √Doliodus, n. g. for the above species; TRAQUAIR, op. cit. x, 1893, p. 145.

# INCERTÆ SEDIS.

- JAEKEL, JB. Mineral. 1892, i, pp. 145-151, figs., in reply to some criticisms of A. S. Woodward, enters into details on the structure of various Ichthyodorulites.
- Acanthodes semistriatus, n. sp., A. S. Woodward, Geol. Mag. (3) ix, 1892, p. 3, pl. i, fig. 3, Lower Devonian, Campbelltown, New Brunswick.
- Gyracanthus incurvus, Traq.: note and figure by A. S. WOODWARD, t. c. p. 2, pl. i, figs. 4 & 5.
- Diplacanthus horridus, n. sp., A. S. WOODWARD, t. c. p. 482, pl. xiii, fig. 1, Upper Devonian, Scaumenac Bay, Canada.
- Cheiracanthus? costellatus, n. sp., TRAQUAIR, op. cit. x, 1893, p. 146, Devonian of Canada.
- Protodus, n. g., for Elasmobranch teeth representing a specialized form of a very simple type; A. S. WOODWARD, op. cit. ix, 1892, p. 1. P. jezi, n. sp., id. t. c. p. 2, pl. i, fig. 1, Lower Devonian, Campbelltown, New Brunswick. Remarks by Traquair, op. cit. x, 1893, p. 145.

# CYCLOSTOMI.

- J. Beard, Anat. Anz. viii, 1892, pp. 59 & 60, gives an abstract of his observations on small specimens of Myxine glutinosa, and on the hermaphroditism of Lampreys.
- B. H. BURNE, P. Z. S. 1892, pp. 706-708, pl. xlvii (1893), writes on the presence of a branchial basket in Myxine glutinosa.
- G. RETZIUS, Biol. Fören. iii, 1892, pp. 81–84, pl. iii, describes the caudal skeleton of Myxine glutinosa.
- G. RETZIUS, op. cit. iv, 1892, pp. 36-41, figs., writes on the termination of the spine and notochord in the tail of Petromyzon fluviatilis.
- On the nerve-terminations in the skin of *Petromyzon*; G. RETZIUS, Biol. Unters. (2) iii, 1892, pp. 37-40, pl. xiii.
- B. HATSCHEK, Anat. Anz. viii, 1892, pp. 89-91, and Verh. Anat. Ges. 1892, pp. 147-161, figs., supplements and corrects some of his observations on Metamery in Ammocates.
- P. Owsjannikow's paper on the embryology of Petromyzon fluviatilis is translated in Ann. N. H. (6) xi, 1893, pp. 34-43.
- B. Lwoff, Biol. Centralbl. xiii, 1893, pp. 45-48, summarises his researches on the segmentation of the *Petromyzon* ovum.
- R. ALCOCK, P. Cambr. Phil. Soc. vii, 1892, pp. 252-255, writes on the digestive processes of Ammocates.
- E. CAVAZZANI, Arch. Ital. Biol. xviii, 1892, pp. 182-186, relates his experiments on the poisonous properties of the blood of Petromyzon marinus.
- Lampetra spadicea, Bean, described and figured by Bean, Nat. Mex. (2) ii, 1892, pp. 171 & 172, pl. viii: note by Bean, P. U. S. Nat. Mus. xv, 1892, p. 283.

# LEPTOCARDII.

- J. W. VAN WIJHE, Anat. Anz. viii, 1893, pp. 152-172, has notes on the morphology of Amphioxus.
- B. HATSCHEK, Anat. Anz. viii, 1892, pp. 89-91, and Verh. Anat. Ges. 1892, pp. 136-147, figs., writes on Metamery.
- E. ROHDE, SB. Ak. Berlin, 1892, pp. 659-664, compares the muscles and nerves of *Amphioxus* with those of *Mermis*.
- J. B. Platt, Anat. Anz. vii, 1892, pp. 282-284, figs., writes on fibres connecting the central nervous system and chords.
- T. BOVERI, Morph. JB., Anat. v, 1892, pp. 429-510, figs., pls. xxxi-xxxiv, deals with the excretory tubules of the urogenital system.
- T. Boveri, Anat. Anz. vii, 1892, pp. 170-181, figs., writes on the genital glands.
- E. B. WILSON, t. c. pp. 732-740, figs., writes on multiple and partial development.
- B. Danilewsky, Arch. ges. Phys. li, 1892, pp. 393-400, writes on the physiology of the central nervous system.
- Branchiostoma elongatum, Sund., recorded from California by EIGEN-MANN, Am. Nat. xxvi, 1892, p. 70.

# VI. TUNICATA.

BY

# PROFESSOR W. A. HERDMAN, D.Sc., F.R.S.

# I.—LIST OF PUBLICATIONS.

- BLES, E. J. Notes on Plankton observed at Plymouth during June— September, 1892. J. Mar. Biol. Ass. (n.s.), ii, No. 4, Nov. 1892, pp. 340-343.
- BUTSCHLI, O. Einige Bemerkungen über die Augen der Salpen.
   Zool. Anz. xv, Jhg., No. 401, pp. 349-353.
- 3. Garstang, W. Attempt to elucidate the real structure and relations of Moss's Polystigmatic Appendicularian. Tr. Biol. Soc. Liverp. vi, pp. 57-69, pl. v.
- On the development of the stigmata in Ascidians. P. R. Soc. li, pp. 505-513.
- GÖPPERT, E. Untersuchungen ü. d. Sehorgan d. Salpen. Morph. JB. 1892, pp. 250-294, pls. viii-x.
- HERDMAN, W. A. Notes on the structure of Oikopleura. Tr. Biol. Soc. Liverp. vi, pp. 40-56, pls. i-iv.
- Note on the Geographical distribution of Tunicata. Brit. Assoc. Rep. for 1892 (Edinburgh meeting), pp. 787 & 788.
- Note on Atrial Tentacles in Ascidians. Brit. Assoc. Rep. for 1892 (Edinburgh meeting), pp. 788 & 789.
- Notes on the Collections made during the Cruise of the s.y.
   "Argo" up the West Coast of Norway in July, 1891. Tr. Biol.
   Soc. Liverp. vi, pp. 70-93, pls. vi & vii.
- 10. —. A functional Hermaphrodite Ascidian. Nature, xlvi, p. 561.
- HJORT, JOHAN. Zum Entwicklungscyclus der zusammengesetzten Ascidien. Zool. Anz. xv, No. 400, pp. 328–332.
- JOURDAIN, S. De la déglutition chez les Synascidies. Bull. Soc. Philom. iv, No. 1, pp. 35 & 36.
  - 1892. [vol. xxix.]

- Julin, Ch. Les Ascidiens des côtes du Boulonnais. I. Recherches sur l'anatomie et l'embryogénie de Styelopsis grossularia—Introduction. Bull. Sci. xxiv, pp. 1-52.
- KOWALEVSKY, A. Einige Beiträge zur Bildung des Mantels des Ascidien. Mém. Ac. Pétersb. vii, xxviii, No. 10, pp. 1-20, pls, i & ii.
- LAHILLE, F. Classification des Tuniciers (groupes primordiaux).
   Naturaliste, 1 Mars, 1892, 12 pp.
- LOHMANN, H. Vorbericht über die Appendikularien der Plankton-Expedition, in Reisebeschreibung der Plankton-Expedition von Dr. Otto Krümmel. Kiel u. Leipzig, 1892, pp. 139-149.
- METCALF, MAYNARD M. The Anatomy and Development of the Eyes and Subneural Gland in Salpa. Johns Hopk. Univ. Circ. xi, No. 97, pp. 78 & 79 (April, 1892).
- Nott, J. T. On the Composite Ascidians of the North Shore Reef. Tr. N. Z. Inst. xxiv, pp. 305-334, pls. xxiv-xxx.
- OKA, A. Die periodische Regeneration der Oberen Körperhälfte bei den Diplosomiden. Biol. Centralbl. 12 bd, Nos. 9 & 10, pp. 265-268.
- Ueber die Knospung der Botrylliden. Z. wiss. Zool. bd. 54, pp. 521-547, pls. xx-xxii.
- 21. Oka, A., and Willey, A. On a new genus of Synascidians from Japan. Q. J. Micr. Sci. xxxiii, pp. 313-323, pla xvii & xviii.
- Pizon, A. Développement de l'organe vibratile chez les Ascidies composées. C.R. cxiv, No. 5, pp. 237-239.
- Histoire de la blastogenèse chez les Botryllidés. Ann. Sci. Nat. Zool. xiv, Nos. 1, 2, 3.
- Développement du système vasculaire colonial chez les Botryllidés.
   Bull. Soc. Philom. iii, No. 4, pp. 183-186.
- SALENSKY, W. Ueber die Thätigkeit der Kalymmocyten (Testazellen) bei der Entwicklung einiger Synascidien (m. Taf. xiv u. xv).
   Festschrift zum siebenzigsten Geburtstage Rudolf Leuckarts.
   Leipzig: 1892, pp. 109-120.
- SEELIGER, O. Ueber die erste Bildung des Zwitterapparates in den jungen Pyrosomenstöcken (m. Taf. xxxviii). Festschrift zum siebenzigsten Geburtstage Rudolf Leuckarts. Leipzig: 1892, pp. 374– 384.
- Bemerkungen zu Herrn Prof. Salensky's Beiträge zur Emb. d. Pyrosomen. Zool. Jahrb. bd. v; Zool. Anz. xv, Jhg., No. 385, pp. 78-84.
- SWAINSON, G. A new form of Appendicularian "Haus." Int. J. Micr., January, 1892, 3 pp., pl. v.
- WATT, JAMES. On the structure of Boltenia pachydermatina. Tr. N. Z. Inst. xxiv, pp. 334-348, pls. xxxi-xxxiv.

- Willey, A. Observations on the Post-embryonic development of Ciona intestinalis and Clavelina lepadiformis. P. R. Soc. li, pp. 513-520.
  - ---. [See also OKA & WILLEY.]
- On the development of the hypophysis in the Ascidians. Zool. Anz. xv, No. 400, pp. 332-334.

# II.—ANATOMY, EMBRYOLOGY, AND PHYSIOLOGY.

# (a,) ANATOMY.

NOTT (18) has given a description, with some anatomical and histological detail, of eight new species of Compound Ascidians from New Zealand belonging to the genera Leptoclinum, Polysyncraton, and Cystodytes. The most important form is Polysyncraton, which presents characters intermediate between those of Leptoclinum, M.-Edw., Calocormus, Herdm., and the Diplosomidæ. Nott proposes finally to follow Jourdain in uniting the families Didemnidæ and Diplosomidæ under the term Oligosomidæ, in which he would put Polysyncraton. He gives a new diagnosis of this family Oligosomidæ.

Watt (29) gives an account of the anatomy, and some points in the histology, of *Boltenia pachydermatina*, Herdm., the large common *Boltenia* of New Zealand. He points out that this species agrees with *Culeolus* in having the heart on the right side of the body, and six folds on each side of the branchial sac.

OKA (20) gives an account of the structure of a species of Botryllus from Misaki, Japan.

BUTSCHLI (2) gives an account of the eyes of Salpa, with the view of showing that while in some species the whole eye is simple and non-inverted, in those with horse-shoe eyes the lateral parts become modified so as to be inverted, and this leads to forms with three eyes, which may be compared with the vertebrate pineal eye and pair of lateral eyes.

METCALF (17) gives a preliminary account of his investigations into the atructure and development of the eyes, subneural gland and neighbouring parts in Salpa. He finds that in the chain form of S. pinnata, in addition to the median dorsal eye, there are two pairs of much smaller eyes on the nerve-ganglion. He finds that the dorsal tubercle is innervated by a plexus of nerve cells and fibres. In the development, the cavity of the central nervous system is continuous with the lumen of the ciliated funnel of the dorsal tubercle. He appears to agree with those previous writers who regard the early function of these structures to have been probably the aeration of the brain, and the carrying off of the waste products, while the opening into the pharnyx was probably a sense organ.

GÖPPERT (5) has examined the structure of the eyes in five species of Salpa. The eye arises as a stalked body, projecting anteriorly and ventrally from the middle of the ganglion. Some of its cells become the

retina and a pigment layer; there are also in most species special groups of sensory elements without pigment. The retina may also be differentiated into two layers; and the end of the optic cell which is directed towards the light in some cases represents the nerve ending, and in other cases is the opposite end of the cell. He points out that while the solitary  $Salp\alpha$  are mostly able to look only dorsally, the chain forms are able to look out ventrally in addition. He considers that there is no direct relation between these eyes of  $Salp\alpha$  and the vertebrate eyes, except that both are developed from parts of the central nervous system.

HERDMAN (6) describes the minute structure of a species of Oikopleura from Liverpool Bay, as made out from serial sections. These show, amongst other things, the condition of the endostyle as a diverticulum to a great extent shut off from the branchial sac, the presence of a genital duct, the distribution of the enlarged ectoderm cells and the cuticular test, the exact course of the nerve cord through the posterior part of the body, and, finally, the shapes and relative positions of the alimentary and reproductive viscera.

Garstang (3), from an examination of the descriptions and figures of Moss and Fol, comes to the conclusion that Moss's remarkable polystigmatic Appendicularian (Mossia dolioloides) has no special resemblance to Doliolum but is identical with Fol's Kowalevskia. He gives a new diagnosis of the genus Kowalevskia, and describes the two species K. tenuis and K. mossi.

HERDMAN (8) points out that atrial tentacles have been known in both Simple and Compound Ascidians since 1882, and are now known to occur in four genera, all of these being cases where colonies or aggregations of individuals are formed.

Julin (13) gives the first part of a detailed memoir on the Ascidians of Boulonnais. He commences with Styclopsis grossularia as a type, and goes over a considerable part of its structure.

OKA & WILLEY (21) give a detailed account of the structure of a new Compound Ascidian allied to the genus *Didemnoides*, from Japan. It forms large red lobed colonies, has large tailed larvæ, and a thin layer of calcareous spicules in the test.

# (b.) EMBRYOLOGY.

OKA (20) gives an elaborate account of the development of the buds, and the formation of the colony, in a *Botryllus* from Japan.

SEELIGER (27) corrects some points in Salensky's account of the development of *Pyrosoma*—especially in regard to his criticisms of Seeliger's former papers.

HJORT (11) gives an account of the cycle of development in the Compound Ascidians. He finds, as Willey does also, that both in the larva, and also in the young bud, the hypophysis and the ganglion have a common origin from the neural tube, and that the neural tube opens into

the stomodseum, as originally described by Kowalevsky. Willey considers that the ganglion of the Ascidians seems to bear a similar relation to the hypophysis that the infundibulum does to the hypophysis of the higher Vertebrates. But, while in the latter case both organs are rudimentary and arise independently, in the Ascidians both organs are functional and arise together.

Pizon (22) finds in various families of the Compound Ascidians and in Clavelinidæ that the dorsal tubercle develops as a blind diverticulum from the primitive endodermal vesicle and opens secondarily into the "branchial" vesicle, while its posterior part becomes atrophied. It is not, then, a buccal invagination, and so Pizon rejects the homology with the hypophysis of Vertebrates. He considers that the dorsal tubercle is an ancestral organ, once of great importance, but now disappearing, and with no important function in the existing Tunicata.

OKA (19) finds a species of *Diplosoma*, from Japan, in which the upper half of the body is periodically renewed, the younger thorax gradually taking the place of the older one.

GARSTANG (4) has investigated the development of the stigmata in the branchial sac of Clavelina, Botryllus, Thylacium, and Styela. He finds that in Botryllus the stigmata of the full-grown oozooid are secondary formations due to the subdivision of a series of transversely elongated primary "proto-stigmata." In the buds, however, there are no protostigmata, the stigmata appearing quite independently of one another. In Thylacium and Styela he also finds proto-stigmata, while in Clavelina, as in the buds of Botryllus, the development is secondarily abbreviated. Garstang therefore regards Pyrosoma in which the adult stigmata are elongated transversely as being a primitive type of Caducichordate Tunicata, in which the proto-stigmata have never been subdivided. He also regards Botryllus and Styela as being more primitive in the Ascidiacea than Clavelina.

WILLEY (30) reconsiders his former views in regard to homologies in Amphiozus and the Tunicata, and now submits that the proboscis cavity, endostyle, mouth, and first pair of gill slits in Ascidians correspond to the organs bearing the same names in Amphiozus, while the club-shaped gland of Amphiozus cannot be homologous with the intestine of Ascidians, as suggested by Van Beneden and Julin.

Kowalevsky (14) has made the remarkable discovery that the cells in the Ascidian test are mesoblastic in origin, being derived from the mesenchyme of the larva and making their way through the ectoderm to the exterior during the metamorphosis. The thin test of the free-swimming tailed larva (*Phallusia mammillata*) has no cells. This agrees well with the observations made by various investigators, that in Compound Ascidians the cells of the test act as phagocytes. Salensky (25) has also independently discovered that in *Pyrosoma* the test cells are mesodermal in origin. Julin (13) is of opinion that in *Styclopsis* there are two kinds of cells in the test, the one pigmented red, like the pigmented connective tissue cells throughout the body, and probably of

mesodermic origin, the others colourless, and probably of epiblastic origin from the proliferation of the ectoderm.

# (c.) PHYSIOLOGY.

JOURDAIN (12) has been making experiments on the process of deglutition in Ascidians. He finds that in *Perophora* and *Clavelina* deglutition takes place along the dorsal side of the branchial sac, a little to the right hand side of the series of languets. He considers that one of the functions of the vibratile pit (dorsal tubercle) is to furnish mucus for the purpose of agglutinating the food particles. HERDMAN (10) places on record an instance of a large Ascidian (A. mentula) acting as a functional hermaphrodite and fertilizing its own ova.

# III.—GEOGRAPHICAL DISTRIBUTION.

NOTT (18) records eight species of Compound Ascidians (all new to science) from the north shore reef, Auckland, New Zealand. Only four species of Ascidians had previously been known from New Zealand.

LOHMANN (16) discusses both the horizontal and the vertical distribution of the Appendicularians in the North Atlantic, as brought out by the "Plankton" Expedition.

BLES (1) finds Appendicularia abundant at Plymouth at beginning of July, chiefly Oikopleura cophocerca.

HERDMAN (7), from a consideration of the geographical distribution of the known *Tunicata*, comes to the conclusion that Ascidians attain their greatest development, both in numbers and size, in the colder northern and southern seas.

HERDMAN (9) gives a list of twenty-three species of Simple and Compound Ascidians from the Norwegian seas. Several of them are new to science, but are not yet described in detail.

### IV.—SYSTEMATIC.

#### ASCIDIÆ COMPOSITÆ.

#### DIDEMNIDÆ.

Sarcodidemnoides, n. g., OKA & WILLEY, (21) p. 321.

Sarcodidemnoides misakiense, n. sp., id. ibid.

Leptoclinum niveum, n. sp., Nott, (18) p. 308.

Leptoclinum densum, n. sp., id. t. c. p. 311.

Leptoclinum tuberatum, n. sp., id. t. c. p. 314.

Leptoclinum maculatum, n. sp., id. t. c. p. 316.

Polysyncraton, n. g., NOTT, (18) p. 318 [genus not defined apart from the species.]

Polysyncraton paradoxum, n. sp., id. ibid.

Polysyncraton fuscum, n. sp., id. t. c. p. 321.

# DISTOMIDE.

Cystodytes aucklandicus, n. sp., Norr, (18) p. 323. Cystodytes perspicuus, n. sp. ?, id. t. c. p. 326.

# DIPLOSOMIDÆ.

Diplosoma mitsukurii, n. sp., OKA, (19) p. 266.

#### LARVACEA.

Folia, n. g., LOHMANN, (16) p. 139.
Folia athiopica, n. sp., id. ibid.
Folia gracilis, n. sp., id. t. c. p. 147.
Oikopleura labradoriensis, n. sp., id. t. c. p. 141.
Althoffia, n. g., id. t. c. p. 146.
Althoffia tumida, n. sp., id. t. c. p. 147.
Fritillaria glandulifera, n. sp., id. ibid.



# VII. MOLLUSCA.

BY

B. B. Woodward, F.G.S., F.R.M.S., &c.

In this Record an attempt has been made to repair the more serious of the omissions that occurred in *Mollusca* in Vols. xxvI and xxVII. Accordingly important works by Cossmann, Dall, Heude, and others have been either noticed for the first time, or portions that had been previously overlooked have been now inserted.

The Palæontological papers have not been so exhaustively treated as they were in Vol. XXVIII. The labour of examining the extensive literature of Geology is of itself very great; and as it does not appear necessary for the purposes of Zoology proper that all the Palæontological details should be given, the Recorder, in the absence of an understanding that the Record should serve the purposes of Palæontologists as well as Zoologists, has not felt justified in extending the length of his Record by introducing details that might be looked on as superfluous. Hence, fossil species described as new are mostly omitted from the Systematic part of the Record, and the titles of the papers do not invariably appear in the general list. Under the heading "Geological," in Section II, there will, however, be found references to all important Palæontological works and communications; and the fact of new species being described in them is always noted. New generic and sectional names, however, are fully recorded in the Systematic part.

#### I.—TITLES.\*

- ADAMS, G. W. Land and Freshwater Shells at Karachi. Journ. Conch. vii, pp. 81 & 82.
- ADAMS, L. E. A contribution to the authenticated records of Derbyshire. T. c. p. 77.

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

- 3. Aldrich, T. H. A new Land Shell [Nanina dohertyi] from Sumatra. Naut. vi, p. 90.
- AMALIZKY, W. Ueber die Anthracosien der Permformation Russlands. Palæontogr. xxxix, pp. 125-212, 5 pls. Also separately in Russian: Text 8vo, pls. 4to. Abstract in Protok. Otdyel. Biol. 1891-92, No. 3.
- Ancey, C. F. Description of a new species of Helicidæ [Ptychodon magdalenæ]. Brit. Nat. 1891, pp. 65 & 66.
- 6. —. Diagnoses of Land Shells. Op. cit. 1892, pp. 125-127, 1 fig.
- 7. —. On some Shells from Eastern Bolivia and Western Brazil. Journ. Conch. vii, pp. 90-97, fig.
- 8. —. Descriptions de mollusques nouveaux. Le Nat. 1892, p. 178.
- Études sur la Faune malacologique des Iles Sandwich. Mém. Soc. Zool. v, pp. 708-722.
- ARCHER, F. Supplementary report upon the Testaceous Mollusca of the L[iverpool] M[arine] B[iological] C[ommittee] District. P. Liverp. Soc. vi, pp. 105-121.
- BAICHÈRE, E. Faunule malacologique de Carcassonne. Feuill. Nat. 1892, pp. 63-68.
- BAKER, F. C. Notes on a collection of Shells from the Mauritius; with a consideration of the genus Magilus of Montfort. P. Rochester Acad. ii, pp. 19-40, 1 pl.
- Ballowitz, E. Ueber den feineren Bau der Muskelsubstanzen.
   Die Muskelfaser der Cephalopoden. Arch. mikr. Anat. xxxix, pp. 291-324, 2 pls. Abstr. in J. R. Mier. Soc. pp. 351 & 352.
- BARROIS, J. Anatomie du Cryptazeca monodonta. Rev. Sci. Nat. Ouest, ii, pp. 330 & 331, figs.
- Bednall, W. T. Land and Freshwater Mollusca [of the Elder Exploring Expedition]. Tr. R. Soc. S. Austr. xvi, pp. 62-67, 1 pl.
- Behrendsen, O. Zur Geologie des Ostabhanges der Argentinischen Cordillere. Z. Geol. Ges. xliv, pp. 1-42.
- BERGH, R. Die Nudibranchiata holohepatica porostomata. Verh. z.-b. Wien, xlii (Abh.), pp. 1-16.
- System der Nudibranchiaten Gasteropoden. Semper's Reisen in Archipel Philippinen, Theil 11, Bd. ii, Hft. 18, pp. 995-1165 [completing vol.].
- Opisthobranches provenant des campagnes du yacht l'Hirondelle. Fasc. IV of "Résultats des Campagnes scientifiques . . . par Albert I. . . . de Monaco." Monaco : 1892 [1893 on wrapper], 4to, 35 pp., 4 pls.

- BEUSHANSEN, L. Amnigenia rhenana ein Anodonta ähnlicher zweischaler aus dem rheinischen Mitteldevon. JB. k. preuss. geol. Landesanst. xi, Abhand. pp. 1-10, fig.
- BIERBAUER, B. A Check-list of the Palsozoic Fossils of Wisconsin, Minnesota, Iowa, Dakota, and Nebraska. Bull. Minnesota Acad. iii. Mollusca, pp. 234-245.
- BINNEY, W. G. A Fourth Supplement to the Fifth Volume of the Terrestrial and Air-breathing Mollusks of the United States and adjacent Territories. Bull. Mus. C. Z. xxii, pp. 163-204, 4 pls.
- 22. BITTNER, A. Neue Arten aus der Trias von Balia in Kleimasien. JB. geol. Reichsanst. xlii, pp. 77-89, 2 pls.
- Ueber die systematische Stellung von Mactra und verwandten Gattungen. Verh. geol. Beichsanst. 1892, pp. 232-241.
- BIZET, E. Catalogue des Mollusques observés à l'état vivant dans le Département de la Somme. Part II. Mem. Soc. L. N. Fr. viii, pp. 262-405.

Part I appeared in op. cit. vii.

- BLAKE, J. F. The Evolution and Classification of the Cephalopoda: an account of recent advances. P. Geol. Ass. xii, pp. 275-295.
- 26. BLUM, J. Schlendertage auf Capri. Nachr. mal. Ges. 1892, pp. 23-30.
- Die Schneckenfauna von Richisan im Klönthal (Kanton Glarus). Nachr. mal. Ges. 1892, pp. 127-130.
- BOEHM, G. Lithiotis problematica, Gümbel. Ber. Ges. Freiburg, vi, pp. 65-80, 3 pls.
- Ueber den Fussmuskeleindruck bei Pachyerisma. T. c. pp. 119 & 120.
- Bœttger, O. Drei neue mitteloligocane Mollusken aus deutschem Rupelthon. Mal. Bl. xi, pp. 89-93.
- Zur Kenntniss der Land- und Süsswasser-Mollusken von Nossi-Bé. III. Nachr. mal. Ges. 1892, pp. 53-58.
- Schnecken von der Insel Giura, N.-Sporaden, aus dem phokischen Parnass und aus anderen griechischen Gebieten. T. c. pp. 59-66.
- Die Meeresmollusken der mittleren Liukiu-Inseln. (Vorwort von A. Fritze.) T. c. pp. 153-157.
- Neue Stenogyren aus Westafrika und ein neues Aperostoma aus Honduras. T. c. pp. 202-204.
- Wissenschaftliche Ergebnisse der Reise Dr. J. Valentin's im Sommer, 1890. II. Die Meeresmollusken der Insel Kalymos. Ber. Senck. Ges. 1892, pp. 150-163.
  - \_\_\_\_\_. [See also for Syndosmya schneideri, n. sp., Schneider (337), and for determination and description of species, Simroth (355).]

- 36. Bourguignar, J. R. Iconographie malacologique des Animaux mollusques fluviatiles du Lac Tanganika. Corbeil: 1888, 8vo, 82 pp., 35 pls.
- 37. —. Mollusques de l'Afrique Équatoriale de Moguedouchou . . . au Tanganyika. Paris : 1889, 8vo, 229 pp., 8 pls.
- 38. BOUTAN, L. Sur le système nerveux de la Nerita polita. C.R. cxiv, pp. 1133-1135. Abstr. in J. R. Micr. Soc. 1892, p. 465.
- 39. BOUVIER, E. L. Quelques observations anatomiques sur les Mollusques Gastéropodes. CR. Soc. Biol. iv, pp. 987-992.
- Sur l'organisation des Amphiboles. Bull. Soc. Philom. iv, pp. 146-153.
  - —. [See also Fischer, P., & Bouvier, E. L. (131 & 132).]
- ROYCOTT, A. E. Contributions towards a list of the Mollusca of Herefordshire. Sci. Goss. 1892, pp. 77-79.
- 42. Brancsik, C. Consignatio systematica specierum in itinere Bosnensi anno 1888 . . . collectarum, &c. Evkön Trencsén Varm. xii (1890), pp. 68-76.
- BRAZIER, J. On the synonymy of Helix (Hadra) gulosa, Gould. P. Linn. Soc. N.S.W. vi, pp. 321-328.
- Catalogue of the Marine Shells of Australia and Tasmania.
   Pt. I. Cephalopoda; Pt. II. Pteropoda. Sydney: 8vo, 42 pp.
   Australian Museum Catalogue, No. 15.
- BROEMME, DR. Zur Fauna von Lugano. Nachr. mal. Ges. 1892, pp. 171-173.
- Bronn, H.G. Klassen und Ordnungen des Thier-Reichs . . . Bd. III.
   *Mollusca* . . . neu bearbeitet von Dr. H. SIMROTH. Leipzig : 1892,
   8vo, Lief 1 & 2.
- 47. Brown, Amos P. The development of the Shell in the Coiled Stage of Baculites compressus, Say. P. Ac. Philad. 1892, pp. 136– 141, 1 pl.
- BRUSINA, S. Ueber die Gruppe der Congeria triangularis. Z. geol. Ges. xliv, pp. 488-497.
- Fauna fossile terziaria di Markuševec in Croazia. Con un elenco delle Dreissenside della Dalmazia, Croazia e Slavonia. Glasnik Naravosl. Druzt. vii, pp. 113-210.
- 50. BUCAILLE, E. Catalogue raisonné des Mollusques terrestres et d'eau douce observés dans le département de la Seine-Inférieure . . . revue . . . par R. Fortin. Bull. Soc. Rouen, xxvii, pp. 171–189.
- BUCHNER, O. Die asymmetrie der Gastropoden in ihren Beziehungen und Wirkungen auf die Lebensäusserungen der schalentragenden Schnecken. JH. Ver. Württ. xlviii, pp. 58-92.

- BUCKMAN, S. S. The reported occurrence of Ammonites jurensis in the Northampton Sands. Geol. Mag. 1892, pp. 258-260.
- The Morphology of 'Stephanoceras' zigzac. Q. J. Geol. Soc. xlviii, pp. 447-452, 2 pls.
- 54. A Monograph on the Inferior Oolite Ammonites of the British Islands. Pt. vii, pp. 313-344, pls. lvii-lxxvi (Pal. Soc.).
- 55. Bucquoy, E., Dautzenberg, P., & Dollfus, G. Les Mollusques Marins du Roussillon: fasc. 19, Carditdæ, Lasæidæ, Galeommidæ, Cardiidæ; fasc. 20, Cardiidæ (fin), Chamidæ, Isocardiidæ, = Tome ii, pp. 221-320, pls. xxxviii-li.
- CANAVARI, M. Note di Malacologia fossile. Bull. Soc. Mal. Ital. xvi, pp. 65-73, 1 pl.
- CARAZZI, D. La perforazione delle rocce calcaree per opera dei datteri (*Lithodomus dactylus*). Atti Soc. Ligust. iii, pp. 279-297, fig.
- CARUANA-GATTO, A. New Clausilies from Malta. Med. Nat. i, pp. 148 & 149.
- Disappearance of Spondylus gaderopus, L., and other species from Maltese Waters. T. c. p. 165.
- 60. —. Maltese Cacilianella. Op. cit. ii, pp. 226 & 227.

  3 new species are described by WESTERLUND.
- 61. —. Assiminea littorina, Delle Chiaie, in Malta. T. c. p. 245.
- CAZIOT, —. Liste des coquilles marines recueillies à Bandol (pres Toulon). Feuill. Nat. 1892, pp. 148 & 149.
- 63. CHASTER, G. W. Shell hunting in Merionethshire. Journ. Conch. vii, pp. 78 & 79.
- 64. CLARKE, J. M. The discovery of Clymenia in the fauna of the intumescens zone (Naples beds) of Western New York, &c. Am. J. Sci. xliii, pp. 57-62, figs.
- 65. CLESSIN, S. Bythinella bosniensis, n. sp. Mal. Bl. xi, p. 20, fig.
- 66. Clubb, J. A. [See Herdman & Clubb (184).]
- 67. COCKERELL, T. D. A. [Descriptions of alleged new species of Helix and Veronicella, concealed in "Museum Notes".] J. Inst. Jamaica, i, pp. 55, 96, 134.
- Remarks on Australian Slugs. Ann. N. H. ix, pp. 370-372.
   [Reply to Hedley (178).]
- The genera Limax, Arion, and Helix. Conchologist, ii, pp. 28 & 29.
- The British Arionida. T. c. pp. 84 & 85.
   [A criticism of Collings (79).]
- New varieties of American Mollusca. Journ. Conch. vii, p. 39.

- 72. [COCKERELL, T. D. A.] A revised list of the species of British Slugs. Journ. Conch. vii, pp. 66-70.
- Notes on the North American species of Succinea. Naut. vi, pp. 21-23 & 29-31.
- The endemic features of the British Slug-Fauna. Sci. Goss. 1892, pp. 255-259.
- Collinge, W. E. Notes on the variation of the genus Arion. Ann. N. H. ix, pp. 307 & 308.
- Limax maximus, L., and its variety cinereo-niger, Wolf. Ann. N. H. x, pp. 425 & 426.
- 77. Descriptions of a new variety of Arion hortensis, Fér., and A. circumscriptus, Johnst. Conchologist, ii, pp. 26-27.
- 78. Some further remarks on the burrowing habits of certain Land Molluscs. T. c. pp. 29 & 30.
- 79. A review of the Arionida of the British Isles. T. c. pp. 56-66 & 76-83.

[Criticism by COCKERELL, t. c. pp. 84 & 85; reply by COLLINGE, pp. 85 & 86.]

- A catalogue of the Slugs of the British Isles. Brit. Nat. 1892, pp. 176-179.
- CONKLIN, E. G. Cleavage of the ovum in Crepidula fornicata. Zool.
   Anz. xv, pp. 185-188, figs. Abstr. in J. R. Micr. Soc. 1892, p. 464.
- 82. COOKE, A. H. On the origin of the genera of Land and Freshwater Mollusca. Conchologist, ii, pp. 41-48.
- On the geographical distribution of the Land-Mollusca of the Philippine Is., and their relations to the Mollusca of the neighbouring groups. P. Z. S. 1892, pp. 447-469. Abstr. in Nature, xlvi, p. 142.
- COOPER, J. G. Catalogue of the Land and Freshwater Mollusca of Lower California. Zoe, iii, pp. 12-25.
- Cossmann, M. Catalogue illustré des coquilles fossiles de l'Éocène des Environs de Paris. Fasc. v (Pteropoda, Cephalopoda), Supplement. Mem. Soc. Mal. Belg. 1891 [1893], pp. 1-163; also, separately in advance, July, 1892.

The names for new sections proposed in Parts iii & iv, and omitted from the RECORD in 1889-90, have been inserted in the Systematic part this year.

- 86. —. Réponse aux observations de M. É. Vincent sur le "Gilbertia inopinata, Morlet." Bull. Soc. Mal. Belg. 1891 [1893], pp. lxiv-lxvi.
- Révision sommaire de la faune du terrain oligocène marin aux environs d'Étampes (continued). J. de Conch. 1892, pp. 330-375.

- 88. COUPIN, H. Les Mollusques: introduction à l'étude de leur organisation, développement, classification, affinités, et principaux types, vii & 261 pp., figg.
- COUTAGNE, G. Sur la Faune malacologique terrestre des Iles de la Rade de Marseille. C.R. Ass. Fr. Sci. 1891, ii, pp. 546-554.
- Cox, J. C. Descriptions of some new species of Pulmonate Mollusca from Australia and the Solomon Is. P. Linn. Soc. N.S.W. vi, pp. 565-570, 2 pls.
- 91. CRANDALL, O. A. Physa heterostropha. Naut. vi, pp. 20 & 21.
- CRICK, W. D. Bibliography of the Land and Freshwater Mollusca of Northamptonshire. Conchologist, ii, pp. 10-13.
- CROSSE, H. Faune malacologique terrestre et fluviatile de l'île de Portorico. J. de Conch. 1892, pp. 5-71.
- 94. —. Études malacologiques sur des genres nouveaux on peu connus. T. c. pp. 279-292, 1 pl.
  - ---. [See also FISCHER & CROSSE (133).]
- CROSSE, H., & FISCHER, P. Note sur les Mollusques marins du Golfe de Siam (Côte O. du Cambodge). T. c. pp. 71-77.
- Wote sur le genre Holospira, Martens, et sur la distribution géographique des espèces dont il se compose. T. c. pp. 256-279, 1 pl.
- 97. & —. Note sur le Neritina picta, Sow. T. c. pp. 292 & 293.
- & . Diagnoses Molluscorum Reipublicæ Mexicanæ et Guatelmalæ incolarum. T. c. pp. 294-296.
  - ---. [See also FISCHER, P., & CROSSE, H. (133).]
- CUÉNOT, L. Études sur le sang et les glandes lymphatiques dans la série animale (2º Partie: Invertébrés). Arch. Z. éxpér. ix, 1891 (Mollusques), pp. 19-54. Abstr. in J. R. Micr. Soc. 1892, pp. 192 & 193.
- L'excrétion chez les Gastéropodes pulmonés. C.R. cxv, pp. 256-258. Abstr. in J. R. Micr. Soc. 1892, p. 597.
- 100A. —. Études physiologiques sur les Gastéropodes Pulmonés. Arch. Biol. xii, pp. 683-740, 1 pl. Abstr. in J. R. Micr. Soc. 1893, p. 312.
- 101. Dall, W. H. Reports on the Results of Dredging, under the supervision of A. Agassiz, in the Gulf of Mexico (1877-78) and in the Caribbean Sea (1879-80), by the U.S. Coast Survey Steamer "Blake," . . . xxix. Report on the Mollusca. . . . Part II. Gastropoda and Scaphopoda. Bull. Mus. C. Z. xviii, 1889, 492 pp., 31 pls.

- 102. [Dall, W. H.] Contributions to the Tertiary Fauna of Florida. . . . Part II [completing the Gastropoda, Pteropoda, Polyplacophora, and Scaphopeda, Tr. Wagner Inst. iii, pp. 201-446, 1 map, and pls. xiii-xxii.
- 103. —. Note on Cytherea convexa, Say. Naut. vi, pp. 52 & 53.
- 104. —... On some types new to the fauna of the Galapagos Islands. Op. cit. v. pp. 97-99.
- 105. —... Instructions for Collecting Mollusks, and other useful hints for the Conchologist. Bull. U.S. Nat. Mus. xxxix, 56 pp.
  - ---. [For Vitrinella williamsoni and Amphissa bicolor, n. spp., see WILLIAMSON (435).]
  - ---. [For Pleurotoma simpsoni, Tralia minuscula, and Pandora bushiana, n. spp. (1889), see SIMPSON (349).]
- 106. Danilo, F., & Sandri, G. B. Elenco dei molluschi lamellibranchiati dei dintorni di Zara. Glasnik Naravoel. Druzt. vi, 1891, pp. 82-108.
- 107. DAUTZENBERG, P. Révision des Mollusques Marins des Açores. [Fasc. I of Résultats des Campagnes scientifiques accomplies sur son yacht par le Prince...de Monaco.] Monaco: 1889, 4to, 112 pp., 4 pls.
- Description d'un Perideris nouveau [P. lechatelieri] provenant du Dahomey. J. de Conch. 1892, p. 297.
- Description d'une espèce nouvelle du genre Chama [C. nicolloni] provenant des côtes océaniques de France. Bull. Soc. Ouest Fr. ii, pp. 133-135, figg.
  - —. [See also Bucquoy, E., DAUTZENBERG, P., & DOLLFUS, G. (55).]
- DEAN, G. W. More about Unio luteolus and U. radiatus. Naut. v, pp. 112 & 113.
- Catalogue of the Shell-bearing Mollusca of Portage County, Ohio. Am. Nat. xxvi, pp. 11-23.
- 112. Deschamps, E. Description d'une nouvelle espèce d'Unio [U. corbeti] de Ceylan. Bull. Soc. Z. Fr. xvii, pp. 68-69, fig. Dollfus, G. [See Bucquoy, E., Dautzenberg, P., & Dollfus, G. (55).]
- Donald, Miss J. Notes on some new and little-known species of Carboniferous Murchisonia. Q. J. Geol. Soc. xlviii, pp. 562-575, 2 pls.
- 114. Dreger, J. Die Gastropoden von Häring bei Kirchbichl in Tirol. Ann. Hofmuseum Wien, vii, pp. 11-34, 4 pls.
- 115. DROUET, H. Unionida nouveaux ou peu connus. J. de Conch. 1892, pp. 86-94.

- 115A. DROUET, H., & CHAPER, M. Voyage de M. Chaper à Borneo.— Unionidæ. Mém. Soc. Zool. v, pp. 145-155, 2 pls.
- 116. Dubois, R. Anatomie et physiologie comparées de la Pholade dactyle. Ann. Univ. Lyon, ii, fasc. 2, x & 167 pp., 15 pls. Dunker, W. [See Hesse, P. (185).]
- 117. EHRMANN, P. Nachtrag zur Gastropoden-Fauna der Umgegend von Leipzig nebst einigen biologischen Bemerkungen. SB. Ges. Leipzig, xviii, pp. 76-80.
- 118. ERLANGEB, B. von. Beiträge zur Entwicklungsgeschichte der Gasteropoden... Thl. I. Zur Entwicklung von Bythinia tentaculata. MT. z. Stat. Neap. x, pp. 376-407, 2 pls., figg. Abstr. in J. R. Micr. Soc. 1893, p. 313.
- 119. Mittheilungen über Bau und Entwicklung einiger marinen Prosobranchier. I. Ueber Capulus hungaricus. Zool. Anz. xv, pp. 465-468. Abstr. in J. B. Micr. Soc. 1893, p. 163.
- 120. ——. On the paired Nephridia of Prosobranchs, the homologies of the only remaining Nephridium of most Prosobranchs, and the relations of the Nephridia to the Gonad and Genital Duct. Q. J. Micr. Sci. xxxiii, pp. 587-623, 2 pls. Abstr. in J. R. Micr. Soc. 1892, p. 598. ETHERIDGE, R., JUN. [See JACK & ETHERIDGE (193).]
- 121. Eyre, W. L. W. On the Classification of Varieties [of Helix hortensis and nemoralis]. Conchologist, ii, pp. 7-10.
- 122. °FAGOT, P. Histoire malacologique des Pyrénées Françaises et Espagnoles. Liste, &c. Bull. Soc. Ramond, 1891, xxvi, p. 215 (fide Rev. Tr. Sci. xii, p. 393).
- Histoire malacologique de la presqu' île Sainte-Lucie (Aude).
   Bull. Soc. Aude, iii, pp. 173-180.
- 124. FAURE, C. Dell' origine di qualche varietà nei Molluschi. Riv. Ital. Sci. Nat. x, 1891, pp. 89, 90, & 105-107.
- FISCHER, H. Recherches sur la morphologie du Foie des Gastéropodes. Bull. Sci. Fr. xxiv, pp. 260-346, 7 pls. Abstr. in J. R. Micr. Soc. 1892, pp. 596 & 597.
- Note sur l'enroulement de la coquille des embryons de Gastropodes. J. de Conch. 1892, pp. 309-313.
- 127. —... Sur le developpement du Foie chez la Paludine. C.R. Soc. Biol. iii, 1891, pp. 644 & 645.
- 128. FISCHER, P. Catalogue et distribution géographique des Mollusques terrestres, fluviatiles, et marins d'une partie de l'Indo-Chine. Bull. Soc. Autun, iv, pp. 87-276.
- 129. —. Note sur la distribution géographique de l'Ovula carnea, Poiret. J. de Conch. 1892, pp. 77 & 78.

- 129A. [FISCHER, P.] Le Catalogue de la Collection Schlüter. J. de Conch. 1892, pp. 208-212.
- 130. —. Note sur la faune terrestre et fluviatile de l'île d'Hainan (Chine). Deuxième supplément. T. c. pp. 313-315.
- 131. FISCHER, P., & BOUVIER, E. I.. Recherches et considérations sur l'asymétrie des Mollusques univalves. T. c. pp. 117-207, 3 pls.
- 132. & Sur l'enroulement des Mollusques univalves. T. c. pp. 234-245.
- 133. FISCHER, P., & CROSSE, H. Études sur les mollusque terrestres et fluviatiles. Mission Scient. Mexique et Amér. Cent. VII, ii, pp. 313-392, pls. liii & liv. Melaniidæ, Proserpinidæ.
  - —. [See also Crosse, H., & Fischer, P. (95).]
- 134. FISCHER, P., & FISCHER, H. Diagnoses d'espèces nouvelles de Mollusques Céphalopodes recueillis dans le cours de l'Expédition scientifique du Talisman. J. de Conch. 1892, pp. 297-300.
- 135. Folin, Marquis De. Les mollusques speciaux à la région extrême sud-ouest de la France et l'Atlantide. Rev. Sci. Nat. Ouest, ii, pp. 324-329.
- 136. FORD, J. Some remarks on New Jersey Coast Shells. Naut. vi, pp. 25-29, figs.
- 137. Foresti, L. Di alcune varietà della *Melania verrii*, De Stef. Bull. Soc. Mal. Ital. xvi, pp. 74-79, 1 pl.
- 138. —. Di una nuova specie di *Pholadomya* [P. elegantula] pliocenica. T. c. pp. 80-82, 1 pl.
- 139. FUTTEBER, —. Die oberen Kreidebildungen den Umgebung des Lago di Santa Croce in der Venetianer Alpen. Pal. Abh. vi, 124 pp., 12 pls.
- 140. GAIN, W. A. Some remarks on the Colour changes in Arion intermedius, Normand. Conchologist, ii, pp. 55 & 56.
- 141. —. [Notes on Variation.] Brit. Nat. 1892.
- 142. GALLENSTEIN, H. von. Die Schalenformungen der Muscheln des Wörther-See's in Kärnten. Nachr. mal. Ges. 1892, pp. 102-114.
- 143. Galletly, A. The Pearl Molluscs of the Persian Gulf. P. Phys. Soc. Edinb. xi, pp. 30 & 31.
- 144. Gamble, F. W. Observations on two rare British Nudibranchs (Lomanotus genei, Verany, and Hancockia sudactylota, Gosse). Ann. N. H. ix, pp. 378-385, 1 pl.
- 145. Ganong, W. F. Zoological Notes. . . . 1. Mollusca [of New Brunswick]. Bull. Nat. Hist. Soc. New Brunswick, ix, pp. 46 & 47. Addenda to list published 1887.

146. GARSTANG, W. On some new or rare Marine Animals recently discovered on the Coast of Devonshire. Rep. Tr. Devon Ass. xxiv, pp. 377-386.

Mollusca, pp. 383 & 384.

- 147. —. Notes on the Structure and Habits of Jorunna johnstoni. Conchologist, ii, pp. 49-52.
  - GATTO, A. C. C. [See CARUANA-GATTO (58, &c.)]
- 148. GIRARD, A. O. Notice sur les Céphalopodes des côtes de l'Espagne. An. Soc. Esp. xxi, pp. 391-399.
- 149. GODWIN-AUSTEN, H. H. On new species and varieties of the . . . genus Diplommatina from the Garo, Naga, and Munipur Hill-ranges, Assam. P. Z. S. 1892, pp. 509-520.
- 150. Description of a new species of Helix of the subgenus Plectopylis [H. fultoni]. Ann. N. H. 1892, pp. 300 & 301.
- 151. GODRICH, E. S. Note on a large Squid (Ommastrephes pteropus, Stp.). J. Mar. Biol. Ass. ii, pp. 314-321, figg.
- 152. GOODCHILD, J. G. Notes on Carboniferous Lamellibranchs [Cteno-donta and Nucula]. P. Phys. Soc. Edinb. xi, pp. 244 & 245.
- 152A. —. Note on the genus Allorisma, King. T. c. pp. 245 & 246.
- 153. GORJANOVIC-KRAMBERGER, D. Ueber einen tertiären Rudisten [Ceratoconcha costata, n. g. & sp.] aus Podsused bei Agram. Glasnik. Naravosl. Druzt. iv, 1889, pp. 48-55, 1 pl.
- 154. Granger, A. Description d'une nouvelle espéce d'Ampullaire [Ampullaria brohardi]. Le Nat. 1892, p. 97, fig.
- 155. Gregorio, March A. Df. Sul genere Pectunculus e pricipuamente sulle specie viventi Mediterranee e fossili. Nat. Sicil. xi, pp. 89-96 & 106-114.

For comment on this paper, see MONTEROSATO (260).

- Intorno alla nota del March. di Monterosato sul genere Pectunculus. T. c. pp. 209-212.
- 157. —... Nota intorno talune conchiglie Mediterranee del terziario inferiore del littorale Ligure. T. c. pp. 270-272.
- 158. GRIESBACH, —. Ueber das Blut der acephalen Mollusken. Verh. Ges. Deutsch. Naturf. lxiii, pt. 2 (1891), pp. 131-133.
- 159. GRIFFITHS, A. B. Sur une globuline incolore [from the blood of Patella] qui possède une fonction respiratoire. C. R. cxv, p. 259.
- 160. —... Sur une globuline respiratoire contenue dans le sang des Chitons. T. c. pp. 474 & 475.
- 161. —. Sur les tissus nerveux de quelques invertébrés. T. c. pp. 562 & 563.
- 162. —. On the Blood of the *Invertebrata*. P. R. Soc. Edinb. xviii (Blood of the *Mollusca*), pp. 290-292, & xix, pp. 127-130.

- 163. GROBBEN, C. Beiträge zur kenntniss des Baues von Cuspidaria (Neaera) cuspidata, Olivi, nebst Betrachtungen über das System der Lamellibranchiaten. Arb. z. Inst. Wien, x, No. 3, 46 pp., 4 pls. Abstr. in J. R. Micr. Soc. 1893, p. 164.
- 164. Das system der Lamellibranchiaten. Zool. Auz. xv, pp. 371-375.

This is the second section of the preceding. Diverse translations of the table of classification appear in J. R. Micr. Soc. 1892, p. 772, and Nat. Sci. ii, p. 9.

- 165. GUPPY, R. J. L. Note on Bulimus oblongus. P. Z. S. 1892, pp. 271 & 272.
- 166. HALLER, B. Die Anatomie von Siphonaria gigas, Less., eines opisthobranchen gasteropodeu. Arb. z. Inst. Wien, x, pp. 71-100, 3 pls. Abstr. in J. R. Micr. Soc. 1892, p. 770.
- Die morphologie der Prosobranchier . . . 111. Naticiden und Calyptraeiden. Morph. JB. xviii, pp. 451-543, 7 pls. Abstr. in J. R. Micr. Soc. 1892, pp. 769-770.
- 168. HART, H. C. Notes on Marine Mollusca collected on the Coasts of Donegal and Dublin. Zool. xvi, pp. 51-56, 105-107, 139-141, 181-183 & 414.
- 169. HARTMANN, W. D. Description of a new Diplomorpha [D. coxi]. P. Linn. Soc. N.S.W. vi, p. 571, figg.
- Catalogue of the genus Partula [pt. i]. Naut. vi, pp. 73-76, figg.
- HECHT, E. Remarques sur quelques moyens de défense des Éolidiens. C.R. cxv, pp. 746-748. Abstr. in J. R. Micr. Soc. 1893, p. 313.
- 172. HEDLEY, C. List of the Mollusca collected by Sir W. MacGregor on the Fly River. Colonial Reports, No. 6, British New Guinea, 1889-1890 (1891), p. 84.
- 172a. —. Uses of shells among the Papuans. Thompson's British New Guinea. London: 1892, 8vo, pp. 283-285.
- 173. —. [Note on a species of Glaucus (G. atlanticus) frequently washed ashore on the Coast of New South Wales.] P. Linn. Soc. N.S.W. vi, p. 576.
- 174. ——. The Land Molluscan fauna of British New Guinea (Anatomical Supplement) [to paper Zool. Rec. 1891, Moll. p. 12, No. 172]. T. c. pp. 685-698, 5 pls.
- 175. —. Observations on the Charopidæ. Op. cit. vii, pp. 157-169, 2 pls.
- 176. —. On the genus *Perrieria*. T. c. pp. 311-313.
- 177. —... On the structure and affinities of *Panda atomata*, Gray. Rec. Austral. Mus. ii, pp. 26-31, 3 pls.

- 178. [Hedley, C.] Remarks on Australian Slugs. Ann. N. H. ix, pp. 169-171.
  - See Cockerell (68) for reply.
- 179. HEDLEY, C., & MUSSON, C. T. On a collection of Land and Freshwater Shells from Queensland. P. Linn, Soc. N.S.W. vi, pp. 551-564.
- 180. Heimburg, H. von. Abbildungund Beschreibung einer neuen Helix [H. aggiei, Solomon Is.]. Mal. Bl. xi, pp. 158 & 159, figg.
- 181. Hemphill, H. Studies among Mollusks, Instinct and Genera. Zoe, ii, pp. 312-318.
- 182. —. Note on a Californian Loligo [L. stearnsii, n. sp.]. Zoe, iii, pp. 51 & 52.
- 183. HERDMAN, W. A. An additional occurrence of Pleurophyllidia loveni in Britain. Conchologist, ii, p. 52.
- 184. HERDMAN, W. A., & CLUBB, J. A. On the innervation of the Cerata of some Nudibranchiata. Q. J. Micr. Sci. xxxiii, pp. 541-558, 3 pls. Abstr. in J. R. Micr. Soc. 1892, p. 598.
- 185. Hesse, P. Zur Kenntniss der Molluskenfauna Westfalens. (Conchylien, welche sich in der Grafschaft Schaumburg und der Umgegend finden. Von W. Dunker.) J. Ber. Westf. Ver. xix (1891), pp. 79–86.
- 186. Heude, P. M. Notes sur les Mollusques terrestres (et d'eau douce) de la Vallée du Fleuve Bleu. Pp. 125-188, pls. xxxiii-xliii in Mém. Hist. Nat. Empire Chinois, par des Pères de la Compagnie de Jésus. Tom i, pt. 4, 1890.
  - Part only seen and recorded in 1891.
  - ---. [See also MOELLENDORFF (257).]
- 187. HEUSCHER, J. Zur Anatomie und Histologie der Proneomenia sluiteri, Hubrecht. Viert. Ges. Zürich, xxxvii, pp. 148-161, figg. Abstr. in J. R. Micr. Soc. 1892, p. 771.
- 188. Holm, G. Om mynningen hos Lituites, Breyn. Geol. Fören. Stock-holm Förh. xiii, pp. 736-774, 3 pls.
- 189. Om tvenne Gyroceras-formigt böjda Endoceras-arter. Op. cit. xiv, pp. 125-134 & 208-210, 3 pls.
- 190. Horsley, J. W. List of *Mollusca* found at Meiringen, Switzerland. Journ. Conch. vii, p. 32.
- 191. HUDLESTON, W. H. A Monograph of the British Jurassic Gasteropoda. Pt. 1, No. 6. Gasteropoda of the Inferior Oolite (Pal. Soc.), pp. 273-324, pls. xxi-xxvi.
- 192. Hudleston, W. H., & Wilson, E. A Catalogue of British Jurassic Gasteropoda, &c. London: 8vo, xxxiii & 147 pp.
- 192A. ISHIKAWA, C. [Introduction to the Anatomy of Animals. 1. Dipsas plicata, Lea.] In Japanese. Tokio: 1892, 8vo, 35 pp., 6 pls.

- 193. Jack, R. L., & ETHERIDGE, R., JUN. The Geology and Palæontology of Queensland and New Guinea. Text & atlas, 2 vols. Brisbane & London: 1892, 8vo, xxx & 768 pp., 68 pls.
- 194. Janet, C. Note sur trois nouvelles Bélemnites Sénoniennes. Bull. Soc. Géol. xix, pp. 716-721, 1 pl., figg.
- 195. JHERING, H. VON. Morphologic und Systematik des Genitalapparates von *Helix*. Z. wiss. Zool. liv, pp. 386-520, 2 pls. Abstr. in J. R. Micr. Soc. 1892, pp. 768 & 769.
- 196. —. Anodonta und Glabaris (finis). Zool. Anz. xv, pp. 1-5.
- Zur Kenntniss der Gattung Cristaria. Nachr. mal. Ges. 1892, pp. 1-14.
- 198. —. Die Gattung Hyalina. T. c. pp. 132-140.
- 198A. —. Ueber Atopos, Simroth. T. c. pp. 140-144.
- 199. JOBERT, —. De la formation du Byssus chez quelques Mollusques acéphales. Rev. Tr. Sci. xii, pp. 396-398.
- JOHNSON, C. W., & PILSBRY, H. A. A new American Helix [H. surgenti]. Naut. vi, pp. 8 & 9.
  - & —. [See also PILSBRY & JOHNSON (303).]
- 201. Johnston, R. M. Critical observations on recent contributions to our knowledge of the freshwater Shells of Tasmania. Part I. P. R. Soc. Tasm. 1888 (1889). pp. 84-90, 4 pls.

The plates illustrate the author's paper in the same publication for 1877.

- Observations on the variability of the Tasmanian Unio. T. c. pp. 95-96, 2 pls.
- 203. JOUBIN, L. Recherches sur la coloration du tegument chez les Céphalopodes. Arch. Z. expér. x, pp. 277-330, 3 pls. & figg.
- 204. JOUSSEAUME, F. Réflexions sur la Faune malacologique de la Mer Rouge. Ann. Sci. Nat. xii, pp. 343-363.
- 205. Kellog, J. L. Some notes from a study in the Morphology of the Lamellibranchiata. Johns Hopk. Univ. Circ. xi, pp. 80-83. Abstr. in J. R. Micr. Soc. 1892, pp. 353 & 354.
- 206. Keyes, C. R. The Plutyceras group of Palseozoic Gasteropods. Am. Geol. x, pp. 273-277.
- 207. KILIAN, W. Sur quelques Ammonitides appartenant au Muséum d'Histoire Naturelle de Lyon. Arch. Mus. Lyon, v, No. 3, 9 pp., 1 pl.
- Sur quelques Céphalopodes nouveaux ou peu connus de la période secondaire. Bull. Soc. Stat. Isère (4) i, pp. 203-208, 7 pls.
- KITTL, E. Die Gastropoden der Schichten von St. Cassian der südalpinen Trias. II Thiel. Ann. Hofmuseum Wien, vii, pp. 35-97, 5 pls.

- 210. KNîPOVÎCH, N. K émbriologhii Kruilonoghikh [On the Embryology of the *Pteropoda*]. Rev. Sci. Nat. St. Pétersb. 1891, pp. 148-151.
- 211. KOENEN, A. VON. Das norddeutsche Unter-Oligocan und seine Mollusken-Fauna . . . Lief. IV. Rissoidæ-Patellidæ . . . Opisthobranchiata . . . Polyplacophora, Scaphopoda, Pteropoda, Cephalopoda. Abh. Geol. specialkarte Preuss. x, No. 4, pp. 819-1004, pls. liii-lxii.
- 212. Koken, E. Ueber die Gastropoden der rothen Schlernschichten nebst Bemerkungen über Verbreitung und Herkunft einiger triassischer Gattungen. JB. Mineral. 1892, ii, pp. 25-36.

For extended paper, see Woehrmann & Koken (438).

- KORSCHELT, E. On the Development of Dreissena polymorpha,
   Pallas. Ann. N. H. ix, pp. 157-169, figg. Transl. from SB. nat. Fr. 1891, pp. 131-146.
- 214. Beiträge zur Entwicklungsgeschichte der Cephalopoden in Festschr. R. Leuckarts. pp. 347-373, 2 pls. & figg.
- 215. —... Ueber die Differenzierung der Keimblätter bei den Cephalopoden mit Rücksicht auf die Bildung des Darm-cauals und nervensystems. Verh. deutsch. Zool. Ges. ii, pp. 87-92.
  - Kramberger-Gorjanovic, D. [See Gorjanovic-Kramberger (153).]
- 216. Krause, A. Mollusken von Ostspitzbergen. Zool. JB. vi, Syst. pp. 339-376, 3 pls.
- 217. LACAZE-DUTHIERS, H. DE. Observation d'un Argonaute de la Méditerranée. Arch. Z. expér. x, pp. 37-56, fig. Abstr. in Le Nat. 1892, pp. 114 & 115, fig., and J. R. Micr. Soc. 1892, p. 351.
- 218. Lang, A. Lehrbuch der Vergleichenden Anatomie, &c. Abth. iii, Mollusca, pp. 567-870, figg. Jena: 1892, 8vo.
- 219. LETELLIER, —. Sur la fonction urinaire de l'organe de Keber chez les Mollusques acéphales. Bull. Soc. L. Norm. v, pp. 8-12.
- 220. Locard, A. Sur les *Pleurotomide* vivants du groupe du *Clathurella purpurea*. Échange, 1891, vii, pp. 5 & 6.
- 220A. —. Sur une espèce nouvelle du genre Neptunea [N. nicolloni]. T. c. p. 34.
- 220<sub>B</sub>. Espèces nouvelles du groupe du Cytherea rudis, Poli. T. c. p. 69.
- 221. —. Les Anomies des côtes de France. T. c. p. 86.
- 222. Révision des Alexia Françaises. T. c. pp. 131 & 132.
- 223. —. Sur une espèce nouvelle du genre Belgrandia [B. coutagnei]. Op. cit. viii, p. 3.

- 224. [LOCARD, A.] Sur quelques Limnées Françaises du groupe du Limnœa limosa. T. c. pp. 18 & 19.
- 225. —. Les Limnées Françaises du groupe du Limnæa peregra. T. c. pp. 77-79, 90, & 91.
- 226. —... Sur la présence d'une *Mitra* [*M. aquitanica*] Française dans la faune océanique. *T. c.* p. 101.
- LOENS, H. Ergiebige Methode, kleine Wasserschnecken zu fangen. Nachr. mal. Ges. 1892, pp. 66-68.
- 228. —. Succinea oblonga, pfeifferi, putris im Trocknen lebend. Hirschberg in Schlesien. T. c. pp. 130 & 131.
- 229. —... Nachtrag zur Molluskenfauna Westfalens. T. c. pp. 169 & 170.
- 230. Die Gastropodenfauna des Münsterlandes. Mal. Bl. xi, pp. 121-157.
- 231. LOEWINSON-LESSING, F. Les Ammonées de la zone à Sporadoceras münsteri, dans les monts Gouberlinskya Gory. Mém. Soc. Belg. Géol. vi, pp. 15-25, 1 pl.
- LOISEL, G. Sur l'appareil musculaire de la Radula chez les Helix.
   J. de l'Anat. xxviii, pp. 566-572, figg.
- 233. LORY, P. Sur les Hoplites Valanginiens du groupe de Hoplites neocomiensis. Bull. Soc. Stat. Isère (4) i, pp. 229-255.
- 234. MacDonald, R. On the Classification of Varieties [of Helix hortensis and nemoralis]. Conchologist, ii, pp. 27 & 28.
- 235. MACGEORGE, —. Un mollusque nouveau [Thylacodes medusæ] du Japon. Le Nat. 1892, pp. 123 & 124, fig.
- 236. M'Intosh, W. C. Note on the occurrence of *Pleurophyllidia loreni* in Britain. Conchologist, ii, p. 21.
- 237. McMurtrie, J. Eigg Shells: notes on the Land and Freshwater Mollusca of the Island of Eigg. Journ. Conch. vii, pp. 113-119.
- 238. MARSHALL, W. B. Preliminary list of New York Unionida. Bull. New York Mus. Nat. Hist. i, No. 1, 17 pp.
- 239. —. Gould's types of Nassa and Columbella. Naut. vi, pp. 47 & 48.
- 240. MARTENS, E. Die lebenden Mollusken in den Kantonen Appenzell und St. Gallen. Ber. St. Gall. Ges. 1889-90 (1891), pp. 108-132.
- 241. —... Einige neue Arten von Land- und Süsswasser-Mollusken aus Uganda und dem Victoria-Nyansa. SB. nat. Fr. 1892, pp. 15-19.
- 242. . Ueber einige seltenere Conchylien der Mark Brandenburg, insbesondere über Clausilia latestriata, Bielz. SB. nat. Fr. 1892, p. 93.
- Ueber die von Dr. Stuhlmann in Nordostafrika gesammelten Land- und Süsswasser-Mollusken. T. c. pp. 174-181.

- 244. [Martens, E.] Die Beschreibung vier neuer Afrikanischer Conchylien-Arten. T. c. pp. 181-183.
- 245. —. Biologia Centrali-Americana. Mollusca, pp. 97-176, pls. vi-ix.
- 246. MARTINI & CHEMNITZ. Systematisches Conchylien-Cabinet. Lief. 390-395, Bd. iii, Abth. 12, Hft. 52-54: Columbellidæ and Nassidæ, bearbeitet von Dr. W. Kobelt, pp. 1-128, pls. i-xviii. Bd. xi, Abth. 4, Hft. 21: Pholadea bearbeitet von S. Clessin, pp. 25-40, pls. vii-xii.
- MANCARINI, A. I Molluschi conchigliferi delle adiacenze di Ascoli-Piceno. Bull. Soc. Mal. Ital. xvi, pp. 234-260.
- 248. MAZZABELLI, G. Intorno al preteso occhio anale dell larve degli Opistobranchi. Rend. R. Acc. Lincei, 1892, i, pp. 103 & 104. Abstr. in J. R. Micr. Soc. 1892, p. 770.
- 249. MELVILL, J. C. Notes upon Cypræa chrysalis and C. amphithales. Journ. Conch. vii, pp. 120-123.
- 250. MELVILL, J. C., & PONSONBY, J. H. Descriptions of seven new species of Terrestrial Mollusca from South Africa. Ann. N. H. ix, pp. 84-87, figg.
- 251. & —. Descriptions of seventeen new Terrestrial Mollusks from South or Central Africa, in the Collection of E. L. Layard, T. c. pp. 87-94.
- 252. & Descriptions of thirteen new species of Terrestrial and Freshwater Mollusca from South Africa. Op. cit. x, pp. 237-242, 1 pl.
- 253. METCALF, M. M. Preliminary notes upon the Embryology of Chiton. Johns Hopk. Univ. Circ. xi, pp. 79 & 80. Abstr. in J. R. Micr. Soc. 1892, pp. 352 & 353.
- 254. Mikhaklis, E. Opisanie novuikh i maloizvyestnuikh mollyuskov yuzhnagho Altaya i syevernoï Dzhungharii [Descriptions of new or little-known Mollusca from Southern Altai and Northern Dzoongaria].

   Limax natalianus, n. sp. Issued by the Acad. Imp. Sci. St. Petersb. 8vo, 6 pp., figg.
- 255. MILLOT, L. Nôte sur les Céphalopodes Dibranches du Lias supérieur de Sainte-Colombe-lès-Avallon (Yonne), précédée d'un résumé sur la classification des Céphalopodes. Bull. Soc. Autun, iv, pp. 37-57, fig.
- 256. Moebius, K. Eine echte Perle von ungewohnlicher Form und Färbung. SB. nat. Fr. 1892, pp. 1-3, fig.
- 257. MOELLENDOBFF, O. von. Bemerkungen zu P. Heude's notes sur les Mollusques terrestres de la vallée du Fleuve Bleu. Nachr. mal. Ges. 1892, pp. 15-23.
- 258. —. Die Landschneckenfauna der Tenimber Inseln (Timorlaut). T. c. pp. 81-102, 1 pl.

- MOENIG, J. Zur Molluskenfauna im Oberamt Saulgau. JH. Ver. Württ. xlviii, pp. 119-134.
- 260. Monterosato, Marchese T. di. Nota intorno ai *Pectunculus* dei mari d'Europa. Nat. Sicil. xi, pp. 143-155.

  Comment on a paper by Gregorio (155).
- 261. —. Note sur l'Emarginula compressa, Cantraine. J. de Conch.
- 1892, pp. 78-81.

  262. —... Monografia dei Vermeti del Mediterraneo. Bull. Soc. Mal.
- Ital. xvii, p. 7-48, 7 pls.

  263. Morlet, L. Diagnoses Molluscorum novorum in Indo-Chinâ
- collectorum. J. de Conch. 1892, pp. 82–86.
- 264. —. Descriptions d'espèces nouvelles, provenant de l'Indo-Chino. T. c. pp. 315-329, 3 pls.
- 265. MOYNIER DE VILLEPOIX, R. Recherches sur la formation et l'accroissement de la coquille des Mollusques. J. de l'Anat. Phys. xxviii, pp. 461-518 & 582-674, 4 pls.

Also issued as a Thesis, dated 1893.

- Sur la réparation de la coquille chez Helix aspersa. Bull.
   Soc. Z. Fr. xvii, pp. 30 & 31. Abstr. in J. R. Micr. Soc. 1893, p. 30.
- 267. —. Note sur le mode de production des formations calcaires du test des Mollusques. Mém. Soc. Biol. iv, pp. 35-42.
   MUSSON, C. T. [See HEDLEY & MUSSON (179).]
- 268. NEWTON, R. B. On the American Palæozoic Gasteropod, Trematonotus, and its identification in Britain, with description of a new species [T. britannicus]. Geol. Mag. 1892, pp. 337-341 & 525, 1 pl.
- NICOLAS, H. Compléments monographiques des genres Lartetia, Moitessieria, Bithinella, Avenionia, et Acme. Ann. Soc. Agric. Lyon, iv, pp. 27-50, 1 pl.
- Étude . . . sur la Faune malacologique du Danien des Environs de Saint-Remy et Les Baux (Provence). C.R. Ass. Fr. Sci. 1891, ii, pp. 448-457, figg.
- 271. Novák, O. Revision der palæozoischen Hyolithiden Böhmens. Abh. Böhm. Ges. iv, No. 6, 48 pp., 6 pls., figg. in text.
- 272. OLDHAM, C. Additions to the South Devon list of Land and Freshwater Mollusca. Journ. Conch. vii, pp. 108 & 109.
- 273. OPPENHEIM, P. Die Gattungen *Dreyssensia* van Beneden und *Congeria* Partsch, ihre gegenseitigen Beziehungen und ihre Vertheilung in zeit und Raum. Z. geol. Ges. xliii, pp. 923-966, 1 pl.
- 274. . Ueber innere gammenfalten bei fossilen Cerithien und Melaniaden. Op. cit. xliv, pp. 439-446, figg.
- 275. Pallecchi, T. Nota sui cromatofori dei Cefalopodi. Atti Soc. Ligust. iii, pp. 204-214.

- 276. Pantanelli, D. Lamellibranchi pliocenici: Enumerazione e sinonimia delle specie dell' Italia superiore e centrale, Bull. Soc. Mal. Ital, xvii, pp. 49-128. [In progress.]
- 277. Pavlow, A. Argiles de Speeton et leurs équivalents, par A. Pavlow & G. W. Lamplugh. 11. Ammonites de Speeton et leur rapports avec les Ammonites des autres pays. Par A. Pavlow. Bull. Soc. Mosc. 1891 (1892), pp. 455-513, 6 pls. Abstr. in Nature, xlvi, p. 257.
- 278. Pelseneer, P. La classification générale des Mollusques. Bull. Sci. Fr. Belg. xxiv, pp. 347-371, 1 fig.
- 279. —. Le système nerveaux streptoneure des Hétéropodes. C.R. cxiv, pp. 775-777. Abstr. in J. R. Micr. Soc. 1892, p. 352.
- L'Oeil de Scutum. Bull. Soc. Mal. Belg. 1891 [1893], pp. xxviii-xxx.
- L'Hermaphroditisme des Nudibranches sacoglosses (Elysiens, &c.). T. c. p. lv.
- 282. Sur quelques points d'organisation des Nudibranches et sur leur phylogénie. T. c. pp. lxviii-lxxi.
- 283. . Sur la dentrorsité de certains Gastropodes dit "sénestres" (Lanistes, Peraclis, Limacina, et larves des Cymbuliidæ). T. c. pp. xciv-xcvi.
- A propos de l' "Asymétrie des Mollusques univalves." J. de Conch. 1892, pp. 229-233.
- PHISALIX, C. Note sur les Chromatophores des Cephalopodes.
   C.B. Soc. Biol. iv, pp. 442-447, fig.
- 286. PICAGLIA, L. Molluschi terrestri e fluviatili viventi nelle Provincie di Modena e reggio. Catalogo sistematico. Bull. Soc. Mal. Ital. xvi, pp. 83-232.

Includes a table with the distribution of Tertiary species.

- 287. —.. Contributo alla Malacologia fossile dell' Emilia. Molluschi terrestri e fluviatili del Modenese e del reggiano. Atti. Soc. Mod. xi, pp. 157-177 & 226.
- 288. PICARD, K. Ueber Balatonites sondershusanus, n. sp. Z. geol. Ges. xliv, pp. 483-487, 1 pl.
- 289. PILSBRY, H. A. Anatomy of West Indian Helices [Caracolus rostratus, acutus, Hemitrochus salvatoris]. P. Ac. Philad. 1892, pp. 128 & 129, 1 pl.
- 290. —. New and unfigured Unionidæ. T. c. pp. 131 & 132, 2 pls.

  [Notes and figures of Unio quintardii, Cragin, pilsbryi, Marsh, and Arconaia provancheriana, Pilsbry; all old species.]
- 291. A new species of Pachychilus [P. (polygonatus, Lea? var.) rorirosai]. T. c. p. 153, fig.

- 292. [PILSBRY, H. A.] On the anatomy of Sagda, Cysticopsis, Ægista, and Dentellaria. T. c. pp. 213-215, 1 pl.
- A new Marine Gasteropod [Chrysodomus stonei] from New Jersey. T. c. 328 pp., figg.
- 293a. —. Notes on a collection of Shells from the State of Tabasco, Mexico. T. c. pp. 338-341, figg.
- 294. —. Preliminary outline of a new classification of the Helices. T. c. pp. 387-401.
- 295. -—. On a collection of Land Mollusca from the I. of Dominica, West Indies. [With notes on habitats, &c., by G. E. Verrill.] Tr. Conn. Ac. viii [4 pp. Extract].
- Description of a new Hydrobia [H. texana] with notes on other Rissoidæ. P. Davenport Ac. v, 1887, pp. 33 & 34.
- A new species of Zonites [Z. brittsii] from Arkansas. Naut. v, pp. 99.
- 298. —. Note on the group Panda. Op. cit. vi, p. 9.
- 299. —. Observations on the Helices of New Zealand. T. c. pp. 54-57.
- 300. —. On the Amnicoloid genus Lyogyrus, with the description of a new species [L. dalli]. T. c. pp. 61 & 62.
- Observations on the Helicoid group Charopa and allied forms. T. c. pp. 67-69.
- Hadra and Camaena [Reply to Moellendorff, see Zool. Rec. 1891, Moll. p. 19, No. 285]. Nachr. mal. Ges. 1892, pp. 69-73.
  - ---. See also TRYON'S Manual, &c.
  - ---. [See also Johnson & Pilsbry (200).]
- 303. PILSBRY, H. A., & JOHNSON, C. W. Catalogue of Fissurellida of the United States. Naut. v, pp. 102-107 & 113.
- 304. Plate, L. Ueber einiger Organisations verhältnisse der Dentalien. SB. Ges. Marb. 1890 (1891), pp. 26–29.
- 305. —. Ueber den Bau und die systematische Stellung der Solenoconchen. Verh. deutsch. Zool. Ges. i, 1891, pp. 60-63. Abstr. in J. R. Micr. Soc. 1892, p. 465, and Am. Nat. xxvi, pp. 347 & 348.
- 306. —... Ueber den Bau und die Verwandtschaftsbeziehung der Oncidien. Verh. deutsch. Zool. Ges. ii, pp. 30-42.
   With phylogeny of Pulmonata and Opisthobranchiata.
- Weber den Bau und die Verwandtschaftsbeziehungen der Solenoconchen. Zool. JB. v, Anat. pp. 301-86, 4 pls. Abstr. in J. B. Micr. Soc. 1893, p. 31.

- 308. POLLONERA, C. Note su alcuni gruppi di specie del genere Xsrophila. 1. Sul gruppo della X. subprofuga. Boll. Mus. Zool. Tor. vii, No. 128, 18 pp.
  - Ponsonby, J. H. [See Melvill & Ponsonby (250-252).]
- 309. POTEAT, W. L. Notes on the fertility of *Physa heterostropha*, Say. J. Elisha Mitchell Sci. Soc. viii, pp. 70-73. Abstr. in Science, xix, pp. 325 & 326.
- 310. PRUVOT, G. Sur l'embryogénie d'une Proneomenia [P. aglaopheniæ]. C.R. cxiv, pp. 1211-1214. Abstr. in J. R. Micr. Soc. 1892, p. 599.
- 311. RAWITZ, B. Ueber den feineren Bau der hinteren Speicheldrüsen der Cephalopoden. Arch. mikr. Anat. xxxix, pp. 596-611, 1 pl. Abstr. in J. R. Micr. Soc. 1892, p. 595.
- 312. Der mantelrand der Acephalen. Dritte Theil, Siphoniata, &c. Jen. Z. Nat. xxvii, pp. 1-232, 7 pls, and figg. Abstr. in J. R. Micr. Soc. 1892, p. 772.
- 313. Reibisch, P. Die [non marine] conchyliologische Fauna der Galápagos-Inseln. SB. Ges. Isis, 1892, pp. 13-32, 1 pl.
- 314. REIBISCH, TH. Verzeichniss der bisher in den diluvialen Mergeln von Cotta bei Dresden aufgefundenen [non marine] Conchylien. T. c. pp. 8-12.
- 315. Retzius, G. Das sensible nervensystem der mollusken [Limax and Arion]. Biol. Untersuch. N. F. iv, pp. 11-18, 3 pls.
- 316. RIGAUX, E. Notice géologique sur le Bas Boulonnais. Mém. Soc. Acad. Boulogne, xiv [n. spp. described on pp. 104-108, 2 pls].
- 317. RIVERS, J. J. A new volutoid shell [Scaphella (Voluta) arnheimi] from Monterey Bay. Naut. v, pp. 111 & 112.
- 318. ROEBUCK, W. D. Additions to the . . . census of the Land and Freshwater *Mollusca* of Scotland. Ann. Scot. N. H. 1892, pp. 104-107 & 235-238.
- 319. ROGERS, J. On the viviparous nature of Balea. Journ. Conch. vii, pp. 40 & 41.
- 320. Bolle, H. Diagnosen neuer Arten. Nachr. mal. Ges. 1892, pp. 50-52.
- 321. Rosen, O. Beitrag zur Kenntniss der Molluskenfauna Transkaspiens und Chorossans. T. c. pp. 121-126.
- 322. ROSSMARSSLER'S Iconographie der Europaïschen Land- und Süsswasser-Mollusken. Fortgesetzt von D. Kobelt. Neue Folge. Bd. v, pp. 41-118, pls. cxxxi-cl; Bd. vi, pp. 1-48, pls. cli-clx.
- 323. ROTHPLETZ, A. Die Perm-, Trias- und Jura- formation auf Timor und Rotti im indischen Archipel. Palæontogr. xxxix.

  Mollusca, pp. 85-106.

- 324. RUEDIGER, H. Ueber die Silur- Cephalopoden aus den Mecklenburgischen Diluvialgeschieben. Arch. Ver. Mecklenb. xlv, pp. 1-86, 3 pls.
- 325. Sacco, F. I molluschi dei terreni terziarii del Piemonte e della Liguria. . . . Pt. x1. Eulimida e Pyramidellida. Mem. Acc. Tor. xlii, pp. 585-682, 2 pls.
- 326. Sampson, F. A. Mesodon andrewsi in Missouri. Naut. vi, pp. 90 & 91.
- 327. SANDBERGER, F. VON. Verzeichniss der Conchylien des nördlichen Badischen Schwarzwalds. Mal. Bl. xi, pp. 94-100.
  - SANDRI, G. B. [See DANILO & SANDRI (106).]
- 328. SARGENT, F. E. Annotated list of Alabama Land Mollusca. Naut. vi, pp. 76-78.
- 329. SCHARFF, R. F. The Slugs of Ireland. Sci. P. R. Dubl. Soc. vii, pp. 192-195.
  - Abstr. of paper in Sci. Tr. recorded for 1891.
- Some remarks on the Distribution of British Land and Freshwater Mollusca. Conchologist, ii, pp. 1-6.
- 331. —. Note on the affinities of the genera Limax, Arion, and Helix. T. c. pp. 14, 83, & 84.
- The Irish Land and Freshwater Mollusca. Irish Nat. i, pp. 45-47, 65-67, 87-90, 105-109, 135-138, 149-153, & 177-181; also separately, 25 pp. & Index.
- 333. SCHEPMAN, M. M. Land and Freshwater Shells collected by Dr. H. Ten Kate in Scemba, Timor, and other East-Indian Islands. Notes Leyd. Mus. xiv, pp. 145-160, 1 pl.
- 334. Schieldt, R. C. Diffuse pigmentation of the Epidermis [not periostracum] of the Oyster, due to prolonged exposure to the light, &c. P. Ac. Philad. 1892, pp. 350 & 351.
- The Hermaphroditism and Viviparity of the Oysters of the N.W. Coast of the United States. T. c. pp. 351 & 352.
- 336. SCHIELDT, R. C., & RYDER, J. A. On the cause of the Greening of the Oyster, and its presumed Algous endoparasites, T. c. p. 352.
- 337. Schneider, O. Vorläufige mittheilung über die Molluskenfauna von Borkum [with description of Syndosmya schneideri, n. sp., by O. BETTGER]. Nachr. mal. Ges. 1892, pp. 114-117.
- 338. SCHUBERTH, O. Beiträge zur vergleichenden Anatomie des Genitalapparates von Helix mit besonderer Berücksichtigung der Systematik. Arch, f. Nat. lviii, pp. 1-65, 6 pls. Abstr. in J. R. Micr. Soc. 1892, p. 196.
- 339. Scorr, A. [List of Mollusca from the raised beach at Fillyside, between Leith and Portobello. In paper by J. Bennie.] P. R. Phys. Soc. Edinb. xi, pp. 221-237.

- 340. Scott, T. Notes on a collection of Echinoderms and Molluscan Shells found in the Moray Firth District. T. c. pp. 82-84.
- On the Land and Freshwater Mollusca of Bute. P. N. H. Soc. Glasg. iii, pp. 170-181, 1 map.
- 342. Shimek, B. A new species of Freshwater Mollusk [Ancylus obliquus]. Bull. Lab. Iowa, i, pp. 214 & 215, fig.
- 344. SIEMIRADZKI, J. VON. Die oberjurassische Ammoniten-Fauna in Polen. Z. geol. Ges. xliv, pp. 447-482.
- 345. SIMONELLI, V. Sopra le affinità zoologiche della Rothpletzia rudista, Sim. Boll. Soc. geol. Ital. xi, pp. 76-80, figg.
- 346. SIMPSON, C. T. Notes on the *Unionida* of Florida and the Southeastern States. P. U. S. Nat. Mus. xv, pp. 405-436, 26 pls.
- 347. —. Collecting Notes. Naut. vi, pp. 37-40.
- 348. —. On a revision of the American Unionides. T. c. pp. 78-80.
- 349. ——. Contributions to the *Mollusca* of Florida. P. Davenport Acc. v (1887-89), pp. 45-72 & 63°-72°.
  - Descriptions of Pleurotoma simpsoni, Tralia minuscula, and Pandora bushiana, n. spp., by W. H. Dall.
- 350. SIMPSON, G. B. Descriptions of new species of Fossils from the Clinton . . . and Waverly Groups, &c. Tr. Am. Phil. Soc. xvi (1890) [Mollusca], pp. 445-459, figg.
- SIMROTH, H. Ueber einige Raublungenschnecken des Kaukasus.
   Festschr. R. Leuckarts. Leipzig: 1892, 8vo, pp. 48-58, 1 pl., figg.
- 352. . Ueber Kaukasische Limaciden und Testacelliden. Verh. deutsch Zool. Ges. i, 1891, pp. 57 & 58.
- 353. ... Ueber eine anzahl von muscheln welche Herr Pechuël-Losche auf seiner bekannten Expedition an den Congofällen gesammelt hatte. SB. Ges. Leipzig, xviii, pp. 23 & 24.
- 353A. —. Ueber Atopos, &c. T. c. pp. 25-29.
- 354. . Ueber die nackten Limaciden und Testacelliden des Kaukasus. T. c. pp. 40-49.
- 355. ... Ueber die bei Gelegenheit einer Reise nach dem Kaukasus gesammelten Gehäuseschnecken [species determined, and Hyalinia horstii, n. sp., described, by Dr. O. BŒTTGER]. T. c. pp. 49-58.
- 356. —... Ueber eine reihe von Vaginulaarten. T. c. pp. 58-73; Nachtrag, pp. 84-86.
- 357. —. Ueber die pelagischen Gastropodenlarven der deutschen Planktonfahrt. T. c. pp. 98-107.

- 358. [Simroth, H.] Einige Bemerkungen über Bithynella dunkeri, Frauenf. Mal. Bl. xi, pp. 106 & 107, figg.
- 359. —... Hesperarion, eine neue Amerikanische Nachtschneckengattung. T. c. pp. 109-119, figg.
- Einige Bemerkungen zu vorstehendem Aufsatz [Ueber Atopos, Simr., von H. v. Ihering]. Nachr. mal. Ges. 1892, pp. 144–149.
- Einige Punkte aus der œkonomie des Weichthierkörpers, ein Kapitel über Constitution. Leopoldina, 1892 [Mollusca], pp. 122– 125 & 141–144.
  - ---. [See also Bronn, H. G. (46).]
- 362. SLUITER, C. P. Ueber die Bewegung einiger tropischen Mollusken, &c. Tijdschr. Nederl. Dierk. Ver. iii, pp. 170-184, 1 pl.
- 363. SMITH, C. W. Additional Notes on the Land and Freshwater Shells of Cumberland and Westmoreland. Tr. Cumberl. Westmorl. Ass. xvi, pp. 139 & 140.
- 364. SMITH, E. A. On the Land Shells of St. Helena. P. Z. S. 1892, pp. 258-270, 2 pls.
- 365. ——. Descriptions of new species of Shells [Pecten crouchi] from Mauritius and [Mitra fultoni] from California. Ann. N. H. ix, pp. 255 & 256, figg.
- 366. —. On the Shells of the Victoria Nyanza, or Lake Oukéréwé. Op. cit. x, pp. 121-128, 1 pl.
- 366a. ——. Additions to the Shell Fauna of the Victoria Nyanza. T. c. pp. 380-383.
- Further additions to the known Marine Molluscan Fauna of St. Helena. T. c. pp. 129-135.
- 368. ——. Description of a new species of Slug [Apera burnupi] from South Africa. T. c. pp. 465 & 466.
- Motes on Gasteropteron meckelii, Kosse. P. N. H. Soc. Glasg. iii, pp. 258 & 259.
- 370. ——. Description of a new species of Spondylus [S. powelli] and a new Helix [H. hedleyi]. Journ. Conch. vii, pp. 70-73.
- 371. —. Description of a new species of Nucula [N. fultoni], and a list of the species belonging to the subgenus Acila. T. c. pp. 110-112.
- 372. —... Note on a variety and the epidermis [i.e., periostracum] of Cyclophorus sebrinus of Benson. Conchologist, ii, pp. 6 & 7.
- 373. ——. Description of a new species of Acroptychia [A. notabilis]. T. c. pp. 22 & 23.
- 374. —. Notice of an imperforate specimen of Haliotis. T. c. pp. 75 & 76.

- 375. [SMITH, E. A.] Remarks upon the relationship of the Molluscan Fauna of the Red Sea and Mediterranean. Med. Nat. i, pp. 109-111. [Reprint from P. Z. S. See Zool. Rec. 1891, Moll. p. 26, No. 405.]
- 376. Smith, J. List of Shells, &c., observed on the Ardeer and Irvine Beaches, Ayrahire. P. N. H. Soc. Glasg. iii, pp. 243-248.
- 377. SOMERVILLE, J. E. Achatina acicula in a Roman Cemetery at Ventimiglia, Italian Riviera. Journ. Conch. vii, pp. 42 & 43.
- 378. SOWERBY, G. B. Marine Shells of South Africa, &c. London: 4to, pp. iv & 89, 5 pls.
- 379. ——. Descriptions of seven new species of Land Shells from the U.S. of Colombia. P. Z. S. 1892, pp. 296-299, 1 pl.
- Beview of the genus Pyrula (Lamarck), and description of a new species [P. filosa]. Conchologist, ii, pp. 78-75.
- 381. STANDEN, R. Observations on the reproduction of the dart, during an attempt to breed from a sinistral *Helix aspersa*, Müll. Journ. Conch. vii, pp. 33–38.
- 382. STEARNS, R. E. C. Preliminary descriptions of new Molluscan forms from West American regions [Galapagos Is.], &c. Naut. vi, pp. 85–89.
- 383. Stein, J. G. Am. Beiträge zur Mollusken-Fauna Graubünden's.
  11. Nachtrag. J. Ber. Ges. Graub. xxxv, pp. 124-139.
- 384. STERKI, V. Some notes on North American forms of Vallonia.

  Naut. v, pp. 100 & 101.
- 385. ——. Preliminary list of North American Pupidæ (North of Mexica).

  Op. cit. vi, pp. 2-8.
- 386. Hyalinia læviuscula, n. sp. T. c. pp. 53 & 54.
- 387. STONE, W. On Atlantic Crepidulas. T. c. pp. 40-42, figg.
- 388. STRODE, W. S. The *Unionida* of Spoon River, Fulton Co., Illinois. Am. Nat. xxvi, pp. 495-501.
- 389. STRUBELL, B. Landschnecken aus Halmahera [Moluccas]. Nachr. mal, Ges. 1892, pp. 41-50.
- 390. SUTER, H. Contributions to the Molluscan Fauna of New Zealand. Tr. N. Z. Inst. xxiv, pp. 270-278.
- List of the introduced Land and Freshwater Mollusca of New Zealand. T. c. pp. 279-281.
- 391a. List of Land and Freshwater Mollusca doubtful for New Zealand or not inhabiting it [sic: i.e., erroneously attributed to New Zealand]. T. c. pp. 281-283.
- 392. —... Miscellaneous communications on New Zealand Land and Freshwater Mollusca. T. c. pp. 283-286.

- 393. [SUTER, H.] On the Dentition of some New Zealand Land and Freshwater Mollusca, with descriptions of new species. Tr. N. Z. Inst. xxiv, pp. 286-303, 4 pls.
- Conchyliologische Mittheilungen aus Neu-Seeland. Mal. Bl. xi, pp. 101-106, figg.
- Communications Conchyliologiques des Antipodes. 1. Les Pitys de la Nouvelle-Zélande. 2. Un Ancylus de la Nouvelle-Zélande.
   Les Limacidæ et Arionidæ de la Nouvelle-Zélande. J. de Conch. 1892, pp. 245-255.
- 396. SYKES, E. R. On some monstrosities of Littorina rudis, Maton. P. Dorset Field Club, xiii, pp. 191-198, 1 pl.
- Bibliography of the Land and Freshwater Mollusca of Dorset. Conchologist, ii, pp. 24-26.
- Szér, R. Die Molluskenfauna der Umgebung von Güns. Mal. Bl. pp. (27-32 in 1889) 33-41.
- 399. TATE, R. Descriptions of some new species of Marine *Mollusca* from Australia. Tr. R. Soc. S. Austr. xv, pp. 125-132, 1 pl.
- 399A. A third supplement to a list of the Lamellibranch and Paliobranch *Mollusca* of S. Australia. T. c. pp. 133-136.
- 3998. —... The Cambrian Fossils of S. Australia. T. c. pp. 183-189, 1 pl.
- 400. TAUSCH, L. VON. Ueber die Bivalvengattung Conchodus und C. schwageri, n. f. aus der obersten Trias der Nordalpen. Abh. Geol. Reichsanst. xvii, No. 1, 8 pp., 3 pls.
- 401 TAYLOR, G. W. Japanese Limpets. Naut. vi, pp. 69 & 70.
- 402. —. Note on Acmaa saccharina (Linne). T. c. pp. 89 & 90.
- THIELE, J. Beiträge zur kenntniss der Mollusken. I. Ueber das Epipodium. Z. wiss. Zool. liii, pp. 578-590, 1 pl.
- 403a. —. II. Ueber die Molluskenschale. T. c. pp. 220-251, 1 pl. Abstr. in J. R. Micr. Soc. 1893, p. 164.
- Zur phylogenie des Byssus-apparats der Lamellibranchier.
   Verh. deutsch. Zool. Ges. ii, pp. 52-57.
- 405. Das Integument der Chitonen. Biol. Centralbl. xi, 1891, pp. 722-726.
- 406. TIPPENHAUER, L. G. Die Insel Haiti. Leipzig: 1893, 4to [List of Mollusca], pp. 325-327.
- 407. Tomlin, J. B. B. Notes on the Marine Mollusca of the North Wales Coast, with complete lists of the recorded Nudibranchs and Cephalopods. Journ. Conch. vii, pp. 25-31.
- 408. Tomlin, B. [Notes on Risson.] Brit. Nat. 1892, pp. 72-74, 155 & 156, 179 & 180, 257 & 258.

- 409. Tevon. Manual of Conchology...continuation by H. A. Pilsbry. Vol. xiv (pts. 53 & 54), Polyplacophora, 128 pp., 29 pls.
- 410. —... Second Series: Pulmonata, Vol. viii (Helicidæ, Vol. vi), 112 pp., 27 pls. [Oochlostylæ completed—Suppt. to Helix].
- 410a. Tuniot, A. Monstruosités et déformations [of the shell] chez les Mollusques. Bull. Soc. Reims, i, pp. 46-58.
- 411. Tye, G. S. On the periostracum of *Helix arbustorum*, Müll. Conchologist, ii, pp. 53 & 54.
- 412. ULRICH, A. Palaeozoische Versteinerungen aus Bolivien. Steinmann's Beitr. Geol. Palæont. S.-Amer. i, 116 pp., 5 pls. Stuttgart: 1892, 8vo.
- ULRICH, E. O. New Lower Silurian Lamellibranchiata chiefly from Minnesota Bocks. Rept. Geol. Surv. Minnesota, No. 19, pp. 211-248, figg.
- 414. New Lamellibranchiata. Am. Geol. x, pp. 96-104, 1 pl.
- 415. UPHAM, W. Recent fossils of the Harbour and Back Bay, Boston. P. Bost. Soc. xxv, pp. 305-316.
- 416. VAUGHAN, T. W. Mollusks of Dorcheat Bayou and Lake Bisteneau, Louisiana.
  - VERRIL, G. E. [See PILSBRY (295).]
- 417. VIALLANES, H. Recherches sur la filtration de l'eau par les Mollusques et applications à l'Ostréiculture et à l'Océanographie. C.R. cxiv, pp. 1386-1388.
- 418. Vincent, E. Observations sur "Gilbertia inopinata, Mor." Bull. Soc. Mal. Belg. 1891 (1893), pp. lii-lv.

  Reply, see Cossmann (86).
- 419. Description d'une nouvelle espèce de "Pholadidæ" [Aspidopholas stainieri] des Environs d'Anvers. T. c. pp. lxxxvi-xci, figg.
- 420. Contribution à la paléontologie de l'Eocène Belge *Pholadidæ*. Mem. Soc. Mal. Belge, 1891 [1893], pp. 164-168, 1 pl., 1 fig.
- VIZE, J. E. Land and Freshwater Shells [of Hereford]. Tr. Woolhope Nat. 1886-9 (1892), pp. 180-188.
- 422. WACKWITZ, J. Beiträge sur Histologie der Mollusken-Muskulatur, speziell der Heteropoden und Pteropoden. Zool. Beitr. iii, pp. 129–160, 3 pls.
- 423. WALKER, B. The Shell-bearing Mollusca of Michigan. Naut. vi, pp. 13-19, 31-35, 42-47, 63-67, 1 pl.
- 424. Walton, J. The Mollusca of Monroe County, N. Y. P. Rochester Acad. ii, pp. 3-19, 8 pls.
- 425. WARREN, [Miss] A. Contributions towards a list of the Marine Mollusca of Killala Bay, Ireland. Journ. Conch. vii, pp. 98-107.

- 426. WATSON, R. B. The relation of the Land and Freshwater Mollusca of the Madeiran Is. to those known elsewhere. T. c. pp. 1-7.
- 427. —. Note sur l'habitat de l'Ovula carnea, Poiret. J. de Conch. 1892, p. 208,
- 428. WESTERLUND, C. A. Spicilegium Malacologicum. Neue Binnenconchylien in der paläarktischen Region. Verh. z.-b. Wien. xlii, (Abb.), pp. 25-48.
- 429. —. Faunula Molluscorum Hispalensis. An. Soc. Esp. xxi, pp. 381-390.
- Spicilegium Malacologicum. Neue Binnen-conchylien in der paläarktischen Region, III. Nachr. mal. Ges. 1892, pp. 185-201.
  - ——. [For 3 n. spp. of Cacilianella described, see CARUANA-GATTO (60).]
- 431. WHIDBORNE, G. F. A monograph of the Devonian Fauna of the South of England. Vol. ii, pt. 2, pp. 57-88, pls. vi-x (Pal. Soc.).
- 432. WHITEAVES, J. F. The Orthoceratide of the Trenton Limestone of the Winnipeg Basin. Tr. R. Soc. Canada, ix (4), pp. 77-90, 7 pls.
- 433. WILLEM, V. Contributions à l'étude physiologique des organes des sens chez les Mollusques. I. La vision chez les Gastropodes Pulmonés. II. Les Gastropodes Pulmonés perçoivent-ils les rayons ultra-violets? III. Observations sur la vision et les organes visuels de quelques Mollusques, Prosobranches, et Opisthobranches. Arch. Biol. xii, pp. 57-149, 1 pl. Abstr. in J. Micr. Soc. 1892, p. 596.
- 434. WILLIAMS, J. W. The Shells of Stourport. Sci. Goss. pp. 226-228.
- 435. WILLIAMSON, (Mrs.) M. B. An annotated list of the Shells of San Pedro Bay and vicinity. . . . With a description of two new species by W. H. Dall. P. U. S. Nat. Mus. xv, pp. 179-219, 5 pls. WILSON, E. [See HUDLESTON & WILSON (192).]
- Wirén, A. Histologiska meddelanden om Chætoderma nitidulum, Lovén. Biol. Fören. iii, 1890, pp. 37-49.
- 437. WITCHELL, C. A., & STRUGNELL, W. B. The Fauna and Flora of Gloucestershire. Stroud: 1892, 8vo. Mollusca, pp. 171-177.
- 438. WOEHRMANN, S. VON, & KOKEN, E. Die Fauna der Raibler Schichten vom Schlernplateau. Z. geol. Ges. xliv, pp. 167-223, 11 pls. Almost all molluscan. A preliminary note of the new genera of Gastropoda appeared in JB. Mineral. 1892, ii, pp. 25-36. See KOKEN (212).
- 439. WOODWARD, B. B. On the mode of growth and the structure of the Shell in *Velaces conoideus*, Lamk., and other *Neritidæ*. P. Z. S. 1892, pp. 528-540, 2 pls. Abstr. in J. R. Micr. Soc. 1893, p. 30.

- 440. [WOODWARD, B. B.] On the radula of *Paludestrina jenkinsi*, Smith, and that of *P. ventrosa*, Mont. Ann. N. H. ix, pp. 376-378, figg.
- 441. ZIMMERMANN, K. W. Ueber den Kernteilungsmodus bei der Spermatogenese von Helix pomutia. Verh. Anat. Ges. v, 1891, pp. 187-193.

## II.—ANATOMY, PHYSIOLOGY, &c.

### 1. GENERAL.

Comparative anatomy of the Mollusca; LANG (218).-Organization, development, &c., of the Mollusca; Coupin (88).—Some points in Molluscan anatomy; SIMROTH (361).—The asymmetry of Univalve Mollusca is discussed very fully by FISCHER & BOUVIER (131) who in the course of their paper describe and figure the main points in the anatomy and the nervous system of: Chrysodomus contraria, Fulgur perversus, Lanistes bolteniana, and Ampullaria insularum.—Coiling of Univalves: Fischer & Bouvier (132).—Asymmetry of Gastropoda; BUCHNER (51).—Asymmetry of Univalves; Pelseneer (284).—On the dextral nature of certain Gastropoda called "sinistral"; Pelseneer (283).—Anatomical observations on Gastropoda; BOUVIER (39).— Physiological observations on Pulmonata; CUENOT (100A).—Anatomy of Aurinia gouldiana, Trophon (Aspella) scalarioides, Cocculina spinigera, Margarita (Turcicula) imperialis, Pleurotomaria quoyana, P. adansoniana, and of the Pelecypoda Septibranchiata; DALL (101) 1889.—Anatomy of parts of Limax natalianus, n. sp.; Mikhaelis (254).—Anatomy of Ariolimax columbianus, Gould, and Hesperarion [n. g.] niger, Coop.; SIMBOTH (359).—Anatomy of Pseudomilax, Selenochlamys, Trigonochlamys; SIMROTH (351).—Anatomy of Panda atomata, Gray; Hedley (177).—Anatomy of Sagda similis, Cysticopsis tenerrima, Ægista platyomphala, Caracolus orbiculata; PILSBRY (292).—Anatomy of Caracolus rostratus, acutus, Hemitrochus salvatoris; PILSBRY (289).-Anatomy of Cryptazeca monodonta; BARROIS (13). — Anatomy of Amphibola; BOUVIER (40).—Anatomy with figg. of Siphonaria gigas, Less.; HALLER (166), who concludes that Siphonaria is an Opisthobranch, ranking next to Umbrella.—Anatomy and morphology of Naticida and Caluptraida; HALLER (167).—Anatomy of Semisinus, Pachychilus, Ceres, Helicina; FISCHER & CROSSE (133).—Anatomy of Scaphopoda; Plate (304, 305, & 307).—Anatomy and histology of Proneomenia sluiteri; HEUSCHER (187).—Histological notes on Chatoderma nitidulum, Loven; WIREN (436).—Some points in the morphology of Pelecypoda; Kellog (205).— Anatomy of Dipsas plicata, Lea; ISHIKAWA (192A).—Anatomy and physiology of Pholas dactylus; DUBOIS (116).—Anatomy of Cuspidaria (Neaera) cuspidata, Olivi; GROBBEN (163).

#### 2. SHELL AND INTEGUMENT.

Formation and growth of the shell in Mollusca, a résumé, including some new observations, and with a very full bibliography; MOYNIER DE VILLEPOIX (267).—Structure and formation of the Molluscan shell; THIELE (403A).—On the mode of production of the calcareous deposit of the shell of Mollusca; MOYNIER DE VILLEPOIX (267).—Development of shell in coiled stage of † Baculites compressus, Say; Brown (47).— Coiling of shell in Mollusca glossophora; Bowell, Sci. Goss. 1892, pp. 127 & 128.—Note on the coiling of the shell in the embryos of Gastropoda; FISCHER, H. (126).—Periostracum of Helix arbustorum; TYE (411).—Repair of shell in Helix aspersa; MOYNIER DE VILLEPOIX (266). -Periostracum of Cyclophorus zebrinus, Bens.; Smith (372).-Inner palatal folds in fossil Cerithiidæ and Melaniidæ; OPPENHEIM (274).— Mode of growth and structure of the shell in Velates convideus, Lanck., and other Neritidæ; WOODWARD (439).—On the integument of Chitons; THIELE (405).—On the margin of the mantle of Acephala siphoniata; RAWITZ (312).—Growth of the shell of the mussels in Lake Wörther (Carinthia); GALLENSTEIN (142).—Pearl of unusual form and colour; MOEBIUS (256).—On the colouration of the tegument of Cephalopoda; JOUBIN (203): The movements of the chromatophores are due to the nervous excitation of their essentially motile amiboid coloured protoplasm, and not to the presence of special motor muscles.—On the chromatophores of Cephalopoda; Phisalix (285), Pallecchi (275).—Diffuse pigmentation of the epidermis [not periostracum] of the oyster due to prolonged exposure to the light; SCHIELDT (334).

#### 3. MUSCULAR SYSTEM.

Epipodium of Mollusca; THIELE (403).—Histology of the muscular system of Heteropoda and Pteropoda; WACKWITZ (422).—Muscle-fibres of Cephalopoda; BALLOWITZ (12).—Micro-structure of the cartilages of Cephalopoda; Zachariadès, CR. Soc. Biol. iii, p. 454.—Muscles of the radular apparatus in Helix; Loisel (232).

#### 4. DIGESTIVE SYSTEM.

Jaws and radula of Nanina, 2 species, Microcystina sappho, Helicarion musgravi, Cristigibba macgregori, Chloritis, 2 species, Hadra broadbenti, Geotrochus, 6 spp., and Succinea simplex, from British New Guinea; Hedley (174).—The radular apparatus in Helix is actuated by one paired and three unpaired muscles; Loisel (232).—Radulæ and jaws of Helix; Schuberth (338)—Dentition of some New Zealand non-marine Mollusca; Suter (393).—Radulæ of Paludestrina jenkinsi and ventrosa; Woodward (439).—On the minute structure of the posterior salivary glands of the Cephalopoda; Rawitz (311).

#### 5. CIRCULATORY SYSTEM.

Blood and lymphatic glands of *Mollusca*; Cuénot (99).—Analyses of the blood of *Mollusca*; Griffiths (162).—A globulin is obtainable from the blood of *Patella vulgata*, which is colourless and contains no metal; this, which has a respiratory value, it is proposed to call achroglobine; Griffiths (159).—Respiratory globulin in the blood of Chitons; Griffiths (160).—On the blood of the *Pelecypoda*; Griesbach (158).—The respiratory value of Hæmocyanin: Cuénot, C.R. cxv, pp. 127 & 128, finds that the blood of *Heliz pomatia* can absorb more oxygen than an equal volume of water.

### 6. RESPIRATORY SYSTEM.

Structure of the gill in Amusium dalli, Dimya argentea, and Arca ectocomata; DALL (101), 1889, who also discusses Pelseneer's classification of the Pelecypoda.

## 7. EXCRETORY AND SECRETORY SYSTEM.

Excretion in the Pulmonata; CUENOT (100).—The alleged anal eye of larval Opisthobranchs is, in fact, the permanent kidney; MAZZARELLI (248).—On the nephridia of Prosobranchs; ERLANGER (120).—On the urinary function of the organ of Keber in Acephala; Leteller (219).—Fischer, H. (125) treats very fully of the morphology of the liver of Gastropoda, and briefly compares it with that of Brachiopoda, Bryozou, and Rotifera. He distinguishes three stages in its evolution, and rummarizes its principal variations. He does not consider it of practical value in classification.—Development of the liver in Paludina; Fischer, H. (127).—A detailed description of the byssus glauds of Barbatia helblingia is given by Slutter (362).—Formation of the byssus in some Pelecypoda; Jobert (199).—On the phylogeny of the byssus-apparatus of the Pelecypoda; Thiele (404).

#### 8. NERVOUS SYSTEM.

Sensory nervous system of *Mollusca* [Limax and Arion]; RETZIUS (315).—Innervation of the Cerata of some *Nudibranchiata*; HERDMAN & CLUBB (184).—Heteropoda are streptoneurous; Pelseneer (279).—Nervous system of Nerita polita; BOUTAN (38).—Analyses of the nervous tissue of some *Mollusca*; GRIFFITHS (161).

### 9. SENSORY ORGANS.

Vision in the *Pulmonata*, *Opisthobranchiata*, and *Prosobranchiata*; WILLEM (433).—The alleged anal eye of larval Opisthobranchs MAZZARELLI (248) finds is, in point of fact, the permanent kidney.—On the eye of *Scutum*; Pelseneer (280).

Position and character of the Osphradia in Acmea; DALL (101) 1889.

### 10. GENERATIVE ORGANS.

Genitalia of Nanina, 2 species, Chloritis lesi, Hadra broadbenti, and Geotrochus, 6 species, from British New Guinea; Hedley (174).—Anatomy of the genital apparatus of Helix; Schuberth (338).—Morphology and systematic value of the genital apparatus in Helix; Jhering (195).—On the mode of nuclear division in the development of the Spermatozoa of Helix pomatia; Zimmermann (441).—Hermaphroditism of Nudibranchiata sacoglossa; Pelseneer (281).—Oysters of the N. W. Coast of the United States are hermaphrodite and viviparous; Schieldt (335).

### 11. EMBRYOLOGY.


Development of Cephalopoda; Korschelt (214).—Differentiation of the germinal layers in Cephalopoda; Korschelt (215).—Notes on the embryology of the Pteropoda [in Russian]; Knipovich (210).—Development of Bythinia tentaculata; Erlanger (118).—Development of Capulus hungaricus; Erlanger (119).—Cleavage of the ovum in Crepidula fornicata; Conklin (81).—Embryology of Chiton; Metcalf (253).—Embryogeny of Proneomenia; Pruvot (310).—Development of Dreissen[si]a polymorpha; Korschelt (213).

#### 12. PHYLOGENY AND CLASSIFICATION.

Pelseneer (278) classifies the Mollusca as follows:-

I BUSENEER (2	io) crassines me w	Ottuscus da Toriows .—	
	· · · ·		
	Streptoneura	( A spidobranchia	{ Rhipidoglossa. { Docoglossa.
Gastropoda		Ctenobranchia	{ Platypoda. { Heteropoda.
	Streptoneura  Euthyneura	, Opisthobranchia [including Ptero- poda].	∫ Tectibranchia. Nudibranchia.
•		Pulmonata	Stylommatophora. Basommatophora.
Scaphopoda		Solenochoncha.	
Lamellibranchia			ia.

The phylogenetic relationships of these groups are expressed in the following tree:—



Pelseneer concludes that the affinities of the *Mollusca* are to be sought in the *Polychata errantia*, and that, of existing forms, the *Eunicida* seem, from the *ensemble* of their organization, the least removed from the *Mollusca*.

Classification of Cephalapoda; BLAKE (25).—Morphology and Phylogeny of †Stephanocerus sigsac series; BUCKMAN (53).—Phylogenetic relationships of the species of †Olicostephanus from the Specton Clay; PAVLOW (277).

#### VII. MOLLUSCA.

PILSBRY (294) invites criticism on the following proposed classification of the Helices:-

Group I. MACROON.

Genus Acavus.

Genus Stylodonta.

Genus Helicophunta.

Group II. BELOGONA.

Genus Helix.

Genus Leucochroa.

Genus Allognathus.

Genus Cochlostyla.

Genus Polymita.

Genus Hemitrochus.

OGenus Glyptostoma.

Genus Acanthinula.

Genus Valonia.

Group III. TELEOPHALLA.

Genus Sagda.

Genus Cysticopsis.

Group IV. EPIPHALLOPHORA.

Genus Caracolus.

Genus Camana.

Genus Camanella.

Genus Obba.

Genus Chloritis.

Genus Papuina.

OGenus Planispira.

Group V. HAPLOGONA.

Genus Polygura.

Genus Eudodonta.

Genus Patula.

Genus Trochomorpha.

Genus Anoglypta.

Group VI. POLYPLACOGNATHA.

Genus Punctum.

Genus Laonia.

### GENERA OF DOUBTFUL POSITION.

In addition to those marked o above, the systematic position of the following genera is doubtful :--

> Strobilops. Ampelita.

Pedinogyra.

Polygyratia. Macrocyclis. Salaropsis.

VON JHERING (195) recognizes 7 genera in Helicidæ, which he arranges phylogenetically thus:-





PLATE (306), p. 42, thus expresses the phylogeny of the *Pulmonata* and *Opisthobranchiata*:—



The Nudibranchiata are Pleurobranchæa specialized by the loss of a gill. The Tritoniidæ are the oldest forms, from these spring the Doridiidæ (through the Polyceridæ) and the Eolidiidæ, of which last the Elysiæ (or Sacoglossa) are specialized forms; Pelseneer (282).

BERGH (17) classifies the Gasteropoda Nudibranchiata as follows:-

## A. NUDIBRANCHIATA CLADOHEPATICA.

Fam. Aeolidiadæ. Fam. Tethymelibidæ. Fam. Lomanotidæ. Fam. Dotonidæ.

Fam. Dendronotida. Fam. Bornellida. Fam. Scyllaeidæ, Fam. Phylliroidæ. Fam. Pleurophyllidiadæ. Fam. Pleuroleuridæ, Fam. Tritoniadæ.

#### B. NUDIBRANCHIATA HOLOHEPATICA.

Fam. Dorididæ cryptobranchiatæ.

Subfam. Bathydorididæ. Subfam. Hexabranchidæ.

Subfam. Archidoridida.

Subfam. Discodorididæ.

Subfam. Diaululidæ.

Cuolam. Diagrama

Subfam. Cadlinidæ.

Subfam. Kentrodorididæ.

Subfam. Platydoridida.

Subfam. Chromodorididæ.

Subfam. Miamirida.

(Dorididæ cryptobranchiatæ).

Fam. Porostomata.

Subfam. Doriopsidæ.

Subfam. Phyllidiada.

Fam. Dorididæ phanerobranchiatæ.


Subfam, Polyceradæ.

Subfam. Goniodoridæ.

(Dorididæ phanerobranchiatæ).

Fam. Corambidæ.

Systematic position of Naticidæ and Calyptræidæ; Haller (167).— Tables of phylogenetic relationships of Eulimidæ and Pyramidellidæ from the Tertiaries of riedmont and Liguria; Sacco (325). The Scaphopoda are considered by PLATE (307) to have branched off from the Prorhipidoglossa after the Pelecypoda and before Patella; he expresses his conclusions thus:—



Classification of the *Pelecypoda*, based partly on the gill-structure, partly on the hinge characters; GROBBEN (163, 164).

Origin of the genera of non-marine Mollusca; COOKE (82).—Origin of some varieties of Mollusca; FAURE (124).—Classification of varieties of Helix hortensis and H. nemoralis; EYRE (121), MacDonald (234).

## 13. STRAY NOTES.

Habits of the Mediterranean Argonaut with figure of position when swimming; LACAZE-DUTHIERS (217) .- Burrowing habits of certain land Molluscs; Collinge (78).—Perforation of limestone rocks by Lithodomus; CARAZZI (57).—Comparison of movements of rotation in Articulata and Mollusca; Petit, Act. Soc. L. Bord. xliv, pp. 349-351.-Fertility of Physa heterostropha, Say; POTEAT (309).—Effects on a Snail of a median section of its brain; Petit, Act. Soc. L. Bord. xlv, pp. 119 & 120, 1 pl.—Phosphorescence in Latia lateralis, Gould; SUTER (394).—Filtration of water by Mollusca; Viallanes (417).—Appearance of Mollusca in isolated ponds; REID, Tr. Norw. Soc. v, pp. 279 & 280. -Locomotion (Bewegung) of certain tropical Mollucca, more especially byssiferous examples; SLUITER (362).—Disappearance of Spondylus goderopus and other species from Maltese waters; CARUANA-GATTO (59). -W. A. GAIN, Brit. Nat. 1892, has a series of notes on the subject of varieties. He begins by deprecating the naming of slight variations from the normal forms, p. 15: in another part, p. 283, he places on record two worthless variations as new to Britain.—Monstrosities and deformations

in the Molluscan shell; Tuniot (410A).—Monstrosities of Littorina rudis, Maton; Sykes (396).—Effects of freezing and desiccation on Snails; Kochs, Biol. Centralbl. xii, pp. 337 & 338.—Dissolution of the shells of Mollusca in the digestive cavity of fish and the bearings of this on fossilization; Chaper, Bull. Soc. Géol. xx, pp. 114-117.—"Instinct and genera"; Hemphill (181).—Parasites of freshwater Mollusca of Cairo; Sonsino, in Festschr. B. Leuckarts, pp. 134-146, 1 pl.—Cercarideum of Helix hortensis and its development into Distomum; Blochmann, CB. Bakt. Parasit. xii, pp. 649-652.—Mollusca consumed as food by Birds; Loens, Zool. Gart. xxxiii, pp. 49-51.

### 14. Economics.

Uses of shells among the Papuans; Hedley (172A).—Paludina juponica, Mart., for sale in the Chinese market, San Francisco; Wood, Naut. v, p. 114.—Japanese Anodonta, brought to San Francisco markets; Wood, Naut. vi, p. 51.—Freshwater Mollusca in the kitchen middens of Florida; Moore, Am. Nat. xxvi, p. 914.—Application to Oyster culture of the filtration of water by Mollusca; Viallanes (417).

### 15. Collection, Preservation, Museums, &c.

Instructions for collecting, &c.; Dall (105).—Method of taking Water-snails; LOENS (227).—Collecting notes; Simpson (347).—Shell collecting at Eastport, U.S.; ROPER, Naut. vi, pp. 49 & 50.—Notice of the collection of H. A. Ward; PILSBRY, Science, xx, p. 369.—Schlüter's Catalogue a simple list: the generic and subgeneric values attributed to him cannot stand; FISCHER (129A).

### III.—DISTRIBUTION.

#### GEOGRAPHICAL.

#### NON-MARINE MOLLUSCA.

#### a. PALEARCTIC ZONE.

New palearctic species; Westerlund (428 & 430).

# 1. Septentrional Region.

British Isles: The Conchological Society's List, Journ. Conch. vii, pp. 49-66.—Remarks on distribution in the British Isles; SCHARFF (330).—Catalogues of British Slugs; Collinge (80): summary of ib.; Cockerell (72 & 74).—Review of British Arionida; Collinge (79): and criticism by Cockerell (70).—A list of species and varieties of non-marine Mollusca, alleged to be peculiar to the British Islands, occurs

in Wallace's Island Life, 2nd edition, pp. 357-359, republished by the compiler; Cockerell, Nature, xlvi, pp. 76 & 77: see also Scharff, t.c., p. 173.—Cumberland: non-marine of; Smith (363).—Derbyshire: additions to list; Adams (2).—Devon: additions to the S. Devon list; Oldham (272).—Dorset: bibliography of non-marine forms; Sykes (397).—Gloucestershire: list; Witchell & Strugnell (437).—Herefordshire: list; Boycott (41), Vize (421).—Merionethshire: specimens taken in; Chaster (63).—Northamptonshire: bibliography; Crick (92).—Sussex: bibliography, Conchologist, ii, pp. 30-33.—Westmoreland: non-marine of; Smith (363).—Worcestershire: shells of Stourport; Williams (434).

Scotland: additions to the Scotch fauna; ROEBUCK (318).—Species in Bute; Scott (341).—List of shells from Eigg; McMurtre (237).

Ireland: Irish non-marine; SCHARFF (332).—Planorbis riparius, West, added to the British fauna from Lough Swilly; MILNE, Irish Nat. 1892, p. 192 [misidentification, = P. nautileus, Linn.].—Helix lampra, alleged new species [= H. ericetorum]; WESTERLUND (428).

Channel Isles: Pupa ringens in Guernsey; MARQUAND, Journ. Conch. vii, p. 44.

Belgium: Anodonta Ihotellerii, n. sp.; DROUET (115).

France: Mollusca of the Seine-Inférieure; BUCAILLE (50).—A lexia armoricana, n. sp. [?], W. coast of France; Locard (222).—Mollusca of the Department of the Somme; BIZET (24).—List of bivalve Mullusca collected in the Seine near Elbeuf [of no value]; Lancelevée, Bull. Soc. d'Elbeuf, x, pp. 51 & 52.

Germany:—Baden: list of the shells of the North Baden Black Forest; 
Sandberger (327).—Bavaria: Unio clessini, n. sp., from the Danube 
[i.e. the Donan] at Lapingen; Drouer (115).—Borkum Is.: list of 30 
species, including Alexia myosotis, Drap., new to Germany, and Assiminea 
grayana, Leech; Schneider (337).—Brandenburg: occurrence of Helice 
ericetorum and H. candicans at Landsberg-a-Warthe; Krause, SB. nat. 
Fr., 1892, p. 141; rare Mollusca of Mark Brandenburg; Martens (242).—Saxony; supplement to list of Mollusca from near Leipzig; Ehrmann 
(117),—Silesia; notes on 6 species; Loens (228).—Westphalia: 
Molluscan fauna; Hesse (185).—Gastropoda of Münsterland; Loens 
(230).—Supplemental list of 30 species; Loens (229).—Wurttemburg: 
Molluscan fauna of Saulgau; Moenig (259).

Sweden: Hyalinia upsalensis, n, sp., WESTERLUND (428).

Switzerland: Helix (Patula) chaperi, n. sp., Westerlund (428).—Recent Mollusca of Appensell and St. Gallen; Martens (240).—Molluscan fauna of Graubundens; Stein (383).—Mollusca found at Meiringen; Horsley (190).—List of 46 species, with varieties, from Richisau (Canton Glarus); Blum (27).—Mollusca of Pays-d'Enhaut, Vaudois; Pitter & Gétaz, Bull. Soc. Vaud. xxvii, pp. 209 & 210.

# 2. Circum-Mediterranean Region.

Species of Helix (Xerophila), with 20 new species; Pollonera (308). Spain: Helix, 4 new species; Cionella disparata, n. sp.; Westerlund (428).—Helix, 5 new species: Pupa retracta, Buliminus hispalensis, Cionella virginea, Physa clathrata, Melanopsis harpa, Neritina mixta, n. spp.; Westerlund (430).—Mollusca of the Pyrenees, French and Spanish; Fagor (122).

France: Mollusca peculiar to the extreme S. W. of France; Folin (135).—Pupa, 4 new species from the Pyrenean Depts.; Westerlund (428).—Malacological fauna of Carcasonne; Baichère (10).—Mollusca of the Peninsular of Sainte-Lucie (Aude); Fagor (123).—Alleged new species of Lartetia, Moitessieria, Bithinella, and Acme, from near Avignon; Nicolas (269).—Belgrandia coutagnei, n. sp. [?], Avignon; Locard (223).—Terrestrial Mollusca of other Isles in the roadstead of Marseilles; Coutagne (89).

Italy: Pupa honesta, n. sp.; Westerlund (430).—Helix neptuni, n. sp.; Westerlund (428).—Catalogue of species in and around the Province of Modena; Picaglia (286).—List of species from Lake Lugano and its shores; Broemme (45).—Species from Ascoli-Piceno; Mascarini (247).

Sardinia: Helix modestissima, n. sp.; WESTERLUND (428),

Capri Is.: List of 26 species; BLUM (26).

Sicily: Hyalinia, 2 new species; Helix musthorella, n. sp.; WESTER-LUND (428).—Pupa refuga, Pomatias circticus, n. spp.; WESTERLUND (430). —Pseudamnicola nebrodensis, n. sp. [= Vestitu, Benoit]; KOBELT, in ROSSMAESSLER (322) v.

Malta: Clausilia melitensis, n. sp., CARUANA-GATTO (58); Coccilianella, new species, id. (60); Occurrence of Assiminea littorina, id. (61).—Cionella, 3 new species; Westerlund (430).

Austro-Hungary:—Carinthia: Assodonta, 2 new species [?]; KOBELT, in Rossmaessler (322) vi: Growth of the shell of the mussels in Lake Wörther; Gallenstein (142).—Carniola: Clausilia vestigans, n. sp., Westerlund (428).—Dalmatia: Pupa appelinsi, n. sp.; Westerlund (428).—Hungary: Mollusca of the environs of Güns, 69 species; Szép (398).—Siebenburgen: Clausilia denbeli, n. sp.; Westerlund (428).—Bosnia: List of species with Zonites tenerrimus, Helix osoria, Delima optabilis, n. spp. (1890), Brancsik (42); Bythinella bosniensis, n. sp., Clessin (65); Clausilia calabacensis, n. sp., Westerlund (428); Helix broemmei. n. sp., from Acarnania, Kobelt, in Rossmaessler (322) v; Pseudammicola euboica, n. sp., Eubosa; Kobelt, in Rossmaessler (322) v.

Greece: Unio thiessæ, n. sp., Anodonta, 3 new species; DROUET (115).—
Helix giurica and Clausilia broemmei, n. spp., with notes on other species
and some new varieties from Greece and Giura Is.; BŒTTGER (32).
—Corfu: Unio helenæ, n. sp.; DROUET (115).—Sofrano Is. [? Sofrana
Nisia]: Buliminus mestus, n. sp.; WESTERLUND (428).

Crete: Hyalinia spratti, Helix medea, n. spp.; WESTERLUND (428).

Roumania: Buliminus montandoni, n. sp., from Dobrudscha; WESTER-LUND (428).

Caucasus: rapacious Mollusca; SIMROTH (351): Limacidæ, with 4 new species, and Testacellidæ; SIMROTH (352): Testaceous Mollusca, with Hyalinia horsti, n. sp.; BETTGER in SIMROTH (355).

Asia Minor: Buliminus movradi, ponsonbyi, n. spp.; Westerlund (428).

Syria: Helix, 2 new species, Clausilia calopleura, n. sp.; WESTER-LUND (428).

Palestine: Helix patriarcharum, Buliminus granulatus, n. spp.; Westerlund (428).

Egypt: Helix, 2 new species; WESTERLUND (428).—Libya: Helix apaturia, n. sp.; WESTERLUND (428).

Tunis: Helix hypsa, n. sp.; WESTERLUND (428).

Algeria: Helix, 4 new species, Buliminus djurdjurensis, n. sp., Cionella, 2 new species; Westerlund (428).

Morocco: Cionella extrema, n. sp.; Westerlund (428).

## 3. Central Asiatic Region.

Transcaspia: 11 forms enumerated, including 3 new species and 2 new varieties; Rosen (321).—Turkestan: Limax natalianus from Semiretschje; Mikhaklis (254).

# 4. Chinese Region.

Mollusca from the Valley of the Blue River, with a large number of new species; Heude, 1890 (186); Moellendorf (257) criticises Heude.—Hainan Is.: 4 species added to fauna; Fischer (130).

# 5. Japanese Region.

Liu-Kiu Is.: List of 19 species from the Middle Is., Ennea densecestata and Trochomorpha fritsei, n. spp.; BETTGER (33).

# 6. Atlantidean Region.

Madeira: relation of the non-marine Mollusca to those found elsewhere; Watson (426).

## b. PALEOTROPICAL AFRICAN ZONE.

Twelve new species from N. E. Africa; MARTENS (243).

# 7. Central African Region.

Becuanaland: Buliminus layardi, n. sp.; Melvill & Ponsoner (251).—Tanganika: illustrations of freshwater Mollusca from Lake Tanganika; Bourguignat (36) 1888: Cyclostoma burtoni, n. sp.;

ANCEY (5).—Uganda: Helix bellula and Ennea stuhlmanni, n. spp.; MARTENS (241).—Usambara: Ennea grossa, n. sp.; MARTENS (244).—Victoria Nyansa: Limnæa nyansæ, Physa trigona, n. spp., Vivipara, 3 new species from Victoria Nyansa; MARTENS (241): Paludina constrica and Spatha subæquilatera, n. spp.; MARTENS, Conch. Mittheil. iii, 1889: shells from Victoria Nyansa, with Viviparus, 3 new species; Smith (366): ib. with Ampullaria, 2 new species, Planorbis victoriæ and Sphærium nyansæ, n. spp.; Smith (366a).

# 8. West African Region.

Pseudoglessula, n. sect., Stenogyra, with 2 new species; BETTGER (34).—Angola: Buliminus quisqualis, n. sp.; Melvill & Ponsonby (251).—Cameroons: Achatina pyramidella, Ennea conospira, Cyclophorus preussi, n. spp.; Martens (244).—Congo: Stenogyra badia, n. sp.; Martens, Conch. Mittheil. iii, 1889.—Dahomey: Perideris lechatelieri, n. sp.; Dautzenberg (108).—Damaraland: Pupa ovampoensis, Stenogyra chapmani, Cionella ovampoensis, n. spp., from Ovampo-Land; Melvill & Ponsonby (251).—Gaboon: Achatina dimidiata, n. sp.; Martens, Conch. Mittheil. iii, 1889.—Senegambia: Ennea kendigiana, n. sp.; Rolle (320).—St. Helena: land shells, with 11 new species; Smith (364).

## 9. South African Region.

Ennea, 7 new species, Erope, 2 new species, Vitrina, 2 new species, Macrocyclis, 2 new species, Helix farquhari, n. sp., H. (Pella), 11 new species, H. (Trochozonites) dioryæ, n. sp., Planorbis segmentina, n. sp.; Melvill & Ponsonby (250-252).—Apera burnupi, n. sp.; Smith (368).—Ennea consobrina, n. sp.; Ancey (6).

# 10. Malagasy Region.

Madagascar: Acroptychia notabilis, n. sp.; Smith (373): Nossi-Bé Is., 3 new species; Bœttger (31).—Mauritius: notes on shells from the; BAKER (11).

#### c. Palæotropical Oriental Zone.

## 12. Indian Region.

Specimens from Karachi; ADAMS (1).—Ceylon: Vaginula sarasinorum from Candy; SIMBOTH (356): Unio corbeti, n. sp., from Lake Kaudy; DESCHAMPS (112).

# 13. Indo-Chinese Region.

Catalogue with geographical distribution of Indo-Chinese species; Fischer (128): 5 new species, land Mollusca; Morlet (264).—Assam:

Helix fultoni, n. sp., from the Khasi Hills (?); GODWIN-AUSTEN (150): Diplommatina, 26 new species from the Garo, Naga and Munipur Hills; GODWIN-AUSTEN (149).—Cambodia: Ampullaria brohardi, n. sp.; GRANGER (154).—Laos and Tonkin: 10 new species; MORLET (263 & 264).—Singapore: Vaginula carusi, n. sp.; SIMBOTH (356).

## 14. Indo-Malay Region.

List of Indo-Malay genera which extend to the Philippines; COOKE (83).—Borneo: Unio, 6 new species, Pseudodon, 2 new species; DROUET & CHAPER (115A).—Java: Vaginula, 5 new species; SIMROTH (356).—Somba, Timor, &c.: Non-marine Mollusca, 6 new species; SCHEPMAN (333).—Sumatra: Nanina dohertyi, n. sp.; ALDRICH (3): Vaginulu sumatrensis and weberi, n. spp.; SIMROTH (356).—Timor Laut: list of 20 species including 15 new; MOELLENDORFF (258).

# 15. Philippine Region.

Philippines: geographical distribution of the land forms; COOKE (83): Vaginula chavesin, n. sp., from Mindanao; SIMROTH (356): Unio serradelli, n. sp.; DROUET (115).

### d. Australian Zone.

# 16. Austro-Malay Region.

Celebes: Vaginula graffi, n. sp.; SIMROTH (356).—Saleijer Is.: Vaginula vivipara, n. sp.; SIMROTH (356).—Moluccas: 9 species from Halmahera (3 new species Helix, 2 new species Leptopoma); STRUBELL (389): Vaginula bocasei, n. sp.; SIMROTH (356): Acochlidium amboinense and paradonum, nom. nud. for two freshwater species from Amboina; STRUBELL, SB. niederrhein. Ges. xlix, p. 62.—New Guinea: list of a few species collected by Sir W. MacGregor on the Fly River; Hedley (172): Jaws, radulæ, and genitalia of some British New Guinea land Mollusca, Hedley (174); Phelix hedleyi, n. sp., SMITH (370).—Solomon Is.: Helix, 2 new species, Bulim [ul]us hobsoni, n. sp., Hedleya macleayi, n. g. & sp.; Cox (90): Helix aggiei, n. sp.; Heimburg (180): Caliacis exigua, Ad. & Angas, erroneously described as from these Is., is an Australian form and a syn. for Perrieria australis, Forbes sp.; Hedley (176).

# 17. Australian Region.

Remarks on the slugs; Hedley (178), Cockerell (68): Helix everardensis, elderi, n. spp., Elder River, Central Australia; Bednall (14).—Queensland: notes on a collection of shells, with 3 new species; Hedley & Musson (179).—Tasmania: observations on freshwater shells; Johnston (201): variability of the Tasmanian Unio; id. (202).

## 18. Austro-Polynesian Region.

New Hebrides: Diplomorpha cozi, n. sp.; HARTMAN (169).

## 19. Polynesian Region.

Sandwich Is.: revision of the genera Pupa (with 3 new species), Cacilianella (1 new species), and Helicter (2 new species); ANCEY (8A).

## 20. Neozelanian Region.

New Zealand: lists of Mollusca; SUTER (390): list of introduced species; id. (391): list of doubtful species and forms erroneously recorded; id. (391A): miscellaneous notes; id. (392): dentition of some forms with Amphidoxa olivacea and Phrixgnathus transitans, n. spp.; id. (393): notes on land Mollusca; SUTER (394): notes on New Zealand Helices; PILSBRY (299, 410): notes on Pitys, occurrence of Ancylus, on the Limacidæ and Arionidæ; SUTER (395): Ptychodon magdalenes, n. sp., 1891; ANCEY (5).

### e. NEANTARCTIC ZONE.

## 21. Patagonian Region.

Paraguay: Vaginula, 3 new species; SIMROTH (356).

# 22. Chilian Region.

Chili: Vaginula chilensis, n. sp.; SIMROTH (356).

# f. NEOTEOPICAL ZONE.

# 23. Peruvian Region.

Galapagos Is.: Non-marine Mollusca, with Bulimulus, 1 new sect. and 14 new species, Pupa, 2 new species, Succinea wolfs and Helicina wolfs, n. spp.; Reibisch, P. (313): Zonites baueri, Leptinuria chathamensis, Helicina nesiotica; Dall (104): Bulimus habeli, Onchidium lesliei, n. spp.; Stearns (382).

## 24. Columbian Region.

United States of Columbia, 7 new species; Sowerby (379).—New Granada: Helix sieversi, n. sp.; Martens, Conch. Mittheil. iii (1889).

# 25. Brazilian Regi n.

Bolivia: Happia dallianu, Guppya anguina, Cyclophorus orbignyi, Odontostomus lemoinei, n. spp., and Helicina, 3 new species; ANCEY (7):

Nenia orbignyi, n. sp.; ANCEY (6): Cyans orbignyi, Nenia orbignyi, Odontostomus lemoinsi, n. spp.; ANCEY (8).—Brazil: Systrophia alcidiana, Bulimulus germaini, Helicina leucozonalis, n. spp.; ANCEY (7).

## 26. Mexican Region.

Melaniidas and Proserpinidas; FISCHER & CROSSE (133).—Central America: Omphalina 2, Hyalinia 3, Guppya 4, Pseudohyalina cidariscus, Helix altispira, and Polygyra 2 new species; Martens (245).—California: Catalogue of non-marine species of Lower California; Cooper (84).—Guatemala: Neritina sargi and Unio usuniasintas, n. spp.; Crosse & FISCHER (98).—Honduras: Aperostoma goldfussi, n. sp.; Bættger (34).—Mexico: Bulimulus chaperi and Anodonta chapalensis, n. spp.; Crosse & FISCHER (98): Patula intonsa, n. sp., from Orizaba; PILSBRY, in TRYON'S Manual (410): Eucalodium compactum and Potamanax rovirosai, n. subg. & n. sp., from Tabasco; PILSBRY (293A): Pachychilus rovirosai, possible new species; PILSBRY (291).

## 27. Caribbean Region.

West Indies: anatomy of Caracolus, 2 species, and Hemitrochus salvatoris; PILSBRY (289).—Dominica Is.: notes on a collection of land Mollusca, including Vaginula punctatissima, Semp., new to loc.; PILSBRY (295).—Haiti: list of Mollusca; TIPPENHAUER (406).—Jamaica: Helix vendryesi, Veronicella jamaicensis, virgata, dissimilis, alleged new species; COCKERELL (67).—Portorico: catalogue with 130 species; CROSSE (93).

## g. NEARCTIC ZONE.

## 28. American Region.

N. American forms of Vallonia; STERKI (384): Pupida of N. America; STERKI (385): North American species of Succinea; Cockerell (73): notes on Unio luteolus and radiatus; DEAN (110).—Alabama: list of land Mollusca; Sargent (328): Helix sargenti, n. sp.; Johnson & Pilsbry (200).—Arkansas : Zonites britteii, n. sp.; Pilsbry (297).— Florida: list of Mollusca (Pleurotoma simpsoni, Tralia minuscula, and Pandora bushiana, n. spp. by DALL) SIMPSON (349): Lyogyrus dalli, n. sp.; PILSBRY (300): Florida Unionida with Unio subluridus, n. sp.; SIMPSON (346).—Illinois: Unionida of Spoon River; Strode (388).—Iowa: Ancylus obliquus, n. sp.; Shimek (342).—Louisiana: Mollusca of Dorcheat Bayou and Lake Bisteneau; VAUGHAN (416).—Michigan: shell-bearing Mollusca; WALKER (423). - Missouri: occurrence of Mesodon andrewsi; SAMPSON (326).—New York: Mollusca of Monroe County; Walton (424): preliminary list of the Unionida; Marshall (238).—Ohio: catalogue of non-marine Testacea of Portage County; DENN (111).—Texas: Hyalinia laviuscula, n. sp.; Sterki (386); Hydrobia texana, n. sp. (1887); PILSBRY (296).

## 29. Californian Region.

Pupida of North America; STERKI (385): North American species of Succinea; Cockerell (73).—California: occurrence of Limax agrestis, Linn.; BAYMOND, Naut. v, p. 101.—Arizona: occurrence of Patula strigosa, Gould; STEARNS, Naut. vi, p. 1.

## 30. Canadian Region.

North American forms of Vallonia; STERKI (384): North American species of Succinea; COCKERELL (73).—New Brunswick: addenda to the Mollusca; GANONG (145).

### MARINE MOLLUSCA.

### 1. Arctic Province.

Mollusca from East Spitzbergen, 71 species, including Pleuroleura walteri, n. sp.; KRAUSE (216).

#### 2. Boreal Province.

[None.]

### 3. Celtic Province.

British Isles: Mollusca of the Liverpool Marine Biological Committee District; Archer (9): Nudibranchia at Puffin Is.; Herdman, Tr. Liverp. Biol. Soc. v (1891), p. 31: Observations on Lomanotus genei, Verany, and Hancockia sudactylota, Gosse; Gamble (144): occurrence of Pleurophyllidia loveni in Britain; M'Intosh (236), Herdman (183), Griesbach, Conchologist, ii, p. 72: Rare Mollusca on the Devon Coast; Garstang (146): marine invertebrate fauna of Plymouth; Garstang, J. Mar. Biol. Ass. ii, pp. 335-337: Mollusca of N. Wales Coast; Tomlin (407): Mollusca of the Moray Firth; Scott (340): list of shells on Ardeer and Irvine beaches, Ayrshire; Smith, J. (376): Mollusca from the Coasts of Donegal and Dublin; Hart (168): list of species from Killala Bay, Ireland; Warren (425).

France: Mollusca of the Department of the Somme; BIZET (24).

Germany: Syndosmya schneideri, n. sp., from Borkum Is.; BETTGER in Schneider (337).

### 4. Lusitanian Province.

Distribution of Ovula carnea, Poiret; FISCHER, P. (129), WATSON (427).—Atlantic: Cratena fructuosa, n. sp., and Pleurobranchillus morosus, n g. and sp.; Bergh (18).—Azores: Mollusca obtained by the Prince of Monaco; Dautzenberg (107) 1889; many new species: Euplocamus

atlanticus, n. sp.; Bergh (18).—Maderia: Spondylus powelli, n. sp.; SMITH (370).—West Coast of France: Mitra aquitanica, found on the West Coast of France; LOCARD (226): Clathurella bourguignati, servaini, n. spp., 1891; LOCARD (220): Neptunea nicolloni, n. sp.; LOCARD (220A): Anomiida with Anomia boletiformis. n. LOCABD (221): Chama nicolloni, n. sp.; DAUTZENBURG (109).—Spain: Cephalopoda of the Spanish Coasts; GIRARD (148).-Mediterranean: Vermetida, with 3 new species of Vermetus; Monterosato (262).— South Coast of France: Molluscan fauna of Bandol (near Toulon); CAZIOT (62): Anomiida; LOCARD (221).—List of Mollusca of the neighbourhood of Zara; Danilo & Sandri (106).—Kalymos Is.: Pollia reticulata, asona, Alvania rufopunctata, Gibbula eliadum, and valentini, n. spp.; BETTGER (35).—Malta: Disappearance of Spondylus gaderopus, &c.; CARUANA-GATTO (59).

## 5. Aralo-Caspian Province.

[None.]

## 6. West African Province.

Mogador: Lepidopleurus granoliratus, n. sp.; TRYON'S Manual (409).—Octopus sponsalis and ergasticus, n. spp., W. coast, off the Sahara littoral; FISCHER, P. & H. (134).—Senegambia: Pisania scholvieni and Turritella maltsani, n. spp.; Rolle (320).

# 7. South African Province.

Catalogue, with 1 new genus and 43 new species; Sowerby (378).—St. Helena: additions to the fauna, with Jeffreysia atlantics and Tellimya producta, n. spp.; SMITH (367).

## 8. Indo-Pacific Province.

Mauritius: notes on shells; Baker (11): Pecten crouchi, n. sp.; Smith (365).—Red Sea: remarks on the fauna; Jousseaume (204).—Persian Gulf: pearl Mollusca; Galletly (143).—India: Nucula fultoni, n. sp., Bay of Bengal; Smith (371).—Indo-China: Catalogue with geographical distribution of Indo-Chinese species; Fischer (128).—Gulf of Siam, W. of Cambodia, 27 species; Crosse & Fischer (95).—Hongkong: Ischnochiton asperior and bisculptus, n. spp.; Tryon's Manual (409): Pyrula filosa, n. sp.; Sowerby (380).—Liu-Kiu Is.: list of 167 species, Monilea fritzei new; Bættger (33).—Philippines Is.: Ischnochiton solidior, n. sp.; Tryon's Manual (409).—Australia: Cephalopoda and Pteropoda; Brazier (44).

#### 9. Australo-Zelandic Province.

Australia: Ischnochiton cariosus and haddoni, n. spp.; Tryon's Manual (409).—Descriptions of 1 new genus and 11 new species of marine

Mollusca from South Australia; TATE, (399) 3rd Suppl. to list of Australian Pelecypoda; id. (399A).—Glaucus atlanticus, Forster, washed ashore, New S. Wales; Hedley (173).—Cephalopoda and Pteropoda of Australia and Tasmania; Brazier (44).—Tasmania: Callochiton lobatus, n. sp.; Tryon's Manual (409): Columbella unisulcata, n. sp.; Kobelt, in Martini & Chemnitz (246).—New Zealand: Spongiochiton productus, n. sp.; Tryon's Manual (409).

## 10. Japonic Province.

Japan: Volutharpa salmiana, n. sp.; Rolle (320): Thylacodes medusæ, n. sp.; MacGeorge (235): Japanese Patellidæ; Taylor (401): Lepidopleurus curvatus and (?) Ischnochiton asperior, n. sp.; Tryon's Manual (409).—Bonin Is.: Ischnochiton multidentatus, n. sp.; Tryon's Manual (409).

## 11. Aleutian Province.

[None.]

## 12. Californian Province.

Fisurellidæ of the United States; PILSBRY & JOHNSON (303).—California: Loligo stearnsii, n. sp. [P]; HEMPHILL (182): Margarita (Turcicula) bairdii, n. sp.; DALL (101) 1889: Uvanilla regina, n. sp.; STEARNS (382): Chætopleura gemmea, n. sp., and Ischnochiton, 9 new species; TRYON'S Manual (409).—Lower California: Mitra fultoni, n. sp.; SMITH (365): Lepidopleurus rugatus, n. sp., and Ischnochiton, 2 new species; TRYON'S Manual (409).—Monterey Bay: Scaphella arnheimi, n. sp.; RIVERS (317).—San Pedro Bay: catalogue of shells from (Vitrinella williamsoni and Amphissa bicolor, n. spp., DALL); WILLIAMSON (435).

### 13. Panamic Province.

Ecuador: Littorina atyphus, n. sp., Manta; STEARNS (382).—Galapagos Is.: Nitidella incerta and Littorina galapagiensis, n. spp.; STEARNS (382).

14. Peruvian Province.

[None.]

15. Antarctic Province.

[None.]

16. Patagonian Province.

Gorriti Is.: Chatopleura armillata, n. sp.; TRYON'S Manual (409).

### 17. Caribbean Province.

Descriptions of species dredged in the Gulf of Mexico, including many new; Dall (101) 1889: Fissurellidæ of the United States; Pilsbry & Johnson (303): List of Mollusca (with Pleurotoma simpsoni, Tralia minuscula, Natica fordiana, and Pandora bushiana, n. spp., by Dall); Simpson (349).—Florida Keys: Ischnochiton floridanus, n. sp.; Tryon's Manual (409).—Haiti Is.: list of Mollusca; Tippenhauer (406).

### 18. Transatlantic Province.

Descriptions of species dredged, including many new; Dall (101) 1889: Fissurellidæ of the United States; Pilsbry & Johnson (303).—New Jersey: Coast shells, with figs.; Ford (136): Chrysodomus stonei, n. sp.; Pilsbry (293): Crepidulæ, Stone (387).

#### GEOLOGICAL.

Summary of Palæontological work done during 1890: Cephalopoda, by E. Haug, Gastropoda, by M. Cossmann; Ann. Geol. Univ. pp. 957-1024.

Palsontology of New Guinea; ETHERIDGE, in THOMPSON'S "British New Guinea," London, 1892, 8vo, pp. 208-215.

Fossil types in the Bath Museum; P. Bath N. H. Soc. vii, pp. 270-279.

## Quaternary.

Pleistocene non-marine Mollusca found in the Admiralty section; ABBOTT, P. Geol. Ass. xii, p. 356.—List of Mollusca from the Pleistocene deposits at West Wittering, Sussex; REID, Q. J. Geol. Soc. xlviii, p. 357.—Notes on marine shells from the Post-pliocene deposits of Manchester; ROEDER, Tr. Manch. Geol. Soc. xxi, pp. 607-619.-Mollusca from raised beach at Fillyside, between Leith and Portobello, N.B.; SCOTT (339).—Malacological fauna of recent deposits at Brussels; RAEYMARKERS, Bull. Soc. Mal. Belg. 1891 [1893], pp. vii-xi.—Postglacial Mollusca, with Rissoa purva, at Aland, Sweden; MADSEN, Geol. Fören. Stockholm Förhandl. xiv, pp. 585-590.—List of non-marine Mollusca from the alluvial marks of Cotta, near Dresden; Reibisch, T. (314).—Silurian Cephalopoda from the alluvial beds of Mecklenburg; RUEDIGER (324).—Pliocene and quaternary Molluscan fauna of the environs of Oran; Pallary, CR. Ass. Fr. Sci. 1891, ii, pp. 383-387, figs.—Recent fossils of the Harbour and Back Bay, Boston; UPHAM (415).—Fossils of the Loess of E. Nebraska and W. Iowa; SHIMEK, Bull. Lab. Iowa, i, pp. 200-214, ii, pp. 89-93.—Post-tertiary shells collected in a dry salt lake near Eddy, Mexico; STERKI, Rep. Geol. Surv. Texas, iii, pp. 263-265.—Post-tertiary Mollusca of Queensland; JACK & ETHERIDGE (193) pp. 640-646.

### 2. TERTIARY.

Belgium: Aspidopholas stainieri, n. sp., from the Tertiary of Antwerp; VINCENT (419).—France: marine Mollusca of the Basin of the Rhone; Fontannes, Études strat. and paléont. periode tort. dans le Bassin du Rhone, x, Paris, 1892, 8vo.—Germany: older Tertiary fauna of the neighbourhood of Berlin, with new species; KOENEN, JB. k. preuss. geol. Landesanst. xi, Abh. pp. 257-276, figg.—Austro-Hungary: Gastropoda from Häring, Tirol, with 15 new species; DREGER (114): fossil fauna of Markuševec in Croatia, 49 new species, Baglivia, n. g.; BRUSINA (49): Tertiary Rudistes, Ceratoconcha costata, n. g. & sp., 1889, from Agram; Gorjanović-Kramberger (153). — Italy: Upper Tertiary Pectunculus; GREGORIO (155): Non-marine Mollusca of the Province of Modena; PICAGLIA (286).—Tertiary species (many said to be new) of Eulimida and Pyramidellidæ from Piedmont and Liguria; SACCO (325).—Sardinia: Spirulirostrina lovisatoi, n. g. & sp.; CANAVABI (56).—Molluscan fauna of Florida, pt. ii, completing the Gastropoda, Pteropoda, Polyplacophora, and Scaphopoda, with very many new species, and Glyptostyla panamensis, n. g. & sp., from the Tertiary beds of Panama; Dall (102).—New Guines: Tertiary Mollusca; JACK & ETHERIDGE (193) pp. 692-694.

Pliocene: Lists of pliocene Mollusca from the environs of Montpellier; VIGUIER, C.R. Ass. Fr. Sci. 1891, ii, pp. 405-416, 1 pl.—Italy: Pliocene Lamellibranchiata from upper and central Italy; PANTANELLI (276).—Pliocene Mollusca of Cerreto-Guidi and confines, with Conus caroli, Arca ida, and Lepton stefanii, n. spp.; FUCINI, Boll. Soc. geol. Ital. x, pp. 59-86, 1 pl.—Pliocene of Bologna, with Pholadomya elegantula, n. sp.; FORESTI (138).—New Zealand: Pliocene Mollusca; HUTTON, Nature, xlvi, p. 474 [Abstr.].

Miocene: Germany: Pelecypoda from the Miocene of Dingden, with many new species; Lehmann, Verh. Ver. Rheinl. xlix, pp. 198-241, 1 pl.—Austro-Hungary: Sarmatic-miocene shells of Oststeiermark; Hilber, MT. Ver. Steierm. xxviii, pp. 235-248, 1 pl.—Pontic fauna of Southern Hungary, with Cardium semseyi, n. sp.; Halaváts, MT. Ung. geol. Anst. x, No. 2.—Prepontic fauna of Agram; Gorjanović-Kramberger, Glasnik Naravoal. Družt. v, 1890, pp. 154-164, 1 pl.—Miocene nou-marine Mollusca from Samos; Stefani in 'Samos.' By C. de Stefani, C. J. Forsyth Major and W. Barbey. Lausanne: 1891 (1892), 4to, pp. 76-81, 1 pl.

Oligocene: France: revision of the Oligocene fauna of Étampes (contd.) with Odontostomia lamberti and Truncatella bezançoni, n. spp.; Cossmann (87): Fauna of the Oligocene of Narbonne; Viguier, Bull. Soc. Béziere, xiv, p. 72.—Germany: Upper Oligocene fossils of the Malliss Brickearths; Geinitz, Arch. Ver. Mecklemb. xlvi, pp. 65-67, 71 & 72: Natica geinitzi, Calliostoma jetschini, Yoldia beyrichi, from the Rupelian of Germany; Bettger (30): Lower Oligocene Mollusca of N. Germany, with many new species; Koenen (211).—Albania: Oligocene Mollusca; Dreyer, JB. geol. Reichsanst. xlii, pp. 339 & 340.

Eccene: Belgium: Eccene Pholadida; VINCENT (420).—France: Eccene Mollusca from the environs of Paris, with some new genera and sections and many new species; Cossmann (85): lists of Eccene fossils collected near Reims on different excursions; Bull. Soc. Reims, i, pp. 62-64, 84 & 85, 91, 94-97.—Germany: new locality for non-marine Mollusca from the Vincentinian Eccene; Oppenheim & Beyrich, Z. geol. Ges. xliv, pp. 500-503.—Italy: Lower Tertiary Mollusca of the Ligurian littoral; Gregorio (157).

### 3. SECONDARY.

Secondary fossils from the Balkans, with many new species; Vankov, Rad. jugoslav. Akad. cxi, pp. 78-102: Secondary fossils from St. Philippe-de-Benguéla, with Desmoceras cuvervillei, Hamites tropiculis and Natica gabonensis, n. spp.; Stanislas Meunier, Le Nat. 1892, pp. 113 & 114, figg.: new genera and new species of Mollusca from the Secondary rocks of the Argentine Cordilleras; Behrendsen (15): Secondary fossils from Queensland with Pseudavicula and Maccoyella, new genera and many new species; Jack & Etheridge (193), pp. 319 & 32), 388 & 389, 445-503, 561-574: Unio eyrensis, Tate, and sp.? from the Mesozoic of South Australia; Etheridge in Brown, Reports on the Coalbearing area near Leigh's Creek [Govt. Rept.] 1891, pp. 11 & 12; Upper Rognacian of Les Baux, with 3 new species of land shells; Nicolas (270).

Cretaceous: Mollusca from the Cretaceous of Schleswig-Holstein with Dimyodon boehmi, n. sp.; Stolley, MT. min. Inst. Kiel. i, pp. 229-244: fossils of the Cretaceous of the Venetian Alps with new Caprinida; Boehm, Ber. Ges. Freiburg, vi, pp. 139-148, figg.: Cretaceous Mollusca from the S. E. of Spain, with new species of Mortoniceras, &c.; Nicklès, Annales Hébert, i, pp. 187-202, figg.: Cretaceous forms from Albania; Dreger, JB. geol. Reichsanst. xlii, pp. 337-339: list of Cretaceous fossils from the Podol'ski Govt.; Radkevich, Zapiski. Kiev. xi (1891), pp. 82 & 83: Cypraa from the Fox Hills Group; Campbell, Naut. vi, p. 50: on the occurrence of Actinocamax quadratus and Belemnitella mucronata; Strombeck, Z. geol. Ges. xliii, pp. 919-922.

Upper Cretaceous: Upper Senonian fossils of the Eifel; Vogel, Verh. Ver. Rheinl. xlix, pp. 18-87, figg.: palæontology of the Upper Senonian of Irnich, Eifel, with many new species; Vogel, Verh. Ver. Rheinl. xlix, pp. 32-106, 1 pl.: Upper Cretaceous Mollusca of the Venetian Alps, with many new species and Cornucaprina and Orthopsychus n. gg.; Futterer (139): 3 new species of Belemnites (Actinocamax) from the Senonian; Janet (194).

Lower Cretaceous: Ammonites from S. W. France, with 3 new species; KILIAN (208): Neocomian of Dîmbovicioara, Roumania, with descriptions of Molluscan species, none new; Uhlig, JB. geol. Reichsanst. xli, pp. 216-234: Valanginian Hoplites of the group H. neocomiensis; Lory (233).

Tithonian: List of fossils from Aizy-sur-Noyarcy (Isere); Gevrey, Bull. Soc. Stat. Isere (4) i, pp. 49-55.

Jurassie: Catalogue of British Jurassic Gastropoda; Hudleston & Wilson (192): Jurassic Mollusca of Yorkshire, list; Fox-Strangways, Mem. Geol. Surv. U. K., Jurassic Rocks, ii, pp. 45-117, 150-185, 204-245, 249, & 250: Specton Clay Ammonites, with several new species; Paylow (277).—Jurassic Mollusca of Boulogne, with new species; Rigaux (316): Upper Jurassic fossils from Grenoble; Paquier, Bull. Soc. Stat. Isère (4) i, pp. 33-44: Harpoceras mathayense, n. sp., Upper Callovian of Doubs; Kilian (208): Renggerithone [Oxfordian] fauna of Kandern, Baden; Lent, MT. Badischen geol. Landesanst, ii, pp. 623-639: Upper Jurassic Ammonites of Poland, Oppelia stolzmanni, Cardioceras suessi, Persphinctes, 5 new species, Proplanulites teisseyrei, m. spp.; Siemiradzki (344): Mollusca from the Middle Jurassic of Castel Tesino in the S. E. Tirol; Boese & Finkelstein, Z. geol. Ges. xliv, pp. 280-284: Jurassic Mollusca, with many new species from the Is. of Timor and Rotti; Rothpletz (323).

Inferior Oolite: Gastropodu of the British Isles, several new species; HUDLESTON (191): Ammonites of the British Islands, with alleged new species; BUCKMAN (54).

Northampton Sands: occurrence of Ammonites jurensis; Buckman (52).

Liassic: List of fossils from the Lias of Northamptonshire; J. Northampt. Soc. 1892, pp. 45-49: Cephalopoda of the Upper Lias of Sainte-Colombe-lès-Avallon (Yonne); MILLOT (255): Mollusca from the Liassic Ironstone near Helmstedt, with new species; Wollemann, Verh. Ver. Rheinl. xlix, pp. 107-124: Liassic fossils from Sette, with Gervilia ombonii and Pecten taramelli, n. spp.; NEGRI, Boll. Soc. geol. Ital. x, pp. 309-331, 1 pl.

Trias: Triassic Mollusca of the Duchy of Sachsen-Altenburg; GEINITZ, MT. Osterlande, v, pp. 184-189: Gastropoda of the St. Cassian beds, Pt. II, with 2 new genera and 32 new species; KITTL (209): Trias of the Northern Alps, notes on Con-hodus, with C. schwageri, n. sp.; TAUSCH (400): Mollusca from the Alpine Trias, with new genera and species; KOKEN (212) & WOEHRMANN & KOKEN (438): Preliminary remarks on the Cephalopod fauna of the Himalayan Trias; Mojsisovics, SB. Ac. Wien, ci, Abth. i, pp. 372-378; Chemnitzia manzavinii and Schizodus segrii, n. spp., Trias of Balia, Asia Minor; BITTNER (22): Timor and Rotti, Triassic Mollusca, with many new species; ROTHPLETZ (323).

Muschelkalk: Fossils of the Muschelkalk, with Beneckeia cognata, n. sp.; WAGNER, Z. geol. Ges. xliii, p. 898, fig.: Balatonites sondershusanus, n. sp., from the Muschelkalk near Sondershausen; PICARD (288).

### 4. PRIMARY.

Palæozoic Hyolithidæ of Bohemia revised, with new genera and new species; Novák (271).—Check-list of Palæozoic Mollusca from Wisconsin, Minnesota, Iowa, Dakota, and Nebraska; BIERBAUER (20).—

Palsozoic species of the *Platyceras* group; KEYES (206).—11 new species of *Pelecypoda*, 6 new species of *Platyceras*, from the Clinton, Lower Helderberg, Chemung, and Waverly groups of Pennsylvania; SIMPSON (350).—Middle Devonian and Permo-Carboniferous *Mollusca* from Queensland, with *Peltopecten* and *Merismopteria*, n. gg., and many new species; JACK & ETHERIDGE, (193) pp. 69 & 264-296

Permian: Permian Mollusca of the Duchy of Sachsen-Altenburg; Geinitz, MT. Oesterlande, v, pp. 168 & 174-177: Anthracosiæ from the Permian of Russia, Palæomutela and Oligodon, n. gg., with many species; Amalizky (4): Mollusca, with new species from the Permian of the Is. of Timor and Rotti; Rothpletz (323): Fossils from Lower Magnesian Limestone of Iowa, with new species of Straparollus, Raphistoma, and Cyrtoceras; Calvin, Bull. Lab. Iowa, ii, pp. 189-193: Lower Magnesian Limestone fossils from Iowa, with new species of Straparollus, Raphistoma, and Cyrtoceras; Calvin, Am. Geol. x, pp. 144-148.

Carboniferous: Carboniferous Murchisonia, 6 new species; DONALD (113): Carboniferous Pelecypoda—Ctenodonta, Nucula, and Allorisma; GOODCHILD (152, 152A).

Devonian: Devonian fauna of S. England, with new species; Whidborne (431): Devonian Mollusca of the Duchy of Sachsen-Altenburg; Geinitz, MT. Osterlande, v, pp. 168 & 174-177: Ammonites of the Sporadoceras muensteri Zone (Upper Devonian) of Orenburg, with Clymenia, 3 new species; Loewinson-Lessing (231): Amnigenia rhenana, n. sp, from the Middle Devonian of Germany; Beushausen (19): Devonian of the Eastern Alps, with Platyceras conoideum, n. sp.; Frech, Z. geol. Ges. xliii, p. 678: Devonian species from Manitoba, with many new, and Astralites, n. g. (incerta sedis); Whiteaves, Contrib. Canadian Palseont. i, pp. 292-346, figg.: New Devonian fossils of Iowa—Schizodus symmetricus, Holopea grandis, Straparollus lativalvis, bicarinatus, tricarinatus, Bucania perornata, Cyclostoma, n. spp.; Calvin, Bull. Lab. Iowa, i, pp. 176-181, figg.: Clymenia neapolitana, n. sp., Upper Devonian (Naples Beds) of Western New York; Clarke (64): Devonian of Bolivia, with 6 new species; Ulrich (412).

Silurian: Trematonatus from the Wenlock Limestone, near Dudley; Newton (268): Silurian Cephalopoda from the alluvial beds of Mecklenburg, with 3 new species of Endoceras; Ruediger (324): Swedish Silurian, Endoceras hircus and schmidti, n. spp.; Holm (189): Swedish Silurian species of Lituites, with 3 new species, and Cyclolituites lynceus, n. sp.; Holm (188): Orthoceratidæ of the Trenton Limestone of the Winnipeg Basin; Whiteaves (432): New Lower Silurian Pelecypoda of the U. S., with Cleionychia, n. g., and 1 new species, Cypricardites, 3 new species, Technophorus, 3 new species, Tellinomya longa, n. sp.; Ulrich (414): New Lower Silurian Pelecypoda, with Plethocardia, n. g., mostly from Minnesota; Ulrich (413): Raphistoma brownii, n. sp., and Orthoceras sp.?, from the Silurian of South Australia; Etheridge in Brown, Reports on the Coalbearing area near Leigh's Creek [Govt. Rep.], 1891, pp. 9 & 10.

Cambrian: Swedish fossils from the Cambrian Sandstone, with Dentalium? cambricum and Hyvlithus insularis, n. spp.; Moberg, Geol. Fören. Stockholm Förhandl. xiv, p. 117, figg.: Cambrian fossils of South Australia, with 3 new species of Mollusca; Tate, Tr. R. Soc. S. Austr. xv, pp. 183–185.

### IV.—SYSTEMATIC.

- Classification of the *Mollusca*; Pelseneer, Bull. Sci. Fr. Belg. xxiv, pp. 347-371, 1 fig.
- Introduction to the study of the *Mollusca*; COUPIN (88).—Comparative Anatomy of the *Mollusca*; LANG (218).—New edition of BRONN (46) by H. SIMROTH, Lief 1 & 2, Historische Uebersicht.
- On the asymmetry of univalve Mollusca; FISCHER & BOUVIER, J. de Conch. 1892, pp. 117-207, pls.; Pelseneer, t. c. pp. 229-233.—Coiling of univalves; FISCHER & BOUVIER, t. c. pp. 234-245.—See also under Gastropoda.

## I. CEPHALOPODA.

- Résumé of the classification of Cephalopoda, with a note on the Cephalopoda Dibranchiata of the Upper Lias of Sainte-Colombe-lès-Avallon (Yonne); MILLOT, Bull. Soc. Autun, iv, pp. 37-57, figg.—Evolution and classification [chiefly of fossil forms]; BLAKE, P. Geol. Ass. xii, pp. 275-295.—Cephalopoda of the Spanish Coast; GIRARD, An. Soc. Esp. xxi, pp. 391-399.
- Catalogue of the Cephulopoda of Australia and Tasmania; BRAZIER (44).

  On the chromatophores of Cephulopoda; Phisalix, C.R. Soc. Biol. iv, p. 442; Pallecchi, Atti Soc. Ligust. iii, pp. 204-214.—Colouration of the tegument; Joubin, Arch. Z. expér. x, pp. 277-330, figg.
- On the structure of the posterior salivary glands of the Cephalopodu; RAWITZ, Arch. mikr. Anat. xxxix, pp. 596-611, 1 pl.
- Development of the Cephalopoda; KORSCHELT (214).—Differentiation of the germinal layers; KORSCHELT, Verh. Deutsch. Zool. Ges. ii, pp. 87-92.

# DIBRANCHIATA.

### OCTOPODA.

- Argonauta: a living specimen and its habits described, with fig. of position when swimming; LACAZE-DUTHIERS, Arch. Z. expér x, pp. 37-56.
- Octopus sponsalis, p. 297, ergasticus, p. 298, W. Coast of Africa, off the Sahara littoral, P. & H. Fischer, J. de Conch. 1892, figg.

### DECAPODA.

- Table of chief characters of Oigopsid Cephalopods; Goodrich, J. Mar. Biol. Ass. ii, pp. 318-321.
- †Actinocamax grossouvrei, p. 716, toucasi, p. 719, alfridi, p. 720, from the Senonian, Janet, Bull. Soc. Géol. xix, figg., n. spp.
- Loligo stearnsii, possible new species, California? Hemphill, Zoe, iii, pp. 50 & 51.
- Ommastrephes pteropus, Stp., notes on; GOODRICH, J. Mar. Biol. Ass. ii, pp. 314-321, figg.
- † Spirulirostrina, n. g. for S. lovisatoi, n. sp., Tertiary of Sardinia; CANA-VARI, Bull. Soc. Mal. Ital. xvi, p. 65, fig.

## AMMONEA.

- †Amaltheus (?) attenuatus, n. sp., Neocomian of the Argentine Cordilleras; Behrendsen, Z. geol. Ges. xliv, p. 17, fig.
- †Ammonites jurensis: note on occurrence in Northampton Sands; Buck-Man, Geol. Mag. 1892, pp. 258-260.
- †Aspidoceras bodenbenderi, n. sp., Tithonian of the Argentine Cordilleras; Behrendsen, Z. geol. Ges. xliv, p. 13, fig.
- †Baculites compressus, Say, development of shell in coiled stage; Brown, P. Ac. Philad. 1892, p. 136.
- † Balatonites sondershusanus, n. sp., Muschelkalk near Sondershausen; PICARD, Z. geol. Ges. xliv, p. 483, figg.
- †Cardioceras suessi, n. sp., Upper Jurassic of Poland; SIEMIRADZKI, Z. geol. Ges. xliv, p. 455.
- †Clymenia (Cyrtoclymenia) neapolitana, Upper Devonian of Western New York, Clarke, Am. J. Sci. xliii, p. 57, fig.; †C. barbaræ, p. 20, inostranzevi, p. 21, dubia, p. 22, Upper Devonian of Orenburg, Loewinson-Lessing, Mém. Soc. Belg. Géol. vi, figg. : n, spp.
- † Harpoceras (Ludwigia) mathayense, v. sp., Upper Callovian of Doubs; Kilian, Bull. Soc. Stat. Isère (4) i, p. 204, figg.
- † Hoplites neocomiensis: on the species of the group from the Valanginian Marls of Southern Dauphiné and Haute-Provence; Loky, Bull. Soc. Stat. Isère (4) i, pp. 229-255.
- †H. angulatiformis, p. 16, neumayri, p. 17, n. spp., Neocomian of the Argentine Cordilleras; Behrendsen, Z. geol. Ges. xliv, figg.
- †Lytoceras stephanense, p. 2, depersti, p. 4, n. spp., Lower Cretaceons, S.W. of France; Kilian, Arch. Mus. Lyon, v, No. 3, fig.
- † Olcostephanus subdivided into the following groups: Virgatites, Craspedites, Polyptychites, Holcodiscus, Astieria, Simbirskites; PAVLOW, Bull. Soc. Mosc. 1891 (1892), p. 113.

- † Oppelia bukowskii, p. 450, stolsmanni, p. 452, n. spp., Upper Jurassic of Poland; SIEMIRADZKI, Z. geol. Ges. xliv.
- †Perisphincies: list of species with many new forms from the Upper Jurassic of Poland; Siemiradzki, Z. geol. Ges. xliv, pp. 455-478.
- † Phylloceras goreti, Lower Cretaceous of S. W. France, Kilian, Arch. Mus. Lyon, v, No. 3, fig.; † P. homophylum, Inferior Colite of the Argentine Cordilleras, Behrendsen, Z. geol. Ges. xliv, p. 6, fig.: n. spp.
- †Proplanulites teisseyrei, n. sp., Upper Jurassic of Poland, SIEMIRADZKI, Z. geol. Ges. xliv, p. 480.
- †Stephanoceras zigzac series, Morphology; Buckmann, Q. J. Geol. Soc. xlviii, pp. 447-452, 2 pls. To certain of the forms the author gives the names S. subprocerum, clausiprocerum, crassizigzag, pseudoprocerum.

## TETRABRANCHIATA.

### NAUTILIDÆ.

- †Cyclolituites lynceus, Silurian of Sweden, Holm, Geol. Fören. Stockholm Förhandl. xiii, p. 766, fig., n. ap.
- † Endoceras angelini, p. 27, schlieffeni, p. 30, angustisiphonatum, p. 37, Silurian species from the diluvial deposits of Mecklenburg, Ruediger, Arch. Ver. Mecklenb. xlv, figg.; † E. hircus, p. 127, schmidti, p. 129, Holm, Geol. Fören. Stockholm Förhandl. xiv, figg.; † E. crassisiphonatum, Trenton Limestone of the Winnipeg Basin; Whiteaves, Tr. R. Soc. Canada, ix, 4, p. 79, figg.: n. spp.
- †Gonioceras lambii, n. sp., Trenton Limestone of the Winnipeg Basin, Whiteaves, Tr. R. Soc. Canada, ix, No. 4, p. 86, fig.
- †Lituites: notes on the mouth of; Holm, Geol. Fören. Stockholm Förhandl. xiii, pp. 736-774, figg.
- †L. tornquisti, p. 742, discors, p. 759, præcurrens, p. 763, Silurian of Sweden, Holm, Geol. Fören. Stockholm Förhandl. xiii, figg., n. spp.
- †Orthoceras semiplanatum, p. 81, selkirkense, winnipegense, p. 82, n. spp., Trenton Limestone of the Winnipeg Basin, Whiteaves, Tr. R. Soc. Canada, ix, 4, figg.
- † Poterioceras gracile, n. sp., Trenton Limestone of the Winnipeg Basin, Whiteaves, Tr. R. Soc. Canada, ix, No. 4, p. 87, fig.
- †Sactoceras canadense, n. sp., Trenton Limestone of the Winnipeg Basin, Whiteaves, Tr. B. Soc. Canada, ix, No. 4, p. 85, fig.

[PTEROPODA: see OPISTHOBRANCHIATA.]

## II. GASTROPODA.

Anatomical observations; BOUVIER, C.R. Soc. Biol. iv, p. 987.—Asymmetry of Gastropoda; BUCHNER, JH. Ver. Württ. xlviii, pp. 58-92.

—Coiling of the shell in embryonic Gastropoda; FISCHER, H., J. de Conch. 1892, pp. 309-313.

See also ante, GENERAL.

Researches on the morphology of the liver of Gastropoda; FISCHER, H., Bull. Sci. Fr. Belg. xxiv, pp. 260-346, 7 pls.

Pelagic Gastropod larvæ of the Plankton Expedition; Simroth, SB. Ges. Leipzig, xviii, pp. 98-107.

† Monograph of British Jurassic Gastropoda; HUDLESTON (191).

†Catalogue of British Jurassic Gastropoda; Hudleston & Wilson (192).

## PULMONATA.

Physiological observations; Cuénot, Arch. Biol. xii, pp. 683-740, 1 pl.: vision; Willem, t. c. pp. 57-122, fig.—Exerction in the *Pulmonata*; Cuénot, C.B. cxv, pp. 256-258.

Remarks on Australian alugs; HEDLEY, Ann. N. H. ix, p. 169; COCK-ERELL, t. c. p. 370.

### STYLOMMATOPHORA.

### TESTACELLIDÆ.

List of British species; Collinge, Brit. Nat. 1892, p. 178.

Testacellidæ of the Caucasus, notes on; SIMROTH, Verh. Deutsch. Zool. Ges. i (1891), pp. 57 & 58.

(Ærope) Helix trimeni, eumacta, South Africa, MELVILL & PONSONBY, Ann. N. H. x, p. 237, figg., n. spp.

Anceya admirabilis, E. Centr. Africa, BOURGUIGNAT, (37) 1889, p. 119, fig., n. sp.

Apera burnupi, S. Africa, SMITH, Ann. N. H. 1892, p. 466, n. sp.

Colpanostoma leroyi, N'gourou, E. Centr. Africa, BOURGUIGNAT, (37) 1889, p. 48, fig., n. sp.

Edentulina grandidieri, E. Centr. Africa, Bourguignat, (37) 1889, p. 142, fig., n. sp.

Elma mitis, China, HEUDE, (186) p. 152, fig., n. sp.

Ennea consobrina, Albany, S. Africa, ANCEY, Brit. Nat. 1892, p. 125; E. (microstrophia) densecostata, Middle Liu-Kiu Is., Bættger, Nachr. mal. Ges. 1892, p. 159; E. fargesiana, China, Heude, (186) p. 152; fig.; E. stuhlmanni, Uganda, Martens, SB. nat. Fr. 1892, p. 16; E. limbata, excavata, p. 178, planidens, (Ptychotrema) runsorana, p. 179, N. E. Africa, Martens, SB. nat. Fr. 1892; E. grossa, Usambara, conospira, Cameroons, Martens, SB. nat. Fr. 1892, p. 182; E. thelodonta, p. 85, munita, dolichoskia, p. 86, bowkera, maria, p. 92, aperostoma, scrobiculata, p. 93, S. Africa, Melvill & Ponsonby,

- Ann. N. H. ix, figg.; E. (Uniplicaria) kendigiana, Gorée, Senegambia, Rolle, Nachr. mal. Ges. 1892, p. 50 : n. spp.
- Hapalus conoideus, n. sp., Butumbi, N. E. Africa, MARTENS, SB. nat. Fr. 1892, p. 177.
- Marconia, n. g., type Ennea lata, Smith, with M. gibbosa, recta, n. spp., E. Centr. Africa; BOURGUIGNAT, (37) 1889, pp. 135-138, figg.
- Selenochlamys: description and anatomy; SIMROTH, (351) p. 54.
- [Streptaxis] Happiu dalliana, E. Bolivia, ANCEY, Journ. Conch. vii, p. 90;
  [S.] Gonaxis bloyeti, E. Centr. Africa, BOURGUIGNAT, (37) 1889,
  p. 134, fig.; S. simonianus, theæcola, China, Heude, (186) p. 151,
  figg.; S. dugasti, Tonkin, Morlet, J. de Conch. 1892, p. 82: n. spp.
  Streptostile costulata. n. sp., Butumbi, N. E. Africa, Martens, SB. nat.
  Fr. 1892, p. 178.
- Tayloria jouberti, n. sp., Nyantaga, E. Centr. Africa, BOURGUIGNAT, (37) 1889, p. 47, fig.

## LIMACIDA.

- List of British naked species; Collinge, Brit. Nat. 1892, pp. 178 & 179. Limacidæ of the Caucasus, notes on; Simroth, Verh. Deutsch. Zool. Ges. i, 1891, pp. 57 & 58. Limacidæ of New Zealand; Suter, J. de Conch. 1892, pp. 250-255.
- Agriolimax subugrestis, p. 41, agresticulus, n. spp., Caucasus, Simroth, SB. Ges. Leipzig, xviii.
- Charopidæ should form a subfam. of Zonitidæ, and not be placed with Helicidæ; Hedley, P. Linn. Soc. N.S.W. vii, pp. 157-169, figg.
- Euplecta orientalis, n. sp., Timor Laut Is.; Moellendorff, Nachr. mal. Ges. 1892, p. 54.
- Guppya anguina, E. Bolivia, Ancey, Journ. Conch. vii, p. 91; G. championi, p. 119, pittieri, biolleyi, p. 121, orosciana, p. 123, Central America, Martens, Biol. Centr. Am. Moll. figg.: n. spp.
- Helicarion musgravi, Hedley, jaw and radula; Hedley, P. Linn. Soc. N.S.W. vi, p. 687, figg.
- H. tenimbericus, n. sp., Timor Laut Is., Moellendorff, Nachr. mal. Ges. 1892, p. 83.
- Hyalinia: notes on the genus; JHERING, Nachr. mal. Ges. 1892, pp. 132-140.
- H. horsti, Caucasus, Bœttger in Simroth, SB. Ges. Leipzig, xviii, p 50;
  H. glomerula, Guatemala, hoffmanni, Costa Rica, p. 115, stolli, Guatemala, p. 118, Martens, Biol. Centr. Am. Moll. figg.; H. gomezi, Bogota, Sowerby, P. Z. S. 1892, p. 298, fig.; H. læviuscula, Texas, Sterki, Nant. vi, p. 53; H. (Polita) upsaliensis, p. 25, Upsala, bellis, p. 26, senilis, Sicily, (Mesomphix) spratti, Crete, p. 44, Westerlund, Verh. z.-b. Wien (Abh.), xlii: n. spp.
  [See also infrà, Zonites.]
- Limax: sensory nervous system; RETZIUS, Biol. Untersuch. N. F. iv, pp. 11-18, figg. Limax, Arion, and Helix: note on the affinities of; SCHARF, Conchologist, ii, pp. 14, 83, & 84; COCKERELL, t. c. pp.

- 28 & 29. L. agrestis, Linn, in California; RAYMOND, Naut. v, p. 101. L. maximus, L., and its var. cinereo-niger, Wolf, notes on; Collinge, Ann. N.H. x, pp. 425 & 426.
- L. natalianus, n. sp., Semiretechje, Turkestan, Mikhaelis (254).
- Lytopelte transcaspia, besttgeri, n. spp., Transcaspia; Rosen, Nachr. mal. Ges. 1892, pp. 123-124.
- Macrochlamis simplex, n. sp., China; HEUDE, (186) p. 135, fig.
- (Macrocyclis) Helix canotera, liparoxantha, n. spp., S. Africa; Melvill & Ponsonby, Ann. N. H. x, p. 238, fig.
- Mesolimax reibischi, n. sp., Caucasus, Simroth, SB. Ges. Leipzig, xviii, p. 43.
- Microcystina sappho, Brazier, jaw and radula; Hedley, P. Linn. Soc. N.S.W. vi, p. 686, figg.
- Moaria chaperiana, n. sp., Usagara, Bourguignat, (37) 1889, p. 14.
- Nanina hunsteini, Smith, divisa, Forbes, jaws, radulæ, and genitalia; HEDLEY, P. Linn. Soc. N.S.W. vi, pp. 685 & 686, figg.
- N. (Ariophanta) dohertyi, Sumatra, Aldrich, Naut. vi, p. 90; N. hypochlora, Cape of Good Hope; Melvill & Ponsonby, Ann. N. H. ix, p. 87, fig.; N. cochlostyloides, Scemba, Schepman, Notes Leyd. Mus. xiv, p. 148, fig.: n. spp.
- Omphalina modesta, p. 110, salkeanu, p. 112, sculpta, p. 113, n. spp., Mexico; Martens, Biol. Centr. Am. Moll., figg.
- Paralimax reibischi, n. sp., Caucasus, Simeoth, SB. Ges. Leipzig, xviii, p. 42.
- Pseudohyalina cidariscus, n. sp., S. E. Mexico, Martens, Biol. Centr. Am. Moll. p. 126, fig.
- Pseudomilax: description and anatomy; SIMROTH, (851) p. 52.
- Sitala brancsiki, filomarginata, n. spp., Is. Nossi-Bé (Madagascar), Bærrger, Nachr. mal. Ges. 1892, pp. 53 & 54.
- Trigonochlumys: description and anatomy; SIMROTH, (351) p. 55.
- Trochonanina leroyi, p. 18, bloyeti, p. 21, n. spp., Usagura, Bourguignat, (37) 1889, figg.
- (Trochozonites) Helix dioryx, n. sp., South Africa, MELVILL & PONSONBY, Ann. N. H. ix, p. 89, fig.
- Vitrina fuscicolor, p. 240, chrysoprasina, p. 241, n. spp., South Africa, Melvill & Ponsonby, Ann. N. H. x, figg.
- Vitrinoides: n. subg. ? of Limax; Simhoth, SB. Ges. Leipzig, xviii, p. 41.
- Xesta micholitsi, n. sp., Timor Laut Is., Mobllendorff, Nachr. mal. Ges. 1892, p. 84, fig.
- Zonites tenerrimus, Bosnia, Brancsik, Evkön. Trencsen Varm. kii, 1890, p. 68; Z. caroliniensis, North Carolina, Cockerell in Binney, Bull. Mus. C. Z. xxii, p. 167, fig.; Z. (Hyalinia) baueri, Galapagos Is., Dall, Naut. v, p. 98; Z. shepardi, p. 167, diegoensis, p. 168, California, Hemphill in Binney, Bull. Mus. C. Z. xxii, figg.; Z. brittsii, Arkansas, Pilsbry, Naut. v, pp. 99: n. spp. [See also supra, Hyalinia.]

## HELICIDE.

New classification of the *Helices* proposed by PILSBRY, P. Ac. Philad., 1892, pp. 387-404.—Supplement to monograph on *Helix*, PILSBRY in TRYON'S Manual, 2nd Series, viii, pp. 55-78.

## ARIONIDÆ.

- List of British Species; Collinge, Brit. Nat. 1892, p. 178.—Review of the British Arionidæ; Collinge, Conch., ii, pp. 56-66 & 76-83: criticised by Cockerell, t. c. pp. 84 & 85: reply; Collinge, t. c. 85 & 86.
- Arionida of New Zealand; SUTER, J. de Conch. 1892, pp. 250-255.
- Arion, Limax, and Helix, note on the affinities of; SCHARFF, Conchologist, ii, pp. 14, 83 & 84: COCKERELL, t. c. pp. 28 & 29.
- Arion: variation of the genus; Collinge, Ann. N. H. ix, pp. 307 & 308: sensory nervous system; RETZIUS, Biol. Untersuch. iv, pp. 11-18, fig.
- Arion hortensis, Fér, and circumscriptus, Johnst., new vars.; Collinge, Conchologist, ii, pp. 26 & 27. A. intermedius, Normand, remarks on colour changes; GAIN, Conchologist, ii, pp. 55 & 56.
- Hesperarion, n. g. for Ariolimax niger, Coop.; SIMROTH, Mal. Bl. xi, p. 111.

### BULIMIDÆ.

- Bulimus oblongus, note on; GUPPY, P. Z. S. 1892, pp. 271 & 272.
- B. (Rachis) jouberti, p. 61, pachistoma, p. 62, elongatulus, p. 63, E. Cent.
  Africa, Bourguignat, (37) 1889, figg.; B. (Amphidromus) columellaris, Timor Laut Is., Moellendorff, Nachr. mal. Ges. 1892, p. 98, fig.; B. guentheri, U. S. of Colombia, Sowerby, P. Z. S. 1892, p. 296, fig.; B. (Mastus) hispalensis, Spain, Westerlund, An. Soc. Esp. xxi, p. 387: n. spp.

#### COCHLOSTYLA GROUP.

- Cochlostyla: monograph completed; PILSBRY in TRYON'S Manual, 2nd Series, viii: sections Orthostylus, Hypselostyla, Eudoxus, Phengus, Canistrum, Prochilus, and Chrysallis.
- C. (Corasia) tenimberica, n. sp., Timor Laut Is., Moellendorff, Nachr. mal. Ges. 1892, p. 97, fig.

#### HELICIDÆ 8.8.

- Hadra and Camaena, reply to Moellendorff; PILSBRY, Nachr. mal. Ges. 1892, pp. 69-73.
- Heliz, Arion, and Limaz, note on the affinities of; SCHARFF, Conchologist, ii, pp. 14, 83 & 84: COCKERELL, t. c. pp. 28 & 29.
- Helix: muscular apparatus of the radula; Loisel, xxviii, pp. 566-572, figg.: anatomy of the genital apparatus with notes on the radulæ and jaws; Schuberth, Arch. f. Nat. lviii, pp. 1-65, figg.: morphology of the genital apparatus; Jhering, Z. wiss. Zool. liv, pp. 386-520, figg.
- Panda: note on the group. PILSBRY, Naut. vi, p. 9, considers it to be a development from the Bulimoid branch instead of the Helicoid stem of Helicoidæ.

- Plecteulota, n. subsect. of Eulota, type E. goniostoma, MOELLENDORFF. [See infrà, Palssotropical Oriental Species, p. 61.]
- Suteria, n. sect. of Helix, type H. ide, Gray; PILSBRY, Naut. vi, p. 56, and TRYON's Manual, 2nd Series, viii, p. 69.
- Thalassohelix, n. sect. of Helix, type H. zelandiæ; PILSBRY, Naut. vi, p. 56, and TRYON'S Manual, 2nd Series, viii, p. 71.
- Xerophila: note on some groups of species of; Pollonera, Boll. Mus. Zool. Tor. vii, No. 128.

### (PALÆARCTIC SPECIES.)

- Helix aspersa: on the repair of the shell; MOYNIER DE VILLEPOIX, Bull. Soc. Z. Fr. xvii, p. 30. H. aspersa, Müll., reproduction of dart; STANDEN, Journ. Conch. vii, pp. 33-38. H. arbustorum, Müll., note on the periostracum; Tye, Conchologist, ii, pp. 53 & 54. H. ericetorum, Müll., and H. candicans, Ziegl., at Landsberg-a-W.; KRAUSE, SB. nat. Fr. 1892, p. 141. H. hortensis and nemoralis, classification of varr.; Eyre, Conchologist, ii, pp. 7-10; MACDONALD, t. c. pp. 27 & 28. H. pomatia, var.; Gredler, Nachr. mal. Ges. 1892, p. 174. H. (Ægista) platyomphala, Mildff, anatomy; Pilsbry, P. Ac. Philad. 1892, p. 214, fig.
- H. (Cressa) giurica, Giura I. [Grecian Arch.], BETTGER, Nachr. mal. Ges. 1892, p. 59; H. osoria, Bosnis, Brancsik, Evkön. Trencsén Varm. xii, 1890, p. 69; H. secundaria, p. 136, serpestes, turbinella, p. 137, vicinella, languescens, molina, p. 138, sapeca, sterilis, p. 143, rhinocerotica, galera, p. 144, subsquamulata, ingloria, vitreola, ternaria, anomia, p. 145, alveolus, p. 146, China, Heude (186), figg.; H. (Campylea) broemmei, Klissura Pass, in Acarnania (Greece), KOBELT in Rossmaessler, (322) v, p. 108, fig.; H. (Xerophila) mentita. Palermo, p. 2, ilvatica, Piedmont, p. 3, trimetina, Is. Pianosa, bulteata, Tudu Mannu, p. 4, warnierella, p. 5, cliens, p. 6, parca, p. 7, Fernana, arida (Monter. MS.), Palermo, p. 7, stossiciana, Trieste, p. 9, trinacrina, Palermo, nerusia, Grasse, p. 11, mendica, Rapallo, p. 14, janalis, Genova, senensis, Siena, p. 15, pistoriana, Pistoia, margarita. Oran. elusatica, S. France, p. 16, bigeronica, Lourdes, festa, Tunis, roucayroli, Algeria, p. 17 [alleged n. spp.], POLLONERA, Boll. Mus. Zool. Tor. vii, No. 128; H. (Xerophila) mutua, pumila, adina, horridula, embryonata, (Macularia) leucochila, Spain [alleged n. spp.], WESTER-LUND, Nachr. mal. Ges. 1892, pp. 187-191; H. (Xerophila) mutua, p. 385, pumila, horridula, embryonata, p. 386, (Macularia) leucochila, p. 387, Spain [alleged n. spp.], WESTERLUND, An. Soc. Esp. xxi; H. (Xerophila) patriarcharum, Palestine, clausella, Spain, bougsoulensis, Algeria, p. 28, mesopotamica, Syria, batuensis, marconi, Algeria, promissa, Spain, p. 29, pinguis, Alexandria, p. 30, masthorella, Sicily, modestissima, Sardinia, neptuni, Italy, p. 31, euaxes, Gibraltar, poichila, Algeria, amblia, Spain, p. 32, lampra [= ericetorum], Ireland, p. 33, mexensis, Egypt, (Pomatia) beilanica, Syria, (Cressa) medea, Crete, hypsa, Tunis, p. 45, (Xeroleuca) apaturia, Libya, p. 46 [alleged n. spp.] WESTERLUND, Verh. z.-b. Wien, xlii: n. spp.

## (PALÆOTROPICAL AFRICAN SPECIES.)

- Helix melvilliana, n. n. for H. chapmani, preocc., W. Africa; ANCEY, Brit. Nat. 1892, p. 126.
- H. bellula, Uganda, Martens, SB. nat. Fr. 1892, p. 15; H. karewia, Karewia, N. E. Africa, Martens, SB. nat. Fr. 1892, p. 175; H. farquhari, Port Elizabeth, Melvill & Ponsonby, Ann. N. H. x, p. 240, fig.; H. (Pella) trichosteiroma, epetrima, p. 84, lygaa, p. 85, rhysodes, p. 87, tuguriolum, erateina, p. 88, bathycoele, p. 89, S. Africa, Melvill & Ponsonby, Ann. N. H. ix, figg.; H. (Pella) actinotricha, p. 238, burnupi, conisalea, p. 239, minythodes, p. 240, S. Africa, Melvill & Ponsonby, Ann. N. H. x, figg.: n. spp.

### (PALÆOTROPICAL ORIENTAL SPECIES.)

H. (Plectopylis) fultoni, Khasi Hills?, Assam, Godwin-Austen, Ann. N. H. 1892, p. 300; H. (Eulota) hemisphaerica, p. 90, tenimberica, bitaeniata, p. 91, fig. 1, (Plecteulota) goniostoma, p. 92, fig., telitecta, p. 93, fig., Timor Laut Is., Moellendorff, Nachr. mal. Ges. 1892; H. (Chloritis) rhodochila, micholitzi, tenuitesta, Timor Laut Is., Moellendorff, Nach. mal. Ges. 1892, pp. 95 & 96, figg.; H. (Plectotropis) hyperteleia, Laos, p. 82, (Phania) dugasti, Tonkin, p. 83, Morlet, J. de Conch. 1892; H. (Ohloritis) remoratrix, Laos, Morlet, J. de Conch. 1892, p. 317, fig.; H. supracostulata, Soemba, Schepman, Notes Leyd. Mus. xiv, p. 150, fig.: n. spp.

#### (Australian and Polynesian Species.)

- Helix (Chloritis) leei, Cox, chloritoides, Pilsbry, jaws and radulæ, with genitalia of leei; Hedley, P. Lind. Soc. N.S.W. vi, pp. 687 & 688, figg. H. (Cristigibba) macgregori, Hedley, jaw and radula; Hedley, P. Lind. Soc. N.S.W. vi, p. 687, figg. H. (Geotrochus) brumeriensis, Forbes, louisiadensis, Forbes, rollsianus, Smith, woodlarkianus, Souv., trobriandensis, Hedley, boyeri, Fisch. & Bern., jaws, radula, and genitalia; Hedley, P. Lind. Soc. N.S.W. vi, pp. 689-691, figg. H. (Hadra) gulosa, Gould, synonymy of; Brazier, P. Lind. Soc. N.S.W. vi, pp. 321-328. H. (Hadra) broadbenti, Brazier, jaw, radula, and genitalia; Hedley, P. Lind. Soc. N.S.W. vi, p. 688, figg. H. (Panda) atomata, Gray, note on structure and affinities; Hedley, Rec. Austral. Mus. ii, pp. 26-31, 3 pls.
- H. (Hadra) everardensis, elderi, Elder River, Central Australia; Bednall, Tr. R. Soc. S. Austr. xvi, pp. 64 & 65, figg.; H. (Hadra) oscarensis, p. 565, derbyi, p. 566, Solomon Is., Cox, P. Linn. Soc. N.S.W. vi, figg.; H. aggiei, Solomon Is., Heimburg, Mal. Bl. xi, p. 158, figg.; H. (Geotrochus) hedleyi, New Guinea?, Smith, Journ. Conch. vii, p. 72; H. (Planispira) halmaherica, p. 44, (Geotrochus) chondrodes, p. 45, (Albersia) pseudocorasia, p. 47, Halmahera [Moluccas], Strubell, Nachr. mal. Ges. 1892; H. (Amphidora) oliracea, Auckland, Suter. Tr. N. Z. Inst. xxiv, p. 290, fig.: n. spp.

## (NEOTROPICAL SPECIES.)

- H. (Caracolus) rostratus, acutus, anatomy; PILSBRY, P. Ac. Philad. 1892, pp. 128 & 129, fig. H. (C.) orbiculata, Fér., anatomy; PILSBRY, P. Ac. Philad. 1892, p. 214, fig. H. (Cysticopsis) tenerrima, C. B. Ad., anatomy; PILSBRY, P. Ac. Philad. 1892, p. 214, fig. H. (Hemitrochus) salvatoris, anatomy; PILSBRY, P. Ac. Philad. 1892, p. 129, fig. H. (Sagda) similis, C. B. Ad., anatomy; PILSBRY, P. Ac. Philad. 1892, p. 213, fig.
- H. creveauxiana, W. Columbia, ANCEY, Bull. Soc. Mal. Fr. vii, 1891, p. 152; H. (Systrophia) alcidiana, Brazil, ANCEY, Journ. Conch. vii, p. 91; H. (Lucerna) vendryesi, Jamaica, Cockerell, J. Inst. Jamaica, i, p. 55 [nom. preocc. and sp. not new, = H. sloaneana, Shuttlw. var.]; H. altispira, Honduras, Martens, Biol. Centr. Am. Moll. p. 156, fig.; H. (Polygyra) dissecta, p. 167, richardsoni, p. 168, Mexico, Martens, Biol. Centr. Am. Moll. figs.; H. sieversi, New Granada, Martens, Conch. Mittheil. iii, 1889, p. 7, fig.: n. spp.

### (NEARCTIC SPECIES.)

- [Helix] Vallonia: notes on N. American forms; STERKI, Naut. v, pp. 100 & 101. H. (Mesodon) andrewsi in Missouri; SAMPSON, Naut. vi, p. 90.
- H. sargenti, n. sp., Woodville, Alabama, Johnson & Pilsbry, Naut. vi, p. 8.

### (FOSSIL FORMS.)

- Grandipatula, n. sect. of Helix, type † H. hemisphærica, Mich.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iv, 1889, p. 354.
- Sagdellina, n. sect. of Helix, type † H. chevallieri, Cossm.; Cat. Coqu. Foss. Eocene Paris, iv, 1889, p. 353.

### (PHENACOHELICIDÆ.)

Under Phenacohelicidæ, SUTER (Tr. N. Z. Inst. xxiv, p. 271) includes the following genera of New Zealand heliciform Mollusca:—Phacussa, Thalassia, Gerontia, Psyra, Pyrrha, Therasia, Phenacohelix, Patulopsis [nom. nov. for Charopa, Hutton], Amphidoxa, and Calymna. The family is characterized as animal heliciform, possessing a caudal gland; jaw ribbed, stegognath; teeth of radula show all intermediate forms, from aculeate to quadrate; shell, that of Helix, flat or elevated, umbilicated or imperforate; peristome acute, aperture without lamellæ; periostracum smooth or ribbed, seldom hairy. Phacussa stands nearest to Limacidæ, Amphidoxa and Calymna to Patulidæ. [Pilsbry, P. Ac. Philad. 1892, pp. 401 & 402, includes this group in the genus Endodonta, as restricted by him.]

### PATULA GROUP.

- Aeschrodomus, n. n. for Thera (sect. of Patula); PILSBRY, Naut. vi, p. 55, note.
- Charopa and allied forms: notes on; PILSBRY, Naut. vi, pp. 67-69.
  Charopids are not Helicids, but should form a subfamily of Zonitida; 4 species are redescribed and figured; HEDLEY, P. Linn. Soc. N.S.W. vii, pp. 157-169.
- Diaglyptus, n. sect., type P. pagodiformis, Smith (= P. bilamellata, Pfr.); Pilsbry in Tryon (410) p. 86.
- Endodonta leimonias, Gray, poscilestica, Pf., are removed to Phrixgnathus (Laoma); Suter, Tr. N. Z. Inst. xxiv, p. 271.
- Helenoconcha, n. sect., type P. bilamellata, Sow.; PILSBRY in TRYON (410) p. 91.
- Patula: Supplement to monograph; PILSBRY in TRYON (410). P. strigosa, Gould, in Arizona; STEARNS, Naut. vi, p. 1.
- P. (Discus) brunnescens, Timor Laut Is., MOELLENDORFF, Nachr. mal. Ges. 1892, p. 87; P. (Microphysu) intonsa, Mexico, PILSBRY in TRYON (410) p. 111, figg.; P. persoluta, latissima, p. 261, vernoni, pseustes, p. 262, minutissima, leptalea, p. 264, St. Helena, SMITH, P. Z. S. 1892, figg.; [P.] Helix chaperi, Switzerland, WESTERLUND, Verh. z.-b. Wien, xlii, p. 27; n. spp.
- Phrizgnathus transitans, New Zealand; SUTER, Tr. N. Z. Inst. xxiv, p. 297, fig.
- Pitys: notes on; Suter, J. de Conch. 1892, pp. 245-248.
- Ptychodon magdalenæ, n. sp., New Zealand, Ancey, Brit. Nat. 1891, p. 65.

  Trochomorpha fritzei, n. sp., Middle Liu-Kiu Is., Bættger, Nachr. mal.

  Ges. 1892, p. 160.

### BULIMULIDA.

- Amphidromus latestrigatus, p. 151, Soemba, reflexilabris, p. 152, Timor, Schepman, Notes Leyd. Mus. xiv, figg., p. spp.
- Bulimulus germaini, Brazil, Ancey, Journ. Conch. vii, p. 92; B. (Placostylus) hobsoni, Solomon Is., Cox, P. Linn. Soc. N.S.W. vi, p. 567, figg.; B. chaperi, Mexico, Crosse & Fischer, J. de Conch. 1892, p. 296; B. (Naesiotus) invalidus, venustus, p. 17, pallidus, p. 18, cinereus, ventrosus, p. 19, acutus, p. 20, curtus, nudus, p. 21, wolfi, p. 22, simrothi, p. 23, (Pleuropyrgus) terebra, p. 24, lina (Pelecostoma, n. sect.), canaliferus, p. 25, cymatoferus, p. 26, Galapagos Is., Reibisch, SB. Ges. Isis, 1892, figg.; B. subtruncatus, turtoni, St. Helena, Smith, P. Z. S. 1892, p. 266, figg.; B. koppeli, da-costa, glandiniformis, Bogota, Sowerby, P. Z. S. 1892, p. 297, figg.; B. (Pleuropyrgus) habeli, Galapagos Is., Stearns, Naut. vi, p. 86 (& v, pp. 98 & 99): n. spp.
- Diplomorpha coci, n. sp., New Hebrides, HARTMAN, P. Linn. Soc. N.S.W. vi, p. 571, fig.
- Partnla: catalogue of the genus begun by HARTMAN, Naut. vi, pp. 73-76. Pelecostoma, n. sect. of Bulimulus [q.v.].

### BULIMINIDE and PUPIDE.

Pupidæ of North America [= U. S.]; STERKI, Naut. vi, pp. 2-8.

Balea, viviparous nature of; Rogers, Journ. Conch. vii, p. 40.

Buliminidius lurvatus, pupatus, n. spp., China, Heude (186) p. 146, figg.

Buliminopsis quaternarius, conoidius, n. spp., China, Heude, (186) p. 147,

figg.

Bulim [in] us lourdeli, p. 51, bridouxi, p. 53, Usagara, Bourguignat, (37) 1889, figg.; Buliminus aureus, p. 148, meronianus, meleagrinus, p. 149, guilcherianus, vidianus, antisecalinus, amedeanus, loliaceus, p. 150, China, Heude (186), figg.; B. trapezoideus, Mt. Runsoro, N. E. Africa, Martens, SB. nat. Fr. 1892, p. 176; B. layardi, Kobis (Becuanaland), quisqualis, Moçambie (Angola), lamoensis, Lamo, E. Africa, Melvill & Ponsonby, Ann. N. H. ix, p. 90, figg.; B. (Amphiscopus) continens, Transcaspia, Rosen, Nachr. mal. Ges. 1892, p. 125; B. (Mastus) hispalensis, Spain, Westerlund, Nachr. mal. Ges. 1892, p. 192; B. (Napaeus) djurdjurensis, Algeria, (Chondrulus) montandoni, Dobrudscha, movradi, Asia Minor, p. 35, granulatus, Samaria, p. 36, ponsonbyi, Lycia, p. 47, (Mastus) mestus, Sofrano Is., p. 46, Westerlund, Verh. z.-b. Wien, xlii: n. spp.

Clausilia (Nenia) orbignyi, Bolivia, Ancey, Brit. Nat. 1892, p. 126, Le Nat. 1892, p. 178, figured in Journ. Conch. vii, p. 97; C. (Albinaria) broemmei, Chelmos, near Kalavryta (Greece), BŒTTGER, Nachr. mal. Ges. 1892, p. 64; C. (Delima) optabilis, Bosnia, Brancsik, Evkön. Trencsen Varm. xii (1890) p. 73; C. (Papillifera) melitensis, Matta, Cahuana-Gatto, Med. Nat. i, p. 148; C. retorta, p. 153, friniana, p. 160, loloensis, p. 161, China, Heude (186), figg.; C. (Phædusa?) massiei, Laos, Morlet, J. de Conch. 1892, p. 83; C. paviei, p. 319, dautzenbergi, p. 320, Laos, Morlet, Journ. Conch. 1892, figg.; C. magistra, Bogota, Sowerby, P. Z. S. 1892, p. 298, fig.; C. (Alopia) deubeli, Siebenburgen, p. 38, (Delima) calabacensis, Greece, p. 39, (Cristataria) calopleura, Syria, (Cusmicia) vestigans, Carniolian Alps, p. 41, Westerlund, Verh. z.-b. Wien, xlii: n. spp.

Cocliazis australis is generically distinct from the type C. layardi, and is referrable to Perrieria, Tap.-Can. C. exigua, Adams, is a syn. of C. australis, and is not a Solomon Is. species; Hedley, P. Linn. Soc. N.S.W. vii, pp. 311-313.

Eucalodium compactum, n. sp., Tabasco, Mexico, Pilsbry, P. Ac. Philad. 1892, p. 338, fig.

Holospira: notes on the genus and its distribution; CROSSE & FISCHER, J. de Conch. 1892, pp. 256-279, 1 pl.

Odontostomus lemoinei, n. sp., Bolivia, ANCEY, Le Nat. 1892, p. 178, and Journ. Conch. vii, p. 93, fig.

Parknodus leroyi, p. 65, sesamorum, p. 66, n. spp., E. Centr. Africa, BOURGUIGNAT, (37) 1889, figg.

Perrieria: notes on the genus; Hedley, P. Linn. Soc. N.S.W. vii, pp. 311-313.

- Pupa monadicula, China, Heude, (186) p. 152, fig. P. ringens, in Guernsey; Marquand, J. Conch. vii, p. 44. P. coloradensis and ingersolli; for notes purporting to be descriptions of these nom. nud., see Cockerell, Brit. Nat. 1891, pp. 100 & 101.
- P. acanthinula, p. 709, lyonsiana, p. 713, magdalenæ, p. 716, Sandwich Is., Ancey, Mém. Soc. Zool. v; P. anodonta, Queensland, Hedley & Musson, P. Linn. Soc. N.S.W. vi, p. 558, fig.; P. elizabethensis, Port Elizabeth, ovampoensis, Ovampoland (Damaraland), Melvill & Ponsonby, Ann. N. H. ix, p. 91, fig.; P. (Leucochila) munita, clausa, Galapagos Is., Reibisch, SB. Ges. Isis, 1892, p. 27, fig.; P. turtoni, obliquicostata, St. Helena, Smith, P. Z. S. 1892, p. 268, figg.; P. (Torquilla) retracta, Spain, Westerlund, An. Soc. Esp. xxi, p. 388; P. (T.) refuga, homala, Sicily, p. 192, retracta, Spain p. 194, (Pupilla) honesta, Italy, p. 194, Westerlund, Nachr. mal. Ges. 1892; P. (Torquilla) domicella, Lourdes, hetaera, Pyrénées Orientales, p. 36, occidentalis, Hautes-Pyrénées, migma, Pyrénées Orientales, appeliusi, Dalmatia, p. 37, Westerlund, Verh. z.-b. Wien, xlii: n. spp.
- Ruchisellus burtoi, p. 69, ledoulzi, p. 70, n. spp., E. Centr. Africa, Bourguignat, (37) 1889, figg.
- Tomigerus (?) perexilis, n. sp., St. Helena, SMITH, P. Z. S. 1892, p. 267, fig.

#### STENOGYRIDA.

- Achatina bloyeti, p. 82, arctespirata, p. 83, randabeli, p. 84, E. Centr. Africa, Bourguignat, (37) 1889, figg.; A. (Burtoa) bridouxiana, p. 92, sebasmia, p. 93, pethericki, p. 95, lavigeriana, p. 96, E. Centr. Africa, Bourguignat, (37) 1889, figg.; A. (Burtopsis) jouberti, E. Centr. Africa, Bourguignat, (37) 1889, p. 99, fig.; A. (Limicolariu) charbonnieri, p. 104, megalæa, p. 105, coulboisi, p. 106, dromauxi, p. 107, sepulcralis, p. 108, lamellosa, p. 109, E. Centr. Africa, Bourguignat, (37) 1889, figg.; A. (Serpæa) pintoi, E. Centr. Africa, Bourguignat, (37) 1889, p. 86, fig.: [alleged n. spp.]. A. (Perideris) lechatelieri, Dahomey, Dautzenberg, J. de Conch. 1892, p. 297; A. stuhlmanni, Ituri, N. E. Africa, Martens, SB. nat. Fr. 1892, p. 176; (Homorus) pyramidella, Cameroons, Martens, SB. nat. Fr. 1892, p. 181; A. dimidiata [preocc., Smith, 1878], Gaboon, Martens, Conch. Mittheil. iii, 1889, p. 17, fig.: n. spp. [See also infrå, Stenogyra.]
- [Cazcilianella] Achatina acicula, in a Roman cemetery; Somerville, Journ. Conch. vii, pp. 42 & 43.
- Cacilianella baldwini, Sandwich Is., ANCEY, Mém. Soc. Zool. v, p. 718;
  C. gattoi, melitensis, polloneræ, Malta, Westerlund in Caruana-Gatto, Med. Nat. ii, p. 227: [alleged n. spp.].
- Cionella ovampoensis, Ovampoland (Damaraland), MELVILL & PONSONBY, Ann. N. H. ix, p. 91, fig.; C. (Ferussacia) virginea, p. 195, Spain, (Cacilianella) gattoi, p. 195, melitensis, pollonere, p. 196, Malta, 1892. [VOL. XXIX.]

WESTERLUND, Nachr. mal. Ges. 1892; C. (Ferussacia) extrema, Marocco, stenophya, bourlieri, Algeria, (Hohenwarthia) disparata, Barcelona, p. 43, WESTERLUND, Verh. z. b. Wien, xlii: C. (Ferussacia) virginea, Spain, WESTERLUND, An. Soc. Esp. xxi, p. 388: n. spp.

Cryptazeca monodonta, anatomy; BARROIS, Rev. Sci. Nat. Ouest, ii, pp. 330 & 331 : figg.

Glessula paviei, n. sp., Indo-China, Morlet, J. de Conch. 1892, p. 321, fig.

Leptinaria chathamensis, n. sp., Galapagos Is., DALL, Naut. v, p. 98.

Pseudoglessula, n. sect. of Stenogyra, for P. heteracra, subcrenata, W. Africa, n. spp.; Biettger, Nachr. mal. Ges. 1892, p. 202.

Subulma cylindracea, jouberti, E. Centr. Africa, BOURGUIGNAT, (37) 1889, p. 115, figg.; S. paucispira, Karewia, N. E. Africa, MARTENS, SB. nat. Fr. 1892, p. 177: n. spp.

Stenogyra. [See Pseudoglessula, n. sect.]

Stenogyra [or rather Achatina] leroyi, p. 110, grandidieriana, p. 111, E. Cent. Africa, Bourguignat, (37) 1889, figg.; S. palus, China, Heude (186) p. 151, fig.; S. badia, Congo, Martens, Conch. Mittheil. iii, 1889, p. 8, fig.; S. cacuminata, p. 85, South Africa, chapmani, p. 90, Ovampoland (Damaraland), Melvill & Ponsonby, Ann. N. H. ix, fig.: n. spp.

Zua sinensis, n. sp., China; HEUDE (186), p. 151, fig.

### HELICTERIDE and Succineide.

Amastra frosti, n. sp., Sandwich Is.; Ancey, Mém. Soc. Zool. v, p. 719. Curelia sinclairi, n. sp., Sandwich Is.; Ancey, Mém. Soc. Zool. v, p. 720. Succinea: living in dry localities; Loens, Nachr. mal. Ges. 1892, pp. 130 & 131. S. simplex, Pfeiff., jaw and radula; Hedley, P. Linn. Soc. N.S.W. vi, p. 691, figg.

S. (Brachyspira) decussata, Timor Laut Is., Mokllendorff, Nachr. mal. Ges. 1892, p. 99; S. wolf, Galapagos Is., Reibisch, SB. Ges. Isis, 1892, p. 28, fig.: n. spp.

#### VAGINULIDE and ONCIDIDE.

Notes on the Vaginulidæ; SIMROTH, Verh. Deutsch. Zool. Ges. i (1891), pp. 58-60.

Atopos, Simr.; notes on; JHERING, Nachr. mal. Ges. 1892, pp. 140-144: [with reply by Simroth, t. c. pp. 144-149]. Atopos, description of; Simroth, SB. Ges. Leipzig, xviii, p. 25. [Abs. of previous paper in Zool. Rec. 1891, Moll. p. 26, No. 400.]

Prisma, n. sect. for Vaginula tourannensis, Soul., prismatica, Tapp.-Canef., and heynemanni, nom. nud.; Simkoth, Z. wiss. Zool. lii, p. 596, fig.

Vaginula carusi, Singapore, chavesin, Miudanao, sarasinorum, Candy, bocagei, Moluccas, &c., strubelli, newtoni, platei, marshalli, Java.

sumatrensis, Sumatra, graffi, Celebes, vivipara, Saleijer Is., weberi, Sumatra, cockerelli, Java, chilensis, Chili, jordani, paraguensis, taylori, Paraguay, Simroth, SB. Ges. Leipzig, xviii, names given, pp. 85 & 86, descriptions, pp. 65–73, n. spp.

Veronicella jamaicensis, n. n. for V. kraussii, Fér., of Semper, p. 55, V. virgata, p. 96, dissimilis, p. 134, Cockerell, J. Inst. Jamaica, i:

alleged n. spp.

Oncidiidæ: on the structure and relationship of the; PLATE, Verh. Deutsch. Zool. Ges. ii, pp. 30-42.

Onchidium lesliei, n. sp., Galapagos Is., STEARNS, Naut. vi, p. 87.

### BASOMMATOPHORA.

### AURICULIDÆ.

- Alexia myosotis, Drap., in Is. of Borkum [first record for Germany]; SCHNBIDER, Nachr. mal. Ges. 1892, p. 116.
- A. armoricanu, n. sp. [?], W. Coast of France; LOCARD, Échange, vii, 1891, p. 132.
- Anelasma, n. sect. of Traliopsis, type † T. lemoinei, n. sp., Cossmann, Cat. Coqu. Foss. Eccene Paris, iv, 1889, p. 343: [altered to Micrelasma, Harris & Burrow, 1891].
- Semiauricula, n. sect. of Auricula, type † A. adversa, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iv, 1889, p. 341.
- Tralia (Alexia?) minuscula, Florida; DALL in SIMPSON, P. Davenport Ac. v, 1889, p. 69.

### LIMNAIDE and PHYSIDE.

- Ancylus in New Zealand; SUTER, J. de Conch. 1892, pp. 248-250.
- A. obliquus, n. sp., Lincoln, Iowa, Shimek, Bull. Lab. Iowa, i, p. 214, fig.
- Latia lateralis, Gould, phosphorescence in; SUTER, Mal. Bl. xi, p. 105.
- Linnaa limosina, conglobata, putriformis, lata, oppressa, pp. 18 & 19, strangulata, p. 78, alleged new French species; Locard, Échange, viii.
- L. nyansæ, Victoria Nyanza, MARTENS, SB. nat. Fr. 1892, p. 16, n. sp.
- Physa heterostropha, Say, notes on; CRANDALL, Naut. vi, p. 20: fertility of; POTEAT; J. Elisha Mitchell Sci. Soc. viii, pp. 70-73.
- P. trigona, Victoria Nyanza, Martens, SB. nat. Fr. 1892, p. 17; P. (Bulinus) clathrata, Spain, Westerlund, Nachr. mal. Ges. 1892, p. 197; P. (Bulinus) clathrata, Spain, Westerlund, An. Soc. Esp. xxi, p. 388: n. spp.
- Physopsis bloyeti, n. sp., E. Cent. Africa; BOURGUIGNAT, (37) 1889, p. 160, fig.
- Planorbis riparius, West., added to Brit. Fauna, from Lough Swilly; Milne, Irish Nat. 1892, p. 192. [Subsequently withdrawn, = P. nautileus.]

P. (Segmentina) emicans, S. Africa, Melvill & Ponsonby, Ann. N. H. x, p. 241, fig.; P. victoriæ, Victoria Nyadza, Smith, Ann. N. H. 1892, p. 383; † P. borellii, verticellus, p. 126, sabljari, p. 127, ptycophorus, lazici, p. 128, marinkovici, p. 129, Tertiary of Markuševec (Croatia), Brusina, Glasnik Naravosl. Druzt. vii: n. spp.

### SIPHONARIIDE and AMPHIBOLIDE.

Amphibola: anatomy; BOUVIER, Bull. Soc. Philom. iv, pp. 146-153.
Siphonaria is an Opisthobranch; Haller, Arb. z. Inst. Wien, x, pp. 71-100, figg., and should be placed next Umbrella.

 gigas, Less., anatomy, with figures; Haller, Arb. z. Inst. Wien, x, pp. 71-100.

## OPISTHOBRANCHIATA.

Vision in Opisthobranchiata; WILLEM, Arch. Biol. xii, pp. 123-149, figg. On the anal eye of the larvæ of Opisthobranchs; MAZZARELLI, Rend. R. Ac. Lincei, 1892, i, p. 103.

Opisthobranchs obtained on the voyages of the 'Hirondelle'; BERGH (18).

## NUDIBRANCHIATA.

Synopsis of families and classification; Bergh (17).—Organisation and phylogeny of the Nudibranchiata; Pelseneer, Bull. Soc. Mal. Belg. 1891 [1893], pp. lxviii-lxxi —Nudibranchiata holohepatica porostomata, synopsis; Bergh, Verh. z.-b. Wien, xlii (Abh.) pp. 1-16.—Nudibranchiata sacoglossa, are hermaphrodite; Pelseneer, Bull. Soc. Mal. Belg. 1891 [1893], p. lv.—Innervation of the cerata of some Nudibranchiata; Herdman & Clubb, Q. J. Micr. Sci. xxxiii, pp. 541-558, figg.—Notes on the Nudibranchiata found at Puffin I; Herdman, Tr. Liverp. Biol. Soc. v (1891) pp. 31-34.

#### ANTHOBRANCHIATA.

Emplocamus atlanticus, n. sp., Azores, BERGH, (18) p. 12, fig.

Jorunna johnstoni, notes on structure and habits; GARSTANG, Conchologist,
ii, pp. 49-52.

## INFEROBRANCHIATA.

Pleuroleura walteri, n. sp., Spitzbergen, Krause, Zool. JB. vi, Syst. p. 366, figg.

Pleurophyllidia loveni, Bergh, in Britain; M'Intosh, Conchologist, ii, p. 21: addition occurrence; Herdman, t. c. p. 52; Greisbach, t. c. p. 72.

### POLYBRANCHIATA.

- Cratena fructuosa, n. sp., Saragossa Sea, BERGH, (18) p. 4, fig.
- Eolididæ, remarks on some means of defence amongst the; HECHT, C.R. cxv, pp. 746-748.
- Glaucus atlanticus, Forster, note on specimens washed ashore on the coast of New South Wales; Hedley, P. Linn. Soc. N.S.W. vi, p. 576.
- Hancockia eudactylota, Gosse, observations on; Gamble, Ann. N. H. ix, pp. 378-385, figg.
- Lomanotus genei, Verany, observations on; Gamble, Ann. N. H. ix, pp. 378-385, figg.

### TECTIBRANCHIATA.

## (REPTANTIA.)

### ACTÆONIDÆ.

- Acteon delicatus, E. Coast U. S., Dall, Bull. Mus. C. Z. xviii (1889), p. 41, fig.; A. monterosatoi, Azores, Dautzenberg, (107) 1889, p. 20, figs.: n. spp.
- Bulimactzon, n. sect. for † Actzon bernayi, n. sp., Cossmann, Mem. Soc. Mal. Belg. 1891 (1892) p. 76.
- Crenilabium, n. sect. of Actaon, type † A. aciculatus, n. sp., Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 302.
- Leucotina elongata, n. sp., Port Elizabeth, Sowerby, (378) p. 52, fig.
- Ovulacteon, n. g., for O. meekii, n. sp., Caribbean; Dall, Bull. Mus. C. Z. xviii (1889), pp. 42 & 43, fig.
- Semiacteon, n. sect. of Acteon, type † Tornatella sphæricula, Desh.; Coss-Mann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 304.

### TORNATINIDE and SCAPHANDRIDE.

- Acrotrema, n. sect. of Cylichna, type †Bulla cylindroides, Desh.; Coss-MANN, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 313.
- Cylichna verrillii, Coast of N. Carolina, Dall, Bull. Mus. C. Z. xviii (1889), p. 54; C. richardi, chevreuxi, Azores, Dautzenberg, (107) 1889, p. 23, figg.: n. spp.
- Tornatina mariei, p. 21, protracta, p. 22, Azores, DAUTZENBERG (107) 1889, figg.; T. (Coleophysis) perplicatus, Caribbean, DALL, Bull. Mus. C. Z. xviii (1889), p. 45: n. spp.
- Utriculus (vortex var.?) domitus, n. sp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), p. 47, fig.
- Volvula bushii, aspinosa, n. spp., E. Coast U. S., Dall, Bull. Mus. C. Z. xviii (1889), p. 51.

## BULLIDA, RINGICULIDA, GASTROPTERIDA, and PHYLINIDA.

Bulla krebsii, Guadelupe, p. 56, clausa, Florida, p. 57, DALL, Bull. Mus.
C. Z. xviii (1889); B. guernei, Azores, DAUTZENBERG, (107) 1889,
p. 24, figg.: n. spp.

- Gasteropteron meckelii, Kosse, notes on; Smith, P. N. H. Soc. Glasg. iii, p. 258.
- †Gilbertia: notes on; Vincent, Bull. Soc. Mal. Belg. 1891 [1893] p. lii; Cossmann, t. c. p. lxiv.
- Philine infundibulum, p. 57, planata, p. 58, n. spp., E. Coast U. S., and Caribbean, Dall, Bull. Mus. C. Z. xviii (1889).

## PLEUROBRANCHIDÆ and UMBRELLIDÆ.

- Pleurobranchillus, n. g., p. 27, with P. morosus, n. sp., p. 28, Atlantic; Bergh, (18) fig.
- Umbraculum (Hyalopatina) rushii, n. sp., Bahamas, Dall, Bull. Mus. C. Z. xviii (1889), p. 61.
- Siphonaria is placed amongst Opisthobranchiata, next to Umbrella, by HALLER, Arb. z. Inst. Wien, x, pp. 71-100.

## (NATANTIA or PTEROPODA.)

- Muscular system of *Pteropoda*; Wackwitz, Zool. Beitr. iii, p. 129.—
  Notes on the embryology of the *Pteropoda*, Rev. Sci. Nat. St.
  Pétersb. 1891, p. 148 [Rusaian].—Catalogue of the *Pteropoda* of
  Australia and Tasmania; Brazier (44).
- † Conularia baini, n. sp., Palseozoic of Bolivia, ULRICH, (412) p. 36, fig. Hyolithidæ: Palseozoic Hyolithidæ of Bohemia revised; Novák, Abb. Böhm. Ges. iv, No. 6, with Ceratotheca, n. g., type Hyolithes aduncus, Barr., and 4 new species. Bactrotheca, n. g. for B. deleta, n. sp., and Hylithes teres, Barr. Pterygotheca, n. g. for P. barraudei, n. sp., and Orthotheca, n. g. & spp.
- † Hyolithes schencki, n. sp., Palseozoic, Bolivia, ULRICH, (412) p. 37, fig.

# [NUCLEOBRANCHIATA: see Prosobranchiata.]

## PROSOBRANCHIATA.

Vision in the *Prosobranchiata*; WILLEM, Arch. Biol. xii, pp. 123-149, figg.—Observations on the nephridia; ERLANGER, Q. J. Micr. Sci. xxxiii, pp. 587-623, figg.

## NUCLEOBRANCHIATA.

The Heteropoda are streptoneurous and are simply Prosobranchs whose exterior aspect has become modified by pelagic life; Pelseneer, C.B. cxiv, pp. 775-777.—Muscular system of Heteropoda; Wackwitz, Zool. Beitr. iii, p. 129.

### PECTINIBRANCHIATA.

## TOXOGLOSSA.

# TEREBRIDE and CONIDE.

- Hemiconus, n. sect. of Conus, type † C. stromboides, Lamk.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 230.
- Terebra (Acus?) benthalis, p. 65, Havana, nassula, limatula, p. 66, n. spp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889).

## PLEUROTOMIDÆ.

- Apiotoma, n. sect. of Pleurotoma, type †P. pirulata, Desh.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 259.
- Bela grimaldi, n. sp., Azores, Dautzenberg, (107) 1889, p. 26, figg.
- Borsonia (Cordieria) rouaultii, n. sp., Barbadoes, Dall., Bull. Mus. C. Z. xviii (1889), p. 98, fig.
- Clathurella watsoni, Azores, DAUTZENBERG, (107) 1889, p. 29, figg.; O. bourguignati, servaini [? new], W. Coast France, LOCARD, Échange, vii (1891), pp. 5 & 6: n. spp.
- Defrancia ponsonbyi, n. sp., Port Elizabeth, Sowerby, (378) p. 7, fig.
- Drillia tryonii, albicoma, Caribbean, p. 83, alesidota, hab.?, p. 85, eucosmia, Grenada, haliostrephis, Gulf of Mexico, p. 86, acestra, Havana, p. 87, acrybia, Florida, tristichia, Gulf of Mexico, p. 88, pagodula, Caribbean, p. 90, premorra, Gulf of Mexico, p. 93, apynota, p. 96, moseri, p. 97, E. Coast U. S., Dall, Bull. Mus. C. Z. xviii (1889), figg., n spp.
- Eopleurotoma, n. sect. of Pleurotoma, type †P. multicostata, Desh.; Coss-Mann, Cat. Coq. Foss. Eocene Paris, iv (1889), p. 265.
- Epalxis, n. sect. of Dolicotoma, type †Pleurotoma crenulata, L.; Coss-Mann, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 250.
- Genota viabrunnea, Caribbean, DALL, Bull. Mus. C. Z. xviii, p. 80, fig.
- Hemipleurotoma, n. sect. of Pleurotoma, type †P. archimedis, Bell.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 260.
- Mangilia (Cythara) bartlettii, cymella, Caribbean, p. 101, (Daphnella) corbicula, reticulosa, Barbadoes, &c., p. 103, pompholyx, Barbadoes, p. 104, retifera, p. 105, calyx, p. 107, E. Coast U. S., sofia, Guadelupe, p. 107, (Glyphostoma) phalera, Trinidad, p. 111, (Mangilia) halitropis, Santa Cruz, p. 114, scipio, Caribbean, p. 117, toreumata, Dominica, p. 118, (Pleurotomella) edgariana, Curaçao, p. 121, catasarca, Caribbean, p. 124, hadria, E. Coast U. S. p. 125, Dall, Bull. Mus. C. Z. xviii (1889), figg., n. spp.
- Mitromorpha biplicata, Barbadoes, Dall., Bull. Mus. C. Z. xviii (1889), p. 165, fig.; M. volva, Port Elizabeth, Sowerby, (378) p. 7, fig.: n. spp.
- Oxyacrum, n. sect. of Pleurotoma, type +P. obliterata, Desh.; Cossmann, Cat. Coqu. Foss. Eoceue Paris, iv (1889), p. 270.

- Pleurotoma periscelida, p. 74, (Leucosyrinx) tenoceras, p. 76, E. Coast U. S., (Ancistrosyrinx) radiata, p. 78, Caribbean, Dall, Bull. Mus. C. Z. xviii, figg.; P. (Mangilia?) simpsoni, Florida, Dall in Simpson, P. Davenport Ac. v (1887), p. 54; †P. haeringensis, Tertiary of Häring, Dreger, Ann. Hofmuseum Wien, vii, p. 28, fig.: n. spp.
- Systemope, n. sect. of Raphitoma, type †R. polycolpa, n. sp.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 289.

## CANCELLARIIDA.

- Admetula, n. sect. of Cancellaria, type †C. evulsa; Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 224.
- Benthobia, n. g., for B. tryonii, n. sp., N. Carolina; Dall, Bull. Mus. C. Z. xviii (1889), pp. 131 & 132, fig.
- Cancellaria (Trigonostoma) agassizii, E. Coast U. S., p. 130, microscopica, Caribbean, p. 131, Dall, Bull. Mus. C. Z. xviii (1889), figg., n. spp.
- Plesiocerithium, n. g., type † Cancellaria magloirei, Mell.; COSSMANN, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 228.
- Sceltella, n. sect. of Cancellaria, type † C. quantula, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 222.

#### RHACHIGLOSSA.

### OLIVIDE and MARGINELLIDE.

- Aucilla pura, n. sp., Port Elizabeth, Sowerby, (378) p. 17, fig.
- † ? Ancillaria olivaeformis, n. sp., Tertiary of Häring, DREGER, Ann. Hofmuseum Wien, vii, p. 26, fig.
- Marginella cassis, p. 137, fig., Gulf of Mexico, styria, Sombrero Is., p. 140, Dall, Bull. Mus. C. Z. xviii; M. electrina, Port Elizabeth, Sowerby, (378) p. 21, fig.: n. spp.

#### Volution and Mitrion.

- Mitra aquitanica, found on the W. Coast of France; LOCARD, Échange, viii, p. 101.
- M. (Costellaria?) styria, p. 159, fig., rushii, p. 160, triphonia, p. 161, Caribbean, (Thatia?) torticula, p. 162, Havana, (Conomitra) blakeana, Yucatan Strait, p. 163, Dall, Bull. Mus. C. Z. xviii (1889), figg.; M. fultoni, Lr. California, Smith, Ann. N. H. ix, p. 256, fig.: n. spp.
- Scaphella (Aurinia) robusta, p. 153, gouldiana [with its anatomy], p. 154, Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), figg.; S. (Voluta) arnheimi, Monterey Bay, Calif., Rivers, Naut. v, p. 111: n. spp.
- Voluta verconis, S. Australia, TATE, Tr. R. Soc. S. Austr. xv, p. 125, fig.; † V. stromboides, Tertiary of Häring, Dreger, Ann. Hofmuseum Wien, vii, p. 25, fig.: n. spp.

## FASCIOLARIIDE and TURBINELLIDE.

- Fusciolaria (Mesorhytis) meekiana, n. sp., Gulf of Mexico, Dall, Bull. Mus. C. Z. xviii (1889), p. 172, fig.
- Fusus timessus, p. 166, eucosmius, p. 167, halistreptus, benthalis, p. 168, amiantus, apynotus, p. 169, alcimus, p. 170, ceramidus, amphiurgus, p. 171, Dall, Bull. Mus. C. Z. xviii (1889), figg.; F. azoricus, Azores, Dautzenberg, (107) 1889, p. 32, figg.; †F. mittereri, Tertiary of Häring, Dreger, Ann. Hofmuseum Wien, vii, p. 23, fig.: n. spp.
- †Glyptostyla, n. g. for G. panamensis, n. sp., Tertiary of Isthmus of Darien; Dall, Tr. Wagner Inst. iii, pp. 232 & 233, figg.
- Latirulus, n. sect. of Latirus, type † Fusus subaffinis, d'Orb.; COSSMANN, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 169.
- Pseudolatirus, n. sect. of Streptochetus, type †S. mellevillei, n. sp.; Coss-MANN, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 170.
- Turbinella truncata, n. sp., Port Elizabeth, Sowerby, (378) p. 17, fig.

### BUCCINIDÆ.

- Chrysodomus (Sipho) rushii, Florida, C. (?) globulus, Little Bahama Bank, Dall, Bull. Mus. C. Z. xviii, p. 175, fig.; C. (Sipho) stonei, New Jersey, Pilsbry, P. Ac. Philad. 1892, p. 328, figg. : n. spp.
- Columbellisipho, n. sect. of Sipho, type †Fusus hordeolus, Lamk.; Coss-Mann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 146.
- Cominella semisulcata, C. (?) sulcata, n. spp., Port Elizabeth, Sowerby, (378) pp. 10 & 11, figg.
- Coptochetus, n. sect. of Siphonalia, type † Fusus scalaroides, Lamk.; Coss-MANN, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 153.
- Endopachychilus, n. sect. of Tritonidea, type †Purpura crassilabrum, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 136.
- Hindsia grimaldi, n. sp., Azores, DAUTZENBERG, (107) 1889, p. 33, figg.
- Liomesus (?) stimpsoni, n. sp., Carolina, Dall, Bull. Mus. C. Z. xviii (1889), p. 176, fig.
- Nassaria bushii, columbellata, p. 182, grayi, p. 183, n. spp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), figg.
- Neptunea nicolloni, Loc., W. Coast of France; Locard, Échange, vii (1891), p. 34.
- Pisania scholvieni, n. sp., Njaning, Senegambia, Rolle, Nachr. mal. Ges. 1892, p. 51.
- Pollia reticulatu, asona, n. spp., Kalymos Is., BŒTTGER, Ber. Seuck. Ges. 1892, p. 153.
- Tortisipho, n. sect. of Sipho, type + Fusus jucundus, Desh.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 147.
- Volutharpa salmiana, n. sp., Japan, Rolle, Nachr. mal. Ges. 1892, p. 52.

### NASSIDÆ and COLUMBELLIDÆ.

Nassa: Gould's types not lost; MARSHALL, Naut. vi, p. 47.

Nassa scissurata, Carribbean, Dall, Bull. Mus. C. Z. xviii (1889), p. 185;
N. crawfordi, Port Elizabeth, Sowerby, (378) p. 13, fig.: n. spp.

- Columbellidæ: monograph by Kobelt, pp. 1-128, pls. i-xviii; MARTINI & CHEMNITZ (246).
- Columbella: Gould's types not lost; MARSHALL, Naut. vi, p. 47.
- C. (Astyris) profundi, Havana, Dall, Bull. Mus. C. Z. xviii (1889), p. 192, fig.; C. (Amphissa) bicolor, San Pedro Bay, Dall in Williamson, P. U. S. Nat. Mus. xv, p. 213, fig.; C. (Mitrella) unisulcata, p. 119, Tasmania, Kobelt in Martini & Chemnitz (246); C. capensis, algoensis, South Africa, Sowerby, (378) p. 22, figg.; C. (Nitidella) incerta, Galapagos Is., Stearns, Naut. vi, p. 88; C. cominellaformis, S. Australia, Tate, Tr. R. Soc. S. Austr. xv, p. 126, fig.: n. spp.

## MURICIDE and CORALLIOPHILIDE.

- Coralliophila lactuca, n. sp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), p. 220, fig.
- Eupleura stimpsonii, n. sp., Barbadoes, Dall, Bull. Mus. C. Z. xviii (1889), p. 204.
- Magilus, Montf., revision of the genus; BAKER, P. Rochester Acad. ii, pp. 37-40, fig.
- Murex (Phyllonotus) hystricinus, Caribbean, p. 200, (Pteronotus) phaneus,
  Florida, p. 201, tristichus, Cuba, p. 202, (Muricidea) philippiana,
  Caribbean, p. 213, Dall, Bull. Mus. C. Z. xviii (1889), figg.; M.
  (Ocinebra?) crawfordi, babingtoni, South Africa, Sowerby, (378)
  p. 2, figg.; M. guembeli, Tertiary of Häring, Dreger, Ann. Hofmuseum Wien, vii, p. 24, fig.: n. spp.
- Trophon (Aspella) scalarioides, anatomy; DALL, Bull. Mus. C. Z. xviii (1889), p. 209.
- T. (Boreotrophon) lacunellus, p. 205, actinophorus, p. 206, Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), figs.; T. dabneyi, p. 36, droueti, p. 37, Azores, Dautzenberg, (107) 1889, figg.: n. spp.

### TÆNIOGLOSSA.

# TRITONIDE, CASSIDIDE, DOLLIDE, and CYPREIDE.

- Amphiperas (Simnia) piragua, p. 235, aureocincta, p. 236, n. spp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889).
- † Cassidaria haeringensis, n. sp., Tertiary of Häring, DREGER, Ann. Hofmuseum Wieu, vii, p. 21, fig.
- †Cassis fuchsi, n. sp., Tertiary of Häring, Dreger, Ann. Hofmuseum Wien, vii, p. 21, fig.
- Cypræa chrysalis, Kiener, and C. amphithales, Melv., notes on; Melville, Journ. Couch. vii, pp. 120-123.
- Dolium (Eudolium) verrillii, n. sp., Grenada, Dall, Bull. Mus. C. Z. xviii (1889), p. 233, fig.
- Lampusia? pharcida, n. sp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), p. 227, fig.

- Monocirsus, n. sect. of Triton, type +T. carinulatus, n. sp., Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 112.
- Ovula carnea, Poiret: distribution; FISCHER, J. de Conch. 1892, pp. 77 & 78; WATSON, t. c. p. 208.
- Pyrula: review of the genus; Sowerby, Conchologist, ii, pp. 73-75. P. filosa, n. sp., Hong Kong, Sowerby, Conchologist, ii, p. 74.
- † Tritonium haeringense, n. sp., Tertiary of Häring, DREGER, Ann. Hofmuseum Wien, vii, p. 22, fig.

## STROMBIDE and APORRHAIIDE.

- † Alaria acuta, n. sp., Neocomian of the Argentine Cordilleras, BEHREND-SEN, Z. geol. Ges. xliv, p. 19, fig.
- Amplogladius, n. sect. of Gladius, type † Rostellaria athleta, d'Orb.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 90.
- Ectinochilus, n. sect. of Rimella, type †Strombus canalis, Lam.; Coss-MANN, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 87.
- †Rostellaria haueri, n. sp., Tertiary of Häring, Dreger, Ann. Hofmuseum Wien, vii, p. 18, fig.
- Semiterebellum, n. sect. of Gladius, type † Rostellaria marceauxi, Desh.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 91.
- Wateletia, n. sect. of Gladius, type †Rostellaria geoffroyi, Wat.; Coss-MANN, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 90.

# CERITHIIDE, PLANAXIDE, and TRICHOTROPIDIDE.

- Cerithiidæ: inner palatal folds of fossil forms; OPPENHEIM, Z. geol. Ges. xliv, pp. 439-446.
- Aneurychilus, n. sect. of Sandbergeria, type †Cerithium secale, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 34.
- Cerithiopsis mataru, martensii, p. 255, acontium, p. 256, n. spp., Caribbean, Dall, Bull, Mus. C. Z. xviii (1889), fig.
- Cerithium foveolatum, n. sp., Port Elizabeth, Sowerby, (378) p. 36, fig.
- Exechestoma, n. sect. of Potamides, type †Cerithium angulosum, Lamk.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 71.
- Granulolabium, n. sect. of Potamides, type † Cerithium plicatum, Desh.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 76.
- Mellevillia, n. sect. of Fastigiella, type †Cerithium gibbosulum, Mell.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 36.
- Orthochilus, n. sect. of Planaxis, type †Cerithium bezançoni, De R.; Coss-MANN, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 7.
- †Pseudoscalites, n. g. for P. elegantissimus (Klipst MS.), n. sp.; KITTL, Ann. Hofmuseum Wien, vii, p. 66, fig.
- Semiverlagus, n. sect. of Cerithium, type C. vertagus, Lin.; COSSMANN, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 28.
- Tiarella, n. sect. of Lovenella, type † Cerithium pupina, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 50. [Altered to Cyrbusiu, Harris & Burrows, 1891.]

- Trachyschænium, n. sect. of Lovenella, type † Cerithium alveolatum, Desh.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iv (1889), p. 47.
- Triforis (Inella) sarissa, Barbadoes, p. 247, (Sychar) samanæ, San Domingo, p. 248, n. spp., Dall., Bull. Mus. C. Z. xviii (1889).
- Tylochilus, n. sect. of Potamides, type †Cerithium tuba, Desh.; Coss-MANN, Cat. Coqu. Foss. Eocene Paris, iv (1889), p. 76.

### VERMETIDE and TURRITELLIDE.

- Acrocoslum, n. sect. of Mathildia, type †M. bouryi, n. sp., Cossmann, Cat. Coqu. Foss. Eccene Paris, iii (1888), p. 315.
- Muthilda barbadense, p. 266, fig., Barbadoes, rushii, scitula, p. 267, n. spp., E. Coast U. S., Dall, Bull. Mus. C. Z. xviii (1889).
- † Siliquaria (?) triadica, n. sp., Trias of St. Cassian, Kittl, Ann. Hof-museum Wien, vii, p. 56, fig.
- Thylacodes medusæ, n. sp., Japan, MacGeorge, Le Nat. 1892, p. 123, fig. Turritella (Torcula) acropora, Transatlantic and Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), p. 264; T. maltzani, Gorée, Senegambia, Rolle,

Nachr. mal. Ges. 1892, p. 51; † T. paedopsis, subtilestriata, fasciata, p. 55, abbatis, p. 56, Trias of St. Cassian, KITTL, Ann. Hofmuseum Wien, vii, figg.: n. spp.

Vermetidæ of the Mediterranean, with Vermetus spirintortus, p. 22, simulans, p. 25, serpuloides, p. 41, n. spp.; Monterosato, Bull. Soc. Mal. Ital. xvii.

## PSEUDOMELANIIDÆ and MELANIIDÆ.

- †Angularia, n. g., with A. marginata, n. sp., Alpine Trias; Koken, JB. Mineral. 1892, ii, p. 32; Woehrmann & Koken, Z. geol. Ges. xliv, p. 198, fig.
- †Chemnitzia manzavinii, Trias of Balia, Asia Minor, BITTNER, JB. geol. Reichsanst. xlii, p. 87, fig.; †C. longiscata, solida, Alpine Trias, WOEHRMANN & KOKEN, Z. geol. Ges. xliv, p. 199, fig.: n. spp.
- † Hypsipleura, n. g, with H. cathedralis, n. sp., Alpine Trias; Koken, JB. Mineral. 1892, ii, p. 32; Woehrmann & Koken, Z. geol. Ges. xliv, p. 201, fig.
- † Katosira, n. g., with K. fragilis, abbreviata, n. spp., Alpine Trias; Koken, JB. Mineral. 1892, ii, p. 31; Woehrmann & Koken, Z. geol. Ges. zliv, p. 205, figg.
- †Loxonema pyrgula, lineatum, n. spp., Alpine Trias, Woehrmann & Koken, Z. geol. Ges. xliv, p. 202, figg.
- Melania verrii, De Stef., varieties; Foresti, Bull. Soc. Mal. Ital. xvi, p. 74, figg.
- M. jacquetiana, joretiana, terminalis, dolium, M. (?) rotundata, p. 166, China, Heude (186), figg.; M. tornata, Albert Nyanes, Martens, SB. nat. Fr. 1892, p. 181; M. ten-katei, Soemba, Schepman, Notes Leyd. Mus. xiv, p. 156, fig.: n. spp.

- Melaniida: inner palatal folds of fossil forms; OPPENHEIM, Z. geol. Ges. zliv, pp. 439-446.
- Melanopsis harpa, Spain, Westerlund, Nachr. mal. Ges. 1892, p. 199;
  M. harpa [= costata var.], Spain, Westerlund, An. Soc. Esp. xxi,
  p. 389; †M. pentagona, p. 138, stricturata, p. 139, bogdanowi, p. 142,
  blanchardi, p. 143, Tertiary of Markuševec (Croatia), Brusina,
  Glasnik. Naravosl. Druzt. vii: n. spp.
- † Melanoptychia paradoza, p. 144, rarinodosa, p. 145, n. spp., Tertiary of Markuševec (Croatia), Brusina, Glasnik. Naravosl. Druzt. vii.
- Potamanax, n. subg. of Pachychilus, for P. rovirosai, n. sp., Tabasco, Mexico; Pilsbry, P. Ac. Philad. 1892, p. 341, fig.
- Pachychilus: anatomy; Fischer & Crosse, (133) pp. 322-327.
- P. (polygonatus, Lea, ? var.) rovirosai, Mexico, Pilsbry, P. Ac. Philad. 1892, p. 153, fig. : possible n. sp.
- † Pustularia, n. g. for Chemnitzia alpina, Eichw., Alpine Trias; Koken, JB. Mineral. 1892, ii, p. 32; Woehrmann & Koken, Z. geol. Ges. xliv, p. 203, fig.
- Plotia leroyi, p. 185, bloyeti, p. 186, n.spp., E. Centr. Africa, BOURGUIGNAT, (37) 1889.
- Semisinus: anatomy; FISCHER & CROSSE, (133) pp. 315-317.
- † Undularia, n. g. for Turritella carinata, Münst, Alpine Trias; Koken, JB. Mineral. 1892, ii, p. 31; Woehrmann & Koken, Z. geol. Ges. xliv, p. 200, fig.
- † Zygopleura, n. g., with Z. spinosa, n. sp., and Coronaria, n. sect., with Z. coronata, n. sp., Alpine Trias; Koken, JB. Mineral. 1892, ii, p. 30; WOEHRMANN & KOKEN, Z. geol. Ges. xliv, pp. 203-5, figg.

### LITTORINIDE and FOSSARIDE.

- Cirsope, n. sect. of Lacuna, type †L. marginata, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 275.
- Cithna jeffreysi, n. sp., Azores; DAUTZENBERG, (107) 1889, p. 44, figg.
- Entomope, n. sect. of Lacuna, type †Litiopa klipsteini, Cossm.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 273.
- Forsarus (Gottoina) bella, compacta, n. spp., Caribbean; Dall, Bull. Mus. C. Z. xviii (1889), pp. 272 & 273, figg.
- †Lacuna karreri, L.(?) incrassata, n. spp., Trias of St. Cassian, KITTL, Ann. Hofmuseum Wien, vii, p. 97, figg.
- Littorina rudis, Maton, monstrosities; Sykes, P. Dorset Field Club, xiii, pp. 191-198, figg.
- L. (Tectarius) galapagiensis, p. 87, Galapagos Is., atyphus, p. 88, Manta, Ecuador, n. spp., Stearns, Nant. vi.
- Medoriopsis, n. sect. of Lacuna, type †L. effusa, Desh.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iii (1888), p. 270.
- Prosthenodon, n. sect. of Littorina, type †L. mondonta, Desh.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iii (1888), p. 261.

- †Pseudofossarus, n. n., = Fossarus, Laube, non Philippi; Koken, JB. Mineral. 1892, ii, p. 27; Woehrmann & Koken, Z. geol. Ges. xliv, p. 191, fig.
- † Purpurina vaceki, n. sp., Trias of St. Cassian; KITTL, Ann. Hofmuseum Wien, vii, p. 64, fig.
- † Tretospira, n. g., allied to Purpurina, with T. multistriata, n. sp., Alpine Trias; Koken, JB. Mineral. 1892, ii, p. 32; Woehrmann & Koken, Z. geol. Ges. xliv, pp. 196 & 197, fig.

#### SOLARIIDÆ.

- Fluxina discula, n. sp., Dominica; DALL, Bull. Mus. C. Z. xviii (1889), p. 273, figg.
- Homalogyra ornata, n. sp., Azores, Dautzenberg, (107) 1889, p. 46, figg. Solarium peracutum, sigsbeei, n. spp., Caribbean; Dall, Bull. Mus. C. Z. xviii (1889), p. 275, figg.

## JEFFREYSIIDE, LITIOPIDE, and RISSOIIDE.

- Alvania rufopunctata, Kalymos Is., BŒTTGER, Ber. Senck. Ges. 1892, p. 158; A. poucheti, p. 49, lamellata, p. 50, Azores, DAUTZENBERG, (107) 1889, figg.: n. spp.
- Benthonella, n. g. for B. gaza, fischeri, nisonis, Caribbean; Dall, Bull. Mus. C. Z. xviii (1889), pp. 282-285.
- † Chevallieria, n. g., type † C. labrosa, n. sp., Eocene of Paris, Coss-MANN, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 244, fig.
- Diala infrasulcata, dubia, n. spp., Port Elizabeth, Sowerby, (378) p. 27, figg.
- Hemibia sublevis, p. 168, carectorum, luteola, crassa, longiscuta, costulata, p. 169, n. spp., China, Heude, (186) figg.
- Jeffreysia atlantica, n. sp., St. Helena, Smith, Ann. N. H. 1892, p. 130.
- Onoba moreleti, n. sp., Azores, DAUTZENBERG, (107) 1889, p. 52, figg.
- Rissoa: notes on; Tomlin, Brit. Nat. 1892, pp. 72-74, 155 & 156, 179 & 180, 257 & 258.
- R. precipitata, Yucatan Strait, Dall. Bull. Mus. C. Z. xviii (1889), p. 279, fig.; R. guernei, p. 47, obesula, p. 48, Azores, Dautzenberg, (107) 1889, figg.; R. (Alvania) argentea, (Cingula) capensis, Port Elizabeth, Sowerby, (378) p. 38, figg.; † R. tirolensis, Alpine Trias, Woehrmann & Koken, Z. geol. Ges. xliv, p. 207, fig.: n. spp.
- Setia roseotincta, n. sp., Azores, DAUTZENBERG, (107) 1889, p. 53, figg.

## HYDROBIIDE and PALUDINIDE.

- † Acrophylictis, n. g., type † Bithinia eugenei, Desh., Eccene of Paris; Cossmann, Cat. Coqu. Foss. Eccene Paris, iii (1888), p. 233, fig.
- † Baglivia, n. g. for B. rugosula, p. 146, goniogyra, strongylogyra, p. 147, streptogyra, ambigua, p. 148, n. spp., Tertiary of Markuševec (Croatia), Brusina, Glasnik Naravosl. Drust. vii.
- Belgrandia coutagnei, n. sp., Avignon, Locard, Echange, viii, p. 3.

- Bythynella dunkeri, Frauenf., remarks on; Simroth, Mal. Bl. xi, pp. 106 & 107, figg.
- B. bosniensis, Bosnia, CLESSIN, Mal. Bl. xi, p. 120, fig.; B. depereti, deformata, lartetiæformis, ventricosa, anglesiana, (Lartetia) pontmartiniana, boulayi, chobauti, chantrei, gabilloti, duhameli, collieri, cureti, (Moitessieria) collieri, Aviguon, NICOLAS, Ann. Soc. Agric. Lyon, iv, pp. 34-48, figg.; B. scitula, Tertiary of Markuševec (Croatia), BRUSINA, Glasnik Naravoel. Drust. vii, p. 154: n. spp.
- Bythinia tentaculuta, development; ERLANGER, MT. z. Stat. Neap. x, pp. 376-407, fig.
- B. delavayana, p. 170, toucheana, chinensis, scalaris, p. 171, spiralis, p. 172, p. spp.. China, Heude, (186) figg.
- † Caspia dybowskii, p. 155, obtusa, acicula, p. 156, incerta, p. 157, n. spp., Tertiary of Markuševec (Croatia), Brusina, Glasnik Naravosl. Druzt. vii.
- Hydrobia texana, Texas, PILSBRY, P. Davenport Ac. v (1887), p. 33; † H. vidovici, taediosa, p. 149, atropida, p. 151, monotropida, ditropida, p. 152, polytropida, p. 153, Tertiary of Markuševec (Croatia), BRUSINA, Glasnik Naravosl. Druzt. vii: n. spp.
- Hypsobia, n. g. for H. humida, China; HEUDE, (186) p. 173, figg.
- Lacunopsis dugasti, n. sp., Laos, Morlet, J. de Conch. 1892, p. 84.
- † Micromelania sulcata, lineolata, p. 161, gracilis, p. 162, u. spp., Tertiary of Markuševec (Croatia), Brusina, Glasnik Naravosl. Druzt. vii.
- Puludestrina jenkinsi, Smith, and ventrosa, Mont., difference between the radulæ; WOODWARD, Ann. N. H. ix, pp. 376-378, figg.
- Paludina: development of the liver; H. FISCHER, CR. Soc. Biol. iii, p. 644.
  Paludina diminuta, leucostoma, fluminalis, catayensis, p. 175, ventricosa, aubryana, dispiralis, p. 175, purificata, fantozatina, p. 175, lapillorum, demolita, deformis, p. 177, China, Heude, (186), figg.; P. constricta, Victoria Nyanza, Martens, Conch. Mittheil. iii (1889), p. 16, fig.; P. laosiensis, Indo China, Morlet, J. de Conch. 1892, p. 85: n. spp. [See also infra, Viviparu.]
- Polycirsus, n. sect. of Hydrobia, type †Bythinia tuba, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 219.
- † Prososthenia croutica, n. sp., Tertiary of Markuševec (Croatia), BRUSINA, Glasnik Naravosl. Druzt. vii, p. 163.
- Pseudamnicola euboica, Euböa, p. 45, nebroden-is [= vestita, Benoit, 1882], Sicily, p. 50, n. spp., Kobelt in Rossmaessler, (322) v, figg.
- Pyrgulopsis scalariformis, Wolf, note on; Shimek, Bull. Lab. Iowa, ii, p. 168.
- Rivularia, n. g., for R. glandina, subelliptica, p. 178, elongata, ovum, globosa, p. 179, China; Heude (186), figg.
- Stenothyra toucheana, China, HEUDE, (186) p 173, figg., n. sp.
- Vivipara phthinotropis, p. 17, trochlearis, and costulata, p. 18, Victoria Nyanza, Martens, SB. nat. Fr. 1892; V. victoria, jucundus, cepoides, Victoria Nyanza, SMITH, Ann. N. H. x, pp. 124 & 125, figg.: n. spp. [See also supra, Paludina.]

## VALVATIDE, AMPULLARIIDE, and Assimineide.

- Ampullaria brohardi, Cambodia, Granger, Le Nat. 1892, p. 97, fig.;
  A. nyanza, gordoni, Victoria Nyanza, Smith, Ann. N. H. x,
  p 382: n. spp.
- Assiminea littorina, Delle Chiaie, in Malta; CARUANA-GATTO, Med. Nat. ii, p. 245.
- Chlorostracia massiei, n. sp., Laos, Morlet, J. de Conch. 1892, p. 84.
- Lyogyrus, notes on, and L. dalli, n. sp., Florida; PILSBRY, Naut. vi, pp. 61 & 62.
- Meladomus [= Lanistes] bloyeti, nitidissimus, p. 171, deguerryanus, p. 175, alexandri, p. 177, n. spp., E. Centr. Africa, BOURGUIGNAT, (37) 1889, figg.
- † Orygoceras corniculum, p. 169, filocinctum, cultratum, cnemopsis, p. 171, n. spp., Tertiary of Markuševec (Croatia), Brusina, Glasnik Naravoal. Druzt. vii.
- Valvata revoili, E. Centr. Africa, Bourguignat, (37) 1889, p. 189, fig.; †V. cyclostrema, leptonema, Tertiary of Markuševec (Croatia), Brusina, Glasnik Naravosl. Druzt. vii, p. 167: n. spp.

### CYCLOPHORIDÆ.

- Alycoeus expansus, dolichodeiros, p. 129, dolomiticus, monadicus, p. 130, n. spp., China, Heude (186), figg.
- Aperostoma (Amphicyclotus) goldfussi, n. sp., Honduras, Betttgen, Nachr. mal. Ges. 1892, p. 203.
- Cy. lophorus zebrinus, Bens., on a variety, and on the periostracum of; SMITH, Conchologist, ii, p. 6.
- C. orbignyi, E. Bolivia, Ancey, Journ. Conch. vii, p. 94; C. elatior, papillaris, Butumbi, N. E. Africa, Martens, SB. nat. Fr. 1892, p. 180; C. preussi, Cameroons, Martens, SB. nat. Fr. 1892, p. 183; C. tornatus, Tonkin, Morlet, J. de Conch. xxxii, p. 322, fig.; †C. matheroni, Upper Rognacian of Les Baux, Nicolas, C.R. Ass. Fr. Sci. 1891, ii, p. 452, fig.: n. spp.
- Cyclotus erroneus, China, Heude, (186) p. 128, figg.; †C. nicolasis, Upper Rognacian of Les Baux, Nicolas, C.R. Ass. Fr. Sci. 1891, ii, p. 453, fig.; C. soembaensis, Soemba, Schepman, Notes Leyd. Mus. xiv, p. 158, fig.; C. filo-liratus, Bogota, Sowerby, P. Z. S. 1892, p. 298, fig.: n. spp.
- Diplommatina decorosa, p. 510, garoensis, elongata, p. 511, chennelli, butleri, p. 512, ambigua, commutata, dohertyi, p. 513, thomsoni, nengloensis, distincta, p. 514, khunhoensis, lapillus, compacta, p. 515, unicrenata, japvoensis, animula, p. 516, subrubella, subtilis, delicata, p. 517, munipurensis, venustula, domuncula, p. 518, succinea, concinna, gibberosa, p. 519, Godwin-Austen, P. Z. S. 1892; D. egregia, Queensland, Hedley & Musson, P. Linn. Soc. N.S.W. vi, p. 561, fig.; D. intermedia, abbreriata, minuscula, China, Heude, (186) p. 131, figg.: p. spp.

- Geothauma, n. g., type Opisthostoma grandispinosum, Godwin-Austen; Crosse, J. de Conch. 1892, pp. 282-284, fig.
- Hedleya, n. g. for H. macleayi, n. sp., allied to Pupina? Solomon Is.; Cox, P. Linn. Soc. N.S. W. vi, p. 568, figg.
- Hungerfordia: notes on; CROSSE, J. de Conch. 1892, pp. 280-282, fig.
- Leptopoma halmahericum, crenilabre, n. spp., Halmahera [Moluccas], STRUBELL, Nachr. mal. Ges. 1892, pp. 48 & 49.
- Maizania, n. g. for M. olivacea, n. sp., E. Centr. Africa; BOURGUIGNAT, (37) 1889, p. 149, fig.
- † Megalomastoma julliani, n. sp., Upper Rognacian of Les Baux; NICOLAS, C.B. Ass. Fr. Sci. 1891, ii, p. 454, fig.
- Opisthoporus hyperborealis, n. sp., China, Heude, (186) p. 128, figg.
- Paxillus borealis, n. sp., China, HEUDE, (186) p. 131, fig.
- Pomatias (Auritus) eircticus, n. sp., Sicily, WESTERLUND, Nachr. mal. Ges. 1892, p. 198.
- Procyclotus, n. g. for Asiatic species of Cyclotus, type C. gassiesiamus, Crosse; Fischer, Bull. Soc. Autun, iv, p. 183.
- Pupina costata, n. sp., Queensland, Hedley & Musson, P. Linn. Soc. N.S.W. vi, p. 559, fig.
- Spiraculum massiei, n. sp., Laos, Morlet, J. de Conch. 1892, p. 85.

## CYCLOSTOMATIDE, ACICULIDE, and TRUNCATELLIDE.

- Acme allardi, alleged n. sp., Avignon, NICOLAS, Ann. Soc. Agric. Lyon, iv, p. 49, fig.
- Acroptychia notabilis, Madagascar, SMITH, Conchologist, ii, p. 23.
- Cyclostoma burtoni, Tanganyka, ANCEY, Brit. Nat. 1892, p. 126, fig.; C. leroyi, E. Centr. Africa, BOURGUIGNAT, (37) 1889, p. 152, figg.; †C. jagici, Tertiary of Markuševec (Croatia), BRUSINA, Glasnik Naravoal. Druzt. vii, p. 172: n. spp.
- Tropidophora freyi, n. sp., Is. Nossi-Be [Madagascar], BETTGER, Nachr. mal. Ges. 1892, p. 55.
- †Truncatella bezançoni, n. sp., Oligocene of Etampes; Cossmann, J. de Conch. xxxii, p. 358, fig.

## HIPPONYCIDE and CAPULIDE.

- Amalthea [= Hipponyx] benthophila, n. sp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), p. 289, fig.
- Calyptraida: anatomy and morphology; HALLER, Morph. JB. xviii, pp. 489-518.
- Capulus hungaricus: structure and development; ERLANGER, Zool. Anz. xv. pp. 465-468.
- C. (Hyalorisia) galea, n. sp., Barbadoes, Dall, Bull. Mus. C. Z. xviii (1889), p. 288, fig.
- Crepidula fornicata: cleavage of the ovum; CONKLIN, Zool. Anz. xv, pp. 185-188, figg.
  - 1892. [VOL. XXIX.]

- Crepidulæ of the New Jersey Coast; STONE, Naut. vi, pp. 40-42, figg.
- † Delphinulopsis, Laube, is restored as a genus by KITTL, Ann. Hofmuseum Wien, vii, p. 58.
- †D. laubei, n. sp., Trias of St. Cassian, KITTL, Ann. Hofmuseum Wien, vii, p. 62, fig.
- Platyceras: notes on Palæozoic species of the group; KEYES, Am. Geol. x, pp. 273-277.
- †Rothpletzia rudistu, Sim.: on its zoological affinities; SIMONELLI, Boll. Soc. geol. Ital. xi, pp. 76-80, figg.

## Naricide, Naticide, Occorythide, and Adeorbiide.

- Adeorbis (Clathrella) naticoides, n. sp., N. Carolina, Dall, Bull. Mus. C. Z. xviii (1889), p. 278.
- Crommium, n. sect. of Ampullina, type †A. willemeti, Desh., Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 177.
- Dalium, n. g. for D. solidum, n. sp., Grenada, Dall, Bull. Mus. C. Z. xviii (1889), p. 230, fig.
- Dialytostoma, n. sect. for Escharella [altered 1891 to Micreschara], type + Fossarus fischeri, De Laub.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 185.
- † Natica (Naticina) geinitzi, Rupelian of Germany, BŒTTGER, Mal. Bl. ix, p. 89; N. castrensis, p. 293, (Neverita) nubila, p. 294, Caribbean, Dall, Bull. Mus. C. Z. xviii (1889); N. (Lunatia) fordiana, Florida, Simpson, P. Davenport Ac. v (1889), p. 72°; † N. bodenbenderi, Inf. Oolite of the Argentine Cordilleras, BEHRENDSEN, Z. geol. Ges. xliv, p. 13, fig.; † N. berwerthi, p. 86, transiens, p. 89, Trias of St. Cassian. Kittl, Ann. Hofmuseum Wien, vii, figg.: n. spp.
- † Naticella (?) anomala, n. sp., Trias of St. Cassian, Kittl, Ann. Hofmuseum Wien, vii, p. 71, fig.
- Naticidæ: anatomy and morphology; HALLER, Morph. JB. xviii, pp. 451-489, figg.
- †Neritaria, n. g., with N. similis, n. sp., Alpine Trias, Woehrmann & Koken, Z. geol. Ges. xliv, p. 192, figg.
- Macromphalina, n. sect. for Escharella [altered 1891 to Microschara], type +Sigaretus problematicus, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 184.
- Micromphalina, n. sect. of Escharella [sltered 1891 to Micreschara], type †Lacuna elegans, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 187.
- † Ptychostoma stachei, washneri, p. 95, mojsisovicsi, fasciatum, p. 96, n. spp., Trias of St. Cassian, Kittl, Ann. Hofmuseum Wien, vii, figg.
- Sigaretopsis, n. sect. of Natica, type †B. infundibulum, Wat.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 172.
- Sigaretus minor, (Eunaticina) carolinensis, n. spp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), p. 297.

#### PTENOGLOSSA.

#### SCALARIIDE.

- Aclis egregia, nucleata, Caribbean, Dall. Bull. Mus. C. Z. xviii (1889), p. 325, figg.; A. tenuistriata, Port Elizabeth, Sowerby, (378) p. 28, fig.: n. spp.
- Bifidoscala, n. sect. of Scalaria, type †S. lemoinei, De Boury; Coss-Mann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 135.
- † Chilocyclus, Bronn: restored as a genus by KITTL, Ann. Hofmuseum Wien, vii, p. 52.
- Scala scipio, apiculata, p. 310, babylonia, p. 311, retifera, p. 312, frielei, rushii, sericifila, p. 313, nitidella, muscapedia, p. 314, belaurita, p. 316, contorquata, p. 318, polacia, p. 319, aurifila, p. 322, concava, p. 323, discoboluria, p. 324, Transatlantic and Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), figg.; S. rodleri, subulata, Tertiary of Häring, Dreger, Ann. Hofmuseum Wien, vii, pp. 14 & 15, figg.; †S. (?) triadica, p. 45, S. damesi, p. 51, Trias of St. Cassian, Kittl, Ann. Hofmuseum Wien, vii, figg.: n. spp.

#### GYMNOGLOSSA.

## EULIMIDE and PYRAMIDELLIDE.

- Phylogeny of the species of Eulimidæ and Pyramidellidæ from the Tertiary strata of N. Italy; SACCO, Mem. Acc. Tor. xlii, pp. 585-682.
- Auriculina lucida, n. sp., Port Elizabeth, Sowerby, (378) p. 37, fig.
- Cingulina acutilirata, n. sp., Port Elizabeth, Sowerby, (378) p. 27, fig.
- Cyclodostomia, n. subg. of Odostomia, with n. spp., Tertiary of N. Italy; Sacco, Mem. Acc. Tor. xlii, p. 628.
- Diptychus, n. sect. of Syrnola, type †Pyramidella clandestina, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 95.
- Eulima (Leiostraca) fusus, Caribbean, DALL, Bull. Mus. C. Z. xviii (1889), p. 329, fig.; E. langleyi, Port Elizabeth, Sowerby, (378) p. 27, fig.: n. spp.
- Eulimella (Stylopsis) resticula, n. sp., Florida Keys, Dall, Bull. Mus. C. Z. xviii (1889), p. 338.
- Hordeulima, n. subg. of Eulima, with †H. hordeola, n. sp., Tertiary of N. Italy; Sacco, Mem. Acc. Tor. xlii, p. 599, fig.
- Loxoptyxis, n. sect. of Syrnola, type +S. conulus, n. sp.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iii (1888), p. 103.
- Macrodostomia, n. subg. Odostomia, with n. spp., Tertiary of N. Italy; SACCO, Mem. Acc. Tor. xlii, p. 625, fig.
- Margineulima, n. sect. of Eulima, type † E. fullax, Desh.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iii (1888), p. 121.
- Miralda crispa, n. sp., Port Elizabeth, Sowerby, (378) p. 37, fig.
- † Odontostomia lamberti, n. sp., Oligocene of Etampes, Cossmann, J. de Conch. 1892, p. 347, fig.

- Peristichia, n. g., for P. toreta, Florida Keys, agria, Cape Hatteras; DALL, Bull. Mus. C. Z. xviii (1889), pp. 339 & 340.
- Pharcidella, n. sect. of Pyramidella [q. v.].
- Ptycheulimella, n. subg. of Eulimella, with n. spp., Tertiary of N. Italy; SACCO, Mem. Acc. Tor. xlii, p. 641.
- Pyramidella (Pharcidella) folinii, n. sp., Barbadoes, Dall, Bull. Mus. C. Z. xviii (1889), p. 334.
- Pyrgolampros, n. subg. of Turbonilla, with n. spp., Tertiary of N. Italy; Sacco, Mem. Acc. Tor. xlii, p. 667, fig.
- †Spica, ? n. g., with S. monterosatoi, n. sp., Tertiary of N. Italy; SACCO, Mem. Acc. Tor. xlii, p. 645, fig.
- Strioturbonilla, n. subg. of Turbonilla, with n. spp., Tertiary of N. Italy; Sacco, Mem. Acc. Tor. xlji, p. 676, fig.
- Sulcosubularia, n. subg. of Eulima, with †S. taurinensis, n. sp., Tertiary of N. Italy; Sacco, Mem. Acc. Tor. xlii, p. 599, fig.
- Sulcoturbonilla, n. subg. Turbonilla, for †S. turricula, Eich.; Sacco, Mem. Acc. Tor. xlii, p. 674, fig.
- Turbonilla belotheca, p. 335, curta, p. 337, Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), figg.; T. guernei, p. 60, coarctata, p. 61, Azores, Dautzenberg, (107) 1889, figg.; T. argentea, læricostata, tegulata, Port Elizabeth, Sowerby, (378) p. 25, figg.; T. crenulifera, S. Australia, Tate, Tr. R. Soc. S. Austr. xv, p. 126, fig. : n. spp.
- Turritodostomia, n. subg. of Odostomia, type † O. plicata, Mont.; SACCO, Mem. Acc. Tor. xlii, p. 623, fig.

## SCUTIBRANCHIATA.

#### RHIPIDOGLOSSA.

## PROSERPINIDE and Helicinide.

Ceres: anatomy; Fischer & Crosse, (133) pp. 379-381.

Cyane orbignyi, n. sp., Bolivia, ANCEY, Le Nat. 1892, p. 178.

Helicina: anatomy; FISCHER & CROSSE, (133) pp. 390-392.

H. leucozonalis, Brazil, bourguignatiana, E. Bolivia, p. 95, lirifera, sulfurea, E. Bolivia, p. 96, Ancey, Journ. Conch. vii; H. (Idesa) nesiotica, Galapagos Is., Dall, Naut. v, p. 97; H. wolfi, Galapagos Is., Reibisch, SB. Ges. Isis, 1892, p. 29, fig.: n. spp.

Heudeia: notes on the genus and its allies; CRUSSE, J. de Conch. 1892, pp. 284-288, figg.

### NERITIDE and NERITOPSIDE.

- On the mode of growth and the structure of the shell in *Neritidæ*, especially in *Velates conoideus*, Lamk.; WOODWARD, P. Z. S. 1892, pp. 528-540.
- † Hologyra, n. g., with H. alpina, carinata, n. spp., Alpine Trias; WOEHRMANN & KOKEN, Z. geol. Ges. xliv, pp. 193 & 194, fig.

- † Nuticopsis (?) ladina, dianae. p. 75, N. zitteli, laubei, p. 76, altoni, p. 78, gaderana, p. 79, kinkelini, p. 82, uhligi (Klipst. MS.), involuta, p. 83, telleri, p. 84, n. spp., Trias of St. Cassian, Kittl, Ann. Hofmuseum Wien, vii, figg.
- Nerita polita: nervous system; Boutan, C.B. cxiv, p. 1133.
- Neritina picta, Sow., note on; CROSSE & FISCHER, J. de Conch. 1892, pp. 292 & 293.
- N. (Clithon) rhyssodes, Is. Nossi-Bé [Madagascar], Bœttger, Nachr. mal. Ges. 1892, p. 57; N. sargi, Guatemala, Crosse & Fischer, J. de Conch. 1892, p. 296; N. (Theodoxus) mixta, Spain, Westerlund, Nachr. mal. Ges. 1892, p. 200, and An. Soc. Esp. xxi, p. 389; †N. imitans, Trias of St. Cassian, Kittl, Ann. Hofmuseum Wien, vii, p. 41, fig.: n. spp.
- † Neritodonta cunici, p. 177, serrulata, lamellata, p. 178, n. sp., Tertiary of Markuševec (Croatia), Brusina, Glasnik Naravosl. Druzt. vii.
- † Palaeonarica, n. g.. p. 42, type Naticella concentrica, Münst., P. constricta, p. 42, cancellata, p. 44, n. spp., Trias of St. Cassian, KITTL, Ann. Hofmuseum Wien, vii, figg.
- † Platychilina, n. g., with P. woehrmanni, n. sp., Alpine Trias; WOEHR-MANN & KOKEN, Z. geol. Ges. xliv, pp. 195 & 196, fig.
- † Velates conoideus, Lamk., structure and growth of shell; WOODWARD, P. Z. S. 1892, pp. 528-540.

## TURBINIDE and TROCHIDE.

- Buthybembix, Watson emend., notes on; CROSS, J. de Conch, 1892, pp. 288-292, figg.
- Calliostoma corbis, p. 365, orion, p. 367, (Eucasta) indiana, p. 368, (Eutrochus) sayanum, p. 370, benedicti, p. 371, cinctellum, p. 372, (Dentistyla) sericifilum, p. 373, Transatlantic and Caribbean, Dall, Bull. Mus. C. Z. xviii (1889) figg.; C. farquhari, Port Elizabeth, Sowerby, (378) p. 43, fig.; †P. jetschini, Rupelian of Germany, Bættger, Mal. Bl. xi, p. 91: n. spp.
- Cirsochilus, n. sect. of Collonia, type † Delphinula striata, Desh.; Coss-MANN, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 76.
- Clanculus waltonæ, n. sp., Port Elizabeth, Sowerby, (378) p. 45, fig.
- Cyniscella, n. sect. of Collonia, type †Cyclostoma cornu-pastoris, Lamk.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iii (1888), p. 79.
- Dillwynella, n. subg. of Ethalia [q.v.].
- Ethalia reclusa, p. 361, suppressa, solida, (Dillwynella) modesta, p. 362, n. spp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), figg.
- Eucasta, n. sect. of Calliostoma [q.v.].
- Euchelus guttarosea, n. sp., Caribbean, DALL, Bull. Mus. C. Z. xviii (1889), p. 382, fig.
- Gaza fischeri, n. sp., Santa Lucia, DALL, Bull. Mus. C. Z. xviii (1889), p. 355, fig.

- Gibbula eliaduru, p. 159, valentini, p. 160, Kalymos Is., Bettger, Ber. Senck. Ges. 1892; G. biporcata, Cape Town, Sowerby, (378) p. 67: n. spp.
- Homalochilus, n. sect. for †Collonia miliaris, n. sp., Eocene Paris; Coss-Mann, Mem. Soc. Mal. Belg. 1891 (1892), p. 44.
- Margarita (Turcicula) imperialis, anatomy; Dall, Bull. Mus. C. Z. xviii (1889), pp. 376 & 377.
- M. srythrocoma, p. 375, Caribbean, (Turcicula) bairdii, p. 378, n. spp., S. California, Dall, Bull. Mus. C. Z. xviii (1889), figg.
- Monilea fritzei, n. sp., Middle Liu-Kiu Is., BŒTTGER, Nachr. mal. Ges. 1892, p. 166.
- Norrisella, n. sect. of Norrisia, type Turbo pygmaus, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 62.
- † Oncospira legayi, n. sp., Oxfordian of Boulogue, RIGAUX, Mém. Soc. Acad. Boulogne, xiv, p. 108, fig.
- Otaulax, n. sect. of Leptothyra, type † Turbo inermis, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 72.
- Periaulax, n. sect. of Eumargarita, type †Solarium spiratum, Lamk.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 66.
- Phasianella azorica, n. sp., Azores, DAUTZENBERG, (107) 1889, p. 61, figg. Phorculus, n. sect. of Gibbula, type †Turbo fraterculus, Desh.; Coss-
- MANN, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 61.

  Pseudodiloma, n. sect. of Gibbula, type † Trochus mirabilis, Desh.; Coss-
- MANN, Cat. Coqu. Foss. Eccene Paris, iii (1888), p. 60.
- Solariella fusco-maculata, n. sp., Port Elizabeth, Sowerby, (378) p. 44, fig.
- Tectariopsis, n. sect. of Turbo, type † T. henrici, Caill.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iii (1888), p. 71.
- † Trochus demersus, n. sp., Tertiary of Häring, Dreger, Ann. Hofmuseum Wien, vii, p. 14, fig.
- Turbo (Leptothyra) philipiana, linnei, Caribbean, Dall., Bull. Mus. C. Z. xviii (1889), p. 353, figg.; † T. tuberculosus, Tertiary of Häring, DREGER, Ann. Hofmuseum Wien, vii, p. 13, fig.: n. spp.
- Umbonium bairdii, n. sp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), p. 359, figg.
- Uvanilla regina, n. sp., West Coast U. S.; STEARNS, Naut. vi, p. 85.

# DELPHINULIDE, CYCLOSTREMATIDE, and Cocculinide.

- Cocculina spinigera, anatomy; DALL, Bull. Mus. C. Z. xviii (1889), p. 349. Cyclostrema turbinum, p. 393, Havana, pompholyx, Gulf of Mexico, cistronium, N. Carolina, p. 394, granulum, Samana Bay, (Granigyra) limatum, Cuba, p. 395, DALL, Bull. Mus. C. Z. xviii (1889), figg.; C. rotundata, inflata, planulata, Port Elizabeth, Sowerby, (378) pp. 45 & 46, figg.: n. spp.
- † Delphinula purkeri, n. sp., Oxfordian of Boulogne, RIGAUX, Mem. Soc. Acad. Boulogne, xiv, p. 107, fig.

- Lintia bairdii, p. 389, miniata, variabilis, p. 390, (Lippistes) acrilla, p. 391, amabilis, p. 392, n. spp., Carribbean, Dall, Bull. Mus. C. Z. xviii (1889), figg.
- Megatyloma, n. sect. of Tinostoma, type † T. wateleti, Desh.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii, p. 50.
- Vitrinella multicarinata, E. Coast U. S., DALL, Bull. Mus. C. Z. xviii (1889), p. 392; V. williamsoni, San Pedro Bay, DALL in WILLIAMSON, P. U. S. Nat. Mus. xv, p. 202, fig. : n. spp.

# HALIOTIDE, PLEUROTOMARIIDE, and BELLEROPHONTIDE.

- Haliotis: imperforate specimen; SMITH, Conchologist, ii, pp. 75 & 76.
- Murchisonia (Goniostropha) hibernica, tatei, p. 563, (Hypergonia) pentonensis, p. 569, kirkbyi, p. 570, plana, p. 571, (Calocaulus?) tuedia, p. 572, n. spp., Carboniferous of British Is., Donald, Q. J. Geol. Soc. xlviii, figg.
- Pleurotomaria quoyana and adansoniana, description of the animal and anatomy; Dall, Bull. Mus. C. Z. xviii (1889), pp. 397-403.
- †P. kayseri, n. sp., Palæozoic of Bolivia, Ulrich, (412) p. 40, fig.
- Schismope fayalensis, n. sp., Azores, DAUTZENBERG, (107) 1889, p. 65, figg.
- † Trematonotus britannicus, Wenlock Beds, near Dudley, proposed as a new species; Newton, Geol. Mag. 1892, p. 339: withdrawn as a synonym of T. dilatatus; Newton, t. c. p. 525.

## FISSURELLIDÆ.

- Fissurellidæ of the United States: catalogue of; Pilsbry & Johnson, Naut. v, pp. 102-107, & 113.
- Emarginula compressa, Cantraine, note on; Monterosato, J. de Conch. 1892, pp. 78-81.
- E. (Rimula) frenulata, W. Florida, Dall, Bull. Mus. C. Z. xviii (1889), p. 406, fig.; †E. kittli, Tertiary of Häring, Dreger, Ann. Hofmuseum Wieu, vii, p. 13, fig. : n. spp.
- Entomella, n. sect. of Emarginula, type †E. clypeata, Lam.; Cossmann, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 39.
- Fissurella (Glyphis) fluviana, n. sp., Caribbean, Dall, Bull. Mus. C. Z. xviii (1889), p. 408, fig.
- Puncturella watsoni, p. 403, Caribbean, P. (?) erecta, p. 405, n. spp., N. Carolina, Dall., Bull. Mus. C. Z. xviii (1889).
- Scutum: note on the eye of; Pelseneer, Bull. Soc. Mal. Belg. 1891 [1893]. p. xxviii. [S. =] Parmophorus australis, figured; Boutan, Rev. Biol. v, pl. ix (p. 69).

#### DOCOGLOSSA.

## ACMRIDE and PATELLIDE.

Acmaa: position of osphradia; Dall, Bull. Mus. C. Z. xviii (1889), p. 410. A. saccharina, Linn., note on; Taylor, Naut. vi, p. 89.

Patellidæ of Japan; TAYLOR, Nant. vi, pp. 69 & 70.

Patella vulgata, respiratory globulin from the blood; GRIFFITHS, C.B. cxv, p. 259.

### INCERTA SEDES.

† Astralites, n. g. for A. fimbriatus, n. sp., Devonian, Manitoba; Whiteaves, Contr. Canad. Pal. i, p. 323, fig. In some characters this genus resembles Uvanilla, Onustus, or Pleurotomaria, but differs in the presence of a conspicuous fold upon the columella.

## III. POLYPLACOPHORA.

- Monograph begun by PILSBRY (409), who subdivides group as far as done into:—1. LEPIDOPLEURIDE, (Lepidopleurus, Hanleya, Hemiarthrum, Microplax). 2. CHITONIDE, Ischnochitoninæ (Leptoplax, Spongiochiton, Chætopleura, Tonicella, Schizoplax, Callochiton, Ischnochiton), Chitoninæ, and Acanthopleurinæ.
- The integument of Chitons; THIELE, Biol. Centralbl. xi (1891), pp. 722-726. Respiratory globulin in the blood of Chitons; GRIFFITHS, C.R. cxv, pp. 474 & 475. Embryology of Chiton; METCALF, Johns Hopk. Univ. Circ. xi, p. 79.
- Callochiton princeps, p. 50, Hab. ?, lobatus, p. 53, Tasmania (Carpenter MS.), n. spp., PILSBRY in TRYON'S Manual, xiv, fig.
- Chætopleura gemmea, p. 31, Monterey, California, armillata, p. 39, Is. Gorriti (Carpenter MS.), n. spp.; Pilsbry in Tryon's Manual, xiv, fig.
- Cyanoplax, n. subg. of Toniciella; PILSBRY in TRYON'S Manual, xiv, p. 44.
- Ischnochiton: PILSBRY in TRYON'S Manual, xiv, divides into subgenera as follows:—Stenochiton, Stenoplax, Ischnoplax, Heterozona, Tachydermon, Ischnochiton, Ischnoradsia.
- floridanus, p. 58, West Key, Florida, fallax (Carpenter MS.), p. 59,
   Monterey, California, acrior (ib.), p. 61, Lower California, conspicuus (ib.), p. 63,
   California and Lower California, cariosus (ib.), p. 65,
   Australia, solidior (ib.), p. 76,
   Philippines, asperior (ib.), p. 77,
   Hong Kong or Hakodade, perornatus (ib.), p. 77,
   hab. ?, multidentatus (ib.), p. 85,
   Bonin Is., haddoni (ib.), p. 88,
   Port Jackson, N.S.W., bisculptus (Carpenter MS.), p. 119,
   Hong Kong ?, newcombi (ib.), p. 120,
   Santa Catalina Is., radians (ib.), p. 121,
   Monterey,
   California, aureotinctus (ib.), p. 123,
   Santa Catalina Is., decipiens (ib.), p. 123,
   Monterey,

- California, corrugatus (ib.), p. 123, California, cooperi (ib.), p. 127, California, sinudentatus (ib.), p. 128, Monterey, California, n. spp., PILSBRY in TRYON'S Manual, xiv, figg.
- Lepidopleurus rugatus, p. 11, Lower California, granoliratus, p. 14, Mogador, curvatus, p. 16, Japan (Carpenter MS.), n. spp., Pilsbry in Tryon's Manual, xiv, figg.
- Lepidozona, n. sect. of subg. Ischnockiton (s. s.); PILSBRY in TRYON'S Manual, xiv, pp. 86 & 125 [at first, p. 55, given as sect. of subg. Ischnoradsia].
- Leptochiton pergranatus, n. sp., Dominica, DALL, Bull. Mus. C. Z. xviii (1889), p. 414.
- Notoplaz floridanus, n. sp., Florida Keys, DALL, Bull. Mus. C. Z. xviii (1889), p. 416.
- Radsiella, n. sect. of subg. Ischnochiton (a. s.); PILSBRY in TRYON'S Manual, xiv, pp. 54 & 86.
- Spongiochiton productus (Carpenter MS.), n. sp., New Zealand, PILSBRY in TRYON'S Manual, xiv, p. 26, fig.

## IV. APLACOPHORA.

- Chætoderma nitidulum, Lovén: histological notes; Wiren, Biol. Fören. Förhandl. iii (1890), pp. 37–49.
- Proneomenia sluiteri: anatomy and histology; HEUSCHER, Viert. Ges. Zurich, xxxvii, pp. 148-161, figg.; P. aglaopheniæ, embryogeny of; PRUVOT, C.B. cxiv, pp. 1211-1214.

## V. SCAPHOPODA.

- Structure and relationship of the Solenoconcha: anatomy of Dentalium dentale, and notes on that of Siphonentalis affinis and Cadulus subfusiformis; Plate, Zool. JB. v, Anat. pp. 301-386, figg.: also id. Verh. Deutsch. Zool. Ges. i (1891), pp. 60-63. [For conclusions as to the phylogenetic relationships, see ante, p. 36.]
- Cadulus acus, fig., San Domingo, minusculus, Cape Hatteras, n. spp., Dall., Bull. Mus. C. Z. xviii (1889), p. 432.
- Dentalium: notes on its organization; PLATE, SB. Ges. Marb. 1890 (1891), pp. 26-29.
- D. callipeplum, p. 419, matara, p. 420, calamus, p. 421, taphrium, p. 422, carduus, p. 423, gouldii, p. 424, callithrix, p. 427, Transatlantic and Caribbean, DALL, Bull. Mus. C. Z. xviii (1889), figg.; †D. haeringense, Tertiary of Häring, DREGER, Ann. Hofmuseum Wien, vii, p. 12, fig.: n. spp.
- Lavidentalium, n. sect. of Dentalium, type +D. incertum, Desh.; Coss-MANN, Cat. Coqu. Foss. Eocene Paris, iii (1888), p. 11.
- Lobantale, n. zect. of Dentalium, type †D. duplex, Defr.; Cossmann, Cat. Coqu. Foss. Eccene Paris, iii (1888), p. 11.

## VI. PELECYPODA.\*

New classification proposed by GROBBEN (163, 164).—Some notes on the morphology of the *Pelecupoda*; Kellog, Johns Hopk. Univ. Circ. xi, pp. 80-83.—On the margin of the mantle of *Acephala siphoniata*; Rawitz, Jon. Z. Nat. xxvii, pp. 1-232, figg.—On the phylogeny of the Byssus-apparatus; Thiele, Verh. Deutsch. Zool. Ges. ii, pp. 52-57.

## SEPTIBRANCHIATA.

- Anatomy of the Septibranchiata discussed; Dall, Bull. Mus. C. Z. xviii (1889), pp. 441-452.
- Cuspidaria (Neaera) cuspidata, Olivi, structure; GROBBEN, Arb. z. Inst. Wien, x, No. 3, pp. 1-31, figg.

## EULAMELLIBRANCHIATA.

#### ANATINACEA.

- †Allorisma re-defined; GOODCHILD, P. Phys. Soc. Edinb. xi, pp. 245 & 246.
- Ectorisma, n. g., closely related to Pholadomya, for E. granulata, n. sp., S. Australia; Tate, Tr. R. Soc. S. Austr. xv, p. 127, fig.
- Pandora (Kennerlia) bushiana, n. sp., Florida, DALL in SIMPSON, P. Davenport Ac. v (1889), p. 71.
- †Pholadomya elegantula, n. sp., Pliocene of Savena, near Bologna, FORESTI, Bull Soc. Mal. Ital. xvi, p. 81, fig.
- †Pleuromya gottschei, n. sp., Inf. Oolite of the Argentine Cordilleras, Behrendsen, Z. geol. Ges. xliv, p. 11, fig.
- † Thracia aequilatera, n. sp., Neocomian of the Argentine Cordilleras, Behrendsen, Z. geol. Ges. xliv, p. 21, fig.

#### PHOLADACEA.

- Monograph by Clessin, pp. 25-40 & pls. vii-xii; Martini & Chemnitz (246).
- † Aspidopholus stainieri, n. sp., Tertiary of Antwerp, Vincent, Bull. Soc. Mal. Belg. 1891 [1893], p. lxxxvii, fig.
- †Jouannetia cochlearella, n. sp., Eccene of Belgium, Vincent, Mem. Soc. Mal. Belg. 1891 [1893], p. 166.
- Pholas dactylus: anatomy and physiology; Dubois, Ann. Univ. Lyon, ii, fasc. 2.
- Pholadidæ, from the Eccene of Belgium; Vincent, Mem. Soc. Mal. Belg. 1891 [1893], pp. 164-168.

<sup>\*</sup> Arranged according to Pelseneer's scheme, vide Woodward, Ann. N. H. zi, p. 158.

#### MYACEA.

†Corbula bodenbenderi, p. 19, inflata, nana, p. 20, n. spp., Neocomian of the Argentine Cordilleras, Behrendsen, Z. geol. Ges. xliv, figg.

#### CARDIACEA.

- Ceratoconcha, n. g. for C. costata, n. sp., Tertiary of Agram; GORJANO-VIC-KRAMBERGER, Glasnik Naravosl. Druzt. iv (1889), p. 50, fig.
- Chama nicolloni, n. sp., West Coast of France, DAUTZENBERG, Bull. Soc. Ouest France, ii, p. 133, figg.
- †Cornucaprina, n. g., for Schiosia carinata, G. Böhm; FUTTERER, Pal. Abh. vi, p. 87, fig.
- †Limnocardium jagici, p. 186, kosici, p. 188, margaritaceum, p. 190, n. spp., Tertiary of Markuševec (Croatia); Brusina, Glasnik Naravosl. Druzt. vii.
- † Orthoptychus, n. g., for O. striatus, n. sp., Upper Cretaceous of the Venetian Alps; FUTTERER, Pal. Abh. vi, p. 91, figg.
- †Schiosia, n. sect. for S. schiosensis, n. sp., Cretaceous of the Venetian Alps; BOEHM, Ber. Ges. Freiburg, vi, p. 11, with other new species of Caprinidæ.
- Sphærium nyanzæ, n. sp., Victoria Nyanza, Smith, Ann. N. H. x, p. 383.

#### VENERACEA.

- †Conchodus: notes on the genus, with C. schwageri, n. sp., Upper Trias of the Northern Alps; TAUSCH, Abh. Geol. Reichsanst. xvii, No. 1, 8 pp., 3 pls.
- Corbicula trajecta, n. n. for C. tongkingensis, Clessin; FISCHER, Bull. Soc. Autun, iv, p. 240.
- †Cyprina argentina, n. sp., Neocomian of the Argentine Cordilleras; Behrendsen, Z. geol. Ges. xliv, p. 22, fig.
- Cytherea convexa, Say, preocc. by Brongniart, for a Paris Basin species; C. sayana, Conrad, stands; Dall, Naut. vi, p. 52.
- C. gracilenta, rugata, nitidula, n. spp. ?, W. Coast of France, Locard, Échange, vii (1891), p. 69.
- † Isocardia koeneni, n. sp., Neocomian of the Argentine Cordilleras, Behrendsen, Z. geol. Ges. xliv, p. 22, fig.
- Isocardiidæ, n. n. for Cyprinidæ; Bucquoy & others, (55) p. 313.
- Meroë ovalis, n. sp., Port Elizabeth, Sowerby, (378) p. 59, fig.
- †Pachyerisma: on the scar of the pedal muscle; BOEHM, Ber. Ges. Freiburg, vi, pp. 119 & 120.
- Petricola ponsonbyi, n. sp., Port Elizabeth, Sowerby, (378) p. 61, fig.
- † Ptychomya koeneni, n. sp., Neocomian of the Argentine Cordilleras, Behrendsen, Z. geol. Ges. xliv, p. 23, fig.

#### TELLINACEA.

Donax bipartitus, n. sp., South Africa, Sowerby, (378) p. 58, fig.

Herouvalia defined; Cossmann, Mem. Soc. Mal. Belg. 1891 (1892), p. 28.

Mactra: on the systematic position of Mactra and its allies; BITTNER, Verh. geol. Reichsanst. 1892, pp. 232-241.

Mactra capensis, n. sp., Port Elizabeth, Sowerby, (378) p. 55, fig.

Syndosmya schneideri, n. sp., Borkum Is., BETTGER in SCHNEIDER, Nachr. mal. Ges. 1892, p. 116.

Tellina (Macoma) crawfordi, rousi, n. spp., Port Elizabeth, Sowerby, (378) p. 57, figg.

## SUBMYTILACEA.

# (Marine Forms.)

- † Anthracosia venjukowi, p. 155, subnucleus, p. 156, loewinsoni, p. 157, oviformis, truncata, obscura, p. 158, n. spp., Permian of Russia, Amalizky, Palssontogr. xxxix, figg.
- † Carbonicola subovalis, p. 146, striata, p. 148, substegocephalum, p. 149, recta, indeterminata, p. 150, nova, p. 152, n. spp., Permian of Russia, AMALIZKY, Palsoontogr. xxxix, figg.
- Cardita eluta, Port Elizabeth, Sowerby, (378) p. 63, fig.; C: gemmulifera, S. Australia, Tate, Tr. B. Soc. S. Austr. xv, p. 130, fig.: n. spp.

Carditella rugosa, n. sp., South Africa, Sowerby, (378) p. 63, fig.

Felania subradiata, n. sp., Port Elizabeth, Sowerby, (378) p. 62, fig.

Lucina per-obliqua, p. 128, paupera, p. 129, n. spp., S. Australia, TATE, Tr. R. Soc. S. Austr. xv, figg.

† Myoconcha parvula, n. sp., Alpine Trias, Woehrmann & Koken, Z. geol. Ges. xliv, p. 177, fig.

Mysella ovalis, n. sp., S. Australia, TATE, Tr. B. Soc. S. Austr. xv, p. 128, fig.

Neocardia, n. g. for N. agulata, n. sp., Port Elizabeth, Sowerby, (378) p. 63, fig.

†Opis pellati, n. sp., Oxfordian of Boulogne, RIGAUX, Mém. Soc. Acad. Boulogne, xiv, p. 107, fig.

†Plethocardia, n. g. for P. umbonata and suberecta, n. spp.; Ulrich, Rep. Geol. Surv. Minnesota, No. 19, pp. 243-245, figg.

Tellimya producta, n. sp., St. Helens, Smith, Ann. N. H. x, p. 130.

†Trigonodus costatus, p. 186, minutus, p. 187, n. spp., Alpine Trias, Woehrmann & Koken, Z. geol. Ges. xliv, figg.

## (Freshwater Forms.)

# Unionide and Dreissensiide.

Anodonta: notes on; JHERING, Zool. Ans. xv, pp. 1-5.

A. chapalensis, Mexico, CROSSE & FISCHER, J. de Conch. 1892, p. 295;
A. graca, p. 90, quellenneci, p. 91, lepida, p. 92, Lake Copais, Greece.

- lhotellerii, Belgium, DROUET, J. de Conch. 1892; A. recurvirostris, (Küster MS.) gallensteini, Carinthia, Kobelt in Rossmaessler, (322) vi, p. 47, figg.: n. spp.
- Congeria and Dreissensia, their relationship and distribution in space and time; OPPENHEIM, Z. geol. Ges. xliii, pp. 923-966, 1 pl. C. triangularis, notes on the group; BRUSINA, Z. geol. Ges. xliv, pp. 488-497.
- †C. harnesi, p. 492, ornithopsis, p. 495, BRUSINA, Z. geol. Ges. xliv; †C. mytilopsis, ramphora, p. 182, doderleini, p. 183, gitneri, p. 184, Tertiary of Markuševec, BRUSINA, Glasnik Naravosl. Druzt. vii: n. spp.
- Cristaria [= Dipsas], note on; JHERING, Nachr. mal. Ges. 1892, pp. 1-14.
- Dipsas plicata, Lea: anatomy; ISHIKAWA (192A).
- Dreissensia and Congeria, their relationship and distribution in space and time; OPPENHEIM, Z. geol. Ges. xliii, pp. 923-966, 1 pl. Dreissen[si]a polymorpha, development; Korschelt, Ann. N. H. ix, pp. 157-169, figg.
- D. massiei, n. sp., Laos, Morlet, J. de Conch. 1892, p. 85.
- † Dreissensiidæ: fossil species of Dalmatia, Croatia, and Slavonia; Brusina, Glasnik Naravosl. Druzt. vii, pp. 195-199.
- Glabaris: notes on; JHERING, Zool. Anz. xv, pp. 1-5.
- Margarituna margaritifer, Linn., notes on a specimen of; Moebius, SB. nat. Fr. 1892, p. 92.
- †Najadites verneuili, p. 187, fischeri, subcastor, p. 191, okensis, p. 193, parallela, sibirzewi, p. 194, indeterminata, dubia, monstrum, p. 195, n. spp., Permian of Russia; AMALIZKY, Palsontogr. xxxix, figg.
- † Oligodon, n. g., p. 179, for O. kingi, p. 181, geinitzi, zitteli, p. 182, n. spp., Permian of Russia, Amalizky, Palseontogr. xxxix, figg.
- † Palæomutela, n. g., p. 159, for P. verneuili, p. 163, subparallela, solenoides, p. 164, trapezoidalis, compressa, p. 165, lunulata, semilunulata, p. 166, keyserlingi, p. 168, orthodonta, ovalis, p. 169, subovalis, trigonalis, p. 170, irregularis, golowkinskiana, p. 171, rectangularis, obliqua, triangularis, p. 174, vaga, parva, p. 175, elegantissima, p. 176, murchinoni, plana, lævis, p. 177, crassa, gorbatowi, p. 178, curiosa, p. 179, n. spp., Permian of Russia, Amalizky, Palæontogr. xxxix, figg.
- Pseudodon crassus, æneolus, Borneo; DROUET & CHAPER, Mém. Soc. Zool. v, pp. 151 & 152, figg. (see also J. de Conch. 1892, pp. 93 & 94). Spatha subaequilatera, n. sp., Victoria Nyanza, MARTENS, Conch. Mittheil. iii (1889), p 18, fig.
- Spathella bloyeti, p. 198, spathuliformis, p. 199, E. Centr. Africa, n. spp., Bourguignat, (37) 1889, figg.
- Unionidæ: on a revision of the American Unionidæ; SIMPSON, Naut. vi, pp. 78-80: preliminary list of New York species; MARSHALL, Bull. New York Mus. Nat. Hist i, 17 pp. Unionidæ of New York State; MARSHALL, Bull. New York Mus. Nat. Hist. i, No. 1. Unionidæ of Spoon River, Illinois; STRODE, Am. Nat. xxvi, pp. 495-501.

- Unio: variability of the Tasmanian; Johnston, P. R. Soc. Tasm. 1888 (1889), pp. 95 & 96.
- U. luteolus and radiatus, more notes on; DEAN, Naut. v, p. 112. U. vagulus, n. n. for U. subtrigonus, Sow., non Deshayes; FISCHER, Bull. Soc. Autun. iv, p. 223. U. quintardii, Cragin, pilsbryi, Marsh, and (Arconuia) provancheriana, Pilsbry, figured; PILSBRY, P. Ac. Philad. 1892, pp. 131 & 132, 2 pls.
- U. usumasinta, yzabalensis, Guatemala, Crosse & Fischer, J. de Conch. 1892, pp. 294; U. corbeti, Lake Kaudy, Ceylon, Deschamps, Bull. Soc. Z. Fr. xvii, p. 68, fig.; U. serradelli, Philippines, p. 86, helenæ, Corfu, p. 87, thiessea, Greece, p. 88, clessini, Danube [i.e. the Douau] at Lauingen, p. 89, Drouet, J. de Conch. 1892; U. lugens, p. 147, saccellus, p. 148, lingulatus, p. 149, radulosus, p. 150, trompi, p. 153, fulvaster, p. 154, Borneo, Drouet & Chaper, Mém. Soc. Z. Fr. vi, figg.; U. dugasti, Laos, Morlet, J. de Conch. 1892, p. 86; U. subluridus, Orange Springs, Volusia County, Florida, Simpson, P. U. S. Nat. Mus. xv, p. 432, fig.: n. spp.

## PSEUDOLAMELLIBRANCHIATA.

#### OSTREIDE and AVICULIDE.

- †Avicula kokeni, n. sp., Alpine Trias, Woehrmann & Koken, Z. geol. Ges. xliv, p. 175, fig.
- †Cleionychia, n. g., type Ambonychia lamellosa, Hall, with C. rhomboidea, n. sp., Treuton Limestone of U. S.; Ulrich, Am. Geol. x, p. 97, fig.
- †Lithiotis problematica, Gümb., notes on; BOEHM, Ber. Ges. Freiburg, vi, pp. 65-80, figg.
- † Maccoyella, n. g., type Avicula barklyi, Moore, Rolling Downs formation of Queensland; JACK & ETHERIDGE, (193) p. 451, fig.
- † Merismopteria, n.g., type Pterinea macroptera, from the Permo-Carboniferous of Queenslaud; JACK & ETHERIDGE, (193) p. 271, fig.
- Ostrea: notes on oysters; Schieldt (334, 335, 336).
- † Perna nana, n. sp., Inferior Oolite of the Argentine Cordilleras, Beh-BENDSEN, Z. geol. Gos. xliv, p. 11, fig.
- † Pinna tomassi, n. sp., Alpine Trias, Woehrmann & Koken, Z. geol. Ges. xliv, p. 177, fig.
- † Posidonomya steinmanni, n. sp., Inferior Oolite of the Argentine Cordilleras, Behrendsen, Z. geol. Ges. xliv, p. 10, fig.
- † Pseudavicula, n. g., type Lucina anomala, Moore, from the Rolling Downs Beds of Queensland; JACK & ETHERIDGE, (193) p. 449, fig.

#### PECTINACEA.

- Amusium dalli: gill structure; DALL, Bull. Mus. C. Z. xviii (1889), p. 433.
- † Deltopecten, n. g., type Pecten illawarensis, Morris, intermediate between

- Pecter and Aviculopecter, from the Permo-Carboniferous of Queensland; JACK & ETHERIDGE, (193) p. 269, fig.
- Dimya argentea: gill structure; Dall., Bull. Mus. C. Z. xviii (1889), p. 433.
- Pecten crouchi, Mauritius, SMITH, Ann. N. H. ix, p. 255, fig.; † P. zitteli, Alpine Trias, Woehrmann & Koken, Z. geol. Ges. xliv, p. 173, fig.: n. spp.
- Spondylus gæderopus, L., disappearance from Maltese Waters; CARUANA-GATTO, Med. Nat. i, p. 165.
- S. powelli, n. sp., Madeira, SMITH, Journ. Conch. vii, p. 70.

## FILIBRANCHIATA.

- Anomiidæ of the Coast of France, with Anomia boletiformis, n. sp. [?]; LOCARD, Échange, vii (1891), p. 86.
- Arca ectocomata: gill structure; DALL, Bull. Mus. C. Z. 1889, p. 433.
- Barbatia helblingia: foot and byssus glands figured; Sluiter, Tijdschr. Nederl. Dierk. Ver. iii, pp. 170-184.
- Lithodomus dactylus, on the boring powers of; CARAZZI, Atti Soc. Ligust. iii, p. 279-297, fig.
- L. projectuns, cuneiformis, S. Australia, n. spp., TATE, Tr. R. Soc. S. Austr. xv, pp. 131 & 132, figg.
- † Myophoria (?) plana, n. sp., Alpine Trias, Woehrmann & Koken. Z. geol. Ges. xliv, p. 183, fig.
- Myrina crenatulifera, n. sp., S. Australia, TATE, Tr. R. Soc. S. Austr. xv, p. 131, fig.
- Pectunculus: species living in the Mediterranean, and fossil in the Upper Tertiary strata of Italy; GREGORIO, Nat. Sicil. xi, pp. 89-96, 106-114, & 209-212; Monterosato, t. c. pp. 143-155.
- †Schizodus negrii, n. sp., Trias of Balia, Asia Minor, BITTNER, JB. Geol. Reichsanst. xlii, p. 86, fig.
- †Trigonia densestriata, Inferior Oolite of the Argentine Cordilleras, Behrendsen, Z. geol. Ges. xliv, p. 12, fig.; †T. transatlantica. p. 27, anguste-costata, p. 28, Upper Cretaceous of the Argentine Cordilleras, Behrendsen, Z. geol. Ges. xliv, figg.; †T. seeleyi, Lower Oolites of Boulogne, Bigaux, Mein. Soc. Acad. Boulogne, xiv, p. 106, fig. : n. spp.

### PROTOBRANCHIATA.

- Nucula: list of species belonging to the subgenus Acila; SMITH, Journ. Conch. vii, pp. 110-112.
- N. (Acila) fultoni, n. sp., Bay of Bengal, SMITH, Journ. Conch. vii, p. 111.

- † Nuculites beneckei, p. 46, kruegeri, p. 48, n. spp., Paleozoic of Bolivia, ULRICH (412), figg.
- †Palæoneilo forbesi, n. sp., Palæozoic of Bolivia, Ulrich, (412), p. 42, fig.
- † Yoldia beyrichi, n. sp., Rupelian of Germany, BŒTTGER, Mal. Bl. xi, p. 92.

### INCERTA SEDES.

† Annigenia rhenana, n. sp., Middle Devonian of Germany, BEUSHAUSEN, JB. Preuss. geol. Landesanst. xi, Abhandl. p. 1, fig.

# VIII. BRACHIOPODA.

BY

B. B. WOODWARD, F.G.S., F.R.M.S., &c.

# I.—TITLES.\*

- 1. Beecher, C. E. Development of the *Brachiopoda*. Pt. 11. Classification of the stages of growth and decline. Am. J. Sci. xliv, pp. 133-154, 1 pl.
- Ueber die Entwickelung der Brachiopoden. JB. Mineral. 1892, i, pp. 178-197, 1 pl.

Translation of article in Am. J. Sci. 1891. See Zool. Rec. 1891.

- BIERBAUER, B. A Check-list of the Palæozoic Fossils of Wisconsin, Minnesota, Iowa, Dakota, and Nebraska. Bull. Minnesota Acad. iii. [Brachiopoda, pp. 226-234.]
- BITTNER, A. Neue Arten aus der Trias von Balia in Kleissasien.
   JB. geol. Reichsanst. xlii, pp. 77-89, 2 pls.
- Brachiopoden der Alpinen Trias. Nachtrag i. Abh. geol. Reichsanst. xii, No. 2, 38 pp., 4 pls.
- BLOCHMANN, F. Ueber die Anatomie und die verwandtschaftlichen Beziehungen der Brachiopoden. Arch. Ver. Mecklenb. xlvi, pp. 37-50.
- Untersuchungen über den Bau der Brachiopoden. 1. Die Anatomie von Crania anomala, O. F. M. Jena: 1892, 4to, 65 pp., 7 pls.

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

 BOESE, E., & FINKELSTEIN, H. Die mitteljurassischen Brachiopoden-Schichten bei Castel Tesino im östlichen Südtirol. Z. geol. Ges. xliv, pp. 265-302.

Brachiopoda begin p. 285.

- Calvin, S. Two unique Spirifers from the Devonian Strata of Iowa. Bull. Lab. Iowa, ii, pp. 165-167, 1 pl.
  - CLARKE, J. M. [See HALL & CLARKE (17).]
- CRANE, MISS A. Recent observations on the anatomy and development of the Brachiopoda. Nat. Sci. 1892, pp. 603-611.
- Derby, O. A. Nota sobre a geologia e paleontologia de Matto Grosso. [Brachiopoda with Notothyris (?) smithii and Centronella margarida, n. spp.] Arch. Mus. R. Jan. ix, pp. 59-88.
- 12. ETHERIDGE, R., JUN. On the occurrence of an Orthis [O. leviensis], allied to O. actonia, J. de C. Sby., and O. flabellulum, J. de C. Sby., in the Lower Silurian rocks of Central Australia. In BROWN Reports on the Coal-bearing area near Leigh's Creek [Govt. Rept.], 1891, pp. 13 & 14, figg.
- 13. —. The *Pentamerida* of New South Wales. Rec. Geol. Surv. N.S.W. iii, pp. 49-60, 2 pls.
  - FINKELSTEIN, H. [See BOESE & FINKELSTEIN (8).]
- FISCHER, P., & CEHLERT, P. Mission scientifique du Cap Horn (1882-83)—Brachiopodes. Bull. Soc. Autun, v, pp. 254-334, 5 pls., text illust.
- & —. Brachiopodes de l'Atlantique Nord. [Fasc. iii of Résultats des campagnes scientifiques accomplies sur son yacht par le ...Prince de Monaco.] Monaco: 1892, 4to, 30 pp., 2 pls.
- —— & ——. Sur l'évolution de l'appareil branchial de quelques Brachiopodes. C.R. cxv, pp. 749-751.
- HALL, J., & CLARKE, J. M. An introduction to the study of the genera of Palsozoic Brachiopoda. Pt. I. Geol. Surv. State of New York. Palsontology viii, pp. xvi & 367, 41 pls.
- Johnston, R. M. Additions to the list of Tasmanian Fossils [Brachiopoda] of Upper Palssozoic age. P. B. Soc. Tasm. 1889 (1890), pp. 137-141, 1 pl.
- JOUBIN, L. Recherches sur l'anatomie de Waldheimia venosa, Sol. Mém. Soc. Zool. v, pp. 554-583, figg.
- KAYSER, E. Beiträge zur Kenntniss der Fauna der Siegenschen Grauwacke. JB. k. preuss. geol. Landesanst. 1890, pp. 95–107, 5 pls.
- 21. MATTHEW, G. F. Trematobolus: an articulate Brachiopod of the inarticulate Order. Can. Rec. v, pp. 276-279, fig.

- Newton, R. B. On the occurrence of *Ohonetes pratti*, Davidson, in the Carboniferous rocks of Western Australia. Geol. Mag. 1892, pp. 468, 469, & 542-544, 1 pl.
  - ŒHLERT, D. P., & FISCHER, P. [See FISCHER & ŒIILERT (14-16).]
- RIGAUX, E. Notice géologique sur le Bas Boulonnais. Mém. Soc. Acad. Boulogne, xiv [n. spp. described on pp. 102-108, 2 pls.].
- ROTHPLETZ, A. Die Perm-, Trias- und Jura-formation auf Timor und Rotti im Indischen Archipel. Palmontogr. xxxix, Permian Brackiopoda, pp. 76-85, figg.
- 25. Schellwien, E. Die Fanna des Karnischen Fusilinenkalks... Thl. I.... Brachiopoda. Palmontogr. xxxix, pp. 1-56, 8 pls. Schuchert, C. [See Winchell & Schuchert (31).]
- SIMPSON, G. B. Descriptions of new species of fossils from the Clinton...and Waverley groups, &c. Tr. Am. Phil. Soc. xvi (1890) [Brachiopoda, pp. 437-445, figg.]
- 27. Sowerby, G. B. Marine shells of South Africa, &c. 4to, London. [6 spp. enumerated at p. 67.]
- Ulrich, A. Palæozoische Versteinerungen aus Bolivien. Steinmann's Beitr. Geol. Palæont. S. Amer. i, 116 pp., 5 pls. Stuttgart: 1892, 8vo.
- 29. WALKER, J. F. The discovery of *Terebratulina substriata*, Schlotheim, in Yorkshire. Geol. Mag. 1892, p. 364.
- 30. On Yorkshire Thecidea. T. c. p. 548.
- WINCHELL, N. H., & SCHUCHERT, C. Preliminary descriptions of new Brachiopoda from the Trenton and Hudson River groups of Minnesota. Am. Geol. ix, pp. 284-294.

# II.—ANATOMY, &c.

Anatomy and relationships of the *Brachiopoda*; BLOCHMANN (6), who considers them closely related to *Phoronis*.—Recent observations on the anatomy and development of the *Brachiopoda* [a summary]; CRANE (10).—Anatomy of *Orania anomala*, O.F.M.; BLOCHMANN (7).—Anatomy with figg. of *Waldheimia venosa*, Solander; JOUBIN (19).

Development of *Brachiopoda*, stages of growth and decline; Brecher (1).

Evolution of the branchial apparatus of Brachiopoda; FISCHER & CEHLERT (16).

Observations on the characters of the shells of Palæozoic *Brachiopoda*, and such structures in the recent forms as throw light upon them are given in HALL & CLARKE (17).

### III.—DISTRIBUTION.

#### A.—GEOGRAPHICAL.

English Channel: Brachiopoda off the Coast of the Department of the Somme; Bizer, Mém. Soc. L. N. Fr. viii, pp. 390-392.

North Atlantic forms; FISCHER & ŒHLERT (15).

South Africa: 6 species recorded from; Sowerby (27).

Australia: third supplement to list of Paliobranchs; Tate (see ante, Moll. 399a).

Cape Horn species; FISCHER & ŒHLERT (14).

#### B.—GEOLOGICAL.

Summary of the paleontology of 1890, Brachiopoda by D. P. ŒILERT, Ann. Géol. univ. Paris, 1890 (1892), pp. 1025-1039.

Fossil types in the Bath Museum, P. Bath N. H. Soc. vii, pp. 279-281. Tertiary species from Queensland, 1 n. sp., JACK & ETHERIDGE, (unte, Moll. 193) pp. 561-574.

Eccene Brachiopoda, from the environs of Paris; Mem. Soc. Mal. Belg. 1891, pp. 15-20.

Secondary Brachiopoda, from the Balkans, with new species; VANKOV, Rad. jugoslav. akad. cxi, pp. 92 & 93.—Secondary Brachiopoda, from Queensland and New Guinea, with new species; JACK & ETHERIDGE, (ante, Moll. 193) pp. 442-444.

Cretaceous Brachiopoda of Schleswig-Holstein; STOLLY, MT. min. Inst. Kiel, i, pp. 245-248.—Upper Senonian Brachiopoda of the Eifel; Vogel, Verh. Ver. Rheinl. xlix, pp. 87-90.—3 species quoted from the Cretaceous of the Podol'ski Government; RADKEVICH, Zapiski Kiev, xi (1891) p. 83.

Jurassic Brachiopoda of Yorkshire, List; Fox-Strangways, Mem. Geol. Surv. U. K. Jurassic Rocks, vol. ii, pp. 40-45, 148-150, & 201-204. —Middle Jurassic Brachiopoda from Castel Tesino, in S. E. Tirol; BOESE & FINKELSTEIN (8).—Rhynconella pellati, n. sp., Oxfordian of Boulogne; RIGAUX (23).

Liassic Brachiopoda from the Ironstone, near Helmstedt, with new species; WOLLEMANN, Verh. Ver. Rheinl. xlix, pp. 125-146.

Triassic Brachiopoda of the Duchy of Sachsen-Altenburg; GEINITZ, MT. Osterlande, v, pp. 189 & 190.—Triassic species from the Alps, with many new forms, and Dinarella, n. g.; BITTNER (4).—Waldheimia bukowskii, Spiriferina moscai, and Spirigera tricuspii, n. spp., Trias of Balia, Asia Minor; BITTNER (5).

Palæozoic Brachiopoda of the State of New York (Inarticulata and Articulata, Orthis to Productus), with many new genera and new species; HALL & CLARKE (17).—Check-list of the Palæozoic Brachiopoda from Wisconsin, Minnesota, Iowa, Dakota, and Nebraska; BIERBAUER (3).—New Brachiopoda from the Palæozoic of Roulogne; RIGAUX (23).—Permian and Devonian Brachiopoda of the Duchy of Sachsen-Akenburg;

Geinitz, MT. Osterlande, v, pp. 169, 177 & 178.—Upper Palæozoic Brachiopoda of Tasmania; Johnston (18).—Middle Devonian and Permo-Carboniferous species from Queensland, with new and indeterminate species; Jack & Etherioge (ants, Moll. 193) pp. 64-68, 225-264.

Permian Brackiopoda from Timor and Rotti, with new species; ROTH-PLETZ (24).

Carboniferous: Brachiopoda of the Carinthian Fusilina Limestone, with many new species; Schellwien (25).—Grauwacke Fauna of Siegen, with Discina, 2 new species; Kayser, JB. k. preuss. geol. Landesanst. xi, Abbandl. pp. 95-107, figg.—Chonetes pratti from the Carboniferous of W. Australia; Newton (22).

Devonian Brachiopoda of the Eastern Alps, with new species; FRECH, Z. geol. Ges. xliii, pp. 672-680.—Devonian Brachiopoda from Matto Grosso: Notothyris (?) smithii and Centronella (?) margarida, n. spp.; Derby (11).—Devonian of Bolivia, with 7 new species; Ulrich (28).—Devonian species from Manitoba, with 2 new species; Whiteaves, Contr. Canad. Pal. i, pp. 281-291, figg. — 9 new species from the Clinton, Lower Helderberg, Chemung and Waverley Groups of Pennsylvania: Simpson (26).—Spirifer urbana and macbridei, from the Devonian of Iowa, figured by Calvin (9).—Terebratula iowensis, n. sp., Devonian, Iowa; Calvin, Bull. Lab. Iowa, i, p. 174, fig.—Fauna of the St. John Group, Canada, with new species of Brachiopoda; Matthew, Tr. R. Soc. Canada, ix (4), pp. 39-50.—Trematobolus, n. g., St. John Group, New Brunswick; Matthew (21).—Lower Devonian Brachiopoda, with 4 new species; Béclard, Bull. Soc. Belg. Géol. v, pp. 96-102, figg.

Siluro-Devonian Pentamerida of Australia; ETHERIDGE (13).

Silurian: New species from the Trenton and Hudson River Groups of Minnesota, with Hullina, n. g., Winchell & Schucher (31).—Orthis leviensis, n. sp., and flabellulum, J. de C. Sby., from the Lower Silurian of Central Australia; Etheridge (12).

Cambrian fossils of S. Australia, with Orthisina compta and Orthis (?) peculiaris, n. spp.; TATE, Tr. R. Soc. S. Austr. xv, p. 185.

#### IV.—SYSTEMATIC.

#### INARTICULATA.

† Barroisella, n. g., type Lingula subspatula, Meek & W.; HALL & CLARKE, (17) p. 62, figg.

Crania anomala, O.F.M., anatomy; BLOCHMANN (7).

Discina siegenensis, p. 95, anomala, p. 96, n. spp., Grauwacke of Siegen, KAYSER, JB. k. preuss. geol. Landesanst. 1890, figg.

† Discinopsis, n. g., type Acrotreta? gulielmi, Matth.; HALL & CLARKE, (17) p. 105, figg.

†Lindstræmella, n. subg. of Discina, type L. aspidum, n. sp., Palæozoic of New York State; Hall & Clarke, (17) p. 134, figg.

- † Chlertella, n. subg. of Discina, type D. pleurites, Meek; HALL & CLARKE, (17) p. 133.
- †Roemerella, n. subg. of Discina, type D. grandis, Vanux.; HALL & CLARKE, (17) p. 137, fig.
- †Tomasina, n. g., type Lingula criei, Dav.; HALL & CLARKE, (17) p. 65, figg.
- † Trematobolus, n. g. for T. insignis, St. John Group of St. Martins, New Brunswick; Matthew, Can. Rec. v, p. 276, fig. This genus is connected with the Obolidæ, but differs in having articulate valves. It may be looked on as combining the features of Siphonotreta, Schizambon, and Obolus in its ventral valve, with those of Obolus, Obolella, and Orthis in its dorsal valve.

#### ARTICULATA.

#### PRODUCTIDAL.

- †Chonetes pratti, Davida, from Carboniferous of W. Australia; Newton, Geol. Mag. 1892, pp. 468, 469, & 542-544, fig.
- †C. douvillei, Devonian of Boulogne, RIGAUX, Mém. Soc. Acad. Boulogne, xiv, p. 104, fig.; †C. arcei, p. 77, ruecki, p. 79, stuebeli, p. 80, Devonian of Bolivia, ULRICH, (28) figg.: n. spp.
- † Chonopertus, n. g., type Chonetes fischeri, Norw.; HALL & CLARKE, (17) p. 312, fig.
- † Chonostrophia, n. g., type Chonetes reversa, Whitf.; HALL & CLARKE, (17) p. 310, fig.

#### STROPHOMENIDE.

- †Billingsella, n. g., type Orthis pepina, Hall; HALL & CLARKE, (17) p. 230, figg.
- †Christiania, n. g., type Leptæna subquadruta, Hall; HALL & CLARKE, (17) p. 298, fig.
- † Dalmanella, n. sect. of Orthis, group of O. testudinaria, Dalm.; Hall & Clarke, (17) p. 205, fig.
- † Dinorthis, n. sect. of Orthis, group O. pectinella, Emm.; HALL & CLARKE, (17) p. 195, fig.
- †Herbertella, n. sect. of Orthis, group O. sinuata, Hall; HALL & CLARKE, (17) p. 198, fig.
- † Heterorthis, n. sect. of Orthis, group O. clytie, Hall; HALL & CLARKE, (17) p. 202, figg.
- †Kayserella, n. g., type Orthis lepida, Schnur.; HALL & CLARKE, (17) p. 259, fig.
- † Leptella, n. g., type Leptæna sordida, Billings; Hall & Clarke, (17) p. 293, fig.

- †Orthidium, n. g., type Orthis gemmicula, Billings; Hall & Clarke, (17) p. 244, fig.
- †Orthis leviensis, n. sp., from the Lower Silurian of Central Australia; ETHERIDGE, (12) p. 24, figg.
- † Orthotichia, n. sect. of Orthis, group O. ? morganiana, Derby; Hall & Clarke, (17) p. 213, figg.
- †Plæsiomys, n. sect. of Orthis, group O. subquadrata, Hall; Hall & Clarke, (17) p. 196, fig.
- † Plectorthis, n. sect. of Orthis, group of O. plicatella, Hall; Hall & Clarke, (17) p. 194, figg.
- † Polytoechia, n. subg. of Clitambonites, type Hemipronites apicalis, Whitf.; HALL & CLARKE, (17) p. 239, fig.
- † Protorthis, n. g., type Orthis billingsi, Hartt.; HALL & CLARKE, (17) p. 231, fig.
- † Rafinesquina, n. g., type Leptana alternata, Conr.; HALL & CLARKE, (17) p. 281, fig.

#### SPIRIFERIDAL

- †Athyris betencourti, n. sp., Devonian, Boulogne; RIGAUX, Mém. Soc. Acad. Boulogne, xiv, p. 103, fig.
- †Cyrtina lonquetii, n. sp., Carboniferous of Boulogne; RIGAUX, Mém. Soc. Acad. Boulogne, xiv, p. 105, fig.
- † Meristella riskowskyi, n. sp., Devonian of Bolivia; ULRICH, (28) p. 64, figg.
- † Spirifer urbana and macbridei, Devonian, Iowa; figured by CALVIN, Bull. Lab. Iowa, ii, pl. xii.
- †S. barroisi, Devonian, Boulogne, RIGAUX, Mém. Soc. Acad. Boulogne, xiv, p. 102, fig.; †S. chuquisaca, Devonian of Bolivia, ULRICH, (28) p. 65, figg.: n. spp.
- † Spiriferina moscai, n. sp., Trias of Balia, Asia Minor; BITTNER, JB. geol. Reichsanst, xlii, p. 80, fig.
- †Spirigera tricupii, n. sp., Trias of Balia, Asia Minor; BITTNER, JB. geol. Reichsanst. xlii, p. 82, fig.
- † Thecocyrtella n. n. for Cyrtotheca, Bittner non Salter; BITTNER, Abh. geol. Beichsanst. xvii, No. 2, p. 15.

#### RHYNCHONELLIDÆ.

- †Pentameridæ of New South Wales; ETHERIDGE, Rec. Geol. Surv. N.S.W. iii, pp. 49-60, 2 pls.
- †Rhynchonella symptychos, Middle Jurassic of S. E. Tirol, Boese & Finkelstein, Z. geol. Ges. xliv, p. 300, fig.; R. lemeslii, Devonian, p. 103, pellati, Oxfordian, p. 106, of Boulogne, Rigaux, Mém. Soc. Acad. Boulogne, xiv, figg.: n. spp.

## TEREBRATULIDÆ.

- † Centronella (?) margarida, Devonian, Matto Grosso, DERBY, Arch. Mus. R. Jan. ix, p. 84; C. silvetii, p. 51, arcsi, p. 53, Devonian of Bolivia, ULRICH, (28) figg.: n. spp.
- † Dinarella, n. g., with D. haueri, n. sp., Trias of the Alps, BITTNER, Abh. geol. Reichsanst. xvii, No. 2, p. 24, figg.
- Hallina, n. g. for H. saffordi, and nicolleti, n. spp., Lower Silurian of Minnesota, Winchell & Schuchert, Am. Geol. ix, pp. 291-293.
- † Notothyris (?) smithii, n. sp., Devonian of Matto Grosso, DERBY, Arch. Mus. R. Jan. ix, p. 81, fig.
- † Terebratula haasi, p. 287, vespertilio, p. 290, Middle Jurassic of S. E. Tirol, n. spp., Boese & Finkelstein, Z. geol. Ges. xliv, figg.
- † Terebratulina substriata, Schloth., in Yorkshire; WALKER, Geol. Mag. 1892, p. 364.
- Waldheimia venosa, Soland., anatomy with figg.; Joubin, Mém. Soc. Zool. v, pp. 554-583.
- † W. bukowskii, Trias of Balia, Asia Minor, BITTNER, JB. geol. Reichsanst. xlii, p. 79, fig.; † W. tesinensis, Middle Jurassic of S. E. Tirol, BOESE & FINKELSTEIN, Z. geol. Ges. xliv, p. 285, fig.: n. spp.

#### THECIDEIDE.

† Thecidea: on Yorkshire species; WALKER, Geol. Mag. 1892, p. 548.

# IX. BRYOZOA (POLYZOA).

BY

B. B. WOODWARD, F.G.S., F.R.M.S., &c.

(With kind assistance from R. KIRKPATRICK, Esq., F.Z.S., &c.)

## I.—TITLES.

- 1. AMI, H. M. Notes and descriptions of some new or hitherto unrecorded species of Fossils from the . . . Ordovician rocks of the Province of Quebec . . . Bryosoa. Can. Rec. v, pp. 96-103.
- BIERBAUER, B. A Check-list of the Palæozoic Fossils of Wisconsin, Minnesota, Iowa, Dakota, and Nebraska. Bull. Minnesota Acad. iii. Polysoa, pp. 222-226.
- Bizet, E. Catalogue des Mollusques observés à l'état vivant dans le Département de la Somme. Mém. Soc. L. N. Fr. viii. Bryozoa, pp. 393-402.
- Braem, F. Die keimblätter der Bryozoen-Knospe. Zool. Anz. xv, pp. 113-115.

For reply see DAVENPORT (5).

 DAVENPORT, C. B. The germ-layers in Bryozoan buds. Zool. Anz. xv, pp. 261-263.

A reply to Braem (4).

- Demade, P. Le Statoblaste des Phylactolémates (Bryozoaires d'eau douce) étudié chez l'Alcyonella fungesa et la Cristatella mucedo. La Cellule, viii, pp. 333-378, 2 pls.
- Hallez, P. Draguages effectués dans le Pas-de-Calais... 1888 et 1889. Rev. Biol. ii (1889) [Bryozoa], pp. 37 & 38.
- Addition à la liste des Bryozoaires du Boulonnais Op. cit. iv (1891), pp. 119 & 120.
- Deuxième supplément à la liste des Bryozoaires du Boulonnais.
   Op. cit. ▼ (1892), pp. 123 & 124.

- HARMER, S. F. On the nature of the Excretory Processes in Marine Polyzoa. Q. J. Micr. Sci. xxxiii, pp. 123-167, 2 pls.
- Helm, S. Cordylophora lacustris and five forms of Animal Life. J. N. Y. Micr. Soc. viii, pp. 43-50, figg.
- Hennig, A. H. Studier öfver Bryozoerna i Sveriges Kritsystem.
   Cheilostomata. Act. Lund. xxviii, No. 11, 51 pp., 2 pls.
- Hincks, T. Contributions towards a general history of the Marine Polyzoa, 1890-91. Appendix [contd.]. Ann. N. H. ix, pp. 327-334.

Species described in the main work are still misquoted as "n. sp."

- The Polyzoa of the St. Lawrence: a study of Arctic forms.
   p. 149-157, 1 pl.
- 15. KOROTNEV, A. Po voprosu razvītiya pryesnovodnuikh mshanok.— Sur la question du développement des Bryozoaires d'eau douce. Zapiski Kiev. x, pp. 393-410, 2 pls.
- Kraepelin, K. Die Deutschen Süsswasser-Bryozoen. II. Entwickelungsgeschichtlicher Theil. Abh. Geb. Hamb. xii, No. 2, 67 pp., 5 pls.
- Namias, I. Su alcune forme Briozoarie del Mar Rosso. Atti Soc. Mod. xi, pp. 74-77.
- NEVIANI, A. Contribuzione alla conoscenza dei Briozoi fossili Italiani—Briozoi post-pliocenici del sottosuolo di Livorno. Boll. Soc. geol. Ital. x, pp. 99-148, 1 pl.
- Vine, G. R. Notes on *Polyzoa* found at Cleethorpes in 1875, 1879, and 1882. Naturalist, 1892, pp. 5-11.
- Notes on Polyzoa of the Zones of the Upper Chalk. Rep. Brit. Ass. 1891 (1892), pp. 656-659.
- WATERS, W. North Italian [fossil] Bryozoa. Pt. II. Cyclostomata.
   Q. J. Geol. Soc. xlviii, pp. 153-162, 1 pl.
- Observations on the Gland-like Bodies in the Bryozoa.
   J. L. S. xxiv, pp. 272-278, 1 pl. Abstr. in J. R. Micr. Soc. 1892, p. 777.

# II.—ANATOMY, &c.

Observations on the gland-like bodies in the Bryozoa; WATERS (22).—Nature of the excretory processes in Marine Bryozoa; HARMER (10).

Development of freshwater Bryozoa; KOROTNEV (15).—Development of German freshwater Bryozoa; KRAEPELIN (16).—DEMADE (6) concludes that the membrane of the statoblast in Alcyonella fungosa and

Cristat lla mucedo is composed of cells which undergo cellular differentiation, whilst the central mass is a tissue during all the periods of its development.

On the germ-layers in Bryozoan buds; BRAEM (4), DAVENPORT (5).

#### III.—DISTRIBUTION.

#### A.—GEOGRAPHICAL.

Lincolnshire: Bryozoa found at Cleethorpes; VINE (19).

Norway: list of Bryozoa dredged on the coast; HERDMANN, P. Liverp. Biol. Soc. vi, pp. 90 & 91.

Boulonnais: list of species; Hallez (7-9).

Somme: Bryozoa of the Department of; BIZET (3).

Red Sea: some species from; Namias (17).

New Jersey: Urnatella walkerii, n. sp. [?], and Octocella libertus, n. g. [?], n. sp. [?]; Helm (11).

List, with figg., of Bryozoa of the St. Lawrence; HINCKS (14).

#### B.—GEOLOGICAL.

Palsontological summary for 1890: Bryozoa, by G. F. Dollfus, Ann. Géol. univ. Paris, 1890 (1892), pp. 1039-1049.

Post-pliocene forms from the neighbourhood of Livorno, with 4 new species; NEVIANI (18).

Tertiary Cyclostomata of N. Italy, 22 species, including Diastopora brendolensis, Defrancia brendolensis, and Lichenopora (?) incrustans, new; WATERS (21).

Cretaceous: Bryozoa from the Upper Chalk; VINE (20).—Cretaceous Bryozoa of Sweden, with 13 new species; Hennig (12).—Bryozoa from the Upper Senonian of Irnich, Eifel; Vogel, Verh. Ver. Bheinl. xlix, pp. 93-99, fig.—Upper Senonian Bryozoa of the Eifel; Vogel, Verh. Ver. Rheinl. xlix, pp. 91-99.—Bryozoa from the Cretaceous of Schleswig-Holstein; Stolly, MT. min. Inst. Kiel, i, pp. 244 & 245.—1 species from the Rolling Downs Beds of Queensland; Jack & Etheridge, (ante, Moll. 193) pp. 441 & 442.

Jurassic species from Yorkshire (List); Fox-Strangways, Mem. Geol. Surv. U. K., Jurassic Rocks, ii, pp. 148 & 200,

Palæozoic: Check-list of Palæozoic Bryozoa from Wisconsin, Minnesota, Iowa, Dakota, and Nebraska; BIERBAUER (2).

Permian Bryozoa from the Duchy of Sachsen-Altenburg; GEINITZ, MT. Osterlande. v. p. 179.

Permo-Carboniferous species, Polypora? smithii, n. sp., from Queensland; JACK & ETHERIDGE, (ante, Moll., 193) pp. 216-225.

Devonian species from Manitoba, with *Pinacotrypa marginata*, n. sp.; WHITEAVES, Contr. Can. Pal. i, pp. 277-281, figg.

Ordovician species from the province of Quebec, with Dicranopora parva, Diplotrypa quebecensis, and Monotrypa incerta, n. spp.; AMI (1).

## IV.—SYSTEMATIC.

(†) is prefixed to fossil forms.

#### ENTOPROCTA.

- Urnatella walkerii, New Jersey; Helm, J. N. Y. Micr. Soc. viii, p. 47, fig., alleged new species; it appears to be a *Pedicellina*, but it is not stated whether it is furnished with a stolon or not.
- Rhabdopleura: cf. VERMES, sub HEMICHORDATA, where this and Cephalodiscus will in future be included with Balanoglossus.

#### ECTOPROCTA.

#### CYCLOSTOMATA.

- †Crisia destefanii, n. sp., Post-pliocene of Livorno; Neviani, Boll. Soc. geol. Ital. x, p. 135, fig.
- † Defrancea brendolensis, n. sp., Tertiary of N. Italy, WATERS, Q. J. Geol. Soc. xlviii, p. 161, fig.
- †Diustopora brendolensis, n. sp., Tertiary of N. Italy, WATERS, Q. J. Geol. Soc. xlviii, p. 155, fig.
- †Dicranoporu parva, n. sp., Ordovician of Quebec, AMI, Can. Rec. v, p. 99.
- †Liplotrypa quebecensis, n. sp., Ordovician of Quebec; AMI, Can. Rec. v, p. 100.
- † Entalophora d'anconæ, n. sp., Post-pliocene of Livorno, NEVIANI, Boll. Soc. geol. Ital. x, p. 143, fig.
- † Idmonea targionii, n. sp., Post-pliocene of Livorno, NEVIANI, Boll. Soc. geol. Ital. x, p. 139, fig.
- † Lichenopora (?) incrustans, n. sp., Tertiary of N. Italy, WATERS, Q. J. Geol. Soc. xlviii, p. 161, fig.
- †Monotrypa incerta, n. sp., Ordovician of Quebec, Ami, Can. Rec. v, p. 101.

#### CTENOSTOMATA.

Octocella, n. g. [?] for O. libertas, n. sp. [?], New Jersey; Helm, J. N. Y. Micr. Soc. viii, p. 48, fig. [May be one out of several genera of Ctenostomata, e.g., Cylindroecium?, Bowerbankia?—not properly defined.]

#### CHILOSTOMATA.

- †Cribrilina lævis, quadrisulcata, n. spp., Cretaceous of Sweden, HENNIG, Act. Lund. xxviii, No. 11, pp. 37 & 38, figg.
- Heteroæcium, n. g. for Membranipora amplectens, Hincks; Hincks, Ann. N. H. ix, p. 332.

- † Lepralia impressipora, n. sp., Cretaceous of Sweden, Hennig, Act. Lund. xxviii, No. 11, p. 43, fig.
- † Membraniporu excuvata, n. sp., Cretaceous of Sweden, Hennig, Act. Lund. xxviii, No. 11, p. 18, fig.
- † Membraniporella aurita, juvensis, n. spp., Cretaceous of Sweden, Hennig, Act. Lund. xxviii, No. 11, pp. 38 & 39, figg.
- † Monoporella inflata, n. sp., Cretaceous of Sweden, Hennig, Act. Lund. xxviii, No. 11, p. 40, fig.
- † Macronella hians, vesperugo, n. spp., Cretaceous of Sweden, Hennig, Act. Lund. xxviii, No. 11, pp. 44 & 45, figg.
- † Schizoporella carinata, n. sp., Cretaceous of Sweden, Hennig, Act. Lund. xxviii, No. 11, p. 42, fig.
- †Smittia adae, n. sp., Post-pliocene of Livorno, NEVIANI, Boll. Soc. geol. Ital. x, p. 127, fig.
- †Steganoporella rhomboidalis, n. sp., Cretaceous of Sweden, HENNIG, Act. Lund. xxviii, No. 11, p. 36, fig.
- † Periteichisma smitti, p. 25, ignabergense, p. 28, n. spp., Cretaceous of Sweden, Hennig, Act. Lund. xxviii, No. 11, figg.

#### PHYLACTOLÆMATA.

The statoblast has been studied in Alcyonella fungosa and Cristatella mucedo by DEMADE, La Cellule, viii, pp. 333-378, 2 pls. [For conclusions, see unte, Anatomy, &c.]

## [VERMIFORMIA.]

[Forms sometimes grouped under this heading will for the future be classed with VERMES (q.v.).]



# X. CRUSTACEA.

# (Including GIGANTOSTRACA AND PANTOPODA.\*)

By R. I. Pocock.

#### ARRANGEMENT OF RECORD:

CRUSTACEA, S. S.
Titles of Papers, p. 1.
Subject-Index, p. 13.
Systematic, p. 16.
PANTOPODA, p. 32.
GIGANTOSTRACA, p. 33.

# I.—TITLES OF PAPERS.

- Alcock, A. (1) On the Stridulating Apparatus of the Red Ocypode Crab. Ann. N. H. (6) x, p. 336.
- ALLEN, E. J. (1) Preliminary account of the Nephridia and Bodycavity of the Larva of *Palamonetes varians*. P. R. Soc. lii, pp. 338-342.
- —. (2) On the Minute Structure of the Gills of Palamonetes varians. Q. J. Micr. Sci. xxxiv, pp. 75-84, 1 pl.
- Ami, H. M. Additional Notes on the Geology and Palsontology of Ottawa and its Environs. Tr. Ottawa Nat. Club, vi, No. 5, p. 77. Ostracoda.

<sup>\*</sup> Since the GIGANTOSTRACA and PANTOPODA form part of the Marine fauna, and are largely, for this reason, studied systematically by students of Carcinology rather than of Arachnology, it has been considered advisable, on grounds of expediency, to transfer them to that section of the Zool. Rec. which deals with Crustacea.

- Andrews, E. A. Notes on the Fauna of Jamaica. Johns Hopk. Univ. Circ. xi.
  - Crustacea, p. 75.
- Aurivillius, C. W. S. (1) Analyse d'un mémoire intitulé du déguisement des Décapodes Oxyrhynques à l'aide d'adaptations singulières du corps. Ann. Sci. Nat. xiii, pp. 341-348.
- —. (2) Neue Cirripeden aus dem Atlantischen Indischen und Stillen Ocean. Œfv. Ak. Förh. 1892, No. 3, pp. 123-134.
- Barrois, T. Liste des Phyllopodes recueillis en Syrie. Rev. Biol. v, pp. 24-29.
- BATESON, W. Supernumerary Claws in Crabs. P. Z. S. 1892, p. 76.
- Beneden, P. J. van. (1) Le mâle de certains Caligidés et un nouveau genre de cette famille. Bull. Ac. Belg. xxiii, p. 220, &c.
- ——. (2) Quelques nouveaux Caligidés de la côte d'Afrique et de l'Archipel des Açores. Op. cit. xxiv, p. 241, &c.
- —. (3) Une nouvelle famille dans la tribu des Schizopodes. Bull. Ac. Belg. (3) xxii, pp. 444-459.
- BENEDICT, J. E. Decapod Crustacea of Kingston Harbour. Johns Hopk. Univ. Circ. xi, p. 77.
- —. (2) Corystoid Crabs of the Genera Telmessus and Erimacrus.
  P. U. S. Nat. Mus. xv, pp. 223-230, pls. xxv-xxvii.
- —... (3) Preliminary descriptions of Thirty-seven New Species of Hermit Crabs, of the genus Eupagurus. Op. cit. No. 887, pp. 1-26.
- Bergh, R. S. (1) Die Drehung des Keimstreifens und die Anlage des Dorsalorgans bei Gammarus pulex. Zool. Anz. xv, p. 268.
- Bernard, H. M. (1) The Apodida: a Morphological Study. Nature Series. London (Macmillan & Co.): 1892.
- —. (2) The Apodemes of Apus and the Endophragmal System of Astacus. Ann. N. H. (6) x, pp. 68-74, pl. v.
- Bertkau, P. Lebende Branchipus grubei. Verh. Ver. Rheinl. SB. xlviii, pp. 54 & 55.
- Birge, E. A. List of Crustacea Cladocera from Madison, Wisconsin. Tr. Wisconsin Acad. viii, p. 393, &c., 1 pl.
- BLANCHARD, R., & RICHARD, J. Faune de lacs salés d'Algérie. Cladocera et Copépodes. Mém. Soc. Z. Fr. iv, No. 5, pp. 512-535 (1891).
  Omitted from previous Record.
- BOLIVAR, —. Liste de las Crustacéos de España y Portugal del Museo de Madrid. Act. Soc. Esp. xxi, p. 124.
- Bordage, —. Myologie des Crustacés Décapodes. CR. Soc. Philom. No. 19, July, pp. 1 & 2 (1892).

- BOUVIER, E. L. (1) Sur le développement embryonnaire des Galathéides du genre Diptychus. C.R. oxiv, p. 767.
- ——. (2) Sur la graisse du foie des Crustacés Décapodes. Bull. Soc. Philom. (8) iii, pp. 170-174.
- ——. (3) Observations sur les mœurs des Pagures, faites au laboratoire maritime de Saint-Vaast-la-Houque pendant le mois d'Août, 1891; Op. cit. iv, pp. 1-9.
- ——. (4) Les Glaucothoés sont-elles des Larves de Pagures? Ann. Sci. Nat. xii, pp. 65–82.
- —... (5) Sur les Paguriens recueillis dans le Mer Rouge, par M. le Dr. Jousseaume. C.R. Soc. Philom. No. 8, p. 1, 1891–92. No new species.
- Brady, G. S. A Revision of the British Species of Freshwater Cyclopidæ and Calanidæ. Nat. Hist. Tr. North Durham, xi, pt. 1, p. 68.
- Bronn, —. Klassen und Ordnungen des Thier-Reichs. Arthropoda, v, pts. 32-34 [On the organization of the Decapod Crustacea], pp. 821-896.
- BROOKS, W. K., & HERRICK, F. H. Embryology and Metamorphosis. Johns Hopk. Univ. Circ. xi, pp. 65-71.
  - On Stenopus, Alpheus, and Gonodactylus.
- CALDÉRON Y ARANA, SALV. Notas Carcinologicas. Act. Soc. Esp. xx, pp. 108-110.
- CAMERA, C. Ricerche sui Copépodi liberi del Piemonte. Boll. Mus. Zool. Tor. vii, No. 120, pp. 1-13.

No new species.

- CAMERANO, L. Recherches sur la force absolue des Muscles des Crustacés Décapodes. Arch. Ital. Biol. xvii, pp. 212, &c.
- CANO, G. Sviluppo postembrionale dei Cancridi. Bull. Ent. Ital. xxiii, pp. 146-158.
- CANU, E. Les Copépodes du Boulonnaise, morphologie, embryologie, taxonomie. Lille (Danel): 1892, 354 pp.
- °CAPANNI, V. La Dafnia. Reggio Emilia, Tip. Artigianelli, 1892.
- CAR, L. Ein Beitrag zur Kenntniss der Copepoden von Triest. Glasnik Naravosl. Druzt. v, pts. 1-4, p. 105, pls. i-iii (1890).

No new species.

- CHEVREUX, E. (1) (Quatrième campagne de l'Hirondelle 1888) Hyule grimaldi et Stenothoe dollfusi, n. spp. Bull. Soc. Z. Fr. xvi, pp. 257-262.
- ——. (2) Vibilia erratica, Amphipode pélagique nouveau du littoral des Alpes Maritimes. Op. cit. xvii, p. 32.
- ---. (3) Sur le male adulte d'Hyperia schizogeneios, Stebbing. T. c. p. 233.
  - 1892. [VOL. XXIX.]

- CHEVREUX, E., & BOUVIER, E. L. (1) Voyage de la goëlette "Melita." Note préliminaire sur les Paguriens. Bull. Soc. Z. Fr. xvi, pp. 252, &c. (1891).
- ---- & ----. (2) Paguriens de la "Melita." Mém. Soc. Zool. v, pp. 8:-141, 3 pls.
- —— & ——. (3) Perrierella crassipes, espèce et genre nouveaux d'Amphipode des côtes de France. Bull. Soc. Z. Fr. xvii, p. 50.
- CHEVREUX, G., & GUERNE, J. DE. (1) Sur un espèce nouvelle de Gammarus du lac d'Annecy et sur les Amphipodes d'eau douce de la France, C.B. cxiv, p. 1286.
- & ——. (2) Description de Gammarus delebecquei, n. sp., du lac d'Amiens, suivie de quelques remarques sur les Amphipodes d'eau douce de la France. Bull. Soc. Z. Fr. xvii, p. 136.
- CHILTON, C. (1) Notes on some New Zealand Amphipoda and Isopoda.
  Tr. N. Z. Inst. xxiv, pp. 258-269.
- ——. (2) On a tubicolous Amphipod from Port Jackson. Rec. Austral. Mus. ii, No. 1, pp. 1–6.
- ---. (3) A new species of Munna from New Zealand. Ann. N. H. (6) x, p. 1, pls. i & ii.
- CHYZER, C. Ueber die Estherien Ungarns. Math. Nat. Ber. Ung. x, pp. 119-135.

No new species.

CLAUS, C. (1) Ueber die gattung Miracia, Dana, mit besonderer Berücksichtigung ihres Augen-Baues. Arb. z. Inst. Wien, ix, pp. 267-284.

[Omitted from Zool. Rec. xxviii.]

——. (2) Beiträge zur Kenntnis der Susswasser-Ostracoden. Op. cit. x, pp. 147-216 (1892).

Anatomical and Systematic.

CUÉNOT, L. Les organes Phagocytaires chez quelques invertébrés. Arch. Z. expér. p. ix.

On Astacus.

- CUNNINGHAM, J. T. Observations on the Larvæ of Palinurus vulgaris. Rep. Brit. Ass. 1891, p. 867.
- Dahl, F. Die Gattung Copilia (Saphirinella). Zool. Jahrb. vi, pp. 499-522, pl. xxiv.

No new species,

- Della Valle, A. Intorno agli organi di escrezioni di alcuni Gammarini. Bull. Soc. Nat. Napoli (1) iii, pl. i, pp. 269-272.
- DE MAN, J. G. (1) Decapoden des Indischen Archipels in Max Weber's Zool. Ergebnisse, &c. Vol. 11. Leiden (E. J. Brill): 1892.

- [Dz Man, J. G.] (2) Carcinological Studies in the Leyden Museum. Notes Leyd. Mus. xiv, pp. 225-264.
- Demoor, J. Recherches sur la marche des Crustacés (concluded). Arch. Z. exper. (2) ix, pp. 497-502.
- Dollfus, A. (1) Crustacés Isopodes terrestres du voyage de M. Ch. Alluaud dans le territoire d'Assinie. Ann. Soc. Ent. Fr. lxi, pp. 385-390, pl. vii.
- —. (2) Catalogue raisonné des Isopodes terrestres de l'Espagne. Au. Soc. Esp. (2) i, pp. 161-190.
- —. (3) Note sur les Isopodes terrestres et fluviatiles de Syrie. Rev. Biol. iv, pp. 1-15 (author's copy).
- ——. (4) Tableaux Synoptiques de la Faune Française. Le genre Armadillidium (Crustacés Isopodes terrestres). Feuill. Nat. xxii, No. 259, pp. 1-18.
- —... (5) Sur deux Isopodés terrestres d'Islande recueillis par M. Gaston Burdett. Bull. Soc. Z. Fr. xvii, p. 231.
- DUBRANT, H. Pond Life Studies. Cypris tristriata. Sci. Goss. 1892, pp. 267-269.
- EDWARDS, A. MILNE, & BOUVIER, E. L. (1) Observations préliminaires sur les Paguriens recueillis par les expéditions du Travailleur et du Talisman. Ann. Sci. Nat. xiii, pp. 185–226.
- & —. (2) Note sur un Pagure des grandes profondeurs de la mer (*Parapagurus pilosimanus*). Congr. Internat. Zool. 1892, pl. i, pp. 1-15.
- —— & ——. (3) Sur le développement embryonnaire des Galathéides abyssaux du genre Diptychus. C.B. 1891-92, No. 10, p. 1.
- ETHERIDGE, R. (1) On *Leaia mitchelli*, Eth., from the Upper Coal Measures of the Newcastle District. P. Linn. Soc. N.S.W. vii, pp. 307 & 308.
- —... (2) Notes on Queensland Cretaceous Crusticea. P. Linn. Soc. N.S.W. vii, pp. 305 & 306.
- —... (3) Hymenocaris salteri, McCoy. Rec. Geol. Surv. N.S.W. 1892, pt. 1, pp. 5-8, pl. iv.
- \*Forbes, S. A. On some Lake Superior Entomostraca. Rep. U. S. Fish. Comm. xv (1891), pp. 701-718 [Omitted from Zool. Rec. xxviii.]
  4 new species.
- Fréderice, L. Nouvelles recherches sur l'autotomie chez le Crabe. Arch. Biol. xii, pp. 169, &c.
- Garstane, J. W. (1) Notes on the Marine Invertebrate fauna of Plymouth for 1892. J. Mar. Biol. Ass. ii, No. 4. Crustacea, p. 337.

- [GARSTANG, J. W.] (2) On some New or Rare Marine Animals recently discovered on the Coast of Devonshire. Tr. Devon Ass. xxiv, Crustacea, pp. 384 & 385.
- GAVINO, C. Crustacei Brachiuri ed Anomuri raccolti nel viaggio della "Vettor Pisani" intorn al globo. Boll. Soc. Nat. Napoli (1) iii, pt. 1, pp. 79-105, & pt. 2, pp. 169-288.
- GIARD, A., & BONNIER, J. Sur le Cerataspis petiti, Guér., et sur la position systématique du genre Cerataspis, Gray. C.B. cxiv, p. 1029.
- GIESBRECHT, W. Systematik und Faunistik der Pelagischen Copepoden der Golfes von Neapel, &c. Berlin (B. Friedländer & Sohn): 1892.
- Gourret, C. Sur un variété de Nebalia bipes. Rec. Z. Suisse, v, p. 93.
- Geobben, C. (1) Ueber die Stammesverwandtschaft der Crustaceen. Verh. z.-b. Wien, xlii, pp. 11, &c.
- ---... (2) Zur Kenutniss der Stammbaumer und der Systems der Crustaceen. SB. Ak. Wien, 1892, pl. i, pp. 237, &c.
- GROOM, T. (1) On the Early Development of Cirripedia. P. R. Soc. lii, pp. 158-162.
- ---. (2) On the Orientation of Sacculina. P. Camb. Phil. Soc. vii, p. 160.
  - [Omitted from Zool. Rec. xxviii.]
- Gueene, Jules de. (1) Sur deux Phyllopodes nouveaux pour la faune des eaux-douces de Madagascar. Bull. Soc. Ent. Fr. 1892, p. l.
- ——. (2) Sur Cryptopus defrancii, Latr., Myside nouveau pour la faune der Açores. T. c. p. lvi.
- ——. (3) Un Ostracode nouveau pour la faune française : La distribution geographique de Cypris bispinosa, Luc. Rev. Biol. iv, pp. 518 & 519
- GUERNE, J. DE, & RICHARD, J. (1) Sur la faune des eaux douces de l'Islande. C.R. cxiv, p. 310.

#### Entomostraca.

- ——. (2) Cladocères et Copépodes d'eau douce des environs de Rufisque. Mém. Soc. Zool. v, pp. 526-538.
- Häcker, V. (1) Ueber specifische Variation bei Arthropoden im Besonderen über die Schutzanpassungen bei Krabben. Ber. Ges. Freiburg, vi, pt. 3, pp. 90-100.
- ——. (2) Die Eibildung bei Cyclops und Canthocamptus. Zool. Jahrb. v, pp. 211–248, 1 pl.
- HANSEN, H. J. Rhizorhina ampelisca, n. g. & sp. En ny til Herpyllobiida, n. fam. hövende Copepod, snyltende paa Ampelisca lavigata, Lilljb. Ent. Med. 1892, pp. 207-234, pl. iii.

- Heim, F. (1) Études sur le sang des Crustacés Décapodes suivies d'un Essai sur le role dès pigments. Ann. Soc. Ent. Fr. lxi, pp. 155-270.
- ——. (2) Sur les pigments des œufs de Crustacés. C.R. Soc. Biol. 1892, p. 467.
- —... (3) Sur la matière colorante bleue du Sang des Crustacès. C.R. exiv, p. 771.
- ---. (4) Les œufs des Crustacés contenant des ferments actifs. Bull. Soc. Ent. Fr. 1891, pp. exvi.
  - [Omitted from Zool. Rec. xxviii.]
- (5) Etudes sur le sang der Crustacés Décapodes. Recherches sur les Diptero-carpacées. Paris (Chamerot & Renouard): 1892.
- HENNEGUY, F., & THÉLOHAN, P. Sur un Sporozoaire parasite des muscles des Crustacés Décapodes. C.R. Soc. Biol. 1892, p. 585, and C.R. cxiv, p. 1552; Trans. Ann. N. H. (6) x, p. 342.
- HILGENDORF, F. (1) Eine neue ostafrikanische Susswasser-Krabbe (Telphusa emini). SB. nat. Fr. 1892, No. 1, pp. 11-13.
- ----. (2) Eine neue Brachynotus-Art aus Aden (Br. harpax). Op. cit. No. 4, pp. 37-40.
- IDE, M. (1) Glandes cutanées à canaux intracellulaires chez les Crustacés Edriophthalmes. La Cellule, vii, pp. 345-372. Abstr. in J. R. Micr. Sci. 1892, pt. 3, p. 364.
- ——. (2) Le tube digestif des Edriophthalmes. La Cellule, viii, pp. 97-204.
- Імног, О. Е. Die Zusammensetzung der pelagischen Fauna der Süsswasserbecken. Biol. Centralbl. xii, pp. 171, &c. Crustacea, pp. 178-182.
- ISHIKAWA, C. Studies on Reproduction Elements. 1. Spermatogenesis, Ovogenesis, and Fertilization in *Diaptomus*. J. Coll. Sci. Japan, v, pp. 1-34.
- IVES, J. E. . . . . Crustaceans collected by the West Greenland Expedition of 1891. P. Ac. Philad. 1891, pl. iii, p. 481.
  [Omitted from Zool. Rec. xxviii.] No new species.
- JOLYET, E., & VIALLANES, H. Recherches physiologiques sur le système nerveux accélérateur et modérateur du cœur chez le Crabe. Ann. Sci. Nat. xiv, pp. 387-404.
- JONES, T. RUPERT, & WOODWARD, HENRY. A Monograph of the British Palsozoic Phyllopoda (Phyllocarida, Packard). Pt. ii. Some bivalved and univalved species. Palsontographical Society for 1892, pp. 73-124.
- JONES, T. RUPERT, & KIRKBY, J. W. On Carboniferous Ostracoda from Mongolia. Ann. N. H. (6) ix, pp. 302-307, pl. xvi.

  No new species.

- JOURDAIN, S. Sur quelques points de l'embryogénie de l'Oniscus murarius, Cuv., et du Porcellio scuber, Leach. C.R. exiv, p. 428.
- KAFKA, J. Die Fauna des Böhmischen Teiche. Arch. naturw. Landesforsch. Böhmen, viii, No. 2, pp. 1-115.
  Crustacea, pp. 110-112.
- KAUFMAN, A. Ueber die Gattung Acanthopus und eine neue Susswassercytheride. Zool. Anz. xv, pp. 393-395.
- KERHERVÉ, L. B. De l'apparition provoquée des éphippies chez les Daphnies (Daphnia magna). Mém. Soc. Zool. v, pp. 227-236.
- KLOCKE, E. Ein neuer Pleuroxus. Zool. Anz. xv, pp. 188 & 189.
- Kochs, W. Versuche über die Künstliche vermehrung Kleiner Crustaceen. Biol. Centralbl. xii, p. 599.
- KŒHLER, M. Recherches sur la cavité générale et sur l'appareil excréteur des Cirripedes. C.R. cxiv, p. 1214.
- Kelbel, K. (1) Ein neuer ostasiatisches Flusskrebs. Anz. Ak. Wien, xviii, pp. 176 & 177.
- (2) Beiträge zur Kenntniss der Crustaceen der Canarischen Inseln.
   Ann. Hofmuseum Wien, vii, No. 3, pp. 105-116.
   2 new species.
- Kojevnikov, G. La faune de la mer Baltique orientale et les problèmes des explorations prochaines de cette faune. Congr. Internat. Zool. 1892, pt. 1, p. 132.
- Kowalevsky, A. Ein Beitrag zur Kenntniss des Excretions-organe des Pantopoden. Mem. Ac. Pétersb. xxxviii, No. 12, pp. 1-8, 1 pl.
- Krause, Aurel. Neue Ostrakoden aus Märkischen Silurgeschieben. Z. geol. Ges. 1892, pp. 383-399, pls. xxi & xxii.
- LANDÉ, A. Quelques remarques sur les Cyclopides. Mém. Soc. Zool. v, pp. 156-173.
- LANG, ARNOLD. Lehrbuch der Vergleichenden Anatomie, &c. Jena (Fischer): 1892.
- LEIENKLAUS, E. Ostracoden des nordwestdeutschen Tertiärs. CB. Ver. Rheinl. 1892, pt. 1, pp. 58 & 59.

No new species.

- LEBEDINSKI, J. [On the Development of Eriphia.] Zapiski. Novoross. Obsch. Estestv. xvii, pp. 137-256.
  - Printed entirely in Russian.
- McMurrich, J. P. The Formation of the Germ-layers in the Isopod Crustacea. Zool. Anz. xv, p. 271.
- MARCHAL, P. (1) Note préliminaire sur l'appareil excréteur des Pagurides et des Palinurides. Bull. Soc. Z. Fr. xvi, p. 57 (1891).
- —... (2) Sur un Pagure habitant une coquille sénestre (Neptunea contraria, Chem.). T. c. p. 267.

- [MARCHAL, P.] (3) Recherches anatomiques et physiologiques sur l'appareil excréteur des Crustacés Décapodes. Arch. Z. expér. (2) x, pp. 57-275, 9 pls.
- MARSH, C. D. On the Deep Water Crustacea of Green Lake. Tr. Wisconsin Acad. viii, pp. 211-213.

No new species.

- MAUPAS, M. Sur le *Belisarius viguieri*, nouveau Copépode d'eau douce. C.R. cxv, p. 132.
- Moniez, R. Faune des lacs salés d'Algérié (Ostracodes). Mém. Soc. Zool. iv, pp. 246-257.

[Omitted from Zool. Rec. xxviii.] 2 new species.

Nathorst, —. Om några mollusker och Ostrakoder från grartara sötvattensflagringar i Ryssland och Tyskland. Œfv. Ak. Förh. 1892, p. 425.

No new species.

- NORMAN, A. M. (1) British Schizopoda of the Families Lophogastridæ and Euphausidæ. Ann. N. H. (6) ix, pp. 454-464.

  No new species.
- —. (2) On British Mysidæ, a family of Crustacea Schizopoda. Op. cit. x, pp. 143-166 & 242-263.

2 new species and 1 new genus.

- ---. (3) Rare Crustacea on the Yorkshire Coast. Naturalist, 1892, pp. 175 & 176.
  - Occurrence of Thysanoessa longicaudata and Nematoscelis megalops.
- NUSBAUM, J. Zur Kenntniss der Wurmerfauna und Crustaceenfauna Polens. Biol. Centralbl. xii, pp. 54, &c. Cyclops, p. 57. No new species.
- ORTMANN, A. (1) Die Decapoden-Krebse des Strassburger Museums. No. III. Die Abtheilungen der Reptantia, Boas: Homaridæ, Loricata, and Thalassinidea. Zool. Jahrb. Syst. vi, pp. 1-57.
- —... (2) No. 1v. Die Abtheilungen Galatheidea und Paguridea. T. c. pp. 241-326.
- ——. (3) Die Decapoden-Krebse des Strassburger Museums. No. v. Die Abtheilungen Hippidea, Dromidea, und Oxystomata. T. c. pp. 532-588, 1 pl.
- Osorio, B. (1) Note sur quelques espèces de Crustacés de l'île St. Thomé, ilet des Rolas et Angola. J. Sci. Lisb. (2) ii, pp. 140 & 141. No new species.
- Carriero de la contribuição para a fauna carcinologica da ilha de St. Thomé. T. c. pp. 199-201.

1 new species.

- OSTROUMOF?, A. Note sur la distribution de Balanus eburneus, Gou'd. Zool. Anz. p. 160.
- PARKER, W. N. Note on abnormalities in the Crayfish. Ann. N. H. (6) ix, pp. 181 & 182.
- POPPE, S. A., & RICHARD, J. Description du Diaptomus schmakeri, n. sp., recueilli par M. Schmacker dans le lac Tahoo (China). Bull. Soc. Z. Fr. xvii, p. 149.
- POUCHET, G. Contribution à l'histoire des Cyamus. J. de l'Anat. Phys. 1892, pp. 99-108.
- RANCE, C. E. DE. Note on the discovery of Estheria minuta (var. Brodeiana) in the New Red Sandstone, Rep. Brit. Ass. 1891, p. 644.
- RATH, O. VOM. Ueber die von C. Claus beschriebene Nervenendigung in den Sinneshaaren der Crustaceen. Zool. Anz. xv, p. 96.
- RATHBUN, MARY J. Catalogue of the Crabs of the family Periceridæ, in the U. S. National Museum. P. U. S. Nat. Mus. xv, pp. 231-277, pls. xxviii-xl.
- RICHARD, J. (1) Sur l'œil latéral des Copépodes du genre Pleuromma. Zool. Anz. xv, p. 400.
- —... (2) Sur la présence d'un Cysticercoid chez un Calanide d'eau douce. Bull. Soc. Z. Fr. xvii, p. 17.
- —. (3) Sur l'identité des genres Ilyopsyllus, Brady & Robertson, et Abacola, Edwards; description de I. jousseaumei, n. sp. T. c. p. 69.
- —. (4) Sur quelques Entomostracés de l'île d'Elbe el de l'île de Monte Cristo. T. c. p. 225.
  - No new species.
- —. (5) Sur la distribution géographique des Cladocera. Congr. Internat. Zool. 1892, pt. 1 (second half), pp. 9-23.
- —. (6) Grimaldina brazzai, Guernella raphaelis, Monodaphnia mocquerysi, Cladocères nouveaux du Congo. Mém. Soc. Zool. v, pp. 213-226.
- —. (7) Recherches sur le système glandulaire et sur le système nerveux des Copépodes libres d'eau douce suivies d'une revision des espèces de ce groupe qui vivent en France. Ann. Sci. Nat. xii, pp. 113-260.
  - No new species.
- RISTORI, G. (1) Note di Carcinologie pliocenica. Atti Soc. Tosc. Proc. Verb. viii, pp. 86-89.
- •——. (2) Resti di Crostacei nel pliocene dell' Isola di Pianosa. T. c. p. 90.

- Robertson, D. (1) A second contribution towards a catalogue of the *Amphipodu* and *Isopoda* of the Firth of Clyde and West of Scotland. P. N. H. Soc. Glasg. iii, pt. 3, pp. 199-223.
- —. (2) On some recent Marine Ostraceda, dredged in Granton Harbour. T. c. pt. 2, pp. 196-198.
  No new species.
- ROBINSON, MARGARET. On the Nauplius-Eye persisting in some Decapoda. Q. J. Micr. Sci. xxxiii, p. 283, 1 pl.
- SARS, G. O. An Account of the Crustacea of Norway, with short descriptions and figures of all the Species. Vol. 1. Amphipoda; parts 10-15. Families Amphilochida, Stenothoida, Leucothoida, and Œdicerida.
- SAUVAGE, H. E. Note sur les Crustacés Jurassiques supérieurs du Boulonnais. Ann. Sci. Nat. xii, 83 pp.
  6 new species of Decapoda.
- SCHALFÉEF, P. Carcinologische Bemerkungen aus dem Zoologischen Museum der Kais. Akademie der Wissenschaften. Bull. Pétersb. xxxv, pp. 331-342.
- SCHMEIL, OTTO. Deutschlands freilebende Süsswasser Copepoden: pt. 1, Cyclopidæ. Bibl. Zool. xi, 8 pls.

  No new species.
- Schneider, A. Sur l'existence d'artères laterales chez les Amphipodes. Tabl. Zool. ii, p. 213, pl. xxxv.
- Scott, T. (1) On some new and rare Crustacea from the Firth of Forth. Ann. N. H. (6) x, p. 201, pls. xv & xvi.
  - 3 species recorded, of which one, a Lichomolgus, is new.
- ----. (2) Notes on Rhincalanus gigas, Brady, and Ectinosoma atlanticum, Brady & Robertson. P. N. H. Soc. Glasg. iii, p. 264.
- —. (3) The Land and Freshwater Crustucea of the district around Edinburgh. P. B. Soc. Edinb. 1892, pp. 74-81.
- —... (4) Is Amphithopsis latipes a commensal? Ann. Scott. Nat. Hist. Soc. 1892, p. 141.
- —. (5) Diaptomus castor in the Braid Ponds, near Edinburgh. T. c. p. 202.
- —. (6) Modiolicola insignis in the Firth of Forth. T. c. p. 272.
- ——. (7) Iliocryptus sordidus in the Loch End Loch, Edinburgh. T. c. p. 273.
- —. (8) Argissa (Syrrhoe) hamatipes, Norman, in the Firth of Forth. T. c. p. 272.
- ——. (9) Notes on some Freshwater, Brackishwater, and Marine Entomostraca, new to the Fauna of Orkney. P. N. H. Soc. Glasg. iii, pt. 2, pp. 91-100.

- Scott, Andrew. Notes on Cecrops latreillei, Leach, and Læmargus muricatus, Kr. T. c. p. 266.
- Scott, T. & A. New and rare Crustacea from the East Coast of Scotland. Ann. Scott. Nat. Hist. Soc. 1892, No. 3, pp. 149-156.
- Scourfield, D. J. On British Cladocera. J. Quek. Club (2) v, p. 63.
- SHARP, B. On Hippa emerita. P. Ac. Philad. 1892, pp. 327 & 328.
- Sowinsky, B. Matériaux pour servir à l'étude des Crustacés d'eau douce dans la partie sud-occidentale de la Russie. Zapiski. Kiev, xi, pp. 107-171.
  - Printed in Russian.
- St. HILAIRE, C. DE. Sur la résorption chez l'écrevisse. Bull. Ac. Belg. xxiv, pp. 506, &c.
- STUHLMANN, F. Zur Kenntniss der Fauna central-africanischer Seen. Zool. Jahrb. vi, pp. 152-154.
- THOMSON, G. M. On the Occurrence of Two Species of Cumacea in New Zealand. J. L. S. xxiv, p. 263.
- ULRICH, E. O. New Lower Silurian Ostracoda. No. 1. Am. Geol. x, No. 5, pp. 263-270, pl. x.
- Viallanes, M. H. (1) Recherches anatomiques et physiologiques sur l'œil composé des Arthropodes. 1. La morphologie de l'œil de la Langouste. Ann. Sci. Nat. xiii, pp. 349-384, 2 pls.
- —. (2) Contributions à l'histologie du système nerveux des Invertébrés. La lame ganglionnaire de la Langouste. T. c. pp. 385-398.
- VIGUIER, C. L'héliotropisme des Nauplius. C.B. cxiv, p. 1489.
- Voigt, W. Synapticola teres, n. g. & sp., ein parasitischen Copépode aus Synapta kefersteinii. Z. wiss. Zool. liii, pp. 31-42.
- WAGNER, C. Études sur le développement des Amphipodes : cinquième partie, Développement de la *Melita palmata*. Bull. Mosc. 1891 (n.s.) v, p. 401, pls. ix & x.
- WALKER, A. O. (1) Note on some Habits of Crustacea. Tr. Biol. Soc. Liverpool, iv, p. 84 (1890).
- —... (2) Report on the Higher Crustacea collected [in Liverpool Bay] during 1889. T. c. p. 239.
- ---... (3) Revision of the *Podopthalmuta* and *Cumacca* of Liverpool Bay. T. c. pp. 26-104.

  No new species.
- —. (4) The Lysianassida of the British Sessile-eyed Crustacea, Bate & Westwood. Ann. N. H. (6) ix, pp. 134-138.
- Weber, Max. Die Süsswasser-Crustaceen des Indischen Archipels, nebst bemerkungen über die Susswasser-Fauna im Allgemeinen. In Max Weber's Zool. Ergebnisse, &c. Vol. 11. Leiden: E. J. Brill.

- Weldon, W. F. R. (1) Certain Correlated Variations in Crangon rulgaris. P. R. Soc. li, pp. 2-21.
- —. (2) The Formation of Germ-layers in Crangon vulgaris. Q. J. Micr. Sci. xxxiii, pp. 343-363, 3 pls.
- Weltner, W. Nachtrage zur Fauna von Helgoland. III. Zur Cirripedien fauna von Helgoland. Zool. Jahrb. vi, pp. 453-455.
- WHITEAVES, J. F. Description of a New Genus and Species of Phyllocarid Crustacea from the Middle Cambrian of Mount Stephen. Can. Rec. v, No. 4, pp. 205-208.
- WOOD-MASON, J., & ALCOCK, A. (1) Natural History Notes from H.M. Marine Steamer "Investigator," &c. Ann. N. H. (6) ix, pp. 265-275, 2 pls.
- & —. (2) Natural History Notes, &c. T. c. pp. 358-370.
- WOODWARD, H. Notes on a New Decapodous Crustacean, *Prosopon etheridgei*, H. Woodw., from the Cretaceous of Queensland. P. Linn. Soc. N.S.W. vii, pp. 301-304, pl. iv.
- ZSCHOKKE, F. Dei Fortpflanzungsthätigkeit der Cladoceren der Hochgebirgseen. Festschr. z. 70. Geburts. R. Leuckart. pp. 396-404.

# II.—SUBJECT INDEX TO PAPERS ON ANATOMY, PHYSIOLOGY, EMBRYOLOGY, DISTRIBUTION, &c.

#### 1. EMBRYOLOGY and METAMORPHOSIS.

Pigment of the eggs in *Decupoda*; Heim (2).—Active ferments in eggs of *Crustacea*; Heim (4).—Genesis of the sperm and ova and fertilization in *Diaptomus*; ISHIKAWA.—Structure of ova in *Cyclops* and *Canthocamptus*; HÄCKER.

Reproductive activity of Cladocera; ZSCHOKKE.

Parthenogenesis in Daphnia; KERHERVÉ.

Artificial reproduction of small Crustacea; Kochs.

Development of Eriphia; Lebedinski.—Embryology and Metamorphoses of Stenopus, Alpheus, and Gonodactylus; Brooks & Herrick. Development of embryo of Diptychus (Galutheidæ); Bouvier (1), and Edwards & Bouvier.—Formation of germ layers in Crangon vulgaris; Weldon.—Development and torsion of the germinal streak in Gammarus; Bergii (1 & 2).—Germ-layers in Isopoda; McMurricii.—Embryology of Oniscidæ; Jourdain.—Early development of Cirripedia; Groom.—Postembryonal development of Crabs; Cano.—Development of Amphipoda; Wagner.—On the larvæ of Palinurus; Cunningham.

#### 2. CIRCULATORY SYSTEM.

Blood of Decapoda; Heim (1, 3, 5). — Arteries of Amphipoda; Schneider.

#### X. CRUSTACEA.

### 3. MUSCULAR SYSTEM.

Muscles of Decapoda; Bordage.—Strength of muscles in Decapoda; Camerano.

# 4. NUTRITION, EXCRETORY ORGANS, &c.

Digestion of the Crayfish; St. HILAIRE.—On the fat of the liver in Decapoda; BOUVIER.—On the Digestive Tract of Edriophthalmia; IDE.

Excretory Organs: On the body-cavity and excretory organs of Cirripedia; Kehler.—On the body-cavity and Nephridia of the larva of Palæmonetes; Allen.—Excretory organs of Gammarus; Della Valle.—Excretory organs of Paguridæ and Decapoda; Marchal (1, 3).

# 5. NERVOUS SYSTEM AND ORGANS OF SENSE.

On the nervous system of the heart in the Crab; JOLYET & VIALLANES.

—Nervous terminations of the sensory hairs; RATH.—Nervous and glandular systems of freshwater Copepoda; RICHARD.—Nervous system of Palinurus; VIALLANES.

Lateral eyes of Copepoda; RICHARD.—Structure of eyes of Miracia; CLAUS.—Nauplius-eye of Decapods; ROBINSON.—Eye of Palinurus; VIALLANES.

#### 6. Anatomy and Miscellaneous Subjects.

Phylogeny and classification of Crustacea; BERNARD, GROBBEN (1, 2). General comparative anatomy; LANG.—Organization of Decapoda; BRONN.—Internal skeleton of Apus and Astacus; BERNARD.—Morphology of freshwater Ostracoda; CLAUS. — Cutaneous glands of Edriophthalmia; IDE.—Structure of branchise in Palæmonetes; ALLEN.

Abnormalities of Crayfish; PARKER.—Additional claws in Crabs; BATESON.

Variations in Crangon vulgaris; WELDON.

Orientation of Sacculina; GROOM (2). — Heliotropism of Nauplius; VIGUIER.

On the presence of Cysticercoid in freshwater Calanida; RICHARD. Phagocytes of Astacus; Curnot.

Parasites (Sporozoa) in the muscles of Decapoda; Henneguy & Thélohan.

Habits, Instincts, &c.: Movements of Crustacea; Demoor.—Decorative instinct of Oxyrhyncha; Aurivillius (1).—Protective adaptation in Crabs; Häcker.—Autotomy of Crabs; Frédericq.—Pagurus in a sinistral shell; Marchal.—Habits of Gelasimus; Alcock (2): Ocypoda; Alcock (1): of Pagurus; Bouvier (3): of Hippa emerita; Sharp: of Oniscus, &c.; Walker (1).—History of Cyamus; Poucher.—Natural History of Daphnia; Capanni.—Remarks on the Cyclopida; Landé.—Living Branchipus; Bertkau.

#### 7. PALEONTOLOGY.

Paleozoic Phyllopoda; JONES & WOODWARD. Middle Cambrian: Phyllopoda; WHITEAVES.

Lower Silurian: Ostracoda; ULRICH.—Silurian: Ostracoda; KRAUSE.

Carboniferous Ostracoda from Mongolia; Jones & Kirkby. Upper Jurassic Crustacea of the Boulonnais: Sauvage.

Pliocene Crustucea; RISTORI (1 & 2). Tertiary Ostracoda; LEIENKLAUS.

### 8. FAUNISTIC.

Pelagic elements in the fauna of freshwater lakes; IMHOF.

Observations on marine and freshwater Crustaceans; MAX Weber.

# a. Holarctic Region.

Britain: Mysidæ; NORMAN (1,2): freshwater Cyclopidæ and Calanidæ; BRADY: Cladocera; Scourfield.—Edinburgh and the Firth of Forth: Scott (1,3,5-8).—Granton: Ostracoda; Robertson (2).—Firth of Clyde: Amphipoda and Isopoda; Robertson (1).—Orkney: Entomostraca; Scott (9).—Yorkshire: rare Crustacea; NORMAN (3).—Liverpool Bay: Podophthalmata and Cumacea; Walker (2, 3).—Plymouth and Coast of Devonshire: Crustacea; Garstang (1 & 2).

Iceland: Isopoda; Dollfus (5): freshwater forms; Guerne & Richard.

Norway: Amphipoda; SARS.

Heligoland: Cirripedia; WELTNER.

Eastern Baltic: KOJEVNIKOV.

Germany: free-living, freshwater Copepoda; SCHMEIL.

France: freshwater Copepoda; BICHARD (7): Ostracoda; GUERNE (3): Isopoda; DOLLFUS (4): Amphipoda; CHEVREUX & BOUVIER (3), CHEVREUX & GUERNE (1, 2).

Hungary: Ostracoda; CHYZER.—Bohemia: pond-life; KAFKA.

Poland: NUSBAUM.

Russia: Ostracoda; Nathorst.—S. W. of Russia: freshwater Crustacea; Sowinsky.—Black Sea; Ostroumoff.

Mediterranean, &c.: Payuridæ; Edwards & Bouvier.—Island of Elba and Monte Christo: Entomostraca; Richard (4).—Triest: Copepoda; Car.—Gulf of Naples: Copepoda; Giesbrecht.

Spain: Isopoda; Dollfus (2).—Spain and Portugal; higher Crustacea; BOLIVAB.

Algeria: Ostracoda; Moniez: Cladocera and Copepoda; Blanchard & Richard.

Syria: Phyllopoda; BARROIS: Isopoda; DOLLFUS (3).

Azores: Oryptopus recorded; Guerne (2):

#### X. CRUSTACEA.

Canary Islands: KŒLBEL.

Senegal: Entomostraca; Guerne & Richard (2).

Japan: ORTMANN (1-3). West Greenland: IVES.

Wisconsin: Cladocera; BIRGE: Deep-water Crustacea of Green Lake;

MARSH.

# b. Ethiopian Region.

African Lakes; STUHLMANN.

St. Thomas (Gulf of Guinea) and Angola: Osobio.—Assinie (W. Africa): Isopoda; Dollfus (1).

Red Sea: Paguridæ; BOUVIER (5). Madagascar: Phyllopoda; GUERNE (1).

# c. Indo-Pacific Region.

Indian Seas: WOOD-MASON & ALCOCK.—Indian Archipelago: Freshwater forms; MAX WEBER.—Indo-Malaya and Austro-Malaya: Decapoda; DE MAN.—New Zealand: Cumacea; Thomson.

# d. Neotropical Region.

Jamaica: higher Crusticea; Andrews, Benedict (1). Chili: freshwater Entomostraca; RICHARD (8).

# III.—SYSTEMATIC.

# MALACOSTRACA.

# PODOPHTHALMA.

#### DECAPODA.

#### BRACHYURA.

#### INACHIDE.

Simocarcinus pusillus, Asaba, pp. 55 & 174, figg. 3 & 4, GAVINO, Bull. Soc. Nat. Napoli. (1) iii, n. sp.

#### Periceride.

The family characterized and divided into the three subfamilies—Pericerina, Othonima, and Mithracina; all the genera and all the species known to the author characterized; RATHBUN.

Libinia emarginata, Leach, p. 235, pl. xxxi, fig. 2, dubia (M.-Edw.), p. 237, fig. 1, N. America, setosa, Lock. Lower California, p. 238, pl. xxviii; RATHBUN, P. U. S. Nat. Mus. xv.

- L. macdonaldi, Gulf of California, p. 238, pl. xxix, spinimana, Carolina, &c., p. 240, pl. xxx, mexicana, Gulf of California, p. 242, pl. xxxi, fig. 3, RATHBUN, P. U. S. Nat. Mus. xv, n. spp.
- Pericera cornuta, var. celata, M.-Edw., Florida, &c., p. 244; RATHBUN, P. U. S. Nat. Mus. xv.
- P. triangulata, Gulf of California, p. 246, pl. xxxii, fig. 1, atlantica, Gulf of Mexico, p. 247, contigua, Gulf of California, p. 247, pl. xxxii, fig. 2, RATHBUN, P. U. S. Nat. Mus. xv, n. spp.
- Macrocaloma camptocera, Stmp., Florida, p. 249, pl. xxxiii, fig. 2, eutheca, Stmp., Florida, p. 251; RATHBUN, P. U. S. Nat. Mus. xv.
- M. tenuirostris, Jamaica, p. 252, pl. xxxiii, fig. 1, RATHBUN, P. U. S. Nat. Mus. xv, n. sp.
- Podohuenia, n. g. for P. erythrea, n. sp., Massaua; GAVINO, Boll. Soc. Nat. Napoli (1) iii, p. 180, fig. 5.
- Tiurinia depressa, Stmp., Ousima, p. 276, pl. xl, fig. 1, spinigera, Stmp., Ousima, p. 277, pl. xl, fig. 2; RATHBUN, P. U. S. Nat. Mus. xv.
- Othonia aculeata, Gibbes, Florida, p. 255, pl. xxxiv, figg. 1 & 2, cherminieri, Schr., W. Indies, &c., p. 255, pl. xxxiv, figg. 3 & 4; RATHBUN, P. U. S. Nat. Mus. xv.
- O. carolinensis, Charleston, p. 256, pl. xxxv, figg. 1 & 2, nicholsi, California, p. 257, pl. xxxv, fig. 3, rotunda, Florida, p. 258, pl. xxxvi, fig. 1, RATHBUN, P. U. S. Nat. Mus. xv, n. spp.
- Mithrax spinipes, Bell, p. 259, acuticornis, Stmp., Gulf of Mexico, p. 260, pl. xxxvii, fig. 1, forceps, M.-Edw., N. America, Brazil, &c., p. 269; RATHBUN, P. U. S. Nat. Mus. xv.
- M. pilosus, Bahamas, p. 262, pl. xxxix, hemphilli, Florida, p. 263, pl. xxxvii, fig. 2, sinensis, California, p. 266, pl. xxxviii, fig. 2, bahamensis, p. 267, pl. xxxviii, fig. 1, braziliensis, Brazil, p. 268, pl. xxxvii, fig. 2, RATHBUN, P. U. S. Nat. Mus. xv; M. trigonopus, Panama, p. 183, GAVINO, Boll. Soc. Nat. Napoli (1) iii : n. spp.
- Mithraculus tumidus, Payta, p. 186, GAVINO, Boll. Soc. Nat. Napoli (1) iii, n. sp.

#### CORYSTIDÆ.

- Telmessus cheiragonus, Til., syn. serratus, White, &c., Oregon, &c., p. 224, pls. xxv & xxvi, figg. 2-4, acutidens, Stmp., Japan, p. 228, pl. xxvi, fig. 1; Benedict, P. U. S. Nat. Mus. xv.
- Erimacrus, n. g., p. 229, for Platycorystes isenbeckii, Brandt, Unalaska, &c., p. 229, pl. xxvi, figg. 5-7, & pl. xxvii; Benedict, P. U. S. Nat. Mus. xv.

# CANCRIDE and TELPHUSIDE.

- Euryetisus, n. g., p. 88, for E. deplanatus, n. sp., Singapore, pp. 81 & 200; Gavino, Boll. Soc. Nat. Napoli (1) iii.
- Actumnus targionii, Pernambuco, pp. 89 & 209, fig. 13; Gavino, Boll. Soc. Nat. Napoli (1) iii, n. sp.

- Eucratoplax spinidentata, Jamaica, Benedict, Johns Hopk. Univ. Circ. xi, p. 77, n. sp.
- Pilumnus tridentatus, Maitl., is referable to the genus Heteropanope, the species figured and described, p. 228, pl. vii, fig. 1; DE MAN, Notes Leyd. Mus. xiv.
- Pilumnus sluiteri, Batavia, p. 283, pl. 1, fig. 2, DE MAN, in Max Weber's Zool. Ergeb. ii, n. sp.
- Telphusa eminii, n. sp., E. Africa, pp. 11-13, HILGENDORF (1). T. granulata, n. sp., Tjibodas, p. 290, pl. ii, fig. 5; larnaudi, n. var. brevimarginata, Sumatra and Java, p. 294, pl. iv, fig. 6; celebensis, n. sp., with 2 new vars., lokaensis and pareparensis, Celebes, p. 297, pls. xvii & xviii, fig. 17; De Man, in Max Weber's Zool. Ergeb. ii.
- Geotelphusa picta, von Martens, Java, p. 234, pl. viii, fig. 2; transversa, von Mart., syn. crassa, M.-Edw., Fiji I., p. 241; DE MAN, Notes Loyd. Mus. xiv.
- G. loxophthalma, Borneo, p. 245, pl. vii, fig. 3, DE MAN, Notes Leyd. Mus. xiv, n. sp.

# GRAPSIDÆ.

- Brachynotus harpax, Aden, HILGENDORF, pp. 37-40, SB. nat. Fr. 1892, n. sp.
- Areograpsus jamaicensis, Jamaica, Benedict, Johns Hopk. Univ. Circ. xi, p. 77, n. sp.
- Glyptograpsus spinipes, Isola delle Perli, Gavino, pp. 92 & 241, fig. 15, Boll. Soc. Nat. Napoli (1) iii, n. sp.
- Dotilla wickmanni, Celebes, p. 308, pl. xviii, fig. 8, De Man, in Max Weber's Zool. Ergeb. ii, n. sp.
- Metaplax crassipes, n. sp., Makassar, p. 325, pl. xix, fig. 12, DE MAN, in Max Weber's Zool. Ergeb. ii.
- Ptychognathus dentatus, n. sp., Celebos, p. 318, pl. xviii, fig. 9; riedelii, n. var. pilosa, Flores, p. 323; De Man, in Max Weber's Zool. Ergeb. ii.
- Sesarma eydouxi, M.-Edw., p. 248, recta, Rand., Surinam, p. 249, pl. x, fig. 4; angusticeps, Dana, syn. ricordi, M.-Edw., p. 253, fig. 5; elongata, A. M.-Edw., p. 256; DE MAN, Notes Leyd. Mus. xiv.
- S. curaçaoensis, n. sp., Curaçao, p. 257, fig. 6; DE MAN, Notes Leyd. Mus. xiv. S. bidentata, n. sp., Jamaica, p. 77; BENEDICT, Johns Hopk. Univ. Circ. xi. S. melissa, de Man, var. moluccensis, Flores, p. 328; mæschii, n. sp., Sumatra, p. 331, pl. xx, fig. 14; frontalis, A. M.-Edw., Flores, p. 334, pl. xx, fig. 13; weberi, n. sp., Flores, p. 338, fig. 15; DE MAN, in Max Weber's Zool. Ergeb. ii. S. crassipes, n. sp., Pernambuco, barbimana, n. sp., Payta, pp. 93, 244 & 245; GAVINO, Boll. Soc. Nat. Napoli (1) iii.
- G. oscsarma, n. subg. of Sesarma, p. 341, nodulifera, n. sp., Java, p. 342, pl. xx, fig. 16; sylvicola, n. sp., Sumatra, p. 345, pl. xx, fig. 18; maculata, n. sp., Flores, p. 347, pl. xxi, fig. 19; DE Man, in Max Weber's Zool. Ergeb. ii.

#### OXYSTOMATA.

This group with its three sections, the *Dorippinea*, Calappinea, and the Leucosinea, pp. 550, &c.; ORTMANN, Zool. Jahrb. vi.

#### CYCLODORIPPIDA:

[New family, p. 582, ORTMANN, Zool. Jahrb. vi.]

Cyclodorippe dromioides, Japan, pp. 559, pl. xxvi, fig. 5, uncifera, Japan, p. 560, fig. 6, Ortmann, Zool. Jahrb. vi, n. spp.

#### HIPPIDEA.

The group, with its two families, Hippidæ and Albuniidæ, characterized, p. 531-538; ORTMANN, t. c.

#### DROMIIDEA.

- The group, with three families, Homolidæ, Dynomenidæ, n. fam., and Dromiidæ, characterized, pp. 538-542; ORTMANN, Zool. Jahrb. vi.
- Cryptodromia canaliculata, n. var. ophryoessa, Japan, pp. 545 & 546; Ortmann, Zool. Jahrb. vi.
- Dicranodromia döderleini, n. sp., Japan, p. 549, pl. xxvi, fig. 4, Ort-MANN, Zool. Jahrb. vi.

#### CALAPPIDÆ.

Calappa japonica, n. sp., Japan, p. 566, pl. xxvi, fig. 8, Ortmann, Zool. Jahrb. vi.

### MATUTIDE.

Notopus novemdentatus, n. sp., New Caledonia, p. 574, pl. xxvi, fig. 11, ORTMANN, Zool. Jahrb. vi.

#### LEUCOSIIDE.

- Cryptocnemus obolus, n. sp., Japan, p. 576, pl. xxvi, fig. 12, ORTMANN, Zool. Jahrb. vi.
- Ebalia longimanus, Japan, p. 579, pl. xxvi, fig. 13, scabriuscula, Japan, p. 580, fig. 14, conifera, Japan, p. 580, fig. 15, Ortmann, Zool. Jahrb. vi, n. spp.
- Philyra heterograna, Japan, p. 582, pl. xxvi, fig. 17, syndactyla, Japan, p. 583, pl. xxvi, fig. 18, Orthann, Zool. Jahrb. vi, n. spp.

#### ANOMUR.1.

### GALATHEIDEA.

This group characterized, with the families Ægleidæ, Chirostylidæ, n., Galatheidæ, and Porcellanidæ; Ortmann, Zool. Jahrb. vi, pp. 241-326,

# PORCELLANIDÆ.

- Pachycheles lavidactylus, n. sp., Brazil, p. 26, pl. xii, fig. 1, ORTMANN, Zool. Jahrb. vi.
- Polyonyx carinatus, Loo Choo, p. 268, pl. xii, fig. 2, ORTMANN, Zool. Jahrb. vi, n. sp.
- Porcellana pulchellula, n. sp., St. Lorenzo, p. 260, GAVINO, Boll. Soc. Nat. Napoli (1) iii.

#### CHIROSTYLIDÆ.

[New family, p. 244, ORTMANN, Zool. Jahrb. vi.]

Chirostylus, n. g., p. 246, for C. dolichopus, n. sp., Japan, p. 246, p. xi, fig. 2; Ortmann, Zool. Jahrb. vi.

#### GALATHEIDE.

- Uroptychus japonicus, Japan, p. 248, pl. xi, fig. 3, ORTMANN, Zool. Jahrb. vi, n. sp.
- Galathea: synopsis of the species, pp. 248 & 249; ORTMANN, Zool. Jahrb. vi.
- G. affinis, n. sp., Fiji I., p. 252, pl. xi, fig. 9, ORTMANN, Zool. Jahrb. vi.
- Munida heteracantha, Japan, p. 255, pl. xi, fig. 12, ORTMANN, Zool. Jahrb. vi. n. sp.
- Munidopsis taurulus, Japan, p. 256, pl. xi, fig. 13, ORTMANN, Zool. Jahrb. vi; M. polymorpha, Canary Is, p. 109, pl. x, figg. 3-16, Kælbel, Ann. Hofmuseum Wien, vii: n. spp.
- Galucantha camelus, Japan, p. 257, pl. xi, fig. 14; ORTMANN, Zool. Jahrb. vi, n. sp.

#### PORCELLANIDE.

Petrolisthes reissii, Ecuador, p. 260, pl. xi, fig. 15, leporinoides, S. Sea, p. 263, ORTMANN, Zool. Jahrb. vi, n. spp.

#### PAGURIDEA.

This tribe contains the families Parapagurida, Pagurida (with subfams. Pagurina, Diogenina, Eupagurina), Canobitida, and Lithodida, pp. 269-273; ORTMANN, Zool. Jahrb. vi.

# PARAPAGURIDE.

Parapagurus abyssorum, Hend., is synonymous with P. pilosimanus, Smith, p. 204; EDWARDS & BOUVIER, Ann. Sci. Nat. xiii.

Chiroplatea scalata, Gulf of Mexico, p. 275, pl. xii, fig. 4; ORTMANN, Zool. Jahrb. vi, n. sp.

# PAGURIDE.

- Anapagurus pusillus, Hend., n. var. japonicus, Japan, p. 296, pl. xii, fig. 11; Ortmann, Zool. Jahrb. vi.
- A. brecicarpus, Spain, p. 215, bicorniger, Cadiz, p. 215, EDWARDS & BOUVIER, Ann. Sci. Nat. xiii; A. curridactylus, Dakar, p. 253, CHEVREUX & BOUVIER, Bull. Soc. Z. Fr. xv, and Mém. Soc. Zool. v, p. 91, pl. ii, figg. 2-8: n. spp.
- Calcinus talismani, Cape Verde, p. 225, EDWARDS & BOUVIER, Ann. Sci. Nat. xiii, n. sp.
- Catapaguroides, n. g. for C. microps, Spain, p. 211, megalops, Marocco, p. 213, acutifrons, Canaries, p. 213, n. spp., Edwards & Bouvier, Ann. Sci. Nat. xiii.
- Clibanarius melitai, Dakar, p. 255, Bull. Soc. Z. Fr. xvi, and p. 135, pl. iv, figg. 3-6, Mém. Soc. Zool. v; senegulensis, Dakar, p. 256, t. c. and p. 131, pl. iv, figg. 7-11, t. c., Chevreux & Bouvier: n. spp.
- Diogenes denticulatus, n. sp., Rufisque, CHEVREUX & BOUVIER, p. 254, Bull. Soc. Z. Fr. xvi, and p. 122, pl. iii, figg. 16-20, Mém. Soc. Zool. v.
- Eupagurus lavimanus, Tahiti, p. 302, pl. xii, fig. 13; dubius, Japan, p. 307. pl. xii, fig. 14; triserratus, Japan, p. 308, pl. xii, fig. 15; similis, Japan, p. 310; barbatus, Japan, p. 311; samoensis, Samoa, p. 312. pl. xii, fig. 17; obtusifrons, Japan, p. 313, pl. xii, fig. 18; ophthalmicus, Japan, p. 314, pl. xii, fig. 19, ORTMANN, Zool. Jahrb. vi; E. variabilis, Gulf of Gascony, p. 217; pubescentulus, Marocco, p. 219; irregularis, Cape Blanco, p. 220; pulchellus, Cape Verde, p. 221; ruber, ? Gulf of Gascony, p. 222, EDWARDS & BOUVIER, Ann. Sci. Nat. xiii; triangularis, Dakar, p. 253, Bull. Soc. Z. Fr. xvi, and p. 93, pl. ii, figg. 9-15, Mém. Soc. Zool. v; inermis, Dakar, p. 254, t. c. and p. 109, pl. iii, t. c. figg. 1-5; minimus, Dakar, p. 106, pl. ii, figg. 21-26, CHEVREUX & BOUVIER; E. alaskensis, Alaska, p. 2, aleuticus, Aleutian Is., p. 3; patagoniensis, Patagonia, p. 4; smithi, Gulf of California, p. 4; impressus, Florida, p. 5; floridanus, Florida, p. 5; exilis, La Plata, p. 6; albus, California, p. 6; gladius, California, p. 7; defensus, Gulf of Mexico, p. 7; capillatus, Alaska, p. 8; brandti, dalli, Aleutian Is., p. 9; tanneri, p. 10; confragosus, p. 11; cornutus, p. 12; townsendi, p. 13; rathbuni, p. 14, Alaska; minutus, Gulf of California, p. 14; purpuratus, Galapagos I., p. 15; hemphilli, California, p. 16; beringanus, Aleutian Is., p. 17; newcombei, British Columbia, p. 17; undosus, Bering Sea, p. 18; setosus, minutus, p. 19; gilli, p. 20, Alaska, curaçacensis, Curaçao, p. 21; californiensis, mexicanus, roscus,

California, p. 22; corallinus, Florida, p. 23; coronatus, California, p. 24; varians, Gulf of California, p. 24; cervicornis, p. 25; parvus, p. 26, California; hispidus, Cuba, p. 26, Benedict, P. U. S. Nat. Mus. xv; n. spp.

Nematopagurus, n. g., p. 209, for N. longicornis, n. sp., Mediterranean, &c.; EDWARDS & BOUVIER, p. 210, Ann. Sci. Nat. xiii.

Paguristes polythophilus, p. 277, pl. xii, fig. 5, acanthomerus, p. 279, pl. xii, fig. 6, kagoshimensis, p. 281, pl. xii, fig. 8, Japan, Ortmann, Zool. Jahrb. vi; P. hispidus, Liberia, p. 208, Edwards & Bouvier, Ann. Sci. Nat. xiii: n. spp.

#### LITHODIDÆ.

Lithodes turritus, Japan, p. 320, pl. xii, fig. 26, ORTMANN, Zool. Jahrb. vi, n. sp.

Hapalogaster grebnitzkii, Behring Is., p. 335, fig. 3, brandti, Sitka, p. 336, fig. 1, Schalfeef, Bull. Pétersb. xxxv, n. spp.

Placetron, n. g., p. 339, for P. wosnessenskii, n. sp., Kadjak, p. 339, fig. 6; Schalfeef, Bull. Pétersb. xxxv.

#### MACRURA.

The groups Homaridea, Loricata, and Thalassinidea, characterized. The Homaridea, containing the Homaridæ and Astacidæ; the Loricata, containing the Palinuridæ and Scyllaridæ; and the Thalassinidea, containing the Axiidæ, Thalassinidæ, Calocaridæ (n. fam.), and Callianassidæ; ORTMANN, Zool. Jahrb. vi.

#### ASTACIDÆ.

Astacus similis, Corea; KŒLBEL, Anz. Ak. Wien, xviii, pp. 176 & 177, n. sp.

#### PALINURIDÆ.

The known genera of the family characterized; ORTMANN, pp. 14 & 15, Zool. Jahrb. vi.

Avus, n. g., syn. Linuparis, Gray; ORTMANN, p. 21, Zool. Jahrb. vi.

Senex, n. g., syn. Panulirus, Gray: synopsis of the species; ORTMANN, pp. 22 & 23, Zool. Jahrb. vi.

Puer, n. g., syn. Panulirus, Bate in part, for angulatus, Bate, and pellucidus, n. sp., Japan, p. 37, pl. i, fig. 3; ORTMANN, Zool. Jahrb. vi.

#### AXIIDE.

Eiconaxius farress, Japan, pp. 49 & 50, pl. i, fig. 4, Ortmann, Zool. Jahrb. vi, n. sp.

# CALLIANASSIDÆ.

- Gebia littoralis, Risso, Naples, &c., p. 53, pl. i, fig. 6, major, De Haan, Japan, p. 54, pl. i, fig. 7, barbata, Str., Samoa, p. 54, pl. i, fig. 8, isodactyla, n. sp., Red Sea, p. 55, pl. i, fig. 9; Ortmann, Zool. Jahrb. vi.
- Callianassa subterranea, Mont., Mediterranean Sea, p. 55, pl. i, fig. 10, n. var. japonica, Japan, fig. 10a, diademata, n. sp., Africa, p. 56, pl. i, fig. 11; ORTMANN, Zool. Jahrb. vi.

# ATYIDÆ.

- Atya moluccensis, De Haan, Java, Sumatra, Celebes, &c., p. 357, pl. xxi, fig. 20, brevirostris, n. sp., Flores, p. 360, pl. xxi, fig. 21; DE MAN, in Max Weber's Zool. Ergeb. ii.
- Caridina typus, M.-Edw., Flores, Celebes, p. 367, pl. xxi, fig. 22; weberi, n. sp., Flores, &c., p. 371, pl. xxii, fig. 23, var. sumatrensis, n. var., Sumatra, p. 375, pl. xxii, fig. 33G; parvirostris, n. sp., Flores, p. 375, pl. xxii, fig. 24; lævis, Hell., Java, p. 376, pl. xxiii, fig. 27; pareparensis, n. sp., Celebes, p. 379, pl. xxii, fig. 25; multidentata, n. sp., Celebes, p. 380, pl. xxii, fig. 26; serratirostris, n. sp., Flores, p. 382, pl. xxiii, fig. 28, n. var. celebensis, Celebes, p. 385, pl. xxiii, fig. 28; wyckii, Hickson, Celebes, p. 386, pl. xxiv, fig. 29, n. var. gracilipes, Celebes, p. 393; brevicarpalis, Celebes, p. 397, pl. xxiv, fig. 30, n. var. endehensis, p. 399, pl. xxiv, fig. 30e; gracilirostris, n. sp., Celebes, p. 399, pl. xxv, fig. 31; De Man, in Max Weber's Zool. Ergeb. ii; japonica, n. sp., Japan, p. 261, pl. ix, figg. 7 & 8; De Man, Notes Leyd. Mus. xiv.

#### PALÆMONIDÆ.

- Alpheus gracilidigitus, Miers, loc. ?, p. 406, pl. xxv, fig. 32; DE MAN, in Max Weber's Zool. Ergeb. ii.
- Hippolyte ponapensis, Ortm., is a synonym of Hetairocaris orientalis, De Man; De Man, p. 263, Notes Leyd. Mus. xiv.
- Palæmon weberi, n. sp., Celebes, p. 421, pl. xxv, fig. 33; dispar, v. Martens, Celebes, &c., p. 427, pl. xxvi, fig. 34; sundaicus, Heller, Flores, Celebes, p. 437, pl. xxvi, fig. 35; elegans, n. sp., Java, p. 440, pl. xxvi, fig. 36; equidens, Dana, Sumatra, p. 453, pl. xxvi, fig. 36; javanicus, Hell., Sumatra, p. 457, pl. xxvi, fig. 38; horstii, n. sp., Celebes, p. 460, pl. xxvii, fig. 39; scabriculus, Hell., Celebes, p. 462, pl. xxvii, fig. 41; endehensis, n. sp., Flores, p. 465, pl. xxvii, fig. 42; modestus, n. sp., Flores, p. 469, pl. xxvii, fig. 43; pilimanus, De Man, Sumatra, p. 471, pls. xxvii & xxviii, fig. 44, n. var. leptodactylus, Java, p. 476 pl. xxviii, figg. 44 & k-i; latimanus, v. Mart., syn. euryrhynchus, Ortm., Flores, Timor, &c., p. 477, pl. xxviii, fig. 45; placidus, n. sp., Sumatra, p. 483, pl. xxviii, fig. 46; placidulus, n. sp., Flores, &c., p. 489, pl. xxviii, fig. 48; lampropus, n. sp., Celebes, p. 493, pl. xxix

fig. 49; bariensis, n. sp., Flores, p. 490, pl. xxix, fig. 50; lepidacty-loides, n. sp., Flores, p. 497, pl. xxix, fig. 51; lepidactylus, Thal., Celebes, p. 501, fig. 52; latidactylus, Thal., Flores, p. 504, pl. xxix, fig. 52; De Man, in Max Weber's Zool. Ergeb. ii.

#### PENÆIDÆ.

Penœus semisulcatus, De Haan, Celebes, p. 510; indicus, M.-Edw., syn. merguiensis, De Man, Celebes, p. 511, pl. xxix, fig. 3; monoceros, Fabr., Celebes, p. 513, pl. xxix, fig. 54; DE MAN, in Max Weber's Zool. Ergeb. ii.

#### PSALIDOPODIDÆ.

[N. fam., p. 265; WOOD-MASON & ALCOCK, Ann. N. H. (6) ix.]

Psalidopus, n. g., p. 266, for P. huxleyi, Andaman Sea, p. 273, pl. xiv, figg. 1, 2, 7, spiniventris, p. 274, pl. xiv, figg. 3-6, and pl. xv, figg. 1-10, n. spp., Andaman Sea; WOOD-MASON & ALCOCK, t. c.

#### ACANTHEPHYRIDE.

Acanthephyra sanguinea, n. sp., p. 358, fig. 1; armata, M.-Edw., p. 359, fig. 2; eximia, Smith, p. 361, fig. 3; brachytelsonis, Bate, p. 363, fig. 4; curtirostris, W.-M., p. 364, fig. 5, Indian Ocean; Wood-Mason & Alcock, Ann. N. H. (6) ix.

#### SCHIZOPODA.

NORMAN, Ann. N. H. (6) ix, pp. 454-464, and x, pp. 143-166, 242-263, characterizes this group with the families, sub-families, genera and species that are known to occur in Britain.

#### CRYPTOPODIDÆ.

[N. fam., p. 444, &c., Beneden, Bull. Ac. Belg. xxii.]

Cryptopus, syn. Cerataspis, referred to the Penæidæ, p. 1029; GIARD & BONNIER, C.R. exiv.

# Mysidæ.

Mysidopsis gibbosa, Sars, p. 164, pl. x, fig. 8; angusta, Sars, p. 165, fig. 9, British Seas; hibernica, n. sp., Valentia, p. 165, pl. ix, figg. 1-5; NORMAN, Ann. N. H. (6) ix.

Schistomysis, n. g., p. 254, for S. parkeri, n. sp., Devon, p. 256, pl. x, figg. 1-7; NORMAN, Ann. N. H. (6) ix.

Heteromysis formosa, Smith, syn. norvegica, Sars, British Seas, p. 158, pl. ix, figg. 6-11; NORMAN, Ann. N. H. (6) ix.

Erythrope elegans, Sars, p. 161, pl. x, fig. 10; NORMAN, Ann. N. H. (6) ix.

#### EDRIOPHTHALMA.

#### CUMACEA.

- Cyclaspis levis, Dunedin, p. 264, pls. xvi & xvii, figg. 1-26, Thomson, J. L. S. xxiv, n. sp.
- Diastylis neo-zelandica, Dunedin, Otago Bay, p. 268, pl. xviii, figg. 1-11; Thomson, J. L. S. xxiv, n. sp.

#### AMPHIPODA.

#### Amphilochidæ.

- Astyra abyssi, Boeck, Norway, p. 214, pl. lxxiii; SARS, Amphipoda of Norway.
- Amphilochoides, n. g., p. 220, for Amphilochus odontonyx, Boeck, p. 221, pl. lxxv, fig. 2; and A. pusillus, n. sp., p. 222, pl. lxxvi, fig. 1, Norway; SARS, Amphipoda of Norway.
- Amphilochus manudens, Sp. Bate, p. 217, pl. lxxiv; tenuimanus, n. sp., p. 218, pl. lxxv, fig. 1, Norway; Sars, Amphipoda of Norway.
- Gitana sarsii, Boeck, syn. A. sabrinæ, p. 228, pl. lxxviii, fig. 1; abyssicola, n. sp., p. 229, pl. lxxviii, fig. 2; rostrata, Boeck, p. 230, pl. lxxix, fig. 1, Norway; Sars, Amphipoda of Norway.
- Gitanopsis, n. g., p. 223, for G. bispinosa, Boeck, p. 224, pl. lxxvi, fig. 2; inermis, Sars, p. 225, pl. lxxvii, fig. 1, and G. arctica, n. sp., p. 227, pl. lxxvii, fig. 2, Norway; Sars, Amphipoda of Norway.
- Stegoplax longirostris, Sars, Norway, p. 232, pl. lxxix, fig. 2; SARS, Amphipoda of Norway.

# GAMMARIDE.

Gammarus delebecquei, from the Lake of Annecy, p. 1286, CHEVREUX & DE GUERNE, Bull. Soc. Z. Fr. xvii, & C.B. cxiv, n. sp.

#### LEUCOTHOIDÆ.

Leucothoe spinicarpa, Abild., p. 283, pls. c & ci, fig. 1, lilljeborgi, Boeck, p. 284, pl. ci, fig. 2; SARS, Amphipoda of Norway. L. incisa, n. n. for L. furina, Chevreux, from Cumbrae, p. 217; ROBERTSON, P. N. H. Soc. Glasg. iii.

#### LYSIANAXIDÆ.

- Synonymy of British Lysianaxida; WALKER, Ann. N. H. (6) ix, pp. 134-138.
- Pararistias, n. g., p. 201, for Lysianassa audouinianus, Bate; Robertson, P. N. H. Soc. Glasg. iii.
- Perrierella, n. g. for P. crassipes, n. sp., France, p. 50; Chevreux & Bouvier, Bull. Soc. Z. Fr. xvii.
- Socarnes erythrophthalmus, Cumbrae, p. 200, Robertson, P. N. H. Soc. Glasg. iii, n. sp.

#### ŒDICERIDÆ.

- Aceros phyllonyx, Sars, p. 338, pls. cxix & cxx, fig. 1, Norway; Sars, Amphipoda of Norway.
- Bathymedon, n.g., p. 332, for B. longimanus, Boeck, p. 333, pl. cxvii; saussurei, Boeck, p. 335, pl. cxviii, fig. 1; obtusifrons, Hans., p. 336, pl. cxviii, fig. 2, Norway; SARS, Amphipoda of Norway.
- Halicreion longicaudatus, Boeck, p. 322, pl. cxiii, fig. 2, Norway; Sans, Amphipoda of Norway.
- Halimedon mülleri, Boeck, p. 327, pl. cxv; acutifrons, n. sp., p. 329, pl. cxvi; megalops, Sars, p. 330, pl. cxvi, fig. 2; brevicalcar, Goës, p. 331, pl. cxvi, fig. 3; Norway; SARS, Amphipoda of Norway.
- Monoculodes carinatus, Sp. Bate, p. 295, pl. cv; tessellatus, Schn., p. 297, pl. cvi, fig. 1; borealis, Boeck, p. 298, pl. cvi, fig. 2; pallidus, n. sp., p. 299, pl. cvi, fig. 3; norvegicus, Boeck, p. 301, pl. cvii, fig. 1; falcatus, n. sp., p. 302, pl. cvii, fig. 2; tuberculatus, Boeck, p. 303, pl. cvii, fig. 3; latimanus, Goës, p. 304, pl. cviii, fig. 1; kroyeri, Boeck, p. 305, pl. cviii, fig. 3; longirostris, Goës, p. 306, pl. cviii, fig. 3; packurdi, Boeck, p. 307, pl. cix, fig. 1; tenuirostratus, Boeck, p. 309, pl. cix, fig. 2; Norway; Sars, Amphipoda of Norway.
- Monoculopsis, n. g., p. 310, for longicornis, Boeck, p. 311, pl. cx, fig. 1, Norway; Sars, Amphipoda of Norway.
- Œdiceropsis brevicornis, Lillj., p. 325, pl. cxiv, Norway; SARS, Amphipoda of Norway.
- Œdiceros saginatus, Kröy., p. 288, pl. cii; borealis, Boeck, p. 290, pl. ciii, fig. 1, Norway; Sars, Amphipoda of Norway.
- Parædiceros, n. g., p. 291, lynceus, Sars, p. 292, pl. ciii, fig. 2, & pl. civ, fig. 1; propinquus, Göes, p. 293, pl. civ, fig. 2, Norway; Sars, Amphipoda of Norway.
- Periculodes, n. g., p. 312, for P. longimanus, Sp. Bate, p. 313, pl. cx, fig. 2, & pl. cxi, fig. 1, Norway; SARS, Amphipoda of Norway.
- Pontocrates norvegicus, Boeck, p. 315, pl. cxi, fig. 2, Norway; SARS, Amphipoda of Norway.
- Synchelidium, n. g., p. 317, for brevicarpum, Sp. Bate, p. 318, pl. cxii, fig. 1; haplocheles, Grube, p. 319, pl. cxii, fig. 2; intermedium, n. sp., p. 320, pl. cxiii, fig. 1, Norway; SARS, Amphipoda of Norway.

#### ORCHESTIDÆ.

Orchestia floresiana, p. 562, martensii, p. 564, Flores, parvispinosa, Java, p. 566, montana, Celebes, p. 567, Max Weber, Zool. Ergeb. ii, n. spp. Hyale grimaldii, p. 257, figg. 1-5, Chevreux, Bull. Soc. Z. Fr. xvi, n. sp.

# STENOTHOIDE.

Cressa dubia, Sp. Bate, p. 278, pl. xcviii, fig. 2, & pl. xcix, fig. 1; minuta, Boeck, p. 280, pl. xcix, fig. 2, Norway; SARS, Amphipoda of Norway. Metopa alderi, Sp. Bate, p. 250, pl. lxxxvi; spectabilis, Sars, p. 251,

pl. lxxxvii; boeckii, n. sp., syn. M. bruzelii, Boeck; borealis, Sars, in part, p. 252, pl. lxxxviii; borealis, Sars, p. 254, pl. lxxxix, fig. 1; rubrovittata, Sars, p. 255, pl. lxxxix, fig. 2; pusilla, n. sp., p. 256, pl. xc, fig. 1; longicornis, Boeck, p. 258, pl. xc, fig. 2; tenuimana, n. sp., p. 259, pl. xci, fig. 1; affinis, Boeck, p. 260, pl. xci, fig. 2; bruzelii, Goës, p. 261, pl. xcii, fig. 1; sinuata, n. sp., syn. bruzelii, Hans., in part, p. 263, pl. xcii, fig. 2; propinqua, n. sp., p. 264, pl. xciii, fig. 1; leptocarpa, Sars, p. 265, pl. xciii, fig. 2; sölsbergi, Schneider, p. 266, pl. xciv, fig. 1; invalida, n. sp., syn. M. alderi, Boeck, in part, p. 267, pl. xciv, fig. 2; pollexiana, Sp. Bate, p. 269, pl. xcv; robustu, n. sp., p. 270, pl. xcvi, fig. 1; palmata, n. sp., p. 272, pl. xcvi, fig. 2; longimana, p. 273, pl. xcvii, fig. 1; neglecta, Hansen, p. 274, pl. xcvii, fig. 2; nasuta, Boeck, p. 276, pl. xcviii, fig. 1: from Norway; Sars, Amphipoda of Norway.

Metopella, n. g., suggested as new generic name for Metopa longimana, Sp. Bate, p. 274; Sars, Amphipoda of Norway.

Probolium gregarium, Sars, p. 245, pl. lxxxiv; calcaratum, Sars, p. 247, pl. lxxxv, Norway; Sars, Amphipoda of Norway.

Stenothoe marina, Sp. Bate, p. 236, pl. lxxx; microps, n. sp., p. 237, pl. lxxxi, fig. 1; tenella, Sars, p. 238, pl. lxxxi, fig. 2; monoculoides, Mont., p. 240, pl. lxxxii, fig. 1; brevicornis, Sars, p. 241, pl. lxxxii, fig. 2; megachir, Boeck, p. 242, pl. lxxxiii, Norway; SARS, Amphipoda of Norway. S. dollfusi, n. sp., Azores, p. 260, figg. 6-10, Chevreux, Bull. Soc. Z. Fr. xvi.

# Aliarum Familiarum.

Hyperia schizogeneios, Stebbing, adult & described; Chevreux, Bull. Soc. Z. Fr. xvii, p. 233.

Pardaliscoides, n. g. for P. tenellus, n. sp.; STEBBING, Rep. on the Amphipoda of the "Challenger," p. 1725 (1888).

#### ISOPODA.

#### ASELLIDÆ.

Asellus coxalis, Syria, p. 13, pl. ii, fig. 11, DOLLFUS, Rev. Biol. iv, n. sp. Jaropsis neo-zelandica, Akaroa, p. 267, CHILTON, Tr. N. Z. Inst. xxiv, n. sp. Munna neo-zelandica, New Zealand, p. 2, pls. i & ii, figg. 1-15; CHILTON, Tr. N. Z. Inst. xxiv, n. sp.

#### Bopyridæ.

Probopyrus giardi on Palamon placidus, Sumatra, p. 557, pl. xxx, fig. 1, Max Weber, Zool. Ergeb. ii, n. sp.

Palegyge borrei, Giard & Bonnier, on Palemon dispar, Flores, p. 557, pl. xxx, fig. 5; bonnieri, n. sp., on P. lar, Celebes, p. 558, pl. xxx, fig. 4; fluviatilis, n. sp., on P. lampropus, Celebes, p. 559; de mani, n. sp., on P. spinimanus, Sumatra, p. 560; Max Weber, Zool, Ergeb. ii.

#### ONISCIDÆ.

- Armadillidium scabrum, n. sp., Seville, p. 165, with fig., Dollfus, Ann. Soc. Esp. (2) i. A. sanctum, n. sp., Syria, p. 4, pl. i, fig. 2; granum, n. sp., Syria, p. 5, fig. 3; Dollfus, Rev. Biol. iv. A. granulatum, Br., serratum, B.-L., nasatum, B.-L., maculatum, Risso, simoni, Dollf., assimile, B.-L., esterelanum, Dollf., zenckeri, Br., pictum, Br., pulchellum, Br., vulgare, Latr., sordidum, Dollf., opacum, Kochs, alpinum, Dollf., quinquepustulatum, B.-L., depressum, Br., album, Dollf., from France, figured and described, pp. 8-18, figg. 1-26; Dollfus, Feuill. Nat. xxii.
- Armadillo officinalis, n. var. syriaca, p. 2, albomarginatus, n. sp., Syria, p. 3; Dollfus, Rev. Biol. iv.
- Mesarmadillo, n. g., p. 385; alluaudi, p. 386, pl. vii, fig. 1, marginatus, p. 387, fig. 2, tuberculatus, p. 387, fig. 3, Assinie, Dollfus, Ann. Soc. Ent. Fr. lxi, n. spp.
- Metoponorthrus rufocinctus, Corunna, p. 184, fig.; Dollfus, Ann. Soc. Esp. (2) i; M. pica, Syria, p. 10, pl. ii, fig. 9, trifasciatus, Tiberias, p. 11, pl. ii, fig. 10; Dollfus, Rev. Biol. iv: n. spp.
- Philoscia minima, Granada, p. 187, fig., Dollfus, Ann. Soc. Esp. (2) i, n. sp.
- Porcellio magnificus, n. sp., Alcazaba, p. 270, fig., succinctus, B.-L., p. 171, fig.; P. nicklesi, p. 171, fig., bolivari, p. 172, fig., expansus, p. 174, fig., pyrenœus, p. 175, fig., de bueni, p. 176, fig., amænus, p. 178, fig., hispanus, p. 179, fig., nigro-granulatus, p. 182, fig., Spain, n. spp.; Dollfus, Ann. Soc. Esp. (2) i. P. inconspicuus, Dead Sea, p. 5, pl. i, fig. 4, fissifrons, Syria, p. 7, pl. i, fig. 5, barroisi, Dead Sea, p. 7, pl. i, fig. 6, contractus, Syria, p. 8, pl. ii, fig. 7; P. (Leptotrichus) pulchellus, Syria, pl. ii, fig. 8; Dollfus, Rev. Biol. iv.: n. spp.

Synarmadillo, n. g., p. 388, for S. clausus, n. sp., Assinie, p. 388, pl. vii, fig. 4; Dollfus, Ann. Soc. Ent. Fr. lxi.

# CYMOTHOIDE, &c.

- Ichyoxenus jellinghausii, Herkl., on species of Puntius, Sumatra, p. 545;
  MAX WEBER, Zool. Ergeb. ii.
- Lironeca sulcata, Canary Is., p. 105, pl. x, figg. 1 & 2, Kœlbel, Ann. Hofmuseum Wien, vii, n. sp.
- Rocinela typus, M.-Edw., Sumatra, p. 553; MAX WEBER, Zool. Ergeb. ii. Sphæroma egregia, Akaroa, p. 269, CHILTON, Tr. N. Z. Inst. xxiv, n. sp. Tachæa lucustris, Sumatra, p. 551, MAX WEBER, Zool. Ergeb. ii.

# ENTOMOSTRACA.

#### PHYLLOPODA.

#### A. Recent Forms.

- Apus (Lepidurus) glacialis, n. var. spitzbergensis, Spitzbergen, p. 291, BERNARD, The Apodidæ.
- Branchipus ledoulzi, Syria, p. 25, fig. 1, BARROIS, Rev. Biol. v. n. sp.
- Estheria hierosolymitans and gihoni, figured and described, pp. 34-37; BARROIS, Rev. Biol. v.

# B. Fossil Forms.

- Anomolocaris, n. g., p. 205, canadensis, p. 206, WHITEAVES, Can. Rec. v, n. sp.
- Aptychopsis burraudeana, n. sp., Moffat Beds, p. 101, pl. xv, fig. 1; cordiformis, n. sp., Lower Wenlock, p. 103, pl. xv, fig. 2; lata, n. sp., p. 104, pl. xv, fig. 6; ovata, n. sp., Dumfries, p. 108, fig. 4; subquadrata, n. sp., Upper Silurian, p. 110, fig. 20; oblata, n. sp., Gala group, p. 111, figg. 18, 21, & 23; Jones & Woodward, Palmont. Soc.
- Discinocaris ovalis, p. 121, pl. xvi, fig. 20, undulata, p. 122, pl. xvi, fig. 24, Jones & Woodward, Palscont. Soc., n. spp.
- Peltocaris patula, Moffat Series, p. 116, pl. xvi, figg. 10 & 11, carruthersii, syn. aptychoides, Woodw., Moffat Series, p. 116, pl. xvii, fig. 7, JONES & WOODWARD, Palseont. Soc., n. spp.

#### CLADOCERA.

- Alona cilio-caudata, S. W. Russia, p. 159, pl. iii, figg. 33-36, SOWINSKY, Zapiski. Kiev, xi. A. lepida, Wisconsin, p. 393, pl. xiii, fig. 19, BIRGE, Tr. Wisconsin Acad. viii: n. spp.
- Grimaldina brazzai, Congo, p. 214, figg. 1-3, RICHARD, Mém. Soc. Zool. v, n. sp.
- Guernella raphaelis, p. 218, figg. 4-6, RICHARD, Mém. Soc. Zool. v, n. sp.
- Latonopsis occidentalis, Wisconsin, p. 383, pl. xiii, figg. 1-5, BIRGE, Tr. Wisconsin Acad. viii, n. sp.
- Macrothrix chevreuxi, Rufisque, p. 530, Guerne & Richard, Mém. Soc. Zool. v, n. sp.
- Moina dubia, Rufisque, p. 527, GUERNE & RICHARD, Mém. Soc. Zool. v, n. sp.
- Moinodaphnia mocquersyi, p. 222, figg. 7 & 8, RICHARD, Mém. Soc. Zool. v, n. sp.
- Pleurozus truncatus, n. var. Poppei, p. 189; KLOCKE, Zool. Anz. xv.

#### OSTRACODA.

- Cypris ungulata, Algeria, p. 247, blanchardi, p. 252, Moniez, Mém. Soc. Zool. iv, n. spp.
- Leucocythere, n. g., for L. mirabilis, p. 394; KAUFMAN, Zool. Anz. xv, n. sp.
- Pachycypris leuckarti, p. 57, incisa, p. 59, S. America, Claus, Arb. z. Inst. Wien, x, n. spp.

#### COPEPODA.

- Argulus indicus, Java, p. 544, fig. 1, MAX WEBER, Zool. Ergeb. ii, n. sp.
  - Arietellus, n. g. for A. setosus, n. sp., Naples, p. 415; GIESBRECHT, Copepoden des Golfes von Neapel.
  - Belisarius vignieri, p. 132, MAUPAS, C.R. cxv, p. 132, n. sp.
- Caligus dakari, p. 243, pl. i, figg. 1-4, Beneden, Bull. Ac. Belg. xxiii, n. sp.
  - Caligera difficilis, n. sp., p. 258, p. iv, Beneden, Bull. Ac. Belg. xxiii.
  - Calina brachyura, p. 249, pl. ii, BENEDEN, Bull. Ac. Belg. xxiii, n. sp.
  - Candace varicans, Naples, p. 439, GIESBRECHT, Copepoden des Golies von Neapel, n. sp.
  - Canthocamptus yahiai, Algeria, p. 516, pl. vi, fig. 42, Blanchard & Richard, Mém. Soc. Zool. iv, n. sp.
  - Chiridius, n. g. for C. poppei, n. sp., Naples, p. 224; GIESBRECHT, Copepoden des Golfes von Neapel.
  - Chlamys, n. g. for C. incisus, n. sp., Dakar Bay, pp. 227-231, pl. ii, figg. 1-10; Beneden, Bull. Ac. Belg. xxiii.
  - Copilia (Saphirinella), genus monographed, with chart of geographical distribution; Dall, Zool. Jahrb. vi.
  - Cyclops hamatus, kiewensis, brevisetosus, palustris, poggenpolii, uljanini, trouchanowi, kovostyschewi, n. spp. from S. W. Russia, mentioned, but apparently not described; Sowinsky, Zapiski. Kiev, xi. C. dybowskii, n. sp., Germany, p. 163, Landé, Mém. Soc. Zool. v.
  - Dactylopus jugurtha, n. sp., Algeria, p. 519, pl. vi, figg. 26-41, Blan-Chard & Richard, Mém. Soc. Zool. iv.
  - Diaptomus sancti-patricii, Connemara, p. 99, pl. xiv, figg. 5-7, hircus, Orkney, p. 100, pl. x, figg. 2-4, Brady, Nat. Hist. Tr. North Durham, xi; D. schmackeri, Lake Tahoo (China), p. 149, Poppe & Richard, Bull. Soc. Z. Fr. xvii; n. spp.
  - Euchata acuta, p. 262, spinosa, p. 263, Naples, GIESBRECHT, Copepoden des Golfes von Neapel, n. spp.
  - Gryropeltis ranarum, Central Africa, pp. 152-154; STUHLMANN, Zool. Jahrb. vi, n. sp.
  - Hemicalanus acutifrons, fertilis, ornatus, spiniceps, Naples, pp. 398 & 399, Giesbrecht, Copepoden des Golfes von Neapel, n. spp.

- Herpyllobiidæ, n. fam. for Rhizorhina, n. g., Ent. Med. 1892, pp. 207-231.
- Ilyopsyllus, syn. Abacola, jousseaumei, n. sp., p. 69; RICHARD, Bull. Soc. Z. Fr. xvii.
- Laophonte mohammed, Algeria, p. 526, Blanchard & Richard, Mém. Soc. Zool. iv, n. sp.
- Lichomolgus agilis, Firth of Forth, p. 201, pl. xv, figg. 1-14, Scott, T., Ann. N. H. x; L. aberdonensis, E. of Scotland, p. 149, pl. vi, figg. 1-12, Scott, T. & A., Ann. Scott. Nat. Hist. Soc. 1892: n, spp.
- Mesochra lybica, Algeria, p. 523, pl. vi, fig. 16, BLANCHARD & RICHARD, Mém. Soc. Zool. iv, n. sp.
- Metridia normani, Faroe Channel, p. 345, GIESBRECHT, Copepoden des Golfes von Neapel, n. sp.
- Monstrilla gracilicauda, longiremis, Naples, p. 589, GIESBRECHT, Copepoden des Golfes von Neapel, n. spp.
- Möbianus, n. g., for M. gyrans, n. sp., Naples, GIESBRECHT, p. 205, Copepoden des Golfes von Neapel.
- Nogagus angustatus, BENEDEN, Bull. Ac. Belg. xxiv, n. sp.
- Oncea minuta and subtilis, Naples, p. 603, GIESBRECHT, Copepoden des Golfes von Neapel, n. spp.
- Pandarus cranchii, male described and figured, p. 221, pl. i, figg. 1-4; Beneden, Bull. Ac. Belg. xxiii.
- Pupulina flores, p. 254, pl. iii, BENEDEN, Bull. Ac. Belg. xxiv, n. sp.
- Rhizorhina, n. g. (Herpyllobiidæ) for R. ampeliscæ, n. sp., Denmark, pp. 231 & 232, pl. iii; HANSEN, Ent. Med. 1892.
- Rotania, n. g. for R. flava, n. sp., Naples; GIESBRECHT, p. 616, Copepoden des Golfes von Neapel.
- Sapphirina lacters, pyrosomatis, maculosa, Naples, GIESBRECHT, pp. 641-643, Copepoden des Golfes von Neapel, n. spp.
- Scolecithrix vittata, profunda, longipes, dentata, tenuiserrata, dubia, auropecten, Naples, Giesbrecht, Copepoden des Golfes von Neapel, n. spp.
- Synapticola, n. g. for S. teres, n. sp., on Synapta kefersteinii, pp. 31-42; Voigt, Z. wiss. Zool. liii.
- Thaumaleus claparedii and reticulatus, Naples, GIESBRECHT, p. 585, Copepoden des Golfes von Neapel, n. spp.
- Xanthocalanus agilis and minor, Naples, GIESBRECHT, pp. 286–292, Copepoden des Golfes von Neapel, n. spp.

#### CIRRIPEDIA.

#### LEPADIDE.

- Alepas japonica, Japan, p. 125, Aurivillius, Œfv. Ak. Förh. 1892, n. sp.
- Dichelaspis cor, S. Africa, p. 124, bullula, Java, p. 125, Aurivillius, Œfv. Ak. Förh. 1892, n. spp.

- Lepas testudinata, S. Africa, p. 123, AURIVILLIUS, Œfv. Ak. Förh. 1892, n. sp.
- Oxynaspis patens, W. Indies, p. 125, AURIVILLIUS, Œfv. Ak. Förh. 1892, n. sp.
- Pacilasma vagans, off Nautilus, p. 124, Aurivillius, Œfv. Ak. Förh. 1892, n. sp.
- Scalpellum gemma, Greenland, scorpio, China Sea, p. 126, gibberum, La Plata, p. 127, calcaratum, Pacific Ocean, p. 128, septentrionale, North Sea, p. 128, obesum, North Sea, p. 129, erosum, N. W. Atlantic, p. 129, aduncum, on a Phoxichilidium, p. 130, luridum, Baffin Bay, p. 130, grönlandicum, Greenland, prunulum, W. Indies, p. 131, galea, La Plata, p. 132, stratum, W. Indies, p. 132, Aurivillius, Œfv. Ak. Förh. 1892, n. spp.

# LITHOGLYPTIDÆ (n. fam.).

Lithoglyptes, n. g., for L. indicus, p. 133, bicornis, p. 134, ampulla, p. 134, Javan Seas, Aurivillius, Œfv. Ak. Förh. 1892, n. spp.

# PANTOPODA.

#### TITLES.

- CARPENTER, G. H. (1) Report on the *Pycnogonida* collected in Torres Straits by Prof. A. C. Haddon. Sci. P. R. Dubl. Soc. vii (n.s.) pp. 552-558, pl. xxii.
- —. (2) Supplement. Op. cit. viii, pp. 21-26, pl. ii.
- GAUBERT, P. Autotomie chez les Pycnogonides. Bull. Soc. Z. Fr. xvii, p. 224.
- IVES, J. E. A new species of Pycnogonum from California. P. Ac. Philad. 1892, p. 142.
- JOURDAIN, —. Affinites Zoologiques des Pantopodes. C.R. Soc. Biol. 1892, p. 945.
- \*Morgan, T. H. The relationships of the Sea Spiders. Biol. Lect. Marine Biol. Lab. Woods Holl. pp. 142-167. (Cf. Zool. Anz. xv, p. 43.)
- Schimkewitsch, W. Sur les Pantopodes recueillis par M. le lieutenant G. Chierchia pendant le voyage de la corvette "Vettor Pisani" en 1882-1885. Atti Soc. Rom. Mém. (4), vi, pp. 329-347 (1889). [Vide Zool. Rec. xxv, Arachnida, p. 24.]

TOPSENT, E. Les Pycnogonides provenant des campagnes du yacht "l'Hirondelle" [Gulf of Gascoyn, Azores, &c.]. Bull. Soc. Z. Fr. xvi, p. 176.

No new species.

#### SYSTEMATIC.

- Ammothea wilsoni, Isl. Chonas, Schimkewitsch, Atti Soc. Rom. Mém. (4) vi, p. 336, figg. 16-23, n. sp.
- Ascorhynchus tenuirostris, Torres Strait, p. 555, pl. xxii, figg. 7-14, Carpenter, Sci. P. R. Dubl. Soc. (n.s.) vii, n. sp.
- Pallenopsis fluminensis, Koroy, Abrolhos I., p. 339, figg. 29-31, Schimke-Witsch, Atti Soc. Rom. Mém. (4) vi.
- Parapallene, n. g. for Pallene australiensis, Hoek, and P. haddonii, n. sp., Torres Straits; Carpenter, p. 553, pl. xxii, figg. 1-6, Sci. P. R. Dubl. Soc. (n.s.) vii.
- Phoxichilidium longicolle, Dohrn, Porto Lagunas, Schimkewitch, p. 343, Mém. R. Linc. (4) vi.
- Pycnogonum stearnsi, California, IVES, P. Ac. Philad. 1892, pp. 142-144, pl. x, n. sp.
- Rhopalorhynchus clavipes, Torres Straits, CARPENTER, p. 24, pl. ii, figg. 1-10, Sci. P. R. Dubl. Soc. (n.s.) viii, n. sp.
- Tanystylum dohrni, Abrolhos I., p. 329, figg. 1-4, calicirostris, Gulf of Panama, p. 331, figg. 5-7, chierchiæ, Chonos, p. 333, figg. 8-11, Schimkewitsch, Atti Soc. Rom. Mém. (4) vi, n. spp.

# GIGANTOSTRACA.

# (A.) LIMULUS.

- Bernard, H. M. The Apodidæ: a Morphological Study. London: Macmillan & Co., 1892.
- Bouvier, E. L. Observations sur l'anatomie du système nerveux de la Limule polyphème (*Limulus polyphemus*). Bull. Soc. Philom. (8) iii, pp. 187-198.
- KINGSLEY, J. S. The Embryology of Limulus. J. Morph. vii, No. 1, pp. 35-66, pls. v & vi.
- KISHINOUYE, K. On the Development of Limulus longispina. J. Coll. Sci. Japan, v, pt. 1, pp. 53-100, 7 pls.
- SMITH, H. M. Notes on the King Crab Fishery of Delaware Bay. Bull. U. S. Fish. Comm. ix, pp. 363-370.
- VIALLANES, M. H. Études Histologiques et Organologiques sur les centres nerveux et les organes des sens des animaux articulés.

  6. Le cerveau de la Limule (L. polyphemus). Ann. Sci. Nat. xiv, pp. 405, &c.

# (B.) TRILOBITES.\*

- COLE, J. A. The Story of Olenellus. Nat. Science, i, No. 5, pp. 340, &c.
- ETHERIDGE, R., & MITCHELL, J. The Silurian Trilobites of New South Wales, with references to those of other parts of Australia. P. Linn. Soc. N.S.W. vi, pp. 311-320, pl. xxv.
- Proetus bowningensis, Mitch., p. 313, fig. 1; rattei, n. sp., p. 316, fig. 2; australis, n. sp., p. 318, fig. 3.
- GÜRICH, —. Ueber eine Cambrische Trilobiten-Fauna bei Sandomir. J. Ber. Schles. Ges., No. 69, 1891, p. 55.
- MATTHEWS, E. F. Notes on Cambrian Fauna. Can. Rec. v, pp. 247-258.
- The following Trilobites described:—Protolenus, Matth., p. 248, paradoxoides, fig. 1; elegans, fig. 2; zoppii, fig. 3. Ellipsocephalus galeatus, n. sp., p. 249, fig. 4; articephalus, Matth., pp. 250-252, fig. 5; hoffi, fig. 6.
- POMPECKI, J. F. Bemerkungen über das Einrollungsversmögen der Trilobiten. JH. Ver. Württ. xlviii, pp. 93-101.

<sup>\*</sup> No special attention has been devoted to this Section.

# XI. ARACHNIDA.

BY

# R. I. Pocock.

#### ARRANGEMENT OF RECORD.

A few changes have been made in the treatment of this section of the The first part contains a complete list of the ZOOLOGICAL RECORD. titles. The second is devoted to papers treating of the anatomy, embryology, physiology, habits, &c. The third part deals similarly with distribution, and the fourth with the purely systematic aspect of the subject. In the latter the sequence of the orders adopted is the same as that put forward in the Ann. N. H. for Jan., 1893; while the families of the larger groups, such as the Aranew and the Acari, and the genera that they contain, have been, for the sake of convenience, arranged alphabetically.

For reasons already stated under the heading Crustacea, the Pantopoda and the Gigantostraca have been transferred from the Arachnida to that section of the RECORD which deals with the Crustacea.

I. TITLES OF PAPERS, p. 1.

II. ANATOMY, EMBRYOLOGY, PHYSIOLOGY, HABITS, &c., p. 11.

III. DISTRIBUTION, p. 13.

IV. SYSTEMATIC

(A.) Scorpiones, p. 14.

(B.) Pedipalpi, p. 14.

(C.) Araneæ, p. 14. (D.) Pseudoscorpiones, p. 33. (E.) Opiliones, p. 34.

(F.) Acari, p. 35.

# I. TITLES OF PAPERS.\*

Notes on the Fauna of Jamaica. Johns Hopk. Andrews, E. A. Univ. Circ. xi.

Arachnida, p. 75.

Balzan, L. Voyage de M. E. Simon au Venezuela: Chernetes (Pseudoscorpiones). Ann. Soc. Ent. Fr. lx, pp. 497-552, pls. 9-12.

<sup>\*</sup> An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

<sup>1892. [</sup>vol. xxix.]

- BANKS, N. (1) Mimiery in Spiders. P. E. Soc. Wash. ii, No. 2, pp. 174-176.
- ---. (2) A new genus of Phalangiida. T. c. pp. 249 & 250.
- —. (3) On Prodidomus rufus, Hentz. T. c. pp. 259-261.
- ——. (4) The Spider Fauna of the Upper Lake, Cayuga Basin. P. Ac. Philad. 1892, pt. 1, pp. 11–81.
- ---. (5) Notes on the *Dysderida* of the United States. Canad. Ent. xxiii, pp. 207-209.
- Batelli, A. Notes anatomofisiologiche sugli *Ixodini*. Bull. Soc. Ent. Ital. xxii, pp. 218-235.
- BECKER, L. Sur le *Pholcus phalangioides*, Fuesely, 1775. Ann. Soc. Ent. Belg. xii, pp. 539 & 540.
- Berlese, A. Acari, Myriopoda et Scorpiones hucusque in Italia reperta. Pts. lxi-lxx, with a special unnumbered part devoted to the Acari of the group Mesostigmata. Padova, 1892.
- BERNARD, H. M. (1) Are the Solpugidæ poisonous? Nature, xlvi, p. 223.
- —. (2) The Apodida, a Morphological Study. Nature Series. London (Macmillan & Co.): 1892.
- ---. (3) Some Observations on the Relation of the Acarida to the Arachida. J. L. S. xxiv, pp. 279-291, pl. xx.
- ----. (4) An endeavour to show that the trachese of the Arthropoda arose from setiparous sacs. Zool. Jahrb. v, p. 511.
- BERTKAU, P. (1) Ueber Sinnesorgane in den Tastern und dem ersten Beinpaare der Solpugiden. Zool. Anz. xv, p. 10.
- ——. (2) Nachträgliche Bemerkungen zu meiner Mittheilung über Sinnesorgane in den Tastern und dem ersten Beinpaare der Solpugiden. T. c. p. 110.
- —... (3) Zu J. Lebedinsky "Die Entwicklung der Coxaldrüse bei Phalangium." 1. c. p. 177.
- —. (4) Ueber das Chiracanthium nutrix. SB. Ver. Rheinl. 1892, p. 102.
- ——. (5) Bau der Giftdrüse einheimischer Spinnen. CB. Ver. Rheinl. 1892, pt. 1, pp. 59 & 60.
- Birula, —. Beiträge zur Kenntniss der anatomischer Baues der Geschlectsorgane bei den Galeodiden. Biol. Centralbl. xii, p. 687.
- BLANCHARD, R. [Remarks upon the observations of F. Heim upon the red pigment of *Trombidium*, cf. Heim (2).] Bull. Soc. Ent. Fr. 1892, p. l.
- Bolivar, -. Sociabilidad de las arañas. Act. Soc. Esp. xxi, p. 22.

- <sup>o</sup>Brandes, G. Ueber die *Argasida*. CB. Ver. Sachr. Thür. 1892, pp. 17-19 & 22.
- Buysson, H. [Une rectification (Chorioptes doit être substitué à Symbiotes).] Bull. Soc. Ent. Fr. 1891, pp. clx & clxi.
- CAMBOUÉ, R. P. (1) Aranéides utiles e nuisibles. Bull. Soc. Acclim. 1887, pp. 1-10.
- —. (2) La soie d'Araignée. Rev. Sci. Nat. Appliq. 1892, No. 6, pp. 1-8.
- —. (3) Sur l'atilisation de deux Aranéides séricigènes de Madagascar. Op. cit. 1889, No. 5, pp. 1-3.
- CAMBRIDGE, O. P. (1) On a new Spider from Calcutta. Ann. N. H. (6) x, p. 417, pl. xxii.
  - A species of Ariannes.
- —... (2) On the British Species of False-Scorpions. P. Dorset Field Club, xiii, p. 199, &c.
- CAMBRIDGE, F. O. P. (1) British Spiders. Brit. Nat. (Supp.) pts. xiv, xv, xvii, xx, & xxii.
  - Drassus, Guaphosa, and Clubiona.
- ---. (2) New and Obscure British Spiders. Ann. N. H. (6) x, pp. 384-397, pls. xx & xxi.
- CANESTRINI, E. (1) Sopra due nuove specie di Fitoptus. Bull. Soc. Ven.-Trent. v, No. 2, pp. 79 & 80.
- (2) Sopra due nuove specie di Fitoptus. Op. cit. xii, p. 377.
- (3) Sopra due nuovi Fitoptidi. T. c. p. 379.
- ---. (4) Sopra tre nuove specie di *Fitoptidi* Italiani. Atti Ist. Venet. l, pp. 837-839 (1891-92).
- —. (5) Sopra due nuove specie di Fitoptidi Italiani. Op. cit. li, pp. 5 & 6 (1892-93).
- CARLINI, A. DE. Artropodi di Val Vigazzo. Bull. Soc. Ent. Ital. 1892, pt. 1: Aruneæ and Opiliones, pp. 90 & 91.
- CARPENTER, G. H. (1) The Walk of Arthropods. Nat. Sci. i, p. 676, &c.
- —... (2) Notes on the Development and Structure of Arachnids. Op. cit. No. 7, pp. 522-527.
- CAUSARD, M. Sur la circulation du sang chez les jeunes Araignées. C.R. cxiv, p. 1035.
- \*CHYZER, C., & KULCZYNSKI, L. Araneæ Hungariæ, &c. Berlin (R. Friedländer): 1892.
- CHYZER, C. Ueber eine neue Spinnenfauna Ungarns. Math. Nat. Ber. Ung. x, pp. 108-118.
  - Notes upon synonyms, &c., of certain species described in the above cited work on Hungarian Spiders, by Chyzer & Kulczynski.

#### XI. ABACHNIDA.

- Costa, A. Sugli effetti del veleno della Scorpione Tunisino (Buthus tunstanus) nell'uomo. Rend. Acc. Nap. (2) vi, pp. 144-146.
- •CURTICE, C. The Biology of the Cattle Tick. J. Comp. Med. Vet. Arch. 1891, pp. 313-319.
- CURTIS, J. S. A New Jumping Spider. Zoe, iii, pp. 332-337.
- Damin, N. (1) O pauku: *Uloborus walckenaeri* (Latr.). Glasnik Naravosl. Druzt. vi, pp. 220-223, pl. iv.
- —. (2) Iz zivota pauku Argiope bruennichi (Scop.). Op. cit. vii, pts. 1-6, pp. 211-220, pl. vi (1892).
- DISTANT, W. L. Are the Solpugida poisonous? Nature, xlvi, p. 247.
- Du Roselle, —. Contributions a la Faune locale—Arachnides. Bull. Soc. L. Nord. France, x, pp. 340-343 (1890-91).
- ENOCK, F. Additional Notes and Observations on the Life-history of Atypus piceus. Tr. E. Soc. 1892, pp. 21-26.
- FLOGEL, J. H. L. Ueber das Nervensystem von Demodex folliculorum. Verh. Ges. Nat. Ärzte, lxiv, pp. 120 & 121.
- FOCKEU, H. Étude sur quelques Galles de Syrie. Rev. Biol. iv, pp. 152-160 & 231-234.
- Fox, W. H. A List of the Spiders from Indiana. P. E. Soc. Wash. ii, No. 2, p. 224.
- GARMAN, H. (1) The Origin and Development of Parasitism among the Sarcoptidæ. Ins. Life, iv, pp. 182-187.
- . (2) American Phytoptocecidii. Psyche, vi, pp. 241-246, 1 pl.
- GAUBERT, P. (1) Sur les muscles des membres et sur l'homologie des articles des Arachnides. Bull. Soc. Philom. (8) iv, No. 1, pp. 31-33.
- —. (2) Sur un ganglion nerveux des pattes du Phalangium opilio. C.B. exv, p. 960.
- —... (3) Sur l'autotomie choz les Araignées. Bull. Soc. Philom. (8) iv. pp. 78 & 79.
- —. (4) Observations sur les pièces buccales des Aranéides. T. c. pp. 80 & 81.
- ——. (5) Recherches sur les organes des sens et sur les systèmes tégumentaires, glandulaires, et musculaires des appendices des Arachnides. Ann. Sci. Nat. xiii, pp. 1–184, 4 pls.
- Göldi, E. A. Zur Orientierung in der Spinnenfauna Brasiliens. MT. Osterlande, v, pp. 200-248.
- GOYEN, P. On New Zealand Araness. Tr. N. Z. Inst. xxiv, pp. 253-257.
- GREEN, E. E. Remarkable Weapons of Defence. Nature, xlvii, p. 199. Distastefulness of Holothyrus coccinsila.

- Grevé, C. Beobachtungen an zwei lebenden Arthropoden aus Mittel-Amerika. Zool. Jahrb. vi, p. 461.
  - On a Scorpion (Centrurus biaculeatus).
- HASSELT, A. W. M. VAN. (1) Waarnemingen omtrent de biologie van Cælotes atropos, Walck. Tijdschr. Ent. xxxiii, pp. xiv-xvi.
- ——. (2) Over de parasieten der Spinnen. Op. cit. xxxv, pp. xxxviiixli.
- HEIM, F. [Sur le pigment de la larve de Trombidium fuliginosum.] Bull. Soc. Ent. Fr. 1892, pp. xlix-l.
- Howard, L. O. Hymenopterous Parasites of Spiders. P. E. Soc. Wash. ii, No. 3, pp. 290, &c.
- IMHOF, O. E. Die Zusammensetzung der pelagischen Fauna der Süsswasserbecken. Biol. Centralbl. xii, pp. 171, &c.
- JAWOROWSKI, A. (1) Ueber die Extremitäten, deren Drüsen und Kopfsegmentierung bei Trochosa singoriensis. Zool. Anz. xv, p. 197.
- Carachiden und Insekten. Kosmos Lemberg, 1891, ii, pp. 1-15, 4 figures.
- Johnson, H. P. Amitosis in the Embryonal Envelopes of the Scorpion. Bull. Mus. C. Z. xxii, pp. 127-160.
- JONES, K. H. A few Remarks on our Commonest Spiders. Sci. Goss. 1892, pp. 73-75.
- JOURDAIN, S. Sur le mode de fixation des larves parasites hexapodes des Acariens. C.R. cxv, p. 621.
- KAPKA, J. Die Fauna des Böhmischen Teiche. Arch. naturw. Landesforsch. Böhmen, viii, No. 2, pp. 1-115.

  Hydrachnidæ, pp. 107-109.
- KARPELLES, L. Ueber einen eigenthümlichen Parasiten der Krontaubo. SB. z.-b. Wien, xlii, pp. 46 & 47.
- KARSCH, F. (1) Arachniden von Ceylon und von Minikoy gesammelt von den Herren Doctoren P. und F. Sarasin. B. E. Z. xxxvi, pt. ii, pp. 267-310, 3 pls.
- —... (2) Ueber Cryptostemma, Guér., als einzigen recenten Auslaüfer der fossilen Arachnoideen-Ordnung der Meridogustra, Thor. Op. cit. xxxvii, pp. 25-32, pl. iv.
- —. (3) Nachschrift zu meinem Aufsatze über Cryptostemma, Guér. T. c. p. 64.
- Kellicott, D. S. Hypoderas columbæ: a Note. Ins. Life, v, pp. 77 & 78.
- Kennel, J. v. Dei Verwandtschaftsbeziehungen und die Abstammung der Tardigraden. SB. Ges. Dorp. 1891, pp. 504-512.

- KEYSERLING, E. (1) Die Spinnen Amerikas. Brasilianische Spinnen. Nürnberg (Bauer & Raspe): 1891.
  - Not seen by Recorder in 1891.
- •—. (2) Die Spinnen Amerikas—Epeiridæ. Nürnberg (Bauer & Raspe): 1892, pt. iv, No. 1.
- KIEFFER, J. J. Des Acarocécidies de Lorraine. Feuill. Nat. xxii, pp. 97-104, 118-129, 141-147, & 162-165.
- Kishinouyž, K. On the Lateral Eyes of Spiders. J. Coll. Sci. Japan, v, pp. 101-103.
- KCENIKE, F. (1) Anmerkungen zu Piersig's Beitrag zur Hydrachnidenkunde. Zool. Anz. xv, p. 263.
- ——. (2) Zwei neue Hydrachniden-Gattungen aus dem Rhätikon. T. c. pp. 320 & 325.
- KOJEVNIKOV, G. La faune de la mer Baltique orientale et les problèmes des explorations prochaines de cette faune. Congr. Internat. Zool. 1892, pt. i, pp. 132, &c.
- KOVALEVSKY, A. Sur les organes excréteurs des Arthropodes terrestres. T. c. pp. 187, &c.
  - Aranew, p. 216; Scorpiones, p. 228.
- Kramer, P. Zur Entwicklungsgeschichte und Systematik der Süsswassermilben. Zool. Anz. xv, p. 149.
- LAURIE, M. On the Development of the Lung-books of Scorpio fulvipes. Zool. Anz. 1892, No. 386, pp. 101-105.
- LAVERAN, A. Acariens de l'oreille chez le lapin, paraphlégie reflexe. C.R. Soc. Biol. 1892, p. 169.
- LEBEDINSKY, J. Die Entwicklung des Coxaldrüse bei *Phalangium*. Zool. Anz. xv, p. 131.
- Lewis, R. T. (1) Notes on the process of Oviposition as observed in a species of Cattle Tick. J. R. Micr. Soc. 1892, No. 4, pp. 449-454, 1 pl.
- —. (2) Note on a species of *Ixodes*. J. Quek. Club (2) v, pp. 10-14, 1 pl.
- MARCHAL, P. La glande coxale du Scorpion et ses rapports morphologiques avec les organes excréteures des Crustacés. C.R. cxv, p. 191. [Translated in Ann. N. H. (6) x, p. 338.]
- Marx, G. (1) A contribution to the Study of the Spider Fauna of the Arctic Regions. P. E. Soc. Wash. ii, No. 2, p. 186.
- —. (2) Notes on the Classification of the Ixodida. T. c. p. 232.
- —. (3) Contributions to the Knowledge of the Life-history of Arachnida. T. c. p. 252.
  - Observations on Thelyphonus and Datames.

- [Marx, G.] (4) [Remarks upon the Distribution of *Drasside* and *Attidee*.]

  T. c., No. 2, p. 251.
- —... (5) A list of the Aranea of the District of Columbia. T. c. p. 148.
- —. (6) On the Morphology of the Ticks. Op. cit. ii, No. 3, pp. 271-287.
- MAZZA, F. Contribuzione all anatomia macro- e microscopica del Pentastomum moniliforme, Dies. Atti Univ. Genova, 1891, pp. 1-45.
- McCook, H. C. Drezelia, a new genus of Spiders. P. Ac. Philad. 1892, p. 127.
- MEADEN, C. W. Bite of the Tarantula (Mygale) Spider. Trinidad Nat. Club, i, No. 5, pp. 127 & 128.
- MEGNIN, P. (1) Acariens des oreilles chez le chat, le furet, et lo chien. C.R. Soc. Biol. 1892, p. 125.
- ——. (2) Un dernier mot sur la question de l'épilepsie de nos Carnassiers domestiques. T. c. p. 142.
- C. (3) Les Acariens parasites. Paris (Gauthier-Villars et fils): 1892. (Cf. Zool. Anz. xvi, p. 54.)
- MICHAEL, A. D. (1) On the variations in the internal anatomy of the Gammarina, especially in that of the genital organs, and in their mode of coition. Tr. L. S. v, pt. 9, pp. 281-324.
- —. (2) Ticks. Nat. Sci. i, No. 3, pp. 200-204.
- MONIEZ, R. (1) Note sur l'Arrhenurus decurtator, n. sp. Rev. Biol. iv, p. 520.
- —... (2) Mémoire sur quelques Acariens et Thysanoures parasites ou commensaux des Fourmis. T. c. pp. 377-389.
- —... (3) Contribution à l'histoire naturelle du Tyroglyphus mycophagus, Mégn. Mém. Soc. Zool. v, pt. 4, pp. 584-601.
- MORIN, J. Studien über Entwicklung der Spinnen. (Printed in Russian.) Zapiski Novoross. Obsch. Estestv. 1888, pp. 1-112, 4 pls.
- Mosley, S. L. Yorkshire Galls. Naturalist, 1892, pp. 273-278.
- NALEPA, A. (1) Neue Gallmilben. SB. Ak. Wien, 1892, p. 16.
- ---. (2) Neue Gallmilben. T. c. p. 128.
- —... (3) Neue Arten der gattung Phytoptus, D.:j., und Cecidophyes, Nal. T. c. p. 155.
- —. (4) Neue Gallmilben. T. c. pp. 190-192.
- —. (5) Tegonotus, ein neues Phytoptiden genus. Zool. Jahrb. vi, p. 327, 1 pl.
- PECKHAM, G. W. & E. G. Ant-like Spiders of the family Attida.

  Vol. II. of the Occasional Papers of the Nat. Hist. Soc. Wisconsin.

  Milwaukee: 1892, pp. 1-83, pls. i-vi.

#### XI. ARACHNIDA.

- Piersig, R. (1) Beiträg zur Hydrachnidenkunde. Zool. Auz. xv, p. 151.
- —. (2) Beiträge zur Kenntniss der im Susswasser lebenden Milben. T. c. pp. 338 & 341.
- —. (3) Eine neue Hydrachniden-gattung aus dem sächsischen Erzgebirge. T. c. p. 408.
- POCOCK, R. I. (1) Report upon a small collection of Scorpions sent to the British Museum by Mr. Edgar Thurston, of the Government Central Museum, Madras. J. A. S. (Bombay) vii, No. 3, pp. 295-312.
- ——. (2) Descriptions of two new genera of Scorpions, with Notes upon some species of *Palamnœus*. Ann. N. H. (6) ix, pp. 38-49, pl. iiib.
- ——. (3) Description of a new Trap-door Spider from Ceylon. T. c. pp. 49-51.
- —. (4) Supplementary Notes on the Arachnida and Myriopoda of the Mergui Archipelago, &c. J. L. S. xxiv, pp. 316-318.
- —... (5) Liphistius and its bearing upon the Classification of Spiders. Ann. N. H. (6) x, pp. 306-314.
- ——. (6) Arachnida and Myriopoda in W. L. Distant's Naturalist in the Transvaal. London (R. H. Porter): 1892, pp. 179-184.
- —. (7) On the . . . . . . Arachnida collected by Dr. Anderson in Algeria and Tunisia. P. Z. S. 1892, pt. 1, pp. 24–28.
- PUGA-BORRE, F. El Lathrodectus formidabilis de Chile. Act. Soc. Sci. Chili, ii, pt. 1, pp. 3-176, & pt. 2, pp. 271-296.
- Purcell, F. (1) Bau der Phalangiden-augen. Verh. deutsch. Zool. Ges. 1892, pp. 141 & 142.
- —... (2) Ueber den Bau und die Entwicklung der Phalangiden-augen. Zool. Anz. xv, p. 461.
- RAILLIET, A., & CADIOT, —. Observations et expériences sur l'otocariase symbiotique des Carnassiers. C.R. Soc. Biol. (9) iv, No. 5, pp. 104-108.
- RAILLIET, A. (1) Simples remarques historiques sur l'otocariase des Carnivores. Op. cit. 1892, p. 126.
- —. (2) Recherches sur la transmissibilité de la gale du Chat et du Lapin due au Sarcoptes minor, Fürst. Op. cit. (9) iv, No. 4, pp. 315-319.
- BATZ, S. VON. Von der aktiven Wanderung der Pentastomum denticulatum. CB. Bakt. Parasit. xii, pp. 329-333.
- Roselle, Du. [See Du Roselle.]

- SCHLECHTENDAL, D. von. (1) Die Gallbildungen deutscher Gefässpflanzen. J. Ber. Ver. Zwickau, 1891 [1892], pp. 1-10.
- ---. (2) Ueber das Eierlegen der Ixodes Weibchen. T. c. pp. 11-14.
- Schneider, A. Mélanges Arachnologiques. Pt. 1. Sur l'endothélium des vaisseaux. Pt. 11. Sur le poumon des Aranéides, 3 pls. Pt. 111. Sur le système artèriel du Scorpion. Pt. 1v. Système artèriel des Aranéides. Tabl. Zool. ii, pp. 135–198.
- SIMON, E. (1) Liste des Arachnides recueillis en Syrie par M. le docteur Barrois. Rev. Biol. v, pp. 80-84.
- —. (2) [Descriptions de deux Arachnides nouveaux du Portugal.]
  Bull. Soc. Ent. Fr. 1892, pp. exciv & exev.
- ——. (3) Liste des espèces de la famille des Aviculariides qui habitont l'Amerique du Nord. Appendice. Liste des Aviculariides qui habitent le Mexique et l'Amerique centrale. Act. Soc. L. Bord. xliv, pp. 307-339.
- —. (4) Descriptions d'espèces et de genres nouveaux de la famille des Aviculariidæ. Ann. Soc. Ent. Fr. lxi, pp. 271-284.
- —. (5) Arachnides des Iles Philippines. T. c. pp. 35-52.
- ——. (6) Histoire naturelle des Araignées. Vol. i, pt. 1, pp. 1-254. Paris : 1892.
- --- (7) Arachnides d'Islande. Bull. Soc. Ent. Fr. lx, p. clxxvi.
- SLINGERLAND, M. V. The Pear-leaf Blister Mite (*Phytoptus pyri*). Ins. Life, v, No. 2, pp. 104-106.
- SMALIAN, —. Argas reflexus in Aschersleben. CB. Ver. Sachs. Thür. 1892, pp. 18, 19, 29, & 30.
- SPENCER, W. BALDWIN. The Anatomy of Pentastomum teretiusculum, Baird. Q. J. Mier. Sci. xxxiv, pp. 1-73, 9 pls.
- STILES, C. W. The Histology of Boophilus. Abstract. P. E. Soc. Wash. ii, No. 3, pp. 303 & 304.
- Stoll, Otto. Arachnida acaridea in Biologia Centrali-Americana. Godman & Salvin, London, March, 1890; Jan. 1893.
- Diagnoses of new species, with list of the species of Acari hitherto described from Mexico and Central America.
- STRUBELL, A. Zur Entwicklungsgeschichte der Pedipalpen. Zool. Anz. xv, pp. 87 & 89. Trans, in Ann. N. H. (6) x, p. 419.
- THORELL, T. (1) Étude sui Ragni Malesi e Papuani. Part IV. Ann. Mus. Genov. xxxi, pp. 1-490, 1891-1892.
- —. (2) On some Spiders from the Andaman Islands, collected by E. W. Oates, Esq. Ann. N. H. (6) ix, pp. 226-237.

- [Thorell, T.] (3) Novæ species Aranearum a Cel. Th. Workman in ins. Singapore collectse. Bull. Ent. Ital. xxiv, pt. 3, pp. 211-282.
- TROUESSART, E. Considérations générales sur la classification des Acariens, suivies d'un essai de classification nouvelle (concluded). Rev. Sci. Nat. Ouest, ii, pp. 20-54.
- URQUHART, A. T. (1) Catalogue of the Described Species of New Zealand Araneidæ. Tr. N. Z. Inst. xxiv, pp. 220-230.
- ---. (2) Descriptions of New Species of Araneæ. T. c. pp. 230-253.
- Vejdovsky, J. F. (1) Sur la question de la segmentation de l'œuf et la formation du blastoderme des Pseudoscorpionides. Congr. Internat. Zool. ii, No. 1, pp. 120-125.
- ——. (2) Sur un organe embryonnaire des Pseudoscorpionides. T. c. pp. 126-131.
- WAGNER, J. Zur Entwicklungsgeschichte der Milben. Furchung des Eies, Entstehung der Keimblätter und Entwicklung der Extremitäten bei Ixodes. Zool. Anz. xv. p. 316.
- WAGNER, F. von. Biologische Notis. T. c. p. 434. Tipula and Pseudoscorpion.
- WARBURTON, C. Spiders from Madeira. Ann. N. H. (6), x, pp. 216-228, 1 pl.
- WEED, C. M. (1) The Ash-grey Harvest Spider. Am. Nat. xxvi, p. 32.
- ---. (2) The Striped Harvest Spider: a Study in variation. T. c. pp. 999-1008.
- —. (3) The Cattle Tick. T. c. pp. 530 & 531.
- ----. (4) Harvest Spider. Notes. T.c. p. 786.
- ---. (5) Protective Resemblance in Trombidium. T. c. p. 786.
- —. (6) Some Florida Spiders. T. c. p. 873.
- Wood, F. H. Notes on Spiders. J. Northampt. Soc., No. 51, pp. 81-91.
- WORKMAN, T. & M. E. Malaysian Spiders: pt. 1. Belfast (Ireland): 1892.
- Coloured figures, with drawings in outline, of eyes, sexual organs, &c., accompanied by brief descriptions of structure, form of snare, habits, &c.
- ZOGRAF, N. Sur l'origine et les parentes des Arthropodes, principalement des Arthropodes trachéates. Congr. Internat. Zool. 1892, pt. i, pp. 278, &c.

# II.—ANATOMY, EMBRYOLOGY, PHYSIOLOGY, HABITS, &c.

# ARACHNIDA (as a whole).

Notes on development and structure of Arachnida; CARPENTER .-Relationship between the Arachnida and other Arthropoda; BERNARD (2).—Relation of the Acari to other Arachnida; BERNARD (3).

Muscles of the limbs and homology of the limb segments in Arachnidu; GAUBERT (1).

Trachese, origin of; BERNARD (4).

Organs of touch and integumentary glands of Arachnida; GAUBERT (5).

#### SCORPIONES.

Excretory organs; KOVALEVSKY.—Arterial system; SCHNEIDER.

Observations on a living Scorpion; GREVÉ.

Poison of Buthus; Costa.

Amitosis in the embryonal envelopes; JOHNSON.—Development of the lung-books in Scorpio fulvipes; LAURIE.

Coxal gland; MARCHAL.

#### PEDIPALPI.

Habits of Thelyphonus; MARX (3).—Development of Thelyphonus; STRUBELL.

#### ARANEÆ.

Habits: Wood,—Remarks upon common Spiders; Jones.—Autotomy: GAUBERT (3).—Sociability; BOLIVAR.—Habits of Atypus; ENOCK: of Uloborus and Argiope; DAMIN.—On Lathrodectus formidabilis; PUGA-BORRE.—On Chiracanthium; BERTKAU (4).—Biology of Calotes atropos; HASSELT .- Notes on Pholcus; BECKER.

Mimicry: Banks (1).—Marptusa imitating flies; Goyen, p. 254.

Parasites of Spiders; HASSELT (2), HOWARD.

Utilisation of Spiders' silk; CAMBOUÉ (2 & 3).—Useful and hurtful Spiders; Camboué (1).

Classification of Araneæ; POCOCK (5), SIMON (6).

Development of Spiders; MORIN.—Development of the lateral cycs;

Segmentation of the head and the form of the appendages in young Trochosa; JAWOROWSKI.

Poison glands; BERTKAU (5), -Effects of the bite of Mygale; MEADEN.

#### XI. ABACHNIDA.

Arterial system; Schneider.—Circulation of blood in young Spiders; Causard.

Mouth parts; GAUBERT (4).—Excretory organs; KOVALEVSKY.—Lung-books; SCHNEIDER.

#### SOLIFUGÆ.

Habits of Datames; MARX (3).

Poison glands; BERNARD (1), DISTANT.—Sense organs; BERTKAU (1 & 2).—Generative organs; BIRULA.

#### PSEUDOSCORPIONES.

Pseudoscorpion and Tipula, Habits; WAGNER, F. Embryology; VEJDOVSKI.

#### OPILIONES.

Coxal glands of *Phalangium*; Bertkau (3), Lebedinsky.—Nervous system of the feet of *Phalangium*; Gaubert (2).—Eyes of *Phalangiida*; Purcell (1 & 2).

#### ACARI.

On Ticks; MICHAEL (2).—Morphology of Ticks; MARX (6).

Acari in Ants' nests; Moniez (2).—Life-history of the Cattle-Tick; Curtice.—Natural history of Tyroglyphus; Moniez (3).—The mode of fixation of the hexapodous larve of Acari; Jourdain.—Oviposition of Ixodes; Lewis (1), Schlechtendal (2).—Freshwater Mites; Piersig (1, 2, 3).—On Phytopius pyri; Slingerland.

Distastefulness of *Holothyrus*; GREEN.—Protective resemblances of *Trombidium*; WEED.

Pigment of Trombidium; HEIM, BLANCHARD.

Origin of parasitism in Sarcoptidæ; GARMAN (1).—Parasites of Pigeons; KARPELLES, KELLICOTT.—Effects of Acari in ears of Mammals; LAVERAN, MEGNIN (1 & 2), BAILLIET (1 & 2), RAILLIET & CADIOT.

Classification of Acari; TROUESSART: of Ixodidæ; MARX (2).

Histology of *Boopilus*; STILES.—Anatomy of *Gamasida*; MICHAEL (1).—Anatomy and physiology of *Ixodes*; BATELLI.—Nervous system of *Demodex*; FLOGEL.

Development of Leodes, WAGNER, J.: of freshwater Mites, KRAMER.

#### TARDIGRADA.

Affinites of Tardigrada; KENNEL.

#### PENTASTOMIDA.

Anatomy of Pentastomum; MAZZA. — Anatomy of Pentastomum teretiusculum; SPENCER.

Migration of Pentastomum ; RATZ.

# III.—DISTRIBUTION.

On the distribution of Attide and Drasside; MARX (4).

# Holarctic Region.

Arctic Regions: Spiders; MARX (1).—Iceland: Arachnida; SIMON (7).

Britain: Pseudoscorpiones; O. P. CAMBBIDGE (2), Aranea; F. O.

CAMBRIDGE.—Yorkshire: Galls; Mosley. France: Spiders; Du Roselle.

Lorraine: Phytoptus-galls; KIEFFER.

Germany: Phytoptus-galls; SCHLECHTENDAL (1).

Eastern Baltic: Halacari; KOJEVNIKOV.

Hungary: Spiders; CHYZER.—Bohemia: Hydrachnida; KAFKA.
Italy: Phytoptidæ; CANESTBINI (1-5), Araneæ and Opiliones; CARLINI.

Portugal: Spiders; SIMON (2).

Algeria and Tuni sia: Arachnida; Pocock (7).

Madeira: Spiders; WARBURTON.

Syria: Phytoptid-galls: FOCKEU.—Arachnida: SIMON (1).

N. America: Aviculariida; Simon (3).—Phytoptid-galls; Garman (2).—Opiliones; Weed (1, 2, & 4).

Columbia: Spiders; MARX (5).

Cayuga Lake: Spiders; Banks (4). United States: Dysderidæ; Banks (5).

Indiana: Spiders; Fox. Florida: Spiders; WEED (6).

# Neotropical Region.

Central America and Mexico: Aviculariida; SIMON (3): Acari, STOLL.

Jamaica: Andrews.

Venezuela: Pseudoscorpiones; BALZAN. Brazil: Spiders; GÖLDI, KEYSERLING.

# Australian Region.

New Zealand: Spiders; URQUHART (1 & 2), and GOYEN.

# Oriental Region.

India: Scorpions; POCOCK (1).—Ceylon: Arachnida; KARSCH (1).—Andaman Is.: Spiders; THORELL (2).—Mergui Archipelago; Arachnida; POCOCK (4).—Singapore: Spiders; THORELL (3).—Malaysia: Spiders; WORKMAN.—Malaysia and Papua: Spiders; THORELL (1).—Philippine Islands: Arachnida; SIMON (5).

# Ethiopian Region.

Transvaal: Arachnida; POCOCK (6).

#### IV.—SYSTEMATIC.

## (A.) SCORPIONES.

- Isometrus thurstoni, Madras, Sheveroy Hills, pp. 297-300, Pocock, J. A. S. B. vii, n. sp.
- Archisometrus scaber, n. sp., Madras, with notes upon tricarinatus, Simon, and shoplandii, Oates, pp. 300-303; Pocock, J. A. S. B. vii.
- Buthus martensii, Karsch, ? syn. B. tamulus, Fabr., p. 303; Россск, J. A. S. B. vii.
- B. tadmorensis, Palmyra, p. 84, Simon, Rev. Biol. v, n. sp.
- Heterocharmus, n. g., p. 46, for H. cinctipes, n. sp., ? India or Ceylon, pp. 47 & 48, pl. iiis, figg. 2 & 2b; Pocock, Ann. N. H. (6) ix.
- Stenochirus, n. g., of Buthidæ, p. 305, for S. sarasinorum, n. sp., Peradenia, p. 306, pl. xii, fig. 30; Karsch, B. E. Z. xxxvi.
- Hormurus læviceps, Pocock, Tranquebar and Sheveroy Hills, discussed; Pococκ, pp. 303 & 304, J. A. S. B. vii.
- Cheloctonus, n. g., for O. jonesii, n. sp., Transvaal, pp. 44-46, pl. iiiB, fig. 1; Рососк, Ann. N. H. (6) ix.
- Palamneus spinifer, Hempr. & Ehrb., syn. petersii, Thorell, 1876, from Calcutta, Singapore, &c., and thorellii, n. sp., syn. bengalensis, Sim., and petersii, Thorell, 1889, from Burma, pp. 38-43; Pocock, Ann. N. H. (6) ix.
- Scorpio swammerdami, Simon, syn. S. afer, Koch, p. 304; fulvipes, Koch, ? syn. afer, Herbst, described, pp. 304-307; phipsoni, n. sp., Madras and Sheveroy Hills, pp. 307-309; cæsar, C. Koch, scaber, Thorell, bengalensis, C. Koch, indicus, Linn., and humilis, Simon, discussed, pp. 309-312; Pocock, J. A. S. B. vii.

#### (B.) PEDIPALPL

#### PEDIPALPI, AMBLYPYGI.

- According to Simon, Ann. Soc. Ent. Fr. lxi, pp. 45-52, this suborder contains the single family Tarantulidæ, which is divided into the Charontinæ, Phryniscinæ, and Tarantulinæ. The Charontinæ contain Charon, Catageus, and the new genera Charinus, based upon australianus, L. Koch, and Sarax, of which brachydactylus, Simon, is the type, p. 48. The Phryniscinæ contain Phryniscus and Damon, and the Tarantulinæ, Admetus and Tarantula, p. 51, pl. ii, figg. 9-16.
- Surax, n. g., p. 43, for S. brachydactylus, n. sp., in Cave, Luzon; Simon, Ann. Soc. Ent. Fr. lxi.

#### (c.) ARANEÆ.

#### Agalenidæ.

Camptostichomma, n. g., p. 295, for C. manicatum, n. sp., Ceylon, p. 296, pl. xii, fig. 19; Karsch, B. E. Z. xxxvi.

- Cicurina creber, p. 26, pl. i, fig. 76, placida, p. 27, fig. 77, Cayuga Lake, Banks, P. Ac. Philad, 1892, n. spp.
- Celotes longitarsus, Em., p. 24, pl. iv, fig. 73; Banes, P. Ac. Philad. 1892.
- C. fidelis, p. 24, pl. 1, fig. 72, altilis, p. 25, pl. i, fig. 74, & pl. iv, fig. 74, lineatus, p. 25, gnavus, p. 26, Cayuga Lake, BANKS, P. Ac. Philad. 1892, n. spp.
- Cybœus giganteus, Cayuga Lake, BANKS, P. Ac. Philad. 1892, p. 23, pl. 1, fig. 71, n. sp.

#### ATTIDE.

- Admestina wheeleri, Peck., p. 78, Cayuga Lake, Banks, P. Ac. Philad. 1892.
- Attus coronatus, Madagascar, Cambour, p. 4, Bull. Soc. Acclim. 1887;

  A. maderiana, Madeira, p. 217, pl. xiv, fig. 1, Warburton, Ann. N. H.

  (6) x: n. spp.
- Bathippus, n. g., for B. macilentus, n. sp., Sumatra, pp. 402-406; THORELL, Ann. Mus. Genov. xxxi.
- Bellota, n. g., p. 67, formica, Tacz., Venezuela, Peru, p. 68, pl. v, fig. 5; Peckham, Nat. Hist. Soc. Wisconsin, ii.
- Bianor leucosticius, pp. 252-256, balius, pp. 256-259, incitatus, pp. 259-262, Sumatra, Thorell, Ann. Mus. Genov. xxxi, n. spp.
- Bindax, n. g. for B. chalcocephalus, Thor., p. 411; THORELL, Ann. Mus. Genov. xxxi.
- Bocus, n. g., p. 38, for B. excelsus, n. sp., Luzon, p. 39, pl. iii, fig. 4; Peckham, Nat. Hist. Soc. Wisconsin, ii.
- Bootes ephippiger, Sim., Sumatra, described, pp. 375-379; Thorell, Ann. Mus. Genov. xxxi.
- Chrysilla, n. g., p. 315, for C. delicata, n. sp., p. 316; THORELL, Ann. Mus. Genov. xxxi.
- C. debilis, p. 319, dories, p. 321, Sumatra, Thorell, Ann. Mus. Genov. xxxi, n. spp.
- Cocalus ramipalpis, Sumatra, p. 353, THORELL, Ann. Mus. Genov. xxxi, n. sp.
- Carrhotus, n. g. for C. viduus, C. Koch, Java, p. 407; THORELL, Ann. Mus. Genov. xxxi.
- Dendryphantes insignis, pl. v, fig. 28, ornatus, pl. iv, fig. 29, & pl. v, fig. 29, exiguus, pl. v, fig. 30, pp. 74 & 75, Cayuga Lake, Banks, P. Ac. Philad. 1892; D. ansolus, California, pp. 332-337, Curtis, Zoe, iii: n. spp.
- Descanso, n. g., p. 72, for D. ragus, n. sp., Santarem, p. 72, pl. v, fig. 9; PECKHAM, Nat. Hist. Soc. Wisconsin, ii.
- D. chapado, Brazil, p. 73, pl. vi, fig. 1, PECKHAM, Nat. Hist. Soc. Wisconsin, ii, n. sp.
- Dischiropue, n. g., p. 299, for D. alticinctus, n. sp., Ceylon, p. 300, pl. xii, fig. 26; KARSCH, B. E. Z. xxxvi.

- Emertonius, n. g., p. 54, for E. exasperans, n. sp., Java, p. 54, pl. iv, fig. 8; PECKHAM, Nat. Hist. Soc. Wisconsin, ii.
- Epocilla, n. g., p. 309, for E. pratextata, n. sp., Java, p. 310; THORELL, Ann. Mus. Genov. xxxi.
- Erica, n. g., p. 55, for E. eugenia, n. sp., Brazil, p. 55, pl. iv, fig. 9; PECK-HAM, Nat. Hist. Soc. Wisconsin, ii.
- Euophrys giebelii, Java, pp. 357-360, Thorell, Ann. Mus. Genov. xxxi, n. sp.
- E. pygæa, Thor., figured, pl. 6; Workman, Malaysian Spiders, pt. 1.
- Fluda, n. g., p. 57, for F. narcissa, n. sp., Brazil, p. 57, pl. iv, fig. 10; F. ruficeps, Tacz., Peru, p. 58, pl. iv, fig. 3; Peckham, Nat. Hist. Soc. Wisconsin, ii.
- Gelotia, n. g., pp. 344 & 345; G. frenata, Sumatra, pp. 345-348, bimaculata, Sarawak, pp. 348-351, Thorell, Ann. Mus. Genov. xxxi, n. spp.
- Hazarius scylax, pp. 413-417, simonis, pp. 418-423, workmanii, pp. 423-426, coprea, pp. 431-436, sobarus, pp. 436-440, Sumatra, dispalans, pp. 440-443, mccookii, pp. 443-449, Java, Thorell, Ann. Mus. Genov. xxxi; H. arcigerus, Ceylon, p. 302, pl. xii, fig. 29, Karsch, B. E. Z. xxxvi: n. spp.
- Harmochirus nervosus, Sumatra, pp. 246-250, Thorell, Ann. Mus. Genov. xxxi, n. sp.
- Hermosa, n. g., p. 53, for H. volatilis, n. sp., Madagascar, p. 53, pl. iv, fig. 7; PECKHAM, Nat. Hist. Soc. Wisconsin, ii.
- Homalattus latidens, Dol., Java, pp. 262-268; bufo, Dol., Sumatra, pp. 275-278, described; Thorell, Ann. Mus. Genov. xxxi.
- H. mordax, Java, pp. 268-275, nobilis, Celebes, pp. 278-282, THORELL, Ann. Mus. Genov. xxxi; H. laminatus, Dambulla, p. 298, pl. xii, fig. 23, reductus, p. 298, fig. 24, cingulatus, p. 299, fig. 25, Ceylon, Karsch, B. E. Z. xxxvi: n. spp.
- Hyllus giganteus, C. Koch, walcknaerii, White, Celebes, mutillarius, C. Koch, Sumatra, lacertosus, C. Koch, n. var., borneensis, Borneo, pp. 379-386; Thorell, Ann. Mus. Genov. xxxi.
- Icius formosus, p. 76, pl. v, fig. 31, næstus, p. 77, pl. v, fig. 33, Cayuga Lake, Banks, P. Ac. Philad. 1892; I. discatus, Ceylon, p. 301; Karsch, B. E. Z. xxxvi: n. spp.
- Iola, n. g., p. 75, for I. cowani, Madagascar, p. 75, pl. vi, fig. 3, Peckham, Nat. Hist. Soc. Wisconsin, ii, n. sp.
- Janigera melanocephala, C. Koch, Sumatra, pp. 224-239; Thorell, Ann. Mus. Genov. xxxi.
- Keyserlingella, n. g., p. 69, for K. perdita, New Grenada, p. 70, pl. v, fig. 7, cara, Guatemala, p. 71, pl. v, fig. 8, n. spp.; Peckham, Nat. Hist. Soc. Wisconsin, ii.
- Leptorchestes chrysopogon, Simon, Europe, p. 48, pl. iv, fig. 5, mutilloides, Lucas, Algiers, p. 49, berolinensis, C. Koch, Europe, p. 50; PECKHAM, Nat. Hist. Soc. Wisconsin, ii.

- L. georgius, Madagascar, p. 52, pl. iv, fig. 6, PECKHAM, Nat. Hist. Soc. Wisconsin, ii, n. sp.
- Mavia vittata, C. Koch, Penang, pp. 335-338; THORELL, Ann. Mus. Genov. xxxi.
- M. glaucochira, Sumatra, pp. 325-328, gemmans, Sumatra, pp. 329-332, marxii, Java, pp. 332-335, cypria, Java, pp. 339-341, laticlavia, Sumatra, pp. 341-343, Thorell, Ann. Mus. Genov. xxxi, n. spp.
- Marpissa armifera, Albany (N. Z.), p. 248, nemoralis, Stratford, p. 250, URQUHART, Tr. N. Z. Inst. xxiv; M. grantii, Madeira, p. 217, pl. xiv, figg. 2 & 3, WARBURTON, Ann. N. H. (6) x: n. spp.
- Marptusa nannodes, pp. 283-288, hians, pp. 289-292, decorata, pp. 292-295, demissa, pp. 295-300, oppressa, pp. 300-302, Sumatra, formosissima, Borneo, pp. 303-306, Thorell, Ann. Mus. Genov. xxxi; M. marina, Otago, pp. 253-255, pl. xix, fig. 1, Goyen, Tr. N. Z. Inst. xxiv: n. spp.
- M. rupicola, Hentz, Cayuga Lake, p. 79, pl. iv, fig. 49; BANKS, P. Ac. Philad. 1892.
- Marengo, n.g., p. 66, for M. crassipes, Ceylon, p. 66, pl. v, fig. 4; Peckham, Nat. Hist. Soc. Wisconsin, ii.
- Martella, n. g., p. 45, for M. pottsii, n. sp., Guatemala, p. 45, pl. iv, fig. 2; Peckham, Nat. Hist. Soc. Wisconsin, ii.
- M. maria, Chapoda (Brazil), p. 46, pl. iv, fig. 3, opica, Bahia, p. 47, pl. iv, fig. 4, PECKHAM, Nat. Hist. Soc. Wisconsin, ii, n. spp.
- Nicylla, n. g., p. 449, for N. sundevallii, n. sp., Sumatra, pp. 450-456; Thorell, Ann. Mus. Genov. xxxi.
- Orcevia, n. g., pp. 360 & 361, for O. keyserlingii, pp. 361-365, eucola, pp. 365-369, Thorrell, Ann. Mus. Genov. xxxi, n. spp.
- Paradamætas formicina, Peckh., Guatemala, p. 74, pl. vi, fig. 2; Peck-Ham, Nat. Hist. Soc. Wisconsin, ii.
- Phidippus minutus, Cayuga Lake, p. 74, pl. v, fig. 27, BANKS, P. Ac. Philad. 1892, n. sp.
- Plezippus gausapatus, Sumatra, pp. 370-374, Thorell, Ann. Mus. Genov. xxxi; P. cumulatus, Ceylon, p. 301, pl. xii, fig. 28, Karsch, B. E. Z. xxxvi; P. sylvarus, Stratford (N. Z.), p. 252, URQUHART, Tr. N. Z. Inst. xxiv: n. spp.
- Polichus, n. g. for Gelotia bimaculata, if generically distinct from Gelotia frenata, p. 351; THORELL, Ann. Mus. Genov. xxxi.
- Salticus formicarius, De G., Europe, p. 16, pl. i, fig. 1, ichneumon, Simon, Zanzibar, p. 17, pl. i, fig. 7, sphippiatus, Hentz, Eastern United States, p. 18, fig. 2, plataleoides, Cambr., Ceylon, p. 33, pl. iii, fig. 1; PECKHAM, Nat. Hist. Soc. Wisconsin, ii.
- formosus, pp. 230-233, pecterosus, with n. var. sternodis, Sumatra, pp. 233-239, alticeps, pp. 239-242, leptognathus, Java, pp. 242-246, Thorell, Ann. Mus. Genov. xxxi; modestus, Andaman Is., p. 235, Thorell, Ann. N. H. (6) ix; S. centralis, Central America, p. 19, pl. i, fig. 6, armatus, Lima, p. 20, pl. vii, fig. 6, desertus, Algiers, p. 21, pl. ii, fig. 6, tristis, Simon, Arabia, p. 22, pl. iii, fig. 2, simplex,

- Madagascar, p. 23, pl. i, fig. 4, augustus, Madagascar, p. 24, pl. i, fig. 5, electricus, Madagascar, p. 25, pl. i, fig. 3, gracilis, Madagascar, p. 26, pl. ii, fig. 1, robustus, Burma, p. 27, pl. ii, fig. 2, nemorensis, Burma, p. 28, pl. ii, fig. 3, dubius, p. 29, pl. ii, fig. 4, niger, p. 30, pl. ii, fig. 7, edentulus, p. 31, pl. ii, fig. 5, belticosus, p. 32, pl. ii, fig. 11, Manilla, providens, p. 34, pl. iii, fig. 3, imbellis, p. 36, pl. ii, fig. 10, spissus, p. 37, pl. ii, fig. 8, Ceylon, PECKHAM, Nat. Hist. Soc. Wisconsin, ii, n. spp.
- Sarinda, n. g., p. 40, for S. nigra, Chapoda, Brazil, p. 40, pl. iii, fig. 5, lata, Rio Janeiro, p. 41, pl. iii, fig. 6, n. spp., Peckham, Nat. Hist. Soc. Wisconsin, ii.
- Semora, n. g., p. 68, for S. napæa, n. sp., Rio Janeiro, p. 69, pl. v, fig. 6; PECKHAM, Nat. Hist. Soc. Wisconsin, ii.
- Simonella americana, Peckh., Guatemala, p. 81, pl. vii, fig. 3, myrmeciæformis, Tacz., Brazil, p. 81, pl. vii, fig. 4, bicolor, n. sp., Venezuela,
  p. 83, pl. vii, fig. 5; Peckham, Nat. Hist. Soc. Wisconsin, ii.
- Synageles dalmatensis, Keys., Europe, p. 60, venator, Luc., Algiers, p. 61, pl. v, fig. 1, albotrimaculatus, Luc., Algiers, p. 62, scorpiona, Hentz, New York, p. 6, pl. v, fig. 1, picata, Hentz, United States, p. 64, pl. v, fig. 2, americana, n. sp., E. United States, p. 65, pl. v, fig. 1; PECKHAM, Nat. Hist. Soc. Wisconsin, ii.
- Synemosyna capito, pp. 215-217, debilis, pp. 218 & 219, n. spp., Java, Thorell, Ann. Mus. Genov. xxxi. S. formica, Hentz, United States, p. 77, pl. vi, fig. 4, hentzii, p. 78, pl. vii, fig. 2, lauretta, p. 79, pl. vii, fig. 1, Chapoda, n. spp.; Peckham, Nat. Hist. Soc. Wisconsin, ii.
- Tapinattus, n. g. for melanognathus, Luc., p. 306; THORELL, Ann. Mus. Genov. xxxi.
- Toxeus mandibularis, n. sp. ?, Java, pp. 220-224; THORELL, Ann. Mus. Genov. xxxi.
- Viciria hasseltii, Thor., Celebes, pp. 389-392, albo-limbata, Sim., Sumatra, terebrifera, n. sp, Sumatra, pp. 397-401; Thorell, Ann. Mus. Genov. xxxi.
- Zuniga, n. g., p. 42, for Z. magna, Santarem, p. 43, pl. iv, fig. 1, severa, Caraccas, p. 44, pl. iii, fig. 7, n. spp., Peckham, Nat. Hist. Soc. Wisconsin, ii.

#### CTENIDÆ.

- Acanthoctenus dimidiatus, Sumatra, pp. 142-145, latus, Sarawak, pp. 146-148, Thorell, Ann. Mus. Genov. xxxi: n. spp. According to Simon, Hist. Nat. des Araignées, p. 229, this genus is the sole representative of the subfamily Acanthactenina of the family Zoropsida.
- Ctenus valvularis, Van Hass., Sumatra, pp. 135-139, pulvinatus, n. sp., Sarawak, pp. 139-142; Thorell, Ann. Mus. Genov. xxxi.
- C. (syn. Phoneutria, Microctenus, Anabita, Isoctenus) nigriventer, Rio Grande, p. 144, pl. iv, fig. 98; vehemens, Espirito Santo, p. 145,

fig. 99, velox, Neu Freiburg, p. 147, fig. 100, brevipes, p. 148, fig. 101, minusculus, p. 149, fig. 102, longipes, p. 150, fig. 103, taniatus, p. 151, fig. 104, Rio Grande, griseus, Taquara, p. 152, fig. 195, medius, Rio Grande, p. 153, fig. 106, KEYSERLING, Spinnen Amerikas, 1891, n. spp. According to KEYSERLING Ctenus is placed in the subfamily Ctenius of the family Drassida.

Nydia, n. g., p. 130, for N. punctata, n. sp., Sumatra, p. 131; Thorell, Ann. Mus. Genov. xxxi.

## DICTYNIDE (Or AMAUROBIIDE).

- Amaurobius occidentalis, n. sp., Portugal, p. exciv, Simon, Bull. Soc. Ent. Fr. 1892; A. jheringii, n. sp., Rio Grande, p. 154, pl. iv, fig. 107, KEYSERLING, Spinnen Amerikas. A. similis, Blk., and fenestralis, Ström., figured and compared, p. 389, pls. xxi, figg. 10 & 11; F. CAMBRIDGE, Ann. N. H. (6) x.
- Atelolathys, n. g. for A. varia, n. sp., Ceylon, p. 243; SIMON, Hist. Nat. des Araignées.
- Auximus, n. g. for Amaurobius dentichelis, Sim., p. 239; Simon, Hist. Nat. des Araignées.
- Calleva, n. g. for C. paupercula, n. sp., Argentine, p. 239; Simon, Hist. Nat. des Araignées.
- Dictyna foxii, Marx, p. 28, pl. i, fig. 78; BANKS, P. Ac. Philad. 1892.
- D. maxima, p. 28, pl. i, fig. 79, & pl. iv, fig. 79, decorata, p. 28, pl. i, fig. 81, & pl. iv, fig. 81, dubia, p. 29, pl. i, fig. 82, Cayuga Lake, Banks, P. Ac. Philad. 1892, n. spp. D. philoteichus, McCook, is synonymous with D. civica, Luc., p. 235; Simon, Hist. Nat. des Araignées.
- Neophanes, Marx, p. 29, pallidus, Marx, Cayuga Lake, p. 29, pl. iii, fig. 86; Banks, P. Ac. Philad. According to Simon, Hist. Nat. des Araignées, p. 243, this genus is a synonym of Scotolathys of Simon.
- Protadia, n. g. for Dictyna patula, Sim., p. 239; Simon, Hist. Nat. des Araignées.
- Titanæca sequerai, Portugal, p. cxcv, Simon, Bull. Soc. Ent. Fr. 1892;
  T. luteipes, Rio Grande, p. 156, pl. iv, fig. 108, Keyserling, Spinnen
  Amerikas: n. spp. Simon, Hist. Nat. des Araignées, p. 237, unites
  this genus with Amaurobius.

## DRASSIDÆ.

## Subfam. DRASSINI (? all the genera).

- Agraca ornata, Cayuga Lake, p. 23, pl. i, fig. 68, BANKS, P. Ac. Philad. 1892, n. sp.
- Chiracanthium subflavum, Blck., syn. edentulum, L. Koch, keyserlingii, Beck., Brazil, p. 37, pl. i, fig. 15; KEYSERLING, Spinnen Amerikas.
- C. montanum, Neu Freiburg, p. 38, fig. 16, brevipes, Serra Vermella, p. 39, fig. 16a, KEYSERLING, Spinnen Amerikas, n. spp.

- Clubiona tibialis, Em., p. 20, pl. i, fig. 62, abbotti, Koch, p. 21, fig. 65; BANKS, P. Ac. Philad. 1892.
- C. pygmæa, p. 21, fig. 64, lenta, p. 21, fig. 66, Cayuga Lake, BANKS, P. Ac. Philad. 1892; C. chevronia, Canterbury (N. Z.), p. 231, viridicoma, Mt. Cook, p. 233, URQUHART, Tr. N. Z. Inst. xxiv: n. spp.
- Drassus humilis, n. sp., Cayuga Lake, p. 20, pl. i, fig. 60, Banks, P. Ac. Philad. 1892; D. brasilianus, Rio Grande, p. 34, pl. i, fig. 13, agelastus, Taquara, p. 35, fig. 14, Keyserling, Spinnen Amerikas: n. spp.
- Echemus pulcher, Rio Grande, p. 31, pl. i, fig. 10, major, Rio Grande, p. 32, fig. 11, medius, Fugenda Sergio Potta de Castro, p. 33, fig. 12, KEYSERLING, Spinnen Amerikas, n. spp.
- Eilica, n. g., p. 29, for E. modesta, n. sp., Blumenau, p. 30, pl. i, fig. 9; KEYSERLING, Spinnen Amerikas.
- Goldia, n. g., p. 45, for G. obscura, n. sp., Espirito Santo, p. 45, pl. i, fig. 20; KEYSERLING, Spinnen Amerikas.
- Gnaphosa humilis, Cayuga Lake, p. 19, pl. i, fig. 59, BANKS, P. Ac. Philad. 1892; G. barroisi, Syria, p. 81, Simon, Rev. Biol. v: n. spp.
- Gytha, n. g., p. 27, for G. obscura, n. sp., Rio Grande, p. 28, pl. i, fig. 8;
  KEYSERLING, Spinnen Amerikas.
- Hypsinotus vitiosus, Rio Grande, p. 51, pl. ii, fig. 24, parvus, Serra Vermella, p. 52, pl. ii, fig. 25, alticeps, Espirito Santo, p. 53, fig. 26, fuvipes, Rio Minas, p. 54, fig. 27, botucatensis, Sao Paulo, p. 55, fig. 28, gracilis, Rio Grande, p. 56, fig. 29, nitens, Neu Freiburg, &c., p. 57, fig. 30, KEYSERLING, Spinnen Amerikas.
- Leptodrassus seriatus, Ceylon, p. 294, pl. xi, fig. 16, KARSCH, B. E. Z. xxxvi, n. sp.
- Micaria formicoides, Cayuga Lake, p. 14, pl. i, fig. 51, BANKS, P. Ac. Philad. 1892, n. sp.
- Odo pulcher, Palmeiras, p. 42, pl. i, fig. 18, similis, Corcovado, KEYSER-LING, Spinnen Amerikas, n. spp.
- Onychocryptus, n. g., p. 294, for O. mutillarius, n. sp., Ceylon, p. 295, pl. xi, fig. 17; KARSCH, B. E. Z. xxxvi.
- Phrurolithus minutus, p. 22, pl. i, fig. 67, palustris, p. 23, fig. 70, Cayuga Lake, Banks, P. Ac. Philad. 1892, p. spp.
- Prosthesima rufula, p. 17, pl. i, fig. 55, frigida, p. 17, fig. 56, immacula, p. 18, fig. 58, blanda, p. 18, fig. 57, minima, p. 19, fig. 69, Cayuga Lake, BANKS, P. Ac. Philad. 1892, n. spp.
- Radulphius, n. g., for R. laticeps, Serra Vermella, p. 47, pl. i, fig. 21, bicolor, Neu Freiburg, p. 48, fig. 22, Keyserling, Spinnen Amerikas, n. spp.
- Stasina nalandica, Nalanda (Ceylon), p. 293, pl. xi, fig. 15, KARSCH, B. E. Z. xxxvi, n. sp.
- Teminius brasilianus, Rio Grande, p. 40, pl. i, fig. 17, KEYSERLING, Spinnen Amerikas, n. sp.
- Thargalia agilis, p. 15, pl. i, fig. 52, perplexa, p. 15, fig. 53, fullax, p. 16, fig. 54, Cayuga Lake, Banks, P. Ac. Philad. 1892, n. spp.

Trachelas flavipes, Rio Grande, p. 59, pl. ii, fig. 31a, foraminosus, Rio Grande, p. 60, fig. 32, gracilis, Blumenau, p. 61, fig. 33, rugosus, Serra Vermella, &c., p. 62, fig. 34, robustus, Neu Freiburg, p. 64, fig. 35, vitiosus, Serra Vermella, p. 65, fig. 36, Keyserling, Spinnen Amerikas, n. spp.

#### Subfam. ANYPHÆNINI.

- Abuzaida, n. g., p. 132, for A. striata, n. sp., Neu Freiburg, p. 133, pl. iv, fig. 91; KEYSERLING, Spinnen Amerikas.
- Anyphæna censoria, Serra Vermella, p. 84, pl. ii, fig. 48, procera, p. 86, fig. 49, affabilis, Brazil, p. 87, fig. 50, lenis, Neu Freiburg, p. 88, fig. 51, pygmæa, p. 90, fig. 52, longipes, p. 91, fig. 53, virgata, p. 92, fig. 54, Brazil, cita, Palmeiras, p. 94, fig. 55, violens, Serra Vermella, p. 95, fig. 56, subrubra, Blumenau, p. 97, fig. 57, helvola, Rio Grande, p. 98, fig. 58, rubro-maculata, p. 99, fig. 59, cluro-vittata, Neu Freiburg, p. 100, fig. 60, unicolor, p. 101, fig. 61, striolata, p. 102, fig. 62, robusta, p. 103, fig. 63, grisea, p. 104, fig. 64, taniata, p. 105, fig. 65, fortis, Fazenda Calvario, p. 106, formosa, Corcovado, p. 107, fig. 67, imbecilla, Corcovado, p. 108, fig. 68, angusticeps, p. 109, fig. 69, suspiciosa, p. 110, fig. 70, glabra, p. 111, fig. 71, Serra Vermella, tensa, p. 112, fig. 72, rubella, p. 113, fig. 73, Neu Freiburg, parvula, Serra Vermella, p. 114, fig. 74, tensipes, Neu Freiburg, p. 115, fig. 75, frenata, p. 116, fig. 76, punctata, p. 117, fig. 77, opertanea, p. 118, p. 78, gentilis, p. 119, fig. 79, concolor, p. 120, fig. 80, Neu Freiburg, recentissima, Blumenau, p. 121, fig. 81, tenuipes, Serra Vermella, p. 122, fig. 82, pellucida, Neu Freiburg, p. 122, fig. 83, variabilis, p. 123, fig. 84, diversicolor, p. 124, fig. 85, Neu Freiburg, KEYSEBLING, Spinnen Amerikas, n. spp.
- Arachosia freiburgensis, Neu Freiburg, p. 126, pl. iv, fig. 86, honesta, Bio Grande, p. 127, fig. 87, KEYSERLING, Spinnen Amerikas, n. spp.
- Aysha, n. g., for A. prospera, Rio Grande, p. 129, pl. iv, fig. 88, gracilipes, p. 130, fig. 89, fulviceps, p. 131, fig. 90, Rio Grande, KEYSERLING, Spinnen Amerikas, n. spp.
- Gayenna x-signata, p. 138, pl. iv, fig. 64, marginata, p. 139, fig. 95, fuscotæniata, p. 140, fig. 96, Rio Grande, Keyserling, Spinnen Amerikas, n. spp.
- Samuza, n. g., p. 134, for S. præsignis, p. 135, pl. iv, fig. 22, minuta, p. 136, fig. 93, Rio Grande, Keyserling, Spinnen Amerikas, n. spp.

#### Subfam. CORINNINÆ.

Castianeira (syn. Corinna), valida, Espirito Santo, p. 68, pl. ii, fig. 37, varia, Blumenau, p. 69, fig. 38, pyriformis, Neu Freiburg, p. 70, fig. 39, obscura, Espirito Santo, p. 71, fig. 40, vittata, Blumenau, p. 73, fig. 41, maculata, Neu Freiburg, p. 74, fig. 42, brevis, Corcovado, p. 75, fig. 43, Keyserling, Spinnen Amerikas, n. spp.

Tylophora onerosa, Palmeiras, p. 76, pl. ii, fig. 44, KEYSERLING, Spiunen Amerikas, n. sp.

#### Subfam. MYRMECIINÆ.

- Myrmecia vertebrata, W., and nigra, Perty, Brazil, pp. 78-80, pl. ii, figg. 45 & 46; Keyserling, Spinnen Amerikas.
- M. obscura, Rio Janeiro, p. 81, pl. ii, fig. 47, Keyserling, Spinnen Amerikas, n. sp.

#### DYSDERIDÆ.

- Nops meridionalis, n. sp., Taquara, p. 24; KEYSERLING, Spinnen Amerikas.
- Segestriu suterii, Canterbury, p. 280, URQUHART, Tr. N. Z. Inst. xxiv, n. sp.

## EPEIRIDE (ARGIOPIDE or EUETRIIDE).

- Argiope multi-fasciata, Singapore, p. 226, THORELL, Bull. Soc. Ent. Ital. xxiv; A. plagiata, Minikoy, p. 279, pl. x, fig. 5, trivittata, Ceylon, p. 280, fig. 6, Karsch, B. E. Z. xxxvi: n. spp.
- Argyroepeira fibulata, Singapore, p. 224, THORELL, Bull. Soc. Ent. Ital. xxiv, n. sp.
- Bunocrania? picta, Singapore, p. 251, THORELL, Bull. Soc. Ent. Ital. xxiv, n. sp.
- Epeira beccarii, Thor., described and figured, with snare, pls. v & va; Workman, Malaysian Spiders.
- E. anaspasta, p. 234, perpolita, p. 237, Singapore, Thorell, Bull. Soc. Ent. Ital. xxiv; E. obtusata, Ceylon, p. 283, pl. x, fig. 7, Karsch, B. E. Z. xxxvi; E. invisibilis, Dunedin, p. 239, sublutia, Dunedin, p. 241, simulata, Stewart I., p. 242, ventriosa, Raglan, p. 243, lucisca, Hastwell, p. 245, Urquhart, Tr. N. Z. Inst. xxiv: n. spp.
- E. (Cyclosa) oatesii, Andaman Is., p. 228, Thorell, Ann. N. H. (6) ix; E. confragosa, p. 239, pellux, p. 243, micula, p. 245, Singapore.
- E. (Cyrtophora) eczematica, Singapore, p. 231, Thorell, Bull. Soc. Ent. Ital. xxiv: n. spp.
- Drexelia, n. g., p. 127, for Epeira directa, Hentz., syn. rubella, Hentz., tetragnathoides, Cambr., deludens, Marx, p. 128; McCook, P. Ac. Philad. 1892.
- Gastracantha doriæ, and brevispina, figured on pls. vii & viii; WORKMAN, Malaysian Spiders.
- Meta adeperenta, Ceylon, p. 284, pl. x, fig. 8; KARSCH, B. E. Z. xxxvi, n. sp.
- Milonia obtusa, Singapore, p. 248, Thorell, Bull. Soc. Ent. Ital. xxiv, n. sp.

Nephila transraalica, Pretoria, p. 180, pl. v, fig. 4, Pocock, in Distant's Naturalist in the Transvaal, n. sp.

Poltys apiculatus, Singapore, p. 228, THORELL, Bull. Soc. Ent. Ital. xxiv,

n. sp.

Tetragnatha pallida, Cayuga Lake, p. 51, pl. v, fig. 88, Banks, P. Ac. Philad. 1892; T. geniculata, Ceylon, p. 286, armatu, Passelawa, p. 287, planata, Jaffna, p. 288, determinata, Minikoy, p. 288, foveata, Jaffna, p. 289, Karsch, B. E. Z. xxxvi: n. spp.

Zilla x-notata, Cl., and atrica, C. Koch, compared and figured, p. 395, pl. xx, figg. 3 & xxi, and 8 & 9; F. Cambridge, Ann. N. H. (6) x.

## ERESIDE, FILISTATIDE and HYPOCHILIDE.

Ectatostica, n. g. for Hypochilus davidi, Sim.; Simon, Hist. Nat. des Araignées.

Filistata garciai, Grotto de San Mateo (Philippines), p. 37, Simon, Ann. Soc. Ent. Fr. xli, n. sp.

Stegodyphus sarasinorum, Peradenia, p. 275, KARSCH, B. E. Z. xxxvi, n. sp.

#### LEPTONETIDE.

Synopsis of the genera, p. 39; SIMON, Ann. Soc. Ent. Fr. lxi.

Psiloderces, n. g., p. 40, for egeria, n. sp., Grotto of Calapnitan, p. 40, Simon, t. c.

Theoclia microphthalma, n. sp., Grotto of Antipolo (Philippines), p. 40; Simon, Ann. Soc. Ent. Fr. lxi.

#### LYCOSIDÆ.

Dolomedes lepidus, Sumatra, pp. 149-151, THORELL, Ann. Mus. Genov. xxxi, n. sp.

Diapontia uruguayensis, Keys., Rio Grande, p. 257, pl. x, fig. 193; Keyserling, Spinnen Amerikas.

Lycosa pusiola, Sumatra and Sarawak, pp. 157-161, sumatrana, Sumatra, pp. 161-166, pinangensis, Pinang, pp. 166-169, ragula, pp. 169-173, rabulana, pp. 173-176, ipnochæra, pp. 176-178, Sumatra, Thorell, Ann. Mus. Genov. xxxi; L. rava, Rio Grande, p. 264, pl. x, fig. 199, Keyserling, Spinnen Amerikas; L. similis, pl. ii, fig. 30, rufiventris, pl. iii, fig. 35, humilis, fig. 36, crudelis, fig. 37, immaculata, pl. v, fig. 38, exitiosa, pl. 1, fig. 39, oblonga, pl. iii, fig. 40, pp. 64-68, Cayuga Lake, Banks, P. Ac. Philad. 1892; L. maura, Mt. Cook, p. 246, Urquhart, Tr. N. Z. Inst. xxiv: n. spp.

Lycosella, n. g., pp. 178 & 179, for L. tenera, with varr. bisulcata, pp. 179-183, minuta, pp. 183-185, n. spp., Sumatra, Thorell, Ann. Mus. Genov. xxxi.

- Passiena, n. g., p. 186, for P. spinicrus, n. sp., Pinang, pp. 186-190; THORELL, Ann. Mus. Genov. xxxi.
- Pardosa annulata, pl. i, fig. 41, venusta; fig. 42, gracilis, figg. 43 & 50, mæsta, pl. iii, fig. 44, obsoleta, fig. 45, pp. 68-71, Cayuga Lake; BANKS, P. Ac. Philad. 1892, n. spp.
- Pirata montanoides, pl. i, fig. 46, agilis, fig. 47, exigua, fig. 48, pp. 71 & 72, Cayuga Lake, Banks, P. Ac. Philad. 1892; P. velox, Rio Grande, p. 265, fig. 200, KEYSERLING, Spinnen Amerikas: n. spp.
- Tarantula fragilis, n. sp., Sumatra, pp. 154-157, Thorell, Ann. Mus. Genov. xxxi; thorellii, Keys., syn. sternalis, Bert., Rio Grande, p. 259, pl. x, fig. 195; nervosa, Rio Grande, p. 260, fig. 196, venefica, Rio Grande, p. 261, fig. 197a, auricoma, Taquara, p. 262, fig. 198, n. spp., Keyserling, Spinnen Amerikas.
- Tetragonophthalma obscurum, Taquara, p. 256, pl. x, fig. 192, KEYSERLING, Spinnen Amerikas, n. sp.
- Trochosa inops, Sumatra, pp. 151-154, Thorell, Ann. Mus. Genov. xxxi; pictipes, Rio Grande, p. 266, pl. x, fig. 201, Keyserling, Spinnen Amerikas, n. spp.

#### Œcobiidæ.

Omanus maculatus, n. sp., Blumenau, &c., p. 160, pl. v, fig. iii; KEYSER-LING, Spinnen Amerikas. According to SIMON, Hist. Nat. des Araignées, p. 247, this genus is a synonym of *Œcobius*.

#### OXYOPIDE.

Oxyopes ceylonicus, n. sp., Ceylon, p. 290, pl. xii, fig. 22, Karsch, B. E. Z. xxxvi; O. lineatipes and gemellus, Pinang, figured by Workman, Malaysian Spiders, lineatipes, C. Koch, Sumatra, pp. 190-194; O. javanus, Java, pp. 195-198, longinguus, ? Indo-Malaysia, pp. 198-200, annulipes, Sumatra, pp. 200-202, providens, Sumatra, pp. 202-204, setipes, Sarawak, pp. 204-206, obtusus, Sumatra, pp. 207-209, fronto, Sumatra, pp. 209-211, latifrons, Sumatra, pp. 211-213, n. spp., Thorell, Ann. Mus. Genov. xxxvi; O. constrictus, Rio Grande, p. 268, pl. x, fig. 202, rubrosignatus, Rio Grande, p. 270, pl. x, fig. 203, n. spp., Keyserling, Spinnen Amerikas.

#### PHILODROMIDE.

- Philodromus signifer, pl. iii, fig. 20, gracilis, fig. 21, unicolor, fig. 22, ornatus, fig. 24, placidus, fig. 25; BANKS, P. Ac. Philad. 1892.
- P. minutus, pl. v, fig. 85, minusculus, pl. ii, fig. 39, exilis, pl. ii, fig. 40, pp. 59-63, Cayuga Lake, BANKS, P. Ac. Philad. 1892; P. meridionalis, Rio Grande, p. 251, pl. x, fig. 190, KEYSERLING, Spinnen Amerikas: n. spp.

Thanatus taquara, Taquara, p. 252, pl. x, fig. 191, KEYSERLING, Spinnen Amerikas, n. sp.

### PROLCIDE.

- Pholcus bicornutus, in Caves, Luzon, p. 41, pl. ii, figg. 3 & 4, SIMON, Ann. Soc. Ent. Fr. lxi; imbecillus, Serra Vermella, p. 170, pl. v, fig. 115, cæruleus, p. 171, fig. 116, luteus, p. 171, fig. 117, togatus, p. 172, fig. 118, cyaneo-maculatus, p. 173, fig. 119, Brazil, altiventer, Serra Vermella, p. 175, fig. 120, cyaneo-tæniatus, Brazil, p. 176, fig. 121, Keyserling, Spinnen Amerikas: n. spp.
- Spermophora estebani, in Cave, Luzon, p. 42, Simon, Ann. Soc. Ent. Fr. lxi; S. maculata, Blumenau, p. 177, pl. vi, fig. 122, unicolor, Serra Vermella, p. 178, fig. 123, bicornis, Sao Paulo, p. 179, fig. 124, KEYSERLING, Spinnen Amerikas: n. spp.
- Calapnita, n. g. for C. vermiformis, n. sp., in Cave, Luzon, p. 42; Simon, Ann. Soc. Ent. Fr. lxi.

### PALPIMANIDE and PRODIDOMIDE.

- Jheringia, n. g. Palpimanidæ, p. 25, for J. lutea, n. sp., Rio Grande, p. 26; KEYSERLING, Spinnen Amerikas.
- Storena annulipes, Singapore, p. 209, THORELL, Bull. Soc. Ent. Ital. xxiv, n. sp.

#### SCYTODIDE.

- Scytodes annulata, Serra Vermella, p. 166, pl. v, fig. 113; Keyserling, Spinnen Amerikas, n. sp.
- Pandeus, n. g., p. 167, for P. caruleus, n. sp., Blumenau, p. 168, pl. v, fig. 114; KEYSERLING, Spinnen Amerikas.

#### SPARASSIDÆ.

- Angœus rhombipes, Sumatra, p. 67; THORELL, Ann. Mus. Genov. xxxi, n. sp.
- Dolothymus, n. g., pp. 62 & 63, for D. pallidus, n. sp., Sumatra, pp. 64-67; THORELL, Ann. Mus. Genov. xxxi.
- Heteropoda boiei, Dol., syn. flavimana, Simon, Sumatra, Java, pp. 15 & 16;
  H. thoracica, C. Koch, Padang, &c., pp. 24-26; THORELL, Ann.
  Mus. Genov. xxxi.
- H. imbecilla, Padang, pp. 16-19, leptoscelis, Sumatra, pp. 19-22, sumatrana (with n. var. montana), pp. 26-31, signata, pp. 32-34, Sumatra, obtusa, Sarawak, pp. 34-37, Thorell, Ann. Mus. Genov. xxxi; H.

- umbrata, pl. xi, fig. 11, eluta, fig. 12, p. 291, subtilis, p. 292, fig. 13, Ceylon, Karsch, B. E. Z. xxxvi: n. spp.
- Holconia beccarii, Sumatra, pp. 46-49, Thorell, Ann. Mus. Genov. xxxi, n. sp.
- Libania, n. g., pp. 49-51, for L. scabricula, n. sp., with var. sulcata, Sumatrs, pp. 51-56; lævis, pp. 56 & 57, annulata, pp. 58-60, armillata, pp. 60-62, Sumatra, Thorell, Ann. Mus. Genov. xxxi, n. spp.
- Palystes incanus, Sarawak, pp. 37-40, THORELL, Ann. Mus. Genov. xxxi, n. sp.
- Sarotes impudicus, Andaman Is., p. 233, THORELL, Ann. N. H. (6) ix; S. ferox, loc. ?, p. 40, THORELL, Ann. Mus. Genov. xxxvi: n. spp.
- Theleticopis orichalcea, Sim., Sarawak, pp. 44-46; Thorell, Ann. Mus. Genov. xxxvi.
- Urgulania, n. g., p. 11, for U. borneensis, n. sp., Sarawak, p. 12, THORELL, Ann. Mus. Genov. xxxvi.
- Zatapina, n. g. for Delena plumipes, Dol., p. 290 (referred to the family Clubionidæ); KARSCH, B. E. Z. XXXVI.

#### STEPHANOPIDE.

- Stephanop[o]idæ. New family based upon Stephanopis, and allied to the Thomisidæ and Sparassidæ (Heteropodidæ), p. 5; THORELL, Ann. Mus. Genov. xxxi.
- Regillus, Cambr., remarks on, p. 6; R. cinerascens, Dol., Ajur Mantjur, pp. 6-11; Thorell, Ann. Mus. Genov. xxxi.

#### THERAPHASIDE OF AVICULARIDE.

- Acanthoscurria minor, Auss., Guyana, p. 281; SIMON, Ann. Soc. Ent. Fr. lxi.
- A. maga, S. America, p. 280, musculosa, p. 281, insubtilis, Bolivia, p. 282, ferina, Amazons, p. 282, Simon, Ann. Soc. Ent. Fr. lxi, n. spp.
- Accola cavicola, Grotto of San Mateo, p. 35, eaca, Grotto d'Antipolo, p. 35, Simon, Ann. Soc. Ent. Fr. lxi, n. spp.
- Actinoxia, n. g. for A. versicolor, n. sp., California, p. 318; Simon, Act. Soc. L. Bord. xliv.
- Agathostola, n. g. for A. strabo, n. sp., Columbia, p. 163; Simon, Hist. Nat. des Araignées.
- Apostichus, n. g. for A. atomarius, p. 317, n. sp.; Sinon, Act. Soc. L. Bord. xliv.
- A. clathratus, p. 318, California, SIMON, Act. Soc. L. Bord. xliv, n. sp.
- Arbanitis huttonii, Cambr., Dunedin, p. 255, pl. xix, figg. 2 & 3; Goyen, Tr. N. Z. Inst. xxiv.
- Bothriocyrtum, n. g., p. 314, for Cteniza californica, Cambr., Texas, California, &c., p. 315, and B. fabrile, n. sp., Mexico, p. 327; Simon, Act. Soc. L. Bord. xliv.

- Brachybothrium (syn. Nidivalvata, Atk.) robustum, Virginia, &c., p. 311, Simon, Act. Soc. L. Bord. xliv, n. sp.
- Brackythele longitarsis, Sim., Texas, &c., p. 319; Simon, Act. Soc. L. Bord. xliv.
- Brackypelma, n. g. for B. emilia, White, Mexico, p. 338; SIMON, Act. Soc. L. Bord. xliv.
- Chilobrachys, n. g., p. 270, for C. nitelinus, n. sp., Ceylon, p. 271, pl. x, fig. 2; Karsch, B. E. Z. xxxvi.
- Cratorrhagus, n. g. for C. cervinus, n. sp., Mexico, p. 330; Simon, Act. Soc. L. Bord. xliv.
- Crypsidromus bolivianus, Bolivia, p. 278, SIMON, Ann. Soc. Ent. Fr. lxi;
  C. perfidus, p. 14, funestus, p. 15, Taquara, Keyserling, Spinnen
  Amerikas: n. spp.
- Cyclocosmia theveneti, California, p. 313, Simon, Act. Soc. L. Bord. xliv, n. sp.
- Cyclosternum obscurum, Mexico, p. 330, Simon, Act. Soc. L. Bord. xliv, n. sp.
- Cyrtauchenias sebra, Zululand, p. 272, Simon, Ann. Soc. Ent. Fr. lxi, n. sp.
- Cyrtosternum meridionale, Taquara, p. 6, KEYSERLING, Spinnen Amerikas, n. sp.
- Diplothele halyi, Ceylon, p. 123, Simon, Hist. Nat. des Araignées, n. sp. Encyocrates, n. g., p. 279, for E. raffrayi, Madagascar, p. 280; Simon, Ann. Soc. Ent. Fr. lxi, n. sp.
- Ephebopus, n. g. for Mygale murinus, Walck., p. 155; Simon, Hist. Nat. des Araignées.
- Eudiplura, n. g. for Diplura rogenhoferi, Auss., p. 179; Simon, Hist. Nat. des Araignées.
- Eurypelma steindachneri, Auss., Texas; hentsi, Gir., syn. mordax, Auss., Texas, &c., p. 322; vagans, Auss., syn. duponti, Becker, Mexico, &c.; Simon, Act. Soc. L. Bord. xliv.
- E. jheringii, p. 19, pl. i, fig. 4, vitiosa, p. 21, pl. i, fig. 5, Taquara, Keyserling, Spinnen Amerikas; E. rusticum, Arizona, p. 323, helluo, Lower California, p. 324, marxi, California, &c., lanceolatum, Nicaragua, p. 335, caniceps, Mexico, p. 336, serratum, Mexico, panamense, p. 337, Panama, Simon, Act. Soc. L. Bord. xliv; E. guyanum, Guyana, p. 167, Simon, Hist. Nat. des Araignées: n. spp.
- Evagrus rubrigularis, Idaho, p. 320, SIMON, Act. Soc. L. Bord. xliv, u sp. Grammostola, n. g. for Eurypelma pulchripes, Simon, Paraguay, p. 163; SIMON, Hist. Nat. des Araignées.
- Hapalopus villosus, Taquara, p. 7, pl. i, fig. 2, KEYSERLING, Spinnen Amerikas, n. sp.
- Hapalothele (Fufius) lanicia, p. 283, garleppi, p. 284, Bolivia, Simon, Ann. Soc. Ent. Fr. lxi, n. spp.
- Haploclastus, n. g. for H. cervinus, n. sp., S. India, p. 152; SIMON, Hist. Nat. des Araignées.

- Haplopelma, n. g. for Selenocosmia doriæ, Thor., p. 151; SIMON, Hist. Nat. des Araignées.
- Heligmomerus, n. g., p. 90, for H. taprobanicus, Kandy, prostans, S. India, p. 91 (in note), Simon, Hist. Nat. des Araignées, n. spp.
- Homecomma texense, Texas, p. 320, Sinon, Act. Soc. L. Bord. xliv, n. sp.
- Homeoplacis, n. g. for H. pentodon, Amazons, p. 275, Simon, Ann. Soc. Ent. Fr. lxi, n. sp.
- Homostola, n. g., p. 271, for H. vulpscula, Zululand, p. 271, Simon, Ann. Soc. Ent. Fr. lxi, n. sp.
- Hysterocrates, n. g. for Phoneyusa greshoffi, Simon, p. 153; Simon, Hist. Nat. des Araignées.
- Idiops germaini, Rio Grande, p. 92, SIMON, Hist. Nat. des Araignées, n. sp. Ischnocolus pilosus, Rio Grande, p. 9, gracilis, Monte Verde, p. 11, rubropilosus, Neu Freiburg, p. 12, jansirus, Rio Janeiro, p. 13, KEYSER-LING, Spinnen Amerikas, n. spp.
- Lampropelma, n. g., p. 151, for L. nigerrimum, Ins. Sangir, p. 151, SIMON, Hist. Nat. des Araignées, n. sp.
- Macrothele segmentata, Pinang, p. 284, Simon, Ann. Soc. Ent. Fr. 1x1, n. sp.
- Magulla, n. g. for M. obesa, Theresopolis, p. 276, Simon, Ann. Soc. Ent. Fr. lxi, n. sp.
- Myrmeciophila atkinsoni, Virginia, p. 316, Simon, Act. Soc. L. Bord. xliv, n. sp.
- Nemesia arenicola, Corsica, p. 113, didieri, Algeria, p. 114, Simon, Hist. Nat. des Araignées, n. spp.
- Œcophlæus, n. g. for Œ. cinctipes, n. sp., Ceylon, pp. 49-51, figured, with nest, pl. iiia; Pocock, Ann. N. H. (6) ix: syn. Sason robustum, Cambr.; Simon, Hist. Nat. des Araignées.
- Ornithoctonus, n. g. for O. andersoni, Mergui, pp. 317 & 318, pl. xvii, figg. 1-3, POCOCK, J. L. S. xxiv, n. sp.
- Orphnæcus, n. g. for O. pellitus, Grotto de Calapnitan (Philippines), pp. 36 & 37, Simon, Ann. Soc. Ent. Fr. lxi, n. sp.
- Pachyloscelis crassipes, Rio Grande do Sul, p. 3, pl. i, fig. 1, luteipes, Rio Janeiro, p. 5, KEYSERLING, Spinnen Amerikas, n. spp.
- Paraphysa, n. g. for P. manicata, S. America, p. 166, Simon, Hist. Nat. des Araignées, n. sp.
- Phlogius cunicularius, Pinang, p. 279, Simon, Ann. Soc. Ent. Fr. lxi, n. sp. Phrissæcia, n. g. for P. ecuadorensis, Ecuador, p. 274, Simon, Ann. Soc. Ent. Fr. lxi, n. sp.
- Plagiobothrus, n. g., p. 272, for P. semilunaris, Peradenia, p. 273, pl. x, fig. 3, KARSCH, B. E. Z. xxxvi, n. sp.
- Planadecta, n. g. for P. lyrata, S. America, p. 168, Simon, Hist. Nat. des Araignées, n. sp.
- Porrhothele, n. g. for Mygals antipodiana, Walck., p. 185; Simon, Hist. Nat. des Araignées.
- Pselligmus, n. g. for P. infaustus, Abrolhos, p. 273, Simon, Ann. Soc. Ent. Fr. lxi, n. sp.

- Ruchias, n. g. for Hermacha dispar, Sim., p. 114; SIMON, Hist. Nat. des Araignées.
- Rhechostica, n. g. for Homosomma tenense, Simon, p. 162; Simon, Hist. Nat. des Araignées.
- Scalidognathus, n. g., p. 269, for S. seticeps, Peradenia, p. 270, pl. x, fig. 1, Karsch, B. E. Z. xxxvi, n. sp.
- Scotinæcus, n. g. for Hexathele cinereo-pilosa, Sim.; Simon, Hist. Nat. des Araignées.
- Sipalolasma, n. g., p. 123, for S. ellioti, Ceylon, p. 124; Simon, Hist. Nat. des Araignées, n. sp.
- Stasimopus, n. g., p. 81, for Actinopus caffrus, C. Koch, p. 81; SIMON, Hist. Nat. des Araignées.
- Stenygrocercus, n. g. for Macrothele silvicola, Sim., p. 185; Simon, Hist. Nat. des Araignées.
- Sterrhochrotus, n. g. for Cteniza farghanensis, Cron., p. 97; Simon, Hist. Nat. des Araignées.
- Stichoplastus fabrei, India, p. 278, SIMON, Ann. Soc. Ent. Fr. lxi, n. sp.
- Tapinauchenius cœrulescens, Indian Terr., p. 325, texense, Texas, p. 326, Simon, Act. Soc. L. Bord. xliv, n. spp.
- Thelecoris, Karsch (syn. Entomothele, Simon), sebrina, Polvon [Nicaragua], p. 329, Simon, Act. Soc. L. Bord. xliv, n. sp.
- Tigidia, n. g. for T. mauriciana, Mauritius, p. 274, SIMON, Ann. Soc. Ent. Fr. lxi, n. sp.
- Tmesiphantes, n. g., for T. nubilus, Bahia, p. 277, Simon, Ann. Soc. Ent. Fr. lxi, n. sp.
- Trechona auronitens, p. 16, pl. i, fig. 3, pantherina, p. 18, Taquara, Keyserling, Spinnen Amerikas, n. spp.
- Xenesthis, n. g., p. 332, for X. columbiana, Panama, p. 333, Simon, Act. Soc. L. Bord. xliv, n. sp.

#### THERIDIDE.

- Achæa maxima, Rio Grande, p. 198, pl. vii, fig. 142, signata, Neu Freiburg, p. 199, fig. 143, pulchra, Rio Janeiro, p. 200, fig. 144, Keyserling, Spinnen Amerikas, n. spp.
- Ariannes longissimus, Neu Freiburg, &c., p. 202, pl. vii, fig. 145, KEYSER-LING, Spinnen Amerikas; A. simulans, Calcutta, p. 417, pl. xxii, CAMBRIDGE, Ann. N. H. (6) x: n. spp.
- Argyrodes fusciatus, Singapore, p. 212, Thorell, Bull. Soc. Ent. Ital. xxiv; A. acuminatus, Serra Vermella, &c., p. 207, pl. vii, fig. 149, longicaudatus, Serra Vermella, p. 209, fig. 150, alticeps, Espirito Santo, p. 210, fig. 151, altus, Espirito Santo, p. 211, fig. 152, ululabilis, Taquara, p. 212, fig. 153, striatus, Espirito Santo, p. 213, fig. 154, Keyserling, Spinnen Amerikas, n. spp.
- Bathyphantes parvulus, Westr., and gracilis, Blk., England, pp. 392 & 393, pl. xxi, figg. 6 & 7; F. Cambridge, Ann. N. H. (6) x.

- B. decorata, p. 44, pl. ii, fig. 41, argenteo-maculata, pallida, pl. v, fig. 42, unimaculata, pl. ii, fig. 65, p. 45, inornata, pl. v, fig. 66, tristis, pl. ii, fig. 45, p. 46, Cayuga Lake, Banks, P. Ac. Philad. 1892, n. spp.
- Ceratinella minuta, Em., pl. ii, fig. 60, & pl. iv, fig. 60, bulbosa, Em., p. 32, pl. v, fig. 59, pygmea, Em., p. 32, pl. ii, fig. 57; BANKS, P. Ac. Philad. 1892.
- C. similis, p. 31, pl. v, fig. 61, masta, p. 32, pl. ii, fig. 58, & pl. v, fig. 58, placida, p. 32, pl. ii, fig. 54, formosa, p. 33, pl. ii, fig. 55, annulipes, p. 33, pl. ii, fig. 56, Cayuga Lake, Banks, P. Ac. Philad. 1892, n. spp.
- Ceratinopsis frontatus, Cayuga Lake, p. 33, pl. v, fig. 63, BANKS, P. Ac. Philad. 1892, n. sp.
- Chrysso nigrosternum, Taquara, p. 206, pl. vii, fig. 148, KEYSERLING, Spinnen Amerikas, n. sp.
- Cornicularia formosa, p. 34, pl. v, fig. 35, placida, p. 35, fig. 36, Cayuga Lake, Banks, P. Ac. Philad. 1892, n. spp.
- Diplostyla pallida, pl. ii, fig. 29, alboventris, pl. v, fig. 31, p. 43, Cayuga Lake, Banks, P. Ac. Philad. 1892, n. spp.
- Dipæna tæniatipes, Rio Grande, p. 224, pl. viii, fig. 165, maculata, Espirito Santo, p. 225, fig. 166, obscura, Serra Vermella, p. 226, fig. 167, Keysekling, Spinnen Amerikas, n. spp.
- Erigone rusticella, Rio Grande, p. 238, pl. ix, fig. 180a, nitida, Rio Grande, p. 239, fig. 181a, fucosa, p. 240, fig. 182a, rusticula, p. 240, fig. 183, vegeta, p. 241, fig. 184, Rio Grande, KEYSERLING, Spinnen Amerikas, n. spp.
- Ero goldii, Serra Vermella, p. 218, pl. viii, fig. 159, gracilis, Neu Freiburg, p. 219, fig. 160, ralida, St. Antonio, p. 220, fig. 161, lata, Neu Freiburg, p. 220, fig. 162, KEYSERLING, Spinnen Amerikas, n. spp.
- Euryopis lutea, Serra Vermella, p. 227, pl. ix, fig. 168, maculata, Miracena, p. 227, fig. 169, KEYSERLING, Spinnen Amerikas, n. spp.
- Frontina turbidulu, Espirito Santo, p. 228, pl. ix, fig. 170, armata, Serra Vermella, p. 230, fig. 171, obscura, Espirito Santo, p. 231, fig. 172, cylindrata, Sao Paulo, p. 232, fig. 173, elegans, Rio Negro, p. 233, fig. 174, clava, Serra Vermella, p. 234, fig. 175, longiceps, Serra Vermella, p. 234, fig. 176, decorata, Neu Freiburg, p. 235, fig. 177, rubriceps, Rio Grande, p. 236, fig. 178, Keyserling, Spinnen Amerikas, n. spp.
- Grammonota venusta, Cayuga Lake, p. 34, pl. v, fig. 64, BANKS, P. Ac. Philad. 1892, n. sp.
- Helvibis longicauda, Espirito Santo, p. 204, pl. vii, fig. 146, monticola, Serra Vermella, p. 205, fig. 147, Keyserling, Spinnen Amerikas, n. spp.
- Heribertus flavomaculatus, Serra Vermella, p. 223, pl. ix, fig. 164, Keyserling, Spinnen Amerikas, n. sp.
- Leptyphantes ericaus, Blk., England, p. 390, pl. xx, fig. 4; F. CAMBRIDGE, Ann. N. H. (6) x.

- L. plumiger, Dorset, p. 386, pl. xx, fig. 2, F. CAMBRIDGE, Ann. N. H. (6) x, n. sp.
- Linyphia conferta, Hentz, pl. ii, fig. 38, p. 42, Cayuga Lake; BANKS, P. Ac. Philad. 1892.
- L. variabilis, pl. ii, fig. 28, Cayuga Lake, BANKS, P. Ac. Philad. 1892, n. sp.
- Lophocarenum castuneum, Em., p. 35, pl. iv, fig. 3; BANKS, P. Ac. Philad. 1892.
- L. triste, p. 35, pl. iv, fig. 1, unimaculatum, p. 35, fig. 2, miniatum, fig. 4, venustum, fig. 5, parvum, fig. 6, exiguum, fig. 7, p. 36, crenatoideum, fig. 8, formosum, fig. 10, arvense, fig. 11, p. 37, longius, p. 38, fig. 12, Cayuga Lake; Banks, P. Ac. Philad. 1892, n. spp.
- Microneta latens, pl. v, fig. 46, p. 46, palustris, pl. ii, fig. 47, luteola, pl. ii, fig. 48, flaveola, pl. v, fig. 42, complicata, pl. ii, fig. 50, p. 47, minutissima, pl. ii, fig. 27, frontata, pl. v, fig. 51, gigantea, pl. ii, fig. 52, distincta, pl. ii, fig. 53, p. 48, Cayuga Lake, Banks, P. Ac. Philad. 1892, n. spp.
- Milione, n. g., p. 215, for M. triacantha, n. sp., Singapore, p. 216; Tho-Bell, Bull. Soc. Ent. Ital. xxiv.
- Mimetus atri-cinctus, Stratford (N. Z.), p. 234, URQUHART, Tr. N. Z. Inst. xxiv, n. sp.
- Theridium weberi, Singapore, p. 219, Thorell, Bull. Soc. Ent. Ital. xxiv; T. bellulum, Neu Freiburg, p. 180, pl. vi, fig. 125, unanimum, Neu Freiburg, p. 181, fig. 126, guadripartitum, Sao Paulo, p. 182, fig. 127, subrotundum, Serra Vermella, p. 183, fig. 128, bentificum, Rio Grande, p. 184, fig. 129, tinctorium, Rio Grande, p. 185, fig. 130, querulum, p. 186, fig. 131, obnubilum, Serra Vermella, p. 187, fig. 132, dubiosum, Neu Freiburg, p. 187, fig. 133, nigriceps, Sao Paulo, p. 188, fig. 134, notabile, Serra Vermella, p. 189, fig. 135, triguttatum, Neu Freiburg, p. 190, fig. 136, rotundum, Neu Freiburg, p. 191, fig. 137, pallipes, Serra Vermella, p. 191, fig. 138, vividum, Espirito Santo, p. 192, fig. 139, bicorne, Serra Vermella, p. 193, fig. 140, passivum, Fazenla Calvario, p. 195, fig. 141, Keyserling, Spinnen Amerikas; T. argentatum, Hawera, p. 235, albo-cinctum, p. 236, ampliatum, p. 237, Stratford, Urquhart, Tr. N. Z. Inst. xxiv: n. spp.
- Tmeticus unicorn, obscurus, flaveolus, luxuosus, rusticus, humilis, mæstus, debilis, palustris, distinctus, maculatus, minutus, pl. iv, figg. 13-24, pp. 38-41, gnavus, p. 41, pl. v, fig. 44, Cayuga Lake, Banks, P. Ac. Philad. 1892; T. simplex, Cannock, p. 384, pl. xx, fig. 5, F. Cambridge, Ann. N. H. (6) x: n. spp.
- Totua, n. g. for T. gracilipes, Rio Grande, p. 216, pl. viii, fig. 158; KEYSERLING, Spinnen Amerikas, n. sp.
- Umfila cornuta, Neu Freiburg, p. 222, pl. viii, fig. 163, KEYSERLING, Spinnen Amerikas, n. sp.

#### THOMISIDA.

- Camaricus formosus, Thor., figured on pl. iv, by WORKMAN, Malaysian Spiders.
- Diæa zonura, Sumatra, pp. 95-98, THORELL, Ann. Mus. Genov. xxxi, n. sp.
- Daradius javanus, Java, pp. 78-82, perspicillatus, Sarawak, pp. 82-86, Thorell, Ann. Mus. Genov. xxxi, n. spp.
- Hedana perspicaz, Sumatra, pp. 106-109, ocellata, Java, pp. 109-113, THORELL, Ann. Mus. Genov. xxxi, n. spp.
- Microcyllus, n. g., p. 118, for M. elegans, Sumatra, pp. 119-121, nanus, Java, pp. 121 & 122, Thorell, Ann. Mus. Genov. xxxi, n. spp.
- Misumena foliata, p. 57, pl. iii, fig. 17, & pl. ii, fig. 37, placida, p. 58, Cayuga Lake, Banks, P. Ac. Philad. 1892; M. dierythra, Singapore, p. 92, Thorell, Ann. Mus. Genov. xxxi; M. maculis-sparsa, Taquara, p. 246, pl. x, fig. 186, Keyserling, Spinnen Amerikas; M. clarkii, Madeira, p. 219, pl. xiv, figg. 7 & 8, Warburton, Ann. N. H. (6) x: n. spp.
- Museeus, n. g., pp. 71 & 72, for M. politus, n. sp., Sumatra, pp. 72-75; THORELL, Ann. Mus. Genov. xxxi.
- Narcaus, n. g., p. 79, for N. picinus, n. sp., Java, pp. 100-102; Thorell, Ann. Mus. Genov. xxxi.
- Palæphatus pseudorchestes, Sumatra, p. 127, THORELL, Ann. Mus. Genov. xxxi, n. sp.
- Peltorhynchus, n. g., pp. 113 & 114, for P. rostratus, n. sp., Java, pp. 114-117, Thorell, Ann. Mus. Genov. xxxi; also figured, pl. iii, by Work-Man, Malaysian Spiders.
- Runcinia kinbergii, pp. 86-89, disticta, pp. 89-91, Java, THORRLL, Ann. Mus. Genov. xxxi, n. spp.
- Synema luteovittata, Rio Grande, p. 247, pl. x, fig. 187, KEYSERLING, Spinnen Amerikas, n. sp.
- Tharpyna varica, Java, p. 102, THORELL, Ann. Mus. Genov. xxxi, n. sp.
- Tmarus cques, Java, pp. 75-78, Thorell, Ann. Mus. Genov. xxxi; T. variatus, pl. x, fig. 188, clavipes, p. 280, fig. 189, Taquara, Keyserling, Spinnen Amerikas: n. spp.
- Xysticus distinctus, pl. iii, fig. 89, brunneus, pl. iii, fig. 4, crudelis, fig. 5, transversus, fig. 6, lentus, pl. ii, fig. 67, nervosus, pl. iii, fig. 8, & pl. iv, fig. 84, formosus, pl. iii, fig. 9, pp. 52-56, Cayuga Lake, Banks, P. Ac. Philad. 1892, n. spp.
- Zametopias, n. g., pp. 122 & 123, for Z. speculator, Sumatra, pp. 123-126; Thorell, Ann. Mus. Genov. xxxi, n. sp.

#### ULOBORIDÆ.

\*\*Ebutina, n. g. for \*\*\mathbb{E}\$. binotata, n. sp., Brazil, p. 222, Simon, Hist. Nat. des Araignées. According to Simon, t. c., this genus is the sole representative of the subfamily \*\*\mathbb{E}\text{butinina}\text{ of the family \$Uloborida.}

Hyptiotes analis, n. sp., Kandy, p. 219, Simon, Hist. Nat. des Araignées. According to Simon, t. c., this genus constitutes, with Miagrammopes, the subfamily Miagrammopines of the family Uloborides.

Uloborus guadri-tuberculatus, Singapore, p. 221, THORELL, Bull. Soc. Ent. Ital. xxiv, n. sp.

## ZODARIIDE and ZOROPSIDE.

- Habronestes (Zodariidæ) major, p. 158, pl. iv, fig. 109, minor, p. 159, fig. 110, Blumenau, Keyserling, Spinnen Amerikas, n. spp.
- Recius, n. g. of Zoropside, for Amaurobius crassipes, L. Koch, p. 230, Simon, Hist. Nat. des Araignées.

## (D.) PSEUDOSCORPIONES.

BALZAN, Ann. Soc. Ent. Fr. lx, divides this group into two suborders: the *Panctenodactyli*, containing the *Cheliferidæ*, and the *Hemictenodactyli*, for the *Obisiidæ* and *Pseudobisiidæ*.

#### CHELIFERIDE.

- Atemnus nidificator, n. var. minor, San Esteban, p. 510, pallidus, Sierra Leone, p. 511, pl. ix, fig. 2, subrobustus, Caraccas, p. 512, fig. 3, BALZAN, Ann. Soc. Ent. Fr. lx, n. spp.
- Chernes insuetus, Dover, p. 225, pl. c, fig. 17, dubius, Sherborne, p. 227, fig. 19, Cambridge, P. Dorset Field Club, xiii, n. spp.
- Lamprochernes macrochelatus, Töm., and argentinus, Thor., Venezuela, pp. 513 & 516, pl. ix, figg. 4 & 7; Balzan, Ann. Soc. Ent. Fr. lx.
- L. octentoctus, S. Africa, p. 514, fig. 5, intermedius, Venezuela, p. 515, fig. 6, similis, Manaos, p. 517, fig. 8, venezuelanus, San Esteban, p. 518, fig. 9, ovatus, Caraccas, fig. 10, thorelli, Sumatra, pl. x, fig. 11, p. 519, elegans, Venezuela, p. 520, fig. 12, BALZAN, Ann. Soc. Ent. Fr. lx, n. spp.
- Trachychernes subrudis, Venezuela, p. 521, pl. x, fig. 13, subrotundatus, Venezuela, p. 522, fig. 14, floridæ, Florida, p. 524, fig. 15, bicolor, Venezuela, p. 524, Caraccas, fig. 16, albomaculatus, Venezuela, p. 526, fig. 17, armiger, Pebas, p. 527, fig. 18, Balzan, Ann. Soc. Ent. Fr. lx, n. spp.
- Chelifer canestrinii, Balz., p. 532, fig. 22, meridianus, L. Koch, fig. 23, longichelifer, Balz., Venezuela, p. 534, fig. 26, rufus, Balz., fig. 25; Balzan, Ann. Soc. Ent. Fr. lx.
- C. pekinensis, Pekin, p. 528, pl. xi, fig. 19, simoni, Sierra Leone, p. 529, fig. 20, tenuimanus, Nossi-bé, p. 531, fig. 21, degeneratus, S. California, p. 532, fig. 24, BALZAN, Ann. Soc. Ent. Fr. lx, n. spp.
- Garypus senegalensis, Senegal, p. 535, pl. xii, fig. 27, BALZAN, Ann. Soc. Ent. Fr. lx, n. sp.
- Olpium crassichelatum, Balz., fig. 29, Venezuela; Balzan, Ann. Soc. Ent. Fr. lx.
- O. cordinanum, Venezuela, p. 536, pl. xii, fig. 28, furculiferum, Venezuela, p. 537, fig. 30, BALZAN, Ann. Soc. Ent. Fr. lx, n. spp. 1892. [Vol. XXIX.7]

#### PSEUDOBISHDÆ.

N. fam., p. 539; BALZAN, Ann. Soc. Ent. Fr. lx.

#### PSEUDOBISIINÆ.

N. subfam., p. 539; BALZAN, Ann. Soc. Ent. Fr. lx.

Ideobisium, n. g., p. 539, (Ideoroncus) gracilis, n. sp., Venezuels, p. 540, pl. xii, fig. 31; BALZAN, Ann. Soc. Ent. Fr. lx.

Ideoblothrus, n. subg. of Ideobisium, p. 541, for I. similis, n. sp., Venezuela, p. 541, pl. xii, fig. 32; BALZAN, Ann. Soc. Ent. Fr. lx.

Ideobisium crassimanum, n. sp., Venezuela, p. 542, pl. xii, fig. 33, BALZAN, Ann. Soc. Ent. Fr. lx.

#### MICROCREAGRINÆ.

N. subfam.; BALZAN, Ann. Soc. Ent. Fr. lx, p. 543.

Microcreagris, n. g., p. 543, for M. gigas, n. sp., China, p. 544, pl. xii, fig. 34; Balzan, Ann. Soc. Ent. Fr. lx.

#### OBISIDE.

#### CHTHONIINÆ.

N. subfam., p. 545; Balzan, Ann. Soc. Ent. Fr. lx.

- Chthonius parvidentatus, Balz., p. 547, fig. 36; Balzan, Ann. Soc. Ent. Fr. lx.
- C. (Pseudochthonius, n. subg.) simoni, Caraccas, p. 546, pl. xii, fig. 35, Balzan, Ann. Soc. Ent. Fr. lx, n. sp.

#### (E.) OPILIONES.

- Caddo, n. g., for C. agilis, n. sp., Long I., p. 280; BANKS, P. E. Soc. Wash, ii.
- Gagrella ceylonensis, p. 308, bispinosa, Passelawa, p. 309, Karsch, B. E. Z. xxxvi, n. spp.
- Liobunum vittatum, Say, pp. 999-1008, pla xxvii & xxviii; Weed, Am. Nat. xxvi.
- Phalangium cinereum, Wood, pp. 32-36, pl. i, figg. 1 & 2; WEED, Am. Nat. xxvi.
- Pseudarthromerus, n. g., p. 308, for P. spurius, n. sp., Nalanda (Ceylon), p. 308; Karsch, B. E. Z. xxxvi.
- Zalmozis sorenseni, in caves, Luzon, p. 44, Simon, Ann. Soc. Ent. Fr. lxi, n. sp.
- Oryptostemma, Guér., notes on, with fig.; the genus referred to the order Meridogastra of Thorell; Karscu, B. E. Z. xxxvii, pp. 25-32 & 64.

#### (F.) ACARL

#### BDELLIDÆ.

- Bdella lignicola, G. Can., pt. lxii, No. 7; Berlese, Acari in Italia reperta. Linopodes, pt. lxi, No. 1, motatorius, L., Italy, No. 2, Berlese, Acari in Italia reperta.
- Norneria, pt. lxi, No. 6, gigas, R. Can., No. 7; Berlese, Acari in Italia reperta.
- Notophallus longipilis, R. Can., pt. lxi, No. 4, minor, R. Can., No. 5, Italy; Berlese, Acari in Italia reperta.
- Pronematus bonatii, Can., Italy, pt. lxi, No. 3; Berlese, Acari in Italia reperta.

#### EUPODIDÆ.

Scyphius maniacus, Guatemala, p. 17, pl. vi, fig. 2, Stoll, Acari in Biol. Centr.-Amer., n. sp.

#### GAMASIDÆ.

- Antennophorus, characterized, pt. lxx, No. 1; Berlese, Acari in Italia reperta.
- Celæno, characterized, pt. lxix, No. 3; Berlese, Acari in Italia reperta.
- Celænopsis uropodoides, British Honduras, p. 35, pl. xvi, fig. 4, & pl. xix, fig. 3, megisthanoides, Panama, p. 36, pl. xix, fig. 4, & pl. xx, fig. 1, STOLL, Acari in Biol. Centr.-Amer., n. spp.
- Cyrtolælaps, pt. lxiv, No. 3, falciger, Can., pt. lxiii, No. 4, nemorensis, Koch, No. 5, n. var. placida, No. 6, captator, n. sp., pt. lxviii, No. 8, spiricornis, Can., pt. lxviii, No. 9, cervus, Kram., No. 10; Berlese, Acari in Italia reperta.
- Dinychus, characterized, pt. lxx, No. 9; Berlese, Acari in Italia reperta. Epicrius, characterized, Can. & Fanz., pt. lxix, No. 10; Berlese, Acari in Italia reperta.
- Gamasus, pt. lxviii, No. 1, dentipes, Koch, No. 2, cornutus, Can., No. 3, furcatus, Can., No. 4, crassus, Kram., pt. lxix, No. 6, coleoptratorum, L., No. 7, rubescens, Can., No. 9, attenuatus, Koch, pt. lxiii, No. 7, n. var. minor, No. 8; Berlese, Acari in Italia reperta.
- G. pusillus, pt. lxix, No. 8; BERLESE, Acari in Italia reperta, n. sp.
- Hydrogamasus, n. g., pt. lxviii, No. 5, for H. litoralis, Can., No. 6; Ber-Lese, Acari in Italia reperta.
- Iphidosoma, n. g., pt. lxx, No. 11, for fimetarium, Can., ovatum, n. sp., pt. lxix, No. 5; Berlese, Acari in Italia reperta.
- Leelaps (Iphis) aculeifer, Can., and n. var. longior, No. 10, ostrinus, Koch, pt. lxvi, No. 4, halleri, Can., pt. lxvii, No. 6, bombicolens, No. 8, lignicola, Can., pt. xx, No. 2, krameri, Can., No. 4; L. (s.s.) holostaspoides, Can., pt. lxx, No. 13, uncinatus, Can., No. 15, Italy; Berlese, Acari in Italia reperta.

- L. (Iphis) miles, pt. lxiii, No. 9, semiscissus, No. 7, Italy, BERLESE, Acari in Italia reperta; L. (s.s.) complanatus, p. 383, in nest of Formica fusca, Moniez, Rev. Biol. iv; L. (s.s.) canestrini, pt. lxix, No. 1, myrmecophilus, No. 2, Italy, BERLESE, Acari in Italia reperta: n. spp.
- Leiognathus lacertinus, pt. lxx, No. 3, BERLESE, Acari in Italia reperta, n. sp. Megisthanus armiger, Berlese, Mexico, p. 34, pl. xix, fig. 1; Stoll, Acari in Biol. Centr.-Amer.
- M. giguntodes, Guatemala, p. 31, pl. xviii, figg. 1 & 2, STOLL, Acari in Biol. Centr.-Amer., n. sp.
- Neoberlesia, n. g. for N. equitans, n. sp., pl. lxii, No. 5; BERLESE, Acari in Italia reperta.
- Ophionyssus natricis, Gerv., pt. lxx, No. 12; Berlese, Acari in Italia reperta.
- Podocinum, characterized, pt. lxx, No. 10; Berlese, Acari in Italia reperta.
- Pachylælaps strigifer, Berl., pt. lxiv, No. 4, n. var. siculus, No. 5, pectinifer, Can., No. 6, n. var. siculus, pt. lxv, No. 10, Berlese, Acari in Italia reperta; P. heros, n. var. mexicanus, Mexico, p. 37, pl. xix, fig. 2, Stoll, Acari in Biol. Centr.-Amer.
- Pacilochirus, pt. lxx, No. 11, fucorum, De Geer, emarginatus, Koch, and spinipes, Koch, pt. lxix, No. 4; Berlese, Acari in Italia reperta.
- Polyaspis, characterized, pt. lxx, No. 8; BERLESE, Acari in Italia reperta. Seiodes, pt. lxx, No. 7, lustricinus, pt. lxx, No. 5, BERLESE, Acari in Italia reperta, n. sp.
- Stylorhinus rovennensis, Can., pt. lxx, No. 13; Berlese, Acari in Italia reperta.
- Uropoda spatulifera, n. sp., with Formica rufa, p. 383, Moniez, Rev. Biol. iv; U. lagena, n. sp., tridentina, Can., pt. lxx, No. 14, Berlese, Acari in Italia reporta.
- U. echinata, p. 28, pl. xvi, fig. 2, inequipunctata, p. 29, pl. xvi, fig. 3, discus, p. 29, pl. xvii, fig. 4, Guatemala, centro-americana, Nicaragua, p. 30, pl. xvii, fig. 1, piriformis, Mexico, p. 31, fig. 3, STOLL, Acari in Biol. Centr.-Amer., n. spp.
- Zercon acanthurus, Can., pt. lxx, No. 6; Berlese, Acari in Italia reperta.

#### HYDRACHNIDE.

- Arrhenurus decurtator, p. 520, MONIEZ, Rev. Biol. iv, n. sp.
- Feltria, n. g., p. 323, for F. minuta, n. sp., Rhostia, p. 323; KGENIKE, Zool. Anz. xv.
- Weltina, n. g. for W. microplicus, n. sp., p. 410; PIERSIG, Zool. Anz. xv. Zschokkea, n. g., p. 320, for Z. oblonga, n. sp., Rhostia, p. 32; Konier, Zool. Anz. xv.

#### Ixodiida.

The Ticks of this group are considered by MARX, P. E. Soc. Wash. ii, pp. 271-287, to rank as a suborder of *Acari*, for which the name *Cynorhastea* is proposed. This suborder is divided into two sections:

- Catastomata for the Eschutocephalidæ and Argusidæ, and Antistomata for the Hæmalastoridæ, Izodidæ, and Rhipistomidæ. These families, and all the genera they contain, are characterized.
- Amblyomma mixtum, C. Koch, Mexico, &c., p. 19, pl. xii, figg. 1 & 2; STOLL, Acari in Biol. Centr.-Amer.
- A. foreli, Guatemala, p. 21, pl. xii, fig. 3, & pl. xiv, fig. 3, crassipunctatum, Nicaragua, p. 22, pl. xiv, fig. 1, sabaneræ, Guatemala, p. 23, pl. xiv, fig. 2, STOLL, Acari in Biol. Centr.-Amer., n. spp.
- Boophilus, n. g. for B. bovis, Riley; CURTICE, J. Comp. Med. Vet. Arch. 1891.
- Ixodes vespertilionis, Koch, pt. lxi, No. 9, hexagonus, Leach, No. 10, Italy; BERLESE, Acari in Italia reperta.
- I. varani, Natal, pp. 10-14, LEWIS, J. Quek. Club, v; I. boarum, Guatemala, p. 18, pl. xiii, fig. 1, & pl. xiv, fig. 4, Stoll, Acari in Biol. Centr.-Amer., n. spp.

#### NICOLETIELLIDÆ.

Nicoletiella neotropica, Guatemala, p. 27, pl. xvi, fig. 1, STOLL, Acari in Biol. Centr.-Amer., n. sp.

#### ORIBATIDÆ.

- Belba, pt. lxiii, No. 3, pectinata, Mich., pt. xli, No. 8; Berlese, Acari in Italia reperta.
- Discopoma, Can., pt. lxviii, No. 7; Berlese, Acari in Italia reperta.
- Hermanna, pt. lxvii, No. 10, nana, Nic., pt. lxiii, No. 1; Berlese, Acari in Italia reperta.
- Hoplophora, Koch, pt. lxvii, No. 5, magna, Nic., pt. lxvii, No. 9; Berlese, Acari in Italia reperta.
- H. retalteca, Guatemala, p. 27, pl. xv, fig. 4, STOLL, Acari in Biol. Centr.-Amer., n. sp.
- Oppia tibialis, Nic., pt. lxiv, No. 1, n. var. sicula; ucorum, Koch, No. 2; Berlese, Acari in Italia reperta.
- Oribates mucronatus, Can., pt. lxii, No. 8; Berlese, Acari in Italia reperta.
- O. centro-americana, Panama, Honduras, p. 24, pl. xv, fig. 1, nigrifrons, Belize and Guatemala, p. 25, pl. xv, fig. 2, Stoll, Acari in Biol. Centr.-Amer., n. spp.
- Zetorchestes, pt. lxiii, No. 2; BERLESE, Acari in Itulia reperta.

#### PHYTOPTIDÆ.

- Anthocoptes, n. g. for P. loricatus, galeatus, heteroproctus, &c., p. 16; NALEPA, SB. Ac. Wien, 1892.
- A. platynotus, p. 191, NALEPA, SB. Ak. Wien, 1892, n. sp.
- Cecidophyes syriacus, p. 158, FOCKEU, Rev. Biol. iv; C. minor, p. 16, gemmarum, p. 128; NALEPA, SB. Ak. Wien, 1892: n. spp.

- Monaulax, n. g., p. 191, for M. sulcatus, n. sp., p. 192; NALEPA, SB. Ak. Wien, 1892.
- Oxypleurites, n. subg. of Tegonotus, for O. (T.) trouessarti, p. 330, pl. xiii, figg. 3 & 4, serratus, p. 333, pl. xiii, figg, 7-9, n. spp., and heptacanthus, Nal., p. 335, pl. xiii, figg. 10-12; Nalepa, Zool. Jahrb. vi.
- Phyllocoptes magnirostris, parvus, p. 128, populinus, p. 16, gracilipes, compressus, gigantorhynchus, comatus, p. 191, Nalepa, SB. Ak. Wien, 1892, n. spp.
- Phytocoptes longifilis, p. 380, cytisicola, p. 6, CANESTRINI, Atti Soc. Ven.-Trent. xii; rostratus, Syria, p. 232, fig., FOCKEU, Rev. Biol. iv: n. spp.
- Phytoptus vitalbæ, p. 79, breviceps, p. 80, Canestrini, Bull. Soc. Ven.-Trent. v; P. xylostei, p. 377, peucedani, p. 378, sanguisorbæ, p. 379, Canestrini, Atti Soc. Ven.-Trent. xii; P. lycii, p. 837, carueli, p. 838, effusus, p. 838, Canestrini, Atti Ist. Venet. l; P. eryugii, p. 5, Canestrini, op. cit. li; P. curvatus, p. 152, fig., barroisi, p. 154, fig., ephedræ, p. 155, fig., orientalis, p. 156, fig., fusiformis, p. 231, fig., Syrin, Fockeu, Rev. Biol. iv; P. eucricotes, triradiatus, p. 128, Nalepa, SB. Ak. Wien, 1892; P. pilosellus, tenellus, glaber, gibbosus, alpestris, pp. 190 & 191, Nalepa, t.c.; P. cladophthirus, p. 16, Nalepa, SB. Ak. Wien, 1892: n. spp.
- P. tilia, n. var. liosoma, p. 16; NALEPA, SB. Ak. Wien, 1892.
- Tegonotus, n. gen., pp. 327-329, for T. carinatus, n.sp., p. 329, pl. xiii, figg. 1 & 2; NALEPA, Zool. Jahrb. vi.
- T. fastigatus, p. 332, pl. xiii, figg. 5-7, Nalepa, Zool. Jahrb. vi; T. salicobius, p. 128, Nalepa, SB. Ak. Wien, 1892: n. spp.
- Trimerus trinotus, p. 191, NALEPA, SB. Ak. Wien, 1892, n. sp.

### SARCOPTIDÆ.

- Canestrinia procrusti, Berl., pt. lxv, No. 9; Berlese, Acari in Italia reperta.
- Chorioptes or Symbiotes, question of identity; Buysson, Bull. Soc. Ent. Fr. 1891, p. clx.
- Megninia oscinum, Koch, pt. xlv, No. 3; Berlese, Acari in Italia reperta.
- M. pteroglossorum, Guatemala, p. 40, pl. xxi, fig. 5, STOLL, Acari in Biol. Centr.-Amer., n. sp.
- Proctophyllodes glandarius, Koch, pt. lxv, No. 7; BERLESE, Acari in Italia reperta.
- P. sialiarum, Guatemala, p. 42, pl. xvi, figg. 3 & 4, Stoll, Acari in Biol. Centr.-Amer., n. sp.
- Pterocolus bifurcatus, Trouess., pt. lxv, No. 8; Berlese, Acari in Italia reperta.
- Pterolichus securiger, Rob., pt. lxv, No. 4, proctogamus, No. 5, rhebergi, Can. & Berl., No. 6; Berlese, Acari in Italia reperta.

- P. momotorum, Guatemala, p. 41, pl. xxi, figg. 1 & 2, Stoll, Acari in Biol. Centr.-Amer., n. sp.
- Pteronyssus fuscus, Nitz., pt. lzv. No. 2; Berlese, Acari in Italia reperta.

#### TROMBIDITOR.

- Cheyletiella chanayi, Berl. & Trouess., pt. lxvi, No. 3; Berlese, Acari in Italia reperts.
- Gekobia insignis, pt. lxvi, No. 1, loricata, No. 2, BERLESE, Acari in Italia reperta, n. spp.
- Rhyncholophus regalis, Koch, pt. lxii, No. 6; Berlese, Acari in Italia reperta.

#### TYROGLYPHIDÆ.

- Glyciphagus peregrinus, pt. lxv, No. 1, BERLESE, Acari in Italia reperta, n. sp.
- Tyroglyphus wasmanni, p. 397, Moniez, Rev. Biol. iv, n. sp.



# XII. MYRIOPODA AND PROTOTRACHEATA.

BY

## R. I. Pocock.

#### MYRIOPODA.

I. TITLES, p. 1.

III. DISTRIBUTION, p. 3.

II. ANATOMY, PHYSIOLOGY, PHYLOGENY, &c., p. 3. IV. Systematic, p. 3.

PROTOTRACHEATA, p. 6.

# MYRIOPODA.

#### I. TITLES.

- Berlese, A. Acari, Myriopoda et Scorpiones hucusque in Italia reperta.

  Padova. Pts. lxii, lxiv, lxvi, & lxvii.
- BERNARD, H. M. An endeavour to show that the trachese of the Arthropoda arose from setiparous sacs. Zool. Jahrb. v, p. 511.
- Broleman, H. W. Contributions a la faune Myriapodologique Mediterranéene. Dix espèces nouvelles. Ann. Soc. L. Lyon, xxix, pp. 7-40, pls. i-iii.
- CAMUS, F. Note sur la présence de Geophilus submarinus, &c., sur le côte de Préfaulle, Loire Inférieure. Bull. Sci. Nat. l'Ouest, Nantes, ii, No. 1, pp.
- CHILD, C. M. The Functions of the Nervous System in the Myriopoda.

  Am. Nat. xxvi, pp. 1051-1055.

- · 2 Myr. XII. MYBIOPODA AND PROTOTBACHEATA.
  - HERBST, C. Beiträge zur Kenntniss der Chiloroden. Drüsen: Coxalorgan, Gefasssystem, u. Eingeweidennervensystem, with 5 plates. Bibl. Zool. pt. ix, pp. 1-42.
- KOVALEVSKY, A. Sur les organes excréteurs des Arthropodes terrestres. Congr. Internat. Zool. pt. i, p. 187.
  Myriopoda, p. 205.
- LATZEL, R. Note sur quelques Myriapodes de Tahiti. Bull. Soc. Z. Fr. xvii, No. 7, pp. 185 & 186.
- Pocock, R. I. (1) On the *Myriopoda* and *Arachnida* collected by Dr. Anderson in Algeria and Tunisia. P. Z. S. 1892, pt. i, pp. 24-28.
- —. (2) Supplementary notes on the Arachnida and Myriopoda of the Mergui Archipelago, with descriptions of some new species from Spain and Malaysia. J. L. S. xxiv, pp. 319-326.
- ——. (3) Report upon two collections of Myriopoda sent from Ceylon by Mr. E. E. Green, and from various parts of Southern India by Mr. Edgar Thurston of the Government Central Museum, Madras. J. A. S. (Bombay) vii, No. 2, pp. 131-174.
- ---. (4) Arachuida and Myriopoda in W. L. Distant's Naturalist in the Transvaal. London (R. H. Porter): 1892, pp. 179-184.
- PRENANT, A. Les corps intermédiaires de Flemming dans les cellules séminales de la Scolopendre et de la Lithobie. C.R. Soc. Biol. (9) iv, pp. 172-176.
- Sinclair, F. G. A new mode of respiration in the *Myriopoda*. P. B. Soc., No. 303, pp. 200 & 201; also in Ann. N. H. (6) ix, p. 263.
- Verhoeff, C. (1) Vorläufige Mittheilung über eine neue deutsche Chordeuma-Art. Zool, Anz. xv, p. 109.
- ---. (2) Neue Diplopoden der paläarktischen Region. T. c. pp. 377 & 390, &c.
- —... (3) Ein Beitrag zur Kenntniss der gattung Chordeuma (Diplopoda) und einige Notizen zur deutschen Diplopoden Fauns. B. E. Z. xxxvii, pp. 7-14.
- —... (4) Bearbeitung der von A. F. Moller auf der Insel St. Thomas gesammelten Chilopoden und Diplopoden. T. c. pp. 192 & 193.
- —. (5) Zur Kenntniss der Analpleurendrüsen bei Scolopendriden, T. c. pp. 203-208.
- ZOGRAF, N. Sur l'origine et les parentes des Arthropodes principalement des Arthropodes tracheates. Congr. Internat. Zool. 1892, pt. i, p. 278, &c.

## II.—ANATOMY, PHYSIOLOGY, PHYLOGENY, &c.

Origin of tracheate Arthropods; ZOGRAF.

Origin of tracheal tubes; Bernard.—Respiration of Scutigera; SINCLAIR.

Nervous system, functions of; CHILD.

Seminal cells of Scolopendra and Lithobius; PRENANT.—Pleural glands of Scolopendridæ; VERHOEFF (5).—Glands, coxal organs, &c., of Chilopoda; HERBST.—Excretory organs; KOVALEVSKY.

## III.—DISTRIBUTION.

## Palæarctic Region.

W. Europe: various localities; VERHOEFF (1-3).

Italy: Berlese, Broleman. Coast of France: Camus.

Algeria and Tunisia: Pocock (1).

## Ethiopian Region.

St. Thomas (Gulf of Guinea): VERHOEFF (4). Transvaal: POCOCK (4).

## Oriental and Australian Regions.

India and Ceylon: Pocock (3).

Mergui: Pocock (2).

Malaysia (Siam, Borneo, &c.): POCOCK (2).

Tahiti : LATZEL.

## IV.—SYSTEMATIC.

## CHILOPODA.

#### SCUTIGERIDE.

Scutigera, Lam., characterized with figures of details, pt. lxvi, No. 7; Berlese, Myriopoda in Italia reperta.

#### LITHOBUDE.

Henicops, Newp., characterized, pt. lxvi, No. 8, cæculus, Brol., Milan, No. 9; Berlese, Myriopoda in Italia reperta.

Lithobius lapidicola, Mein., pt. lxvi, No. 10, lucifugus, L. Koch, pt. lxvii, No. 3, dentatus, C. Koch, No. 4, Italy; Berlese, Myriopoda in Italia reperta.

L. acuminatus, Lombardy, p. 7, pl. i, fig. 1, Broleman, Ann. Soc. L. Lyon, xxix; L. scutigeroides, St. Thomas (Gulf of Guinea), p. 201, Verhoeff, B. E. Z. xxxvii: n. spp.

#### SCOLOPENDRIDÆ.

- Cormocephalus pygmæus, Madras, p. 140, Pocock, J. A. S. (Bombay) vii, n. sp.
- Otostigma ceylonicum, splendens, and morsitans compared, p. 140; Pocock, J. A. S. (Bombay) vii. O. productum, Karsch, St. Thomas (Gulf of Guinea), redescribed, p. 199; Verhoeff, B. E. Z. xxxvii.
- O. owenii, Mergui, p. 319, Pocock, J. L. S. xxiv, n. sp.
- Heterostoma spinosum, paucispinosum, and trists compared, pp. 138 & 139; POCOCK, J. A. S. (Bombay) vii.

#### GEOPHILIDÆ.

- Chatechelyne, Mein., characterized, pt. lxvi, No. 5; Berlese, Myriopoda in Italia reperta.
- Dignathodon, Mein., characterized, pt. lxvi, No. 6; Berlese, Myriopoda in Italia reporta.

## DIPLOPODA.

#### POLYXENIDÆ.

Polyxenus ceylonicus, Punduloya (Ceylon), p. 142, Pocock, J. A. S. (Bombay), vii, n. sp.

#### GLOMERIDE.

- Glomeris characterized, pt. lxvii, No. 1, marmorata, C. Koch, pt. lxii, No. 1, varieties, No. 2, hexasticha, Br., No. 3, var. pallidior, No. 4, multistriata, C. Koch, No. 10, aurita, C. Koch, pt. lxiv, No. 7, marginata, Vill., No. 8, connexa, C. Koch, No. 9, var. tenebrosa, Latz., No. 10, from Italy; Berlese, Myriopoda in Italia reperta.
- G. pusilla, No. 9, Italy, Berlese, Myriopoda in Italia reperta; G. occultocolorata, Coimbra, p. 390, Verhoeff, Zool. Anz. xv: n. spp.
- Gervaisia, Waga, characterized, with figures of details, pt. lxvii, No. 2; Berlese, Myriopoda in Italia reperta.
- Zephronia brandti, Humb., syn. Z. chitonoides and rugulosa, Butl., Ceylon, pp. 143 & 144; inermis, Humb., syn. corrugata and leopardina, Butl., Ceylon, pp. 144 & 145; heterostictica, Newp., syn. lutescens and atrisparsa, Butl., Madras, pp. 145-147; POCOCK, J. A. S. (Bombay), vii.

#### Polydesmidæ.

- Polydesmus subulifer, n. sp., syn. P. complanatus, Auct. Ital., p. 10, pl. i, fig. 4; brevimanus, p. 13, fig. 3, fissilobus, p. 216, fig. 2, bigeniculatus, p. 20, fig. 5, n. spp., Lombardy, Broleman, Ann. Soc. L. Lyon, xxix; P. lusitanicus, Coimbra, p. 389, Verhoeff, Zool. Anz. xv, n. sp.
- Brachydesmus insculptus, Algeria, p. 27, with fig., POCOCK, P. Z. S. 1892, n. sp.
- Strongylosoma cingalense and skinneri compared, pp. 150 & 151; POCOCE, J. A. S. (Bombay) vii.
- S. setosum, Mergui, p. 320, Pocock, J. L. S. xxiv; S. greeni, Ceylon, p. 140, pl. ii, fig. 14, phipsoni, Calcutta, p. 151, pl. i, fig. 4, & pl. ii, fig. 13, jerdoni, Madras, p. 152, Pocock, J. A. S. (Bombay) vii; S. lusitanicum, Portugal, p. 383, bertkaui, Coimbra, p. 389, Verhoeff, Zool. Anz. xv: n. spp.
- Leptodesmus tanjoricus, Tanjore, p. 147, pl. i, fig. 3, POCOCK, J. A. S. (Bombay) vii, n. sp.
- Cryptodesmus ceylonicus, p. 153, pl. ii, fig. 2, greeni, p. 154, pl. ii, fig. 3, Ceylon, Pocock, J. A. S. (Bombay) vii, n. spp.
- Pyrgodesmus, n. g., p. 155, for P. obscurus, n. sp., Ceylon, p. 155, pl. ii, fig. 1; Pocock, J. A. S. (Bombay) vii.

## CHORDEUMIDE.

- Atractosoma lombardicum, p. 22, pl. ii, fig. 6, tellinense, p. 25, fig. 7, Lombardy, Broleman, Ann. Soc. L. Lyon, xxix, n. spp.
- Chordeuma germanicum, Rhineland, p. 109, VERHOEFF, Zool. Anz. xv, also p. 10, id. B. E. Z. xxxvii, n. sp.
- Craspedosoma dentatum, p. 28, pl. ii, fig. 8, Lombardy, Broleman, Ann. Soc. L. Lyon, xxix, n. sp.

#### LYSIOPETALIDA.

- Stemmiulus, Gervais, characterized, p. 156; POCOCK, J. A. S. (Bombay) vii.
- S. ceylonicus, Ceylon, p. 157, pl. i, fig. 2, POCOCK, J. A. S. (Bombay) vii, n. sp.

#### POLYZONIDÆ.

Siphonophora humberti, Ceylon, p. 173, Pocock, J. A. S. (Bombay) vii, n, sp.

#### JULIDÆ.

- Julus intermedius, p. 30, pls. ii & iii, fig. 9, cantonii, p. 36, pls. ii & iii, fig. 10, n. spp.; descriptions and figures of varieties of J. luridus, C. Koch, and J. londinensis, Leach, pp. 38 & 39, pls. ii & iii, figg. 12 & 13; BROLEMAN, Ann. Soc. L. Lyon, xxix: J. alemannicus, n. sp., Heidelberg, p. 377; synopsis of the males of the fallax group, p. 379; J. molleri, p. 381, frisioides, p. 383, n. spp., Coimbra; Verhoeff, Zool. Anz. xv.
- Hemipodiulus, n. subg. of Julus, for J. (H.) karschi, n. sp. (? syn. J. lusitanicus, Karsch), Portugal; Verhoeff, p. 380, Zool. Anz. xv.
- Spirostreptus malabaricus, Gerv., Kortallum, pp. 158 & 159, nigrolabiatus, Newp., Madras, p. 159, pl. i, fig. 7, & pl. ii, fig. 5, lankaensis, Humb., Ceylon, p. 165, pl. ii, fig. 6; Pocock, J. A. S. (Bombay) vii.
- S. asthenes, jerdoni, Madras, p. 161, centrurus, p. 162, pl. ii, fig. 4, insculptus, Ceylon, p. 163, pl. i, fig. 8, Pocock, J. A. S. (Bombay) vii; S. bowringii, Siam, p. 321, fig. 1, perakensis, Perak, p. 322, fig. 2, patricii, Batavia, fig. 3, hosei, p. 323, fig. 4, everettii, N. W. Borneo, p. 324, fig. 5, dulitianus, Mt. Dulit, p. 325, fig. 6, baluensis, Mt. Kina Balu, p. 326, fig. 2, Pocock, J. L. S. xxiv; S. transvaalicus, Pretoria, p. 181, fig., Pocock, in Distant's Naturalist in Tranvaal; S. molleri, St. Thomas (Gulf of Guinea), p. 193, pl. v, fig. 199, Verhoeff, B. E. Z. xxxvii: n. spp.
- Odontopyge pretoriæ, Pretoria, p. 183, Pocock, in Distant's Naturalist in Transvaal, n. sp.
- Spirobolus carnifez, Fabr., syn. ruficollis, Newp., Madras, and Ceylon, p. 166, pl. ii, fig. 9, goësii, Porath, syn. dominicz, Pocock, Madras, p. 167; Pocock, J. A. S. (Bombay) vii.
- S. thurstoni, p. 167, pl. i, fig. 9, & pl. ii, fig. 8, uroceros, p. 169, pl. ii, fig. 7, Madras, greeni, pl. i, fig. 10, longicornis, pl. ii, fig. 11, p. 170, longicollis, p. 172, pl. ii, fig. 10, Ceylon, Pocock, J. A. S. (Bombay) vii; S. nannodes, Tahiti, p. 186, LATZEL, Bull. Soc. Ent. Fr. xvii: n. spp.

# PROTOTRACHEATA.

- Bernard, H. M. The Apodida: a Morphological Study. Nature Series. London (Macmillan & Co.): 1892.
- Section xvi of this work is set apart for the consideration of Peripatus.
- DENDY, A. Further observations on the eggs of *Peripatus*. Austr. Ass. Adv. Sci. 1892, pp. 10-14.
- —. On the Oviparity of Peripatus leuckartii. P. R. Soc. Vict. iv, pt. i, p. 31.

- [DENDY, A.] Further Notes on the Oviparity of the larger Victorian *Peripatus*, generally known as *P. leuckartii*. P. Linn. Soc. N.S.W. vii, pt. ii, pp. 267-276; also Ann. N. H. (6) x, p. 136.
- FLETCHER, J. J. A Viviparous Australian Peripatus (P. leuckartii, Sänger). P. Linn. Soc. N.S.W. vii, pp. 179-196.
- GRABHAM, M., & COCKERELL, T. D. A. Peripatus rediscovered in Jamaica. Nature, xlvi, p. 514.

  P. jamaicensis. n. sp.
- Pocock, R. I. Peripatus from St. Vincent. T. c. p. 100. Rediscovery of P. juliformis.



# XIII. INSECTA.

By D. SHARP.

THE number of titles in this RECORD is 1026, the number for 1891 being 974.

During the year commencements have been made with two works that will, no doubt, be of much use when completed, viz., Dalla Torre's (198) Catalogue of Hymenoptera and Kirby's Catalogue of Moths (496). The latter work resembles its well-known predecessor, the "Synonymic Catalogue of Butterflies," except that it has not been thought necessary to give the synonymy of the European species in full: a decision that will, no doubt, be considered wise in view of the fact that this volume extends, even as it is, to nearly 1000 pages.

Two parts of the Entomology of Grandider's work on Madagascar have been published. They both relate to Hymenoptera. One of them, by Forel (287), is an excellent systematic treatise on the Ants: the other is by DE Saussure (793); it treats of the rest of the Aculeate Hymenoptera, and proposes some modifications in the classification of the Fossoria. Some Terebrantia are figured on the plates, but are not alluded to in the letterpress.

Considerable progress has been made with the Biologia Centrali-Americana. But to us in Britain, perhaps, the most interesting faunistic work of the year is the first volume of Hampson's Moths (384), forming part of the valuable "Fauna of British India," and published under the auspices of the Secretary of State for India. This volume is well executed, and will augment the good reputation of the series of which it forms a part. All entomologists will wish that the other orders of Insects could be dealt with in due course, but we fear that material sufficient for the purposes of such a work does not yet exist in collections.

The papers by Forel (293) and Wroughton (1018) on the Ants of India should be noticed, not only on account of the amount of information they give on the subject, but also as showing what an extensive field of work for entomologists exists in India.

MEYRICK has given us another instalment (603) of his arrangement of Lepidoptera, on the basis of wing-neuration; and Spuler also gives us

(881) a study on this subject, but finds himself not quite able to agree with his predecessors in this field of enquiry.

HOFFBAUER has made a very interesting contribution (405) to a know-ledge of the structure of the elytra of *Coleoptera*: this memoir, and RILEY'S study of *Pronuba* and *Yucca* (742), will, no doubt, be considered as amongst the most attractive of the papers enumerated in this RECORD. So much interest, too, attaches to the conclusions announced in GRASSI'S summary of his observations (359) on *Termitidæ*, that entomologists will wait for further details with some impatience.

The most interesting Insects described in 1892 are probably the two Blattids discovered by SIMON in caves in the Philippine Islands, and described by BOLIVAR (710) as forming the new genus he calls Nocticola.

HENKING (396) discusses a considerable variety of points in the early stages of development of the embryo, including the development in the parthogenetic eggs of some *Hymenoptera*; and we also record what we fear must be the last (355) of Graber's series of papers on the comparative embryology of Insects.

We owe thanks to TASCHENBERG for his apparently exhaustive treatise (915) on the history of the literature of Parthenogenesis.

The Recorder regrets he can form no opinion as to PORTSCHINSKY'S remarks on the biology of *Diptera* (698), and on the origin of striking coloration in *Lepidoptera* (700), owing to their being written entirely in the Russian language.

## I.—TITLES.\*

- ABEILLE, E. DE PERRIN. Notices entomologiques. Rev. d'Ent. xi, pp. 62-67. [Coleoptera.]
- Deux nouvelles espèces de Clérides. Bull. Soc. Ent. Fr. lxi, p. ccxxxvi. [Coleoptera.]
- ADELUNG, N. VON. Beiträge zur Kenntniss des tibialen Gehörapparates der Locustiden. Z. wiss. Zool. liv, pp. 316-349, pls. xiv & xv. [Orthoptera.]
- ALPHERAKY, S. Lepidoptera nova a Gr. Grum-Grshimailo in Asia centrali novissime lecta. Hor. Ent. Ross. xxvi, pp. 444-459.
  - ---. [See also Romanoff (779).]
- Aldrich, J. M. New species of Phora. Canad. Ent. xxiv, pp. 142– 146. [Diptera.]
- A new genus and species of Tabanida. Psyche, vi, p. 236.
   [Diptera.]

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

- Allard, E. Collection d'insectes formée dans l'Indo-Chine par M. Pavie, Consul de France au Cambodge. Coléoptères. Phytophages. N. Arch. Mus. Paris (3) iii, pp. 229-234.
- ALLUAUD, C. Deux nouvelles espèces de Coléoptères. Bull. Soc. Ent. Fr. lxi, pp. cexxxviii & cexxxix.
- Anderson, E. Notes on some Victorian Lepidoptera. Vict. Nat. ix, pp. 80-93.

Some interesting remarks on habits and variation.

- Andersson, J. Bidrag till Kännedomen om Nagelspinnarens (Aglia tau, L.) utvecklingshistoria. Ent. Tidskr. xiii, pp. 93-96. [Lepidoptera.]
- 11. André, E. Species des Hymenoptères. 41 Fasc. pp. 89-144, pls. iv, viii, & x.

Continuation of Chrysidida, by R. du Buysson.

- Matériaux myrmécologiques. Rev. d'Ent. xi, pp. 45-56.
   [Hymenoptera.]
- Voyage de M. Chaper à Borneo. Catalogue des Fourmis et description des espèces nouvelles. Mém. Soc. Zool. v, pp. 46-55. [Hymenoptera.]
- Angell, G. W. J. Two new species of Coleoptera. Ent. News, iii, p. 84.
- Arribalzaga, F. L. Dipterologia Argentina (Continuación). An. Soc. Arg. xxxiii, pp. 51-59, 111-122, 188-200, 236-254, and xxxiv, pp. 33-46, 173-192, & 241-280.
- 16. ASHMEAD, W. H. The Insect collections in the Berlin Museum. P. E. Soc. Wash. ii, pp. 205-211.

141,000 species of Hexapods.

- Notes on the genus Melittobia. T. c. pp. 228-232. [Hymenoptera.]
- On the discovery of the genus Eunotus in America. T. c. pp. 287 & 288. [Hymenoptera.]
- 18A. —. Notes on the genus *Enicocephalus*, Westwood, and description of a new species from Utah. *T. c.* pp. 328-333. [*Rhynchota.*]
- Notes on the Eucharids found in the United States. T. c. pp. 354-358. [Hymenoptera.]
- Discovery of the genus Cratæpus, Förster, in America, and the description of a new species. Canad. Ent. xxiv, p. 309. [Hymenoptera.]
- Three new Pambolids from the United States. Psyche, vi, pp. 289 & 290. [Hymenoptera.]

- [ASHMEAD, W. H.] Two new and remarkable Ophionids. Ent. News, iii, p. 105. [Hymenoptera.]
- Aurivillius, C. Nordens fjarilar. Handbok i Sveriges, Norges, Danmarks och Finlands Macrolepidoptera. 4to, pp. lii & 277, 50 plates and 35 woodcuts. Stockholm, 1888–1891.

He adopts three divisions for *Lepidoptera*, viz., Neolepidoptera, Mesolepidoptera, and Palæolepidoptera.

- Verzeichniss einer vom Herrn Fritz Theorin aus Gabun und dem Gebiete des Camerunflusses heimgebrachten Schmetterlingssammlung. II. Heterocera. Ent. Tidskr. xiii, pp. 181-200. [Lepidoptera.]
- Collection d'insectes formée dans l'Indo-Chine par M. Pavie,
   Consul de France au Cambodge. Coléoptères, Curculionines. N.
   Arch. Mus. Paris (3) iii, pp. 205-224.
- Eine neue Palæarktische Eulengattung Ent. Tidskr. xiii, p. 285. [Lepidoptera.]
- Leddjurens synförmåga. T. c. pp. 171-180.
   Relates chiefly to Exner's views on vision.
- Om slökorn-flugan (Oscinis frit, L.) ett gif-akt till sveriges Kornodlare. T. c. pp. 209-224. [Diptera.]
- 29. AUSTAUT, L. Deux Sphingides nouveaux de l'Asie orientale. Le Nat. 1892, pp. 68 & 69. [Lepidoptera.]
- 30. Baker, G. T. B. Notes on Lyozna (recte Thecla) rhymnus, tengstræmii, and pretiosa. Tr. Ent. Soc. 1892, pp. 27-31, pl. ii. [Lepidoptera.]
- Balbi, E. Diagnosi e descrizioni di due nuove specie di Coleotteri.
   Riv. Ital. Sci. Nat. xii, pp. 49 & 50.
- Contributione alla Fauna coleotterologica d'Europe. Cf. Wien. ent. Z. xi, p. 251.
- 33. Banks, N. A synopsis, catalogue, and bibliography of the Neuropteroid Insects of Temperate North America. Tr. Am. Ent. Soc. xix, pp. 327-373.
- 34. BARRETT, C. G. The Lepidoptera of the British Islands, London, 8vo.

Intended to include all the British Lepidoptera. Parts I-v have been at present published.

- BATAILLON, E. Sur le déterminisme physiologique de la metamorphose chez le ver-à-soie. C.R. cxv, pp. 61-64.
- 36. BATES, H. W. Additions to the Longicornia of Mexico and Central America, with remarks on some of the previously-recorded species. By the late Henry Walter Bates, with an Introduction by Frederick DuCane Godman. Tr. Ent. Soc. 1892, pp. 143-183, pls. v-vii. [Coleoptera.]

- [Bates, H. W.] Coléoptères du Bengale occidental. Seconde liste des Carabida. Ann. Ent. Belg. xxxvi, pp. 230-233.
- Viaggio de Leonardo Fea in Birmania e regioni vicine. XLIV,
   List of the Carabida. Ann. Mus. Genov. (2) xii, pp. 267-428.
   [Coleoptera.]
- The Naturalist on the River Amazons: reprint of the unabridged edition, with a memoir of the author by Edward Clodd. London, 8vo, 1892.
- Bateson, W. On Variation in the Colour of Cocoons of Eriogaster lanestris and Saturnia carpini. Tr. Ent. Soc. 1892, pp. 45-52. [Lepidoptera.]

Summary in J. R. Micr. Soc. 1892, p. 469.

- 41. —. On Variation in the Colour of Cocoons, Pupse and Larvee: further experiments. T. c. pp. 205-214. [Lepidoptera.]
- Bateson, W., & Brindley, H. H. On some cases of Variation in Secondary Sexual Characters, statistically examined. P. Z. S. 1892, pp. 585-594. [Coleoptera, Orthoptera.]
- BAUDI, F. Di una nuova specie del genere Tritoma, Geoffr. Nat. Sicil. xi, p. 121. [Coleoptera.]
- 44. BECKER, A. Neue Pflanzen- und Insektenentdeckungen in der Umgegend von Sarepta, und Zusammenstellung der Raupen und Käfer die nur von einer Pflanzenart, und zwei, drei, Pflanzenarten leben, die aber zu einer Familie gehören. Bull. Soc. Mosc. 1892, pp. 62-70.
- BECKER, TH. Eine neue Leptis aus der Schweiz. Wien. ent. Z. xi, p. 23. [Diptera.]
- 46. Beddard, F. E. Animal Coloration: an account of the principal facts and theories relating to the colours and markings of animals. London, 8vo, 288 pp., iv pls.

Includes much Entomology.

- Bedel, L. Synopsis des grands Hydrophiles. Rev. d'Ent. x, pp. 306-322. [Coleoptera.]
- Description de deux Coléoptères hétèromères du nord de l'Afrique. Ann. Mus. Genov. (2) x, pp. 794 & 795.
- Synopsis des Chironitis, Onitis et Bubas de la région méditerranéenne: extrait de la monographie des Onitides de G. van Lansberge. Abeille, xxvii, pp. 257-280. [Coleoptera.]
- Revision des Scarabæus palæarctiques. T. c. pp. 281-288.
   [Coleoptera.]
- Renseignements sur les mœurs et le développement des Meloidæ. T. c. pp. 235-250 & 304. [Coleoptera.]

- 52. Behr, H. H. Some Biological Peculiarities of the Phylloxera and a Method for utilising them for the Protection of Vineyards. Zoe, ii, pp. 305-310.
- 53. Belon, M. J. Viaggio di Leonardo Fea in Birmania e regioni vicine. xxxviii, Lathridiida. Ann. Mus. Genov. (2) x, pp. 877-880. [Coleoptera.]
- 54. Benton, F. How Far do Bees Fly? Ins. Life, iv, pp. 319-321.
  [Hymenoptera.]
- 55. Benzi, A. Contribuzione allo studio degli *Imenotteri* del Modenese e particolarmente delle Specie del Genere *Nomada*, con la descrizione d'una nuova Specie. Atti Soc. Mod. (3) xi, pp. 213-225.
- BERG, C. Nova Hemiptera, faunarum Argentines et Uruguayensis (Continuación.) An. Soc. Arg. xxxiii, pp. 5-11, 43-50, 65-72, 97-104, 151-165, and xxxiv, pp. 82-96, & 193-205.
- Eolus pyroblaptus, un nuevo destructor del trigo. T. c. pp. 60-62. [Coleoptera.]
- Canabalismo entre insectos. Op. cit. xxxiv, pp. 236-238.
   Bergenstamm, J. E. v. [See Brauer & Bergenstamm (119).]
- Berger, R. Beiträge zum Melanismus der Schmetterlinge. Soc. Ent. vi, pp. 180, &c.

States that melanism is much extending amongst Lepidoptera on the Continent of Europe.

- BERGROTH, E. Ueber Lasiopsis, Er., und verwandte Gattungen.
   Wien. ent. Z. xi, pp. 98-100. [Colsoptera.]
- —. Beitrag zur Tipuliden-Fauna der Schweiz. MT. Ges. Bern. 1890, pp. 131-138. [Diptera.]
- Description d'un genre nouveau de la sous-famille des Plataspidina. Bull. Soc. Ent. Fr. lxi, p. celv. [Rhynchota.]
- Plataspidinæ quatuor novæ. Wien. ent. Z. xi, pp. 171-173. [Rhynchota.]
- 64. —. Notes on the Nearctic Aradida. P. E. Soc. Wash. ii, pp. 332-337, and Additional note, pp. 337 & 338. [Rhynchota.]
- 65. —. Aradidæ novæ. Rev. d'Ent. xi, pp. 259-261. [Rhynchota.]
- 66. —. Aradidi dell' isola di Engano raccolti dal Dott. Elio Modigliani. Ann. Mus. Genov. (2) xii, pp. 806 & 807. [Rhynchota.]
- 67. —. Aradides du Bengale boréal. Ann. Ent. Belg. xxxvi, pp. 503. & 504. [Rhynchota.]
- 68. —. Rhynchota ethiopica. T. c. pp. 160-162.

- 69. [Bergroth, E.] Viaggio di Leonardo Fea in Birmania e regioni vicine. XLVII, Commentarius secundus de Aradidis in Burma et Tenasserim a L. Fea collectis. Ann. Mus. Genov. (2) xii, pp. 710– 717. [Rhynchota.]
- Nouvelle espèce du genre Megymenum. Ann. Ent. Belg. xxxvi, p. 504. [Rhynchota.]
- On an Indian Ant-mimicking Hemipteron. Ent. M. M. xxviii, p. 107. [Bhynchota.]
- 72. —. A new genus of Eusthening. T. c. p. 158. [Rhynchota.]
- BERNARD, H. M. An endeavour to show that the trachese of the Arthropoda arose from setiparous sacs. Zool. Jahrb. Anat. v, pp. 511– 524.
- 74. Berthoumieu, V. Nouvelles espèces d'Ichneumonides. Rev. d'Ent. xi, pp. 37-44. [Hymenoptera.]
- 75. BEUTENMÜLLER, W. On the earlier stages of some species of North American Moths. Bull. Am. Mus. iv, pp. 65-70. [Lepidoptera.]
- List of types of some species of Lepidoptera, described by Grote and Robinson, in the American Museum of Natural History. T. c. pp. 59-64.
- Some notes on transformations of Australian Lepidoptera, made by the late Henry Edwards, with notes and additions. T. c. pp. 71-74.
- Notes on the transformations of some species of Lepidoptera.
   By Henry Edwards and S. L. Elliot, edited with additions by. T. c. pp. 75-80.
- Bezzi, M. Contribuzione alla fauna Ditterologica della provincia di Pavia. Parte seconda. Bull. Ent. Ital. xxiv, pp. 64-84 & 97-151.
- Di alcuni Ditteri raccolti nel paese dei Somali dall' Ingegnere
   Bricchetti Robecchi. Ann. Mus. Genov. (2) xii, pp. 181-196.
- 81. BIDENKAP, O. Undersögelser over *Diptera brachycera* i Jarlsberg og Laurvigs amt Sommeren 1891. Ent. Tidskr. xiii, pp. 225-246.
- 82. BIGNELL, G. C. A new species of Cynipide. Ent. M. M. xxviii, p. 176. [Hymenoptera.]
- 83. Bigot, J. M. F. Diptères nouveaux ou peu connus. 37e partie. xLvI. Bombylidi. Ann. Soc. Ent. Fr. lxi, pp. 321-376.
- Catalogue of the *Diptera* of the Oriental region. Part 11,
   J. A. S. B. lxi, pp. 135-178; Part 111, t. c. pp. 178-236.
- Descriptions de Diptères nouveaux. Mém. Soc. Zool. v, pp. 602-691.
- 86. Description de trois nouveaux Diptères de l'Inde. Bull. Soc. Z. Fr. 1892, pp. 210 & 211.

The locality is Mahé.

- [Bigot, J. M. F.] Nova genera *Dipterorum*. Wien. ent. Z. xi, pp. 161 & 162.
- 88. Binet, A. Les racines du nerf alaire chez les Coléoptères. C.R. cxiv, pp. 1130-1132.
  - Summary in J. R. Micr. Soc. 1892, p. 471.
- Sur la structure d'un ganglion nerveux d'Insecte. Bull. Soc. Ent. Fr. lxi, p. clxxii.
  - ---. [See also Henneguy & Binet (399).]
- BLACKBURN, T. Notes on Australian Coleoptera, with descriptions of new species. Part x. P. Linn. Soc. N.S.W. (2) vi, pp. 479-550.
- 91. —. Notes on Australian Coleoptera, with descriptions of new species. Part x1. Op. cit. vii, pp. 65-152.
- Notes on Australian Coleoptera, with descriptions of new species. Part XII. T. c. pp. 283-300.
- Further Notes on Australian Coleoptera, with descriptions of new genera and species. xI. Tr. R. Soc. S. Austr. xv, pp. 20-73.
- Further Notes on Australian Coleoptera, with descriptions of new genera and species. XII. T. c. pp. 207-261.
- 95. —. Scientific Results of the Elder Exploring Expedition. Coleoptera. Op. cit. xvi, pp. 16-61.
- BLAISDELL, F. E. Notes on Histeridæ observed in San Diego County.
   Zoe, iii, pp. 337-339. [Coleoptera.]
- A New Species of Coleoptera from California. Ent. News, iii, p. 51.
- 98. —..... New Coleoptera from California. T. c. pp. 241-243.
- BLANDFORD, W. F. H. Insects injurious to Conifera. J. R. Horticult. Soc. xiv, pp. 1-29 (separate copy).
- Sugar-cane Borers in the West Indies. Kew Bull. 1892, pp. 153-178, pl.
- 101. BLANCHARD, R. Sur les Œstrides américains dont la larve vit dans la peau de l'homme. Ann. Soc. Ent. Fr. lxi, pp. 109-154. [Diptera.]
- 102. —. Note additionelle sur les Œstrides américains dont la larve vit dans la peau de l'homme. Bull. Soc. Ent. Fr. lxi, pp. ccix-ccxi. [Diptera.]
- 103. BLATCH, W. G., & HORNER, A. C. Description of a New Species of Rhizophagus. Ent. M. M. xxviii, p. 303. [Coleoptera.]
- 104. BLATCHLEY, W. S. Two New Orthoptera from Indiana. Canad. Ent. xxiv, pp. 26 & 27.
- 105 —. Some Indian Acridiida. II. T. c. p. 28. [Orthoptera.]

- 106. [Blatchley, W. 8.] Entomologizing in Mexico. Ent. News, iii, pp. 111-114 & 131-136.
- 107. BLATHWAYT, L. Forficula auricularia. Int. J. Micr. (3) ii, pp. 7-16, pls. i & ii. [Orthoptera.]
- 108. BLATTER, P. Sur l'histologie des organes annexes de l'appareil mâle chez la *Periplaneta orientalis*. C.R. cxv, pp. 1332-1334.
- 109. BLOCHMANN, F. Ueber das Vorkommen von Bakterienähnlichen Gebilden in den Geweben und Eiern verschiedener Insekten. CB. Bakt. Parasit. zi, pp. 234-240.
- BOLIVAR, J. La Langosta migratoria de Chile. Act. Soc. Chili, ii, pp. 196-199. [Orthoptera.]
  - ---. [See also RAFFRAY & BOLIVAR (710).]
- BORRIES, H. De Danske Lophyrus-Arter og deres udbredelse. Ent. Med. iii, pp. 97-124, pl. ii. [Hymenoptera.]
- 112. Bourgeois, J. Lycides nouveaux on peu connus du Musée civique de Gênes. Deuxième mémoire, deuxième partie. Sous-tribu des Lycini (genuini). Ann. Mus. Genov. (2) xii, pp. 495-516. [Coleoptera.]
- 113. —... Coléoptères du Bengale occidental. Dascillides et Malacodermes. Ann. Ent. Belg. xxxvi, pp. 234-238.
- Descriptions de trois espèces nouvelles de Lycides de Borneo.
   Notes Leyd. Mus. xiv, pp. 39-42. [Coleoptera.]
  - ---. [See also DISTANT (210).]
- 115. Brancsik, K. Beiträge zur Kenntniss Nossibé's und dessen Fauna nach Sendungen und Mittheilungen des Herrn P. Frey. Jahresb. Ver. Trencsin. xiii, pp. 122-167.
- 116. Brauer, F. Über die aus Afrika bekannt gewordenen Œstriden und insbesondere über zwei neue von Dr. Holub aus Südafrika mitgebrachte Larven aus dieser Gruppe. SB. Ak. Wien, ci, pp. 4-16, pl. i. [Diptera.]
  - Summary in J. R. Micr. Soc. 1892, p. 600.
- 117. —. Ueber das sogenannte Stillstandstadium in der Entwicklung der Æstriden-Larven. Verh. z.-b. Wien, xlii, pp. 79-84. [Diptera.]
  - Summary in J. R. Micr. Soc. 1892, p. 358.
- 118. —... Ansichten über die Gattung Pachystylum, Macq., und Rückblicke auf die in den Denkschriften der Kais. Akademie erschienenen "Vorarbeiten zu einer Monographie der Muscaria, &c., von Prof. Fr. Brauer u. J. Edl. v. Bergenstamm." SB. Ak. Wien, ci, pp. 593-607. [Diptera.]

- 119. Brauer, F., & Bergenstamm, J. E. von. Die Zweiflügler des Kaiserlichen Museums zu Wien. v. Vorarbeiten zu einer Monographie der Muscaria schizometopa (exclusive Anthomyida). Pars 11. Denk. Ak. Wien, lviii, pp. 305-446.
- 120. Brendel, E. On the *Pselaphid* genus *Trimium*. Tr. Am. Ent. Soc. xix, pp. 165-168. [Coleoptera.]
- 121. —. Rhexidius. Ent. News, iii, pp. 11-13. [Coleoptera.]
- 122. Brenske, E. Beitrag zur Kenntniss der Gattungen Lepidiota und Leucopholis. Berl. Ent. Zeit. xxxvii, pp. 33-62. [Coleoptera.]
- 123. Die Arten der Coleopteren-Gattung, Brahmina, Bl. T. c. pp. 79-124.
- Neue Arten der Coleopteren-Gattung, Holotrichia (Lachnosterna). T. c. pp. 159-192.
- 125. —... Ueber einige neue Gattungen und Arten der Melolonthiden.
   Ent. Nachr. xviii, pp. 151-159. [Coleoptera.]
   BRINDLEY, H. H. [See BATESON & BRINDLEY (42).]
- 126. Brischke, C. G. A. Einige neue, oder für Westpreussen neue, Hymenopteren und Dipteren. Schr. Ges. Danz. (n.s.) viii, pp. 19-22.
- 127. Bericht über eine Excursion ins Radaunethal bei Babenthal während des Juni 1890. T. c. pp. 23-56. [Insecta.]
- 128. Brisout, C. de Barneville. Description d'une Corticaria nouvelle de France. Rev. d'Ent. xi, p. 68. [Coleoptera.]
- 129. Brodle, W. Canadian Galls and their Occupants. Canad. Ent. xxiv, pp. 12-14.
- Canadian Galls and their Occupants. Eurosta solidaginis,
   Fitch. T. c. pp. 137-139.
- 131. Bromilow, F. Butterflies of the Riviera. A short Account of the Rhopalocera of the Maritime Alps. Nice, 1892, 115 pp., 8vo. [Lepidoptera.]
- 132. BRONGNIART, C. Monographie du genre Palophus (Orthoptères de la famille des Phasmiens). N. Arch. Mus. Paris (3) iii, pp. 193-204, pls. viii & ix.
- 133. —. Collection d'insectes formée dans l'Indo-Chine par M. Pavie, Consul de France au Cambodge. Coléoptères, Longicornes. T. c. pp. 237-254, pl. x.
- 134. —. Monographie du genre Eumegalodon (Orthoptères de la famille des Locustides). T. c. pp. 277-286, pl. xii.
- 135. Bruner, L. Report upon Insect Depredations in Nebraska. Bull. Dep. Agric. Ent. xxvi, pp. 9-12.
- 136. BRUNER, L., COQUILLETT, D. W., & OSBORN, H. Reports on the Damage by destructive Locusts during the season of 1891. Bull. Dep. Agric. Ent. xxvii, pp. 1-64. [Orthoptera.]

- 137. Brunner, C. von Wattenwyl. Notizen über die Orthopteren-Fauna Ceylon's. Ent. Nachr. xviii, pp. 337-340.
- 138. Brunner, C. von Wattenwyl, & Redtenbacher, J. On the Orthoptera of the Island of St. Vincent, West Indies. P. Z. S. 1892, pp. 196-220, pls. xv-xvii.
  - With an introductory notice by D. SHARP.
- 139. Buchenau, F. Die "springenden Bohnen" aus Mexiko. Dritter Beitrag. Abh. Ver. Brem. xii, pp. 277-290.
- 140. Bugnion, E. Résumé des recherches de M. Jules Gonin sur la Métamorphose des Lépidoptères. MT. Schw. ent. Ges. viii, pp. 403-407.
- BUTLER, A. G. On a collection of Lepidoptera from Sandakan,
   N. E. Borneo. P. Z. S. 1892, pp. 120-133, pl. vi.
- 142. —... On the genus Hypocala, a group of Noctuid Moths. Ann. N. H. (6) x, pp. 17-22. [Lepidoptera.]
- 143. —. Description of a new genus and species of African Moths. T. c. pp. 295 & 296. [Lepidoptera.]
- 144. —... On the Noctuid genera allied to Hypætra of Guenée. T. c. pp. 297-300. [Lepidoptera.]
- 145. —. Revision of the *Noctuid* genus *Melipotis*, Hub., with descriptions of two new species. *T. c.* pp. 315-327. [Lepidoptera.]
- 146. Description of a new Moth of the genus Anaphe, from Madagascar, with a note on the natural position of the genus. T. c. p. 398. [Lepidoptera.]
- 147. BUTLER, A. G., & BOTHSCHILD, W. On a new and also on a little-known species of *Pseudacræa* in the collection of the Hon. Walter Bothschild. Tr. Ent. Soc. 1892, pp. 201 & 202, pl. x. [Lepidoptera.]
- 148. Buysson, R. du. Sur les glandes à venin des Ichneumonides. Rev. d'Ent. xi, pp. 257 & 258, pl. i. [Hymenoptera.]
  - ----. [See also André (11).]
- 149. BUYSSON, R. DU, & MARSHALL, T. A. Voyage de M. E. Simon au Venezuela. Hyménoptères. Ann. Soc. Ent. Fr. lxi.
- 150. CALVERT, P. P. Preliminary notes on some African Odonata. Tr. Am. Ent. Soc. xix, pp. 161-164.
- 151. Cameron, P. Hymenoptera orientalis; or contributions to a knowledge of the Hymenoptera of the Indian Zoological region: Part IV. Mem. Soc. Manch. (4) v, pp. 97-137, pl. i.
- 152. CANDÈZE, E. Elatérides recueillis en Birmanie en 1888, par M. L. Fea. 2º article. Ann. Mus. Genov. (2) x, pp. 771-793. [Coleoptera.]

- 153. [CANDÈZE, E.] Elatérides recueillis par M. Modigliani dans l'île d'Engano en Mai et Juin 1891. Op. cit. xii, pp. 795–800. [Coleoptera.]
- 154. —. Viaggio di Lamberto Loria nella Papuasia orientale. VI. Elatérides recueillis par M. Loria en 1889–1891, dans la Nouvelle-Guinée méridionale et régions voisines. T. c. pp. 801–805. [Coleoptera.]
- 155. —, Deuxième note sur les Élatérides de Chota-Nagpore. Ann. Ent. Belg. xxxvi, pp. 480-495. [Coleoptera.]
- 156. —... Trois Elatérides nouveaux. Notes Leyd. Mus. xiv, pp. 9 & 10. [Coleoptera.]
- 157. CARLET, G. Sur le mode d'union des anneaux de l'abdomen (articulation en zigzag) chez les Hyménoptères. C.R. cxiv, p. 766. Summary in J. R. Micr. Soc. 1892, p. 357.
- 158. CARLINI, A. DE. Rincoti raccolti nel paese dei Somali dall' Ing. Bricchetti-Robecchi, con note su altri dell' Africa centrale. Ann. Mus. Genov. (2) xii, pp. 527–538.
- 159. CARPENTER, G. H. Reports on the Zoological collections made in Torres Straits by Prof. A. C. Haddon. Rhynchota from Murray island and Mabuiag. P. R. Dublin Soc. (n.s.) vii, pp. 137-146, pls. xii & xiii.
- 160. CARPENTIER, L. Observations sur les parasites du Liparis salicis, Dup. Bull. Soc. L. Nord. Fr. x, pp. 35-40. [Lepidoptera, Hymeno-ptera.]
- 161. Casey, T. L. Coleopterological Notices. III. Ann. N. York Ac. vi, pp. 9-214.
  - This is in large part devoted to a revision of the N. American Cistelidæ.
- 162. ——. Coleopterological Notices. IV. T. c. pp. 359-712.

  This part is devoted to Rhynchophora.
- 163. CHAMPION, G. C. Note on *Pentaria badia*, Rosh., with description of a second species from the Pyrénées-orientales. Ent. M. M. xxviii, p. 108. [Coleoptera.]
  - ---. [See also Godman & Salvin (347).]
- 164. CHAPMAN, T. A. The life-history of Lampronia capitella. Ent. M. M. xxviii, pp. 297-300. [Lepidoptera.]
- 165. CHATIN, J. Sur l'origine et la formation du revêtement chitineux chez les larves de *Libellules*. C.R. exiv, pp. 1135-1138.
  Summary in J. R. Micr. Soc. 1892, p. 473.
- 166. —. Sur le processus général de la cuticularisation tégumentaire chez les larves de Libellules. Bull. Soc. Philom. (8) iv, pp. 105 & 106.

167. CHAUMAT, H. L'œuvre scientifique du Dr. G. Carlet. Bull. Soc. Isère (4) i, pp. 403-451.

Includes abstracts of each of his memoirs.

- 168. CHITTENDEN, F. H. Notes on the Food Habits of some Species of Chrysomelida. P. E. Soc. Wash. ii, pp. 261-267. [Coleoptera.]
- 169. CHOBAUT, —. Un nouveau Rhipidius du Mont Ventoux. Mem. Ac. Vaucluse, pp. 213-221, pl. [Coleoptera.]
- 170. CHOLODKOVSKY, N. On the Morphology and Phylogeny of Insects. Ann. N. H. (6) x, pp. 429-451.
- A translation of a portion of the paper alluded to last year in Zool. Rec. Ins. Titles (145).
- Zur Kenntniss der männlichen Geschlectsorgane der Dipteren.
   Zool. Anz. xv, pp. 178–180.
- Translation in Ann. N. H. (6) x, pp. 268-270. Summary in J. R. Micr. Soc. 1892, p. 471.
- 172. —. Zur Kenntniss der Coniferen-Laüse. T. c. pp. 66-70 & 73-78. [Rhynchota.]
  - Summary in J. R. Micr. Soc. 1892, p. 600.
- 173. Embrional'noe razvītie prussaka (*Phyllodromia germanica*).

  Trudui St. Petersburg Est. xxii, pp. 39-223, pls. i-vi. [*Orthoptera*.]

  Apparently an important paper, but it is entirely in Russian.
- 174. CHRETIEN, P. Les chenilles de l'Artichaut. Le Nat. 1892, p. 132. [Lepidoptera.]
- 175. Christoph, H. Arctia turbans eine neue sibirische Art. Hor. Ent. Ross. xxvi, p. 460. [Lepidoptera.]
- 176. COBELLI, R. Quattro nuove specie di *Imenotteri*. Verh. z.-b. Wien, xlii, pp. 67-72.
- 177. COCKERELL, T. D. A. The Sugar-cane Borer. Bull. Botan. dept. Jamaica, No. 30, 1892. [Lepidoptera.]
- 178. —. Coccidæ, or Scale Insects. Op. cit. No. 36, pp 5-9.
- 179. —. Museum Notes. J. Inst. Jamaica, i, pp. 134-137. One of the descriptions of *Coccidæ* is by Morgan.
- 180. A Cecid bred from Coccidæ. Ent. xxv, pp. 180 & 181. [Diptera.]
- Comstock, J. H. The Descent of the Lepidoptera. An application of the theory of Natural Selection to Taxonomy. P. Am. Ass. xli, pp. 199-201.
- 182. Constant, A. Liste annotée des Lépidoptères envoyés à la Société d'histoire naturelle d'Autun. Bull. Soc. Autun, v, pp. 15-83. Includes notes on food-plants and localities, chiefly of Micros.
- 183. Cook, A. J. Report of Apicultural Experiments in 1891. Bull. Dep. Agric. Ent. xxvi, pp. 83-92. [Hymenoptera.]

- 184. Coquillett, D. W. Report on the Scale Insects of California. T. c. pp. 13-35. [Coccidæ.]
- 185. —... Revision of the Species of Anthrax from America North of Mexico. Tr. Am. Ent. Soc. xix, pp. 168-187. [Diptera.]
- 186. —. Revision of the Bombylid genus Epacmus (Leptochilus). Canad. Ent. xxiv, pp. 9-11. [Diptera.]
- 187. —... Notes and Descriptions of Bombylidæ. T. c. pp. 123-126. [Diptera.]
- A New Genus of Diptera allied to Rhaphiomidas. T. c. p. 314.
- 189. —.. The Dipterous parasite of Melanoplus devastator in California. Ins. Life, v, pp. 22-24.
- 190. —. A New Dalmannia from California. Ent. News, iii, p. 150. [Diptera.]
  - [See also Bruner, Coquillett, & Osborn (136).]
- Cosmovici, L. C. Contributions à l'étude de la faune entomologique Roumaine. Lépidoptères. Le Nat. 1892, pp. 254-256, 264, & 280.
- 192. Coste, —. On Insect Colours. Nature, xlv, pp. 513-517, 541, & 542.
  - Cf. HOPKINS, t. c. p. 581; also Coste & Meldola, t. c. p. 605.
- 193. COUCKE, E. Les Brachymères de Belgique et des régions voisines.

  Ann. Ent. Belg. xxxvi, pp. 69-79. [Colsoptera.]
- 194. CROISSANDEAU, J. Description d'une nouvelle espèce de Thorectes. Bull. Soc. Ent. Fr. lxi, p. coxxi. [Coleoptera.]
- 195. CUNNINGHAM, D. D. On the Phenomena of Fertilization in Ficus roxburghii, Wall. Ann. R. Botan. Gard. Calcutta, i, App. pp. 11-51, pls. i-v. [Hymenoptera.]
- 196. ODALE, C. W. The Lepidoptera of Dorsetshire. A Catalogue of Butterflies and Moths found in the County of Dorset. Second edition. Dorchester: 76 pp.
- Dalla Torre, K. W. V. Hymenopterologische Notizen. Wien. ent. Z. xi, pp. 89-93.

These notices consist chiefly of new names proposed for synonymical reasons. In consideration of the current publication of the author's Catalogue of *Hymenoptera*, it is not necessary to give these new specific names in our pages.

198. DALLA TORRE, C. G. Catalogus Hymenopterorum hucusque descriptorum systematicus et synonymicus. Vol. vi. Chrysidida (Tubulifera). Leipzig: 1892, 8vo, pp. viii & 118.

- 199. DESBROCHERS, J. DES LOGES. Étude sur les Scolytus d'Europe. Frelon, i, pp. 10-17. [Coleoptera.]
- Essai d'une Révision des Hypomeces et genres voisins. Ourculionides du groupe des Tanymécides. T. c. pp. 45-51. [Coleoptera.]
- Etudes sur les Curculionides d'Europe et des pays limitrophes.
   T. c. pp. 52-107. [Coleoptera.]
- 202. —. Diagnoses d'espèces inédites du genre Apion. T. c. pp. 107
   & 108. [Coleoptera.]
- 203. —. Description d'un Brenthide nouveau des Nouvelles-Hébrides. T. c. p. 109. [Coleoptera.]
- 204. —... Monographie des Cassida de France, suivie d'observations sur le classement des espèces de ce genre au Catalogus Coleopterorum Europæ, Caucasi et Armeniæ rossicæ. T. c. pp. 1-48 (sep. pag.). See also op. cit. ii, pp. 82-87.
- 205. ——. Espèces inédites de Curculionides de l'ancien monde. T. c. i, pp. 110-123; ii, pp. 1-12 & 88-100. [Coleoptera.]
- 206. —. Essai d'une révision des espèces du genre Hyphantus, Curculionide de la tribu des Otiorhynchides. T. c. ii, pp. 12-19. [Coleoptera.]
- Révision des Baridiides d'Europe et des Contrées limitrophes.
   T. c. pp. 19-69. [Coleoptera.]
- 208. —. Tableaux dichotomiques des Curculionides Cossonides d'Europe et des pays limitrophes. T. c. pp. 69-82. [Curculionidæ.]
- 209. —. Revision des Balaninides et des Anthonomides d'Europe et des confins de la Méditerranée, en Afrique et en Asie. T. c. pp. 101-(incomplete). [Coleoptera.]
- 210. DISTANT, W. L. A Naturalist in the Transvaal. London: 1892, 8vo, pp. xvi & 277, pls. v.
- Pp. 187-262 are devoted to lists of Insects and descriptions of new species; these are by the author himself and BOURGEOIS, DURRANT, GAHAN, GORHAM, JACOBY, and DE SAUSSURE. There are also many entomological observations in the narrative.
- 211. A monograph of Oriental Cicadida. Parts v-vII, pp. ix & 97-158, pls. x-xv.

These parts complete the work.

- Contribution to a knowledge of the Homopterous family Fulgoridæ. Tr. Ent. Soc. 1892, pp. 275-286, pl. xiii. [Rhynchota.]
- 213. —... On some undescribed *Cicadidæ*, with synonymical notes. Ann. N. H. (6) ix, pp. 313-327. [Rhynchota.]
- On some undescribed Cicadidæ, with synonymical notes.
   Op. cit. x, pp. 54-67. [Rhynchota.]

- 215. [DISTANT, W. L.] Description of a new species of the Homopterous family Cicadida. T. c. p. 406.
- 216. Contributions to a knowledge of the Entomology of the Transvaal. T. c. pp. 407 & 408. [Coleoptera, Lepidoptera.]
- 217. —. Descriptions of four new species of Scutellerina. Ent. xxv, Supp. pp. 95 & 96. [Rhynchota.]
- Notes on Ethiopian Rhynchota. Ent. M. M. xxviii, pp. 187
   188.
  - ---. [See also GODMAN & SALVIN (347).]
- 219. Dodero, A. Res ligusticæ. xvi. Nota sul genere Trogaster, Sharp. Ann. Mus. Genov. (2) x, pp. 665-668. [Coleoptera.]
- Dognin, P. Description de Lépidoptères nouveaux. Le Nat. 1892, pp. 16, &c., &c.
- 221. DOHBN, C. A. Belonuchus nulli-cedo. S. E. Z. liii, p. 74. [Coleoptera.]
- 222. DOHRN, H. Neue und ungenügend bekannte Phaneropteriden aus dem Malayischen Faunengebiete. T. c. pp. 63-73. [Orthoptera.]
- 223. DOBAN, E. W. The evolution of heat by Silvanus cassiæ. P. E. Soc. Wash. ii, pp. 325-327. [Coleoptera.]
- 224. Douglas, J. W. Notes on some British and Exotic Coccidæ. Ent. M. M. xxviii, p. 105, &c.
  - ---. [See also Morgan (625).]
- 225. Druce, H. Descriptions of new species of Eratina from Tropical America, Ann. N. H. (6) ix, pp. 97-107. [Lepidoptera.]
- 226. —. Description of a new genus and some new species of Heterocera from Central America. T. c. pp. 275-279. [Lepidoptera.]
  - ---. [See also GODMAN & SALVIN (347).]
- 227. DRUCE, H. H. A List of the Lycanidae of the South Pacific Islands, east of the Solomon group, with descriptions of several new species. P. Z. S. 1892, pp. 434-446, pl. xxvii. [Lepidoptera.]
- 228. Dufour, J. Note sur le Botrytis tenella et son emploi pour la destruction des Vers blancs. Bull. Soc. Vaud. xxviii, pp. 49-56.
- Dugès, A. Acanthia inodora. Naturaleza (2) ii, p. 169. [Rhynchota.]
  - DURRANT, J. H. [See DISTANT (210).]
- 230. DUVIVIER, A. Melanges entomologiques. x & xI. Diagnoses de Coléoptères nouveaux du Congo. Ann. Ent. Belg. xxxvi, pp. 56-60 & 163-167.

231. [DUVIVIER, A.] Contributions à la faune entomologique de l'Afrique centrale. II. Note sur les Coléoptères des vallées de l'Itimbiri-rubri et de l'Uelle (régions du Haut-Congo) recueillis en 1890 par MM. Joseph Duvivier et Mils. T. c. pp. 257-383.

A catalogue, with descriptions of several species recently described, as well as some new.

- 232. Les *Phytophages* du Chota-Nagpore. 2º Note. *T. c.* pp. 396-449. [Coleoptera.]
- 233. Duzee, E. P. van. New North-American Homoptera.—IV. Canad. Ent. xxiv, pp. 113-116. [Rhynchota.]
- 234. —. New North-American Homoptera.—v. T. c. pp. 266-268. [Rhynchota.]
- 235. A revision of the N. American species of *Phlepsius*. Tr. Am. Ent. Soc. xix, pp. 63-82, pl. i. [Rhynchota.]
- 236. A synoptical arrangement of the genera of the N. American Jassidæ, with some descriptions of new species. T. c. pp. 295-307. [Rhynchota.]
- 237. The North American Jassidæ allied to Thamnotettia, Psyche, vi, pp. 305-310. [Rhynchota.]
- 238. DYAR, H. G. A new Gluphisia from the Pacific coast. Ent. News, iii, p. 168. [Lepidoptera.]
- 239. —. New forms of Ichthyura. Canad. Ent. xxiv, p. 179. [Lepidoptera.]
- Notes on Cerura, with descriptions of new species. Psyche, vi, pp. 290-292. [Lepidoptera.]
- 241. Eaton, A. E. New species of Ephemeridæ from the Tenasserim Valley. Tr. Ent. Soc. 1892, pp. 185-190. [Neuroptera.]
- 242. —. Notes on some native Ephemeridæ in the Indian Museum, Calcutta. J.A.S.B.lx, pp. 406-413. [Neuroptera.]
- 243. ——. On two new and some other Japanese species of Ephemeridæ. Ent. M. M. xxviii, pp. 302 & 303.
  - ---. [See also GODMAN & SALVIN (347).]
- 244. EBERLI, J. Untersuchungen am Verdanungstraktus von Gryllotalpa vulgaris. Viert. Ges. Zürich, xxxvii, pp. 167-212. [Orthoptera.] Summary in J. R. Micr. Soc. 1892, p. 781.
- 245. Eckstein, K. Der Baumweissling, Aporia cratægi, Hb. Zool. Jahrb. (Syst.) vi, pp. 230-240. [Lepidoptera.]

Summary in J. R. Micr. Soc. 1892, p. 470.

EDWARDS, H. [See BEUTENMÜLLER (77, 78).]

246. EDWARDS, W. H. The Butterflies of North America. Part XIII. [Lepidoptera.]

1892. [VOL, XXIX.]

- 247. [EDWARDS, W. H.] Miscellaneous Notes on Butterflies, Larvæ, &c. Canad. Ent. xxiv, pp. 49-56 & 105-111. [Lepidoptera.]

  ELLIOT, S. [the late]. [See BEUTENMÜLLER (78).]
- 248. ELWES, H. J. On Butterflies collected by Mr. W. Doherty in the Naga and Karen Hills and in Perak. P. Z. S. 1892, pp. 617-664, pls. xliii & xliv. [Lepidoptera.]
- 249. EMERY, C. Le formiche dell'ambra siciliana nel museo mineralogico dell' Università di Bologna. Mem. Acc. Bologn. (5) i, pp. 141-165, pls. i-iii. [Hymenoptera.]
- 250. —. Revision critique des Fourmis de la Tunisie. Exploration scientifique de la Tunisie. Hymenoptères, pp. i, ii, & 1-21.
- 251. —... Origines de la faune actuelle des Fourmis de l'Europe. Bull. Soc. Vaud. xxvii, pp. 258-260. [Hymenoptera.]
- 252. —. Sopra alcune Formiche raccolte dall' Ingegnere L. Bricchetti Robecchi nel paese dei Somali. Ann. Mus. Genov. (2) xii, pp. 110-122. [Hymenoptera.]
- 253. ——. Crematogaster striatulus, n. sp. Bull. Soc. Ent. Fr. lxi, p. liii. [Hymenoptera.]
- 254. —. Diagnoses de cinq nouveaux genres de Formicides. T. c. pp. celxxv-celxxvii. [Hymenoptera.]
- 255. EPPELSHEIM, E. Neue Staphylinen der österreichisch-ungarischen Monarchie und der angrenzenden Länder. Wien. ent. Z. xi, pp. 289-297. [Coleoptera.]
- 256. —... Zur Staphylinen-fauna Turkestan's. Deutsche e. Z. 1892, pp. 321-346. [Coleoptera.]
  ERSCHOFF, N. [See ROMANOFF (779).]
- 257. ESCHERICH, C. Die biologische Bedentung der "Genitalanhänge" der Insekten. (Ein Beitrag zur Bastardfrage.) Verh. z.-b. Wien, xlii, pp. 225-240, pl. iv.
- 258. ESCHERICH, K. Ueber die Gesetzmässigkeit im abändern der Zeichnung bei Insecten. Beitrag zur Theorie von der Entwicklung aus constitutionellen Ursachen. Deutsche e. Z. 1892, pp. 113-130, pl. i. [Coleoptera.]
  - Deals with variation in Zonabris.
- Studien über die Cleriden-Gattung Trichodes, Herbst.
   Wien. ent. Z. xi, pp. 226-230. [Coleoptera.]
- 260. EVERTS, E. Tabellarisch overzicht der in Nederland waargenomen Donaciini. Tijdschr. Ent. xxxv, pp. 31-58. [Coleoptera.]
- 261. FAIRMAIRE, L. Voyage de M. E. Simon au Venezuela. Coléoptères Hétéromères. Ann. Soc. Ent. Fr. lxi, pp. 77-98.
- 262. Coléoptères d'Obock. Troisième partie. Rev. d'Ent. xi, pp. 77-126.

- 263. [FAIRMAIRE, L.] Descriptions de Coléoptères des environs d'Akbès (Syrie). Ann. Ent. Belg. xxxvi, pp. 144-159.
- Descriptions de quelques Coléoptères Argentins. T. c. pp. 242-263.
- 265. —... Nouveau genre de Coléoptère Hétéromère. Bull. Soc. Ent, Fr. lxi, p. vii.
- 266. —. Nouveau genre de Cérambycide. T. c. p. xxi. [Coleoptera.]
- 267. —. Descriptions de deux Coléoptères nouveaux. T. c. p. cix.
- 268, —. Descriptions de quelques Coléoptères de Turkestan. T. c. pp. cxxi-cxxiv.
- Descriptions de quelques Coléoptères de Madagascar. T. c. p. clxviii.
- 270. —. Trois Coléoptères nouveaux. T. c. p. cli.
- 271. Descriptions de quelques Coléoptères nouveaux de Turkestan. T. c. p. clii.
- 272. \*FAUCONNET, L. Faune analytique des Coléoptères de France. Autun, 8vo, 528 pp. Cf. Rev. d'Ent. xi, p. 59.
- 273. FAUST, J. Die Anchoniden-Gruppe. Deutsche e. Z. 1892, pp. 17-60.
  [Coleoptera.]
- 274. —. Einige neue Curculioniden. T. c. pp. 60-64. [Coleoptera.]
- 275. —. Reise von E. Simon in Venezuela. Curculionidæ. S. E. Z. liii, pp. 1-43. Coleoptera.
- 276. —. Notizen über Rüsselkäfer. T. c. pp. 44-52. [Coleoptera.]
- 277. —. Fünf neue Curculioniden von Australien. T. c. pp. 179-184. [Coleoptera.].
- 278. —. Curculioniden aus dem Malayischen Archipel. T. c. pp. 184-227. [Coleoptera.]
  - —. [See also Heyden (402).]
- 279. FENK, C. Mittheilungen über das Auftreten des Nonnenspinners (Liparis monacha) in den Jahren 1890 und 1891 in Süddeutschland und der Ostschweiz. Ber. St. Gall. Ges. 1890-91, pp. 289-306. [Lepidoptera.]
- 280. FERNALD, C. H. New North American Micro-Lepidoptera. Canad. Ent. xxiv, pp. 121, 122, & 178.
- 281. FERRARI, P. M. Res ligusticæ. XIX. Elenco dei Rincoti ligustici (Emitteri e Cicadarii) fin ora osservati. Ann. Mus. Genov. (2) xii, pp. 548-576.
- 282. Finot, A. Orthoptère nouveau du midi de la France. Bull. Soc. Ent. Fr. lxi, p. xxxv.

- 283. FITZ-GIBBON, M. Use of the Hairs of Acronycta alni. Ent. xxv, pp. 39 & 40. [Lepidoptera.]
- 284. FLEISCHER, A. Anthaxia semilimbata, n. sp. Wien. ent. Z. xi, p. 209. [Coleoptera.]
- 285. FLEUTIAUX, E. Note sur les Physodactylini. Ann. Soc. Ent. Fr. lxi, pp. 403-412, pl. viii. [Coleoptera.]
- 286. Forbes, S. A. Bacteria normal to Digestive Organs of Hemiptera. Bull. Illinois Lab. iv, pp. 1-6.
- 287. FOREL, A. Histoire physique, naturelle et politique de Madagascar, publiée par Alfred Grandidier. Volume xx, Histoire Naturelle des Hymenoptères. Deuxième partie. Les Formicides. Paris, 1891, pp. v & 243, pls. i-v.
- Die Nester der Ameisen. Neujahrsbl. Ges. Zurich, xcv, pp. 1-36, pl. [Hymenoptera.]
- 289. Die Ameisen fauna Bulgariens (nebst biologischen Beobachtungen). Verh. z.-b. Wien, xlii, pp. 305-318, pl. v. [Hymenoptera.]
- Motes myrmécologiques. Ann. Ent. Belg. xxxvi, pp. 38-43.
   [Hymenoptera.]
- Quelques Fourmis de la faune méditerranéenne. T. c. pp. 452-457. [Hymenoptera.]
- 292. —. Liste der aus Somaliland von Hrn. Prof. Dr. Conr. Keller aus der Expedition des Prinzen Ruspoli im August und September 1891, zurückgebrachten Ameisen. Mt. Schw. ent. Ges. viii, pp. 349-354. [Hymenoptera.]
- 293. Les Formicides de l'empire des Indes et de Ceylan. Part 1. J. Bomb. Soc. vii, pp. 219-245, pl. A. [Hymenoptera.]
- 294. —. Die Ameisen Neu-Seelands. Mt. Schw. ent. Ges. viii, pp. 331-343. [Hymenoptera.]
- 295. —... Attini und Cryptocerini: Zwei neue Apterostigma-Arten.
  T. c. pp. 344-349. [Hymenoptera.]
- 296. —. Le mâle des Cardiocondyla et la reproduction consanguine perpétuée. Ann. Ent. Belg. xxxvi, pp. 458-461. [Hymenoptera.]
- 297. —. Hermaphrodite de l'Azteca instabilis. Bull. Soc. Vaud. xxviii, pp. 268-270, pl. xvi. [Hymenoptera.]
  —. [See also Keller (475).]
- Fox, W. J. Report on the Hymenoptera collected in West Greenland. P. Ac. Philad. 1892, pp. 133-135.
- 299. —. The North American species of Ceropales. Tr. Am. Ent. Soc. xix, pp. 49-63. [Hymenoptera.]

Ì

- 300. [Fox, W. J.] Monograph of the North American species of Tachytes. T. c. pp. 234-252, pl. xi. [Hymenoptera.]
- 301. —. The North American Pemphredonidæ. T. c. pp. 307-326. [Hymenoptera.]
- 302. —. New North American species of Gorytes. Canad. Ent. xxiv, pp. 149-153. [Hymenoptera.]
- 303. —... Synopsis of the North American species of Astatus, Latr. T. c. pp. 232-235. [Hymenoptera.]
- 304. Hymenopterological notes. Ent. News, iii, pp. 9, 10, & 29.
- 305. Description of Four new species of Fossorial Hymenoptera from California. T. c. pp. 170-172.
- Description of a new solitary Wasp. T. c. p. 197.
   [Hymenoptera.]
- 307. FREY-GESSNER, E. Les Bourdons du Valais. Bull. Soc. Murith. xiii, pp. 32-36. [Hymenoptera.]
- 308. Tables analytiques pour la détermination des Hymenoptères du Valais. T. c. pp. 37-48, pls. i, ii; and op. cit. xvi, pp. 43-112, pl. iii.
- 309. FRIVALDSZKY, J. Coleoptera in expeditione D. Comitis Belæ Széchenyi in China, præcipue boreali, a Dominis Gustavo Kreitner et Ludovico Lóczy anno 1879 collecta. Pars secunda. Term. füzetek, xv, pp. 114-125.
- 310. —. Coleoptera duo nova ex Hungaria. T. c. pp. 132 & 133.
- 311. FROGGATT, W. W. Notes on Australian Cynipidæ, with descriptions of several new species. P. Linn. Soc. N.S.W. (2) vii, pp. 152-156. [Hymenoptera.]
- Catalogue of the described Hymenoptera of Australia: Part II.
   T. c. pp. 205-248.
- 313. Fuchs, A. Macrolepidopteren der Loreley-Gegend. Dritte Besprechung. JB. nass. Ver. xlv, pp. 83-106.
- 314. FYLES, T. W. Further notes on Gelechia gallædiplopappi, and description of a new species of Bracon. Canad. Ent. xxiv, pp. 34 & 35. [Hymenoptera.]
- 315. Gahan, C. J. Additions to the Longicornia of Mexico and Central America, with notes on some previously recorded species. Tr. E. Soc. 1892, pp. 255-274, pl. xii. [Coleoptera.]
- Notes on Longicorn Coleoptera of the group Cerambycina, with descriptions of new genera and species. Ann. N. H. (6) ix, pp. 19-32.
  - —. [See also Distant (210) and Gorham & Gahan (353).]

- 317. GANGLBAUER, L. Ein neuer Anophthalmus aus der Herzegowina. Wien. ent. Z. xi, p. 233. [Coleoptera.]
- 318. —. Ein neues Bembidium aus den Alpes maritimes. T. c. p. 314. [Coleoptera.]
- 319. —. Zwei neue Boreaphilinen. Soc. Ent. vii, p. 17. [Coleoptera.]
- 320. GASPERINI, R. Tenthredinidum species nova. Prilog Fauni dalmatinskih Pauka. [Hymenoptera.] (f. Wien. ent. Z. xi, p. 232.
- 321. GERMAIN, P. Trois mois dans la vallée du Sipotuba (Province de Matto-Grosso, Brésil). Act. Soc. Chili, i, pp. 53-65.
- 322. —. Notes sur les Coléoptères du Chili. Renseignements et observations; descriptions d'espèces nouvelles; rectifications; indications de synonymie. Act. Soc. Chili, ii, pp. 241-261.
  - Gerhardt, J. [See Letzner (551).]
- 323. Gerstaecker, A. Bestimmung der von Herrn Dr. F. Stuhlmann in Ost-Afrika gesammelten *Hemiptera*. JB. Hamb. Anst. ix (2) pp. 45-58.
- 324. Gestro, R. Viaggio di Leonardo Fea in Birmania e regioni vicine. xxxIII. Sul genere Arctolamia. Ann. Mus. Genov. (2) x, pp. 221-224. [Coleoptera.]
- 325. —... Viaggio di Leonardo Fea in Birmania e regioni vicine. xxix. Enumerazione delle *Hispidæ*. T. c. pp. 225-268. [Coleoptera.]
- 326. —. Viaggio di Leonardo Fea in Birmania e regioni vicine. XXXIV. Materiali per lo studio del genere Ichthyurus. T. c. pp. 555-594. [Coleoptera.]
- 327. —. Viaggio di Leonardo Fea in Birmania e regioni vicine. xxxvII. Enumerazione delle Cetonie. T. c. pp. 835-876, pl. ii. [Coleoptera.]
- 328. —. Viaggio di Leonardo Fea in Birmania e regioni vicine. xL. Contribuzione allo studio degli insetti termitofili. T. c. pp. 903-907. [Coleoptera.]
- 829. Appendice ai materiali per lo studio del genere Ichthyurus. T. c. pp. 1022-1048. [Colcoptera.]
- 330. Appunti sul genere Omophron. T. c. pp. 961-964. [Coleoptera.]
- 331. —. Viaggio di Lamberto Loria nella Papuasia orientale. VI. Hispidæ. T. c. pp. 1015-1021. [Colcoptera.]
- 332. —... Nuovi materiali per lo studio degli Anophthalmus italiani. Op. cit. xii, pp. 79-85. [Coleoptera.]
- 333. —. Viaggio di Leonardo Fea in Birmania e regioni vicine. XLVI. Cenno sui Paussidi. T. c. pp. 705-709. [Coleoptera.]

ì

- 334. [GESTRO, R.] Di alcuni *Coleotteri* raccolti nel paese dei Somali dall' Ing. L. Brichetti Robecchi. T. c. pp. 747-790.
- 335. —... Sopra due *Hispidæ* di Engano raccolte dal Dott. Elio Modigliani. *T. c.* pp. 791-794. [Coleoptera.]
- 336. GIARD, A. Sur un Diptère Stratyomide (Beris vallata) imitant un Tenthrède (Athalia annulata Fab.) C.R. Soc. Biol. (n.s.) iv, pp. 43-45.
- 337. Sur un Hémiptère Hétéroptère (Halticus minutus, Reuter) qui ravage les Arachides en Cochin-Chine. T. c. pp. 79-82.
- 338. Sur une Laboulbéniacée, parasite de Mormolyce phyllodes. T. c. pp. 156-158.
- 339. Giglio-Tos, E. Parassitismo di una larva d'Aricia in un Curabo. Ann. Ac. Agric. Torino, xxxiv (pp. 1-13, sep. copy). [Diptera.]
- 340. —. Sul parassitismo di una larva d'Aricia in un Carabo. Boll. Mus. Tor. vii, No. 116. [Diptera.]
   Supplementary Bibliographic List. .
- 341. —. Un nuovo genere di Sirfidi. T. c. No. 117. [Diptera.]
- 342. —. Diagnosi di nuove specie di *Ditteri*. VI & VII, *Sirfidi* del Messico. T. c. Nos. 123 & 132.
- 343. —... Sui due generi di Sirfidi, Rhopalosyrphus ed Omegasyrphus. T. c. No. 118. [Diptera.]
- 344. GILLETTE, C. P. Colorado Cynipidæ. Ent. News, iii, pp. 246-248. [Hymenoptera.]
- 345. Goding, F. W. A synopsis of the subfamilies and genera of the Membracidæ of North America. Tr. Am. Ent. Soc. xix, pp. 253-260. [Rhynchota.]
- 346. —. Studies in North American Membracidæ. Ent. News, iii, pp. 108, 200, & 201. [Rhynchota.]
- 347. GODMAN, F. D., & SALVIN, O. Biologia Centrali-Americana. Parts 99-105.
  - These parts include the following entomological matter :-
    - Coleoptera, ii (1): Trogositidæ, pl. iii.
    - Coleoptera, iv (1): Tenebrionidæ: supp. pp. 477-524, by G. C. CHAMPION.
    - Coleoptera, iv (2): Cantharidæ, &c., pp. 369-448, pls. xvii-xx, by G. C. Champion.
    - Coleoptera, vi (1): supp. pp. 313-374, pls. xlii & xliii, completing the part. Phytophaga, by M. Jacoby.
    - Coleoptera, vii: Coccinellida, pp. 161-176, pl. ix.
    - Lepidoptera-Heterocera, ii, pp. 25-128, pls. xliv-lii. Geometrides, by H. DRUCE.
    - Diptera, iii, pp. 57-89, pls. i & ii. Syrphida, by S. W. WILLISTON.
    - Rhynchota-Heteroptera, pp. 353-368, by W. L. DISTANT.
    - Neuroptera, pp. 1-16, pl. i. Ephemeridæ, by A. E. EATON.

- 348. Gorham, H. S. On three new species of Coccinellidae from Scioa.

  Ann. Mus. Genov. (2) x, pp. 910 & 911. [Coleoptera.]
- 350. Descriptions of Coleoptera collected by Mr. John Whitehead on Kina Balu, Borneo. P. Z. S. 1892, pp. 83-90.
- Coleoptera from Central China and the Korea. Ent. xxv, Supp. pp. 81-85.
- 352. Description of a new species of Elater. T. c. pp. 304 & 305. [Coleoptera.]
  - ---. [See also DISTANT (210) and GODMAN & SALVIN (347).]
- 353. GORHAM, H. S., & GAHAN, C. T. On the Coleoptera collected by Mr. W. Bonny in the Aruwimi Valley, Central Africa. P. Z. S. 1892, pp. 90-96.
- 354. Gould, L. J. Experiments in 1890, 1891, on the colour-relation between certain *Lepidopterous* larvæ and their surroundings, together with some other observations on *Lepidopterous* larvæ. Tr. E. Soc. 1892, pp. 215-245, pl. xi.
- 355. GRABER, V. Beiträge zur vergleichenden Embryologie der Insecten. Denk. Ak. Wien, lviii, pp. 803-866, pls. i-vii.
- 356. Graeser, L. Beiträge zur Kenntniss der Lepidopteren-Fauna des Amurlandes. B. E. Z. xxxvii, pp. 209-234.
- 357. —. Neue Lepidopteren aus Central-Asien. T. c. pp. 299-318.
- 358. GRÄFFE, E. Le api dei dintorni di Trieste. Atti Mus. civico Trieste, viii, pp. 123-142. [Hymenoptera.]
  Includes description of a new Osmia by SCHMIEDEKNECHT.
- 359. Grassi, B. Conclusioni d'una Memoria sulla società dei Termiti.

  Atti Acc. Lincei, 1892, pp. 33-36. [Neuroptera.]
- 360. GRIFFINI, A. Sulla variabilità delle apofisi metasternali nel Dytiscus marginalis, L. Boll. Mus. Zool. Tor. vii, No. 115. [Coleoptera.]
- 361. —. Sui generi Anonconotus, Cam., e Analota, Brunner. Op. cit. No. 125. [Orthoptera.]
- 362. GRIFFITHS, A. B. Recherches sur les couleurs de quelques insectes. C.R. cxv, pp. 958 & 959.
  - Summary in J. R. Micr. Soc. 1892, p. 778.
- 363. —. On the Blood of the *Invertebrata*. P. R. Soc. Edinb. xix. The Blood of Insects, pp. 123-126.
- 364. GRILAT, R. Description d'une nouvelle espèce de Ceutorrhynchus. Échange, No. 32, 1887. [Coleoptera.]

- 365. [Grilat, R.] Description d'une nouvelle espèce de Bythinus de France. Op. cit. No. 58, 1889, p. 78. [Coleoptera.]
- 366. GROUVELLE, A. Voyage de M. E. Simon au Venezuela. Coléoptères. Colydiides. Ann. Soc. Ent. Fr. lxi, pp. 99-102.
- Voyage de M. E. Simon à l'île de Luzon, Philippines. Cucujides. T. c. pp. 285-288. [Coleoptera.]
- Voyage de M. Ch. Alluaud dans le territoire d'Assinie. Nitidulidæ, Colydiidæ, Rhysodidæ. T. c. pp. 292-300. [Coleoptera.]
- 369. —. Viaggio di Leonardo Fea in Birmania e regioni vicine. L. Nitidulides, Cucujides, et Parnides. 2me partie. Ann. Mus. Genov. (2) xii, pp. 833-868. [Coleoptera.]
- 370. —. Clavicornes de Bengale. 2me Liste. Ann. Ent. Belg. xxxvi, pp. 60 & 61. [Coleoptera.]
- 371. —. Deux espèces nouvelles de Coléoptères de la famille des Nitidulidæ. Notes Leyd. Mus. xiv, pp. 43 & 44.
- 372. Trois espèces nouvelles d'Helmides des îles de la Sonde. T. c. pp. 187-189. [Coleoptera.]
- 373. GUILLEBEAU, F. Revision des Phalacrides de la Faune paléarctique. Rev. d'Ent. xi, pp. 141-197. [Coleoptera.]
- 374. Description d'une nouvelle espèce de Limnobius. T. c. p. 324. [Coleoptera.]
- 375. Descriptions de quelques nouvelles espèces de Coléoptères. Echange, i, No. 6, &c.
- 376. —. Description de trois espèces nouvelles de Coléoptères. Pp. 1-3.
  - Published with Echange, No. 19 (1886), with separate pagination.
- 377. —. Note sur quelques Catops nouveaux. Op. cit. 1891, p. 116. [Coleoptera.]
- 378. Descriptions de deux Coléoptères nouveaux. Bull. Soc. Ent. Fr. lxi, pp. cxxxiii & cxxxiv.
- 379. —. Description de deux espèces nouvelles de Coléoptères. T. c. pp. clxxxiii & clxxxiv.
- 380. GUMPPENBERG, —. Systema Geometrarum zonæ temperatioris septentrionalis. Fünfter Theil. N. Acta Ac. L.-C. Nat. cur. lviii, pp. 223-359. [Lepidoptera.] Reviewed by REBEL (714).
- 381. GUTHRIE, L. G. On a Toad Killed by the Larvæ of Blow-flies. Ent. M. M. xxviii, pp. 9-12.
- 382. HAASE, E. Untersuchungen über die Mimicry auf Grundlagen eines naturlichen Systems der Papilioniden. Bibl. Zool., Heft viii, Lief. 4-10, pp. 9-98, pls. v-xiv.

- 383. Hagen, B. Beitrag zur Kenntniss der Rhopuloceren der Insel Banka. Berl. Ent. Zeit. xxxvii, pp. 139-158. [Lepidoptera.]
- 381. Hampson, G. F. The Fauna of British India, including Ceylon and Burma. Published under the authority of the Secretary of State for India in Council. Edited by W. T. Blanford. Moths, Vol. 1. London: 1892, 8vo, xxiii & 527 pp.
- 385. On Stridulation in certain Lepidoptera, and on the Distortion of the Hind Wings in the Males of certain Ommatophorina. P. Z. S. 1892, pp. 188 & 189.
- 386. HANDLIRSCH, A. Monographie der mit Nysson und Bembee verwandten Grabwespen, vi. SB. Ak. Wien, ei, pp. 25-205, pls. i-iii. A monograph of the genus Stizus.
- 387. Ein neuer Sphecius aus Madagaskar, Jahresheft, Ver. Trenesin. xiii, p. 27. [Hymenoptera.]
- 388. HARRINGTON, W. H. Canadian Hymenoptera. No. 1. Canad. Ent. xxiv, pp. 98 & 99.
- 389. A New Ischalia from Vancouver Island. T. c. p. 132. [Coleoptera.]
- 390. HARVEY, F. L. An American Species of Templetonia. Ent. News, iii, p. 57. [Thysanura.]
- 391. —. A New Smynthurus. T. c. p. 169. [Thysanura.]
- 392. HEIDEMANN, O. Note on the Food-plants of some Capsidae from the vicinity of Washington. P. E. Soc. Wash. ii, pp. 224-227. [Rhynchota.]
- 393. Heim, F. Sur le centre réflexe du saut chez la Sauterelle. Bull. Soc. Ent. Fr. lxi, p. xcviii. [Orthoptera.]
- 394. HELLER, K. M. Eine neue Calandriden-art der Gattung Paratasis, Chev., aus Java. Notes Leyd. Mus. xiv, pp. 269-272. [Coleoptera.]
- 395. —. Ein beachtenswerther Fall von Mimicry. Ent. Nachr. xviii, pp. 183-185.
  - A case of supposed "aggressive mimicry."
- 396. Henking, H. Untersuchungen über die ersten Entwicklungsvorgauge in den Eiern der Insekten. 111. Specielles und Allgemeines. Z. wiss. Zool. liv, pp. 1-274, pls. i-xii.
- Methoden bei entwicklungsgeschichtlichen Untersuchungen an Insecteneiern. Z. wiss. Mikr. viii, pp. 156-167.
- 398. Henneguy, L. F. Contribution à Pembryogénie des Chalcidiens. C.R. exiv, pp. 133-136. [Hymenoptera.]
- Translated in Ann. N. H. (6) x, pp. 271 & 272. Summary, J. R. Micr. Soc. 1892, p. 357.

- 399. Henneguy, F., & Binet, A. Contribution à l'étude microscopique du système nerveux larvaire de Stratiomys longicornis, Sc. Ann. Soc. Ent. Fr. lxi, pp. 309-316, pl. vi. [Diptera.] Summary in J. R. Micr. Soc. 1892, p. 356.
- 400. HENSHAW, S. Report upon the Gypsy-moth in Massachusetts. Bull. Dep. Agric. Ent. xxvi, pp. 75-82. [Lepidoptera.]
- Hering, E. Platyptilia ochrodactyla, Hb., dichrodactyla, bertrami, und borgmanni. S. E. Z. liii, pp. 269-279.
- 402. HEYDEN, L. VON. Beitrag zur Coleopteren-Fauna von Turkestan. Unter Mitwirkung der Herren J. FAUST, A. KUWERT, und J. WEISE. XIV, Deutsche e. Z. 1892, pp. 99-104; xv, t. c. pp. 105-110.
- 403. HEYLAERTS, F. J. M. Heterocera exotica, nouvelles espèces des Indes orientales neerlandaises (Suite). Ann. Soc. Ent. Belg. xxxvi, pp. 7-15 & 43-48. [Lepidoptera.]
- 404. HEYMONS, —. Entwicklung der Cölomsäcke von Phyllodromia germanica. Verh. deutsch. Zool. Ges. 1892, pp. 139-141.
- 405. HOFFBAUER, C. Beiträge zur Kenntniss der Insektenflügel. Z. wiss. Zool. liv, pp. 579-630, pls. xxvi & xxvii.

Describes a variety of glandular structures in the interior of elytra of Coleoptera.

- 406. Holland, W. J. New species of Neptis from Africa. Ent. News, iii, pp. 248-250, pl. ix. [Lepidoptera.].
- 407. —. Descriptions of some new species of African Lepidoptera. Ann. N. H. (6) x, pp. 284-294.
- 408. —... Descriptions of some new species of African Lepidoptera. Ent. xxv, Supp. pp. 89-95.
- 409. —. Notes upon the transformations of some African Lepidoptera. Psyche vi, pp. 213-216, pl. v.
- 410. —. The life-history of Spalgis s-signata, Holl. T. c. pp. 201-203, pl. iv. [Lepidoptera.]
- 411. HOLMGREN, E. Histologiska studier öfver några Lepidopter-larvers digestions kanal och en del af deras körtelartade Bildningar. Ent. Tidskr. xiii, pp. 129-170, pls. i-vi.
- 412. HONRATH, E. G. Neue Rhopalocera, x. B. E. Z. xxxvi, pp. 429-440. [Lepidoptera.]
- 413. —. Eine neue Notodonta. T. c. p. 445. [Lepidoptera.]
- 414. HORMUZAKI, C. v. Lepidopterologische Beobachtungen in der Bucovina. Ent. Nachr. xviii, pp. 305-321.
- 415. —. Zur systematischen Begrenzung der Gattungen Satyrus, Latr. und Epinephele, Hübn. T. c. pp. 354-358. [Lepidoptera.]

- 416. HORN, G. H. Studies in Chrysomelidæ. Tr. Am. Ent. Soc. xix, pp. 1-17. [Coleoptera.]
- 417. —. Notes on Amara, subg. Triæna. T. c. pp. 18 & 19. [Coleoptera.]
- 418. —. Random studies in N. American Coleoptera. T. c. pp. 40-48.
- 419. —. The Eumolpini of Boreal America. T. c. pp. 195-234. [Coleoptera.]
- Wariations of Color-markings in Coleoptera. Ent. News, iii, pp. 25-28, pl. i.
- 421. Horn, W. Fünf dekaden neuer Cicindeleten. Doutsche e. Z. 1892, pp. 65-91. [Coleoptera.]
- 422. Die Cicindeliden des Wiener Hof-Museums. T. c. pp. 92-98. [Coleoptera.]
- 423. Drei neue afrikanische Cicindeliden. T. c. pp. 220 & 221. [Coleoptera.]
- 424. —. Nachträge zur Monographie der Cicindeliden-Gattung Collyris (Chaudoir, 1864). T. c. pp. 353-368. [Coleoptera.]
- 425. Eine neue Cicindela aus Sao Paulo. T.c. p. 368. [Coleoptera.]
- 426. Einige neue Cicindeliden Arten und Varietaten. T. c. pp. 369-372. [Coleoptera.]
  - Horner, A. C. [See Blatch & Horner (103).]
- 427. HORVÁTH, G. Sur l'existence des séries parallèles dans le cycle biologique des *Pemphigiens*. C.R. cxiv, pp. 842-844. [Rhynchota.] Summary in J. R. Micr. Soc. 1892, p. 357.
- 428. . Hemiptera nonnulla nova Asiatica. Term. füzetek, xv, pp. 134-137.
- 429. —. Uebersicht der Hemipteren-Gattung Campylostira, Fieb. Wien. ent. Z. xi, pp. 309-311. [Rhynchota.]
- 430. —. Nouvelle révision du genre Scolopostethus. Rev. d'Ent. xi, pp. 253-256. [Rhynchota.]
- Chasses hivernales dans le midi de la France. T. c. pp. 128–140. [Rhynchota.]
- 432. HOWARD, L. O. The Biology of the Hymenopterous Insects of the Family Chalcidide. P. U. S. Nat. Mus. xiv, pp. 567-588.
- 433. —... Insects of the Subfamily Encyrtinæ with branched antennæ. Op. cit. xv, pp. 361-369, pls. xlvi & xlvii. [Hymenoptera.]
- 434. —. The Habits of Melittobia. P. E. Soc. Wash. ii, pp. 244-248. [Hymenoptera.]
- The Hymenopterous Parasites of Spiders. T. c. pp. 290-802, pl. ii.
- 436. —. A Note on the Parasites of the Coccidæ. T. c. pp. 351-353.

- 437. Hubbard, H. G. Notes on the Larva of Amphizoa. Ins. Life, v, pp. 19-22. [Coleoptera.]
- 438. —. Description of the Larva of Amphizoa lecontei. P. E. Soc. Wash. ii, pp. 341-346, pl. iii. [Coleoptera.]
- 439. —. The Life-history of Xenos. Canad. Ent. xxiv, pp. 257-261. [Coleoptera.]
- 440. Hudson, G. V. An Elementary Manual of New Zealand Entomology. London: 1892, 8vo, 128 pp., xx pls.
- 441. HULST, G. D. New Species of Pyralidæ. Canad. Ent. xxiv, pp. 59-63. [Lepidoptera.]
- 442. Huth, E. Die Futterpflanzen des Seidenspinners. Helios, x, pp. 51-53. [Lepidoptera.]
- 443. Jacobsohn, G. Beitrag zur Systematik des Geotrypini. Hor. Ent. Ross. xxvi, pp. 245-257. [Colsoptera.]
- 444. —... Analytische Übersicht der bekannten Donacia- und Plateumaris-Arten der alten Welt. T. c. pp. 412-437. [Coleoptera.]
- 445. —... Conspectus specierum generis *Haltica*, Geoffr., ex Asia media hucusque notarum. *T. c.* pp. 462–467. [Coleoptera.]
- 446. Jacoby, M. Viaggio di Leonardo Fea in Birmania e regioni vicine. Li. Description of the New Genera and Species of the Phytophagous Coleoptera. Ann. Mus. Genov. (2) xii, pp. 869-999.
- Descriptions of some New Genera and Species of Phytophagous Coleoptera from Madagascar. P. Z. S. 1892, pp. 564-579, pl. xxxix.
- 448. ——. Descriptions of some New Species of *Phytophagous Coleoptera*, and synonymic notes. Ent. xxv, Supp., pp. 86-88.
  - ---. [See also DISTANT (210) and GODMAN & SALVIN (347).]
- 449. Jacquet, E. Recherches et remarques sur l'organisation extérieure et la classification des *Coléoptères*, pp. 1-31 & table.

Includes an attempt to define the "series": issued with Échange, Nos. 37-42.

450. ——. Tableaux analytiques pour la détermination des Rhynchophores de la Faune française, pp. 1-34.

This fragment—its condition as such being due to the decease of the author—was issued with Nos. 42-45 of Échange.

- 451. JAKOWLEFF, B. E. Generis Lethrus, Scop., species novæ vel parum cognitæ. Hor. Ent. Ross. xxvi, pp. 182-199. [Coleoptera.]
- 452. —. Die Peritrechus-Arten der russichen Fauna. T. c. pp. 228-231. [Rhynchota.]
- 453. JAKOWLEW, A. Diagnoses Tenthredinidarum novarum ex Rossia europæa, Sibiria, Asia media et confinium. T. c. pp. 1-62. [Hymenoptera.]

- 454. Janson, O. E. On Taniodera quadrilineata and some allied species. Notes Leyd. Mus. xiv, pp. 55-60. [Coleoptera.]
- 455. JAWOROWSKI, A. Prof. Dr. Veit Graber. Ein Nachruf. Wien. ent. Z. xi, pp. 253-262.

Includes list of his writings, and a portrait.

- 456. Jenkinson, W. Secreting Glands in the Feet of Flies. Sci. Goss. 1892, pp. 85-87. [Diptera.]
- 457. JOHANSEN, H. Ueber die Entwicklung des Imago-auges von Vanessa. Zool. Anz. xv, pp. 353-355.

Summary in J. R. Micr. Soc. 1892, p. 779.

458. Joseph, G. L'influence de l'éclairage sur la disjonction des organes visuels, leur reduction, leur atrophie complète et leur compensation chez les animaux cavernicoles. Bull. Soc. Zool. Fr. 1892, pp. 121-125.

Several cases in Insecta are discussed.

- 459. KANNEGIETER, J. Z. Eenige opmerkingen over het Verzamelen en conserveeren van Insecten in de Tropen. Tijdschr. Ent. xxxv, pp. 59-72.
- 460. Karsch, F. Insecten von Baliburg (Deutsch-Westafrika) gesammelt von Herrn Dr. Eugen Zintgraff. Ent. Nachr. xviii, pp. 161-183.
- 461. —. Vorläufige Beschreibung von drei neuen Lepidopteren von Bismarckburg im Togolande (Deutsch-Westafrika). T. c. pp. 113-117.
- 462. —. Abantis bismarcki eine neue Hesperiide von Bismarckburg im Togolando (Deutsch-Westafrika). T. c. p. 228. [Lepidoptera.]
- 463. —. Caprona adelica eine neue Hesperiide von der deutschen Forschungs-station Bismarckburg im Togolande (Westafrika). T. c. pp. 241-244. [Lepidoptera.]
- 464. —. Vorläufige Beschreibung einer neuen Saturniide aus dem Kamerungebirge. T. c. pp. 353 & 354. [Lepidoptera.]
- 465. —. Kurze Charakteristik neuer Wanzen aus Kamerun gesammelt durch Herrn Dr. Paul Preuss. T. c. pp. 129-136. [Rhynchota.]
- 466. —. Ueber eine Collection durch Herrn Hans Fruhstorfer auf Java gefangener Æschniden. T. c. pp. 249-255. [Neuroptera.]
- 467. —. Zwei neue chinesische Libellen aus der Familie der Kalopterygiden. Berl. Ent. Zeit. xxxvi, pp. 455 & 456. [Neuroptera.]
- 468. —. Eine neue von Herrn Dr. Paul Preuss bei Buea im Kamerungebirge entdeckte Odonate der Gattung 'Macromia. Op. cit. xxxvii, p. 15. [Neuroptera.]
- 469. —. Fünf neue afrikanische Mantodeen. Ent. Nachr. xviii, pp. 5-13. [Orthoptera.]

- 470. [KARSCH, F.] Kurze Charakteristik neuer Mantodeen aus Kamerun gesammelt von Herrn Dr. Paul Preuss. T. c. pp. 145–150. [Orthoptera.]
- Webersicht der von Herrn Dr. Paul Preuss auf der Barombi-Station in Kamerun gesammelten Locustodeen. Berl. Ent. Zeit. xxxvi, pp. 317-346. [Orthoptera.]
- 472. —... Verzeichniss der von Herrn Dr. Paul Preuss im Kamerungebirge erbeutete Orthopteren. Op. cit. xxxvii, pp. 65-78.
- 473. KATURIC, M. Osservazioni biologiche sulle Formiche. Glasnik Naravosl. Druzt. ii, pp. 105-110. [Hymenoptera.]
- 474. —. Ulteriori osservazioni biologiche sulle Formiche. Op. cit. vi, pp. 14-28. [Hymenoptera.]
- 475. Keller, C. Neue Beobachtungen über Symbiose zwischen Ameisen und Akazien. Zool. Auz. xv, pp. 137-143. [Hymenoptera.] Includes descriptions by Forel. A note by Emery, t. c. p. 237, refers to this. Summary in J. R. Micr. Soc. 1892, p. 359.
- 476. Kellicott, D. S. Notes on the Ægeriidæ of Central Ohio. Canad. Ent. xxiv, pp. 42-47, and 11, t. c. pp. 209-212. [Lepidoptera.]
- 477. —. Notes on Ægeriidæ of Central Ohio. Ins. Life, v, pp. 81-85. [Lepidoptera.]
- 478. Kellogg, V. L. Some notes on the Mallophaga. Tr. Kansas Ac. xii, pp. 46-48. [Neuroptera.]
- Motes on Melitera dentata, Grote. Kansas quart. i, pp. 39-41, pl. viii. [Lepidoptera.]
- 480. KERNER, A. V. MARILAUN. Pflanzenleben. Zweiter Band. Geschichte der Pflanzen. Leipzig, 1891, 8vo, 842 pp.
- Mik, in Wien. ent. Z. xi, pp. 28-30, summarizes the views of this writer on galls.
- 481. KERREMANS, C. Catalogue synonymique des Buprestides decrits de 1758 à 1890. Mem. Soc. Ent. Belg. i, pp. 1-304. [Coleoptera.]
- 482. —. Voyage de M. Ch. Alluaud dans le territoire d'Assinie. Buprestidæ. Ann. Soc. Ent. Fr. lxi, pp. 301-304. [Coleoptera.]
- 483. —. Voyage de M. Simon aux îles Philippines. 3e Mémoire. Buprestidæ. T. c. pp. 21-26. [Coleoptera.]
- 484. —. Viaggio di Leonardo Fea iu Birmania e regioni vicine. XLIX. Buprestides. Ann. Mus. Genov. (2) xii, pp. 809-832. [Coleoptera.]
- 485. —... Nouvelle espèce de Buprestide recueillie dans l'île d'Engano, par Mr. le Dr. E. Modigliani. T. c. p. 1000. [Coleoptera.]
- 486. —. Viaggio di Lamberto Loria nella Papuasia orientale. VII. Buprestides. T. c. p. 1001-1006. [Coleoptera.]

- 487. [Kerremans, C.] Note sur les Chrysodémides africaines attribuées au genre Iridotænia. Ann. Ent. Belg. xxxvi, pp. 49-55. [Coleoptera.]
- 488. Diagnoses de deux Buprestides de la région du lac Tanganyka. T. c. p. 56. [Coleoptera.]
- 489. —. Buprestides de l'Inde. T. c. pp. 171-226. [Coleoptera.]
- 490. KERRY, F. Migration of Pieris brassicæ observed at Harwich. Essex Nat. vii, p. 205. [Lepidoptera.]
- 491. °KESSLER, H. F. Die Ausbreitung der Reblaus Krankheit in Deutschland und deren Bekämpfung, unter Benützung von amtlichen Schriftstücken beleuchtet. Berlin: 1892. Cf. Wien. ent. Z. xi, p. 87.
- 492. KIEFFER, J. J. Beobachtungen über Gallmücken mit Beschreibung einiger neuen Arten. Wien. ent. Z. xi, pp. 212-224, pl. i. [Diptera.]
- 493. —. Zur Kenntniss der Weidengallmücken. Berl. Ent. Zeit. xxxvi, pp. 241-258, pl. ix. [Diptera.]
- 494. —. Mittheilungen über Gallmücken. T. c. pp. 259-266. [Diptera.]
- 495. —. Die Zoocecidien Lothringens. Ent. Nachr. xviii, pp. 43-46, 59-64, & 73-80.
- 496. KIRBY, W. F. A synonymic Catalogue of Lepidoptera Heterocera (Moths). Vol. I. Sphinges and Bombyces. London: 1892, 8vo, pp. xii & 951.
- Descriptions of three new species of Saturniidæ in the collection of the British Museum. Ann. N. H. (6) ix, pp. 173-175, pl. ix. [Lepidoptera.]
- 498. Elementary Text-book of Entomology. Second Edition.

  An appendix and index have been added, occupying pp. 240–281.
  - —. [See also SMITH & KIRBY (858).]
- 499. KLEMENSIEWICZ, S. Eine neue Art der Gattung Ellopia. Verh. z.-b. Wien, xlii, SB. p. 67. [Lepidoptera.]
- 500. KNOTEK, J. Zwei neue Scolytidz aus dem Occupationsgebiete. Wien. ent. Z. xi, pp. 234 & 235. [Coleoptera.]
- Translated from a paper in the Croatian language, published in an almost unknown Museum-journal.
- KOHL, F. F. Neue Hymenopterenformen. Ann. Hofmuseum Wien, vii, pp. 197-234, pls. xiii-xv.
- 502. Kolbe, H. J. Ueber die von Hauptmann Kling und Dr. R. Büttner in Togo (Ober-Guinea) gesammelten melitophilen Lamellicornier. S. E. Z. liii, pp. 125-142. [Coleoptera.]
- 503. —. Ueber Elaphinis atomosparsa, Fairm. T. c. p. 143.
  [Coleoptera.]

- 504. [Kolbe, H. J.] Ueber zwei neue Cicindelen aus Afrika. S. E. Z. liii, pp. 143-145. [Coleoptera.]
- 505. —... Ueber die Gattung Stephanorrhinæ und ihre Arten. T. c. pp. 145-155. [Coleoptera.]
- 506. —. Beiträge zur Kenntniss der Brenthiden. T. c. pp. 162-175. [Coleoptera.]
- Ueber die von Herrn Leopold Conradt in Deutsch-Ostafrika namentlich in der Gebirgslandschaft von Usambara gesammelten melitophilen Lamellicornier. SB. nat. Fr. 1892, pp. 62-75. [Coleoptera.]
- 508. —. Ueber die melitophilen Lamellicornier von Kamerun. T. c. pp. 235-261. [Coleoptera.]
- 509. —. Einführung in die Kenntniss der Insekten. Pts. 1-x. Berlin: 8vo.
  - Commenced in 1889: 528 pp. had appeared at the end of 1892.
- 510. Konow, F. W. Analytische Uebersicht der europäischen Arten der Tenthrediniden-Gattung Schizocera, Latr. Wien. ent. Z. xi, pp. 11-22. [Hymenoptera.]
- 511. KOROTNEFF, A. Histolyse und Histogenese des Muskelgewebes bei der Metamorphose der Insekten. Biol. Centralbl. xii, pp. 261–265. Summary in J. R. Micr. Soc. 1992, p. 468.
- 512. Kowarz, F. Die europäischen Arten der Dipteren-Gattung Lispa, Latr. Wien. ent. Z. xi, pp. 33-54.
- 513. Kraatz, G. Paussus cervinus, n. sp. Deutsche e. Z. 1892, p. 8. [Coleoptera.]
- 514. —. Monographische Revision der Ruteliden-Gattung Popillia, Serville. T. c. pp. 177-192 & 225-306. [Coleoptera.]
- Zwei neue Macronotiden-Gattungen von Java. T. c. pp. 312-314. [Coleoptera.]
- 516. —. Ueber Taniodera-Arten von Java und Malacca. T. c. pp. 315-320. [Coleoptera.]
- 517. —. Paussus opacus, n. sp. T. c. p. 377. [Coleoptera.]
- 518. —. Hadrosticta, n. g., Cetonidarum. T. c. p. 378. [Coleoptera.]
- 519. Krassilstschik, J. Zur Anatomie der Phytophthires. Zool. Anz. xv, pp. 217-223. [Rhynchota.]
  - Summary in J. R. Micr. Soc. 1892, p. 781.
- 520. KRAUSS, H. Dermapteren und Orthopteren aus Tunis. Wien. ent. Z. xi, pp. 143-150.
- 521. Systematisches Verzeichniss der canarischen Dermapteren und Orthopteren mit Diagnosen der neuen Gattungen und Arten. Zool. Anz. xv, pp. 163-171.
  - 1892. [VOL. XXIX.]

- 522. KRIECHBAUMER, —. Blattwespen Studien. Ent. Nachr. xviii, pp. 98-101. [Hymenoptera.]
- 523. Xylonomiden und Pimpliden-Studien. T. c. pp. 211-220.
- 524. Ophioniden-Studien. T. c. pp. 232-235. [Hymenoptera.]
- 525. —. Cryptiden-Studien. T. c. pp. 340-352, 362-365, & 370-373. [Hymenoptera.]
- 526. —. Ein neuer Psilomastus. T. c. p. 101. [Hymenoptera.]
- 527. KRIEGER, R. Zwei Hymenopteren-Zwitter. Z. Naturw. lxv, pp. 137-140.
- 528. KULAGIN, N. Zur Entwicklungsgeschichte der parasitischen Hautflügler. Zool. Anz. xv, pp. 85-87. [Hymenoptera.]
  Summary in J. R. Micr. Soc. 1892, p. 470.
- 529. KÜNCKEL, J. D'HERCULAIS. Le Criquet pèlerin (Schistocera peregrina) et ses changements de coloration; rôle des pigments dans les phénomènes d'histolyse et d'histogenèse qui accompagnent la metamorphose. C.R. exiv, pp. 240-242. [Orthoptera.]
  Summary in J. R. Micr. Soc. 1892, p. 468.
- 530. —.. Sur les changements de coloration et sur le rôle des pigments dans les phénomènes d'histolyse et d'histogenèse qui accompagnent les mues et la metamorphose du Criquet pèlerin. Bull. Soc. Ent. Fr. lxi, pp. xxv-xxvii.
- 531. —. Les invasions des Acridiens, vulgo Sauterelles, en Algérie pendant l'année 1891. C.R. Ass. Fr. Sci. xx, pt. 1, p. 241. [Orthoptera.]
- 532. Les invasions des Acridiens en Algérie en 1891. Observations biologiques et physiologiques. T. c. pt. 2, pp. 554-556. [Orthoptera.]
- 533. Kunze, E. Insects attracted by Fragrance or Brilliancy of Flowers for purposes of Cross fertilization. Canad. Ent. xxiv, pp. 173-177.
  - Kuwert, A. [See Heyden (402).]
- 534. Lameere, A. Nouveaux Lamiides de l'Afrique tropicale. Ann. Ent. Belg. xxxvi, pp. 506-509. [Coleoptera.]
- 535. —... Note sur les Fourmis de la Belgique. T. c. pp. 61-69. [*Hymenoptera*.]
- LAMPA, S. Berättelse till Landtbruksstyrelsen, 1891. Ent. Tidskr. xiii, pp. 1-38.
- 537. —... Agromyza lappæ, Loew. En för Sveriges fauna ny flugart. 1. c. pp. 117-120. [Diptera.]
- Kornflugan, Chlorops pumilionis, Bierk. T. c. pp. 257-274,
   pl. vii. [Diptera.]

``

- 539. LATASTE, F. Mode de preparation et d'emploi d'un ciment adapté au bouchage des flacons des collections d'objets d'histoire naturelle conservés dans l'alcool. (Ciment Emzed.) Act. Soc. Chili, ii, pp. 190-195.
- 540. La question du Criquet voyageur signalé l'an dernier dans les provinces australes du Chili. T. c. pp. 200-209. [Orthoptera.]
- 541. LATTER, O. H. The secretion of potassium hydroxide by *Dicranura vinula* (imago), and the emergence of the imago from the cocoon. Tr. E. Soc. 1892, pp. 287-292. [Lepidoptera.]
- 542. LEECH, J. H. Butterflies from China, Japan, and Corea. Parts 1 & 11. London: 1892, 4to, pp. 1-286, pls. i-xiii. [Lepidoptera.]
- 543. Lefèvre, E. Description of new species of South African Eumolpidæ. Tr. S. African Phil. Soc. vi, pp. 39-43. [Coleoptera.]
- 544. Description d'une nouvelle espèce de Clytrides. Bull. Soc. Ent. Fr. lxi, p. xciv. [Coleoptera.]
- 545. Léon, N. Labialtaster bei den Hemipteren. Zool. Anz. xv, pp. 145 & 146.
- 546. LEPRIEUR, C. E. La chasse aux Coléoptères. 2° edition, revue, augmentée et corrigée. Lyons: 1887, 8vo, 64 pp. Published as supplements to Échange.
- 547. Lesne, P. Collection d'insectes formée dans l'Indo-Chine par M. Pavie, Consul de France au Cambodge. Coléoptères, Anthribines. N. Arch. Mus. Paris (3) iii, pp. 225-228.
- 548. —... Sur un Braconide du genre Perilitus, Nees. Ann. Soc. Ent. Fr. lxi, pp. 305-308, pl. v. [Hymenoptera.]
- 549. LETHIERRY, L. Description d'un nouveau genre d'Homoptères d'Europe, et note sur le Delphax lugubris. Rev. d'Ent. xi, pp. 69 & 70. [Rhynchota.]
- 550. —. Liste d'Hemiptères récoltés à Mahé (Inde) par M. Em. Deschamps. Bull. Soc. Z. Fr. 1892, pp. 207-209.
- 551. LETZNER, K. Verzeichniss der Käfer Schlesiens. Zweite Auflage; Fortgesetzt und vollendet von J. Gerhardt. 8vo, xxvi pp. Published as a supplement to Z. Ent. Bresl. xvii.
- 552. Léveillé, A. Description d'un Temnochilide nouveau. Bull. Soc. Ent. Fr. lxi, p. xcv. [Coleoptera.]
- 553. Lewis, G. Viaggio di Leonardo Fea in Birmania e regioni vicine. xLII. Histeridæ, part 2. Ann. Mus. Genov. (2) xii, pp. 16-39. [Coleoptera.]
- 554. ——. On some Japanese species of Paromalus. Ann. N. H. (6) ix, pp. 32-38. [Coleoptera.]
- 555. On some new species of Histeridæ. T. c. pp. 341-357.
  [Coleoptera.]

- 556. [Lewis, G.] On the Japanese Clerida. Ann. N. H. x, pp. 183-192. [Coleoptera.]
- 557. —. On Eretmotus and Epiechnus (Histeridæ). T. c. pp. 231-236, pl. xix. [Coleoptera.]
- 558. On some new species of *Histerida*, and one new genus. Ent. M. M. xxviii, pp. 102-104. [Coleoptera.]
- 559. —. On a new Onthophilus from Mexico. T. c. p. 124. [Coleoptera.]
- 560. —. Note on Sternocælis, and on one new species. T. c. p. 263-[Coleoptera.]
- 561. —. On a new species of Histeridæ of Bengal. Ann. Eut. Belg. xxxvi, pp. 142 & 143. [Coleoptera.]
- 562. LIEBEL, R. Die Zoocecidien (Pflanzendeformationen) der Holzgewächse Lothringens. Ent. Nachr. xviii, pp. 257-287.
  Arranged phytologically.
- 563. LINDEN, GRÄFIN MARIA V. Beiträge zur Biologie der Phryganeiden. Biol. Centralbl. xii, pp. 523-527. [Neuroptera.]
- 564. LINTNER, J. A. Seventh report on the Injurious and other Insects of the State of New York. Albany: 1891, pp. 201-404. From the 44th Report of the New York State Museum.
- 565. LOWER, O. B. Descriptions of New South Australian Lepidoptera. Tr. B. Soc. S. Austr. xv, pp. 5-17.
- 566. —. Scientific Results of the Elder Exploring Expedition. Lepidoptera. Op. cit. xvi, pp. 10-15.
- 567. LOWNE, T. The Anatomy, Physiology, Morphology and Development of the Blowfly. Part III (Vol. i, pp. 215-350).
  - Devoted chiefly to embryology and development. Completes Vol. i.
- 568. Lucas, T. P. On Queensland and other Australian Lepidoptera, with descriptions of new species. P. Linn. Soc. N.S.W. (2) vi, pp. 277-306.
- 569. —. On twenty new species of Australian Lepidoptera. Op. cit. vii, pp. 249-266.
- 570. LYMAN, H. H. Can Insects survive freezing? Canad. Ent. xxxiv, pp. 1-6.
- 571. Mabille, P. Six Lépidoptères Hétérocères d'Afrique. Bull. Soc. Ent. Fr. lxi, pp. cxxxviii & cxxxix.
- 572. —. Nouvelle espèce du genre Zygana. T. c. p. cl. [Lepidoptera.]
- 573. MABILLE & VUILLOT. Novitates Lepidopterologica. Fasc. 7-9, pls. viii-xii, pp. 49-89.
- 574. MacLachlan, R. Supplementary note on the Neuroptera of the Hawaiian Islands. Ann. N. H. (6) x, pp. 176-178.

- 575. MAGRETTI, P. Di alcune specie d'Imenotteri raccolte dall' Ing. L. Bricchetti Robecchi nel paese dei Somali. Ann. Mus. Genov. (2) x, pp. 950-960.
- 576. —... Viaggio di Leonardo Fea in Birmania e regioni vicine. XLIII. Imenotteri. Parte prima. Mutillidei, Scoliidei, Tifiidei, Tinnidei, colla descrizione di parecchie unove specie. Op. cit. xii, pp. 197-266, pl. v.
- 577. Mally, F. W. Report of progress in the investigation of the Cotton Boll Worm. Bull. Dept. Agric. Ent. xxvi, pp. 45-56. [Lepidoptera.]
- 578. —. An insectivorous primrose (*Enothera speciosa*). P. E. Soc. Wash, ii, pp. 288 & 289.
- 579. Manders, N. The Butterflies of Rawal Pindi and the Murree Hills (Punjab). Ent. M. M. xxviii, pp. 88-92 & 130-132. [Lepidoptera.]
- 580. MARCHAL, P. Notes sur la vie et les mœurs des Insectes. Observations sur l'Ammophila affinis. Arch. Z. expér. (2) x, pp. 23-36. [Hymenoptera.]
  - Summary in J. R. Micr. Soc. 1892, p. 471.
- 581. MARLATT, C. L. A study of the ovipositor in Hymenoptera. P. E. Soc. Wash. ii, pp. 201-205.
- 582. —... Notes on the early stages of three Moths. Tr. Kansas Ac xi, pp. 110-114. [Lepidoptera.]
- 583. Marshall, T. A. On a new genus and species of Belytidæ from New Zealand. Ent. M. M. xxviii, p. 275. [Hymenoptera.]
  - —. [See also Buysson & Marshall (149).]
- 584. MARTEL, V. Les Cécidies des environs d'Elbeuf. Bull. Soc. d'Elbeuf, 1891, pp. 79-134.
  - Arranged according to plants on which the galls occur.
- 585. MARTIN, J. Sur la respiration des larves de Libellules. Bull. Soc Philom. (8) iv, pp. 122-124. [Neuroptera.]
- 586. MARTIN, L. Aus meinen Tagebuche. Berl. Ent. Zeit. xxxvii, pp. 291-298. [Lepidoptera.]
  - Jottings of a collector of Lepidoptera in Sumatra.
- 587. MARTIN, R. Les Trichoptères du département de l'Indre. Rev. d'Ent. xi, pp. 1-23 & 76. [Neuroptera.]
- 588. Les Perlides du département de l'Indre. T. c. pp. 198-201. [Neuroptera.]
- 589. —. Les Psocides du département de l'Indre. T. c. pp. 285-288. [Neuroptera.]

- 590. Maskell, W. M. Further Coccid notes; with descriptions of new species, and remarks on Coccids from New Zealand, Australia, and elsewhere. Tr. N. Z. Inst. xxiv, pp. 1-64, pls. i-xiii.
- 591. . Migrations and new localities of some Coccids. Ent. M. M. xxviii, pp. 69-71.
- 592. —... A new Icerya, and some other Coccids from Australia. T. c. pp. 183 & 184.
- 593. MATTHEWS, A. Notes on Dr. C. Flach's synonymic list of the European Trichopterygidæ. Ann. N. H. (6) ix, pp. 442-448. [Coleoptera.]
- 594. MAYER, P. Zur Kenntniss von Coccus cacti. MT. z. Stat. Neap. x, pp. 505-518, pl. xxxii. [Rhynchota.]
- 595. MEADE, R. H. Speciei novæ Tachinidarum descriptio. Wien. ent. Z. xi, p. 114. [Diptera.]
- 596. —. Description d'un Diptère nouveau. Bull. Soc. Ent. Fr. lxi, p. cxvi.
- 597. —... Annotated list of British Tachinidæ. Ent. M. M. xxviii, pp. 17, &c., &c. [Diptera.]
- 598. MEDINA, —. Nuevas especies de Véspidos de España. Act. Soc. Esp. xxi, pp. 142-144. [Hymenoptera.]
- 599. Meinert, F. Træk af Insektlivet i Venezuela. Ent. Medd. iii, pp. 125-166.
- 600. MERRIFIELD, F. The effects of Artificial Temperature on the Colouring of several species of *Lepidoptera*, with an account of some experiments on the effect of light. Tr. E. Soc. 1892, pp. 33-44. Summary in J. R. Micr. Soc. 1892, p. 469.
- 601. —. Double-broodedness: whether influenced by the state of the food-plant? Ent. M. M. xxviii, p. 20. [Lepidoptera.]
- 602. MEUNIER, F. Aperçu des genres de Dolichopodidæ de l'ambre suivi du Catalogue bibliographique des Diptères fossiles de cette Résine. Ann. Soc. Ent. Fr. lxi, pp. 377-384.
- 603. MEYRICK, E. On the classification of the Geometrina of the European fauna. Tr. E. Soc. 1892, pp. 53-140, pl. iii. [Lepidoptera.] Reviewed by REBEL (714).
- Revision of Australian Lepidoptera. v. P. Linn. Soc. N.S.W.
   vi, pp. 581-678.
- 605. —. On new species of Lepidoptera. Tr. N. Z. Inst. xxiv, pp. 216–220.
- 606. Mik, J. Dipterologische Miscellen. xx. (1 Serie.) Wien. ent. Z. xi, pp. 55-58.
  - Includes a table of contents of the twenty memoirs forming the series.

- 607. [Mik, J.] Dipterologische Miscellen. (2 Serie.) Wien. ent. Z. xi, pp. 116, &c.
- 608. —... Zur Kenntniss der Dipteren Gattung Hilara. T. c. pp. 78–85, [Diptera.]
- Ueber ein neues hochalpines Dipteron aus der Familie der Dolichopodiden. T. c. pp. 279-281.
- 610. . Ueber zwei Cecidomyiden-Gallen aus Tirol. T. c. pp. 306-308, pl. iii. [Diptera.]
- 611. Miskin, W. H. A revision of the Australian Sphingidæ. P. R. Soc. Queensl. viii, pt. iii, pp. [Lepidoptera.]

This and the two following are recorded from separate copies; although taken from Vol. viii of the Queensland Society, Vol. vii does not appear to have yet reached London.

- 612. —. Further note on Australian Sphingidæ. T. c. pp. [Lepidoptera.]
- 613. —. New species of Australian Macro-Lepidoptera. T. c. pp
- 614. MOBERG, J. C. Om en Hemipter från Sveriges Undre Graptoletakiffer. Geol. Fören. Stockholm Förh. xiv, pp. 121-124.
- 615. Mocsáry, A. Hymenoptera in expeditione Comitis Belæ Szechenyi in China et Tibet a Dom G. Kreitner et L. Lóczy anno 1879 collecta. Term. füzetek, xv, pp. 126–131.
- 616. MONIEZ, R. Mémoire sur quelques Acariens et *Thysanoures* parasites ou commensaux des Fourmis. Rev. Biol. iv, pp. 377-391.
- 617. Montandon, A. Deux Hémiptères nouveaux. Rev. d'Ent. xi, pp. 73-76.
- 618. Hémiptères-Hétéroptères nouveaux. T. c. pp. 265-273.
- 619. Hémiptères Plataspides nouveaux. T. c. pp. 273-284.
- 620. Études sur la sous-famille des Plataspidinæ. T. c. pp. 294-308. [Rhynchota.]
- 621. Monti, R. Ricerche microscopiche sul sistema nervoso degli insetti.

  Rend. Ist. Lombardo (2) xxv, pp. 533-540 & 654.
- Reported in Arch. Ital. Biol. xviii, pp. 62-70. 622. MOORE, F. Lepidoptera Indica. Pts. IX-XII.
  - Vol. I is completed, and consists of xii and 317 pp. and 94 pls.
- 623. —. Descriptions of some new species of Asiatic Satu rniida. Ann. N. H. (6) ix, pp. 448-453. [Lepidoptera.]
- 624. MORAWITZ, F. Hymenoptera aculeata, rossica nova. Hor. Ent. Ross. xxvi, pp. 132-181.
- 625. Morgan, A. C. F. A new genus and species of Aleurodidæ. Ent. M. M. xxviii, pp. 29-33, pl. i. [Rhynchota.] The description is partly by Douglas.
- 626. —. Observations on Coccides. No. 9. T. c. pp. 12-16.

- Morgan, A. C. F. [See also Cockerell (179).]
- 627. MORTENSEN, R. C. Lucilia sylvarum, Meig., als Schmarotzer an Bufo vulgaris. Zool. Anz. xv, pp. 193-195. [Diptera.]
- 628. Müggenburg, F. H. Der Rüssel der Diptera pupipara. Arch. f. Nat. lviii, i, pp. 287-332, pls. xv & xvi. Summary in J. R. Micr. Soc. 1893, p. 35.
- 629. MULLER, G. W. Beobachtungen an im Wasser lebenden Schmetterlingsraupen. Zool. Jahrb. Syst. vi, pp. 617-630, pl. xxviii. [Lepidoptera.]
- 630. MURTFELDT, M. E. Entomological notes for the season of 1891. Bull. Dep. Agric. Ent. xxvi, pp. 36-44.
- 631. NERÉN, C. H. Entomologiska Anteckningar. Ent. Tidskr. xiii, pp. 57-68. [Hymenoptera.]
- 632. —. Entomologiska Sommarstudier. T. c. pp. 97-116. [Hymenoptera.]
- 633. Neumoegen, B. Some beautiful new Bombycids from the West and Northwest. Canad. Ent. xxiv, pp. 225-228. [Lepidoptera.]
- Description of a new Cossid from Texas. Ent. News. iii, p. 258. [Lepidoptera.]
- 635. NEVINSON, B. G. Description of three new species of the genus *Phanœus*. Ent. M. M. xxviii, pp. 33-35. [Coleoptera.]
- 636. —... Revised synonymic list of species in the genus *Phanœus*, Macleay. London, 1892, 8vo, 10 pp.
- 637. Newstead, R. On new or little known *Coccida*, chiefly English. Ent. M. M. xxviii, pp. 141-147, pl. ii.
- 638. NICÉVILLE, L. DE. On new and little known Butterflies from the Indo-Malayan region. J. Bomb. N. H. Soc. vii, pp. 322-356, pls. H, I, J. [Lepidoptera.]
- 639. NICOLAS, H. Les *Hymenoptères* et leurs parasites. Échange, 1890 & 1891, pp. 189, &c., &c.
- 640. —. Observations entomologiques et autres faits au sommet du Mont Ventoux. C.R. Ass. Fr. Sci. xx, pt. 2, pp. 566-571. [Hymenoptera.]
  - Effects on Osmia of transplanting it to summit.
- 641. Nolan, E. J. The introduction of the Ailanthus Silkworm Moth. Ent. News, iii, pp. 193-195. [Lepidoptera.]
- 642. NONFRIED, A. F. Neue afrikanische, central-amerikanische und ost-asiatische Melolonthiden und Ruteliden. Berl. Ent. Zeit. xxxvi, pp. 221-240. [Coleoptera.]
- 643. —. Verzeichniss der Rutelidæ beschrieben nach der Herausgabe des Münchener Kataloges. T. c. pp. 347-358 & 449-454. [Coleoptera.]

- 644. [Nonfried, A. F.] Weitere Beiträge zur Käferfauna von Süd-asien und Neu-guinea. Berl. Ent. Zeit. xxxvi, pp. 359-380. [Coleoptera.]
- 645. Monographische Uebersicht der Prioniden-gattung Callipogon, Serv. Op. cit. xxxvii, pp. 17-24, pl. iii. [Coleoptera.]
- 646. —... Verzeichniss der seit 1871 neu beschriebenen Glaphyriden, Melolonthiden und Euchiriden. T. c. pp. 249-290. [Coleoptera.]
- 647. —. Verzeichniss der um Nienghali in Südchina gesammelten Lucanoiden, Soarabæiden, Buprestiden und Cerambyciden, nebst Beschreibung neuer Arten. Ent. Nachr. xviii, pp. 81-95. [Coleoptera.]
- 648. Beiträge zur Coleopteren-fauna von Africa und Madagascar. T. c. pp. 105-111, 117-126, & 136-141.
- 649. Nunney, W. H. The British Perlida or Stone-flies. Sci. Goss. 1892, p. 35, &c.
- 650. OBERTHUR, C. Études d'entomologie. Faunes entomologiques. Descriptions d'insectes nouveaux ou peu connus. 16<sup>me</sup> Livraison, pp. x & 9, pls. i & ii. [Lepidoptera.]
- 651. —... Étude sur une collection de Lépidoptères formée sur la côte de Malabar et à Ceylan par M. Emile Deschamps. Mém. Soc. Zool. v, pp. 237-252.
  - Some notes on metamorphoses.
- 652. OLIVIER, E. Lumpyrides rapportés de Birmanie par M. L. Fea, avec descriptions des espèces nouvelles. Ann. Mus. Genov. (2) x, pp. 595-604. [Coleoptera.]
- 653. ——. Descriptions de deux nouvelles espèces du genre Luciola. T. c. pp. 1010 & 1011. [Coleoptera.]
- 654. Olliff, A. S. Further remarks on the saltbush Scale (Pulvinaria maskelli, Oll.). Agric. Gaz. N.S.W. iii, pp. 176-180. [Coccidæ-Lepidoptera.]
- 655. A new Longicorn Beetle attacking Orange trees. T. c. pp. 895-897, pl. lvii. [Coleoptera.]
- 656. ORMEROD, E. A. Report of observations of injurious insects and common farm pests, with special report on attack of Caterpillars of the Diamond-back Moth, during the year 1891, with methods of prevention and remedy. Fifteenth Report. London: 1892, 8vo, pp. vi & 170.
- 657. OSBORN, H. Insects of the Season in Iowa. Bull. Dep. Agric. Ent. xxvi, pp. 57-62.
  - —. [See also Bruner, Coquillett, & Osborn (136).]
- 659. OSTEN-SACKEN, C. R. Second notice on the Apiocerina. Berl. Ent. Zeit. xxxvi, pp. 311-315. [Diptera.]

- 660. [OSTEN-SACKEN, C. R.] Synopsis of the described genera and species of the *Blepharoceridæ*. T. c. pp. 407-411. [Diptera.]
- 661. —. On the Chaptotaxy of Caccoxenus indagator, Lw. T. c. pp. 411 & 412. [Diptera.]
- 662. Synonymy of Antocha, O.-S., and Orimargula, Mik. T. c. pp. 413-416. [Diptera.]
- 663. —. Additions and corrections to the Catalogue of the described species of South-American Asilidæ, by S. W. Williston. T. c. pp. 417-428. [Diptera.]
- 664. OUDEMANS, J. T. Het prepareeren van Rupsen. Tijdschr. Entxxxv, pp. 27-30, pl. ii. [Lepidoptera.]
- 665. PACKARD, A. S. On the scale-like and flattened hairs of certain Lepidopterous larvæ. Ann. N. H. (6) ix, pp. 372-375.
- 666. On the larva of *Lagoa*, a Bombycine Caterpillar with seven pairs of abdominal legs; with notes on its metameric glandular abdominal processes. Zool. Anz. xv, pp. 229-234. [*Lepidoptera*.]
- 667. PAGENSTECHER, A. Ueber einige neue arten des Pyraliden genus Tetraphana, Ragonot. Deutsche e. Z. Lep. v, pp. 1-4. [Lepidoptera.]
- 668. —. Ueber die Familie der Siculiden. T. c. pp. 5-131. [Lepidoptera.]
  - Summary and statistics by Seitz, S. E. Z. liii, pp. 155 & 156.
- 669. —. Nachträge über die Familie der Siculiden. T. c. pp. 443-449. [Lepidoptera.]
- 670. PATTON, W. H. Notes upon Larradæ. Ent. News, iii, pp. 89 & 90. [Hymenoptera.]
- 671. Pergande, T. Peculiar habit of Ammophila gryphus, Sm. P. E. Soc. Wash. ii, pp. 256-258. [Hymenoptera.]
- 672. PÉRINGUEY, L. Third contribution to the South African Coleopterous Fauna. Tr. S. African Phil. Soc. vi, part 2, pp. 1-94.
- 673. ——. Fourth contribution to the South African Coleopterous Fauna. Description of new Coleoptera in the South African Museum. T. c. pp. 95-134.
- 674. Perkins, R. C. L. Stylopized Bees. Ent. M. M. xxviii, pp. 1-4.
- 675. Petersen, W. Die Entwickelung des Schmetterlings nach dem Verlassen der Puppenhülle. Deutsche e. Z. Lep. iv, pp. 199-214. [Lepidoptera.]
- 676. . Ueber die Ungleichzeitigkeit in der Erscheinung der Geschlechter bei Schmetterlingen. Zool. Jahrb. Syst. vi, pp. 671-679. [Lepidoptera.]

- 677. Pic, M. Cortodera reitteri, n. sp. Échange, 1891, No. 78, p. 43. [Coleoptera.]
- 678. —. Descriptions et corrections. Op. cit. 1892, p. 4. [Coleoptera.]
- 679. —. Petite étude sur le genre Stenopterus, Steph. T. c. pp. 21-23. [Coleoptera.]
- 680. —. Contributions à l'Étude des Anthicides d'Algérie. T. c. p. 102. [Coleoptera.]
- 681. —. Conizonia leprieuri, n. sp. Bull. Soc. Ent. Fr. lxi, p. lii. [Coleoptera.]
- 682. —. Anthicus chobauti, n. sp. T. c. p. lxxv. [Coleoptera.]
- 683. —. Delagrangeus, n. gen. T. c. pp. xciii & xciv. [Coleoptera.]
- 684. —. Descriptions de Coléoptères. T. c. pp. excvi-excviii.
- 685. Description de deux Anthicus nouveaux. T. c. p. cexxi. [Coleoptera.]
- 686. Diagnoses de *Polyarthron* nouveaux. *T. c.* pp. cclix & cclx. [Coleoptera.]
- 687. —. Descriptions de nouveaux Anthicus. T. c. pp. celxviii-celxxii. [Coleoptera.]
- 688. —. Voyage de M. Charles Delagrange dans la Haute-Syrie. Longicornes. Ann. Soc. Ent. Fr. lxi, pp. 413-416. [Coleoptera.]
- 689. —... Anthicides recueillis en Algérie en avril et mai 1892. Rev. Sci. Bourb. v, pp. 212-220. [Coleoptera.]
- 690. —. Sur les Tomoderus, Laf. T. c. pp. 240-242. [Coleoptera.]
- 691. —. Descriptions de Longicornes d'Algérie. Abeille, xxvii, pp. 289 & 290. [Coleoptera.]
- 692. PICTET, A., & SAUSSURE, H. DE. Iconographie de quelques Sauterelles vertes. Geneva: 1892, 4to, 26 pp., 3 pls. [Orthoptera.]
- 693. PLATEAU, F. Une forme speciale de Colonies temporaires de Coccinella septempunctata. Ann. Ent. Belg. xxxvi, pp. 393-396. [Coleoptera.]
- 694. —. Les chenilles carnassières. Le Nat. 1892, pp. 71 & 72. [Lepidoptera.]
- 695. La ressemblance protectrice dans le regne animal. Bull. Ac. Belg. (3) xxiii, pp. 89-135.
- 696. Poll, J. R. H. N. van de. Contributions à la Faune Indo-Chinoise.
   9e Mémoire. Buprestidæ. Ann. Soc. Ent. Fr. lxi, pp. 17-26.
   [Coleoptera.]
- 697. —... On new or little-known Australian Longicornia. I. Tijdschr. Ent. xxxiv, pp. 219–228, pl. xiii. [Coleoptera.]

698. PORTSCHINSKY, J. Biologie des mouches coprophagues et nécrophagues. Première partie. Hor. Ent. Ross. xxvi, pp. 63-131. [Diptera.]

Entirely in the Russian language.

699. —. Diptera europæa et asiatica nova aut minus cognita. T. c. pp. 201-227, pl. i.

In the Russian language, but with Latin diagnoses.

700. —. Lepidopterorum Rossim biologia. III. Coloration marquante et taches ocellées, leur origine et leur développement. Suite. T. c. pp. 258-411, pls. ii & iii.

Entirely in Russian.

 POSTANS, R. B. Butterflies. Tr. Eastbourne Soc. (n.s.) ii, pp. 259-292, pl.

Interesting considerations of a popular character.

- 702. POUJADE, G. A. Collection d'insectes formée dans l'Indo-Chine par M. Pavie, Consul de France au Cambodge. *Lépidoptères*. N. Arch. Mus. (3) iii, pp. 255–276, pl. xi.
- 703. POULTON, E. B. Further experiments upon the colour-relation between certain *Lepidopterous* larvæ, pupæ, cocoons, and imagines and their surroundings. Tr. Ent. Soc. 1892, pp. 293-487, pls. xiv & xv.
- 704. Pratt, A. E. To the Snows of Tibet through China. London: 8vo. Includes information as to localities of several Chinese Lepidoptera.
- PROCHÁZKA, J. Revision der Arten der Coleopteren-Gattung. Hapalus, Fabr. Wien. ent. Z. xi, pp. 263-270.
- 706. PÜNGELER, R. Ueber zwei Psychiden-Arten aus Sizilien. Deutsche e. Z. Lep. v, pp. 133-140. [Lepidoptera.]
- 707. PUTON, A. Descriptions de trois Hémiptères nouveaux. Rev. d'Ent. xi, pp. 71 & 72.
- 708. —. Hémiptères nouveaux ou peu connus. T. c. pp. 24-36.
- 709. RAFFRAY, A. Recherches Anatomiques sur le Pentaplatarthus paussoides. Coléoptère de la famille des Paussides. N. Arch. Mus. (3) iv, pp. 91-102, pl. xiii.
- 710. & BOLIVAR, I. Étude sur les Arthropodes cavernicoles de l'fie de Luzon. Ann. Soc. Ent. Fr. lxi, pp. 27-34, pl. i. [Coleoptera, Orthoptera.]
- RAGONOT, E. L. Trois espèces inédites de Microlépidoptères. Bull. Soc. Ent. Fr. lxi, p. lxxxii.
- 712. Diagnoses d'un genre nouveau et de deux espèces nouvelles de Phycites. T. c. pp. coxxxv & coxxxvi. [Lepidoptera.]

- 713. RATH, O. VOM. Zur Kenntniss der Spermatogenese von Gryllotalpa vulgaris. Arch. mikr. Anat. xl, pp. 102-132, pl. v. [Orthoptera.]
  - Summary in J. R. Micr. Soc. 1892, p. 781.
- 714. Rebel, H. Zwei Geometriden-Classificationen. S. E. Z. liii, pp. 247-268. [Lepidoptera.]
- Meyrick's & Gumppenberg's. Some points in the work of the former are much objected to; he considers it a matter for surprise that Gumppenberg's work should be published by a great Society.
- Beitrag zur Microlepidopteren-fauna des canarischen Archipels. Ann. Hofmuseum Wien, vii, pp. 241-284, pl. xvii.
- Beitrag zur Lepidopteren-fauna Südtirols, insbesondere der Umgebung Bozens. Verh. z.-b. Wien, xlii, pp. 509-536.
- 717. REDTENBACHER, J. Monographische Uebersicht der Mecopodiden. T. c. pp. 183-224, pl. iii. [Orthoptera.]
  - ---. [See also Brunner & Redtenbacher (138).]
- 718. REED, E. C. Entomología Chilena, sinonimia. Act. Soc. Chili, i, pp. 66-69. [Lepidoptera, Diptera.]
- 719. —... Revision de las Abejas Chilenas descritas en la obra de Gay. Op. cit. ii, pp. 223-240. [Hymenoptera.]
- 720. RÉGIMBART, M. Enumération des Dytiscidæ et Gyrinidæ recueillis par M. Leonardo Fea, dans les voyages en Birmanie et regiones voisines. Ann. Mus. Genov. (2) x, pp. 536-554. [Coleoptera.]
- 721. —. Viaggio di Lamberto Loria nella Papuasia orientale. IV Haliplida, Dytiscida, et Gyrinida. T. c. pp. 978-997. [Coleoptera.]
- 722. Description de deux Hydrocoptus (Dytiscidæ) nouveaux. Notes Leyd. Mus. xiv, pp. 11 & 12. [Coleoptera.]
- 723. —. Insectes du Bengale occidental. Hydrocanthares. Ann. Ent. Belg. xxxvi, pp. 112-121. [Coleoptera.]
- 724. REITTER, E. Coleopterologische Notizen. Wien. ent. Z. xi, pp. 25, &c.
- Fünfter Beitrag zur Coleopteren-fauna des russischen Reiches.
   c. pp. 59-68. Sechster Beitrag, t. c. pp. 133-135. Siebenter Beitrag, t. c. pp. 151-154.
- T. c. pp. 93-96.
- 727. —... Ueber eine Lasiopsis-Art aus dem Kaukasus. T. c. p. 101. [Colsoptera.]
- 728. ... Uebersicht der bekannten Arten der Coleopteren-Gattung Apotomus, Ill., aus der paläaretischen Fauna. T. c. pp. 137 & 138.
- 729. . Uebersicht der Arten der Coleopteren-Gattung Monotropus, Er. T. c. p. 142.

- 730. REUTER, O. M. Hemiptera Heteroptera från trakterna Kring Sajanska bärgskedjan. Œfv. Finska Förh. xxxiii, pp. 166-208.
- Voyage de M. E. Simon au Venezuela. Hemiptères-Hétéroptères. 1ere partie. Capsides. Ann. Soc. Ent. Fr. lxi, pp. 391-402.
- 732. A new species of *Helopeltis*. Ent. M. M. xxviii, p. 159 [Rhynchota.]
- 733. Species Capsidarum et Anthocoridarum madagascarienses, descripsit. T. c. pp. 185-187. [Rhynchota.]
- 734. —... Podurider från nordvestra Sibirien. Œfv. Finska Förh. xxxiii, pp. 226-229. [Thysanura.]
- 735. REY, C. Remarques en passant. Échange i, No. 17, &c., &c.

It is not possible to treat the names mentioned in this series of papers as representing new forms, though many of them are probably thought to do so by their author.

- 736. —. Description d'une nouvelle espèce de Curculionide. Op. cit. 1888, No. 45, p. 4. [Coleoptera.]
- RIBBE, C. Anleitung zum K\u00e4fersammeln in tropischen L\u00e4ndern. Berl. Ent. Zeit. xxxvii, pp. 125-138.
- 738. RICHARDSON, N. M. On a case of apparent substitution of a wing for a leg in a Moth. P. Dorset Club, xi, pp. 64-73. [Lepidoptera.]
- 739. —... On some of the chief peculiarities in the *Lepidopterous* fauna of Portland. T. c. pp. 46-63, pl.
- 740. —. Notes on Dorset Lepidoptera in 1891. Op. cit. xiii, pp. 168–177, pl.
- 741. RILEY, C. V. Some interrelations of plants and insects. P. Biol. Soc. Washington, vii, pp. 81-104.
- On certain peculiar structures of Lepidoptera. P. E. Soc. Wash. ii, pp. 305-312.
- 743. —. New species of *Prodoxidæ*. T. c. pp. 312-319. [Lepidoptera.]
- 744. —. Rose Sawflies in the United States. Ins. Life, v, pp. 6-11. [Hymenoptera.]
- The larger Digger-wasp. Op. cit. iv, pp. 248-252. [Hymenoptera.]
- 746. —. On the larva and some peculiarities of the cocoon of Sphecius speciosus. P. E. Soc. Wash. ii, pp. 170-172. [Hymenoptera.]
- 747. A probable Microgaster parasite of Eleodes in the perfect state. T. c. p. 211. [Hymenoptera.]
- 748. —... Mexican Jumping Bean: the determination of the plant. T. c. pp. 178-181. [Lepidoptera.]

- 749. [RILEY, C. V.] Further note on Carpocapsa saltitans and on a new Grapholitha producing Jumping Beans. P. E. Soc. Wash. ii, pp. 213 & 214. [Lepidoptera.]
- 750. ——. Coleopterous larvæ with so-called dorsal pro-legs. T. c. pp. 319-323.
- 751. —... The first larval or post-embryonic stage of the Pea and Bean Weevils. Canad. Ent. xxiv, pp. 185 & 186. [Coleoptera.]
- 752. —. On the nomenclature and on the oviposition of the Bean Weevil. Ins. Life, v, pp. 27-33. [Coleoptera.]
- 753. —. The Ox Bot in the United States. Habits and Natural History of Hypoderma lineata. Op. cit. iv, pp. 302-317. [Diptera.]
- 754. —. The Pea and Bean Weevils. T. c. pp. 297-302. [Coleoptera.]
- 755. RIPPON, R. H. F. Icones Ornithopterorum. Pt. v. London: 1892. [Lepidoptera.]
- 756. ——. Description of a new species of Ornithoptera, of the priamus group, in the collection of the Hon. L. Walter Rothschild. Ann. N. H. (6) x, pp. 193-196. [Lepidoptera.]
- 757. RITSEMA, C. The genus *Helota*, as represented in the Civic Museum of Natural History at Genoa, with descriptions of the new species collected in Burma by Mr. L. Fea. Ann. Mus. Genov. (2) x, pp. 885-902. [Coleoptera.]
- 758. Description of a new species of the Lucanoid genus Cyclommutus, and list of the described species. Notes Leyd. Mus. xiv, pp. 1-6. [Coleoptera.]
- 759. A new Lucanoid Beetle from Java. T. c. p. 31. [Coleoptera.]
- 760. A new Longicorn Beetle. T. c. p. 38. [Coleoptera.]
- 761. —... Cyclommatus squamosus, a new species of Lucanid from Borneo. T. c. p. 54. [Coleoptera.]
- 762. —. Prosopocælus tarsalis, a new Lucanid. T. c. p. 191. [Coleoptera.]
- Three new species of the Longicorn genus Pachyteria. T. c. pp. 213-220. [Colsoptera.]
- 764. —. Two new species of the Longicorn genus Glenea. T. c. xiv, pp. 221-224. [Coleoptera.]
- 765. ROBBE, H. Insectes du Bengale occidental. Liste d'une collection de Lépidoptères recueillis au Bengale occidental; avec une description d'une variété nouvelle et quelques considerations sur des espèces connues. Ann. Ent. Belg. xxxvi, pp. 122-131.
- 766. —. Lépidoptères du Congo. Description de deux nouvelles espèces et de deux nouvelles variétés. T. c. pp. 132-134.

767. RÖBER, J. Beitrag zur Kenntniss der Indo-australischen Lepidopteren-fauna. Tijdschr. Ent. xxxiv, pp. 261-334.

There are plates to this memoir that appear, unfortunately, only in the next volume.

- 768. ROBERTSON, C. Descriptions of new North-American Bees. Am. Nat. xxvi, pp. 267-274. [Hymenoptera.]
- 769. RÖDER, V. v. Ueber die Dipteren-Gattung Lyroneurus, Löw, nebst Beschreibung einer neuen Art. Soc. Ent. vii, p. 81.
- Brachyceræa novum genus Conopidarum. Ent. Nachr. xviii,
   p. 366. [Diptera.]
- Drei neue Dipteren, aus der Sendung des Herrn Baron von Müller, in Melbourne, an das Königliche Naturalien-Cabinet in Stuttgart. S. E. Z. liii, pp. 241-244.
- 772. ROELOFS, W. Description d'une espèce nouvelle du genre Eugithopus. Notes Leyd. Mus. xiv, p. 7. [Coleoptera.]
- 773. Observations sur les espèces du genre Oxyopisthen et des genres voisins. T. c. pp. 33-37. [Coleoptera.]
- 774. Description de deux nouvelles espèces du genre Onychogymnus. T. c. pp. 49-53. [Coleoptera.]
- 775. Description d'un nouveau genre et d'une nouvelle espèce de Curculionides de la tribu des Ulomascides. T. c. pp. 136-138. [Coleoptera.]
- 776. —... Deux nouveaux genres et deux nouvelles espèces du groupe des Rhynchophorides. T. c. pp. 207-212. [Coleoptera.]
- 777. ROGENHOFER, A. F. Ueber die taschenförmigen Hinterleibs-Anhänge der weiblichen Schmetterlinge der Acræiden. Verh. z.-b. Wien, xlii, pp. 579-581. [Lepidoptera.]
- 778. Neue Lepidopteren des k. k. naturhistorischen Hofmuseums. T. c. pp. 571-575.
- 779. ROMANOFF, N. M. Mémoires sur les Lépidoptères. Tome vi. St. Pétersbourg: 1892, 8vo, 770 pp., 16 pls.

The contents are:—1. Alphéraky, S.: Lépidoptères rapportés de la Chine et de la Mongolie par G. N. Potanine; pp. 1-81, pls. i-iii. 2. Staudinger, O.: Die Macrolépidopteren des Amurgebiets; I Theil, Rhopalocera, Sphinges, Bombyces, Noctuæ, pp. 83-658, pls. iv-xiv. 3. Standfus, M.: Lepidopterologisches; pp. 659-669, pl. xv. 4. Erschoff, N.: Verzeichniss von Schmetterlingen aus Central-Siberien; pp. 670-672, pl. xvi.

- 780. Rost, —. Plectes polychrous, n. sp., &c. Deutsche e. Z. 1892, pp. 401 & 402. [Coleoptera.]
- 781. ROSTOCK, M. Capnodes schilleri, eine neue deutsche Perlide. Berl. Ent. Zeit. xxxvii, pp. 1-5, pl. i. [Neuroptera.]

- 782. ROTHSCHILD, W. On a little-known species of Papilio from the Island of Lifu, Loyalty group. Tr. E. Soc. 1892, p. 141, pl. iv. [Lepidoptera.]
- 783. —... Notes on a collection of *Lepidoptera* made by William Doherty in Southern Celebes, during August and September, 1891. Part I. *Rhopalocera*. Deutsche e. Z. Lep. v, pp. 429-442.
  - ---. [See also BUTLER & ROTHSCHILD (147).]
- 784. RÜBSAAMEN, E. H. Mittheilungen über Gallmücken. Verh. z.-b. Ges. Wien, xlii, pl. ii. [Diptera.]
- Weber Gallmückenlarven. Berl. Ent. Zeit. xxxvi, pp. 381-392, pl. xiv. [Diptera.]
- 786. —. Neue Gallmücken und Gallen. T. c. pp. 393-406. [Diptera.]
- 787. Die Gallmücken des Konigl. Museums für Naturkunde zu Berlin. Op. cit. xxxvii, pp. 319-411, pls. vii-xviii. [Diptera.]
- 788. Ruge, —. Tetracha horni, n. sp. Deutsche e. Z. 1892, p. 130. [Coleoptera.]
- 789. RÜHL, F. Die palæarktischen Gross-schmetterlinge und ihre Naturgeschichte. Leipzig: 8vo.
  - To be completed in 75 parts; the first is published.
- Neue europäische Dasypolien. Soc. Ent. vi, pp. 169 & 170.
   [Lepidoptera.]
  - Salvin, O. [See Godman & Salvin (347).]
- Sangagli, G. Echinococco ed Œstrus nell'uomon. Rend. Ist. Lombardo (2) xxv, pp. 1167-1178. [Diptera.]
- 792. SAUNDERS, E. The Hemiptera-Heteroptera of the British Islands. London: 8vo, vi & 350 pp., 32 pls.
- 793. SAUSSURE, H. DE. Histoire naturelle des Hymenoptères. Histoire physique, naturelle et politique de Madagascar, publiée par Alfred Grandidier. Vol. xx, première partie, pp. 177-590, pls. xxi-xxvii.
- Completes the part, the antecedent portion having been recorded last year. The parasitica are not described, though figured on the plates.
- 794. —. Orthoptera Centrali-Americana. Soc. Ent. vii, pp. 121-124.
- 795. —. Note supplémentaire à la Synopsis de la tribu des Sagiens, Orthoptères de la famille des Locustides. Ann. Soc. Ent. Fr. lxi, pp. 5-16.
  - \_\_\_\_. [See also DISTANT (210) and PICTET & DE SAUSSURE (692).]
- 796. SAWTSCHENKO, J. Die Beziehung der Fliegen zur Verbreitung der Cholera. CB. Bakt. Parasit. xii, pp. 893-898.
- 797. Schäff, —. Ueber Insektenreste aus dem Torflager von Klinge, 8B. Ges. Naturf. Berl. 1892, pp. 8-11.
  - 1892. [vol. xxix.]

- 798. SCHAUS, W., JUN. American Lepidoptera; illustrations of new and rare species. Part 1, pp. 1-24, pls. i-iii. London: 1892, 4to.
- 799. Descriptions of new species of Lepidoptera Heterocera from Brazil, Mexico, and Peru. P. Z. S. 1892. Part I, pp. 272-291; Part II, pp. 318-341.
  - Many of the species are figured in the preceding (798).
- 800. SCHLECHTENDAL, D. von. Ueber das Vorkommen fossilen "Ruckenschwimmer" (Notonecten) im Braunkohlengebirge von Rott. Z. Naturw. lxv, pp. 141-143. [Rhynchota.]
- 801. SCHMIDT, J. Neue Histeriden. Ent. Nachr. xviii, pp. 17-29. [Coleoptera.]
- 802. Voyage de M. Ch. Alluaud dans le territoire d'Assinie (Afrique occidentale). Histeridæ. Ann. Soc. Ent. Fr. lxi, pp. 289 291. [Coleoptera.]
  - Schmiedcknecht, O. [See Gräffe (358).]
- 803. SCHULTESS-RECHBERG, A. v. Tiphia picta spec. nov. aus Bulgarien. MT. Schw. ent. Ges. viii, p. 384. [Hymenoptera.]
- 804. Schwarz, E. A. Entomological Club of American Association for the Advancement of Science—Annual Address of the President. Canad. Ent. xxiv, pp. 213-224.
- A consideration of the present position of Coleopterology in N. America.
- 805. —. Note on Phytobius. P. Ent. Soc. Wash. ii, p. 165. [Coleoptera.]
- 806. —... Note on the Food-habits of some Halticids. T. c. pp. 182-184. [Coleoptera.]
- 807. —. Verdigris in Coleoptera. T. c. pp. 220-222.
- 808. —. Time of Flight in Lachnosterna. T. c. pp. 241-244. [Coleoptera.]
- SCHWARZ, O. Revision der paläarktischen Arten der Elateriden-Gattung Melanotus, Esch. Deutsche e. Z. 1892, pp. 145-164.
   [Coleoptera.]
- 810. —. Neue Cardiophorus-Arten. T. c. pp. 383 & 384. [Coleoptera.]
- 811. —... Spheniscosomus, eine neue, bisher verkannte Elateriden-Gattung. Wien. ent. Z. xi, p. 132. [Coleoptera.]
- 812. Scudder, S. H. The songs of our Grasshoppers and Crickets. Rep. E. Soc. Ont. xxiii, pp. 62-78. [Orthoptera.]
- 813. —. New light on the formation of the abdominal pouch in *Parnassius*. Tr. Ent. Soc. 1892, pp. 249-253. [Lepidoptera.]
- 814. —. The Tertiary Rhynchophora in North America. P. Bost. Soc. xxv, pp. 370-386. [Coleoptera.]

- 815. [SCUDDER, S. H.] The services of Edward Burgess to Natural Science. P. Bost. Soc. xxv, pp. 358-364.
- 816. . The Orthopteran genus Hippiscus. Psyche, vi, pp. 265-274, 285-288, 301-305, 317-320, 333-336, 347-350, & 359-363.
- 817. SEITZ, A. Reiseskizzen, I. Singapore. S. E. Z. liii, pp. 52-61. [Lepidoptera.]
- 818. —... Reiseskizzen, II. Lantana. T.c. pp. 228-238. [Lepidoptera.]
  Concerns the attractiveness of Lantana bush to insects in the East.
- Mittheilungen über Mimicry. Verh. deutsch. Zool. Ges. i, 1891, pp. 49-53. [Lepidoptera.]
- 820. —... Mittheilungen von einer Reise nach China und Japan. Op. cit. 1892, pp. 14-18.
- 821. —... Eine zoologische Excursion in die Umgegend von Shanghai. JB. nass. Ver. xlv, pp. 41-48.
- 822. —... Eine entomologische Excursion in die Umgebung von Hiogo. T. c. pp. 51-61.
- Ein Ausflug in die Umgebung von Nagasaki. Ent. Nachr. xviii, pp. 49-54.
- 824. SELYS-LONGCHAMPS, E. DE. Viaggio di Leonardo Fea in Birmania e regioni vicine. XXXII. Odonates. Ann. Mus. Genov. (2) x, pp. 433-518. [Neuroptera.]
- 825. Causeries Odonatologiques. No. 6, Les Gomphines d'Afrique. Ann. Ent. Belg. xxxvi, pp. 86-107. [Neuroptera.]
- 826. Apparition accidentelle de la *Mantis religiosa* et de quelques autres insectes en Belgique. T. c. pp. 495-503.
- 827. Semenow, A. De genre *Fseudochrysis* m. Hor. Ent. Ross. xxvi, pp. 480-491. [Hymenoptera.]
- 828. Revisio Hymenopterorum Musei Zoologici Academiæ Cossareæ scientiarum Petropolitanæ. 1. Genus Cleptes, Latr. Bull. Petersb. (n.s.) ii, pp. 497-504. 11. Genus Abia, Leach; t. c. pp. 509-517. 111. Evaniidæ; op. cit. iii, pp. 9-30.
- 829. —... Chrysididarum species novæ. Bull. Petersb. (n.s.) iii, pp. 71-95. [Hymenoptera.]
- 830. —... Note sur la subdivision du genre Lethrus, Scop., et description de deux espèces nouvelles. Hor. Eut. Ross. xxvi, pp. 232-244. [Coleoptera.]
- 831. De Brenthidarum genere novo palmarctico. T. c. pp. 438-443. [Coleoptera.]
- 832. Conspectus specierum generis Glaresis, Er. T.c. pp. 468-470. [Coleoptera.]

- 833. [Semenow, A.] Notes breves de quibusdam Melolonthides. Hor. Ent. Ross. xxvi, pp. 471-476. [Coleoptera.]
- 834. —. Generis Glaphyrus, Latr., species nova rossica. T. c. pp. 477-479. [Coleoptera.]
- 835. Semper, G. Reisen im Archipel der Philippinen von Dr. C. Semper. Zweiter Theil. Wissenschaftliche Resultate. Fünfter Band. Die Schmetterlinge der Philippinischen Inseln. Beitrag zur Indo-Malayischen Lepidopteren-fauna. Erste Abtheilung, Die Tagfalter. Rhopalocera, part VIII, pp. 271-380, pls. xlvii-xlix.

This part completes the volume, which has a second title page as follows: Die Schmetterlinge der Philippinischen Inseln: Beitrag zur Indo-Malayischen Lepidopteren-fauna von Georg Semper. Erster Band: Die Tagfalter. Rhopalocera.

- 836. SÉNAC, H. Note sur les *Pimelia* des îles Canaries. Ann. Soc. Ent. Fr. lxi, pp. 103-108. [Coleoptera.]
- 837. Senna, A. Contributions to the knowledge of the family Brenthidæ. VIII. Enumeration of the species known as yet from Java. Notes Leyd. Mus. xiv, pp. 161-186. [Coleoptera.]
- 838. Contribuzioni allo studio dei *Brentidi*. Nota ix, Bull. Ent. Ital. xxiv, pp. 26-37. Nota x, t. c. pp. 38-63. Nota xii, t. c. pp. 152-163. [Coleoptera.]
- 839. —... Viaggio di Leonarda Fea in Birmania e regioni vicine. XLV. Brentidi. Ann. Mus. Genov. (2) xii, pp. 429-494. [Coleoptera.]
- 840. SERGI, G. Ricerche su alcuni organi di senso nelle antenne delle Formiche. Bull. Ent. Ital. xxiv, pp. 18-25. [Hymenoptera.] Summary in J. R. Micr. Soc. 1892, p. 780.
- 841. Severin, G. Les collections d'Articulés du Musée royal d'histoire naturelle de Belgique. Aun. Ent. Belg. xxxvi, pp. 469-478. [Coleoptera.]
- 842. Sharp, D. On some Eggs of *Hemiptera*. Tr. E. Soc. 1892, pp. 191-199, pls. viii & ix.
- 843. Descriptions of two new Pselaphida found by Mr. J. J. Walker in Australia and China. Ent. M. M. xxviii, pp. 240-242. [Coleoptera.]
- 844. —... A new species of the Staphylinid genus Trygaus. Notes Leyd. Mus. xiv, p. 61. [Coleoptera.]
- 845. Entomological Pins. Ent. xxv, pp. 54-57 & 119. [See also Brunner & Redtenbacher (138).]
- 846. SICARD, A. J. Weise: Bestimmungs-Tabellen der europäischen Coleopteren: fasc. II, 2º edition. Traduction complète, due en grande partie a M. le Dr. A. Sicard, médecin aide-major de 1º classe, et accompagnée de diverses notes inédites, signées chacune du nom de leur auteur. Abeille, xxviii, pp. 1-80 (still incomplete). [Coleoptera.]

- 847. SKINNER, H. A new species of Pamphila. Ent. News, iii, p. 174.
  [Lepidoptera.]
- 848. —. A new Pamphila. T. c. p. 222. [Lepidoptera.]
- 849. —. A new species of Eudamus. Canad. Ent. xxiv, p. 164. [Lepidoptera.]
- SKINNER, H., & MENGEL, L. W. Greenland Lepidoptera. P. Ac. Philad. 1892, pp. 156-159.
- SLINGERLAND, M. V. The Pearl-tree Psylla. Bull. Cornell Univ. zliv, pp. 161-186. [Rhynchota.]
- 852. SLOANE, T. G. Descriptions of two new species of Carenum from West Australia, with notes of the synonymy and distribution of some previously described species. P. Linn. Soc. N.S.W. (2) vi, pp. 427-429. [Coleoptera.]
- 853. —. Studies in Australian Entomology. No. v. Notes on the subfamily *Broscini* (*Carabida*), with descriptions of new species. *Op. cit.* vii, pp. 45-64. [*Coleoptera.*]
- 854. SLOSSON, A. T. A new Arctia. Ent. News, iii, p. 257. [Lepidoptera.]
- 855. A new Dasylophia from Florida. Canad. Ent. xxiv, p. 139. [Lepidoptera.]
- 856. SMIRENSKY, A. Ueber schädliche Insekten der Umgebung von Kazan. Trudui Kazan. Univ. xxiii, No. 4, pp. 1-18.
  In the Russian language.
- 857. SMITH, H. G. Descriptions of three new species of Butterflies captured by Mr. D. Cator in British North Borneo, in the Collection of Mr. Grose Smith. Ann. N. H. (6) x, pp. 426-428. [Lepidoptera.]
- 858. SMITH, H. G., & KIRBY, W. F. Rhopalocera exotica. Pts. 19-22.

  Vol. I was completed by Part 20. It is proposed to continue the work for several vols.
- 859. SMITH, J. B. List of Lepidoptera of Boreal America. American Entomological Society, 1891, 8vo, 124 pp.
  - A list of 6020 names. For corrections, &c., see DYAR, Canad. Ent. xxiv, pp. 47 & 48, and SMITH, t. c. p. 103.
- 860. —... Synonymical notes on some *Heterocera* in the British Museum. Canad. Ent. xxiv, pp. 133-136. [Lepidoptera.]
- 861. —. New species of Noctuidæ. Ent. News, iii, pp. 250-253, pl. x. [Lepidoptera.]
- 862. —. Note on generic characters in the Noctuidæ. Ent. xxv, pp. 101-105. [Lepidoptera.]
- 863. —. Mr. Butler's Notes on Synonymy of North American Noctuida. T. c. p. 2. [Lepidoptera.]

- 864. [SMITH, J. B.] The Mouth of Copris carolina and Notes on the Homology of the Mandible. P. Am. Ass. xl, p. 324.
- 865. —. The Mouth Parts of Copris carolina, with Notes on the Homologies of the Mandibles. Tr. Am. Ent. Soc. xix, pp. 83-87, pls. ii & iii. [Coleoptera.]
- 866. —. Notes on the Homology of the Hemipterous Mouth. P. Am. Ass. xl, p. 325.
- 867. —... Contributions towards a Monograph of the Noctuide of Boreal America. Revision of the genus Cucullia; revision of the Dicopina; revision of Xylomiges and Morrisonia. P. U. S. Nat. Mus. xv, pp. 33-86, pls. ii & iii. [Lepidoptera.]
- 868. SMITH, T. J. Remarks on Apis mellifica, the Hive Bee. Rep. North Staffordsh. Club, xxvi, pp. 82-91.
- 869. SNELLEN, P. C. T. Aanteekeningen over eene verzameling Lepidoptera in October 1889, van het eiland Flores ontvangen. Tijdschr. Ent. xxxiv, pp. 229-256.
- 870. —. Deudoryx verriculata, n. sp. T. c. pp. 257-260.
- 871. Euplæa perdita, Butl. var. T. c. p. 339. [Lepidoptera.]
- Aanteekeningen over Lepidoptera schadelijk voor het suikerriet. T. c. pp. 341-356, pls. xviii & xix.

The species described and figured in this memoir appear to have been previously dealt with in an inaccessible work as follows:—

- 873. ———— in Mededeelingen van het Proefstation voor Suikerriet in West Java; 1890, pp. 94, &c., pls. i & ii. [Lepidoptera.]
- 874. —... Aanteekeningen over Exotische Lepidoptera. Tijdschr. Ent. xxxv, pp. 1-23, pl. i.
- 875. Boekaankondiging. T. c. pp. 73-84. [Lepidoptera.]

  A review of the second part of Saalmüller's work on the Lepidoptera of

  Madagascar, and includes many synonyms and critical remarks.
- 876. Beschrijving van eenige nieuwe Javaansche dagvlinders. T. c. pp. 133-144, pls. iii & iv (incomplete). [Lepidoptera.]
- Snow, W. A. Notes and descriptions of Syrphidæ. Kansas Quart. i, pp. 33-38, pl. vii. [Diptera.]
- 878. Sonthonnax, L. Description d'une Saturnie nouvelle de la Chine. Échange, 1892, p. 20. [Lepidoptera.]
- 879. SOUTH, R. Notes on British Lepidoptera. Ent. xxv, pp. 29-36, 86-93, & 134-138.

These notes relate to variation in several species.

880. —... The Entomological Club of London. T. c. pp. 4-9.

881. Spuler, A. Zur Phylogenie und Ontogenie des Flügelgeäders der Schmetterlinge. Z. wiss. Zool. liii, pp. 597-646, pls. xxv & xxvi. [Lepidoptera.]

The neuration in the perfect insects of numerous forms, especially of *Tineida*, is discussed, in addition to the phylogeny and to the condition in the pupal state. Summary in J. R. Micr. Soc. 1892, p. 469, and by Seitz, S. E. Z. liii, pp. 238-241.

- 882. Zur Stammesgeschichte der Papilioniden. Zool. Jahrb. Syst. vi, pp. 465–498, pls. xxii & xxiii. [Lepidoptera.]
- 883, STADELMANN, H. Neue Hymenopteren der Zool. Sammlung des Kgl. Museums für Naturkunde zu Berlin. Berl. Ent. Zeit. xxxvii, pp. 235-240.
- 884. STANDFUSS, M. Ueber die Hybridation bei den Insekten. MT. Schw. ent. Ges. viii, pp. 386-396.
  - —. [See also Romanoff (779).]
- 885. STAUDINGER, O. Neue afrikanische Lycaniden. Deutsche e. Z. Lep. iv, pp. 215-223. [Lepidoptera.]
- Neue Arten und Varietäten von Lepidopteren des paläarktischen Faunengebiets. T. c. pp. 224-339.
- 887. —... Neue Arten und Varietäten von paläarktischen Geometriden.

  Op. cit. v, pp. 141-260. [Lepidoptera.]
- 888. —. Zwei neue Charaxes-Arten. T. c. pp. 260-263. [Lepidoptera.]
- 889. . Ueber die Arten der Palla Decius-Gruppe. T. c. pp. 264-268. [Lepidoptera.]
- 890. —. Lepidopteren aus Tunis. T. c. pp. 277-300. Includes some descriptions by RAGONOT.
- 891. —. Lepidopteren des Kentei-Gebirges. T. c. pp. 300-393.
- 892. —. Ornithoptera andromache, n. sp. T. c. p. 393. [Lepidoptera.]
- 893. —. Neue Papilio-Formen aus Südamerika. T. c. pp. 427 & 428. [Lepidoptera.]
- 894. —. Einige neue Rhopaloceren aus Nord-Borneo. T. c. pp. 449-452. [Lepidoptera.]
  - ---. [See also ROMANOFF (779).]
- 895. STEIN, P. Die Anthomyiden der Fallen-Zetterstedt'schen Sammlung. Ent. Nachr. xviii, pp. 321-333. [Diptera.]
- Drei neue merkwürdige Homalomyia-Arten. Wien. ent. Z. zi, pp. 69-77. [Diptera.]
- 897. STEINERT, H. Ueber das Auftreten von Amphidasys betularius, L., ab. doubledayarius, B., in Sachsen, mit Bemerkungen über die Zucht des Thieres. Deutsche e. Z. Lep. v, pp. 424-427. [Lepidoptera.]

- 898. STIERLIN, —. Due nuovi Curculionidi di Sicilia. Nat. Sicil. xi, p. 156. [Coleoptera.]
- 899. ——. Beschreibung einiger neuen Rüsselkäfer. MT. Schw. ent. Ges. viii, pp. 357-366. [Coleoptera.]
- Beschreibung einiger neuen europ. Rüsselkäfer. T. c. pp. 408-412. [Coleoptera.]
- 901. STILES, C. W. On the American intermediate host of Echinorhynchus gigas. Zool. Anz. xv, pp. 52-54. [Coleoptera.] Lamellicorn beetles of the genus Lachnosterna.
- 902. STROBL, G. Die österreichischen Arten der Gattung Hilara, Meig. (Mit Berücksichtigung der Arten Deutschlands und der Schweiz.) Verh. z.-b. Wien, zlii, pp. 85–182. [Diptera.]
- 903. —... Interessante österreichische *Dolichopoden*. Wien. ent. Z. xi, pp. 102-107. [*Diptera*.]
- Untersuchungen über die Gattung Spania, Meig. T. c. pp. 121-125. [Diptera.]
- 905. —. Zwei alpine Sapromyzen. T. c. pp. 155 & 156. [Diptera.]
- Zur Kenntniss und Verbreitung der Phoriden Oesterreichs.
   T. c. pp. 193-204. [Diptera.]
- 907. STUMMER-TRAUNFELS, R. R. v. Vergleichende Untersuchungen über die Mundwerkzeuge der *Thysanuren* und *Collembolen*. SB. Ak. Wien. c, Abth. i, pp. 216–234, pls. i & ii.
  - Summary in J. R. Micr. Soc. 1892, p. 781.
- 908. Swinhoe, C. Catalogue of Eastern and Australian Lepidoptera Heterocera in the collection of the Oxford University Museum. Part 1. Sphinges and Bombyces. Oxford: viii & 324 pp., 8 pls.
- 909. —. New species of *Heterocera* from the Khasia Hills. Part II. Tr. E. Soc. 1892, pp. 1-20, pl. i.
- 910. —. Mimicry in Nature. P. Croydon Club, iii, pp. 332-341.
- 911. SWIERSTRA, K. N. Een hermaphrodiet van Aspilates strigillaria.
  Tijdschr. Ent. xxxiv, p. 338, pl. xvii, fig. 3. [Lepidoptera.]
- 912. Targioni-Tozzetti, A. Aonidia blanchardi, specie nuova di Cocciniglia delle palme da dattero del Sahara. Bull. Ent. Ital. xxiv, pp. 170-188. [Coccidæ.]
- 913. —... Aonidia blanchardi, nouvelle espèce de Cochenille du dattier du Sahara. Mem. Soc. Zool. v, pp. 69-82. [Coccidæ.]
- 914. TARRAL, H. Observations sur les oothèques du Stauronotus maroccanus. Bull. Soc. Béziers, xii, pp. 10-13. [Orthoptera.]
- 915. Taschenberg, O. Historische Entwickelung der Lehre von der Parthenogenesis. Abh. Ges. Halle, xvii, pp. 365-453.

- 916. TEPPER, J. G. O. The *Phaneropterida* of Australia and Polynesia. Tr. R. Soc. S. Austr. xv, pp. 77-113. [Orthoptera.]
- 917. —. The Gryllacridæ and Stenopelmatidæ of Australia and Polynesia. T. c. pp. 137-178. [Orthoptera.]
- 918. Tetens, H. Resultate der anatomischen Untersuchung eines lateralen Zwitters von Smerinthus populi. Berl. Ent. Zeit. xxxvi, pp. 457-466, pl. xiii. [Lepidoptera.]
- 919. THEOBALD, F. V. An account of British Flies (Diptera.). Vol. I. London: 1892, 8vo, xix & 215 pp. Parts I-III were recorded last year.
- 920. Stylopized Bees. Ent. M. M. xxxviii, pp. 40-42.
- 921. THÉRY, A. Description de Coléoptères. Bull. Soc. Ent. Fr. lxi, p. xcvi.
- 922. Description d'un Coléoptère nouveau. T. c. p. cclviii.
- 923. THIERRY-MIEG, P. Descriptions de Lépidoptères nocturnes. Le Nat. 1892, pp. 216, 217, 235, 236, & 262.
- 924. THOMAS, F. Alpine Mückengallen. Verh. z.-b. Wien, xlii, pp. 356-376, pls. vi & vii. [Diptera.]
- 925. THOMAS, F. A. W. Beobachtungen über Mückengallen. Sep. from Wiss. Beilage z. Programme Gym. Gleichense Ohrdruf. pp. 1-16.
  - Account of Cecidomyiid galls, arranged according to plants.
- 926. THOMAS, FR. Die Blattflohkrankheit der Lorbeerbaume. Gartenflora, xl, Heft. 2 & 3. [Rhynchota.]

  Cf. Wien. ent. Z. xi, p. 87.
- 927. THOMSON, C. G. Opuscula entomologia. Fasciculus xvimus. [Hymenoptera.]

Relates to Braconidæ, and to some Ichneumonidæ allied to Anomalon.

- 928. Thomson, J. P. British New Guinea. London: 8vo.
  Pp. 222-272 are occupied by a sketch of the present state of our know-ledge of the entomology of the island.
- 929. TÖMÖSVÁRY, D. E. Golubačkha muha. Glasnik naravosl. Drust. vi, pp. 187–203, pl. ii. [Diptera.]
- 930. TOURNEUX, F. Sur les modifications structurales que présentent les muscles jaunes du Dytique pendant la contraction. J. de l'Anat. Phys. xxviii, pp. 573-581, pl. xxi.
- 931. Townsend, C. H. T. New North American Tachinida. Ent. News, iii, pp. 80, 81, & 129-131. [Diptera.]
- 932. —. New Jamaica Tachinidæ. T. c. p. 146.

- 933. [Townsend, C. H. T.] Notes on North American Tachinide, sens. str., with descriptions of new genera and species. Tr. Am. Ent. Soc. xix, pp. 88-132. [Diptera.]
- 934. —... The North American genera of Calyptrate Muscidæ. Paper 11.
  T. c. pp. 133-144. [Diptera.]
- 935. The North American genera of Nemocerous Diptera. T. c. pp. 144-160.
- 936. —... The North American genera of Calyptrate *Muscidie*. Paper 111, t. c. pp. 273-278. Paper IV, t. c. pp. 279-284. Paper V, t. c. pp. 290-294. [*Diptera*.]
- 937. —... Notes on North American Tachinida, with descriptions of new species. Paper VII, t. c. pp. 284-289. [Diptera.]
- 938. —... Notes on North American Tachinidæ, with descriptions of new genera and species. Canad. Ent. xxiv. Paper v, pp. 64-70 & 77-82. Paper vi, pp. 165-172. [Diptera.]
- 939. A Sarcophagid parasite of Cimbex americana. T. c. p. 126. [Diptera.]
- 940. A preliminary grouping of the described species of Sapromyza of North America, with one new species. T. c. pp. 301-304. [Diptera.]
- 941. —. Biologic notes on New Mexico Insects. T. c. pp. 193-200.
- 942. —. Description of a Sarcophaga bred from Helix. Psyche, vi, p. 220. [Diptera.]
- 943. —. An Aporia bred from Limacodes sp. T. c. p. 275. [Diptera.]
- 944. —. Descriptions of Æstrid larvæ taken from the Jack-rabbit and Cotton-tail. T. c. pp. 298-300. [Diptera.]
- 945. —. Oviposition of a *Homopterous* Insect in Yucca. T. c. p. 353. [Rhynchota.]
- 946. A Tachinid bred from larva of Protoparce jamaicensis. J. Inst. Jamaica, i, p. 70. [Diptera.]
- 947. A Dexiid parasite of a Longicorn Beetle. T. c. pp. 105 & 106. [Diptera.]
- 948. Insects frequenting Yucca blooms. Zoe, iii, pp. 113-115.
- 949. Notes on two Mexican species of Ceroplastes, with a record of parasites reared from one. T. c. pp. 255-257. [Coccidæ, Hymenoptera.]
- 950. TRAIL, J. W. H. The Gall-making Hymenoptera of Scotland (exclusive of those that live on oaks). Tr. & P. Perthsh. Soc. i, pp. 72-90.
  - Includes a list of plants that bear Hymenopterous galls.

- TRYBOM, F. Mermis-larver hos Chironomus. Ent. Tidskr. xiii, pp. 81-92. [Diptera.]
- 952. TRYON, H. Zoology of British New Guinea. Part II. Hemiptera. Ann. Queensland Mus. ii, pp. 13-24.
- 953. —... Colsopters collected by Mr. A. C. English in the St. Joseph river district of British New Guines, under the auspices of his Honour the Administrator. Rep. Administrator Brit. New Guines, ii, app. v, pp. 109-112.

Recorded from a separate copy, paged 1-8.

954. —... Rhopalocera collected in British New Guinea during years 1889-90, under the auspices of his Honour the Administrator. T. c. pp. 112-115. Brisbane, 1890.

Recorded from a separate copy, 8vo, pp. 1-7.

- 955. Turr, J. W. British Noctuce and their varieties. Vol. 11. London: 1892
- 956. URECH, F. Ueber einen grünen Farbstoff in den Flügelchen (nicht in den Schuppen) der Chrysalide von Pieris brassicæ. Zool. Anz. xv, pp. 281-283. [Lepidoptera.]

Summary in J. R. Micr. Soc. 1892, p. 778.

- 957. Beobachtungen über die zeitliche Succession des Auftretens der Farbenfelder auf den Puppenflügelchen von Pieris brassica. T. c. pp. 284-290 & 293-299. [Lepidoptera.]
- 958. . Ueber Eigenschaften der Schuppenpigmente einiger Lepidopteren-species. T. c. pp. 299-306.

Summary in J. R. Micr. Soc. 1892, p. 778.

- 959. VACHAL, J. Halicti, novæ species. Bull. Soc. Ent. Fr. lxi, p. xxii. [Hymenoptera.]
- 960. Hymenoptères récoltés au Soudan. T. c. pp. cxxxv-cxxxvii.
- 961. Verhoeff, C. Zur Kenntniss des biologischen Verhältnisses zwischen Wirth- und Parasiten-Bienenlarven. Zool. Anz. xv, pp. 41-43. [Hymenoptera.]

Summary in J. R. Micr. Soc. 1892, p. 358.

- 962. Die physiologische Bedeutung des Stachelapparates besonders der *Hymenopteren*-Nymphen. T. c. pp. 355-360.
- 963. —... Neue und wenig bekannte Gesetze aus der Hymenopteren-Biologie. T. c. pp. 362-370.

Relates to the numbers and positions of the sexes in their pupal cells, and to the proportions of males to females in certain species of Fossoria.

Summary in J. R. Micr. Soc. 1893, p. 34.

964. — Beiträge zur Biologie der Hymenoptera. Zool. Jahrb. Syst. vi, pp. 680-754, pls. 30 & 31.

Includes discussion on phylogeny of Social Hymenoptera.

- [Verhoeff, C.] Weitere Untersuchungen über den Ausfärbungsprocess. Ent. Nachr. xviii, pp. 54-58.
- 966. —. Ueber einige neue und seltene Fossorien. T. c. pp. 65-72. [Hymenoptera.]
- 967. —... Einige Worte zu H. Friese's Osmien-Studien und über einen Bau von Osmia tridentata. T. c. pp. 225-228. [Hymenoptera.]
- 968. —. Ueber Kämpfende und gesellige Bienenmännchen. T. c. pp. 244-247. [Hymenoptera.]
- 969. —. Eine neue Osmia aus Dalmatien. T. c. pp. 289-291. [Hymenoptera.]
- 970. Zur Entwickelung von Hemerobius subnebulosus und über Verfärbung der Neuropteren. T. c. pp. 279 & 298.
- 971. VERRALL, G. H. Two new English species of Homalomyia. Ent. M. M. xxviii, p. 149. [Diptera.]
- 972. Verson, E. Postlarvale Neubildung von Zelldrüsen beim Seidenspinner. Zool. Anz. xv, p. 216. [Lepidoptera.]
- 973. —... Besitzen die Bauchfüsse der Seidenraupe Haftpapillen? T. c. pp. 279 & 281. [Lepidoptera.] Summary in J. R. Micr. Soc. 1892, p. 780.
- 974. —... Altre cellule glandulari di origine postlarvale. Bull. Ent. Ital. xxiv, pp. 3-17.
  Summary in J. R. Micr. Soc. 1892, p. 779.
- 975. VIALLANES, H. Recherches anatomiques et physiologiques sur l'œil composé des arthropodes. Ann. Sci. Nat. (7) xiii, pp. 349-382, pls. x & xi.
- 976. VILLARD, L. Description d'un Cérambycide nouveau du Japon. Bull. Soc. Ent. Fr. lxi, p. li. [Coleoptera.]
- 977. VONHOF, O. Bienenmass, oder: die Descendenz lehre ist ein falscher Schluss! Auszug aus einer Studie. Bremen: 1891.

  Cf. Ent. Nachr. xviii, p. 237.
- 978. Vuillot, P. Descriptions de Lépidoptères. Bull. Soc. Ent. Fr. lxi, pp. cxl & cxli.
- 979. —. Description d'un Lépidoptère nouveau. T. c. p. cxlviii.
- 980. Descriptions de six Lépidoptères nouveaux. T. c. pp. cxc-cxciv.
  - ---. [See also Mabille & Vuillot (573).]
- WAGNER, F. v. Zur biologie der wilden Bienen. Biol. Centralbl. xii, pp. 572-584. [Hymenoptera.]
   Relates chiefly to Friese's views.

- 982. WARBURG, O. Ueber Ameisenpflanzen. Biol. Centralbl. xii, pp. 129-142.
- An interesting discussion on the symbiosis of plants and ants. Summary in J. R. Micr. Soc. 1892, p. 353.
- 983. WARREN, W. Descriptions of new genera and species of *Pyralidæ*, contained in the British Museum Collection. Ann. N. H. (6) ix, pp. 172-179, 294-302, 389-397, & 429-442, continued from Vol. viii, p. 70. [Lepidoptera.]
- 984. WASMANN, E. Einiges über springende Ameisen. Wien. ent. Z. xi, pp. 316 & 317. [Hymenoptera.]
- Ein neuer Paussus vom Somaliland. MT. Schw. ent. Ges. viii, pp. 355-357. [Coleoptera.]
- 986. —. Zur Biologie einiger Ameisengaste. Deutsche e. Z. 1892, pp. 347-351. [Coleoptera.]
- 987. —. Die internationalen Beziehungen von Lomechusa strumosa. Biol. Centralbl. xii, pp. 584-599 & 633-669.
  - Summary in J. R. Micr. Soc. 1892, p. 780, and 1893, p. 34.
- 988. WATERHOUSE, C. O. Two new Buprestidæ from Damma Island. Ann. N. H. (6) x, pp. 410-412. [Coleoptera.]
- 989. Weber, L. Über die sog. Triungulinus form der Meloëlarven. Ber. Ver. Cassel, 1891-92, Abh. pp. 1-5, pl. [Coleoptera.]
- 990. WEBSTER, F. M. Report of Entomological work of the season of 1891. Bull. Dep. Agric. Ent. xxvi, pp. 63-74.
- 991. —. Food-plants of the Lixi. P. E. Soc. Wash. ii, p. 339. [Coleoptera.]
- 992. WEISE, J. Zwei neue Cassida-Arten. Wien. ent. Z. xi, p. 238. [Coleoptera.]
- 993. —. Kleine Beiträge zur Coccinelliden-Fauna Ost-Afrika's. Deutsche e. Z. 1892, pp. 15 & 16. [Coleoptera.]
- 994. —... Beitrag zur Coleopteren-Fauna von Turkestan. T. c. pp. 131-140.
- 995. —... Coccinelliden aus Osteibirien. T. c. pp. 140 & 141. [Coleoptera.].
- 996. —... Pachygnatha, eine neue Chrysomelen-Gattung. T. c. p. 352. [Ooleoptera.]
- 997. —... Chrysomeliden und Coccinelliden von der Insel Nias, nebst Bemerkungen über andere, meistens Südostasiatische Arten. T. c. pp. 385-400. [Coleoptera.]
- 998. Beschreibung einiger Galeruciden. T. c. pp. 403-405. [Coleoptera.]

- 999. [Weise, J.] Chrysomelen aus dem hohen Caucasus. Deutsche e. Z. 1892, pp. 405 & 406. [Coleoptera.]
- 1000. —. Malthodes brevicollis, Payk. Typ einer Gattung. T. c. pp. 407-410. [Coleoptera.]
- 1001. —. Les Coccinellides du Chota-Nagpore. Ann. Ent. Belg. xxxvi, pp. 16-30. [Coleoptera.]
  - —. [See also HEYDEN (402) and SICARD (846).]
- 1002. WERNER, F. Selbstverstümmelung bei Heuschrecken. Zool. Anz. xv, pp. 58-60. [Orthoptera.]
- 1003. WEYMER, G. Exotische Lepidopteren. vi. Aus dem afrikanischen Faunengebiet. S. E. Z. liii, pp. 79-125.
- 1004. WHEELER, W. M. Concerning the "blood-tissue" of the *Insecta*. Psyche, vi, pp. 216-220, 233-236, & 253-258, pl. vii.
- 1005. WICKHAM, H. F. On the early stages of three North American Coleoptera. Bull. Lab. Iowa, ii, pp. 195-201.
- 1006. —. Collecting in the far North. I. The Alaskan Coast. Ent. News, iii, pp. 139-144. II. The Stikine River. T. c. pp. 256 & 257.
- 1007. —. Report on an Entomological reconnoissance of Southern Alaska and adjacent portions of British Columbia. Bull. Lab. Iowa, ii, pp. 202-233.
- 1008. Description of four Insect monstrosities. Canad. Ent. xxiv, pp. 141 & 142. [Coleoptera.]
- 1009. —. Notes on some Myrmecophilous Coleoptera. Psyche, vi, pp. 321-323.
- 1010. WILLISTON, S. W. Diptera Brasiliana. Kansas Quart. i, pp. 44-46.
- 1011. —. A new species of Criorhinia, and notes on synonymy. Ent. News, iii, p. 145. [Diptera.]
- 1012. —. Note on the habits of Ammophila. T. c. pp. 85 & 86. [Hymenoptera.]
  - ---. [See also GODMAN & SALVIN (347).]
- 1013. Winn, A. F. The Horn Fly. Can. Rec. v, pp. 272-275.
  [Diptera.]
  - First appearance in Canada.
- 1014. Wood, J. H. Our Rush-feeding Coleophoræ. Ent. M. M. xxviii, pp. 117-122, & 169-176. [Lepidoptera.]
- 1015. —. Further notes on our Rush-feeding Coleophoræ. T. c pp. 282-285. [Lepidoptera.]

- 1016. WRIGHT, W. G. Getting Butterfly Eggs. Canad. Ent. xxiv, pp. 70-74. [Lepidoptera.]
- 1017. —. Description of a new Anthocharie. T. c. p. 154. [Lepidoptera.]
- 1018. WROUGHTON, R. C. Our Ants. Part I. J. Bomb. N. H. Soc. vii, pp. 13-60, pls. A, B. Part II. T. c. pp. 175-202, pls. c, D. [Hymenoptera.]
- 1019. WULP, F. M. VAN DER. Einige uitlandische Diptera. Tijdschr. Ent. xxxiv, p. 193, pl. xii.
- 1020. XAMBEU, —. Description de quatre Coléoptères nouveaux et particuliers à notre département. Soc. Agricole Pyrén.-orient. xxxiii, pp. 241-243.
  - The descriptions have appeared some years since elsewhere.
- 1021. —. Mœurs et métamorphoses d'insectes. Ann. Soc. L. Lyon (n.s.), xxxix, pp. 135-194. [Coleoptera.]
- 1022. Mœurs et métamorphoses d'insectes. Soc. Agricole Pyrén.orient. xxxii, pp. 473-484. [Coleoptera.]
- 1023. —. Mœurs et métamorphoses d'insectes. 2<sup>me</sup> Mémoire.
   Coléoptères. Pp. 1-46. Lyon: 1892, 8vo.
   Published as supplement to Échange.
- 1024. Mosurs et métamorphoses d'insectes. 3° Mémoire. Buprestides. Rev. d'Ent. xi, pp. 202-252. [Coleoptera.]
- 1025. YERBURY, J. W. The Butterflies of Aden and neighbourhood. J. Bomb. N. H. Soc. vii, pp. 207-218. [Lepidoptera.]
- 1026. Zoufal, V. Uebersicht der mir bekannten Arten der Coleopteren-Gattung, Tenebrio, L. Wien. ent. Z. xi, pp. 283-286.

## II.—ANATOMY, PHYSIOLOGY, BIOLOGY.

## 1. General, Comparative, and External.

General anatomy of Phylloxera, Krassilstschik (519).—Anatomy of Coccus cacti, and production of wax, Mayer (594).—General anatomy and histology of the Blow-fly, Lowne (567).—The structure of male appendages, Eschebich (257).—Blood-tissue, various points discussed, Wheeler (1004).—Metameric glandular processes in larva of Lagoa, Packard (666).—Mouth-parts of Copris, and homology of mandibles, Smith (864, 865).—Anatomy of mouth of Diptera Pupipara, Müggenburg (628).—Comparative anatomy of trophi of Thysanura and Collembola, Stummer-Traunfels (907).—Glands in pulvilli of Diptera, Jenkinson (456).—Poison glands in Ichneumonida, Buysson (148).—Tegulæ and patagia of Lepidoptera, Riley, P. E. Soc. Wash. ii, p. 310.

Mechanism of suction in *Hymenoptera* and other insects, Janet, p. xcvii, Gazagnaire, p. xcviii, Bull. Soc. Ent. Fr. lxi.

Histology of metamorphosis of Lepidoptera, Bugnion (140).—Histology of elytra of Coleoptera, HOFFBAUER (405).

Physiological nature (Asphyxia) of metamorphosis in the silkworm, BATAILLON (35).—Mode of development of the chitinous envelope in larvæ of Libellulidæ, CHATIN (165).—The process of development of the epiderm in larvæ of Libellulidæ, CHATIN (166).—Post larval formation of cellular glands in Bombyx mori, VERSON (972).—Glandular epigastric cells of postlarval origin in silkworm, VERSON (974).—Survival of freezing, LYMAN (570).

#### 2. NERVOUS SYSTEM, ORGANS OF SENSE, AND PSYCHOLOGY.

Nervous system of *Pentaplatarthus*, Raffeay (709).—Composition of the nerve-tissue in insects, Griffiths, C.R. cxv, p. 563.—Structure of the nerve ganglia, Binet (89).—Terminations of nerves in muscles, Monti (621).—Histology of nervous system of larva of *Stratiomys*, Henneguy & Binet (399).—Innervation of wings and elytra in *Coleoptera*, Binet (88).—Centre of reflex action of leaping in *Orthoptera*, Heine (393).

Exner's views of vision discussed, Aurivillius (27).—Nature of the retinal image of insects, Viallanes (975).—Development of the imaginal eye of Vanessa, Johansen (457).—Sense-organs in antennæ of ants, Sergi (840).—Tibial cars of Locustidæ, Adelung (3).—Cenchri in Hymenoptera and Lepidoptera, Riley, P. E. Soc. Wash. ii, p. 309.—Odoriferous Chalcididæ, Howard, t. c. p. 215.

Construction of nests of ants, Forel (288).—Habits of ants, Wroughton (1018); Katurič (473, 474).—Wasp concealing tile, Blackburn, P. Ent. Soc. 1892, p. xx.—Intelligence of fossorial Hymenoptera, Pergande (671).—Instinct of fossorial Hymenopteron, Ammophila, Williston (1012).—Mode of stinging by Ammophila, Marchal (580).—Instinct of Apis mellifica, Smith (868).—"Assembling" males of Lepidoptera, Sykes, Ent. xxv, p. 84.

Social relations of Termitidæ, GRASSI (359).

Observations on the nature of the relations between Lomechusa and various kinds of ants, Wasmann (987).

Self-mutilation of Orthoptera, Werner (1002).—Music of Orthoptera, Scudder (812).—Insect cannibalism, Berg (58).

#### 3. Muscular System.

Histolysis and histogenesis of muscle, KOROTNEFF (511).—Condition of muscles of *Dytiscus* during contraction, TOURNEUX (930).

# 4. ORGANS AND FUNCTION OF NUTRITION, CIRCULATION, AND RESPIRATION.

Composition of the blood of insects, GRIFFITHS (363).

Digestive system of a Paussid Coleopteron, RAFFRAY (709).—Histology of alimentary canal and its glands, Holmgren (411).—Anatomy and histology of alimentary canal of Gryllotalpa, EBERLI (244).—Bacteria normally present in digestive organs of Hemiptera, Forbes (286).

Movements of the Malpighian tubules, MARCHAL, p. cclvi, Bull. Soc. Ent. Fr. lxi.

Respiration of larva of *Eschna*, Martin (585).—Origin of trachem from setiparous sacs, Bernard (73).—Evolution of heat by beetles, DORAN (223).

#### 5. SEXUAL ORGANS AND EMBRYOLOGY.

Reproductive organs of *Pentaplatarthus*, RAFFRAY (709).—Morphology of ovipositor of *Hymenoptera*, MARLATT (581).—Histology of the annexes of the sexual organs in *Periplaneta*, BLATTER (108).—The radiate bodies in the receptaculum seminis of *Pronuba* and *Prodoxus*, RILEY (742).

Graber (355) deals chiefly with the following points:—Chap. I, abdominal appendages; II, stigmata, tracheæ, Malpighian tubes, salivary glands, &c.; III, blood tissue; IV, the ganglionic chain; V, the genital glands; VI, enteroderm; VII, Keimhüllen.—Early stages in insects of various orders, and in parthenogenetic eggs; structure of micropyle in *Pyrrhocoris*, &c.; comparative embryology of the early stages, Henking (396).—Embryology of viviparous *Aphididæ*, Lemoine, p. clxiv, Bull. Soc. Ent. Fr. lxi.—Embryology and development of the blow-fly, Lowne (567).—Embryology of *Phyllodromia germanica*, Cholodkovsky (173); Heymons (404).—Embryology of *Smicra clavipes*, Henneguy (398).—Preliminary communication on various points in the embryology of parasitic *Hymenoptera*, Kulagin (528).—Mode of preparing eggs for investigation, Henking (397).

Structure of male organs and spermatogenesis in Laphria (Diptera), CHOLODKOVSKY (171); translated in Ann. N. H. (6) x, pp. 268-270.— Spermatogenesis in Gryllotalpu, RATH (713).—Micropyle (?) apparatus on eggs of Hemiptera, SHARP (842).

Wingless males of ants and consanguineous reproduction, Forel., p. 458, Ann. Ent. Belg. xxxvi.

Parthenogenesis: history of knowledge of, Taschenberg (915); in Chalcididæ, HOWARD (432).

Hermaphrodites: of Hymenoptera, Halictus and Macrophya, KRIEGER (527); of Azteca (Formicida), Forel (297); of Chalicodoma muraria, FREY-GESSNER, p. 372, MT. Schw. ent. Ges. viii; of Ocueria dispar, FERNALD, Canad. Ent. xxiv, p. 87; of Aglia tau, var. nigerrima, RUHL, p. 36, Soc. Ent. vii; of Aspilates strigillaris, SWIERSTRA (911).—Anatomy

of lateral hermaphrodite of Smerinthus populi, Tetens (918).—Papilio asterias, ind. of mixed sexes, Edwards, p. 49, Canad. Ent. xxiv.

Sexual structures and hybridism, ESCHERICH (257).—On obtaining hybrids in Lepidoptera, STANDFUSS (884).—Hybrid of Smerinthus ocellatus and populi, Ziegler, Berl. Ent. Zeit. xxxvi, p. xxiii; of Zygæna trigonellæ and transalpina, STRECKFUSS, l. c.

#### 6. GENERAL BIOLOGY AND STRAY NOTES.

Insects on the surface of snow, Ins. Life, iv, pp. 335 & 336.—Aquatic lepidopterous larvæ, MÜLLER (629).—Butterflies entering water, EDWARDS, Canad. Ent. xxiv, p. 16.

Carnivorous caterpillars, Plateau (694).—Secretion of potassium hydroxide by Dicranura, Latter (541).—Parnassius, formation of pouch, Scudder (813), and discussion, P. E. Soc. 1892, p. xxxiii.—Mode of expansion of the wings of Lepidoptera, Petersen (675).—Dorsal prolegs on Coleopterous larvæ, Riley (750).—Mode of succession of cycles of generations in Pemphigiens, Horvath (427).—Food influencing emergence of Lepidoptera, Merrifleld, Ent. M. M. xxviii, p. 20.—Bacterial structures in tissues and eggs, Blochmann (109).—Jumping seeds, Buchenau (139); Jumping seeds, cocoons, &c., Ascherson, SB. Ges. Naturf. 1892, pp. 19 & 20.—Stridulation in Lepidoptera, Hampson (385).—Stridulation of Acherontia, Roth, Ent. Tidskr. xiii, p. 250.—On the function of the striæ of Histeridæ, Lewis (557).—Different times of appearance of the sexes of Lepidoptera, value of, discussed, Petersen (676).—Hours of flight of insects, Schwarz (808).

Cave insects, RAFFRAY & BOLIVAR (710).

Variety of Papilio macareus, NICÉVILLE, J. Bomb. N. H. Soc. vii, p. 345, pl. i, fig. 1.—Albino of Colius, EDWARDS, p. 49, Canad. Ent. xxiv.—Pallid female varr. of species of Colias, Skinner & Mengel, P. Ac. Philad. 1892, pp. 156 & 157.—Colias edusa and helice, proportional numbers, ADYE, P. E. Soc. 1892, p. xxv.—Pieris napi, Irish var., WEIR, Ent. M. M. xxviii, p. 22.—Pieris rapæ, melanic var., Swierstra, pl. xvii, fig. 1, p. 337, Tijdschr. Ent. xxxiv.—Var. of Euplæa perdita, Snellen (871); varr. of Argynnis, Arctia, and Aglia, Büren, MT. Ges. Bern. 1892, p. xiv; var. of Argynnis paphia, Frohawk, Ent. xxv, p. 242: 9 var., Leech, p. 240, pl. xxiii, fig. 2, Butt. China; varieties of Vanessa antiopa, JÄGER, Ent. xxv, p. 43.—Charaxes jasius var., HONRATH, Berl. Ent. Zeit. xxxvi, p. xxiii; Melitæa cynthia and others, variation, Ziegler, l. c.; Zygæna spp., variation, THURAU, t. c. p. xxvi. Var. of Epinephele tithonus, SEEBOLD, p. 467, Berl. Ent. Zeit. xxxvi; varieties of Lepidoptera (Epinephele junira, &c.) in Rhineland, Füchs, JB. nass. Ver. xlv, pp. 85-106; var. of Epinephele hyperanthus, Frohawk, Ent. xxv, p. 215; var. of Lycana adonis, HANBURY, P. E. Soc. 1892, p. xxxiv; of Lycana bellargus, Austen, Ent. xxv, p. 16; of Polyommatus agestis, BARRETT, Ent. M. M. xxviii, p. 245.—Var. in neuration of Pisola (Hesperiida), NICÉVILLE, p. 351, J. Bomb. N. H. Soc. vii. Var. of Sesia formicæformis,

FROHAWK, Ent. xxv, p. 216; of Trochilium apiforme and Zygena cynare, STANDFUSS, pp. 368 & 369, MT. Schw. ent. Ges. viii; of Zygana filipendulæ, Bird, Ent. xxv, p. 194; of Z. loniceræ, Bromilow, t. c. p. 194. Variation in ocelli of Satyrida and formula for expressing it, COCKERELL, Canad. Ent. xxiv, p. 101.—Vanessa urtica, variation in China, LEECH, Butt. China, pp. 258-260.—Variation of Cynthia arsinoë, Röber, p. 302, Tijdschr. Ent. xxxiv.—Variation in males of Teracolus, YERBURY, p. 216. J. Bomb. N. H. Soc. vii.—Variation of secondary male characters in Forficula and Dynastida, statistics, BATESON & BRINDLEY (42); of Noctua festiva, BARRETT, Ent. M. M. xxviii, p. 47. Euchelia jacobea, yellow var. of, Anderson, Ent. xxv, p. 216; of Euproctis fulviceps, P. E. Soc. 1892, p. ii.—Breeding between Arctia lubricipeda and var., HARRISON, P. E. Soc. 1892, p. xxix.—Arctia caja var., LADDIMAN, Ent. xxv, p. 1, fig.; Anderson, t. c. p. 290.—Attempt to obtain varr. of Arctia caja by interbreeding, ARKLE, Ent. xxv, p. 57.—Liparis monacha, variation from parents from different localities, Clark, P. E. Soc. 1892, p. xxx.—Amphidasys betularius var. doubledayarius, breeding from, Steinert (897).

Variation in colour of eggs of Biston hirtaria, ADKIN, Ent. xxv, p. 129, Pearson, t. c. p. 163.—Localised variation of Callimorpha, Mason, Ent. News, iii, p. 52.—Variation and heredity in Pronuba and Prodoxus. BILEY (741).—Variation in Zonabris according to law, Escherich (258).—Seasonal variation in Satyridæ, Moore, Lep. Ind. i, pp. 147, &c., &c.—Extension of melanism on the Continent of Europe, Berger (59).—Amphidasis betularia, increasing extension of var. doubledayaria, HOFFMANN, Ent. M. M. xxviii, p. 48.—Phæism of Java Butterflies. Sanson, Ent. xxv, p. 43.—Local variation, &c., in the Bucovina, HORMUZAKI (414).—Variation of British Noctuce, Tutt (955); in several species of British Lepidoptera (Noctuidae, Geometridae), South (879) .--Trimorphism of Hypocala, BUTLER, Ann. N. H. (6) x, p. 17 .-Variation in colour of Sphingid larva, Beutenmüller, Bull. Am. Mus. iv. p. 65.—Variation of Spilosoma obliqua in Australia, Anderson, Vict. Nat. ix, p. 83.—Variation of neuration in Selidosemidæ, MEYRICK. pp. 124-140, Tr. E. Soc. 1892.—Polia chi, melanism, Porritt, P. E. Soc. 1892, p. ii.—Variation of Catocala nupta, Frohawk, Ent. xxv, p. 243.

Melanic var. of Eupithecia castigata, Tugwell, Ent. xxv, p. 41.—
Boarmia repandata, var., Hall, Ent. M. M. xxviii, p. 267.—Thamnonoma vauraria, remarkable aberration, Vries, p. 24, & Snellen, p. 25, Tijdschr. Ent. xxxv.—Many varr. of Lepidoptera noticed in Ent. Record, iii, passim.

Variation in markings of Coleoptera, Horn (420).—Variation of Cicindela flexuosa, Beuthin, Ent. Nachr. xviii, pp. 333-335; in other Cicindela, Beuthin, t. c. pp. 359-362, 376, & 377.—Variation in structure of Carabus hispanus, Beuthin, p. 211, Ent. Nachr. xviii.—Variation in structure of Dytiscus marginalis, ?, Peytoureau, Bull. Soc. Ent. Fr lxi, p. xlviii.—Variation in metasternal processes of Dytiscus marginalis Griffini (360).—Variation of Gnathocera hyacinthina, Kolbe, S. E. Z liii, pp. 129-131.—Typophorus canellus (Eumolpini), great colour

variation, HORN, p. 208, Tr. Am. Ent. Soc. xix.—Varieties of Coccinellidæ, Weise, pp. 17-24, Ann. Ent. Belg. xxxvi.—System of nomenclature for varr. of Coccinellidæ, Abeille, xxviii, p. 2.

Variation in Opomyza punctella, SINTENIS, p. 177, Wien. ent. Z. xi. Variation in melanic Cicadidæ, DISTANT, p. 111, Mon. Orient. Cicad.

Rudimentary wings of Coleoptera, CASEY, Ann. N. York Ac. vi.—Absence or inferior condition of eyes in cave insects, JOSEPH (458).

Green and brown pups of *Papilio podalirius*, Bonhote, Ent. xxv, p. 44; of *Papilio*, Cockerell, Ent. xxv, p. 93, and Frohawk & Bedford, t. c. p. 120.—Coloration of pups of *Limnas*, Yerbury, J. Bomb, Soc. vii, p. 209.

Origin of striking colour and eye spots in Lepidoptera, Portschinsky (700).—Protective and warning coloration, &c., Beddard (46).—Development of colour in the final metamorphosis, Verhoeff (965, 970).—Composition of pigment of Lepidoptera, Griffiths (362).—Chemistry of pigment in yellow Butterflies, Hopkins, Nature, xlv, p. 197.—Action of chemical agents on scales of some Lepidoptera, Coste (192).—Order (evolutionary) of colours in Lepidoptera, Coste, Ent. xxv, p. 28.—The peculiarities of the scale-pigments in numerous Lepidoptera, Urech (958).—On the process of coloration of the pupal wings of Pieris brassicæ and Vanessa, Urech (957).—A colouring matter in wings of pupa of Pieris brassicæ, Urech (956).—Colour and its relation to histolysis and histogenesis in Orthoptera, Künckel (529).—Changes of colour due to histolysis and histogenesis at the moults of Schistocerca, Künckel (530); cf. also Brongniart, t. c. p. lix.

Lepidoptera choosing protective position, Jeffreys, Ent. xxv, p. 68, Sykes, t. c. p. 69, & Biddle, t. c. p. 94.—Protection of Saturnia pavonia by its spines, Heim, Bull. Soc. Ent. Fr. lxi, p. xlviii.—Palatability of conspicuous larvæ, Gould (354); Perkins, Ibis, 1893, pp. 103 & 104.—Protection of Acronycta alni, Fitz-Gibbon (283).—Kallima, habits of, Fowler, P. E. Soc. 1892, p. xviii.

Resemblances: between various insects, seeds, &c., NICOLAS, Échange. 1892, p. 143; between varr. of different Coccinellida, BEDEL, p. 36, note. Abeille, xxviii; between ant and bug, BERGROTH (71), between two bugs, id. (72); between ant and bug, GERSTAECKER, JB. Hamb. ix. p. 51: of portions of Phyllium to vegetable structures, SHARP, P. E. Soc. 1892. p. xiv; between Gyrophæna and ant, Wickham (1009); between Psyllids and Coccids, MASKELL, Tr. N. Z. Inst. xxiv, p. 3; between Cicadidæ and Sphingidæ, DISTANT, p. 119, Mon. Orient. Cicad.; of Epicopeia to Papilio, Honrath, p. xxi, Berl. Ent. Zeit. 1892; between beetle, Exema gibber, and excrement of caterpillar, HARRINGTON. Ottawa Natural. 1892, p. 84; of beetle's egg to bark on which it is placed, Popenoe, Tr. Kansas Ac. xii, p. 15; resemblance to parts of vegetables, Plateau (695); between Pompilid and Cicindelid, Saussure, p. 338, Grandidier Madagascar, xx, pt. 1; of Lepidopterous larvee to snakes, HART, J. Bomb. N. H. Soc. vii, p. 105; resemblance between Aradida and Pentatomida, BERGROTH (70); Zygana and Tettigonia. HEYLAERTS, p. 47, Ann. Ent. Belg. xxxvi; Diptera and Tenthredinida, GIARD (336).—Resemblance of parasitised mealy-bug to puparium of Diptera, HOWARD, P. E. Soc. Wash. ii, p. 237; between ants and spiders, BANKS, P. E. Soc. Wash. ii, p. 174; grasshoppers agreeing in colour with burnt patches, Thomas, Nature, xlvi, p. 612.—The Yucca moth and bogus Yucca moth, RILEY, P. Biol. Soc. Washington, vii, p. 96.

Chrysalis resembling old man's head, HOLLAND (410).

Mimiery: Seitz (819, 820), Weir, P. E. Soc., 1892, p. xxxiv, Plateau (695), Haase (382), Swinhoe (910).—Aggressive mimiery, Heller (395).—Cases of mimiery enumerated, Heller, Ent. Nachr. xviii, p. 185.—The mimiery of Volucella, Bateson, Nature, xlvi, p. 585, Poulton, xlvii, p. 28, Bateson, t. c. p. 77, Hart, p. 78, Poulton, t. c. p. 126, Romanes, t. c. p. 200.—Volucella and Vespa, Nicolas (639).

Relation of colour of Lepidopterous larvæ, cocoons, and pupæ to surroundings, Poulton (703) and Gould (354).—Colour of pupæ and surroundings, Merrifield, P. E. Soc. 1892, p. xxx.—Temperature and colour in *Lepidoptera*, Merrifield, P. E. Soc. 1892, p. xxiii, and Adkin, t. c. p. xxxi.

Influence of temperature and light, applied in the pupal stage, on the colours of Lepidoptera, MERRIFIELD (600).—Source of colour of cocoons of Saturnia, Eriogaster, and of pupse of Vanessa urtica, BATESON (40, 41).—Cocoons of Halias chlorana, variation in colour of, Ent. Rec. iii, p. 9.

Deformities of some Coleoptera, Planet, Le Nat. 1892, pp. 50, 51.— Deformities in the abdominal segments of Tachinida, Townsend, Ent. News, iii, pp. 166, 167.—Supernumerary and abnormal tarsi, &c., of some Coleoptera, Wickham (1008), cf. also t. c. p. 182.—Zygana filipendula with fifth wing in place of a leg, Richardson (738) with fig.— Telephorus rusticus, additional leg, Adams, P. E. Soc. 1892, pp. iv & xii.— Antenna in Carabus monilis, Pic, p. 258, Rev. d'Ent. xi.—One-eyed bees, Cook, P. Am. Ass. xl, p. 327.—Shortened wings in Agrotis; abnormal antenna in Tragidion, Riley, P. E. Soc. Wash. ii, p. 181.

Competition of & bees, VERHOEFF (968).—Darwinism, VONHOF (977).

—The phylogeny and embryonic morphology of insects, CHOLODKOVSKY (170).—Phylogeny of social Hymenoptera, VERHOEFF (964); phylogeny of Papilionidæ, SPULER (882).—Wing-neuration and phylogeny of Lepidoptera, SPULER (881).—Homology of elytra of Coleoptera, HOFFBAUER (405).

Stylopized bees, Perkins (674), and Theobald (920).—Ants-nest beetles, Wasmann (986), Wickham (1005).—Galls: Brodie (129), Liebel (562), Martel (584), Mik (610).—Classification of galls, Kerner (480).—Galls of Cecidomyiidæ, Kieffer (493), Rübsaamen (786), Thomas (925).—Alpine Cecidomyideous galls, Thomas (924).—Galls of Lorraine, Kieffer (495). —Galls on Muhlenbergia mexicana, insects reared from, Webster, Canad. Ent. p. 243; of Eurostu solidaginis, Brodie (130).

Plants and ants, Keller (475), Warburg (982).

Insects and flowers: Insects attacking man. [See Economic.]

#### III.—FAUNISTIC AND PALÆONTOLOGY.

The numbers in brackets refer to "Titles"; by their means it is intended to give a reference to every memoir containing a notice affecting the entomological fauna of each of the twelve regions here made use of. But synonymical notes and changes of systematic position are not included in these references.

New species, whose locality is not known:—Coleoptera: Rhabdopholis melolonthoides, cf. Scarubæidæ. Hymenoptera: Evania cribrata, cf. Hym. par. Evaniidæ. Lepidoptera: Rhodoneura notula, cf. Pyralidæ, &c. Diptera: Bellardia furcata, Atylotus erraticus, rubrinotatus, BIGOT. Orthoptera: Mecopoda divergens, cf. Locustidæ; Phyllozelus genicularis, n. g. & sp., Brochopeplus reticulatus, n. g. & sp., cf. Locustidæ.

#### 1. ARCTIC AND ANTARCTIC.

West Greenland (Insecta), Fox (298).

Greenland (Lepidoptera), SKINNER & MENGEL (850); (Diptera), BRAUER & BERGENSTAMM (119).

## 2. Insular (including New Zealand).

Iceland (Coleoptera), Sénac, Bull. Soc. Ent. Fr. lxi, p. xxviii.

Madeira (Diptera), BIGOT (85).

Madeira and Azores (Orthoptera), Bolivar, Bull. Soc. Z. Fr. 1892, pp. 46-49.

Canary Is. (Coleoptera), Sénac (836); (Hymenoptera), Forbl (291), Kohl (501); (Lepidoptera), Alluaud, Bull. Soc. Ent. Fr. lxi, p. ccxxxix, Rebel (715); (Diptera), Brauer & Bergenstamm (119); (Orthoptera), Krauss (521).

St. Helena (Odonata), CALVERT (150).

Fiji Is. (Orthoptera), KARSCH (471, p. 343).

Fiji and Pelew Is. (Orthoptera), REDTENBACHER (717).

Loyalty Group (Lepidoptera), ROTHSCHILD (782).

Admiralty and Solomon Is. (Coleoptera), FAUST (278).

New Hebrides (Coleoptera), DESBROCHERS (203).

New Ireland (Lepidoptera), HONRATH (412).

New Caledonia (Rhynchota), DISTANT (214), and MONTANDON (617).

Norfolk I., New Caledonia (Rhynchota), DISTANT (213).

Pitcairn I. (Lepidoptera), WARREN (983).

S. Pacific Is. (Lepidoptera), DRUCE (227).

New Zealand and New Caledonia (Diptera), BIGOT (85).

New Zealand (Insecta), HUDSON (440); Insects introduced, Ins. Life, iv, p. 339; (Coleoptera), HORN (421); Platysoma jejunum withdrawn,

LEWIS (555); (Hymenoptera), FOREL (294), MARSHALL (583); (Lepidoptera), MEYRICK (605), WARREN (983); (Coccidæ), MASKELL (590).

Hawaiian Is. (Neuroptera), MACLACHLAN (574).

Galapagos (Diptera), BRAUER & BERGENSTAMM (119).

#### 3. PALEARCTIC REGION.

### (a.) The Region.

Coleoptera. Balbi (31, 32), Desbrochers (201, 202, 204, 205, 207, 208, 209), Escherich (259), Ganglbauer (319), Guillebeau (373), Jakowleff (451), Reitter (724), Schwarz (809, 810), Stierlin (899, 900, Théry (921).

Hymenoptera. Berthoumieu (74), Handlirsch (386), Jakowlew (453), Kohl (501), Semenow (828, 829).

Lepidoptera. ROMANOFF (779), RÜHL (789), STAUDINGER (886, 887), id. [S. Siberia] (891), WARREN (983).

Diptera. Bigot (83, 85), Brauer & Bergenstamm (119), Mik (608), Portschinsky (699).

Rhynchota. PUTON (707, 708).

### (b.) Collective Europe.

Hymenoptera. André (11), Konow (510), Thomson (927). Diptera. Kowarz (512), Rübsaamen (787). Rhynchota. Horvath (429).

## (c.) British Islands.

Additions.

Coleoptera. BLATCH, novelty, Ent. M. M. xxviii, p. 160.—CHAMPION, novelty, t. c. p. 68.—BLATCH & HORNER, n. sp. (103).

Hymenoptera. BIGNELL, n. sp. (82).

Lepidoptera. Barrett, novelty, Ent. M. M. xxviii, p. 244.—Wood, novelties, n. spp. (1014); id. n. sp. (1015).

Diptera. MEADE, n. sp. (595), n. sp. (596), novelty, n. sp. (597).— VERBALL, n. spp. (971).

Aphaniptera. SAUNDERS, novelty, Ent. M. M. xxviii, p. 66.

Rhynchota. SAUNDERS, novelties, Ent. M. M. xxviii. p. 8; novelty, id. t. c. p. 258.

Coccidæ. Newstead, novelties, n. spp. (637).

Neuroptera. Morton, novelty, Ent. M. M. xxviii, p. 108.

LOCAL LISTS.

Insecta. East Sussex, PORRITT, Ent. M. M. xxviii, p. 227.—Bristol (Phenological), P. Bristol Soc. vii, pp. 10 & 11.—Bayswater, Pascoe, Ent. M. M. xxviii, p. 230.—Withernsea, Baker & Porter, Naturalist, 1892, pp. 296 & 297.

Coleoptera. Cornwall, ISABELL, Rep. Penzauce Soc. 1891-92, p. 376.— Plymouth, KEYS, Ent. M. M. xxviii, p. 23.—New Forest, CHITTY, Ent. M. M. xxviii, p. 24, and t. c. p. 216.—Hastings, Bennett, Ent. M. M. xxviii, p. 193.—Gomshall, Butler, Ent. M. M. xxviii, p. 268.—Wellington, Elton, Ent. M. M. xxviii, p. 288.—Woking, Champion, Ent. M. M. xxviii, p. 193, and t. c. p. 243.—Gloucestershire, Shipp, t. c. p. 288.— Cardiff, Storrie, Rep. Cardiff Soc. xxiv, p. 64.—Cambridge, Fowler, Ent. M. M. xxviii, p. 215.—Withernsea, Fowler, t. c. p. 215.—Doncaster BAYFORD, t. c. p. 267.—Birkenhead, Ellis, Ent. xxv, p. 144.—The Hydradephaga of Lancashire and Cheshire, W. E. SHARP, Naturalist, pp. 189, &c., &c.—Scotland, CHAMPION, Ent. M. M. xxviii, p. 243.— Some additions to Scottish Coleoptera, with notes on species new or rare in the "Solway" district, LENNON & DOUGLAS, Ann. Scot. N. H. i, pp. 107-115.—Kirkcudbright, Douglas, t.c. p. 271.—Loch Awe, CHITTY, Ent. M. M. xxviii, p. 216.—Elginshire and Mid Lothian, Evans, Ann. Scot. N. H. i, p. 78.—Ireland, CUTHBERT, Brit. Nat. ii, p. 29; JOHNSON, Ent. M. M. xxviii, p. 213.—N. Ireland, Johnson, t. c. p. 134.—Donegal, JOHNSON, t. c. p. 310.—Louth, CUTHBERT, Irish Nat. i, p. 158; Dublin, Browne, t. c. p. 189.—Armagh, Johnson, t. c. pp. 14, &c.

Strepsiptera. Surbiton, SAUNDERS, Ent. M. M. xxviii, p. 249; Bruton, EATON, t. c. p. 250.

Hymenoptera. Wotton-under-Edge, Perkins, Ent. M. M. xxviii, p. 135.—Shirley, Rothney, t. c. p. 51.—Rugby, Morice, Rep. Rugby Soc. 1891, pp. 56-60.—Norfolk, Bridgman, Tr. Norw. Soc. iv, p. 690; Fauna and flora of Norfolk: part xii, Phytophagous Hymenoptera (Sawflies), Tr. Norw. Soc. iv, pp. 523-536.—Yorkshire, Nelson, Summerfield, Roebuck, Wilson, Naturalist, 1892, p. 325.—A preliminary list of the Aculeuta of Lancashire and Cheshire; Brit. Nat. ii, pp. 21-23, &c.—Scotland, Trail (950).—Dunbar, Pow, Ann. Scot. N. H. ii, p. 79.—Perthshire, M'Gregor, Tr. Perthsh. Soc. i, pp. 65-72 & 152.

Lepidoptera. The Lepidoptera of Britain, BARRETT (34).-Label list of British Lepidoptera: (first part) issued as a supp. to J. Inst. Cornwall, xi.—Notes concerning local British Butterflies, Goss, Ent. M. M. xxviii, p. 44.—Immigration of Pieris brassicæ, KERRY (490).—Occurrence of Deiopeia pulchella in England, South, Ent. xxv, pp. 153-155, 166-168, & 191.—Ocneria dispar in England, South, Ent. xxv, p. 259.— Colias edusa in England in 1892, Ent. M. M., Ent., Ent. Rec., &c., numerous notices.—Introduction of Callimorpha hera, LUFF, Ent. M. M. xxviii, p. 21.—S. Devon, Brooks, Ent. M. M. xxviii, p. 280; Prideaux, Ent. xxv, p. 221; BIGNELL, t. c. p. 256.—Sidmouth, Wells, Ent. xxv, pp. 67 & 68.—Ilfracombe, BOYD, P. E. Soc. 1892, p. vii.—Exeter, STILL, Ent. M. M. xxviii, p. 132; BARRETT, P. E. Soc. 1892, p. xiv.—Plymouth, Briggs, Ent. xxv, pp. 45-47.—Dartmoor, Still, t. c. pp. 228-230.— Dorset, CAMBRIDGE, Ent. xxv, p. 196; DALE (196), RICHARDSON (740). -Bloxworth, Cambridge, Ent. xxv, pp. 82-84.—Purbeck, Bankes, Ent. M. M. xxviii, pp. 48 & 49, and t. c. pp. 308 & 309.—Lepidopterous fauna of Portland, Richardson (739).—Bournemouth, Adye, P. E. Soc. 1892, p. xxv.—New Forest and Swanage, Alderson, Ent. xxv, pp. 313-315.— Lyndhurst, Jackson, t. c. p. 219.—Freshwater, Stevens, P. E. Soc. 1892, p. xi.—Winchester, NEVINSON, Ent. xxv, p. 169.—Hastings, BLOOMFIELD, Ent. M. M. xxviii, p. 267.—Brighton, FIELD, Ent. xxv, p. 318.—East Sussex, Tugwell, t. c. pp. 156-159.—Folkestone, Partridge, Ent. M. M. xxviii, p. 265.—Shorncliffe, id. t. c. p. 191.—Ashford, Viggers, Ent. xxv, p. 289.—Tunbridge Wells, BEECHING, P. E. Soc. 1892, p. xxv.—Dorking, FLOOD, Ent. xxv, p. 169.—Surbiton, Goss, Ent. M. M. xxviii, p. 246.— Alton, Reid, Ent. xxv, p. 220.—Middlesex, Cockerell, t. c. pp. 115-118, 130-134, 182-185, & 203-207.—Crouch End, James, t. c. pp. 65-67.— Tineina, &c., at Sandridge, GRIFFITH, Tr. Hertf. Soc. vi, pp. 97-102.-Wellington, GHICA, Rep. Wellington Soc. xxiii, p. 49.—Marlborough, MEYRICK, Rep. Marlb. Coll. Soc. xl, pp. 124, &c.—Essex, FITCH, Ess. Nat. vii, pp. 115 & 140; Cole & Argent, t. c. pp. 140 & 141.—Notes on the Lepidoptera of Northamptonshire, EDWARDS, J. Northampt. Soc. 1891, pp. 235, &c., and 1892, pp. 19-23.—Nottinghamshire, BARRETT, Ent. M. M. xxviii, p. 267.—Suffolk, Rendlesham, Ent. xxv, p. 244.— Aldeburgh, CRUTTWELL, Ent. M. M. xxviii, p. 253.—Norfolk, BARRETT, P. E. Soc. 1892, p. xxvii; and Tr. Norw. Soc. iv, pp. 691-701.—Ranworth, WHEELER, Ent. M. M. xxviii, p. 203.—Staffordshire, WOODFORDE, Ent. xxv, p. 241.—North Staffordshire, Daltry, t. c. p. 95.—N. Staffordshire, Daltry, Rep. N. Staffordsh. Club, 1888, p. 11; 1889, p. 12; 1890, p. 15; 1891, p. 14; 1892, p. 74.—Delamere Forest, ARKLE, Ent. xxv, p. 145.—Lincoln, CARR, Ent. M. M. xxviii, pp. 130 & 131.—Halifax, HALLIDAY, Naturalist, 1892, p. 339.—York, Hewett, Ent. xxv, pp. 170-172.—Isle of Man, CLARKE, Brit. Nat. ii, p. 24, and Lioar Man, pp. 287, &c., &c.—Windermere, Moss, Ent. xxv, p. 246.—Westmoreland, NEVINSON, t. c. p. 144.—Carlisle, ROUTLEDGE, t. c. p. 169.—A list of the Macro-Lepidoptera of Balerno, Mid Lothian, Ann. Scot. N. H. i, pp. 39, &c.—Jura, Evans, t. c. p. 141.—Perth, White, t. c. p. 271.— Rannoch, ADKIN, Ent. xxv, pp. 105-110.—Aberdeenshire, REID, Brit. Nat. i, p. 66, &c.; ii, p. 80, &c.—Lochinvar, Beveridge, Ann. Scot. N. H. i, pp. 172-177.—Armagh, Johnson, Ent. M. M. xxviii, p. 217; Mourne district, WATTS, l. c.

Diptera. South Devon, Matthews, Ent. M. M. xxviii, p. 268.—Horrabridge, Dale, t. c. p. 310.—Diptera of Dorsetshire, Dale, Brit. Nat. i, p. 3, &c.; ii, p. 63, &c.—Glanvilles Wootton, Dale, Ent. xxv, p. 72.—Dorset and Hants, Dale, Ent. M. M. xxviii, p. 268.—Sutton Park, Bradley, Ent. M. M. xxviii, p. 110.—Sherwood and Wyre Forests, id. t. c. p. 290.—North Uist, Dale, t. c. p. 50.

Rhynchota. The Hemiptera-Heteroptera of Britain, Saunders (792).—Additions to the list of Hemiptera-Heteroptera collected in the Island of Guernsey, Luff, Ent. M. M. xxviii, p. 7.—Woking, Saunders, t. c. p. 218.—Remarks on some Hemiptera-Heteroptera taken in the neighbourhood of Bath, Blathwayt, P. Bath Club, vi, pp. 315-326. List of terrestrial Hemiptera-Heteroptera taken in the neighbourhood of Bath, id. t. c. pp. 432-435.—Chobham and Surbiton, Saunders, Ent. M. M.

xxviii, p. 290.—Norfolk, Edwards, Tr. Norw. Soc. iv, pp. 702-710.—Chesil Bank,  $Coccid\alpha$ , Dale, Ent. M. M. xxviii, p. 219.

Neuroptera. Localities for Dragon-flies, Sci. Goss. 1892, pp. 17, &c., &c. — Neuroptera observed in the Channel Islands in September, 1891, McLachlan, Ent. M. M. xxviii, pp. 4-6.—A list of the Neuroptera inhabiting the Island of Guernsey, LUFF, Rep. Guernsey Soc. 1891, pp. 155-160.—Guernsey, LUFF, Ent. M. M. xxviii, pp. 72 & 73.—Sark, McLachlan, Ent. M. M. xxviii, p. 74.—Leatherhead, Briggs, t. c. pp. 194 & 268.—Chippenham, King, t. c. p. 246.—Penistone, Porritt, t. c. p. 311.—Ireland, Morton, t. c. pp. 110 & 301.

Orthoptera. Norfolk, SHAW, P. E. Soc. 1892, p. xxvii.

# (d.) North Europe and Siberia [see also (a.) and (b.)].

Insecta. KLÆR, Ent. Tidskr. xiii, p. 92.

Coleoptera. Sweden, Lampa, Ent. Tidskr. xiii, p. 208.—Siberia, BERGROTH (60); REITTER, Wien. ent. Z. xi, pp. 187, 188.—E. Siberia, BRENSKE (123), WEISE (995).

Hymenoptera. Norway, Fortegnelse over nogle for Norges Fauna nye arter af *Phytophage Hymenoptera*, KLER, Ent. Tidskr. xiii, pp. 67-70.—Sweden, NERÉN (631).

Lepidoptera. Scandinavia, Aurivillius (23); Sahlberg, Ent. Tidskr xiii, p. 291.—Roldal, Walker, Ent. xxv, pp. 36-39.—Christiania, Huitfeldt-Kaas, Ent. Tidskr. xiii, pp. 78-80.—Sweden, Ammitzböll, Ent. Tidskr. xiii, p. 73.—Siberia, Christoph (175), Ragonot (712), Romanoff (779).—Amurland, Graeser, (356).—S. Siberia, Lep. see (a.).

Rhynchota. Siberia, JAKOWLEFF (452), BERGROTH (65).

Diptera. Norway, BIDENKAP (81); Sweden, LAMPA (537).—Thysanura.

Siberia, Reuter (734).

# (e.) Central and Eastern Europe [see also (a.) and (b.)].

Insecta. Belgium, Selys (826).—Germany, Brischke (126, 127).—Rheinland, Verhoeff, Ent. Nachr. xviii, p. 3.—Galls of Lorraine, Kieffer (495).—St. Gall, Wegelin, Ber. St. Gall. Ges. 1890-91, pp. 307-335.—Trencsen, Brancsik, Jahresheft. Ver. Trencsin. xiii, pp. 43-46.—Becker, Sarepta (44).

Coleoptera. Denmark, Lovendal, Ent. Medd. iii, pp. 235-240.— Netherlands, Everts (260), and Tijdschr. Ent. xxxv, p. xii.—Belgium, Borre, p. 228, Ann. Ent. Belg. xxxvi; L. Coucke, t. c. p. 79; E. Coucke (193).—Matériaux pour la faune entomologique de la province du Luxembourg, belge. Coléoptères, Troisième centurie. De Borre, Publ. inst. Luxemb, xxi, pp. 81-117.—Schilsky, vii, Beitrag zur Kenntniss der deutschen Käferfauna, Deutsche e. Z. 1892, pp. 193-208.—Brunswick, Heller, Ent. Nachr. xviii, pp. 33-36.—Die Käfer von Nassau und Frankfurt. Sechster Nachtrag. JB. nass. Ver. xlv, pp. 65-82.—Göttingen, Rade, p. 367, Ent. Nachr. xviii.—Catalogue des Coléoptères récoltés à Aigle et environs, Bull. Soc. Murith. xix, pp. 21-60.—Switzerland.

GUILLEBEAU, Echange, i, No. 7, p. 3.—Simplon, BOURGEOIS, p. ccxiv, Bull. Soc. Ent. Fr. 1xi.—Brunn, FLEISCHER, pp. 206, 209, & 211, Wien. ent. Z. xi.—Silesia, LETZNER (551).—Austria, EPPELSHEIM (255).—Hungary, FRIVALDSZKY (310).—Carinthia, REITTER, p. 186, Wien. ent. Z. xi, Weise (1000).—Russia, Semenow (834).—Sarepta, Pic (677).

Hymenoptera. Belgium, Lameere (535).—Germany, Kriechbaumer (523, 524), Verhoeff (966).—Switzerland, Frey-Gessner (307, 308).—Russia, Morawitz (624).

Lepidoptera. Belgium, HIPPERT, pp. 451 & 480, Ann. Ent. Belg. xxxvi; Selys, t. c. p. 391.—Netherlands, Cappel, Tijdschr. Ent. xxxv, p. xv.—Loreley, Füchs (313).— Die Macrolepidopteren der Dresdner Gegend, Steinert, pp. 395-423, Deutsche e. Z. Lep. v.—Switzerland, Rühl, MT. Schw. ent. Ges. viii, p. 367.—Stilfser-Joch, Rühl (790).—Galicia, Klemensiewicz (499).—Bucovina, Hormuzaki (414).—Tyrol, Elwes, P. E. Soc. 1892, p. xxvi.—South Tyrol, Rebel (716).—Carinthia, Horrath (413).

Diptera. Kieffer (492, 493, 494), Meade (596), Mik (609), Strobl (902).—Belgium, L. Coucke, Ann. Ent. Belg. xxxvi, pp. 134-139.—Hertogenwald, E. & L. Coucke, Ann. Ent. Belg. xxxvi, pp. 462-469.—Switzerland, Becker (45), Bergroth (61), Röder, Wien. ent. Z. xi, p. 170.—Germany, Liebel (562), Rübsaamen (786), Stein (896).—Austria, Strobl (903, 905, 906).—Thuringia, Röder, Ent. Nachr. xviii, p. 204.

Rhynchota. Belgium, Fokker, Aun. Ent. Belg. xxxvi, p. 170; Énumération des Hémiptères de Belgique. II. Homoptera: Coubeaux, Ann. Ent. Belg. xxxvi, pp. 34-36, & 80-83.—Catalogus der in Nederland voorkomende Hemiptera: Addenda et corrigenda, Fokker, pp. 357-378, Tijdschr. Ent. xxxiv.—Lorraine, Kieffer, Ent. Nach. xviii, p. 30.—Hungary, Lethierry (549).—Pressburg, Sabrausky, Verh. Ver. Pressburg (n.s.) vii, pp. 16-24.—Duda, Schnabelkerfe Böhmens, cf. Wien. ent. Z. xi, p. 250.—St. Petersburg, Cholodkovsky (172).

Neuroptera. Schwarzwald, McLachlan, Ent. M. M. xxviii, p. 79.—Germany, Rostock (781).

# (f.) France [see also (a.) (b.) and (g.)].

Insecta. Aude, Liste des espèces de Diptères, Hémiptères et Hymenoptères recueillies dans le département et principalement aux environs de Carcassonne, Gavoy, Bull. Soc. Aude, iii, pp. 255-298.—Auvergne, Desbrochers, Frelon, i, pp. 2-10.—Ria, Xambeu, Soc. Agricole Pyrén.orient. xxxiii, pp. 229-240.

Coleoptera. ABEILLE (1), BRISOUT (128), CHOBAUT (169), and Bull. Soc. Ent. Fr. lxi, p. ccxxiv; CROISSANDEAU (194), and p. cclviii, Bull. Soc. Ent. Fr. lxi; GRILAT (365); GUILLEBEAU (375, 376, 377); and Catalogue des Coléoptères du département de l'Ain, Échange, No. 54, &c., &c.; JACQUET, Échange, i, No. 6, p. 3; REY (736); Somme, COLIN, Bull.

Soc. L. Nord. France, x, pp. 212-216, and Delaby, t. c. pp. 308-314; Seine-inf., Étienne, Bull. Soc. d'Elbeuf, p. 61; Catalogue des Coléoptères du Finistère (completed), Hervé, Bull. Soc. Finistère, xii, pp. 81-127; Supplement, pp. 129-132; Soulac, Gouin, Act. Soc. L. Bord. v, p. clxi.

Hymenoptera. Hyménoptères de la Vendée, BLAND, Rev. Sci. Nat. Ouest, pp. 141-149.

Lepidoptera. CHRÉTIEN (174), CONSTANT (182), RAGONOT, p. clxiii, Bull. Soc. Ent. Fr. lxi; Calvados, Dumans, p. lxxii, Bull. Soc. Ent. Fr. lxi.—Elbeuf, Dupont, Bull. Soc. d'Elbeuf, 1891, p. 43.—Alpes maritimes, Bromilow, Ent. xxv, p. 245.—Contribution à la faune des Lépidoptères du sud-ouest de la France: Supplément aux Catalogues de M.M. Roger et Trimoulet, et Catalogue provisoire des Pyrales des environs de Bordeaux, Breiquet, Brown, Act. Soc. L. Bord. xlv, pp. 49-91.

Rhynchota. HORVATH (431), PUTON, Échange, 1890, No. 62, p. 110, and Rev. d'Ent. xi, pp. 31 & 32.—Catalogue des Hémiptères, (Hétéroptères, Homoptères, Psyllides) recueillis dans la département de la Loire-inférieure, DOMINIQUE, Bull. Soc. Sci. Nat. Ouest. France, 1892, pp. 81-130.—Matériaux pour la faune des Hémiptères de l'Oise, CARPENTIER & DUBOIS, Mém. Soc. L. N. Fr. viii, pp. 406-440.

Neuroptera. MARTIN (587, 588, 589).

Orthoptera. Finot (282); Matériaux pour le Catalogue des Orthoptères de la Somme, Bull. Soc. L. N. Fr. x, pp. 69-75.—Loire-inf., Dominique, pp. 146-148.

# (g.) Southern Europe and Mediterranean Basin [see also (a.) and (b.)].

Insecta. Val Vigezzo, Carlini, Bull. Ent. Ital. xxiv, pp. 83-91.— I. of Lampedusa, Ragusa, Nat. Sicil. xi, pp. 234-238.

Coleoptera. ABEILLE (2), CHAMPION (163), LEWIS (557), PIC (678, 684, 687), ZOUFAL (1026).—Spain, GUILLEBEAU (378), BEITTER, p. 239, Wien. ent. Z. xi.—Italy, Dodero (219), Gestro (332).—Val Pesio, Ganglbauer (318).—Corsica, Croissandeau (194), Guillebeau (379).—I. of Capri, Blum, Nachrichtsbl. deutsche Malak. Z. 1892, pp. 27& 28.—Sicily, Baudi (43), Stierlin (898).—Syria and Cyprus, Areille (1).—Bosnia, Pic (690).—S. E. Europe, Reitter (729).—Herzegovina, Knotek (500), Ganglbauer (317).—Persia, Heyden, Deutsche e. Z. 1892, p. 111; Bedel (48).—Algeria, Grilat (364), Guillebeau (374), Lewis (560), Pic (680, 681, 686, 689, 691).—Egypt, Pic (685).—Asia Minor, Fleischer (284), Pic (679), Théry (922).—Syria, Fairmaire (263), Pic (683, 688).

Hymenoptera. EMERY (250), FOREL (291), SAUSSURE (793), VACHAL (959).—Spain, MEDINA (598).—Andalusia, MEDINA, Act. Soc. Esp. xxi, pp. 145-148. — Sardinia, KRIECHBAUMER (526). — Italy, BENZI (55), COBELLI (176).—Trieste, GRÄFFE (358).—Dalmatia, GASPERINI (320), VERHOEFF (969).—Bulgaria, FOREL (289), SCHULTHESS-RECHBERG (803).—Algeria, LESNE (548).—Aden, FOREL (290).

Lepidoptera. Zapater, Catálogo de los Lepidopteros de la provincia de Teruel, Ann. Soc. Esp. xxi, pp. 103-159.—S. France, Nicholson, Ent. M. M. xxviii, p. 270.—N. Italy, Norris, Ent. xxv, pp. 239, &c.—Modena, Pozzi, Note Lepidotterologiche, Atti Soc. Mod. (3) xi, pp. 1-11.—Sardinia, Mabille (572).—Sicily, Püngeler (706).—Roumania, Joannis, Bull. Soc. Ent. Fr. lxi, pp. ccxxxix-ccxli; Cosmovici (191).—Bucharest, Montandon, Bull. Soc. Ent. Fr. lxi, p. xlvi.—Tunis, Ragonot (711); Staudinger (890).—Aden, Butler (145); Yerbury (1025).

Diptera. Pavia, BEZZI (79).

Rhynchota. Italy, Su alcuni Emitteri Eterotteri del Modenese, Testi, Atti Soc. Mod. (3) xi, pp. 203-212. Ferrari (281).—Croatia, Horvath, Glasnik Naravosl. Druzt. vi, pp. 29-49.—Algeria, Montandon (618).—Sahara, Coccide, Targioni-Tozzetti (912, 913); Giard, Bull. Soc. Ent. Fr. lxi, p. celxxiii.—Egypt, Puton, Rev. d'Ent. xi, pp. 32-34.—Syria, Puton (708).

Neuroptera. Contribuzione allo studio dei Pseudoneurotteri del Modenese, Bentivoglio, Atti Soc. Mod. (3) xi, pp. 122-124.—Novak, Terzo cenno (Neuropt. and Orthopt.) sulla Fauna dell' Isola Lesina in Dalmazia, Glasnik Naravosl. Druzt. vi, pp. 50-58.

Orthoptera. Locusts near Florence, Bull. Eut. Ital. xxiv, pp. 164-169.

—Trentino, Cobelli, Verh.-z.-b. Wien, xlii, p. 61.—Novak, Secondo cenno sulla Fauna dell' Isola Lesina in Dalmazia, Glasnik Naravosl. Druzt. v, pp. 119-128.—Bulgaria, Frey-Gessner, MT. Schw. ent. Ges. viii, pp. 397-403.—N. Africa, Krauss (520).—Oran, Olivier, Bull. Soc. Ent. Fr. lxi, p. xlv.

# (h.) Caucasus and West and Central Asia [see also (a.) and (b.)].

Coleoptera. Caucasus, Reitter (727); Rost (780); Weise (999).— W. Aria, Brenske (125); Fairmaire (268, 271); Faust (274); Jacobsohn (445); Prochazka (705); Reitter (725, 726, 728); Schmidt (801); Semenow (831, 832, 833); Weise (992).—Turkestan, Eppelsheim (256); Heyden (402); Weise (994).

Lepidoptera. W. Asia, Alpheraky (4).—Centr. Asia, Graeser (357). Rhynchota. W. Asia, Horvath (429).

# 4. Africa (Ethiopian Region).

Insecta. DISTANT (210, 216).

Rhynchota and Lepidoptera. KARSCH (460).

Coleoptera. ALLUAUD (8); BRENSKE (122); DESBROCHERS (200); DUVIVIER (230, 231); FAIRMAIRE (262); GESTRO (329, 334); GORHAM (348); GORHAM & GAHAN (353); GROUVELLE (368); HORN (421, 423, 426); KERREMANS (482, 487, 488); KOLBE (502, 504, 505, 506, 507, 508); KRAATZ (514); LAMEERE (534); LEFÈVRE (543); LEWIS (558); NON-

FRIED (642, 648); PÉRINGUEY (672, 673); ROELOFS (774, 775); SCHMIDT (801, 802); SENNA (838); WASMANN (985); WEISE (993, 997, 998).

Hymenoptera. André (12); Emery (252, 253); Forel (290, 292); Handlirsch (386); Keller (475); Kohl (501); Magretti (575); Saussure (793); Stadelmann (883); Vachal (960).

Lepidoptera. Aurivillius (24); Butler (143); Butler & Rothschild (147); Holland (403, 407, 408); Honrath (412); Karsch (461, 462, 463, 464); Kirby (497); Mabille (571); Mabille & Vuillot (573); Pagenstecher (668); Robbe (766); Rogenhofer (778); Smith & Kirby (858); Staudinger (885, 888, 889); Vuillot (978, 979, 980); Warren (983); Weymer (1003).

Diptera. BEZZI (80); BIGOT (85); BRAUER (116); BRAUER & BERGENSTAMM (119).

Rhynchota. Bergroth (62, 68); Carlini (158); Distant (210, 213, 218); Gerstaecker (323); Karsch (465); Montandon (618, 619, 620); Reuter (732).

Neuroptera. Calvert (150); Karsch (468); Selys (825); Karsch (469, 470, 471, 472); Pictet & Saussure (692); Redtenbacher (717).

# 5. MADAGASCAR.

Insecta. Nossibé, Brancsik (115).

Coleoptera. FAIRMAIRE (266, 267, 269).—Seychelles, FAIRMAIRE (270); HORN (421); JACOBY (447); KRAATZ (513, 517); LEWIS (555); NONFRIED (642, 648); ROELOFS (774).

Hymenoptera. Forel (287); HANDLIRSCH (387); KOHL (501); SAUSSURE (793).

Lepidoptera. Butler (146); Smith & Kirby (858); Warren (983). Rhynchota. Bergroth (72); Distant (214, 218); Montandon (619); Reuter (733).

Neuroptera. SELYS (825); Seychelles, CALVERT (150).

## 6. TROPICAL AND EASTERN ASIA WITH JAPAN.

For Aden (Hymenoptera and Lepidoptera) see 3 (g.). Insecta. Seitz (820); id., China (821); id., Japan (822, 823).

Coleoptera. Allard (7); Aurivillius (25); Bates (37, 38); Bedel (47); Belon (53); Bergé, pp. 239-241, Ann. Ent. Belg. xxxvi; Bourgeois (113); Brenske (122, 123, 125); Brongniart (133); Candèze (152, 155); Desbrochers (200); Duvivier (232); Fairmaire (267); Faust (276); Frivaldszky, China (309); Gestro (324, 325, 326, 327, 328, 329, 330, 333); Gorham (349, 351); Grouvelle (367, 369, 370); Guillebeau (378); Horn (421, 422, 424); Jacoby (446, 448); Janson (454); Kerremans (484, 489); Kolbe (506); Kraatz (514, 516); Lefèvre (544); Lesne (547); Lewis (553, 555, 558, 561); id., Japan (554, 556); Nonfried (642, 644, 647); Olivier (652); Poll (696):

RÉGIMBART (720, 723); RITSEMA (757); SCHMIDT (801); SCHENFELDT, Ann. Ent. Belg. xxxvi, pp. 107-111; SENNA (838, 839); SHARP (843); VILLARD, Japan (976); WEISE (997, 1001).

Hymenoptera. André (12); Cameron (151); Forel (290); Forel, Ceylon (288, p. 17, 293); Handlesch (386); Jakowlew (453); Kohl (501); Magretti (576); Mocsáry, China and Tibet (615); Rothney, E. India ants, P. E. Soc. 1892, pp. viii-x; Saussure (793); Wroughton, Ants of India (1018).

Lepidoptera. Austaut, Japan (29); Butler, Ceylon (142); Elwes (248); Hampson, Moths of India, vol. i (384); Leech (542); Manders (579); Moore (622, 623); Nicéville (638); Oberthur (650, 651); Poujade (702); Robbe (765); Romanoff, China and Mongolia (779); Seitz (817, 818, 820, 921); Sonthonnax (878); Pekin, Staudinger, Deutsche e. Z. Lep. iv, p. 339; India, Swinhoe (908, 909); Warren (983).

Diptera. BIGOT (83, 84, 85, 86, 87); BRAUER & BERGENSTAMM (119). Bhynchota. BERGROTH (63, 66, 67, 70, 71); DISTANT (211, 212, 213, 217); HORVATH, Trop. Asia and Tibet (428); LETHIERRY (550); MASKELL, Coccidæ (500); MONTANDON (617, 618, 619, 620); REUTER, E. Siberia and Mongolia (730).

Neuroptera. EATON (241, 242); id., Japan (243); KARSCH, China (467); SELYS (824).

Orthoptera. Brunner, Ceylon (137); PICTET & SAUSSURE (692).

# 7. ASIATIC ARCHIPELAGO, INCLUDING PAPUA.

Coleoptera. Bourgeois (112); Borneo, Bourgeois (114); Brenske (122, 123); CANDÈZE (156); id., Engano (153); id., New Guinea (154); DESBROCHERS (200); DOHRN, Philippines (221); FAIRMAIRE, Sumatra (265); FAUST (278); GESTRO (327, 329); id., Engano (335); id., New Guinea (331), and Ann. Mus. Genov. (2) x, pp. 265-268; GORHAM, Borneo (350); GROUVELLE (372); id., Philippines (367); id., Borneo (371); HELLER, Java (394); HORN (421, 424); JACOBY, Sumatra (448); Janson (454); Kerremans, Philippines (483); id., Engano (485); id., New Guinea (486); Kolbe (506); Kraatz (514, 516); id., Java (515); Lewis (555); id., Borneo (558); Nonfried (642, 644); OLIVIER, New Guinea (653); RAFFRAY, Philippines (710); RÉGIMBART (721); id., Borneo (722); RITSEMA (763, 764), and Notes Leyd. Mus. xiv, pp. 139, 143, & 144; id., Java (759, 762); id., Java and Nias I. (758); id., Borneo (761); id., Amboyna (760); Roelofs, Borneo (772); id., Aru Is. (776); Schmidt (801); Senna, Java (837); Sharp, Java (844); TRYON, New Guinea (953); WATERHOUSE, Damma I. (988); Weise (997); id., Celebes (998).

Hymenoptera. André, Borneo (13); Handlirsch (386); Kohl (501); Stadelmann (883).

Lepidoptera. Butler, Borneo (141); Hagen, Banka (383); Heylaerts (403); Honrath (412); Moore (623); Nicéville (638);

PAGENSTECHER (668); RIPPON, Aru Is. (756); RÖBER (767); ROGENHOFER, Sumatra (778); ROTHSCHILD, Celebes (783), and ROTHSCHILD, P. E. Soc. 1892, p. xv; SEMPER, Philippines (835); SMITH, Borneo (857); SNELLEN, Flores (869); id., Java (872, 876); id., Celebes (870); STAUDINGER, Borneo (892, 894); id., Mindoro (888); SWINHOE (908); TRYON, New Guinea (954); WARREN (983).

Diptera. BIGOT (85); BRAUER & BERGENSTAMM (119); BÖDER (771); WULP, Java (1019).

Rhynchota. BERGROTH (63); id., Engano (66); DISTANT (212, 213); id., Borneo (211); id., Timor (215); TRYON, New Guinea (952).

Neuroptera. KARSCII, Java (466); SELYS (824).

Orthoptera. Bolivar (710); Dohrn (222); Pictet & Saussure (692); Redtenbacher (717).

## 8. Australia and Tasmania.

Insecta. Tasmania, WALKER, Ent. M. M. xxviii, p. 153; Notes from Fremantle, &c., id. t. c. p. 55.

Coleoptera. Bedel (47); Scymnus lophanthæ said to be Australian, Ent. News, iii, p. 144, note; Blackburn (90, 91, 92, 93, 94, 95); Faust (277); Lewis (555); Nonfried (644); Olliff (655); Poll (697); Roelofs (776); Schmidt (801); Sharp (843); Sloane (852, 853).

Hymenoptera. FROGGATT (311, 312); HANDLIRSCH (386); KOHL (501); ROTHNEY, P. E. Soc. 1892, p. iii; SAUSSURE (793).

Lepidoptera. Lower (565, 566); Lucas (568, 569); Meyrick (604); Miskin (611, 612, 613); Olliff (654); a list of Victorian *Rhopalocera*, Spry & Anderson, Vict. Nat. ix, pp. 121-123; Swinhoe (908).

Diptera. BIGOT (83, 85); BRAUER & BERGENSTAMM (119); RÖDER (771).

Rhynchota. Murray I. and Mabuiag, CARPENTER (159); DISTANT (212, 213, 214, 217); MASKELL (590, 592).

Orthoptera. REDTENBACHER (717); SAUSSURE (795); TEPPER (916, 917).

## 9. North America.

Insecta. Godman & Salvin (347); Jamesburg, Ent. News, iii, p. 36.—Catalina I., Seavey, Zoe, iii, pp. 262 & 263.—New Mexico, Townsend (941).—Mississippi Bottoms, Weed, Canad. Ent. xxiv, pp. 276-279.—Alaska, Wickham (1007).—Ottawa, numerous records by Harrington and J. F. in Ottawa Natural. 1892. Insects in Cryptoporus valvatus in British Columbia, Hurbard, Canad. Ent. xxiv, pp. 250-256.

Coleoptera. Angell (14); Blaisdell, San Diego, Zoe, iii, pp. 47-49, 102, 252.—Blaisdell (96, 97, 98); Brendel (120, 121); Casey (161, 162); San Francisco, Dunn, Zoe, ii, p. 310; N. Brunswick, Harrington, Canad. Ent. xxiv, p. 112; id. (389); Horn (416, 417, 418, 418); W. Horn (421); Schwarz (805); Wickham (1009).

id., Bull. Lab. Iowa, ii, p. 234; id., Alaska (1006); id., Iowa, Canad. Ent. xxiv, pp. 99-101.

Hymenoptera. Ashmead (17, 18, 19, 20, 21, 22); Brodie (129); Fox (298 to 306), and Ent. News, iii, p. 17; Fyles (314); Gillette (344); Handlirsch (386); Harrington (388); Howard (433, 435); Kohl (501); Patton (670); Robertson (768).

Lepidoptera. Catalogue, SMITH (859).—DYAR (238, 239, 240); EDWARDS (246, 247); FERNALD (280); HULST (441); KELLICOTT (477); NEUMOEGEN (633, 634); NOLAN (641); PAGENSTECHER (668); RILEY (744); SKINNER (847, 848, 849); SLOSSON (854, 855); SMITH (860, 861, 863, 867); WARREN (983); WRIGHT (1017).—Ohio, KELLICOTT, Canad. Ent. xxiv, p. 42.—Adirondack, McKnight, Ent. News, iii, pp. 87-89.—N. Carolina, Mead, Canad. Ent. xxiv, p. 313; Moffat, Canad. Ent. xxiv, pp. 18 & 132.—Alaska, Wright, Ent. News, iii, p. 74.—A list of the Butterflies found at Marshall, Missouri, and vicinity, STALEY, Canad. Ent. xxiv, pp. 201-204.—Colorado, Bruce, Ent. News, iii, p. 136, and Cockerell, p. 192, Canad. Ent. xxiv.—Notes on some of the Butterflies of the Yosemite Valley and adjacent region, Van Dyke, Zoe, iii, pp. 237-241.—Pennsylvania, Ehrman, Ent. News, iii, p. 163.

Diptera. ALDRICH (5, 6); BIGOT (83, 85); BRAUER & BERGENSTAMM (119); Coquillett (185, 186, 187, 188, 189, 190); Hood, Payche, vi pp. 283 & 284; SNOW (877); TOWNSEND (931, 933, 934, 935, 936, 937, 938, 939, 940, 942, 943), and Psyche, p. 220; WILLISTON (1011).

Rhynchota. BERGROTH (64); DUZEE (233, 234, 235, 236, 237); GODING (345, 346).—Fauna ottawaensis. Hemiptera, Ottawa Natural. pp. 25-32.

Neuroptera. Banks, Catalogue (33); Canada, Moffat, Canad. Ent. xxiv, p. 11.—Odonata of Maine, Wadsworth, Ent. News, iii, p. 8, and Harvey, t. c. pp. 92 & 116.

Thysanura. HARVEY (390, 391).

Orthoptera. BLATCHLEY (104, 105); SCUDDER (816).

# 10. CENTRAL AMERICA, INCLUDING MEXICO.

Insecta. GODMAN & SALVIN (347).

Coleoptera. Bates (36); Casey (162); Faust (273); Gahan (315); Horn (426); Kraatz (518); Lewis (559); Nevinson (635); Nonfried (642-645).

Hymenoptera. HANDLIRSCH (386); KOHL (501); SAUSSURE (793).

Lepidoptera. DRUCE (226); PAGENSTECHER (668); RILEY (750); SCHAUS (798, 799); THIERRY-MIEG (923).

Diptera. BIGOT (83, 85); BRAUER & BERGENSTAMM (119); GIGLIO-TOS (342); MIK (606); TOWNSEND (938).

Rhynchota. BERGROTH (65); DISTANT (214); DUGES (229), Orthoptera. PICTET & SAUSSURE (692); SAUSSURE (794).

1892. [VOL. XXIX.] R. 6

#### 11. Antilles.

Insecta. Jamaica, Cockerell (179), and J. Inst. Jamaica, i, pp. 71-74.

—Grenada, Rep. Brit. Ass. 1892, pp. 353 & 354.

Coleoptera. FAUST (273).

Hymenoptera. Jamaica, André (12); Jamaica, Fox, Ent. News, iii, p. 226; Cuba, Ashmead (19).

Lepidoptera. PAGENSTECHER (668); St. Lucia, SCHAUS (798) p. 10; Weir, J. Inst. Jamaica, i, p. 106.—Bahamas, see N. America.

Diptera. Haiti, BIGOT (83, 85); BRAUER & BERGENSTAMM (119).— Jamaica, Cockerell (180); Townsend (931, 932, 946, 947).

Rhynchota. Jamaica, Johnson & Fox, Ent. News, iii, p. 59.

Orthoptera. St. Vincent, BRUNNER & REDTENBACHER (138).

## 12. South America.

Insecta. GODMAN & SALVIN (347); MEINERT (599).

Coleoptera. BEDEL (47); BERG (57); DESBROCHERS (206); FAIR-MAIRE (261, 264); FAUST (273, 275, and 278, p. 203); FLEUTIAUX (285); GAHAN (316); GERMAIN, Chili (322); GROUVELLE (366); HORN (421, 425); LÉVEILLÉ (552); NEVINSON (635); RUGE (788); SENNA, Chili (838).

Hymenoptera. André (12); Buysson & Marshall (149); Forel (290, 295); Handlirsch (386); Kohl (501); Saussure (793); Stadelmann (883).

Lepidoptera. Butler (145); Dognin (220); Druce (225); Kirby (497); Oberthur (650); Pagenstecher (667, 668); Schaus (798, 799); Smith & Kirby (858); Staudinger (893); Thierry-Mieg (923); Vuillot (980); Warren (983); Brief aus Bolivien, Garlepp, Deutsche e. Z. Lep. v, pp. 272-276.

Diptera. Arribalzaga (15); BIGOT (83, 85, 87); BRAUER & BERGENSTAMM (119); MIK (606); OSTEN-SACKEN (663); RÖDER (769); WILLISTON (1010); WULP (1019).

Rhynchota. Berg (56); Bergroth (65); Distant (213, 214); Douglas, Coccids (224); Morgan (625); Reuter (731).

Orthoptera. REDTENBACHER (717).

# PALÆO-ENTOMOLOGY.

[Cf. Emery (249), Meunier (602), Moberg (614), Schäff (797), Schlechtendal (800), Scudder (814).]

# COLEOPTERA.

Coleoptera in peat; SCHAFF (797).

Remarks on, with summary of, the N. American Tertiary Rhynchophorous fauna; SCUDDER (814).

# HYMENOPTERA.

Ectatomma gracile, Sicilian amber; EMERY, p. 145, pl. i, figs. 1 & 2, Mem. Acc. Bologn. (5) i, n. sp.

Ponera? leptocephala, Sicilian amber; EMERY, p. 146, pl. i, figs. 3 & 4, Mem. Acc. Bologn. (5) i, n. sp.

Cataulacus silvestrii, p. 147, pl. i, figs. 5-7, planiceps, p. 148, figs. 8 & 9, Sicilian amber; EMERY, Mem. Acc. Bologn. (5) i, n. sp.

Hypopomyrmex, n. g., near Strumigenys, p. 148, for H. bombiccii, n. sp., p. 149, pl. i, figs. 10 & 11; EMERY, Mem. Acc. Bologn. (5) i.

Acrostigma, n. subg. of Podomyrma, p. 149, for P. (A.) mayri, n. sp., Sicilian amber, p. 150, pl. ii, figs. 12-16; EMERY, Mem. Acc. Bologn. (5) i.

Aëromyrma sophiæ, Sicilian amber; EMERY, p. 151, pl. ii, figs. 17 & 18, Mem. Acc. Bologn. (5) i, n. sp.

Crematogaster pracursor, Sicilian amber; EMERY, p. 152, pl. ii, figs. 19-21, Mem. Acc. Bologn. (5) i, n. sp.

Leptomyrmex maravignæ, Sicilian amber; EMERY, p. 152, pl. ii, fig. 22, Mem. Acc. Bologn. (5) i, n. sp.

Tapinoma minutissimum, Sioilian amber; EMERY, p. 153, pl. iii, figs. 24 & 25, Mem. Acc. Bologn. (5) i.

Technomyrmex deletus, Sicilian amber; EMERY, p. 153, pl. iii, figs. 26-28, Mem. Acc. Bologn. (5) i, n. sp.

Plagiolepis labilis, Sicilian amber; EMERY, p. 154, pl. iii, figs. 29-32, Mem. Acc. Bologn. (5), i, n. sp.

Gesomyrmex corniger, Sicilian amber; EMERY, p. 155, pl. iii, figs. 33-35, Mem. Acc. Bologn. (5) i, n. sp.

Œcophylla sicula, Sicilian amber; EMERY, p. 156, pl. iii, figs. 36 & 37, Mem. Acc. Bologn. (5) i, n. sp.

#### DIPTERA.

Leptidæ in amber; MEUNIER, p. lxxxiii, Bull. Soc. Ent. Fr. lxi (provisional new genus, Palæochrysopila).

Dolichopodidæ found in amber, notes on; MEUNIER (602).

#### RHYNCHOTA.

Protocimez siluricus, Silurian of Sweden; Moberg, Geol. Fören. Stockholm Förh. xiv, p. 121, n. sp.

Notonecta, note on fossil; SCHLECHTENDAL (800).

#### IV.—ECONOMIC.

Reports: LINTNER (564); ORMEROD (656).—Report on the position of economic entomology in Europe; Bos, Tijdschr. Ent. xxxv, pp. xlvii-lxviii: on the ravages of *Xyleborus perforans* in the West Indies; BLANDFORD (100).—Insects injurious to *Conifera* in Britain; BLANDFORD (99).—Various notes on injurious insects in Rep. E. Soc. Ont. xxiii,

by BETHUNE, MOFFAT, FLETCHER. — For various notes on injurious insects, see Ins. Life, iv & v, passim.—Report on economic entomologists' work in N. America in 1891, see Bull. Dep. Agric. Ent. xxvi.

Bruchus obtectus, oviposition, &c.; RILEY (752, 753, 755).—"Entomological Notes" on various insects injurious in Australia; Olliff, in Agric. Gaz. N.S.W. iii, pp. 26, &c., &c.—Injurious insects in Kazan district; SMIRENSKY (856).—Injurious insects in Sweden in 1891; LAMPA (536) pp. 1-38.

Use of hellebore for destroying larvæ; SMITH, Ent. XXV, p. 80. —Destruction of insects by fungi; DECAUX, p. 196, Le Nat. 1892.—Experiments on infecting Chinch-bugs; SNOW, Tr. Kansas Ac. XII, pp. 34 & 119, and Psyche, vi, pp. 225-233.

Destruction of Melolontha in Spain; DELACROIX, Act. Soc. Esp. xxi, pp. 3-7.—Destruction of Melolontha by Botrytis, experiments; DUFOUR (228).—Elaterid injurious to wheat in Argentine Rep.; BERG (57).— Longicorn attacking orange trees in Australia; OLLIFF (655).—Oncoscelis sulciventris injurious to oranges in Australia; Olliff, p. 368, Agric. Gaz. N.S.W. iii.—Carpocapsa pomonella in Australia; Agric. Gaz. N.S.W. iii, p. 504, pl. xxxiii.—Ravages of Phædon cochleariæ; Enock, Ent. xxv, pp. 230-233.—Galeruca xanthomelæna, notes on; Smith, Canad. Ent. xxiv, pp. 246-250.—Selandria cerasi in Australia; Olliff, p. 27, Agric. Gaz. N.S.W. iii.—Habits of Hypoderma lineata; RILEY (754).—Hylemyia nigrescens, destructive to carnations; MACLACHLAN, Ent. M. M. xxviii, p. 135.—Chlorops pumilionis, metamorphoses; LAMPA (538).—Chlorops tæniopus, Meig, ravages in Sweden; HEDSTRÖM, Ent. Tidskr. xiii, pp. 201 -203.—Ravages of Oscinis frit in Sweden; Aurivillius (28).—Horn-fly in Florida; Rolfs, Bull. exp. Station Florida, xvii, pp. 12-14.—Hæmatobia serrata in Canada; WINN (1013).—Lepidoptera injurious to sugar cane in Java; Snellen (872, 873).—Diatræa saccharalis in Jamaica; COCKERELL (177).—Ocneria dispar: special report of the State Board of Agriculture on the work of extermination. Boston: 1892. Cf. Canad. Ent. xxiv, p. 156.—Liparis monacha in Germany; FERNOW, pp. 172, 216, & 237, P. E. Soc. Wash. ii.—Liparis monacha in Germany and Switzerland; FENK (279). - Grapholitha tedella, injurious to Abies excelsa: PUTON, p. x, Bull. Soc. Ent. Fr. lxi.—Ctenopseustis obliquana, ravages in N. Zealand; SMITH, Ent. M. M. xxviii, p. 110.—Lita solanella in America; Ins. Life, iv, p. 241.—Lithocolletis platana, injurious at Perpignan; XAMBEU, p. 244, Soc. Agricole Pyrén.-orient. xxxiii.—Lampronia capitella, life-history; CHAPMAN (164).—Halticus minutus injurious to Arachis hypogæa; Giard (337).—Psylla pyricola, report on; Slingerland (851): P. pyricola in N. America; id. Ins. Life, v, p. 100-104.—Injury to Laurus nobilis by Trioza alacris; THOMAS (926).—Phylloxera, destruction of in vineyards; BEHR (52).—Spread of Phylloxera in Germany, and cost of combating it; KESSLER (491). - Schizoneura lanigera in Australia; Olliff, Agric. Gaz. N.S.W. iii, p. 26.—Locusts near Florence: Bull. Ent. Ital. xxiv, pp. 164-169.—Locust in Algeria in 1891; Kunckel (531, 532).—Destruction of locusts in N. Africa; DECAUX, Le Nat. 1892. pp. 168 & 213.—The locust in Egypt in 1891, and mode of its destruction; Agric. Gaz. N.S.W. iii, pp. 829-832.—Destructive locusts in N. America in 1891; Bruner, Coquillett, & Osborn (136).—Gryllus servillei injurious to trees in Australia; Olliff, Agric. Gaz. N.S.W. iii, p. 270.

Insects and Flowers: Kunze (533).—Fertilization of Ficus roxburghii by Psenes; Cunningham (195).—Pronuba and Yucca; Riley (741).—Insects frequenting Yucca blooms; Townsend (948).—An insectivorous primrose; Mally (577).—Cnicus discolor as an insect trap; Blatchley, p. 310, Canad. Ent. xxiv.—Bees as fertilizers; Benton, Ins. Life, iv, p. 254.—Bees and flowers: constancy to various kinds of flowers; Bulman, Sci. Goss. 1892, p. 98.

Insects attacking man: *Œstrides* in America; Blanchard (101, 102).

—*Œstrus* in man; Sangagli (791).—The bite of *Leptis scolopacea*; Heim, p. c, Bull. Soc. Ent. Fr. lxi, and t. c. pp. clv-clvii.

Cutaneous larvæ; Germain, p. 59, Act. Soc. Chili, i.—Diptera as carriers of infection; Sawtschenko (796).

Collections: Collection and preservation of insects in the tropics; KANNEGIETER (459).—Method of preparing Cecidomyiid larvæ; Thomas, Verh.-z.-b. Wien, xlii, p. 356.—Mode of preparing Lepidopterous larvæ, with fig. of the apparatus; Oudemans (664).—Mode of preparing eggs for investigation; Henking (396).—Collecting and preserving Coleoptera in the Tropics; Ribbe, pp. 125-138, Berl. Ent. Zeit. xxxvii.— Effects of coloured glass in preventing fading of Lepidoptera.—Collections of Coleoptera; Leprieur (546).—Occurrence of verdigris in Coleoptera; Schwarz (807).—The use of silver pins; Sharp, Ent. xxv, p. 54.—Cement for use in closing bottles containing alcohol; Lataste (539).—Relaxing Lepidoptera; Cant, Ent. xxv, p. 119.—Packing specimens of insects for transportation; Canad. Ent. xxiv, pp. 117 & 118.—Notes on killing, preserving, and relaxing insects; Fletcher, Canad. Ent. xxiv, pp. 14-16.—Bait for Lepidoptera; Streckfuss, Berl. Ent. Zeit. xxxvi, p. xix.

Account of the Berlin Museum and the number of species it contains; ASHMEAD (16).—List of types and of rare species of aquatic Coleoptera in the National Museum at Brussels; SEVERIN (841).—Catalogue of Coleopterous larvæ in the collection at Copenhagen; MEINERT, Ent. Medd. iii, pp. 167-205.—Study of the types of Anthomyiidæ in the Zetterstedt Collection at Lund; STEIN (895).—Albarda's Collection of Neuroptera placed in the Leyden Museum; Ent. News, iii, p. 259.

Account of the collections of *Coleoptera* of H. W. Bates; OBERTHUR, p. cviii, Bull. Soc. Ent. Fr. lxi.—Lists of types of Grote & Robinson's *Lepidoptera* in the American Museum; BEUTENMÜLLER (76).

History of the Entomological Club of London; SOUTH (880).

# V.—BIBLIOGRAPHY, NOMENCLATURE, BIOGRAPHY.

- 1. Bibliography. Piller & Mitterpacher: Bedel reproduces the descriptions of Coleoptera described as new in the "Iter per Poseganam;" Abeille, xxvii, pp. 291-300.—Clerck's Icones, account of, by Holland, Canad, Ent. xxiv, pp. 83-87.—Notes on the bees described in Gay's Chili; Reed (719).—Critical and synonymical review of Saalmüller's work on Madagascar Lepidoptera; Snellen (875).—Bibliography of Scandinavian Entomological literature, continued by Bergroth & Aurivillius in Ent. Tidskr. xiii.
- 2. Nomenclature. Rules of nomenclature adopted by the Zoological Congress at Moscow; Zool. Anz. xv, pp, 440-443.—Query as to the mode of procedure in some cases of double employment of names; Dalla Torre, p. 92, Wien. ent. Z. xi.—Variable priority; Beckers, p. 318, Wien. ent. Z. xi.—"Correct" names; Maskell, Tr. N. Z. Inst. xxiv, p. 9.—On the distinction of generic names; Casey, Ann. N. York Ac. vi, pp. 10-12.—Classical emendations; Sharp, Ent. M. M. xxviii, p. 111.—The use of capital small letters for personal specific names; Kraatz, Deutsche e. Z. 1892, p. 380.
- 3. Biographical. Correspondence between Harris, Say, and Pickering (continued); Psyche, vi, pp. 297, 298, & 345.—Biographical notes, with portraits, have appeared in Brit. Nat. ii, of Lord Walsingham, p. 20; S. J. Capper, t. c. p. 60; E. A. Ormerod, t. c. p. 82; J. W. Douglas, t. c. p. 128; G. T. Porritt, t. c. p. 172; D. Sharp, t. c. p. 199; Rev. O. P. Cambridge, t. c. p. 231; H. W. J. Vaughan, t. c. p. 258.

Obituary Notices. Francis Archer; Ent. M. M. xxviii, p. 112, and Ent. xxv, p. 100.—H. W. Bates: by A. R. W., Nature, xlv, p. 398; by E. CLODD (39); by McLachlan, Ent. M. M. xxviii, p. 83; by D. S., with portrait, Ent. xxv, pp. 77-80.—Edward Burgess, with list and account of his writings, Scudder, P. Bost. Soc. xxv, pp. 358-364.—Hermann C. C. Burmeister: Ent. Nachr. xviii, pp. 220-222; Ent. M. M. xxviii, p. 221; Account of his funeral, An. Soc. Arg. xxxiii, pp. 145-150.— G. Carlet: Le Nat. 1892, p. 152; Bull. Soc. Ent. Fr. lxi, p. cxxix; with list and abstracts of his memoirs, CHAUMAT (167).—F. B. Caulfield; Canad. Ent. xxiv, p. 104.—Dr. C. A. Dohrn: Ent. M. M. xxviii, p. 164; by GROTE, Canad. Ent. xxiv, pp. 181, 182.—Prof. Veit Graber, with portrait and list of his writings, by JAWOROWSKI, Wien. ent. Z. xi. p. 253.—George Haggar; Eut. M. M. xxviii, p. 112.—J. T. Harris; Ent. M. M. xxviii, p. 291.—Dr. Ernst Hofman: Wien. ent. Z. xi, p. 120; Deutsche e. Z. Lep. v, pp. 459-463.—Dr. E. Killias; MT. Schw. ent. Ges. viii, pp. 373-375.—Eugène Lemoro; Bull. Soc. Ent. Fr. lxi, p. xcii. -Sir W. Macleay; P. Linn, Soc. N.S.W. (2) vi, pp. 707-716.-Adrien Maurissen, by Selys-Longchamps, Ann. Ent. Belg. xxxvi, p. 389.— N. G. Moe, by Schöyen, with portrait, Ent. Tidskr. xiii, p. 275.— L'Abbé Léon Provancher: Ent. M. M. xxviii, p. 247; by W. H. H.,

Canad. Ent. xxiv, p. 130.—G. Quedenfeldt, by Kolbe, Berl. Ent. Zeit. xxxvii, pp. 241-246, portrait.—Max Quedenfeldt: Berl. Ent. Zeit. xxxvi, p. 473.—Eugène Revelière, by Rey, Échange, 1892, p. 31; Bull. Soc. Ent. Fr. lxi, p. lxii.—Dr. P. Richter; Berl. Ent. Zeit. xxxvi, p. 472.—Max Saalmüller, by Honrath, Berl. Ent. Zeit xxxvi, pp. 468-471; by Heyden, Deutsche e. Z. Lep. v, pp. 453-459.—Charles Smethurst, with portrait, Naturalist, 1892, p. 145.—Bruno Sperrhaken; Deutsche e. Z. Lep. v, p. 463.—W. Thompson; Ent. xxv, p. 328.—H. Tieffenbach; Kraatz, Deutsche e. Z. 1892, p. 12.—H. W. J. Vaughan: Ent. M. M. xxviii, p. 291; Ent. xxv, p. 300; and Rev. A. H. Wratislaw; Ent. xxv, p. 328.

## VI.—SYSTEMATIC.

N.B.—The reference-numbers collected under the headings of the various groups refer to "Titles," and are intended to give, in each case, a complete list of all memoirs containing anything of importance in connection with the group.

# (A.) COLEOPTERA.

Glandular structures in elytra; HOFFBAUER (405). Early stages of embryology; HENKING (396). Series defined; JACQUET (449).

#### CICINDELIDÆ.

[Cf. Blackburn (95), G. H. Horn (420), W. Horn (421 to 426), Kolbe (504), Péringuey (672, 673), Ruge (788).]

List of some recent additions to the Catalogue of Cicindelidæ; HORN, Deutsche e. Z. 1892, pp. 222-224.

For many suggestions as to synonymy and varietal forms in Cicindelidæ, not here specially alluded to, see Horn, iii Beitrag zur Kenntniss der Cicindeleten, Deutsche e. Z. 1892, pp. 209-219.

Nomenclature of varieties discussed; Beuthin, Ent. Nachr. xviii, p. 290.

Caledonica acentra, Chd., = (barayi, Fauv.); Horn, p. 93, Deutsche e. Z. 1892.

Cicindela: variations in markings of N. American; Horn, Ent. News, iii, p. 25, pl. i. Numerous synonymical remarks, pp. 95-98, C. kaleea, Bates, = (mandarina, Fleut.), p. 95, C. tenuisignata, Lec., = (psilogramma, Bates), p. 97, C. interrupta, Fab., = (circumcincta, Cast.), p. 98; Horn, Deutsche e. Z. 1892. C. tenuicincta and latisignata, notes on; Blaisdell, Zoe, iii, pp. 47 & 48. C. discreta, n. varr. reducta, subfasciata; Horn, p. 370, Deutsche e. Z. 1892. C. flexuosa, variation, with many new names; Beuthin, pp. 333-335, Ent. Nachr. xviii. C. litoralis, variation, with new names; id. pp. 359-362, 376, & 377, t. c.

C. (Cratohærea-Odontochila) cratohæroides, Guinea, p. 220, C. (Odontochila) hennebergorum, Njam-Njam, p. 221, Horn, Deutsche e. Z. 1892; C. junkeri. Njam-Njam, Kolbe, p. 144, S. E. Z. liii; C. jucunda, S. W. Africa, Péringuey, p. 4, Tr. S. African Phil. Soc. vi, 2; C. ganglbaueri, Ceylon, Horn, p. 95, Deutsche e. Z. 1892; C. chevrolati, Yucatan, Horn, p. 371, Deutsche e. Z. 1892;

staudingeri, Sao Paulo, Horn, Deutsche e. Z. 1892, p. 368; C. (n. g.?) paradoxa, Ceylon, p. 75, interposita, p. 76, Philippines, guineensis, N. Guinea, affinis, Venezuela, elegantissima, p. 77, catoptroides, Sumatra, longipalpis, Java, p. 78, gestroi, Sumatra, labioænea, Ceylon, p. 79, seriepunctata, dubia, E. India, tenuilineata, Madagascar, p. 80, nivicinctoides, Malay Archipelago, incerta, Philippines, elongata, Celebes, p. 81, differens, Zanzibar, incognita, New Zealand, p. 82, novaseelandica, p. 83, New Zealand, laphyricornis, Caffraria, p. 84, securiformis, Cape of Good Hope, drakei, Mendoza, p. 85, unicolor, Argentine Mountains, despectata, Philippines, plumigera, Trichinopolis, p. 86, C. (Calochroa) ceylonensis, cyanea (Fab.), Bengal, p. 87, C. prodotiformis, W. Africa, p. 88, Horn, Deutsche e. Z. 1892: n. spp.

- Collyris: many synonymical and varietal observations on species of; HORN, Deutsche e. Z. 1892, pp. 354-368. C. filiformis and cribrosa are one species; HORN, p. 92, t. c.
- C. rugei, Mindanao, p. 354, batesi, E. India, p. 355, thoracica, hab. ?, p. 356, labiomaculata, Borneo, p. 357, angularis, p. 358, erichsoni, p. 359, Philippines, spuria, Sumatra, p. 360, speciosula, Palawan, similis, p. 361, chaudoiri, p. 362, affinis, p. 363, Philippines, fleutiauxi, Borneo, p. 364, kraatzi, Java, roeschkei, Calcutta, p. 365, schaumi, Cornwallis, p. 366, Horn, Deutsche e. Z. 1892, n. spp.

Cosmena intermedia, Caffraria, HORN, p. 68, Deutsche e. Z. 1892, n. sp.

Ctenostoma hirsutum, Brazil, HORN, Deutsche e. Z. 1892, p. 66, n. sp.

Derocrania schaumi, Ceylon, Horn, p. 67, Deutsche e. Z. 1892; D. flavicornis, Ceylon, Horn, p. 92, t. c.: n. spp.

Distipsidera gruti, var., = (plustchevskyi, Dok.), and D. fusciata, Motsch., = (volitans, Macl.); HORN, p. 93, Deutsche e. Z. 1892.

Dromica 5-costata, Natal, HORN, p. 67, Deutsche e. Z. 1892, n. sp.

Euryoda similis, Guinea, p. 70, madagascariensis, Nossi-bé, p. 71, Hohn, Deutsche e. Z. 1892, n. spp.

Heptodonta, Hope, = (Eurytarsa, Dok.), Horn, p. 94, Deutsche e. Z. 1892.

Manticora, the species noticed; Péringuey, p. 3, Tr. S. African Phil. Soc. vi, 2: the true species of; Horn, p. 144, Deutsche e. Z. 1892.

Megacephala denticollis, & character; HORN, p. 144, Deutsche e. Z. 1892.

M. kruperi, W. Africa, HORN, p. 91, Deutsche e. Z. 1892; M. angulicollis,
Victoria Nyanza, KOLBE, p. 145, S. E. Z. liii: n. spp.

Megalomma schaumi, Madagascar, fulgens, Bourbon, p. 73, affine, Madagascar, p. 74, Horn, Deutsche e. Z. 1892, n. spp.

Myrmecoptera schaumi, E. Africa, Horn, p. 220, Deutsche e. Z. 1892; M. erikssoni, S.W. Africa, Péringuey, p. 5, Tr. S. African Phil. Soc. vi, 2; M. limpopoiana, junodi, S. Africa, Péringuey, p. 95, t. c.: n. spp.

Odontochila confusa, Dej., and aperta, Kl., distinctions of; Horn, p. 94, Deutsche e. Z. 1892. O. batesi, n. var. semicincta; Horn, p. 69, Deutsche e. Z. 1892.

O. chrysopyga, W. Africa? Horn, p. 70, Deutsche e. Z. 1892, n. sp.

- Omus hornianus, California? HORN, p. 91, Deutsche e. Z. 1892, n. sp Ophryodera rufo-marginata, n. var. erikssoni; Péringuey, Tr. S. Al Phil. Soc. vi, 2, p. 4.
- O. richteri, W. Africa, HORN, p. 72, Deutsche e. Z. 1892, n. sp.
- Oxychila; Cophognathus is a subgen. only, C. opacipennis = (0. 1 Horn); Horn, p. 143, Deutsche e. Z. 1892.
- O. batesi, Sao Paulo, similis, Rio Grande do Sul, Horn, p. 89, Deuts Z. 1892, n. spp.
- Oxygoniola, n. g., near Oxygonia, for O. chamæleon, n. sp., Celebes, j. HORN, Deutsche e. Z. 1892.
- Togohostoma pusillum, n. var. flavomaculatum; Horn, p. 372, Deuts Z. 1892.
- P. bispinosum, bidentatum, Madagascar, Horn, Deutsche e. Z. 1892, n. spp.
- Styphloderma schaumi, Soudan, HORN, p. 369, Deutsche e. Z. 1892,
- Tetracha lucifera, n. var.? erichsoni; HORN, p. 369, Deutsche e. Z. 189
  sobrina not distinct from sommeri; HORN, p. 144, Deutsche e. Z.
- T. njam-njamensis, Africa, thoracica, Amazone, p. 90, Horn, Deuts
   Z. 1892; T. helmsi, W. Australia, BLACKBURN, p. 16, Tr. R. S
   Austr. xvi; T. horni, Peru, Ruge, p. 130, Deutsche e. Z. 1
   n. spp.
- Tetreurytarsa, n. g., for Euryoda tetrasticta; Honn, p. 94, Deutschi 1892.
- Therates labiatus, coracinus, fulvipennis, bidentatus, caligatus are species; Horn, p. 94, Deutsche e. Z. 1892.
- T. schaumi, Borneo, HORN, p. 69, Deutsche e. Z. 1892, n. sp.
- Thopeutica theratoides, n. var. albapicalis, T. eximia, n. var. fulce Horn, p. 74, Deutsche e. Z. 1892.
- T. schaumi, Celebes, Horn, Deutsche e. Z. p. 75, n. sp.
- Tricondyla aptera, varr. (pedestris and chevrolati), T. cyanea va (wallacei); HORN, p. 93, Deutsche e. Z. 1892. T. cyanea, n paradoxa; HORN, p. 371, Deutsche e. Z. 1892.
- T. frontalis, Philippines, HORN, p. 66, Deutsche e. Z. 1892, n. sp.

# CARABIDÆ.

- [Cf. Abeille (1), Bates (37, 38), Blackburn (90, 91, 93, 95), Duv (230), Fairmaire (262, 267, 269, 270), Frivaldszky (309), Geauer (317, 318), Gestro (330, 332, 334), Horn (417, Péringuey (672, 673), Reitter (724, 725, 728), Rost (780), St (852, 853), Tryon (953).]
- .1 bacetus cyclodes, India, BATES, p. 232, Ann. Ent. Belg. xxxinsolatus, illuminans, p. 359, sulculatus, foveifrons, p. 360, ant tatus, p. 362, Burma, BATES, Ann. Mus. Genov. (2) xii: n. spp
- Acanthogenius infuscatus, Burma, BATES, p. 389, Ann. Mus. Genc xii, n. sp.
- Acinopus picipes, metamorphoses; XAMBEU, p. 200, Le Nat. 1892.

- Acupalpus tessellutus, elegantulus, S.W. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2) p. 22; A. sinuellus, Bhamo, Bates, p. 350, Ann. Mus. Genov. (2) xii: n. spp.
- Agastillus, n. subg. of Pterostichus; REITTER, p. 61, Wien. ent. Z. xi.
- Agonocheila fenestrata, S. Australia, BLACKBURN, p. 80, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Allocota arata, Burma, Bates, p. 425, Ann. Mus. Genov. (2) xii, n. sp.
- Amara: subgg. Celia and Triana, study of N. American forms; HORN, pp. 18-40, Tr. Am. Ent. Soc. xix. A. apricaria, metamorphoses; XAMBEU, Mœurs, ii, p. 13.
- (Triæna) belfragei, Texas, Horn, p. 19, Tr. Am. Ent. Soc. xix; A.
   (Celia) femoralis, p. 30, robustula, p. 32, nupera, p. 33, imitatriæ, p. 34, N. America, Horn, t. c.: n. spp.
- Amblystomus rotundiceps, vittatus, India, Bates, p. 231, Ann. Ent. Belg. xxxvi; A. magnus, p. 333, punctatus, tetrastigma, p. 335, Burma, Bates, Ann. Mus. Genov. (2) xii: n. spp.
- Amblytelus brevis, sinuatus, Australia, BLACKBUEN, p. 87, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Anatrichis birmanica, Rangoon, BATES, p. 325, Ann. Mus. Genov. (2) xii, n. sp.
- Anisodactylus basicollis, Seychelles, FAIRMAIRE, p. cli, Bull. Soc. Ent. Fr. lxi, n. sp.
- Anophthalmus apenninus, fig. 3, vacca, fig. 4, ramorinii, fig. 5, pl. iv, figured; Gestro, Ann. Mus. Genov. (2) xii.
- A. launi, p. 80, pl. iv, fig. 1, bensæ, p. 81, fig. 2, Italy, Gestro, t. c.; A. antoninæ, Ordubad, Reitter, p. 60, Wien. ent. Z. xi: n. spp.
- Anoplogenius planicollis, p. 345, patinalis, p. 346, A. (Hemiaulax) dentipennis, p. 347, Burma, BATES, Ann. Mus. Genov. (2) xii, n. spp.
- Anthia omoplata and limbipennis, varr. of described; Péringuey, pp. 15 & 16, Tr. S. African Phil. Soc. vi (2).
- Antisphodrus leptoderus, Margelan, REITTER, p. 67, Wien. ent. Z. xi, n. sp. Apotomus: characters of the palæarctic species; REITTER, p. 137, Wien. ent. Z. xi.
- A. latigena, Turkestan (but on p. 188 corrected to Algeria), Reitter, l. c.; A. hirsutulus, Rangoon, Bates, p. 287, Ann. Mus. Genov. (2) xii: n. spp.
- Arenozenus, n. g., near Loxandrus, for A. harpaloides, n. sp., Burma; Bates, p. 365, Ann. Mus. Genov. (2) xii.
- Aristolebia, n. g., near Libia, for A. quadridentata, n. sp., Burma; Bates, p. 428, Ann. Mus. Genov. (2) xii.
- Bembidium subviride, Macl., = ocellatum, Blackn.; BLACKBURN, p. 98, P. Linn. Soc. N.S.W. (2) vii.
- B. (Pseudolimnæum) doderoi, Val Pesio, GANGLBAUER, p. 314, Wien. ent. Z. xi; B. (Peryphus) subcylindricum, Taschkent, REITTER, p. 59, Wien. ent. Z. xi; B. xanthotelum, Rangoon, BATES, p. 287, Ann. Mus. Genov. (2) xii: n. spp.
- Blechrus discoidalis, Obock, FAIRMAIRE, p. 83, Rev. d'Ent. zi, n. sp.

- Brachichila rugulipennis, Burma, BATES, p. 406, Ann. Mus. Gei xii, n. sp.
- Brachinus nobilis, Dej., = (Pheropsophus alstoni, Per.); PÉR p. 135, Tr. S. African Phil. Soc. vi (2).
- B. concinnus, flavicapillus, p. 393, exquisitus, p. 394, evanescens, cle p. 395, circumtinctus, intactus, p. 396, Burma, Bates, An Genov. (2) xii, n. spp.
- Brachyonychus lævipennis, n. var. parumpunctatus; BATES, p. 29 Mus. Genov. (2) xii.
- Broscini: notes, taxonomic, synonymic, and critical, on Augenera and species; SLOANE, P. Linn. Soc. N.S.W. (2) 45-64.
- Culleida sultana, p. 421, excelsa, doriæ, p. 422, Burma, BATES, Au Genov. (2) xii, n. spp.
- Curabus cebennicus, n. n. for hispanus, auct., nec Fab.; C. hispanus, e (rutilans, auct.); Vuillefroy, p. 320, Ann. Soc. Ent. C. hispanus, n. var. boudeti; Beuthin, p. 211, Ent. xviii.
- Carenum: synonymical notes on, and localities for, numerous: SLOANE, P. Linu. Soc. N.S.W. (2) vi, pp. 429-432.
- C. ignotus, p. 427, habilis, p. 428, W. Australia, SLOANE, P. Lin N.S.W. (2) vi, n. spp.
- Cusnonia albicolon, graciliceps, 381, zanthe, p. 382, Burma, Ann. Mus. Genov. (2) xii, n. spp.
- Catascopus mirabilis, Burma, BATES, p. 409, Ann. Mus. Genov. (n. sp.
- Chlanioctenus, n. subg. of Chlonius, q. v.; BATES, Ann. Mus. Ger xii, p. 309.
- Chlænius rerecundus, Natal, Péringuey, p. 103, Tr. S. African Phil (2); C. rudicollis, Madagascar, Fairmaire, p. clxix, Bull Soc. I lxi; C. erikssoni, p. 18, ovampo, egregius, p. 19, fraternus, p. 2 Africa, Péringuey, Tr. S. African Phil. Soc. vi (2); C. later p. 82, cyaneo-rufus, p. 83, Obock, Fairmaire, Rev. d'Ent. (Homalolachnus) feanus, p. 308, C. (Chlænioctenus) pectinipes, Burma, eneides, Assam, p. 310, C. bhamoensis, p. 311, an p. 312, privatus, p. 315, delicatus, ochroperas, p. 316, pæcilinus, corrosulus, p. 319, cæruleiceps, p. 320, Burma, Bates, Ant Genov. (2) xii: n. spp.
- Clivina aqualis, description amended; BLACKBURN, p. 83, P. Soc. N.S.W. (2) vii.
- C. sagittaria, p. 275, sectifrons, p. 276, læviceps, p. 277, pileolata, scissa, p. 278, scuticeps, p. 280, occipitalis, costulo p. 281, chlorizans, p. 282, Burma, Bates, Ann. Mus. Genxii; C. helmsi, Australia, Blackburn, p. 22, Tr. R. Soc. S. xvi: n. spp.
- Coleolissus, n. subg. of Hypolithus, q.v.; BATES, p. 338, Ann Genov. (2) xii.

- Colpodes muleyitus, cælitis, p. 372, janellus, p. 373, acroglyptus, castaniventris, p. 374, punctulicollis, undulipennis, p. 375, eucuemis, ischioxanthus, p. 376, spinulifer, p. 377, Burma, Bates, Ann. Mus. Genov. (2) xii, n. spp.
- Coriza? araticeps, Obock, FAIRMAIRE, p. 84, Rev. d'Ent. xi, n. sp.
- Creagris binoculus, Burma, hamaticollis, Borneo, Bates, p. 390, Ann. Mus. Genov. (2) xii, n. sp.
- Cyclothorax eyrensis, S. Australia, BLACKBURN, p. 480, P. Linn. Soc. N.S.W. (2) vi, n. sp.
- Cymindoidea planulata, Burma, BATES, p. 417, Ann. Mus. Genov. (2) xii, n. sp.
- Cypholoba amatonga, S. Africa, Péringuey, p. 102, Tr. S. African Phil. Soc. v (2), n. sp.
- Demetrias tweedensis, N. S. Wales, BLACKBURN, p. 67, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Dendrocellus gestroi, Burma, BATES, p. 121, Ann. Mus. Genov. (2) xii, n. sp.
- Dicalus splendidus, larva and pupa described, and figured; Wickham, p. 195, Bull. Lab. Iowa, ii.
- Dischiesus alaticollis, Tenasserim, BATES, p. 302, Ann. Mus. Genov. (2) xii, n. sp.
- Dolichoctis rutilipennis, p. 412, angusticollis, p. 413, incerta, iridea, p. 414, expansicollis, p. 415, Burma, BATES, Ann. Mus. Genov. (2) xii, n. spp.
- Dromius yarra, Victoria, BLACKBURN, p. 71, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Drypta siderea, p. 382, fimbriata, p. 384, Burma, Bates, Ann. Mus. Genov. (2) xii, n. spp.
- Dyschirius aneobrachialis, Burma, BATES, p. 283, Ann. Mus. Genov. (2) xii, n. sp.
- Dystrichothorax, n. g., near Amblytelus, for D. bicolor, p. 88, sloanei, bipunctutus, p. 89, lividus, vicinus, p. 90, n. spp., Australia, Blackburn, P. Linn. Soc. N.S.W. (2) vii.
- Ecnomolaus, n. g., Trigonotomini, p. 357, for E. clivinoides, n. sp., Burma, p. 358, Bates, Ann. Mus. Genov. (2) xii.
- Ectroma beneficum, variation and synonymy discussed; BLACKBURN, p. 73, P. Linn. Soc. N.S.W. (2) vii.
- E. obsoletum, N. S. Wales, id. t. c. p. 72, n. sp.
- Elaphrus dauricus and sibiricus, distinctions noticed; Reitter, p. 188, Wien. ent. Z. xi.
- Endynomena discoidulis, Burma, Bates, p. 423, Ann. Mus. Genov. (2) xii, n. sp.
- Epelyx, n. g., near Amblytelus, for E. lindensis, latus, n. spp., Australia, Blackburn, p. 92, P. Linn. Soc. N.S.W. (2) vii.
- Epicosmus breviformis, mandarinellus, p. 299, latigenis, p. 300, brevisternis, p. 301, gracilipes, p. 302, Burms, Bates, Ann. Mus. Genov. (2) xii, n. spp.

- Eudromus opacus, Diego-Suares, Fairmaire, p. cix, Bull. Soc. En lxi, n. sp.
- Euryaptus, n. g., near Trigonotoma, for E. nigellus, Burma, p. assamensis, Assam, rufipes, Andaman Is., p. 356, n. spp.; Bates Mus. Genov. (2) xii.
- Eurydera rotundata, Madagascar, FAIRMAIRE, p. clxix, Bull. Soc. E lxi, n. sp.
- Eurylychnus victoriæ, Australia, Sloane, p. 52, P. Linn. Soc. N.S. vii, n. sp.
- Euryscaphus sulcicollis, p. 81, chaudoiri, p. 82, Australia; BLACE P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Galerita birmanica, p. 121, feæ, p. 122, Burma, Bates, Ann. Mus. (
  (2) xii, n. spp.
- Gigadema eremita, p. 17, mandibulare, p. 18, Australia, BLACKBUI B. Soc. S. Austr. xv, n. spp.
- Gnathaphanus acutipennis, p. 328, exaratus, rufitactor, p. 329, I BATES, Ann. Mus. Genov. (2) xii; G. aridus, Australia, BLACI p. 22, Tr. R. Soc. S. Austr. xvi: n. spp.
- Graphipterus lugens, Chaud., = (univittatus, Per.); Péringuey, Tr. S. African Phil. Soc. vi, 2.
- G. grandis, p. 7, ornatus, p. 8, vicinis, deceptor, ovipennis, p. 9, agilis simulator, amicus, lætus, p. 11, erikssoni, p. 12, S. W. Africa, I Guey, Tr. S. African Phil. Soc. vi, 2; G. mimus, p. 98, egregiu bezianus, p. 99, fraternus, p. 100, S. Africa, Péringuey, t. c.:
- Haplogaster ampliatus, Burma, BATES, p. 270, Ann. Mus. Genov. (n. sp.
- Harpaliscus, n. g., near Harpalus, p. 340, for H. birmanicus, n. sp., I p. 341; BATES, Ann. Mus. Genov. (2) xii.
- Harpalus? karennius, Burma, BATES, p. 341, Ann. Mus. Genov. (n. sp.
- Hemiaulux, n. subg. of Anoplogenius, q. v.; BATES, Ann. Mus. (2) xii, p. 347.
- Hemichlænius, n. g., near Chlænius, p. 307, for H. microspilus, Burma, p. 308, Bates, Ann. Mus. Genov. (2) xii, n. sp.
- Homethes angulatus, p. 68, gracilis, parvicollis, p. 69, rotundatus, Australia, Blackburn, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Hoplolenus atronitens, Madagascar, FAIRMAIRE, p. clxx, Bull. Soc Fr. lxi, n. sp.
- Hydroporomorpha obockiana, Africa, FAIRMAIRE, p. 86, Rev. d'E n. sp.
- Hypharpax australis and inornatus distinguished; BLACKBURN, P. Linn, Soc. N.S.W. (2) vii.
- H. obsoletus, W. Australia, id. t. c. p. 84, n. sp.
- Hypolithus ornatus, S. W. Africa, Péringuey, p. 21, Tr. S. Africar Soc. vi, 2; H. subtilis, H. (Coleolissus), lamprotus, p. 338, bicol p. 339, viridellus, eulamprus, p. 340, Burma, Bates, Ann. Mus. ( (2) xii, n. spp.

- Hystrichopus agilis, altecostatus, p. 97, porrectus, p. 98, S. Africa, Péringuey, Tr. S. African Phil. Soc. vi, 2, n. spp.
- Lachnoderma? biguttatum, Burma, BATES, p. 424, Ann. Mus. Genov. (2) xii, n. sp.
- Lebia fulvipes, France, Jacquet, p. 2, Échange, 1887, No. 35, and distinctive characters, Pic, Échange, 1891, p. 136; L. chrysis, Taschkent, Beitter, p. 27, Wien. ent. Z. xi; L. mashuana, Mashuanaland, PÉRINGUEY, p. 96, Tr. S. African Phil. Soc. vi, 2; L. cardoni, Bengal, Bates, p. 233, Ann. Ent. Belg. xxxvi; L. karenia, p. 426, maharani, p. 427, Burma, Bates, Ann. Mus. Genov. (2) xii: n. spp.
- Lecamomerus major, striatus, N. S. Wales, BLACKBURN, p. 479, P. Linn. Soc. N.S.W. (2) vi, n. spp.
- Leptorembus kolbei, Congo, Duvivier, p. 56, Ann. Ent. Belg. xxxvi, n. sp.
- Loxandrus, characters and components noticed; Blackburn, p. 95, P. Linn. Soc. N.S.W. (2) vii.
- L. lericollis, p. 96, micans, p. 97, Australia, BLACKBURN, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Lychnus ater, Putz., = (strangulatus, Bates); SLOANE, p. 57, P. Linn. Soc. N.S.W. (2) vii.
- Lymnastis pilosus, p. 296, atricapillus, p. 297, Burma, BATES, Ann. Mus. Genov. (2) xii, n. spp.
- Macrochilus ferrugiaipes, Madagascar, FAIRMAIRE, p. clxviii, Bull. Soc. Ent. Fr. lxi, n. sp.
- Masoreus (Æphnidius) submaculatus, Burma, Bates, p. 405, Ann. Mus. Genov. (2) xii, n. sp.
- Mastax rugiceps, p. 397, alveolatus, carissimus, p. 398, gestroi, p. 399, Burma, Bates, Ann. Mus. Genov. (2) xii, n. spp.
- Mecodema blagravii, characters of, and description; SLOANE, p. 51, P. Linn. Soc. N.S.W. (2) vii.
- Megalonychus birmanicus, dilaticollis, Burma, p. 369, cyanipennis, Assam, p. 370, Bates, Ann. Mus. Genov. (2) xii, n. spp.
- Metabacetus, n. g., near Pediomorphus, for M. immarginatus, n. sp., Burms, Bates, p. 364, Ann. Mus. Genov. (2) xii.
- Metabletus subvittatus, p. 417, cymindulus, p. 418, Burma, Bates, Ann. Mus. Genov. (2) xii, n. spp.
- Microferonia adelaidæ, referred to Anchomenini; BLACKBURN, p. 95, P. Linn. Soc. N.S.W. (2) vii.
- Monacanthonyx, n. g., near Culathus, for M. pocillator, n. sp., Burma; Bates, p. 367, Ann. Mus. Genov. (2) xii.
- Mormolyce: parasitic fungus on; GIARD (338), and Bull. Soc. Ent. Fr. lxi, p. lx.
- Nanochlænius, n. g., Chlæniinæ, p. 321, for N. feæ, n. sp., Burma, p. 322; BATES, Ann. Mus. Genov. (2) xii.
- Nebria araschinica, Ordubad, REITTER, p. 66, Wien. ent. Z. xi, n. sp. Nematopeza auberti, Obock, FAIRMAIRE, p. 84, Rev. d'Ent. x, n. sp.

- Notiophilus melanophthalmus, Schlosser, characters of; HEYDEN, Wien. ent. Z. xi.
- Omophron: list of the described species; Gestro, pp. 953 & 964 Mus. Genov. (2) x.
- O. oberthurii, Burma, GESTRO, p. 962, t. c., n. sp.
- Onycholabis melitopus, Burma, Bates, p. 371, Ann. Mus. Genov. n. sp.
- Oodes conspicuus, Transvaal, Péringuey, p. 20, Tr. S. African Pt vi (2); O. cribristernis, pequensis, p. 323, rhodopus, p. 324, I Bates, Ann. Mus. Genov. (2) xii: n. spp.
- Ophionea bhamoensis, Burma, BATES, p. 380, Ann. Mus. Genov. n. sp.
- Ophoniscus, n. g., near Ophonus, for O. iridulus, p. 337, cribrifron. lithoides, p. 338, Burma, Bates, Ann. Mus. Genov. (2) xii, n.
- Orthogonius quadricollis, p. 400, dispar, p. 401, rustventris, p. 402, latus, p. 403, Burma, Bates, Ann. Mus. Genov. (2) xii, n. spr
- Orthotrichus alternatus, indicus, Burma, Bates, p. 368, Ann. Mus. (2) xii, n. spp.
- Oxycentrus acutulus, omaseoides, Burma, BATES, p. 344, Ann. Mus. (2) xii, n. spp.
- Oxydrepanus birmanicus, Tenasserim, Bates, p. 283, Ann. Mus. (2) xii, n. sp.
- Pentagonica daimiella, Japan, Bates, p. 426, Ann. Mus. Genov. n. sp.
- Percolestus, n. g., Broscini, p. 54, for P. blackburni, n. sp., Victori SLOANE, P. Linn. Soc. N.S.W. (2) vii.
- Percosoma: the species tabulated; SLOANE, p. 58, P. Linn. Soc. (2) vii.
- P. concolor, Victoria, SLOANE, t. c. p. 61, n. sp.
- Perigona nigriceps and pallipennis, distinctive characters; Hor: Tr. Am. Ent. Soc. xix.
- P. bigener, Burma, Bates, p. 379, Ann. Mus. Genov. (2) xii, n. sp. Perileptus melanopygus, Obock, Fairmaire, p. 87, Rev. d'l n. sp.
- Pheropsophus acutecostatus, Madagascar, FAIRMAIRE, p. claviii, Bi Ent. Fr. lxi; P. nanodes, Assam, p. 391, infantulus, Burma, BATES, Ann. Mus. Genov. (2) xii: n. spp.
- Philophlæus pygmæus, ornatus, Australia, Blackburn, p. 19, Tr. S. Austr. xvi; P. monticola, p. 75, sydneyensis, p. 76, laticolli confertus, p. 78, Australia, Blackburn, P. Linn. Soc. N.S.W. (n. spp.
- Piezia albolineata, Wallengr, = (albosignata, Pér.); Péringuey, Tr. S. African Phil. Soc. vi (2).
- P. ovampoensis, S.W. Africa, PÉRINGUEY, Tr. S. African Phil. Soc p. 13, n. sp.
- Pirantillus extensicollis, Burma, Bates, p. 370, Ann. Mus. Genov. n. sp.

- Platymetopus longulus, p. 330, P.? erebius, edentatus, sublavis, p. 331, grandiceps, p. 332, amariformis, p. 333, Burma, gnathaphanoides, Assam, p. 332, Bates, Ann. Mus. Genov. (2) xii, n. spp.
- Platynus reflexus, variation; HAMILTON, p. 157, Canad. Ent. xxiv.
- P. myrmecodes, arizonensis, lunguidus, Arizona, HORN, p. 42, Tr. Am. Ent. Soc. xix, n. spp.
- Plectes platessa, Motsch., noticed, biebersteini, n. var. adelphus; Rost, Deutsche e. Z. 1892, p. 142.
- P. polychrous, Caucasus, Rost, p. 401, Deutsche e. Z. 1892, n. sp.
- Plochionus timidus, habits; MURTFELDT, Canad. Ent. xxiv, p. 279.
- Pacilus ornatus, New Guinea, TRYON (953), n. sp.
- Pogonoglossus carinipennis, Burma, BATES, p. 388, Ann. Mus. Genov. (2) xii, n. sp.
- Polyhirma somalica, Somaliland, Gestro, p. 751, Ann. Mus. Genov. (2) xii; P. edax, p. 13, alstoni, p. 14, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2); P. chaudoiri, p. 100, rutata, p. 101, S. Africa, Péringuey, t. c. : n. spp.
- Pristomachærus quadristigma, p. 303, rubellus, eucharis, p. 305, lebioides, cauliops, p. 306, Burma, Bates, Ann. Mus. Genov. (2) xii, n. spp.
- Promecoderus castelnaui, p. 46, ambiguus, p. 49, Australia, SLOANE, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Pseudotaphoxenus gracilicornis, China, FRIVALDSZKY, p.122, Term. füzetek, xv, n. sp.
- Pterostichus amethystinus, brunneus, castaneus, & characters; brunneus = (scutellaris, Lec.), mancus and diligendus, synonymy; mutus, Say, = (pulrinatus and stenops, Haus.); Horn, p. 41, Tr. Am. Ent. Soc. xix.
- P. (Agastillus, n. subg.) cucujinus, Ordubad, REITTER, p. 61, Wien. ent. Z. xi, n. sp.
- Rhembus rectificatus, p. 325, lærigatus, colossus, p. 326, Burma, BATES, Ann. Mus. Genov. (2) xii, n. spp.
- Rhytisternus splendens, p. 93, cardwellensis, p. 94, Australia, BLACKBURN, P. Linn. Soc. N.S. W. (2) vii, n. spp.
- Surothrocrepis posticalis and calida, distinction from corticalis queried; BLACKBURN, p. 71, P. Linn. Soc. N.S.W. (2) vii.
- Sarticus obscurus, brevicornis, Australia, Blackburn, p. 23, Tr. R. Soc. S. Austr. xvi, n. spp.
- Scotodipnus mayeti, Alpes Maritimes, ABEILLE, p. 62, Rev. d'Ent. xi, n. sp. Siagona simplex, p. 6, australis, p. 7, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2); S. subtilis, angulifrons, p. 284, angustipennis, p. 286, Burma, Bates, Ann. Mus. Genov. (2) xii: n. spp.
- Simous lumpros, Burma, BATES, p. 322, Ann. Mus. Genov. (2) xii, n. sp. Simurus nitidus, p. 407, graciliceps, p. 408, Burma, BATES, Ann. Mus. Genov. (2) xii, n. spp.
- Stenolophus harpaloides, p. 347, rectifrons, p. 348, charis, p. 349, Burma, Bates, Ann. Mus. Genov. (2) xii, n. spp.
- Stereostoma meridionalis, Transvaal, Péringuey, p. 103, Tr. S. African Phil. Soc. vi (2), n. sp.

- Stomonaxus dilaticollis, p. 351, inermis, p. 352, Burma, BATES, An Genov. (2) xii, n. spp.
- Tuchys rittatus, Motsch., redescribed; REITTER, p. 67, Wien. ent T. sellatus, Obock, FAIRMAIRE, p. 87, Rev. d'Ent. xi; T. prolixus, 1 p. 288, euryodes, obsolescens, haliploides, p. 289, remotiporis transumbratus, T. (Barytachys) callispilotus, p. 291, feanus, c p. 292, expansicollis, p. 293, mirabilis, p. 294, nietneri = (Niet.), p. 295, unitarius, p. 296, Burma, BATES, Ann. Mus. Ge xii; T. yarrensis, Victoria, BLACKBURN, p. 20, Tr. R. Soc. 8 xv: n. spp.
- Taniolobus birmanicus, Teinzo, Bates, p. 270, Ann. Mus. Genov. n. sp.
- Taridius birmanicus, Burma, BATES, p. 416, Ann. Mus. Genov n. sp.
- Tetrugonoderus dispur, S. W. Africa, Péringuey, Tr. S. Afric Soc. vi (2), p. 6, n. sp.
- Trechus (Anophthalmus) speiseri, Herzegovina, GANGLBAUER, Wien. ent. Z. xi; T. angelicæ, Ordubad, REITTER, p. 60, W Z. xi; T. birmanicus, p. 297, cauliops, p. 298, Burma, BAT Mus. Genov. (2) xii; T. barbaræ, California, HORN, p. 41, Ent. Soc. xix: n. spp.
- Triwnogenius obockianus, Africa, FAIRMAIRE, p. 85, Rev. d n. sp.
- Trichocarenum, n.g., near Eutoma, p. 20, for T. elderi, n. sp., Desert, p. 21; BLACKBURN, Tr. R. Soc. S. Austr. xvi.
- Trichotichnus birmanicus, Burma, BATES, p. 342, Ann. Mus. Gxii, n. sp.
- Trigonognatha feana, Burma, BATES, p. 365, Ann. Mus. Genov. n. sp.
- Trigonothops longiplaga, pacifica, are probably one species; BLA p. 65, P. Linn. Soc. N.S.W. (2) vii.
- Trigonotoma lamprodera, igneicollis, p. 353, bhamoensis, c p. 354, Burma, iodes, Assam, p. 353, Bates, Ann. Mus. (2) xii, n. spp.
- Triplogenius feanus, p. 356, pequensis, p. 357, Burma, Bates, At Genov. (2) xii, n. spp.
- Zeloticus, n. g., near Chlanius, for Z. tristis, n. sp., S. W. Africa Gury, Tr. S. African Phil. Soc. vi, 2, p. 17, n. sp.
- Zuphium formosum, p. 386, præstans, p. 387, Burma, BATES, An Genov. (2) xii, n. spp.

# AMPHIZOIDE, HALIPLIDE, DYTISCIDE.

[Cf. Hubbard (437, 438), Regimber (720, 721, 722, 723).]

Amphizoa lecontei, imago figured, larva fig., and description Bard, Ins. Life, v, pp. 19-22: larva described and Hubbard, P. E. Soc. Wash. ii, pp. 341-346, pl. iii.

- Huliplus angustifrons, Bengal, RÉGIMBART, p. 112, Ann. Ent. Belg. xxxvi; H. signatipennis, ferruginipes, New Guinea, RÉGIMBART, p. 979, Ann. Mus. Genov. (2) x: n. spp.
- Bidessus guttulatus, Burma, RÉGIMBART, p. 542, Ann. Mus. Genov. (2) x;
  B. loriæ, p. 989, neoguineensis, p. 990, New Guinea, RÉGIMBART,
  Ann. Mus. Genov. (2) x; B. severini, p. 116, indicus, p. 117,
  consanguineus, antematus, p. 118, inconstans, p. 119, livens, p. 120,
  Bengal, RÉGIMBART, Ann. Ent. Belg. xxxvi: n. spp.
- Copelatus speciosus, n. n. for formosus, Rég.; RÉGIMBART, p. 991, Ann. Mus. Genov. (2) x.
- C. gestroi, New Guinea, REGIMBART, p. 991, Ann. Mus. Genov. (2) x, n. sp.
- Cybister owas, n. var. congoana, Duvivier, p. 272, Ann. Ent. Belg. xxxvi. C. hypomelas, New Guines, Régimbart, p. 994, Ann. Mus. Genov. (2), x, n. sp.
- Dytiscus marginalis, variation in coxal processes; GRIFFINI, Boll. Mus. Tor. vii, No. 115.
- Hydaticus epipleuricus, Burma, REGIMBART, p. 545, Ann. Mus. Genov. (2) x; H. suffusus, p. 992, rhantaticoides, p. 993, New Guinea, REGIMBART, Ann. Mus. Genov. (2) x: n. spp.
- Hydrocoptus opatrinus, p. 11, bosschæ, p. 12, Borneo, Régimbart, Notes Leyd. Mus. xiv, n. spp.
- Hydrovatus pinguis, Bengal, Régimbart, p. 114, Ann. Ent. Belg. xxxvi, n. sp.
- Hyphoporus severini, Bengal, RÉGIMBART, p. 115, Ann. Ent. Belg. xxxvi, n. sp.
- Hyphydrus lyratus, n. var. foveolatus; Régimbart, p. 986, Ann. Mus. Genov. (2) x.
- H. flavicans, Bengal, Régimbart, Ann. Ent. Belg. xxxvi, p. 115; H. loriæ, p. 985, gibbicollis, p. 986, curvipes, p. 987, New Guinea, Régimbart, Ann. Mus. Genov. (2) x: n. spp.
- Lacconectes lividus, Burms, RÉGIMBART, p. 544, Ann. Mus. Genov. (2) x, n. sp.
- Laccophilus parvulus, variation noticed; RÉGIMBART, p. 113, Ann. Ent. Belg. xxxvi.
- L. birmanicus, p. 538, planitarsis, p. 539, Burma, Régimbart, Ann. Mus. Genov. (2) x; L. univittatus, auriculatus, New Guinea, Régimbart, p. 981, t. c. : n. spp.
- Mucroporus piceatus, New Guinea, RÉGIMBART, p. 984, Ann. Mus. Genov. (2) x, n. sp.
- Neptosternus speciosus, p. 982, bicinctus, p. 983, New Guinea, REGIMBART, Ann. Mus. Genov. (2) x, n. spp.

#### GYRINIDÆ.

#### [Cf. Régimbart (720, 721).]

Dineutes neoguineensis, New Guinea, REGIMBART, p. 996, Ann. Mus. Genov. (2) x, n. sp.

- Gyrinus smaragdinus, Burma, Régimbart, p. 547, Ann. Mus. Geno x, n. sp.
- Orectochilus apicalis, p. 548, birmanicus, p. 550, cribratellus, p. villosovittatus, p. 552, tomentosus, p. 553, Burma, Régimbart n. spp.
- Porrhorrhynchus depressus, New Guinea, REGIMBART, p. 996, t. c., n

### HYDROPHILIDÆ.

- [G. Bedel (47), Fairmaire (262, 263), Guillebeau (374, Kuwert (402), Péringuey (672, 673).]
- Berosus majusculus and externespinosus, characters of; Blaci p. 207, Tr. R. Soc. S. Austr. xv.
- B. fuscostriatus, p. 87, immaculicollis, p. 88, Obock, FAIRMAIRE d'Ent. xi, n. spp.
- Dibolocelus, n. subg. of Stethoxus; BEDEL, p. 308, Rev. d'Ent. x.
- Eccoptometopus, n.g., near Ochthebius, p. 104, for E. sculpticollis, p. 105, proximus, p. 106, n. spp., S. Africa; Péringury, African Phil. Soc. vi (2).
- Helophorus punctato-sulcatus, Turkestan, Kuwert, p. 102, Deut Z. 1892, n. sp.
- Hydrana capicola, p. 106, extrema, p. 107, S. Africa, Péringuey African Phil. Soc. vi (2), n. spp.
- Hydrobaticus luridus and tristis are one species; BLACKBURN, p. Linn. Soc. N.S.W. (2) vii.
- Hydrochus capensis, Cape Colony, Péringuey, p. 104, Tr. S. . Phil. Soc. vi (2), n. sp.
- Hydrophilus, revision of (under the name Stethozus); BEDEI d'Ent. x, pp. 306-322: H. gansuensis, characters of; SEI p. 163, Wien. ent. Z. xi.
- Hydrous dichromus, Syria, FAIRMAIRE, p. 144, Ann. Ent. Belg. n. sp.
- Limnobius theryi, Algeria, Guillebeau, p. 324, Rev. d'Ent. (Tricholimnobius) grouvellei, Spain, Guillebeau, p. exxxiii, B Ent. Fr. lxi: n. spp.
- Octhebius turkestanicus, Alai Mts., Kuwert, p. 102, Deutsch 1892, n. sp.
- Philydrus suturalis, note on; Sharp, Ent. M. M. xxviiii, p. 1 maculiceps, lavigatus, distinctive characters; Blackburn, Tr. R. Soc. S. Austr. xv.
- Spercheus interruptus, Obock, FAIRMAIRE, p. 88, Rev. d'E S. australis, S. Africa, Péringuey, p. 23, Tr. S. African P vi (2); S. capicola, p. 107, algoensis, p. 108, S. Africa, Pér Tr. S. African Phil. Soc. vi (2): n. spp.
- Stethoxus, rufo-cinctus, Hindostan, p. 309, cavisternum, Tonkin, indicus, Hindostan, p. 310, pedipalpus, Australia, p. 312, si Guyana, p. 313, BEDEL, Rev. d'Ent. x, n. spp.

#### STAPHYLINIDÆ.

- [Cf. Blackburn (93), Dohrn (221), Eppelsheim (255, 256), Fairmaibe (262), Gestro (334), Ganglbauer (319), Reitter (725), Sharp (844), Wasmann (987).]
- Achenium hauseri, Djizak, EPPELSHEIM, p. 336, Deutsche e. Z. 1892, n. sp. Amphicroum adelaidæ, S. Australia, BLACKBURN, p. 23, Tr. R. Soc. S. Austr. xv, n. sp.
- Anthobium sareptanum, Epp., = (starcki, Reit.); Reitter, p. 26, Wien. ent. Z. xi.
- Ateneles pubicollis, n. var. foreli; WASMANN, p. 351, Deutsche e. Z. 1892. Atheta (Hydrosmecta) tenuissima, Europe, Eppelsheim, p. 292, Wien. ent. Z. xi; A. (Homalota) flavida, p. 322, A. (Metaxya) blanda, p. 323, concolor, p. 324, A. (Microdota) turanica, p. 325, Turkestan, Eppelsheim, Deutsche e. Z. 1892: n. spp.
- Belonuchus nullicedo, Philippines, Dohrn, p. 74, S. E. Z. liii, n. sp.
- Bledius simulator, p. 344, immaturus, p. 345, Turkestan, Eppelsheim, Deutsche e. Z. 1892; B. lividipes, Obock, Fairmaire, p. 90, Rev. d'Ent. xi: n. spp.
- Boreaphilus carinthiacus, Europe, GANGLBAUER, p. 17, Soc. Ent. vii, n. sp. Conurus pubescens, n. var. decurtatus; EPPELSHEIM, p. 328, Deutsche e. Z. 1892.
- Coproporus lateimpressus, Margelan, EPPELSHEIM, p. 327, Deutsche e. Z. 1892, n. sp.
- Dilacra fleischeri, Austria, EPPELSHEIM, p. 293, Wien. ent. Z. xi, n. sp.
- Eleusis parva, N. S. Wales, BLACKBURN, p. 24, Tr. R. Soc. S. Austr. xv, n. sp.
- Euryusa laticollis, habits; WASMANN, p. 350, Deutsche e. Z. 1892.
- Fulagria subænea, Taschkent, EPPELSHEIM, p. 326, Deutsche e. Z. 1892, n. sp.
- Geodromicus, distinctions of the sexes; EPPELSHEIM, p. 49, Soc. Ent. vii. Geostiba apfelbecki, Bosnia, EPPELSHEIM, p. 289, Wien. ent. Z. xi, n. sp.
- Hyperomma abnorme, Victoria, BLACKBURN, p. 22, Tr. R. Soc. S. Austr.
- Lathrobium (Lobathrium) cognatum, Turcomania, EPPELSHEIM, p. 355, Deutsche e. Z. 1892, n. sp.
- Leptusa (Sipalia) bosnica, Bosnia, EPPELSHEIM, p. 295, Wien. ent. Z. xi, n. sp.
- Liogluta montivagans, Carpathians, EPPELSHEIM, p. 291, Wien. ent. Z. xi, n. sp.
- Liparocephalus, referred to Aleocharini; CASEY, p. 711, Ann. N. York Ac. vi. L. brevipennis and cordicollis, probably one species; HAMILTON, p. 157, Canad. Ent. xxiv.
- Lomechusa strumosa, its relations with various kinds of ants discussed; Wasmann, Biol. Centralbl. xii, pp. 584, &c.
- Medon propinquus, n. var. nigrinus; EPPELSHEIM, p. 337, Deutsche e. Z. 1892.

- Microglossa lederi, Taschkent, EPPELSHEIM, p. 321, Deutsche e. Z. n. sp.
- Mycetoporus reitteri, Tatra Mts., EPPELSHEIM, p. 296, Wien. ent. M. rubricus, Turkestan, EPPELSHEIM, p. 329, Deutsche e. Z. n. spp.
- Myrmedonia afra, Somaliland, Gestro, p. 753, Ann. Mus. Genov. (n. sp.
- Myrmocia fussi, habits; WASMANN, Deutsche e. Z. 1892, p. 348.
- Niphetodes apfelbecki, Carpathians, GANGLBAUER, p. 17, Soc. E n. sp.
- Oxytelus sexualis, p. 342, subtilis, p. 343, Taschkent, EPPELSHEIM, De e. Z. 1892, n. sp.
- Pæderus debilior, Turkestan, EPPELSHEIM, p. 339, Deutsche e. Z n. sp.
- Philonthus variabilis, Taschkent, EPPELSHEIM, p. 333, Deutsche e. Z. P. flavicornis, Obock, FAIRMAIRE, p. 89, Rev. d'Ent. xi: n. sp. Philorinum hopfigarteni, Siebenburgen, EPPELSHEIM, p. 297, Wie Z. xi, n. sp.
- Platysthetus depravatus, Taschkent, EPPELSHEIM, p. 341, Deutsch 1892, n. sp.
- Protinus limbatus, metamorphoses; XAMBEU, Mœurs, ii, p. 7.
- Quedius crassus, p. 4, cinctus, p. 6, metamorphoses; Xambeu, Moeu Q. capitalis, p. 329, novus, p. 331, Turkestan, Eppelsheim, Deutsch 1892; Q. pictipennis, Victoria, Blackburn, p. 21, Tr. R. S. Austr. xv: n. spp.
- Scimbalium grandiceps, Margelan, REITTER, p. 68, Wien. ent. n. sp.
- Scopæus similis, Taschkent, EPPELSHEIM, p. 337, Deutsche e. Z. S. aliiceps, Obock, FAIRMAIRE, p. 89, Rev. d'Ent. xi; S. fem N. S. Wales, BLACKBURN, p. 22, Tr. R. Soc. S. Austr. xv; n.
- Stenus rugifer, Caa., = (anastomozans), vexatus = (insularis), placi (tumicollis), villosus = (jejunus), milleporus = (sectilifer), diffic (tenuis), nanus = (nanulus), humilis, Er., = (mammops), rigi. (ageus), brumalis = (pauperculus), gratiosus = (hirsutus), Grav., = (haplus and indistinctus, Cas.), umbratilis = (frate pollens = (patens), reconditus = (propinquus), callosus = (va punctatus = (dilutus and obsoletus), hubbardi = (simiolus), l = (leviceps, politulus); Casey, Ann. N. York Ac. vi. p. 711.
- S. posticalis, Taschkent, EPPELSHEIM, Deutsche e. Z. 1892, p. n. sp.
- Tachinus laticollis, larva described; REY, p. clxxxii, Bull. Soc. E1
- Tachyporus nitidulus, n. var. crux; EPPELSHEIM, p. 328, Deutsche 1892.
- Trygæus javanicus, Java, Sharp, p. 61, Notes Leyd. Mus. xiv, n. sp. Xantholinus ochripennis, Turkestan, Eppelsheim, p. 334, Deutsche 1892, n. sp.

# PSELAPHIDÆ.

- [Cf. Brendel (120, 121), Dodero (219), Grilat (365), Guillebeau (376, 379), Pic (687), Raffray (710), Sharp (843), Xambeu (1021).]
- Dimorphic males noticed; Reitter, p. 26, Wien. ent. Z. xi.
- Batrisus adnexus, oculatus, habits; Wasmann, Deutsche e. Z. 1892, p. 350.
- Bibloporus pini, p. clxxxiii, ultimus, p. clxxxiv, Corsica, Guillebeau, Bull. Soc. Ent. Fr. lxi, n. spp.
- Bythinus carinula, Lyons, Rey, Échange, No. 42, 1888, p. 4, n. sp. (other doubtful n. spp. or varr. mentioned with names, l. c.); B. ravouxi, France, Grilat, Échange, No. 58, 1889, p. 78 (but cf. t. c. p. 84); B. croissandeaui, Algeria, Pic, p. cclxxii, Bull. Soc. Ent. Fr. lxi: n. spp.
- Chennium bituberculatum, larva described; XAMBEU, p. 135, Ann. Soc. L. Lyon, xxxix.
- Eudranes, n. g. (Tyrini), p. 241, for E. carinatus, n. sp., N. W. Australia, p. 242; Sharp, Ent. M. M. xxviii.
- Eulasinus, n. g. (Tyrini), for E. walkeri, n. sp., Chusan; Sharp, p. 240, Ent. M. M. xxviii.
- Rhexidius trogasteroides, p. 12, intermedius, p. 13, N. America, BRENDEL, Ent. News, iii, n. spp.
- Trimium durum, costale, p. 166, gracile, laticolle, majus, p. 167, N. America, Brendel, Tr. Am. Ent. Soc. xix, n. spp.
- Trogaster doriæ, Busalla, Dodero, p. 666, Ann. Mus. Genov. (2) x, n. sp. Tychus striola, France, Guillebeau, p. 1, Supp. to Échange, No. 19, 1886.

#### SCYDMÆNIDÆ.

- Cephennium majus, Reitt., and laticolle, Aubé, distinctive characters of; REITTER, p. 137, Wien. ent. Z. xi.
- Neuraphes panormitanus, Sicily, RAGUSA, p. 80, Nat. Sicil. xi, n. sp.

#### Paussidæ.

- [Cf. Blackburn (93), Gestro (333), Kraatz (513, 517), Péringuey (673), Raffray (709), Wasmann (985), Xambeu (1021).]
- Arthropterus foveipennis, S. Australia, p. 24, occidentalis, W. Australia, p. 25, BLACKBURN, Tr. R. Soc. S. Austr. xv, n. spp.
- Pentaplatarthrus paussoides, anatomy, structure of detonating organs; RAFFRAY, N. Arch. Mus. (3) iv, pp. 91-100, pl. xiii.
- Paussus favieri, larva described; XAMBEU, p. 137, Ann. Soc. L. Lyon, xxxix.
- P. cervinus, Madagascar, Kraatz, Deutsche e. Z. 1892, p. 8; P. opacus, Madagascar, Kraatz, t. c. p. 377; P. spinicola, Somaliland, Wasmann, p. 355, MT. Schw. ent. Ges. viii; P. fullax, Transvaal, Péringuey, p. 108, Tr. S. African Phil. Soc. vi, 2: n. spp.
- Pretopaussus, n. g., p. 706, for P. feæ, n. sp., Burma, p. 707; Gastro, Ann. Mus. Genov. (2) xii.

#### SILPHIDE.

- [Cf. Blackburn (93), Fairmaire (263), Guillebeau (375, 377), 1 (418), Péringuey (673), Xambeu (1021).]
- Pinodytini and Lyrosomini, distinctive characters; Horn, p. 44, Tr. Ent. Soc. xix.
- Anemadus strigosus, habits; WASMANN, Deutsche e. Z. 1892, p. 350. Calyptomerus capensis, S. Africa, Péringuey, p. 109, Tr. S. African Soc. vi, 2, n. sp.
- Cutops fuscipes, Men., characters noticed; Reitter, p. 240, ent. Z. xi.
- C. hervei, France, Guillebeau, p. 116, Échange, 1891, n. sp.
- Cutopsimorphus pilosus, larva; XAMBEU, p. 151, Ann. Soc. L. Lyon, 2
- Cholera major, Syria, FAIRMAIRE, p. 144, Ann. Ent. Belg. xxxvi, n. Colon melbournense, Victoria, BLACKBURN, p. 25, Tr. R. Soc. S. 1 xv. n. sp.
- Hydnobius fulvescens, Marseilles, Guillebeau, Échange, i, No. 6, n. sp.
- Necrophorus fossor, p. 147, humator, p. 150, metamorphoses; XA:
  Ann. Soc. L. Lyon, xxxix.
- Pinodytes hamiltoni, pusio, N. America, Horn, p. 45, Tr. Am. Ent xix, n. spp.
- Ptomaphagus nigriclavis, France, GUILLEBRAU, p. 116, Échange, (but 1892, p. 4, this is said to be sericeus, Panz.), n. sp.
- Silpha sinuata, p. 139, carinata, p. 142, metamorphoses; XAMBEU, Soc. L. Lyon, 1892.

#### TRICHOPTERYGIDA.

Critical notice of Flach's synopsis and synonymy of European Trterygide; MATTHEWS, Ann. N. H. (6) ix, pp. 442-448. Bæccera palumboi, Sicily, RAGUSA, p. 255, Nat. Sicil. xi, n. sp.

#### HISTERIDE.

- [Cf. Blackburn (93), Blaisdell (96), Horn (418), Lewis (553 555, 557, 558, 559, 560, 561), Schmidt (801, 802), Wickham (1
- Abræus mikado, Japan, Lewis, p. 356, Ann. N. H. (6) ix; A. stigme Perak, Schmidt, p. 29, Ent. Nachr. xviii: n. spp.
- Acritus shogunus, Japan, tasmaniæ, Tasmania, Lewis, p. 357, Anu. (6) ix, n. spp.
- Apobletes foliaceus and migneauxi, synonymical note; SCHMIDT, p

  Ann. Soc. Ent. Fr. lxi.
- A. nigritulus, Madagascar, Lewis, Ann. N. H. (6) ix, p. 342; A. a Perak, Schmidt, p. 19, Ent. Nachr. xviii; A. nirvana, B Lewis, p. 20, Ann. Mus. Genov. (2) xii: n. spp.
- Carcinops karenensis, Burma, Lewis, p. 31, Ann. Mus. Genov. (2 C. suavis, Sumatra, Schmidt, p. 27, Ent. Nachr. xviii: n. spp.

- Cylistix asiatica, Perak, p. 343, orientalis, Siam, Lewis, Ann. N. H. (6) ix, n. spp.
- Dendrophilus californicus, California, HORN, p. 46, Tr. Am. Ent. Soc. xix, n. sp.
- Eblisia cavipyga, Burma, Lewis, Ent. M. M. xxviii, p. 102; E. monticola, Burma, Lewis, p. 19, Ann. Mus. Genov. (2) xii: n. spp.
- Epiechnus: characters of; Lewis, Ann. N. H. (6) x, p. 23, pl. xix, figs. 8-10.
- E. taprobanæ, Ceylon, p. 355, birmanus, Burma, p. 356, Lewis, Ann. N. H. (6) ix, n. spp.
- Epierus nemoralis, Assam, Lewis, p. 347, Ann. N. H. (6) ix; E. amandus, p. 25, monticola, p. 26, Java, Schmidt, Ent. Nachr. xviii: n. spp.
- Eretmotus: habits of, and notes on specific characters; Lewis, Ann. N. H. (6) x, pp. 233-236, pl. xix, figs. 1-7. E. leprieuri, Mars., = (approximans, Fairm.); Lewis, p. 350, Ann. N. H. (6) ix.
- E. corpulentus, cirtensis, p. 233, kubyliæ, p. 234, bedeli, p. 235, Lewis, op. cit. (6) x, n. spp.
- Hetærius hornii, Wyoming, WICKHAM, Psyche, vi, p. 322, n. sp.
- Hister tropicola, Guinea, p. 23, punctipennis, Sierra Leone, oblongulus, Turkestan, p. 24, Schmidt, Ent. Nachr. xviii; H. robusticollis, Natal, p. 104, Lewis, Ent. M.M. xxviii; H. pransus, p. 24, gentilis, p. 25, stenocephalus, p. 26, sinuaticollis, p. 27, fragosus, p. 28, dentipes, p. 29, ixion, p. 30, striatipennis, p. 31, Burma, Lewis, Ann. Mus. Genov. (2) xii; H. rugistrius, Bengal, fumulus, Sumatra, Lewis, p. 346, Ann. N. H. (6) ix: n. spp.
- Hololepta dilatata, H. (Lioderma) pinguis, Gaboon, SCHMIDT, p. 17, Ent.
  Nachr. zviii; H. fea, Burma, Lewis, p. 17, Ann. Mus. Genov. (2)
  zii; H. pervalida, California, Blaisdell, p. 337, Zoe, iii: n. spp.
- Idister gestroi, Burma, Lewis, p. 18, Ann. Mus. Genov. (2) xii, n. sp.
- Liopygus famelicus, Burma, Lewis, p. 21, Ann. Mus. Genov. (2) xii, n. sp. Notodoma solstitiale, Perak, p. 348, rufulum, Borneo, p. 349, Lewis, Ann. N. H. (6) ix, n. spp.
- Onthophilus tuberculatus, p. 353, sculptilis, p. 354, Burma, Lewis, Ann. N. H. (6) ix; O. julii, Mexico, Lewis, p. 124, Ent. M. M. xxviii: n. spp.
- Pachycrarus ritsema, Mars., = (P. violaceipennis, Lew.); LEWIS, p. 345, Ann. N. H. (6) ix.
- Paratropus sererini, Bengal, Lewis, p. 142, Ann. Ent. Belg. xxxvi, n. sp. Paromalus complanatus, Panz., = (Platysoma theryanum, Reitt.); Reitter, p. 188, Wien. ent. Z. xi.
- P. mendicus, viaticus, p. 33, fujisanus, p. 34, vernalis, tardipes, p. 35, omineus, p. 36, montivagus, p. 37, Japan, Lewis, Ann. N. H. (6) ix; P. indicus, p. 32, rermiculatus, brevipes, p. 33, submetallicus, p. 34, Burma, Lewis, Ann. Mus. Genov. (2) xii; P. ludovici, N. S. Wales, Blackburn, p. 26, Tr. R. Soc. S. Austr. xv: n. spp.
- Platylister, n. g., type Platysoma ovatum, Er.; Lewis, Ent. M. M. xxviii, p. 103.

Platysoma solivayam, Burma, Lewis, p. 21, Ann. Mus. Genov. (2)
P. jejunum, Perak, Lewis, p. 343, Ann. N. H. (6) ix striatisternum, Borneo, Lewis, p. 103, Ent. M. M. xxviii distinctum, Java, p. 19, sincerum, Perak, p. 20, biimpressum, Qu land, p. 21, semilineatum, robustum, Australia, p. 22, Schmidt, Nachr. xviii: n. spp.

Plasius bisinuatus, Aru Is., Schimdt, p. 18, Ent. Nachr. xviii, n. sp. Psiloscelis limatulus, Assam, Lewis, p. 345, Ann. N. H. (6) ix, n. sp. Sternocalis: structure of claws described; Lewis, p. 263, Ent. M. M. x S. riaticus, Algeria, id. l. c., n. sp.

Teretrius walkeri, Tasmania, Lewis, p. 353, Ann. N. H. (6) ix, n. sp. Teretriosoma rajah, Burma, Lewis, p. 37, Ann. Mus. Genov. (2) xii, i Triballus onustus, Zanzibar, opimus, Borneo, Lewis, p. 350, Ann. 1 (6) ix; T. cyclonotus, orphanus, Burma, Lewis, p. 35, Ann. Genov. (2) xii: n. spp.

Trypeticus crassus, Perak, SCHMIDT, p. 28, Ent. Nachr. xviii nemorivagus, Burma, p. 351, prædaceus, Perak, p. 352, Lewis, N. H. (6) ix: n. spp.

Xestipyge fryi, Burma, LEWIS, Ann. N. H. (6) ix, n. sp.

## PHALACRIDÆ.

- [G. BLACKBURN (93), GUILLEBEAU (373, 378), PÉRINGUEY (673).] Revision of the palssarctic species with fresh classification; GUILLEI Rev. d'Ent. xi, p. 141.
- Eustilbus sharpi, Syria, Guillebeau, p. 191, Rev. d'Ent. xi, n. sp. Grourelleus, n. g., for G. prosternalis, n. sp., Saigon, Guillei p. exxxiv, Bull. Soc. Ent. Fr. lxi.
- Helectrus, n. g., for H. brisouti, Jordan, n. sp., Guillebeau, p. 187, d'Ent. xi.
- Litochrus sydneyensis, N. S. Wales, BLACKBURN, p. 26, Tr. R. Sc Austr. xv; L. promontorii, Cape Colony, Pébinguey, p. 110 S. African Phil. Soc. vi, 2: n. spp.
- Litochroides, n. g., for L. sharpi, Djedda, p. 187, simuaticollis, Ism p. 181, n. spp.; Guillebeau, Rev. d'Eut. xi.
- Micromerus, n. g., for Litochrus koltzei, Flach, Guillebeau, p. 189, d'Ent. xi.
- Olibrus desbrochersi, p. 171, selvei, p. 177, bedeli, p. 180, bonnairei, p. 181, viennensis, p. 182, biplagiatus, camptoides, gentilis, p. abeillei, p. 185, Palwarctic region, Guillebeau, Rev. d'Ent O. consanguineus, Cape Colony, Péringuey, Tr. S. African Soc. vi (2): n. spp.
- Phalacrus substriatus, metamorphoses; Guillebeau, p. 194, d'Ent. xi.
- P. confusus, p. 153 (and Échange, 1892, p. 20), grouvellei, insu p. 156, mayeti, championi, p. 158, Europe, Guillebeau, d'Ent. xi, n. spp.

- Pryacoderus, n. g., for P. lemoroi, n. sp., Biskra, Guillebeau, p. 187, Rev. d'Ent. xi.
- Tolyphus sedilloti, Tunis, Guillebeau, p. 163, Rev. d'Ent. xi, n. sp.

# NITIDULIDÆ.

- [Cf. ABEILLE (1), BLACKBURN (93), BLATCH & HORNER (103), FAIRMAIRE (262), GROUVELLE (368, 369, 370, 371), GUILLEBEAU (375, 376), PÉRINGUEY (673), REITTER (725).]
- Amphicrossus opacus, Bengal, GROUVELLE, p. 60, Ann. Ent. Belg. xxxvi, n. sp.
- Amphotis marginata, habits; WASMANN, p. 347, Deutsche e. Z. 1892.
- Aporozoum, n. g., for Coxelus sylvatious, Phil.; GERMAIN, p. 250, Act. Soc. Chili, ii.
- Brachypeplus ater, W. Africa, GROUVELLE, p. 292, Ann. Soc. Ent. Fr. lxi; B. birmanicus, p. 834, fex, p. 835, Burma, GROUVELLE, Ann. Mus. Genov. (2) xii: n. spp.
- Brachypterus obtusus, Marseilles; Guillebeau, p. 2, Supp. to Échange, No. 19, 1886; B. opacus, p. 66, drusus, p. 67, Syria, Abeille, Rev. d'Ent. xi: n. spp.
- Carpophilus (Urophorus) bicolor, heros, p. 836, C. (Ecnomorphus) murrayi, p. 837, Burma, GROUVELLE, Ann. Mus. Genov. (2) xii; C. bosschæ, Borneo, GROUVELLE, Notes Leyd. Mus. xiv, p. 43: n. spp.
- Cryptarcha alluaudi, W. Africa, GROUVELLE, p. 293, Ann. Soc. Ent. Fr. xli; C. bicolor, nigropunctata, p. 854, fee, p. 855, Burma, GROUVELLE, Ann. Mus. Genov. (2) xii: n. spp.
- Cyllodes quadrinotatus undulatus, p. 845, dubius nigropunctatus, p. 846, limbatus, ænescens, p. 847, humeralis, p. 848, Burma, GROUVELLE, Ann. Mus. Genov. (2) xii: n. spp.
- Decamerus and Diontolobus, Sol., belong to Nitidulidæ, not to Dermestidæ; GERMAIN, p. 250, Act. Soc. Chili, ii.
- Epuræa birmanica, consobrina, p. 840, reticulata, p. 841, Burma, GROUVELLE, Ann. Mus. Genov. (2) xii, n. spp.
- Eugoniopus reitteri, Burma, GROUVELLE, p. 852, Ann. Mus. Genov. (2) xii, n. sp.
- Idathina cincta, note on; BLACKBURN, p. 29, Tr. R. Soc. S. Austr. xv.
- Ischana fea, Burma, GROUVELLE, p. 842, Ann. Mus. Genov. (2) xii, n. sp.
- Librodor egregius, Burma, GROUVELLE, p. 856, Ann. Mus. Genov. (2) xii, n. sp.
- Lobostoma, n.g., near Epuræa, p. 90, for L. picea, n. sp, Obock, p. 91; FAIRMAIRE, Rev. d'Ent. xi.
- Meligethes (Acunthogethes) reyi, France, Guillebeau, Échange, i, No. 7, p. 2, n. sp.
- Mimema tabulæ, S. Africa, PÉRINGUEY, p. 111, Tr. S. African Phil. Soc. vi, 2, n. sp.

- Neopallodes deutatus, p. 849, vicinus, p. 850, Burma, GROUVELLE, Mus. Genov. (2) xii, n. spp.
- Notobrachypterus, n. g., near Brachypterus, p. 26, for N. australis, c. p. 27, bifoveatus, nitidiusculus, p. 28, liliputanus, p. 29, n. spp., tralia; Blackburn, Tr. R. Soc. S. Austr. xv.
- Pallodes birmanicus, feα, Burma, GROUVELLE, p. 851, Ann. Mus. G. (2) xii, n. spp.
- Parametopia bosschæ, Borneo, GROUVELLE, p. 44, Notes Leyd. xiv, n. sp.
- Physorhina reitteri, Burma, GROUVELLE, p. 843, Ann. Mus. Genov xii, n. sp.
- Platamartus, n. g., near Cercus, for P. jakolewi, n. sp., Siberia; RET p. 151, Wien. ent. Z. xi.
- Pocadites dubitabilis, Burma, GROUVELLE, p. 843, Ann. Mus. G (2) xii, n. sp.
- Pocadius testaceus, Bengal, GROUVELLE, p. 60, Ann. Ent. Relg. x n. sp.
- Rhizophagus oblongicollis, England, BLATCH & HORNER, Ent. M xxviii, p. 303, n. sp.
- Tetrisus epuracoides, Burma, GROUVELLE, p. 839, Ann. Mus. Genov xii, n. sp.
- Tricanus punctatissimus, p. 852, striatopunctatus, p. 853, Burma, G VELLE, Ann. Mus. Genov. (2) xii, n. spp.

## TROGOSITIDÆ.

[Cf. Léveillé (552), Xambeu (1021).]

Alindria sedilloti, = (sikorai, Kuw.); Léveillé, p. cviii, Bull. Ent. Fr. lxi.

Leptonyza fairmairei, Rio de Janeiro, LEVEILLE, p. zcv, Bull. Soc. Fr. lxi, n. sp.

Nemosoma siculum, Sicily, RAGUSA, p. 193, Nat. Sicil. xi, n. sp.

Trogosita mauritanica, metamorphoses; XAMBEU, p. 153, Ann. Sc Lyon, xxxix.

#### COLYDIIDE.

- [Cf. Blackburn (93), Fairmaire (263), Grouvelle (366, 368), P GUEY (673).]
- Bothrideres capicola, p. 111, distinctus, scutatus, p. 112, S. Africa, P GUEY, Tr. S. African Phil. Soc. vi (2), n. spp.
- Corticus syriacus, Akbès, FAIRMAIRE, p. 144, Ann. Ent. Belg. xxxvi, Ditoma opaca, elongatu, W. Africa, GROUVELLE, p. 296, Ann. Soc. Fr. lxi, n. spp.
- Mychocerus simonis, Venezuela, GROUVELLE, p. 102, Ann. Soc. En lxi, n. sp.
- Orthocerus funicornis, Bed., cf. Acotulus in Tenebrionidæ.

- Puryphus erichsoni, p. 99, serratus, crassus, p. 103, obesus, p. 101, Venezuela, GROUVELLE, Ann. Soc. Ent. Fr. lxi, n. spp.
- Sparactus: Endophiaus flexuosus referred to this; GERMAIN, p. 252, Act. Soc. Chili, ii.
- Tristaria labralis, Victoria, BLACKBURN, p. 30, Tr. R. Soc. S. Austr. xv, n. sp.

# CUCUJIDÆ and RHYSSODIDÆ.

- [Cf. Blackburn (93), Grouvelle (367, 368, 369).]
- Cutogenus feæ, Burma, GROUVELLE, p. 857, Ann. Mus. Genov. (2) xii, n. sp.
- Cerylon alluaudi, p. 297, infimum, striolatum, p. 298, W. Africa, GROU-VELLE, Ann. Soc. Ent. Fr. lxi, n. spp.
- Cryptamorpha macleayi, N.S. Wales, BLACKBURN, p. 31, Tr. R. Soc. S. Austr. xv, n. sp.
- Hyliota fee, p. 858, fallax, p. 859, Burms, GROUVELLE, Ann. Mus. Genov. (2) xii, n. spp.
- Læmophlæus australasiæ, Victoria, BlackBurn, p. 30, Tr. R. Soc. S. Austr. xv; L. rugifrons, mandibularis, p. 860, spinosus, p. 861, Burma, Grouvelle, Ann. Mus. Genov. (2) xii: n. spp.
- Myrabolia lindensis, S. Australia, p. 31, parca, N. S. Wales, p. 32, BLACK-BURN, Tr. R. Soc. S. Austr. xv, n. spp.
- Psammecus simonis, Philippines, GROUVELLE, p. 287, Ann. Soc. Ent. Fr. lxi, n. sp.
- Silvanopsis, n. g., for S. simonis, Philippines, p. 286, raffrayi, Singapore, p. 287, n. spp.; GROUVELLE, Ann. Soc. Ent. Fr. lxi.
- Silvanus longicornis, Philippines, GROUVELLE, p. 285, Ann. Soc. Ent. Fr. lxi; S. birmanicus, feæ, Burma, GROUVELLE, p. 863, Ann. Mus. Genov. (2) xii: n. spp.
- Rhysodes africanus, W. Africa, GEOUVELLE, p. 299, Ann. Soc. Ent. Fr. lxi, n. sp.

## CRYPTOPHAGIDE.

- [Cf. Blackburn (93), Guillebeau (375), Peringuey (673), Thomson (927).]
- Atomaria godarti, Marseilles, Guillebeau, Échange, i, No. 6, p. 3; A. eucalypti, N. S. Wales, Blackburn, Tr. R. Soc. S. Austr. xv, n. sp.
- Canoscelis grandis, Sweden, Thomson, Opusc. ent. xvi, p. 1773,
- Cryptophagus plagiatus, S. Africa, Péringuey, p. 113, Tr. S. African Phil. Soc. vi (2); C. gibbipennis, Victoria, Blackburn, p. 32, Tr. R. Soc. S. Austr. xv : n. spp.
- Emphylus americanus, habits; SCHWARZ, P. E. Soc. Wash. ii, p. 227.

# LATHRIDIIDÆ, MYCETOPHAGIDÆ.

[Cf. Baudi (43), Belon (53), Brisout (128), Reitter (725).]

Colovocera oculata, Burma, Gestro, p. 878, Ann. Mus. Genov. (2) x, r

Corticaria normanna, Calvados, Brisout, p. 68, Rev. d'Ent. xi; C.

pulchra, Taschkent, Reitter, p. 134, Wien. ent. Z. xi: n. spp.

Melanophthalma birmana, Burma, Gestro, p. 879, Ann. Mus. Genov
x, n. sp.

Monotoma punctata, Sicily, RAGUSA, p. 201, Nat. Sicil. xi, n. sp. Tritoma siculæ, Sicily, BAUDI, p. 121, Nat. Sicil. xi, n. sp.

## DERMESTIDE, BYERHIDE.

- [Cf. Blackburn (93, 94), Coucke (193), Frivaldszky (309), Gui beau (376), Tryon (953).]
- Characters of the *Dermestidæ* and *Byrrhidæ* of Belgium; Coupp. 69-79, Ann. Ent. Belg. xxxvi.
- Anthrenus pimpinellæ, n. var. latefasciatus; Reitter, p. 134, W ent. Z. x.
- Attagenus pantherinus, parasitic on Anthophora; Buysson, p. celvii, I Soc. Ent. Fr. Ixi.
- A. gobicola, China, FRIVALDSZKY, p. 122, Term. füzetek, xv, n. sp.
- Trogoderma froggatti, N. S. Wales, Blackburn, p. 34, Tr. R. Soc Austr. xv; T. reitteri, S. Australia, Blackburn, p. 208, t. c. : n.

Dendropinis calva, vestita, New Guinea, TRYON (953), n. spp.

- Limnichus variegatus, Fréjus, Guillebeau, p. 3, Supp. to Échange, 19, 1886, n. sp
- Pedilophorus æneus, F., = (subparallelus, Motsch.); REITTER, p. Wien. ent. Z. xi.'

## PARNIDE, HETEROCERIDE.

- [Cf. Angell (14), Germain (322), Grouvelle (369, 372), Péring (673).]
- Dryops opacus, anescens, Burma, GROUVELLE, p. 865, Ann. Mus. Ge. (2) xii, n. spp.
- Elmis columbiensis, British Columbia, Angell, p. 84, Ent. News, E. anthracinus, trivialis, p. 245, scissicollis, p. 246, nyctelioides, p. Chili, Germain, Act. Soc. Chili, ii : n. spp.
- Macronychus minusculus, Sumatra, GROUVELLE, p. 187, Notes Leyd. A xiv, n. sp.
- Potamophilus fex, longipes, Burma, GROUVELLE, p. 864, Ann. M. Genov. (2) xii, n. spp.
- Sostea birmanica, Tenasserim, GROUVELLE, p. 866, Ann. Mus. Genov. xii, n. sp.
- Stenelmis testacea, birmanicu, Burma, Gestro, p. 867, Ann. Mus. Gei (2) xii; S. bosschæ, Borneo, sulcata, Sumatra, Grouvelle, p. Notes Leyd. Mus. xiv: n. spp.

- Strina promoutorii, S. Africa, Péringuey, p. 113, Tr. S. African Phil. Soc. vi (2), n. sp.
- Heterocerus capensis, meridionalis, S. Africa, PÉRINGUEY, p. 114, Tr. S. African Phil. Soc. vi (2), n. spp.

## LUCANIDE and PASSALIDE.

- [Cf. Fairmaire (263), Nonfried (644, 648), Ritsema (758, 759, 761 762), Tryon (953).]
- List of *Lucanidæ* found in Java, Sumatra, and Nias, RITSEMA, pp. 139-144, Notes Leyd. Mus. xiv.
- Cyclommatus, list of the described species; RITSEMA, Notes Leyd. Mus. xiv, pp. 3-6.
- C. frey-gessneri, Java, RITSEMA, p. 1, t. c.; C. squamosus, Borneo, RITSEMA, p. 45, t. c.: n. spp.
- Dorcus parallelus, costatus, brevis, synonymic note with figg.; Horn, Ent. News, 1892, p. 73, pl. iii.
- Hexarthrius colesi, E. India, NONFRIED, p. 365, Berl. Ent. Zeit. xxxvi, n. sp.
- Lucanus cervus, variation in & mandibles; Bateson & Brindley, P. Z. S. 1892, p. 593.
- Nigidius oberndorferi, Nossibé, Nonfried, p. 118, Ent. Nachr. xviii, n. sp. Phalacrognathus muelleri, habitat, Skuse, p. 20, P. Linn. Soc. N.S.W. (2) vii.
- Platycerus delagrangei, Syria, FAIRMAIRE, p. 145, Ann. Ent. Belg. xxxvi, n. sp.
- Prosopocælus serricornis, n. var. brunneus, Nonfried, p. 119, Ent. Nachr. xviii, &c.
- P. pasteuri, Java, Rithema, p. 31, Notes Leyd. Mus. xi; P. tarsalis, Java, Rithema, t. c. p. 191: n. spp.
- Mastochilus subobliquus, New Guinea, TRYON (953), n. sp.

#### SCARABÆIDÆ.

- [Cf. Bateson & Brindley (42), Bedel (49, 50), Bergroth (60), Blackburn (90, 91, 92, 93, 94, 95), Brenske (122, 123, 124, 125), Casey (161), Croissandeau (194), Distant (210), Duvivier (230), Fairmaire (262, 263, 264, 268), Frivaldszky (309), Gestro (328), Jacobsohn (443), Jakowleff (451), Kraatz (514), Nevinson (635, 636), Nonfried (642, 643, 644, 646, 647, 648), Péringuey (672), Reitter (725, 727, 729), Schwarz (808), Semenow (830, 832, 833, 834), Smith (864, 865), Stiles (901), Xambeu (1021); also Cetoniini, p. 119.]
- Hours of flight of Scarabaida; SCHWARZ (808).

Coprini.

Ammæcius elevatus, metamorphoses; Xambeu, p. 165, Ann. Soc. L. Lyon, xxxix.

- Aphodius fossor, p. 157, alpinus, p. 158, depressus, p. 162, metamorę XAMBEU, Ann. Soc. L. Lyon, xxxix. A. granarius, p. 1, dej p. 16, metamorphoses; XAMBEU, Mœurs, ii.
- A. bidentulus, gibbifrons, Obock, FAIRMAIRE, p. 92, Rev. d'Ent. yorkensis, S. Australia, BLACKBURN, p. 209, Tr. R. S. Austr. xv; A. frenchi, lindensis, S. Australia, BLACKBURN, t. c.: n. spp.
- Atenius mendax, Victoria, torridus, S. Australia, BLACKBURN, p. 36, Soc. S. Austr. xv, n. spp.
- Ateuchetus, n. subg. of Scarabæus; BEDEI., p. 283, Abeille, xxvii.
- Athyreus chalybeatus, Argentine Rep., FAIRMAIRE, p. 242, Ann. Enxxvi, n. sp.
- Autolethrus, n. subg. of Lethrus; Semenow, Hor. Ent. Ross. xxvi, Bolboceras serripes, Fairm., & described; Gestro, p. 756, Ann Genov. (2) xii.
- B. gagarinei, Turkestan, FAIRMAIRE, p. cxxii, Bull. Soc. Ent. I B. gaujani, Obock, FAIRMAIRE, p. 95, Rev. d'Ent. xi; B. Transvaal, DISTANT, p. 191, pl. i, fig. 5, Naturalist Tran n. spp.
- Catharsius insignis, csphalotes, S. W. Africa, Péringuey, p. 27, African Phil. Soc. vi, 2, n. spp.
- Chætopisthes termiticola, Burma, Gestro, p. 904, Ann. Mus. (2) x, n. sp.
- Copris carolina, anatomy of mouth and homology of mandibles; pp. 83-87, pls. ii & iii, Tr. Am. Ent. Soc. xix.
- Geotrupini: notes on Jacobsohn's classification; REITTER, pp. 2; Wien, ent Z. xi.
- Geotrupes: revision of the genera allied to, or subgenera of; sohn, Hor. Ent. Ross. xxvi, pp. 245-257. G. rernalis, var. FAUVEL, p. 57, Rev. d'Ent. xi.
- Glaresis oxiana, W. Turkestan, Semenow, p. 470, Hor. Ent. xxxvi, n. sp.
- Ilybosorus incultus, S. W. Africa, PERINGUEY, p. 31, Tr. S. Africa Soc. vi, 2, n. sp.
- Lethrotrypes, n. subg. of Thorectes; JACOBSOHN, p. 257, Hol Ross. xxvi.
- Lethrulus, n. subg. of Lethrus, SEMENOW, p. 235, Hor. Ent. Ross. x Lethrus, n. subg., see Autolethrus, Lethrulus, Scelolethrus.
- L. anescens, p. cxxi, bradytus, mediocris, p. cxxii, Turkestan, Fair Bull, Soc. Ent. Fr. lxi; L. mandibularis, p. 182, forcipatus, Asia Minor, spinimanus, Askhabad, p. 190, Jakowleff, Hc Ross. xxvi; L. (Heteroplistodus) jakowlewi, Kirghisia, p. L. (Autolethrus) koshanstschikowi, Transcaspian region, p. Semenow, Hor. Ent. Ross. xxvi: n. spp.
- Mendidius feculentus, Obock, FAIRMAIRE, p. 93, Rev. d'Ent. xi, n. 1 Mesotrypes, n. subg. of Thorectes; JACOBSOHN, p. 256, Hol Ross. xxvi.

- Neoctodon, n. n. for Octodon, Lansb.; BEDEL, p. 282, Abeille, xxvii.
- Onitis: synopsis of the Mediterranean species of Chironitis, Onitis, and Bubas; Bedel, pp. 250-280, Abeille, xxvii.
- O. suspectus, amabilis, p. 29, bovinus, exiguus, p. 30, S. W. Africa, PÉRINGUEY, Tr. S. African Phil. Soc. vi, 2; O. intermedius, China, FRIVALDSZKY, p. 122, Term. füzetek, xv: n. spp.
- Onthophagus furciceps, Mars., characters of; HEYDEN, p. 112, Wien. ent. Z. xi. O. crocatus, Muls., = (imitator, Reit.); REITTER, p. 26, Wien. ent. Z. xi. O. ater, n. var., granulipennis, FRIVALDSZKY, p. 123, Term. füzetek, xv. O. adelaidæ, Hope, = (hostilis, Har.); BLACKBURN, p. 283, P. Linn. Soc. N.S.W.(2) vii.
- O. trispinus, Margelan, REITTER, p. 63, Wien. ent. Z. xi; O. sulcicollis, Margelan, REITTER, p. 135, t. c.; O. quadriarmatus, Obock, FAIRMAIRE, p. 91, Rev. d'Ent. xi; O. geelongensis, Victoria, BLACKBURN, p. 34, Tr. R. Soc. S. Austr. xv; O. bluckwoodensis, henleyensis, S. Australia, p. 208, nitidior, N. S. Wales, p. 209, BLACKBURN, t. c.; O. murchisoni, Australia, BLACKBURN, p. 25, op. cit. xvi: n. spp.
- Oxysternon macleayi, Amazons, pteroderum, Brazil, Nevinson, Ent. M. M. xxviii, p. 34, n. spp.
- Phanaus: synonymic list of the species; NEVINSON (636).
- P. flohri, Mexico, NEVINSON, Ent. M. M. xxviii, p. 33, n. sp.
- Proctammodes, n. n. for Proctophanes, Har. (nec Proctophana, Lac.);
  BLACKBURN, p. 37, Tr. R. Soc. S. Austr. xv.
- Psammobius plicatulus, Obock, FAIRMAIRE, p. 95, Rev. d'Ent. xi, n. sp. Rhyssemus asperocostatus, læsifrons, Obock, FAIRMAIRE, p. 94, Rev. d'Ent.
- Scarabæus plausibilis, S. Africa, Peringuey, p. 25, Tr. S. African Phil. Soc. vi (2), n. sp.
- Scelolethrus, n. subg. of Lethrus; Semenow, p. 236, Hor. Ent. Ross. xxvi. Silotrupes epistomalis, note on; Croissandeau, p. lviii, Bull. Soc. Ent. Fr. lxi.
- Thorectes cheroni, Corsica, CROISSANDEAU, p. ccxxi, Bull. Soc. Ent. Fr. lxi, n. sp.
- Trox scaber, metamorphoses; XAMBEU, p. 168, Ann. Soc. L. Lyon, xxxix. T. elderi, tatei, S. Australia, p. 37, eremita, Centr. Australia, quadridens, S. Australia, p. 38, euclensis, W. Australia, augusta, S. Australia, p. 39, relutinus, N. Queensland, p. 40, BLACKBURN, Tr. R. Soc. S. Austr. xv, n. spp.

# Glaphyrini and Melolonthini.

- Catalogue of Glaphyrides, Systellopides, Melolonthides, and Euchirides, described since the Munich Catalogue; Nonfried, Berl. Ent. Zeit. xxxvii, pp. 249-290.
- Classification of Rhizotrogides and Melolonthides; Brenske, pp. 33-38, Berl. Ent. Zeit. xxxvii.
- Ancylonycha holosericea, Redt., referred to Hilyotrogus; BRENSKE, p. 154, Ent. Nachr. xviii.

- A. cochinchina (sic), Cochin China, Nonfried, p. 360, Berl. Ent xxxvi, n. sp.
- Anoxia pilosa, F., = (cinerea, Motsch.); Semenow, p. 476, Hor. Ross. xxvi.
- Brahmina: revision of the genus, referring to it various species in other genera, and merging in it Rhizocolax, Mots., and Crz bius, Kraatz; Brenske, Berl. Ent. Zeit. xxxvii, pp. 79-124 rubretra, Fald., and allied species, notes on; Kraatz, Deut. Z. 1892, p. 309. B. conspersa, sibirica, note on; Brenske, pt. c.
- B. bengalensis, Bengal, NONFRIED, p. 229, Berl. Ent. Zeit. xxx donckieri, Darjeeling, p. 86, castanipes, sibirica, E. Siberia, turkestanica, Turkestan, p. 94, phytaloides, Saigon, p. 98, setosa jeeling, p. 99, cotesi, Sikkim, p. 100, chinensis, China, p. 101, ab Saigon, p. 102, siamensis, Malacca, p. 103, microphylla, turo flabellata, E. India, p. 104, sumatrensis, Sumatra, obscura, 1 Hills, p. 105, cribripennis, Sumatra, p. 106, rugulosa, Borneo, heydeni, Korea, cardoni, p. 108, thoracica, p. 109, E. India, china, p. 112, himalayica, sikkimensis, E. India, p. 113, bur Buroo Is., p. 114, Brenske, Berl. Ent. Zeit. xxxvii: n. spp.
- Byrrhomorpha, n. g., near Heteronyx, p. 109, for B. verres, p. 110, rosa, p. 111, n. spp., Australia; BLACKBURN, P. Linn. Soc. 1 (2) vii.
- Ceraspis elegans, imitatrix, Honduras, Nonfried, p. 224, Berl. Ent xxxvi, n. spp.
- Cheiragra macleayi, Victoria, BLACKBURN, p. 482, P. Linn. Soc. 1 (2) vi, n. sp.
- Colpochila campestris, W. Australia, Blackburn, p. 103, P. Lin N.S.W. (2) vii, n. sp.
- Cryphæobius: validity and characters of the genus; KRAATZ, Deut Z. 1892, pp. 307-311.
- Cyphonotus oryctoides, Transcaspian region, Semenow, p. 474, Ho Ross. xxvi, n. sp.
- Dejeania brenskei, Siam, Nonfried, p. 226, Berl. Ent. Zeit. xxxvi, Diphucephala elegans, Victoria, Blackburn, p. 481, P. Linn. Soc. I (2) vi; D. spreta, W. Australia, Blackburn, p. 99, P. Lini N.S.W. (2) vii: n. spp.
- Empecta cambouei, Madagascar, Nonfried, p. 107, Ent. Nachr. xviii Encya nigra, Madagascar, Nonfried, p. 227, Berl. Ent. Zeit. xxxvi Eupegylis, n. g., Schizonychides, for E. confusa, n. sp., Congo, Duvp. 56, Ann. Ent. Belg. xxxvi.
- Frenchella, n. g., near Colpochila, for F. lubrica, n. sp., V: BLACKBURN, p. 104, P. Linn. Soc. N.S.W. (2) vii.
- Glaphyrus sogdianus, Zerafschan, Semenow, p. 477, Hor. Ent. xxvi, n. sp.
- Haplopsis viridis, Australia, BLACKBURN, p. 109, P. Linn. Soc. 1 (2) vii, n. sp.

- Heptophylla, Motsch., characters amended; Brenske, p. 155, Ent. Nachr. xviii.
- Heteronyx baldiensis, p. 485, terrena, p. 486, incognitus, alpicola, p. 487, tridens, p. 489, consanguineus, p. 490, proditor, p. 492, Australia, BLACKBURN, P. Linn. Soc. N.S.W. (2) vi; H. merus, arcanus, p. 106, protervus, p. 107, Queensland, BLACKBURN, P. Linn. Soc. N.S.W. (2) vii; H. rhinoceros, p. 284, marcidus, p. 285, Australia, BLACKBURN, t. c.; H. severus, alienus, p. 29, fervidus, p. 30, decorus, fruserensis, p. 31, cribriceps, p. 32, siccus, helmsi, p. 33, furvus, p. 34, Australia, BLACKBURN, Tr. R. Soc. S. Austr. xvi: n. spp.
- Molotrichia bicolorea, Heyd., referred to Hilyotrogus, BRENSKE, p. 154, Ent. Nachr. xviii.
- H. farinosa, E. India, NONFRIED, p. 228, Berl. Ent. Zeit. xxxvi; H. glubriclypeata, E. India, cochinchina, Cochin China, p. 161, obscura, Sikkim, p. 162, luta, Hong Kong, richteri, Borneo, p. 163. scabrifrons, Ceylon, p. 164, behrensi, Sumatra, flachi, Manilla, p. 165, staudingeri, Sikkim, p. 166, frontalis, cavifrons, p. 167, sericata, p. 168, sikkimensis, p. 169, biehli, E. India, javana, Java, p. 170, nitida, p. 171, aurosericea, p. 172, E. India, foveolata, Borneo, p. 173, scrobiculata, Sikkim, p. 174, bombycina, Tenasserim, p. 175, iridipennis, nigricollis, E. India, p. 177, singhalensis, Ceylon, p. 178, karschi, E. India, p. 179, parva, Ceylon, sharpi, Borneo, p. 180, longicarinata, kallies, p. 181, standfussi, Sumatra, p. 182, maxillata, Java, p. 183, anthracina, p. 184, cotesi, plagiata, p. 185, E. India, setosa (Walk.), Ceylon, p. 186, bipunctata, Java, burmeisteri, p. 187, philippinea, p. 188, Philippines, inducta (Walk.), Ceylon, p. 189, pygidialis, Sumatra, glubrifrons, Khasia Hills, p. 190, atkinsoni, S. India, p. 191, Brenske, Berl. Ent. Zeit. xxxvii: n. spp.
- Homaloplia ruricola, metamorphoses; XAMBEU, Mœurs, ii, p. 30. H. diabolica, Reitt., = (ursina, Fairm.); Brenske, p. 170, Ann. Ent. Belg. xxxvi.
- H. ursina, mutilata, Syria, FAIRMAIRE, p. 147, Ann. Ent. Belg. xxxvi, n. spp.
- Hoplia fasciculata, p. 108, cornuta, p. 109, Madagascar, Nonfried, Ent. Nachr. xviii; H. siningensis, China, Frivaldszky, p. 123, Torm. füzetek, xv : n. spp.
- Isonychus prasinus, Honduras, Nonfried, p. 223, Berl. Ent. Zeit. xxxvi, n. sp.
- Lacknosterna as intermediate host of a parasite affecting Swine; STILES (901).
- L. senegalensis, Africa, Nonfried, p. 106, Ent. Nachr. xviii, n. sp.
- Lachnota bedeli, Siberia, BERGROTH, p. 98, Wien. ent. Z. xi.
- Lasiopsis, Er., and allied genera, composition of; Bergroth, p. 98, Wien. ent. Z. xi.
- L. bergrothi, Caucasus, Reitter, p. 101, Wien. ent. Z. xi, n. sp.
- Lepidiota pauper, Sumatra, p. 40, siamensis, Siam, p. 41, nonfriedi, Hong Kong, p. 42, crenaticollis, E. India, p. 43, richteri, Himalaya, p. 44,

- L. (Eucirrus) elegans, Borneo, p. 45, L. sus, Malay Penina, h. China, p. 46, ornata, Sumatra, p. 47, africana, Senegal, reuleauxi, New Guinea, p. 49, quedenfeldti, Aduma, labrata, Tep. 50, oberndorferi, Sumatra, p. 51, BRENSKE, Berl. Ent. xxxvii, n. spp.
- Leucopholis rufu, p. 52, pungiena, p. 53, celebensis, Celebes, sumat Sumatra, p. 54, nigra, Borneo, p. 55, lateralis, Malacca, staudingeri, Borneo, p. 57, crassa, Assam, elongata, E. India?, tristis, Malacca, p. 59, Brenske, Berl. Ent. Zeit. xxxvii, n. spp
- Liogenys opacicollis, bidentulus, Argentine Rep., FAIRMAIRE, p. 243 Ent. Belg. xxxvi, n. spp.
- Liparetrus spretus, p. 482, alpicola, p. 483, Australia, BLACKBU Linn. Soc. N.S.W. (2) vi; L. maurus, p. 99, lugens, p. adelaidæ, p. 101, Australia, BLACKBURN, P. Linn. Soc. N.S. vii: n. spp.
- Liparochrus rufus, Australia, BLACKBURN, p. 25, Tr. R. Soc. S. xv, n. sp.
- Machidius tibialis, N. S. Wales, BLACKBURN, Tr. R. Soc. S. xv, n. sp.
- Melichrus, n. g., Rhizotrogides, for M. kolbei, n. sp., Darjeeling: Br p. 156, Ent. Nachr. xviii.
- Melolontha: ravages and numbers of M. vulgaris and hippocas Sweden; LAMPA (536). M. hippocastani, n. var. baicalica; RI p. 152, Wien. ent. Z. xi.
- M. siamensis, Siam; Nonfried, p. 230, Berl. Ent. Zeit. xxxvi, n. : Metabolus flavescens, Turkostan, Brenske, p. 153, Ent. Nachr. xvii Monotropus starcki, Charkow, Reitter, p. 142, Wien. ent. Z. xi, n
- Ocnodus lugubris, W. Australia, BLACKBURN, p. 105, P. Linn. Soc.
- (2) vii; O. spinicollis, p. 27, fallax, ferrugineus, p. 28, At Blackburn, Tr. R. Soc. S. Austr. xvi: n. spp.
- Octoplasia, n. g. for Luchnosterna lineata and princeps, Shp.; B1 p. 152, Ent. Nachr. xviii.
- Polyphylla diffracta, New Mexico, Casey, p. 18, Ann. N. York n. sp.
- Pseudoheteronyx, n. g., near Heteronyx, for P. helwoides, n. sp., Quee Blackburn, p. 111, P. Linn. Soc. N.S.W. (2) vii.
- Pseudopholis, n.g., near Leucopholis, p. 57, for P. squamulosa, n. sp. p. 58; Duvivier, Ann. Ent. Belg. xxxvi.
- Pseudotrochalus brenskei, Zambesi, Nonfried, p. 106, Ent. Nach
- Rhabdopholis melolouthoides, hab. ?, Brenske, p. 157, Ent. Nacl n. sp.
- Rhizocolax senescens, = (Rhizotrogus potanini, Sem.); Semenow Hor. Ent. Ross. xxvi.
- Rhizotrogus cicatricosus, metamorphoses; Xambeu, p. 173, Ann. Lyon, xxxix. R. altaicus, = (Monotropus suwortzewi, Semenow, p. 472, Hor. Ent. Ross. xxvi.

- Rhopea soror, mussoni, N. S. Wales; BLACKBURN, P. Linn. Soc. N.S.W. (2) vii, p. 112, n. spp.
- Schizonycha angustiformis, Obock, aspericollis, Abyssinia, FAIRMAIRE, p. 96, Rev. d'Ent. xi; S. curtula, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. vi, 2: n. spp.
- Sciton, n. g., near Anodontonyx, p. 101, for S. ruber, n. sp., S. Australia, p. 102; BLACKBURN, P. Linn. Soc. N.S.W. (2) vii.
- Serica carneola, p. 34, deceptor, obesa, lucidula, p. 35, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. v, 2; S. subglobosa, Centr. Africa, Nonfried, p. 105, Ent. Nachr. xviii; S. picea, Siam, Nonfried, p. 359, Berl. Ent. Zeit. xxxvi; S. iridescens, Borneo, p. 221, siamensis, Siam, p. 222, Nonfried, Berl. Ent. Zeit. xxxvi: n. spp.
- Symmachia, n. g., near Brahmina, for S. chinensis, n. sp., China; Brenske, p. 151, Ent. Nachr. xviii.
- Tanyproctus rugulosus, Syria, FAIRMAIRE, p. 148, Ann. Ent. Belg. xxxvi, n. sp.
- Thyce blaisdelli, pp. 19 & 214, squamosa, p. 20, California, CASEY, Ann. N. York Ac. vi, n. spp.
- Tricholepis grandis, Cast., = (Leucopholis manilla, Redt.); BRENSKE, p. 61, Berl. Ent. Zeit. xxxvii.
- Triodonta flavimana, Burm., = (difformipes, Fair.); BRENSKE, p. 170, Ann. Ent. Belg. xxxvi.
- T. modesta, S. W. Africa, Peringuey, Tr. S. African Phil. Soc. vi, 2, p. 36; T. difformipes, dispar, Syria, Fairmaire, p. 146, Ann. Ent. Belg. xxxvi: n. spp.
- Trochalus modestus, p. 32, mærens, plagiger, rufovittatus, p. 33, mæstus, plagiatus, p. 34, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. vi. 2, n. spp.

### Rutelini.

- List of Rutelidæ described since the Munich Catalogue; Nonfried (643).

  Adoretus parviceps, Obock, Fairmaire, p. 93, Rev. d'Ent. xi; A. sykoræ,
  Madagascar, kahlei, Zambesi, p. 110, fascicularis, Abyssinia, p. 117,
  Nonfried, Ent. Nachr. xviii; A. flavoritutus, Siam, p. 362, castaneus,
  Borneo, p. 363, griseosetosus, Siam, p. 364, Nonfried, Berl. Ent.
  Zeit. xxxvi: n. spp.
- Anomala obscuripes, Obock, FAIRMAIRE, p. 97, Rev. d'Ent. xi; A. windrathi, Siam, Nonfried, Berl. Ent. Zeit. xxxvi, p. 361; A. viridicostata, p. 86, hirsutala, p. 87, nigripes, p. 88, S. China, Nonfried, Ent. Nachr. xviii; A. australasiæ, Queensland, Blackburn, p. 113, P. Linn. Soc. N.S.W. (2) vii; A. (Euchlora) nigra, p. 234, A. fruhstorferi, p. 235, Java, inscripta, Siam, p. 236, pyropyga, Venezuela, p. 237, Nonfried, Berl. Ent. Zeit. xxxvi: n. spp.
- Anoplognathus brecicollis, p. 493, macleayi, p. 495, Australia, BLACKBURN, P. Linn. Soc. N.S.W. (2) vi, n. spp.
- Areoda magnifica, Brazil, NONFRIED, p. 238, Berl. Ent. Zeit. xxxvi, n. sp.

- Epectinaspis hondura, Honduras, Nonfried, p. 232, Berl. Ent. 2 xxxvi, n. sp.
- Euchlora heydeni, China, FRIVALDSZKY, p. 124, Term. füzetek, n. sp.
- Gnatholabis dorsalis, p. 299, nickerli, p. 301, Trop. Africa, KRA Deutsche e. Z. 1892, n. spp.
- Hadropopillia, n. g., for Popillia splendida, Guér.; KRAATZ, p Deutsche e. Z. 1892.
- Heteroplia siamensis, Siam, Nonfried, p. 362, Berl. Ent. Zeit. x. n. sp.
- Ischnopopillia, n. g., p. 292, for a part of Popillia and I. moorei, p. erythroptera, p. 295, n. spp., E. India; KRAATZ, Deutsche 1892.
- Leucothyreus purpureo-sericeus, Ecuador, Nonfried, p. 239, Berl. Zeit. xxxvi, n. sp.
- Malaia? semperi, Philippines, KRAATZ, p. 178, Deutsche e. Z. 189: ferred below to Pseudomalaia, n. g.), n. sp.
- Parastasia bigibbosa, Siam, Nonfried, p. 238, Berl. Ent. Zeit. x n. sp.
- Phyllopertha latitursis, p. 232, sericeomicans, p. 233, Honduras, NONF Berl. Ent. Zeit. xxxvi, n. spp.
- Pacilosticta, n. g., p. 290, for P. princeps, n. sp., Ceylon, p. 291; KR Doutsche e. Z. 1892.
- Popillia: revision of; KRAATZ, Deutsche e. Z. 1892, pp. 177, &c.
- P. flavotæniata, Cameroons, Nonfried, p. 118, Ent. Nachr. xviii oberthuri, Zanzibar, p. 184, candezei, Fernando-Po, p. 185, wel Angola, p. 187, erythropus, Sierra Leone, p. 191, smaragdina, p. cupripes, lævicollis, p. 229, soror, p. 230, bitacta, p. 233, W. A strigilata, E. Africa, p. 235, signifera, p. 237, opaca, p. 238, W. A perrottetii, p. 244, S. India, semicuea, Kiu-Kiang, p. 251, s. Hong Kong, p. 252, viridula, Thibet, p. 255, straminipennis, p. subquadrata, p. 259, anomaloides, p. 260, 6-maculata, p. 261, (fea, p. 269, testaceipennis, p. 270, flavofusciata, p. 271, Burma, collis, p. 272, nottrotti, p. 273, lævicollis, p. 275, pilicollis, p. 2 India, andamanica, p. 279, Andaman Is., sumatrensis, Sumatra, variabilis, p. 283, picticollis, p. 284, depressiuscula, p. 286, dep. 287, Philippines, Kraatz, Deutsche e. Z. 1892: n. spp.
- Pseudomalaia, n. g., for Popillia pilifera, Burm., tagala, Hell., and ... semperi (Kr., suprà); Kraatz, p. 296, Deutsche e. Z. 1892.
- Rhinyptia dollei, plana, and Amphimallon rostratum, queried as syno-FAIRMAIRE, p. 97, Rev. d'Ent. xi.
- R. testucea, E. India, Nonfried, p. 230, Berl. Ent. Zeit. n. sp.
- Singhala schaiblei, S. China, Nonfried, p. 89, Ent. Nachr. n. sp.
- Spilopopillia, n. g., for Popillia scryuttata, Fairm.; KRAATZ, 1 Deutsche e. Z. 1892.

### Dynastini.

- Aneurystypus collaris, W. Australia, BLACKBURN, p. 286, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Corynophyllus metallicola, N. S. Wales, BLACKBURN, p. 286, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Cyclocephala occipitalis, Argentine Rep., FAIRMAIRE, p. 244, Ann. Ent. Belg. xxxvi, n. sp.
- Dichodontus renkeni, Borneo, Nonfried, p. 365, Berl. Ent. Zeit. xxxvi, n. sp.
- Ligyrus bidentulus, distinctus, Argentine Rep., FAIRMAIRE, p. 244, Ann. Ent. Belg. xxxvi, n. spp.
- Oryctomorphus: notes on the species; GERMAIN, p. 260, Act. Soc. Chili, ii.
- Phyllognathus fortipes, Obock; FAIRMAIRE, p. 98, Rev. d'Ent. xi, n. sp.
- Pseudoryctes tectus, Australia, BLACKBURN, p. 211, Tr. R. Soc. S. Austr. xv, n. sp.
- Xylotrupes gideon, statistics of variation of horns of male; BATESON & BRINDLEY, P. Z. S. 1892, p. 589.

### Cetoniini.

- [Cf. Blackburn (91), Gestro (327), Janson (454), Kolbe (502, 503, 505, 507, 508), Kraatz (515, 516, 518), Nonfried (644, 647, 648), Péringuey (672, 673), Xambeu (1021).]
- Agenius elegans, Cape Colony, Péringuey, p. 115, Tr. S. African Phil. Soc. vi (2), n. sp.
- Anaspilus, n. g., near Cymophorus, p. 140, for A. rufiventris, n. sp., Guinea, p. 141; Kolbe, S. E. Z. liii.
- Anelaphinis, n. g., for Cetonia dominula, Har.; Kolbe, p. 136, S. E. Z. liii.
- Aphanesthes, n. g., near Chordodera, p. 248, for A. pseudincoides, n. sp., Cameroons, p. 249; Kolbe, SB. nat. Fr. 1892.
- Cetonia (Pachnoda) kustai, Centr. Africa, Nonfried, p. 122, Ent. Nachr. xviii; C. montana, Himalaya, Nonfried, p. 371, Berl. Ent. Zeit. xxxvi; C. rhododendri, p. 847, pl. ii, fig. 9, oatesii, p. 848, cariana, p. 849, Burma, Gestro, Ann. Mus. Genov. (2) x: n. spp.
- Cœnochilus leoninus, S. W. Africa, Péringuey, p. 39, Tr. S. African Phil. Soc. vi (2); C. (Xenogenius) conradti, E. Africa, Kolbe, p. 71, SB. nat. Fr. 1892: n. spp.
- Conradtia, n. g., near Diplognatha, p. 69, for C. principalis, n. sp., E. Africa, p. 70; Kolbe, SB. nat. Fr. 1892.
- Corynotrichius, n. g., Trichiides, p. 259, for C. bicolor, n. sp., Cameroons, p. 260; Kolbe, SB. nat. Fr. 1892.
- Cymophorus monticola, E. Africa, Kolbe, p. 71, SB. nat. Fr. 1892; С. toganus, p. 139, flavonotatus, p. 140, Guinea, Kolbe, S. E. Z. liii: п. spp.
- Digenethle calata, Gestro, = (ramulosipennis, Thoms.); Gestro, p. 855, Ann. Mus. Genov. (2) x.

- Diphrontis monticola, Cameroons, Kolbe, SB. nat. Fr. 1892, p. n. sp.
- Diploa, n. g., near Xiphoscelis, for D. probes, n. sp., E. Africa; Ko p. 65, SB. nat. Fr. 1892.
- Diplognatha montana, E. Africa, Kolbe, p. 68, SB. nat. Fr. 1892 preussi, Cameroons, Kolbe, p. 256, t. c.: n. spp.
- Dolichostethus, n. g., for Elaphinis atomosparsa, Fairm.; Kolbe, p. S. E. Z. liii.
- Epistalagma cornuta, n. sp., for an insect which is probably the : E. multiimpressa, Fairm., KRAATZ, Deutsche e. Z. 1892, p. figured, pl. iv, fig. 4.
- Eucosma breviceps, Cameroons, Kolbe, p. 253, SB. nat. Fr. 1: E. flavoguttata, Guinea, Kolbe, p. 132, S. E. Z. liii: n. spp.
- Eudicella smithi, var. trilineata, Quedf., = (thomsoni, Ancey, and quardi, Oberth.); Kolbe, p. 63, SB. nat. Fr. 1892.
- Eudoxazus, n. g., Trichiides, p. 72, for E. conradti, n. sp., E. Africa, p. Kolbe, SB. nat. Fr. 1892.
- Eupachnoda, n. g., for Pachnoda inscripta; Kolbe, p. 253, SB. nat 1892.
- Eutelesthes, n. g., near Plæsiorrhina, for E. luteralis, n. sp., Camero Kolbe, SB. nat. Fr. 1892, p. 252.
- Gnathocera hyacinthina, Kr., = (schlütteri, Nonf., infrå), G. vi. Jans., = (cincta, Kr.); KRAATZ, p. 172, Deutsche e. Z. 1892. hyacinthina, Jans., = (hyacinthina, Kr.); KRAATZ, t. c. p. 415.
- G. augustata, p. 128, flavovirens, p. 129, Guinea, Kolbe, S. E. Z. liii schlütteri, Centr. Africa, Nonfried, p. 121, Ent. Nachr. x n. spp.
- Gnorimus: table of characters of the palsearctic species; ZOUFAL, pp & 242, Wien. ent. Z. xi. G. variabilis, var. heydeni discus BECKERS, p. 315, Wien. ent. Z. xi: larva described; FOWLER, M. M. xxviii, p. 242.
- Hadrosticta, n. g., near Argyripa, for II. viridiflua, n. sp., Central Ame Kraatz, p. 378, Deutsche e. Z. 1892.
- Heterorrhina euryrrhina, p. 838, pl. ii, figs. 1 & 2, leonardi, p. 840, fi & 4, Burma, Gestro, Ann. Mus. Genov. (2) x, n. spp.
- Jothochilus, n. g., Anoplochilinæ, p. 136, for J. undulatus, n. sp., Gu p. 137; Kolbe, S. E. Z. liii.
- Leucocelis parallelocollis, lunicollis, p. 66, annulipes, polysticia, p. 6 Africa, Kolbe, SB. nat. Fr. 1892, n. spp.
- Liotrichius, n. g., Trichiides, p. 73, for L. anomala, n. sp., E. Africa, p. Kolbe, SB. nat. Fr. 1892.
- Lomaptera jamesi, n. varr. lixi, p. 366, nobilitatu, p. 368; Nonfried, Ent. Zeit. xxxvi.
- Macrelaphinis kraatzi, Kolbe, note on; Kraatz, p. 415, Deutsche
- M. buttneri, Guinea, kruatzi, Ashanti, Kolbe, S. E. Z. liii, p. 134, n. Macroma superba, figured, pl. ii, fig. 11; Gestro, Ann Mus. Genov. (

- M. camerunica, W. Africa, Kolbe, p. 258, SB. nat. Fr. 1892; M. exclamationis, klingi, Guinea, Kolbe, p. 138, S. E. Z. liii; M. insignis, Burma, Gestro, p. 852, Ann. Mus. Genov. (2) x; M. feistmanteli, S. China, Nonfried, p. 90, Ent. Nachr. xviii: n. spp.
- Macronota: this name must replace Gymnetis; KRAATZ, Deutsche e. Z. 1892, p. 375.
- M. pulchella, Burma, GESTRO, p. 844, Ann. Mus. Genov. (2) x, n. sp.
- Mausoleopsis argentata, Comores Is., Nonfried, p. 124, Ent. Nachr. xviii, n. sp.
- Mecinonota, n. n. for a part of Mucronota, auct.; KRAATZ, p. 375, Deutsche e. Z. 1892.
- Metallesthes subpilosa, Queensland, Nonfried, Berl. Ent. Zeit. xxxvi, n. sp.
- Microvalgus yilgurnensis, W. Australia, BLACKBURN, p. 115, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Mystroceros: this genus is established on an individual manufactured from two genera; KRAATZ, p. 373, Deutsche e. Z. 1892.
- Oreoderus planicollis, p. 862, humilis, p. 863, humeralis, p. 864, waterhousei, p. 865, rufulus, p. 867, brevipennis, p. 868, maculipennis, p. 869, Burma, borneensis, Sarawak, p. 875, Gestro, Ann. Mus. Genov. (2) x, n. spp.
- Oxythyrea stictica, metamorphoses; XAMBEU, p. 178, Ann. Soc. L. Lyon, xxxix.
- Pachnoda stehlini, n. var. fuliginosa; Nonfried, p. 123, Ent. Nachr. xviii. P. rossi, Cameroons, Kolbe, p. 254, SB. nat. Fr. 1892, n. sp.
- Penthima, n. g., Macronotidarum, p. 313, for P. nigerrima, n. sp., Java, p. 314, Kraatz, Deutscho e. Z. 1892.
- Pilinurgus leveillei, E. India, NONFRIED, p. 372, Berl. Ent Zeit. xxxvi, n. sp.
- Plesiorrhina triplagiata, E. Africa, Kolbe, p. 64, SB. nat. Fr. 1892, n. sp.
  Pleuronota, n. g., Macronotidarum, for P. octomaculata, n. sp., Java,
  KRAATZ, p. 312, Deutsche c. Z. 1892.
- Polystalactica sansibarica, E. Africa, Kolbe, p. 75, SB. nat. Fr. 1892; P. stipatrix, p. 132, contempta, p. 133, Guinea, Kolbe, S. E. Z. liii: n. spp.
- Pygora lenoc. (sic) n. var. brzowskii, Nonfried, p. 120, Ent. Nachr. xviii. Rhomborrhina flammea, n. var. cariana, Gestro, p. 837, Ann. Mus. Genov. (2) x.
- Schizorrhina (Diaphonia) suturata, Thursday Is., Queensland, NONFRIED, p. 370, Berl. Ent. Zeit. xxxvi, n. sp.
- Somalibia, Lansb. = (Dolichostethus, Kolbe); KEAATZ, p. 415, Deutsche e. Z. 1892.
- Stenovalgus, n. g., near Microvalgus, p. 141, for S. carinulatus, n. sp., Guinea, Kolbe, p. 142, S. E. Z. liii.
- Stephanocrates, n. g., near Compsocephalus, for S. preussi, n. sp., Cameroons, Kolbe. p. 240, SB. Nat. Fr. 1892.
- Stephanorrhina, species tabulated, pp. 145, 147, S. guttata, p. varr. hybrida, geminata, Kolbe, pp. 148 & 149, S. E. Z. liii.

- S. temeraria, p. 150, Cameroons, haroldi, Angola, p. 153, Kolbe, S. l liii, n. spp.
- Taniodera, varr. of 4-lineata, rufflesiana, biplugiata, antiqua, and p Guér., 2 described; Kraatz, Deutsche e. Z. 1892, pp. 316-T. quadrilineata, Hope and allies, synonymy; Janson, pp. 5: Notes Leyd. Mus. xiv.
- T. rugosicollis, p. 315, Malacca, quadristrigata, Darjeeling, p. 316, p. fig. 9, borneensis, Borneo, p. 317, Kraatz, Deutsche e. Z. 1
  T. goryi, Java, p. 58, virgata, E. India, p. 59, Janson, Notes I Mus. xiv: p. spp.
- Taurrhina longiceps, Guinea, Kolbe, p. 126, S. E. Z. liii, n. sp.
- Tephraa beinlingi, Zambesi, Nonfried, p. 124, Ent. Nachr. xviii, n. : Tmesorrhina iris, n. var. camerunica; Nonfried, p. 120, Ent. N
- T. alpestris, p. 249, barombina, p. 250, Cameroous, Kolbe, SB. nat 1892, n. spp.
- Trichius fusciatus, n. var. albohirtus; REITTER, p. 152, Wien. ent. Z. T. jansonii, Burma, Gestro, p. 854, pl. ii, fig. 12, Ann. Mus. Genov x, n. sp.
- Trigonophorus foveiceps, figured, pl. ii, figs. 7, 8, Ann. Mus. Genov. ( T. fec, Burms, Gestro, p. 841, pl. ii, fig. 5, Ann. Mus. Genov. (2) x,:
- Valgus longulus, p. 855, stictopygus, p. 857, luctuosus, p. 858, tristis, p. trisinuatus, p. 860, doriæ, p. 870, cristatus, p. 871, Sara beccarii, Celebes, p. 873, sumatranus, p. 874, Sumatra, Gestro, Mus. Genov. (2) x; V. thibetanus, Thibet, Nonfried, p. 372, Ent. Zeit. xxxvi: n. spp.
- Xenogenius, n. subg. of Canochilus, q.v.; KOLBE, SB. Nat. Fr. 1892, 1

### BUPRESTIDE.

- [Cf. Blackburn (90, 92, 93, 94, 95), Fairmaire (262, 271), Fleist (284), Germain (322), Kerremans (481 to 489), Nonfried 647, 648), Péringuey (672, 673), Poll (696), Théry (921), Wainouse (988), Xambeu (1021, 1024).]
- Synonymic catalogue of the family, including 4200 species; Kerres Mem. Soc. Ent. Belg. i.
- Reproduction of descriptions of the early stages, with notes on ha &c.; XAMBEU, Rev. d'Ent. xi, pp. 202-252.
- Acmaodero reflexangula, despecta, distinctions of; Reitter, p. 188, V ent. Z. xi.
- A. bella, p. 126, dives, p. 136, E. Africa, Nonfried, Ent. Nachr. x A. smaragdina, Centr. Africa, Kerremans, p. 56, Ann. Ent. 1 xxxvi; A. suareola, p. 42, fullax, p. 43, S. W. Africa, Péringuey S. African Phil. Soc. vi, 2; A. zambeziana, Limpopo, Péring p. 120, t. c.; A. interrupta, Bongal, Kerremans, p. 175, Ann. Belg. xxxvi: n. spp.

- Edisterna namaqua, p. 115, livida, p. 116, modesta, limbata, p. 117, S. Africa, Péringuey, Tr. S. African Phil. Soc. vi, 2, n, spp.
- Agrilus sinuatus, p. 9, roscidus, p. 17, metamorphoses; XAMBEU, Mœurs, ii.
- A. carifrons, Obock, FAIRMAIRE, p. 100, Rev. d'Ent. xi; A. niveoguitatus, p. 210, birmanicus, p. 211, lafertei, p. 212, decoloratus, p. 213, villosostriatus, with var. carmineus, p. 214, imbellis, inops, p. 215, mandaricus, p. 216, mixtus, p. 217, ineptus, neelgheriensis, p. 218, E. India, Kerremans, Ann. Ent. Belg. xxxvi; A. ostrinus, p. 819, agnatus, p. 820, modicus, p. 821, albopictus, p. 822, dianthus, nubilus, p. 823, inamænus, p. 824, livens, p. 825, Burma, Kerremans, Ann. Mus. Genov. (2) xii; A. riolaceus, cyaneomicans, S. China, Nonfried, p. 91, Ent. Nachr. xviii; A. striatocollis, Philippines, Kerremans, p. 24, Ann. Soc. Ent. Fr. lxi; A. terræreginæ, Queensland, Blackburn, p. 220, Tr. R. Soc. S. Austr. xv: n. spp.
- Amorphosomus pectoralis, E. India, KERREMANS, p. 210, Ann. Ent. Belg. xxxvi; A. spathatum, Burma, KERREMANS, p. 818, Ann. Mus. Genov. (2) xii: n. spp.
- Ancylochira salomonii, Th., synonymy of; HEYDEN, p. 111, Deutsche e. Z. 1892.
- Anthaxia hypomelana, metamorphoses; XAMBEU, p. 190, Ann. Soc. L. Lyon, xxxix. A. morio, Herbst., = (similis, Saund.); REITTER, p. 136, Wien. ent. Z. xi.
- A. semilimbata, Amasia, Fleischer, p. 209, Wieu. ent. Z. xi; A. obockiona, Africa, Fairmaire, p. 99, Rev. d'Ent. xi; A. capitata, morosa, India, Kerremans, p. 173, Ann. Ent. Belg. xxxvi: n. spp.
- Aphanisticus metallescens, E. India, KERREMANS, p. 219, Ann. Ent. Belg. xxxvi; A. cephalicus, Burma, KERREMANS, p. 826, Ann. Mus. Genov. (2) xii; A. perraudierei, Cochin China, Poll., p. 19, Ann. Soc. Ent. Fr. lxi; A. cupricornis, New Guinea, KERREMANS, p. 1004, Ann. Mus. Genov. (2) xii: n. spp.
- Astraus simplex, S. Australia, BLACKBURN, p. 211, Tr. R. Soc. S. Austr. xv, n. sp.
- Belionota sphenopteroides, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. vi, 2, p. 43; B. ignicollis, Anam, Nonfried, p. 374, Berl. Ent. Zeit. xxxvi: n. spp.
- Bubastes ragans, S. Australia, BLACKBURN, p. 213, Tr. R. Soc. S. Austr. xv, n. sp.
- Bubastodes, n. g., near Bubastes, for B. sulcicollis, n. sp., S. Australia, Blackburn, p. 212, Tr. R. Soc. S. Austr. xv.
- Buprestis impressicollis, India, KERREMANS, p. 172, Ann. Ent. Belg. xxxvi, n. sp.
- Capuodis tenebrionis, metamorphoses; Xambeu, p. 183, Ann. Soc. L. Lyon, xxxix.
- Cutozantha bicolor, n. var. cyanura, KERREMANS, p. 171, Ann. Ent. Belg.

- Chulcogenia viriditarsis, W. Africa, Nonfried, p. 125, Ent. N xviii, n. sp.
- Chalcophora patricia, p. 118, confusa, p. 119, S. Africa, Péringue S. African Phil. Soc. vi (2), n. spp.
- Chalcotænia gigantea, New Guinea, Nonfried, p. 374, Berl. Eut. xxxvi; C. cupreosuturata, Abyssinia, Fairmaire, p. 101, Rev. xi: n. spp.
- Chrysobothrys nana, Turkestan, FAIRMAIRE, p. clii, Bull. Soc. En lxi; C. carinata, p. 197, quadraticollis, p. 198, riolacea, p. 198, India, Kerremans, Ann. Ent. Belg. xxxvi: n. spp.
- Chrysochroa limbata, Borneo, Nonfried, p. 373, Berl. Ent. Zeit. 3 n. sp.
- Chrysodema navicularis, p. 21, riolacea, p. 22, hebes, p. 23, Philip Kerremans, Ann. Soc. Ent. Fr. lxi, n. spp.
- Coræbus rubi, metamorphoses; XAMBEU, Mœurs, ii, p. 15.
- C. dorsalis, caruleus, p. 200, cyancopictus, p. 201, chloropictus, mod p. 202, fulgidiceps, smaragdineus, p. 203, E. India, KERREMANS, Ent. Belg. xxxvi; C. feæ, p. 811, fossulatus, p. 812, gestroi, p Burms, KERREMANS, Ann. Mus. Genov. (2) xii: n. spp.
- Cryptodactylus cyaneoniger, p. 207, scutellaris, cuprascens, p. 209, E. 1 Kerremans, Ann. Ent. Belg. xxxvi, n. spp.
- Curis discondulis, W. Australia, BLACKBURN, p. 214, Tr. R. Soc. S. Axv, n. sp.
- Cylindromorphus orientalis, Manilla, KERREMANS, p. 26, Ann. Soc. Fr. lxi, n. sp.
- Cyphogastra plicata, Philippines, Kerremans, p. 23, Ann. Soc. Enlxi; C. abdominalis, Damma I., Waterhouse, p. 412, Ann. 1 (6) x: n. spp.
- Cyria tridens, Australia, BLACKBURN, p. 41, Tr. R. Soc. S. Austin. sp.
- Discoderes quadricornis, Abyssinia, FAIRMAIRE, p. 100, Rev. d'En
- Endelus coraboides, p. 219, aureocupreus, p. 220, E. India, KERREI Ann. Ent. Belg. xxxvi; E. curtus, Tenasserim, KERREMANS, p. Ann. Mus. Genov. (2) xii: n. spp.
- Gulbella globosa, Bengal, KERREMANS, p. 226, Ann. Ent. Belg. x: n. sp.
- Hypostigmodera, n. g., near Stigmodera, for H. variegata, n sp., Quiland; Blackburn, p. 215, Tr. R. Soc. S. Austr. xv.
- Iridotænia cingulata, Zanzibar, KERREMANS, p. 51, Ann. Ent. 1 xxxvi, n. sp.
- Julodie abeillei, Taurus, Therr, p. celviii, Bull. Soc. Ent. Fr. lxi cyphodera, Turkestan, Fairmaire, p. clii, Bull. Soc. Ent. Fr. n. spp.
- Melibaus laripennis, riolaceicollis, W. Africa, KERBEMANS, p. 302, 1 Soc. Ent. Fr. lxi; M. indicolus, p. 204, chrysomelinus, aurarius, p. E. India, KERREMANS, Ann. Ent. Belg. xxxvi; M. delauneyi, 1

- Poll, p. 19, Ann. Soc. Ent. Fr. lxi; M. transversus, p. 814, saundersi = (cupricollis, Saund.), venustus, p. 815, magnificus, p. 816, Burma, Kerbemans, Ann. Mus. Genov. (2) xii: n. spp.
- Melobasis cærulea, Engano, KERREMANS, p. 1000, Ann. Mus. Genov. (2) xii; M. incerta, ignicauda, New Guinea, KERREMANS, p. 1002, Ann. Mus. Genov. (2) xii; M. monticola, p. 496, beltanensis, p. 499, Australia, Blackburn, P. Linn. Soc. N.S.W. (2) vi; M. subcyanea, S. Australia, Blackburn, p. 288, op. cit. (2) vii; M. derbyensis, N. W. Australia, Blackburn, p. 44, Tr. R. Soc. S. Austr. xv; M. elderi, Fraser River, Blackburn, p. 35, op. cit. xvi: n. spp.
- Neocuris: generic characters; BLACKBURN, p. 42, Tr. R. Soc. S. Austr.
- N. dilataticollis, p. 42, nigricans, p. 43, N. S. Wales, BLACKBURN, Tr. R. Soc. S. Austr. xvii, n. spp.
- Neobubastes, n. g., near Bubastes, for N. aureocincta, n. sp., Centr. Australia; Blackburn, p. 213, Tr. R. Soc. S. Austr. xv.
- Paracupta calestis, New Guinea, KERREMANS, p. 1001, Ann. Mus. Genov. (2) xii, n. sp.
- Paratænia, n. g., p. 52, for P. inornata, aspera, p. 54, orbicularis, simplicicollis, p. 55, n. spp., W. Africa, and including Chrysodema chrysochlora, Pall., and opaca, Lansb.: Kerremans, Ann. Ent. Belg. xxxvi.
- Philanthaxia latifrons, India, KERREMANS, p. 173, Ann. Ent. Belg. xxxvi; P. splendida, Saigon, Poll., Ann. Soc. Ent. Fr. lxi, p. 17: n. spp.
- Pacilonota (Lampro) festiva, metamorphoses; XAMBEU, p. 188, Ann. Soc. L. Lyon, xxxix.
- P. magnifica, Bengal, KERREMANS, p. 171, Ann. Ent. Belg. xxxvi, n. sp.
- Polycesta curta, Bengal, KERREMANS, p. 174, Ann. Ent. Belg. xxxvi; P. paulseni, Chili, GERMAIN, p. 243, Act. Soc. Chili, ii: n. spp.
- Pseudochrysodema (?) walkeri, Damma I., WATERHOUSE, p. 411, Ann. N. H. (6) x, n. sp.
- Psiloptera stormsi, Centr. Africa, Kerremans, p. 56, Ann. Ent. Belg. xxxvi; P. hilaris, S.W. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2), p. 41: n. spp.
- Sambus gibbicollis, p. 206, melanoderus, nigritus, p. 207, E. India, Kerremans, Ann. Ent. Belg. xxxvi; S. coloratus, p. 816, femoralis, p. 817, Burma, Kerremans, Ann. Mus. Genov. (2) xii; S. loria, New Guinea, Kerremans, p. 1003, Ann. Mus. Genov. (2) xii: n. spp.
- Sphenoptera potanini, Jak., = (procera, Reit.); REITTER, p. 136, Wien. ent. Z. xi.
- S. latescutata, Turkestau, FAIRMAIRE, p. cliii, Bull. Soc. Ent. Fr. lxi; S. leonensis, W. Africa, p. 137, nervosa, Centr. Africa, p. 138, howa, Madagascar, p. 139, Nonfried, Ent. Nachr. xviii; S. curta, p. 180, impressa, p. 182, inocua, p. 187, depressa, konbirensis, p. 188, crebrepunctata, p. 189, fulgidiceps, p. 190, auricollis, p. 191, deducta, p. 192, gossypii, p. 195, E. India, Kerremans, Ann. Ent. Belg. xxxvi: n. spp.
- Sternocera orissa, n. var. modesta, Péringuey, p. 40, Tr. S. African Phil. Soc. vi (2), n. sp.

Stigmodera minuta, Queensland, p. 44, septemmaculata, S. Australia, p. skusei, N. S. Wales, hostilis, W. Australia, p. 46, sternalis, S. Aust p. 47, macleayi, Queensland, equina, S. Australia, p. 48, quadrine Victoria, p. 49, dispar, p. 50, canaliculata, Queensland, arbor. Australia, p. 51, marcida, W. Australia, p. 52, carpentariæ, Qu land, ornata, p. 53, longula, p. 54, Victoria, Blackburn, Tr. R. S. Austr. xv; S. cara, p. 216, insignis, filiformis, p. 217, capu regia, p. 218, ignea, p. 219, Australia, Blackburn, t. c.; S. e Victoria Desert, Blackburn, p. 36, op. cit. xvi: n. spp.

Strigoptera frenchi, p. 500, australis, marmorata, p. 501, Australia, Bl. Burn, P. Linn. Soc. N.S.W. (2) vi, n. spp.

Trachys fleutiauxi, Hué, Poll, p. 19, Ann. Soc. Ent. Fr. lxi; T. anei p. 221, subviolacea, p. 222, flaviceps, obliqua, p. 223, transversa, b. nata, p. 224, stigmatica, integra, p. 225, Kerremans, Ann. Ent. I xxxvi; T. penicillata, p. 827, virescens, lyra, p. 828, nigra, p. vexator, rudis, p. 830, polita, p. 831, Burma, Kerremans, Ann. Genov. (2) xii; T. rugata, New Guinea, Kerremans, p. 1005, T. viridula, rufescens, Philippines, Kerremans, p. 25, Ann. Soc. Fr. lxi: n. spp.

## THROSCIDÆ, MONOMMIDÆ, EUCNEMIDÆ.

[Cf. Blackburn (90, 93), Rost (780).]

Aspathines ovatus, habits; SCHWARZ, P. E. Soc. Wash. ii, p. 241.

Drupetes sulcatus, Caucasus, Rost, Deutsche e. Z. 1892, p. 402, n. sp. Dromæolus interioris, S. Australia, Blackburn, p. 502, P. Linn. N.S.W. (2) vi. n. sp.

Dyscolocerus heros, N. S. Wales, BLACKBURN, p. 56, Tr. R. Soc. S. A xv, n. sp.

#### ELATERIDE.

- [Cf. Angell (14), Berg (57), Blackburn (90, 92, 93), Candèze (15 156), Casey (161), Fairmaire (262, 270), Fleutiaux (2 Frivaldszky (309), Gorham (352), Péringuey (672), Reij (725), Schwarz (809, 810, 811).]
- Adelocera carinensis, vicina, Burma, CANDÈZE, p. 772, Ann. Mus. Ge (2) x; A. cognata, Eugano, CANDÈZE, p. 796, op. cit. xii : n. spp. Adiaphorus modestus, Bengal, CANDÈZE, p. 488, Ann. Ent. Belg. xi n. sp.
- \*\*Eolus queenslandicus, N. Queensland, BLACKBURN, p. 296, P. Linn. N.S.W. (2) vii; \*\*E. pyroblaptus, Argentine Rep., Berg, p. 61, Soc. Arg. xxxiii: n. spp.
- Agonischius humilis, p. 493, limbatus, p. 494, E. India, CANDÈZE, 1 Ent. Belg. xxxvi; A. lætus, p. 787, diversus, conjugatus, p. simplex, 789, Burma, CANDÈZE, Ann. Mus. Genov. (2) x: n. spp

- Agrypnus fallaciosus, dicergens, Obock, FAIRMAIRE, p. 101, Rev. d'Ent. xi: A. labeculatus, Engano, CANDÈZE, p. 795, Ann. Mus. Genov. (2) xii: n. spp.
- Alaus mahenus, Seychelles, FAIRMAIRE, p. cli, Bull. Soc. Ent. Fr. lxi;
  A. modiglianii, Engano, CANDÈZE, p. 797, Ann. Mus. Genov. (2) xii;
  A. loriæ, New Guinea, CANDÈZE, p. 802, t. c. : p. spp.
- Anchastus bengalensis, India, CANDÈZE, p. 490, Ann. Ent. Belg. xxxvi; A. insulsus, New Guinea, CANDÈZE, p. 804, Ann. Mus. Genov. (2) xii : n. spp.
- Aptopus birmanicus, Burma, CANDÈZE, p. 781, Ann. Mus. Genov. (2) x, n. sp.
- Athous olbiensis, = (chamboreti); BUYSSON, p. xxiv, Bull. Soc. Ent. Fr. lxi. A. proximus and spectabilis, Hampe, are one species; KRAATZ, p. 176, Deutsche e. Z. 1892.
- sanguinicollis, Japan, FRIVALDSZKY, p. 124, Term. füzetek, xv, n. sp.
- Cumpsosternus pasteuri, Nias Is., CANDÈZE, Notes Leyd. Mus. xiv, p. 9, n. sp.
- Curdiophorus rufipes, n. var. persicus; SCHWARZ, p. 381, Deutsche e. Z. 1892.
- C. analis, Asia Minor, pellitus, Turkestan, Schwarz, Deutsche e. Z. 1892; C. macer, Bengal, Candèze, p. 493, Ann. Ent. Belg. xxxvi; C. seminalis, Burma, Candèze, p. 778, Ann. Mus. Genov. (2) x; C. rictoriensis, eucalypti, Australia, Blackburn, P. Linn. Soc. N.S.W. (2) vi, p.: n. spp.
- Cardiotarsus antennalis, Burma, CANDÈZE, p. 780, Ann. Mus. Genov. (2) x, n. sp.
- Chalcolepidius apacheanus, Arizona, Casey, Ann. N. York Ac. vi, p. 16, n. sp.
- Corymbites vireus, n. var. stramineus; CANDÈZE, p. 228, Ann. Ent. Belg. xxxvi. C. caricinus, umbricola, lobatus, characters of; Hamilton, p. 296, Canad. Ent. xxiv.
- C. infirmus, Siberia, REITTER, p. 152, Wien. ent. Z. xi; C. weidtii, British Columbia, ANGELL, p. 84, Ent. News, iii: n. spp.
- Cryptohypnus sericeus, p. 490, pictus, p. 491, E. India, Candèze, Ann. Ent. Belg. xxxvi, n. spp.
- Dactylophysus, n. g., p. 408, for D. mendax, n. sp., Brazil, p. 410, and including Heterocrepidius tibialis, Cand.; FLEUTIAUX, Ann. Soc. Ent. Fr. lxi.
- Diploconus rufulus, p. 781, nutritus, ornatus, serricornis, p. 782, Burma, Candèze, Ann. Mus. Genov. (2) x; D. enganensis, Engano, Candèze, p. 799, op. cit. xii: n. spp.
- Elater miniatus (= pomonæ, auct., nec Steph.), England, Gorham, Ent. xxv, p. 304; E. wentworthensis, N. S. Wales, Blackburn, p. 297, P. Linn. Soc. N.S.W. (2) vii: n. spp.
- Glypheus alpinus, Victoria, BLACKBURN, p. 290, P. Linn. Soc. N.S.W. (2) vii, n. sp.

- Glyphochilus montanus, N. S. Wales, BLACKBURN, p. 293, P. Lini N.S.W. (2) vii, n. sp.
- Glyphonyz suturalis, Bengal, CANDEZE, p. 494, Ann. Ent. Belg. 3 G. correctus, Burma, CANDEZE, p. 789, Ann. Mus. Genov. (n. spp.
- Hemirrhaphes ruficollis, Bengal, CANDEZE, p. 491, Ann. Ent. xxxvi, n. sp.
- Heteroderes canus, Bengal, Candezz, p. 489, Ann. Ent. Belg. n. sp.
- Lacon senilis, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. p. 44; L. tabularius, Bengal, Candèze, p. 483, Ann. Ent. xxxvi; L. collisus, p. 772, gypsatus, p. 773, hilaris, subcostatus, Burma, Candèze, Ann. Mus. Genov. (2) x; L. indutissimus, E Candèze, p. 797, op. cit. xii; L. pupillus, New Guinea, Ca. p. 802, t. c.; L. brightensis, p. 503, murrayensis, lindensis, 1 adelaida, p. 505, duplex, p. 506, eucalypti, p. 507, andersoni, Australia, Blackburn, P. Linn. Soc. N.S.W. (2) vi; L. man. p. 288, squalescens, yilgarnensis, p. 289, Australia, Blackburn, vii: n. spp.
- Ludius clavus, Burma, Candèze, p. 786, Ann. Mus. Genov. (2) x, r Megapenthes aspernendus, Bengal, Candèze, p. 489, Ann. Ent. xxxvi; M. litteratus, Java, remotus, Celebes, Candèze, p. 10, Leyd. Mus. xi; M. musicatus, Engano, Candèze, p. 798, Ann Genov. (2) xii; M. brunniventris, New Guinea, Candèze, j. t. c.: n. spp.
- Melanotus: revision of the palmarctic species; Schwarz, pp. 14 pl. ii, Deutsche e. Z. 1892.
- M. kraatsi, Greece, p. 155, humilis, Alai Mts., p. 156, fragilis, Turp. 160, brunnicornis, China, p. 164; Schwarz, Deutsche e. Z. 18 repletus, p. 783, comatus, conicus, p. 784, Burma, Candèze, Angenov. (2) x; M. hericius, Engano, Candèze, p. 800, op. cn. spp.
- Melanozanthus anticus, Bengal, CANDÈZE, p. 490, Ann. Ent. Belg.

  M. virgatus, p. 777, punctum, p. 778, Burma, CANDÈZE, Ann
  Genov. (2) x; M. palliatus, sexguttatus, Engano, CANDÈZE,
  t. c.: n. spp.
- Meristhus indecorus, Bengal, CANDÈZE, p. 485, Ann. Ent. Belg. n. sp.
- Monocrepidius mitigatus, Engano, Candèze, p. 798, Ann. Mus. Gerxii; M. disjunctus, New Guinea, Candèze, p. 803, t. c.; M. rup. 509, frontalis, p. 510, alpicola, p. 511, macleayi, p. 512, ove baldiensis, p. 514, Australia, Blackburn, P. Linn. Soc. N.Svi; M. ingens, p. 291, terræreginæ, p. 292, picticollis, narrap. 293, mentitor, p. 294, olliffi, p. 295, Australia, Blackburn, vii: n. spp.
- Otocryptus, n. g., forming a new group Otocryptites, p. 486, for O. c. n. sp., Bengal; Candèze, Ann. Ent. Belg. xxxvi.

- Pantolamprus ovampo, S.W. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2) p. 45, n. sp.
- Penia plagiata, Burma, CANDEZE, p. 786, Ann. Mus. Genov. (2) x, n. sp. Pericus discedens, Bengal, CANDEZE, p. 485, Ann. Ent. Belg. xxxvi, n. sp. Physodactylus foreatostriatus, p. 406, oberthuri, niger, p. 407, brasiliensis, p. 408, Brazil, FLEUTIAUX, Ann. Soc. Ent. Fr. lxi, n. spp.
- Silesis inflectus, p. 494, nigriceps, p. 495, Bengal, Candèze, Ann. Ent. Belg. xxxvi, n. spp.
- Singhalenus gibbus, Bengal, CANDÈZE, p. 487, Ann. Ent. Belg. xxxvi, n. sp. Spheniscosomus, n. g., for a part of Melanotus; Schwarz, p. 132, Wien. ent. Z. xi.
- Teslasena, n.g., Physodactylini, for Anelustes femoralis, Luc.; FLEUTIAUX, p. 410, Ann. Soc. Ent. Fr. lxi.

#### DASCILLIDÆ.

- [Cf. Blackburn (90, 92), Bourgeois (113).]
- Dascillus calvescens, ruforillosus, Bengal, BOURGEOIS, p. 234, Ann. Ent. Belg. xxxvi, n. spp.
- Helodes princeps, p. 517, cinctus, olliffi, p. 518, montivagans, p. 519, H. (Cyphon?) pictus, ovensensis, frater, p. 520, adelaidæ, spilotus, p. 521, lindensis, p. 522, Australia, BLACKBURN, P. Linn. Soc. N.S.W. (2) vi; H. (Cyphon) fenestratus, N. S. Wales, BLACKBURN, op. cit. vii: n. spp.
- Macrohelodes, n. g., for M. lucidus, crassus, p. 298, intricatus, gravis, p. 299, n. spp., Australia, and including Helodes princeps (suprd); BLACKBURN, P. Linn. Soc. N.S.W. (2) vii.
- Sclerocyphon, n. g., p. 522, for S. maculatus, n. sp., Victoria, p. 523; BLACKBURN, P. Linn. Soc. N.S.W. (2) vi.
- Scirtes helmsi, Victoria, BLACKBURN, p. 524, P. Linn. Soc. N.S.W. (2) vi, n. sp.

#### MALACODERMATA.

- [Cf. Blackburn (90, 94, 95), Bourgeois (112, 113, 114, 210), Gestro (326, 329, 334), Gorham (210, 350), Olivier (652, 653), Péringuey (672), Weise (1000).]
- List of Lycidæ of Borneo; Bourgeois, p. 42, Notes Leyd. Mus. xiv.
- Calochromus: the structure of max. palpi differs sexually: Blackburn, p. 528, P. Linn. Soc. N.S.W. (2) vi.
- Chauliognathus marginatus, eggs and oviposition; RILEY, p. 186, Canad. Ent. xxiv.
- Cladophorus (Odontocerus) flabellifer, p. 496, sellatus, beccarii, p. 498, quadraticollis, pectinatus, p. 500, dorsalis, p. 501, New Guinea, diabolicus, Amboyna, belzebuth, p. 502, stygius, infernalis, p. 503, acheronticus, p. 504, pallidulus, parallelus, p. 505, rubentipes, p. 506, mandibularis, p. 507, ornaticollis, p. 508, tibialis, rufithorax, p. 509, dimidiatus, p. 510, septemareolatus, p. 511, papuensis, p. 512, acutangulus, obsoletus, p. 513, New Guinea, Bourgeois, Ann. Mus. Genov. (2) xii, n. spp.
  - 1892. [vol. xxix.]

- Dasytes squirensis, helmsi, Australia, BLACKBURN, p. 38, Tr. R. Soc. S. Austr. xvi, n. spp.
- Diaphanes mendax, p. 597, plagiator, p. 598, Burma, OLIVIER, Ann. Mus. Genov. (2) x; D. apicalis, Borneo, Gorham, p. 89, pl. iv, fig. 8, P. Z. S. 1892: n. spp.
- Ditua nigricornis, p. 514, alternata, fluviceps, p. 515, New Guines, Bourgeois, Ann. Mus. Genov. (2) xii, n. spp.
- Ellychnia californica, pupa noticed; WICKHAM, p. 201, Bull. Lab. Iowa, ii.
- Hapalochrus opulentus, S. W. Africa, Péringuey, Tr. S. African Phil, Soc. vi (2), p. 46, n. sp.
- Hedybius amænus, Transvaal, Gorham in Distant, Naturalist Transvaal, p. 197, pl. i, fig. 2, n. sp.
- Heteromastix anticus, p. 221, dilataticollis, p. 222, N. S. Wales, BLACKBURN, Tr. R. Soc. S. Austr. xv, n. spp.
- Ichthyurus urospilus, p. 561, luctuosus, p. 562, feæ, p. 564, longicauda, p. 566, nigromaculatus, p. 569, montanus, p. 571, nigriceps, p. 573, agilis, p. 574, spinicrus, p. 576, pallidus, p. 578, vittatus, p. 579, laniger, p. 581, propomacrus, p. 583, ritsemæ, p. 588, crassicauda, p. 589, laticauda, p. 590, Burma, Gestro, Ann. Mus. Genov. (2) x; I. bourgeoisii, China, p. 1023, mouhoti, p. 1025, hirundo, p. 1026, obscurus, p. 1028, Siam, vandepollii, China, p. 1029, wallacei, Singapore, p. 1031, borneensis, p. 1032, Borneo, insularis, p. 1033, curvicauda, p. 1035, pachygaster, p. 1036, Singapore, macrurus, Cochin China, p. 1037, davidii, Kiang-Si, p. 1038, oberthurii, Burma, p. 1039, malayanus, Java, p. 1041, melanospilus, Sylhet, p. 1042, lucassenii, Borneo, p. 1043, beccarii, Sumatra, p. 1045, abnormis, Darjeeling, p. 1046, afer, Niger, p. 1047, Gestro, t.c.: n. spp.
- Idgia flavibuccis, Bengal, Bourgeois, p. 237, Ann. Ent. Belg. xxxvi, n. sp. Laius femoralis, eyrensis, p. 531, pretiosus, p. 532, Australia, Blackburn, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Lamprophorus morator, Burma, OLIVIER, p. 596, Ann. Mus. Genov. (2) x, n. sp.
- Lampyris bicarinata, Muls., note on; OLIVIER, Bull. Soc. Ent. Fr. lxi, p. xlv.
- Luciola æmula, p. 601, seriata, p. 602, ægrota, p. 603, abscondita, p. 604, Burma, OLIVIER, Ann. Mus. Genov. (2) x; L. cribellata, p. 1010, pupilla, 1011, New Guinea, OLIVIER, t. c., n. spp.
- Lycus distanti, Transvaal, Bourgeois in Distant, Naturalist Transvaal, p. 196, pl. i, fig. 3, n. sp.
- Malachius, internal structure of apices of elytra; HOFFBAUER (405).

  M. inornatus, metamorphoses; XAMBEU, Mœurs, ii, p. 44, and Le Nat. 1892, p. 157.
- Maltharchus, n. g. for Malthodes brevicollis and allies; Weise, p. 408, Deutsche e. Z. 1892.
- Malthodes: characters of the French species with figures; BOURGEOIS, Faune gallo-rhenane, pp. 173-208.

- M. caudatus, Carinthia, Weise, p. 410, Deutsche e. Z. 1892, n. sp.
- Metriorhynchus gigas, p. 524, semicostatus, p. 525, monticola, occidentalis, p. 526, Letus, p. 527, Australia, BLACKBURN, P. Linn. Soc. N.S.W. (2) vi, n. spp.
- Phengodes laticollis, Q noticed; RILEY & ULKE, P. E. Soc. Wash. ii, p. 227.
- Plateros? cribripennis, Bengal, BOURGEOIS, p. 236, Ann. Ent. Belg. xxxvi, n. sp.
- Pyrocælia lumpyroides, p. 598, microceras, cribripennis, p. 599, feai, p. 600, Burma, Olivier, Ann. Mus. Genov. (2) x; P. collaris, Borneo, Gorham, p. 89, pl. iv, fig. 7, P. Z. S. 1892: n. spp.
- Rhagonycha nigripes, metamorphoses; XAMBEU, p. 28, Mœurs, ii.
- Selasia robecchii, Somaliland, Gestro, p. 759, Ann. Mus. Genov. (2) xii, n. sp.
- Selenurus variegatus, p. 220, sydneyanus, p. 221, N. S. Wales, BLACKBURN, Tr. R. Soc. N.S.W. xv, n. spp.
- Silis australis, S. Australia, BLACKBURN, p. 530, P. Linn. Soc. N.S.W. (2) vi, n. sp.
- Telephorus victoriensis, p. 528, guleatus, fusicornis, p. 529, Australia, BLACKBURN, P. Linn. Soc. N.S.W. (2) vi, n. spp.
- Trichalus hypocrita, p. 40, flavidus, p. 41, Borneo, Bourgeois, Notes Leyd. Mus. xiv, n. spp.
- Vesta aurantiaca, figured, pl. iv, fig. 9, noticed, p. 89, GORHAM, P. Z. S. 1892.
- Xylobanus reticulatus, figured, pl. iv, fig. 6, P. Z. S. 1892.
- X. ritsemæ, Borneo, Bourgeois, p. 39, Notes Leyd. Mus. xiv. n. sp.

### CLERIDÆ.

- [Cf. Abeille (2), Blackburn (94), Duvivier (231), Eschericii (259), Fairmaire (262, 263), Gestro (334), Gorham (349), Lewis (556).]
  Callimerus elegans, p. 720, suavis, p. 722, feæ, p. 723, gracilis, faustus,
- P. 724, festivus, p. 725, pretiosus, pallidus, p. 727, latesignatus, p. 728, Burma, Gorham, Ann. Mus. Genov. (2) xii, n. spp.
- Cladiscus obeliscus, Japan, Lewis, p. 185, Ann. N. H. (6) x, n. sp.
- Denops plagiatus, Obock, FAIRMAIRE, p. 102, Rev. d'Ent. xi, n. sp.
- Isoclerus, n. g., near Thaneroclerus, for I. pictus, n. sp., Japan; Lewis, Ann. N. H. (6) x, p. 191.
- Lemidia pictipes, simulans, p. 222, munda, soror, pulchella, p. 223, angustula, leoparda, p. 224, Australia, BLACKBURN, Tr. R. Soc. S. Austr. xv, n. spp.
- Lyctosoma, n. g., near Thaneroclerus, for L. parallelum, n. sp., Japan;
  Lewis, p. 192, Ann. N. H. (6) x.
- Neoclerus, n. g., near Thuneroclerus, for N. ornatulus, n. sp., Japan; Lewis, p. 190, Ann. N. H. (6) x.
- Neohydnus, n. g., near Hydnocera, p. 742, for N. despectus, n. sp., Burma, p. 743; GORHAM, Ann. Mus. Genov. (2) xii.

- Nodepus, n. g., Tillides, p. 719, for N. conicicollis, n. sp., Burma, I Gorham, Ann. Mus. Genov. (2) xii.
- Omnadius parciceps, p. 741, tricinctus, p. 742, Burma, GORHAM, Ant Genov. (2) xii; O. nigromaculatus, Japan, Lewis, p. 187, Ann (6) x: n. spp.
- Opetiopalpus karenicus, Burma, Gorham, p. 746, Ann. Mus. Ger xii, n. sp.
- Opilo foreicollis, Algeria, ABEILLE, p. ccxxxvi, Bull. Soc. Ent. 1
  O. mirus, Somaliland, Gestro, p. 761, Ann. Mus. Genov. (
  O. hypocaustus, Burma, Gorham, p. 730, t. c.; O. longipilis,
  FAIRMAIRE, p. 105, Rev. d'Ent. xi; O. carinatus, niponicus,
  Lewis, p. 186, Ann. N. H. (6) x: n. spp.
- Orthopleura funebris, Syria, FAIRMAIRE, p. 148, Ann. Ent. Belg. xxxv Orthrius fex, p. 736, tarsalis, p. 737, Burma, Gorham, Ann. Mus. (2) xii, n. spp.
- Pelonium lividum, p. 744, discrepens, p. 745, Burma, Gorham, An-Genov. (2) xii, n. spp.
- Phlæocopus mediozonatus, Obock, FAIRMAIRE, p. 103, Rev. d'Ent. ? Pseudachlamys, n. g., p. 297, for P. penicillatus, n. sp., Congo, ; DUVIVIER, Ann. Ent. Belg. xxxvi.
- Spinoza, n.g., near Cymatodera, p. 184, for S. cærulea, n. sp., p. 185, LEWIS, Ann. N. H. (6) x.
- Stigmatium obockianum, Africa, FAIRMAIRE, p. 103, Rev. d'E. S. birmanicum, Burma, GORHAM, p. 740, Ann. Mus. Genov. (: n. spp.
- Tenerus biplagiatus, Obock, FAIRMAIRE, p. 104, Rev. d'Ent. cyaneus, maculicollis, higonius, Japan, Lewis, p. 189, Ann. (6) x: n. spp.
- Thanasimus nigricollis, albomaculatus, Japan, Lewis, p. 187, Ann (6) x, n. spp.
- Thaneroclerus quinquemaculatus, Burma, Gobham, p. 738, Ani Genov. (2) xii; T. aino, Japan, Lewis, p. 190, Ann. N. H. n. spp.
- Tillicera cleroides, p. 731, bibalteata, p. 732, Burma, Gorham, An Genov. (2) xii : n. spp.
- Tillus notatus, Kl., = (lewisii, Kies.); Lewis, p. 185, Ann. N. H. T. pectinicornis, Syris, Abeille, p. ccxxxvi, Bull. Soc. Ent. T. birmanicus, Bhamo, Goriiam, p. 729, Ann. Mus. Genov. (
- Trichodes sipylus and allies, synonymy; ESCHERICH, p. 244, ent. Z. xi; T. syriacus and allies, synonymy; ESCHERICH, Soc. Ent. vii.
- T. suspectus, Syria, p. 226, pulcherrimus, Persia, p. 227, maxim junctus, Syria, p. 229, heydeni, Caucasus, p. 230, ESCHERICH ent. Z. xi, n. spp.
- Xenorthrius, n. g., p. 733, for X. mouhoti, p. 734, subfusciatus, be p. 735, n. spp., Burma; Gorham, Ann. Mus. Genov. (2) xii.

### BOSTRYCHIDÆ, PTINIDÆ, CIOIDÆ.

- [Gf. Casey (161), Blackburn (95), Fairmaire (262, 264), Frivaldszky (309), Germain (322), Xambeu (1023).]
- Notes on the Anobiidæ of Chili; GERMAIN, pp. 254-259, Act. Soc. Chili, ii.
- Apate rufo-coronata, Obock, FAIRMAIRE, p. 104, Rev. d'Ent. xi, n. sp.
- Bostrichus sulcicollis, structural peculiarities and generic position discussed; GERMAIN, p. 259, Act. Soc. Chili, ii.
- Calymmoderus, recharacterised; GERMAIN, p. 258, Act. Soc. Chili, ii.
- Diplocotes niger, Poll, and Diphobia familiaris, Olliff, are one species; BLACKBURN, p. 300, P. Linn. Soc. N.S.W. (2) vii.
- Dysides platensis, Argentine Rep., FAIRMAIRE, p. 245, Ann. Ent. Belg. xxxvi, n. sp.
- Lyctus carolinæ, p. 13, californicus, p. 14, curtulus, p. 15, N. America, Casey, Ann. N. York Ac. vi, n. spp.
- Ocelliger, Phil., = (Eurhopalus, Sol.); GERMAIN, p. 254, Act. Soc. Chili, ii.
- Pachotelus fuscus, Sol., = (Anobium hamorrhoidale, Phil.); GERMAIN, p. 256, Act. Soc. Chili, ii.
- Ptinus sexpunctatus, metamorphoses; Xambeu, p. 36, Mœurs, ii, and NICOLAS, Échange, 1892, p. 143, pl. P. brunneus, metamorphoses; Xambeu, Mœurs, ii, p. 42.
- P. senilis, Obock, FAIRMAIRE, p. 105, Rev. d'Ent. xi, n. sp.
- Sitodrepa panicea, habits; ROTH, Ent. Tidskr. xiii, p. 254.
- Trogoxylon merged in Lyctus; Casey, p. 12, Ann. N. York Ac. vi.
- Xyletinus (Calypterus) chinensis, China, FRIVALDSZKY, p. 124, Term. füzetek, xv, n. sp.
- Xylopertha pustulata, metamorphoses; XAMBEU, p. 66, Le Nat. 1892.
- X. vidua, note on; Blackburn, p. 39, Tr. R. Soc. S. Austr. xvi.
- X. hexacantha, Argentine Rep., FAIRMAIRE, p. 245, Ann. Ent. Belg. xxxvi, n. sp.

#### TENEBRIONIDÆ.

- [Cf. ALLUAUD (8), BLACKBURN (90, 94, 95), BLAISDELL (98), CASEY (161, 162), CHAMPION (347), DISTANT (210), DUVIVIER (230), FAIRMAIRE (261, 262, 263, 264, 265, 267), GESTRO (334), HEYDEN (402), PÉRINGUEY (672, 673), REITTER (725, 726), SENAC (836), ZOUFAL (1026).]
- Notes on habits of *Tenebrionida* of San Diego; Blaisdell, Zoe, iii, pp. 102-108.
- Abantis, n. gen. near Clitobius, p. 109, for A. ænescens, n. sp., Obook, p. 110; FAIRMAIRE, Rev. d'Ent. xi.
- Acotulus (sub Orthocerus) funicornis, Bed., = (oranieneis, Reit.);
  REITTER, p. 26, Wien. ent. Z. xi.
- Acropteron magnicolle, Venesuela, FAIRMAIRE, p. 90, Ann. Soc. Ent. Fr. lxi, n. sp.

- Adelium parallelum, Germ., referred to Seirotruna; BLACKBURN, p. 5 P. Liun. Soc. N.S.W. (2) vi.
- A. simpler, Australia, BLACKBURN, p. 44, Tr. B. Soc. S. Austr. xvi; pustulosum, p. 534, rictoriæ, p. 535, inconspicuum, alpicola, p. 5 tropicum, p. 537, lindense, angulatum, p. 538, æquale, p. 5 Australia, BLACKBURN, P. Linn. Soc. N.S.W. (2) vi : n. spp.
- Adesmia leucostictu, p. 765, nobilis, p. 766, Somaliland, GESTRO, A. Mus. Genov. (2) xii; A. (Onymachris) agilis, S. Africa, Péringu Tr. S. African Phil. Soc. vi (2), p. 120: n. spp.
- Alephus macilentus, Arizona, Casey, p. 61, Ann. N. York Ac. vi, n. sp. Amarygmus natalensis, S. W. Africa, Péringuey, Tr. S. African P. Soc. vi (2), p. 59, n. sp.
- Ambigatus, n. g., near Melanophorus, for A. rufonitens, p. 246, strictico bembidioides, p. 247, n. spp., Argentine Rep.; FAIRMAIRE, Ann. l Belg. xxxvi.
- Amiantus undosus, Transvaal, DISTANT, Naturalist Transvaal, p. 199, j fig. 1, n. sp.
- Anadus posticalis, Venezuela, FAIRMAIRE, p. 89, Ann. Soc. Ent. Fr. n. sp.
- Anepsius montanus, Colorado, Casey, p. 55, Ann. N. York Ac. n. sp.
- Araoschizus mexicanus, Tepetlapa, Champion, p. 491, pl. xxii, fi<sub>i</sub> Biol. Centr. Am. Col. iv (1), n. sp.
- Argoporis rufipes, = (nitida, Casey); CHAMPION, p. 518, Biol. Centr. Col. iv (1).
- A. unicalcurata, tridentata, p. 519, laricollis, p. 520, Mexico, Chamit. c., p. 8pp.
- Arthrodeis byrrhiformis, Obock, FAIRMAIRE, p. 106, Rev. d'Ent. xi plicatus, Somaliland, GESTRO, p. 764, Ann. Mus. Genov. (2) n. spp.
- Aryenis minor, Argentine Rep., FAIRMAIRE, p. 251, Ann. Ent. xxxvi, n. sp.
- Asida: note on the Central American allies or subgenera, p. A. sordida, = (rugosissima, interrupta), clathrata, = (fun p. 492, fallax, = (fuvosa, similata), p. 496, A. collaris, n. n marginicollis, Champ., nec Rosenh., p. 499; CHAMPION, Biol. (Am. Col. iv (1).
- Asida geminata, p. 492, pl. xxii, fig. 8, intricata, p. 493, fig. 9, spinin fig. 11, p. 494, tenebrosa, fig. 12, fasciculata, fig. 21, p. 495, fig. 10, p. 496, moricoides, fig. 14, segregata, p. 497, fæda, fi tarda, fig. 17, p. 498, furcata, fig. 15, p. 499, latissima, fig. 18, ho fig. 13, p. 500, unicostata, fig. 19, p. 501, ingens, fig. 20, p. 503, M CHAMPION, t. c., n. spp.
- Astrotus undatus, pl. xxii, fig. 25, erosus, p. 504, nosodermoides, p. fig. 22, Mexico, Champion, Biol. Centr. Am. Col. iv (1), n. spp.
- Azarelius, n. g., near Gonocnemis, for A. sculpticollis, n. sp., Sun FAIRMAIRE, p. vii, Bull. Soc. Ent. Fr. 1xi.

- Blapstinus coronadensis, California, BLAISDELL, p. 242, Ent. News, iii; B. opatrinoides, pseudoæneus, p. 81, piliferus, infimus, p. 82, Venezuela, FAIRMAIRE, Ann. Soc. Ent. Fr. lxi: n. spp.
- Blepegenes nitidus, Queensland, BLACKBURN, p. 533, P. Linn. Soc. N.S.W. (2) vi, n. sp.
- Branchus opatroides, Mexico, CHAMPION, p. 507, pl. xxii, fig. 24, Biol. Centr. Am. Col. iv (1), n. sp.
- Calymnus longedentatus, Venezuela, FAIRMAIRE, p. 83, Ann. Soc. Ent. Fr. lxi, n. sp.
- Capnisa cribratella, Turkestan, FAIRMAIRE, p. cliii, Bull. Soc. Ent. Fr. lxi, n. sp.
- Carchares, Pasc., = (Podoces, Per.); Peringuey, p. 135, Tr. S. African Phil. Soc. vi, 2.
- Cardigenius densegranatus, Argentine Rep., FAIRMAIRE, p. 246, Ann. Ent. Belg. xxxvi, n. sp.
- Cardiothorax æripennis, N. S. Wales, BLACKBURN, p. 226, Tr. R. Soc. S. Austr. xv, n. sp.
- Catamerus fairmairei, Tabora, Alluaud, p. exxxix, Bull. Soc. Ent. Fr. lxi; C. transvaalensis, S. Africa, Péringuey, p. 125, Tr. S. African Phil. Soc. vi, 2: n. spp.
- Cataphronetis soror, Obock, FAIRMAIRE, p. 110, Rev. d'Ent. xi, n. sp.
- Centrioptera spiculosa, Mexico, CHAMPION, p. 508, Biol. Centr. Am. Col. iv (1), n. sp.
- Centronopus bimaculatus, Mexico, Champion, p. 521, pl. xxiii, fig. 1, Biol. Centr. Am. Col. iv (1), n. sp.
- Chalcopterus eremita, p. 44, meyricki, p. 45, W. Australia, BLACKBURN, Tr. R. Soc. S. Austr. xvi, n. spp.
- Chartopteryx victoriensis, Australia, BLACKBURN, p. 226, Tr. R. Soc. S. Austr. xv, n. sp.
- Choaspes simonis, Venezuela, FAIRMAIRE, p. 80, Ann. Soc. Ent. Fr. lxi, n. sp.
- Cibdelis lævigata, California, Casey, p. 60, Ann. N. York Ac. vi, n. sp.
- Clitobius laripennis, Obock, strongyloides, Zambesi, Fairmaire, p. 111, Rev. d'Ent. xi, n. spp.
- Colpotus magnicollis, Syria, FAIRMAIRE, p. 148, Ann. Ent. Belg. xxxvi, n. sp.
- Conibiosoma laciniata, Arizona, Casey, p. 64, Ann. N. York Ac. vi, n. sp. Cryptoglossa granulifera, Mexico, Champion, p. 508, Biol. Centr. Am. Col. iv (1), n. sp.
- Cyclosattus, n. g., for Eusattus websteri (n. sp. infrå); Casey, p. 710, Ann. N. York Ac. vi.
- Cyrtosoma oblongulum, impressipenne, Venezuela, FAIRMAIRE, p. 89, Ann. Soc. Ent. Fr. lxi, n. spp.
- Dædrosis victoriæ, Australia, Blackburn, p. 533, P. Linn. Soc. N.S.W.

  (2) vi; D. monticola, N. S. Wales, Blackburn, p. 227, Tr. R. Soc. S.

  Austr. xv: n. spp.

- Diastolinus brecicollis, p. 80, impressicollis, p. 81, Venezuela, FAIRMAI Ann. Soc. Ent. Fr. lxi, n. spp.
- Dichtrethus angustipennis, S. W. Africa, Péringuey, Tr. S. African P Soc. vi (2), p. 52, n. sp.
- Elwodes eschscholtzi, = (wickhami); CHAMPION, p. 511, Biol. Centr. I Col. iv (1).
- Eleodes interrupta, California, Blaisdell, Ent. News, iii, p. 241; rileyi, p. 57, nitidus, Arisona, compositus, Texas, p. 58, Casey, A. N. York Ac. vi; E. macrura, p. 511, polita, segregata, p. 513, ol p. 516, peropaca, tesselluta, p. 517, Mexico, Champion, Biol. Ce Am. Col. iv (1): n. spp.
- Emmenastus belti, = (rotundicollis and intermedius), p. 480, alatus, (salvini), p. 482; Champion, Biol. Centr. Am. Col. iv (1).
- E. mexicanus, subapterus, p. 481, seriatus, constrictus, p. 482, stoli p. 483, igualensis, p. 484, Mexico, Champion, Biol. Centr. Am. iv (1), n. spp.
- Emyon scabrosus, S. W. Africa, Péringuey, p. 58, Tr. S. African I Soc. vi (2), n. sp.
- Epairops levigata, variegata, S. W. Africa, Péringuey, Tr. S. Afr Phil. Soc. vi (2), p. 55, n. spp.
- Epitragodes julichi, Texas, Casey, p. 55, Ann. N. York Ac. vi, n. sp. Epitragus: tabulation of the N. American species; Casey, p. 53, A. N. York. Ac. vi. E. canus, figured and noticed, p. 487, pl. 2 fig. 5, E. rorulentus, = (scutatus), p. 486; Champion, Biol. Ce Am. Col. iv (1).
- E. cristatus, p. 485, Mexico, pl. xxii, fig. 4, ruatanensis, Honduras, p. Champion, Biol. Centr. Am. Col. iv (1); E. restitus, Arizona, Ca p. 53, Ann. N. York Ac. vi; E. parumpunctatus, p. 78, angustifor p. 79, Venezuela, Fairmaire, Ann. Soc. Ent. Fr. lxi; E. ær rufocustaneus, p. 247, nitidicollis, crassus, p. 248, nanus, Argen Rep., minutissimus, p. 249, Brazil, Fairmaire, Ann. Ent. Belg. xxi n. spp.
- Eulabis laticornis, California, Casey, p. 59, Ann. N. York Ac. vi, n. & Eusattus websteri, Colorado, Casey, p. 56, Ann. N. York Ac. vi venosus, p. 509, sculptus, mexicanus, obliteratus, p. 510, Mez Champion, Biol. Centr. Am. Col. iv (1): n. spp.
- Glyptotus yucatanus, Mexico, CHAMPION, p. 524, Biol. Centr. Am. Co (1), n. sp.
- Haporema, n. g., ? near Misolampidius, for H. decipiens, n. sp., Obc FAIRMAIRE, p. 109, Rev. d'Ent. xi.
- Hedyphanes koltzei, Turkestan, HEYDEN, p. 108, Deutsche e. Z. 1 n. sp.
- Helaus aridus, p. 41, debilis, lubricus, p. 42, ingens, p. 43, Centr. Austr. Blackburn, Tr. R. Soc. S. Austr. xvi, n. spp.
- Helopinus costatus, = (elegans, Baudi, and misolampoides, Lac.); F. MAIRE, p. 112, Rev. d'Ent. xi.
- Helops cerberus, metamorphoses; XAMBEU, Mœurs, ii, p. 11.

- H. blaisdelli, p. 66, discipula, p. 67, California, cylindriformis, New Mexico, p. 68, CASEY, Ann. N. York Ac. vi, n. spp.
- Himatismus pavesii, Somaliland, GESTRO, p. 769, Ann. Mus. Genov. (2) xii, n. sp.
- Histiaa, n. g., near Anemiu, p. 107, for H. bidentula, n. sp., Obock, p. 108; FAIRMAIRE, Rev. d'Ent. xi.
- Hopatrum elderi, p. 39, longicorne, p. 40, Australia, BLACKBURN, Tr. R. Soc. S. Austr. xvi, n. spp.
- Hoplonyx distinctus, Congo, DUVIVIER, p. 163, Ann. Ent. Belg. xxxvi; H. ingratus, S. W. Africa, Péringuey, p. 59, Tr. S. African Phil. Soc. vi (2); H. vicinus, Transvaal, Péringuey, t. c.: n. spp.
- Hyperops plicicollis, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2), p. 50, n. sp.
- Idiobates, n. g. for Tenebrio castaneus, Knoch; CASEY, p. 62, Ann. N. York Ac. vi.
- Idricus pacificus, S. Africa, Péringuey, p. 122, Tr. S. African Phil. Soc. vi (2), n. sp.
- Isicerdes vicinus, Central America, Champion, p. 524, Biol. Centr. Am. Col. iv (1), n. sp.
- Isostira raucipennis, Queensland, BLACKBURN, p. 225, Tr. R. Soc. S. Austr. xv, n. sp.
- Lasiostola costuligera, Margelan, HEYDEN, p. 107, Deutsche e. Z. 1892, n. sp.
- Leptodes, characters and synonymy; Reitter, pp. 93-96, Wien. ent. Z. xi.
- L. semenowi, Araxes Valley, REITTER, p. 66, Wien. ent. Z. xi; L. heydeni, Turkestan, REITTER, t. c. p. 95: n. spp.
- Licinoma pallipes, Australia, BLACKBURN, p. 542, P. Linn. Soc. N.S.W. (2) vi, n. sp.
- Lyphia striolata, Obock, FAIRMAIRE, p. 111, Rev. d'Ent. xi, n. sp.
- Mesabates spissicornis, p. 479, inæqualis, p. 480, Mexico, Champion, Biol. Centr. Am. Col. iv (1), n. spp.
- Mesostenopa auberti, Obock, FAIRMAIRE, p. 107, Rev. d'Ent. xi, n. sp.
- Micrantereus lavior, Obock, FAIRMAIRE, p. 114, Rev. d'Ent. xi, n. sp.
- Mitys politus, = (lavis); Champion, p. 520, Biol. Centr. Am. Col. iv (1).
- Nosoderma inaquale, Say, = (morbillosum, Sol.); Champion, p. 490, Biol. Centr. Am. Col. iv (1).
- Nuptis laticollis, Panama, Champion, p. 523, Biol. Centr. Am. Col. iv (1), n. sp.
- Odontopezus asper, Usagara, Alluaud, p. ccxxxviii, Bull. Soc. Ent. Fr. lxi, n. sp.
- Œatus similis, Central America, Champion, p. 523, Biol. Centr. Am. Col. iv (1), n. sp.
- Ologlyptus hebes, Mexico, Champion, p. 506, pl. xxii, fig. 23, Biol. Centr. Am. Col. iv (1), n. sp.
- Opatroides curtulus, Syria, FAIRMAIRE, p. 149, Ann. Ent. Belg. xxxvi, n. sp.

- Oxycara breriuscula, Obock, FAIRMAIRE, p. 107, Rev. d'Ent. xi, n. sp. Palembus, n. g. near Oplocephala for P. ocularis, n. sp., Florida; Cas p. 65, Ann. N. York Ac. vi.
- Paranemia, n. g., Trachyscelidarum, for P. schroderi, n. sp., Turkest: HEYDEN, p. 103, Deutsche e. Z. 1892.
- Phymatodes denticollis, Caracas, p. 86, claricornis, Buenos Aires, p. FAIRMAIRE, Ann. Soc. Ent. Fr. lxi, n. spp.
- Pimelia, characters and synonymy of the species of the Canaries; SEN pp. 103-108, Ann. Soc. Ent. Fr lxi. P. thomsoni, redescrib SENAC, p. lxxiii, Bull. Soc. Ent. Fr. lxi.
- P. alluaudi, Canary Is., SENAC, p. 107, Ann. Soc. Ent. Fr. lxi, n. sp.
- Pimeliopsis, n. g. Epiphysides, for P. granulata, n. sp., Mexico, p. 4 pl. xxii, fig. 1; Champion, Biol. Centr. Am. Col. iv (1).
- Platydema contractum, Venezuela, FAIRMAIRE, p. 84, Ann. Soc. Ent. lxi; P. basicorne, Argentine Rep., FAIRMAIRE, p. 250, Ann. I Belg. xxxvi: n. spp.
- Platyphanes creber, Queensland, BLACKBURN, p. 225, Tr. R. Soc Austr. xv, n. sp.
- Pacilesthus ragelineatus, Venezuela, FAIRMAIRE, p. 91, Ann. Soc. Ent. lxi, n. sp.
- Poyonobasis hirsuta, S. W. Africa, Péringuey, p. 51, Tr. S. African I Soc. vi (2), n. sp.
- Pruogena tristis, S. W. Africa, PÉRINGUEY, Tr. S. African Phil. Soc (2), n. sp.
- Psammodes puchysoma, p. 52, errans, p. 53, S. W. Africa, Péringuey, S. African Phil. Soc. vi (2), n. spp.
- Pyres (sub Centronopus) speciosus, Pasc., = (metallicus, Chan Champion, p. 521, Biol. Centr. Am. Col. iv (1).
- Rhicnodus, n. g. near Chatyllus, for R. asper, p. 87, simonis, parcegranstriatopunctatus, p. 88, n. spp., Venezuela; Fairmaire, Ann. Ent. Fr. lxi.
- Rhytidonota absurda, Somaliland, Gestro, p. 768, Ann. Mus. Genov xii, n. sp.
- Rues, n. g. for Helops ovipennis, Casey; Casey, p. 66, Ann. N. J. Ac. vi.
- Saragus tricarinatus, Australia, BLACKBURN, p. 43, Tr. R. Soc. S. Axvi, n. sp.
- Scleron amplicolle, Obock, FAIRMAIRE, p. 108, Rev. d'Ent. xi, n. sp.
- Seirotrana simplex, p. 540, monticola, p. 541, dispar, p. 542, Austr Blackburn, P. Linn. Soc. N.S.W. (2) vi; S. major, N. S. W Blackburn, p. 228, Tr. R. Soc. S. Austr. xv: n. spp.
- Sepidiopsis, n. g., near Sepidium, p. 771, for S. cornigera, p. 772, vil p. 773, n. spp., Somaliland; Gestro, Ann. Mus. Genov. (2) xii.
- Sepidium ovampoense, S. W. Africa, PÉRINGUEY, p. 55, Tr. S. Afr Phil. Soc. vi (2); S. zambezianum, S. Africa, PÉRINGUEY, p. t. c.: n. spp.
- Sepidostenus, = (Sepidiacis); GESTRO, p. 775, Ann. Mus. Genov. (2) x

- Solenomerus relox, S. W. Africa, PÉRINGUEY, p. 58, Tr. S. African Phil. Soc. vi (2), n. sp.
- Stenocara albovillosa, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2), p. 49; S. intermedia, S. Africa, Péringuey, p. 121, t. c.: n. spp.
- Stibia maritimo, California, Casey, p. 52, Ann. N. York Ac. vi, n. sp. Strongylium variolosum, Congo, Duvivier, p. 164, Ann. Ent. Belg. xxxvi; S. cyaneum, S. W. Africa, Péringuey, Tr. S. African Phil.
  - Soc. vi (2); S. picicorne, Venezuela, FAIRMAIRE, p. 92, Ann. Soc. Ent. Fr. lxi; S. platense, Argentine Rep., FAIRMAIRE, p. 251, Ann. Ent. Belg. xxxvi: n. spp.
- Tactoderus subopacus, Obock, FAIRMAIRE, p. 112, Rev. d'Ent. xi, n. sp.
- Tænophthalmus costifer, Turkestan, HEYDEN, p. 108, Deutsche e. Z. 1892, n. sp.
- Talanus parallelus, p. 92, crenulatus, p. 93, Venezuela, FAIRMAIRE, Ann. Soc. Ent. Fr. lxi, n. spp.
- Tenebrio angustus, N. Persia, syriacus, Syria, Zoufal, p. 284, Wien. ent. Z. xi, n. spp.
- Tenesis, n. g., near Amenophis, for T. femoratus, n. sp., Congo, DUVIVIER, p. 163, Ann. Ent. Belg. xxxvi.
- Trachynotus batesi, = (scrobiculatus, Per.), gracilipes, = (attenuatus, Per.);
  Pringuey, p. 135, Tr. S. African Phil. Soc. vi (2).
- T. distinctus, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2), p. 54; T. lightfooti, S. Africa, Péringuey, p. 122, t. c.: n. spp.
- Trigonopus laticollis, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2), p. 57; T. armaticeps, Damaraland, Péringuey, p. 123, t. c.: n. spp.
- Trimytis obovata, pl. xxii, fig. 2, flohri, Mexico, Champion, p. 478, Biol. Centr. Am. Col. iv (1), n. spp.
- Ulosonia ceratodera, Venezuela, FAIRMAIRE, p. 85, Ann. Soc. Ent. Fr. lxi; U. parvicornis, Argentine Rep., FAIRMAIRE, p. 250, Ann. Ent. Belg. xxxvi: n. spp.
- Ulus latus, California, BLAISDELL, p. 243, Ent. News, iii, n. sp.
- Usagaria australis, S. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2), p. 56, n. sp.
- Zialas, n. g., near Gonocnemis, for Z. insolitus, n. sp., Hué; FAIRMAIRE p. ex, Bull. Soc. Ent. Fr. lxi.
- Zopherus venosus is distinct from lævicollis, jourdani, Sall., = (mexicanus, Sol., nec Gray), p. 488, elegans, pl. xxii, fig. 7, nodulosus, fig. 6, figured; Champion, Biol. Centr. Am. Col. iv (1).
- Zophobas morio, variation noticed; Champion, p. 522, Biol. Centr. Am. Col. iv (1).
- Zophosis acuticostu, n. var. pinardi; Thert, p. clxxxi, Bull. Soc. Ent. Fr. lxi.
- Z. ogadenica, Somaliland, Gestro, p. 762, Ann. Mus. Genov. (2) xii; Z. amplicollis, Obock, FAIRMAIRE, p. 106, Rev. d'Ent. xi; Z. rugatipennis, dorsata, p. 47, hypocrita, nana, p. 48, S. W. Africa, PÉRINGUEY, Tr. S. African Phil. Soc. vi (2): n. spp.

### CISTELIDA.

- [Cf. Bedel (48), Casey (161), Fairmaire (261, 262, 263), Reiti (725).]
- Revision of the N. American genera and species; Casey, pp. 69–169, A. N. York Ac. vi.
- Allecula seriatopora, Argentine Rep., FAIRMAIRE, p. 251, Ann. Ent. Be xxxvi, n. sp.
- Amirimus, n. g., p. 155, for a part of Cteniopus, and for A. brunneus, p. 1 concolor, p. 158, nigrescens, convergens, p. 159, n. app., N. Americ Casey, Ann. N. York Ac. vi.
- Apellutus palpalis, ? = (mastersi, Maol.); BLACKBURN, p. 115, P. Lin Soc. N.S.W. (2) vii.
- Cistela forticornis, flavida, Syria, Fairmaire, p. 150, Ann. F Belg. xxxvi; C. ocularis, Obock, Fairmaire, p. 115, Rev. d'I xi: n. spp.
- Cteniopus impressicollis, C.? crassus, Syria, FAIRMAIRE, p. 152, Ann. I Belg. xxxvi, n. spp.
- Dietopsis acuminata, Venezuela, FAIRMAIRE, p. 94, Ann. Soc. Ent. lxi, n. sp.
- Eryx asiatica, Syria, FAIRMAIRE, p. 151, Ann. Ent. Belg. xx n. sp.
- Gonodera delagrangei, p. 150, atronitens, p. 151, Syria, FAIRMAIRE, & Ent. Belg. xxxvi, n. spp.
- Hymenorus infuscatus, picipennis, p. 90, melsheimeri, p. 92, obesus, p. difficiles, p. 94, perforatus, curticollis, p. 95, discrepans, p. 98, apa anus, p. 99, exiguus, p. 100, helvinus, porosicornis, p. 101, interme p. 102, prolixus, p. 103, discretus, p. 105, convexus, p. 106, disser seriatus, p. 109, testaceus, p. 110, rotundicollis, p. 111, inquili fuscicornis, p. 112, nitidipennis, p. 113, inæqualis, p. 114, tene sobrinus, p. 115, floridanus, p. 116, fusculus, p. 117, macer, p. indutus, p. 119, gemellus, p. 121, uniseriatus, p. 122, N. America, Ca Ann. N. York Ac. vi, n. spp.
- Isomira iowensis, discolor, p. 145, tenebrosa, p. 146, luscitiosa, p. 148, :
  ticola, p. 150, oblongula, p. 151, texana, p. 153, N. America, C.
  Ann. N. York Ac. vi, n. spp.
- Lobopoda subcuneata, p. 79, oculatifrons, p. 81, Texas, Casey, Anr York Ac. vi; L. pilosula, Venesuela, Fairmaire, p. 93, Ann. Ent. Fr. lxi: n. spp.
- Mycetochara megalops, p. 129, gilvipes, p. 131, nigerrima, p. 132, l pennis, pacifica, p. 139, procera, p. 140, nevadensis, crassulipes, p. N. America, Casey, Ann. N. York Ac. vi, n. spp.
- ()mophlus lepturoides, metamorphoses; XAMBEU, Mœurs, ii, p. 33.
- O. (Heliotaurus) brisouti, Algeria, p. 794, doriæ, Tunis, p. 795, Bi Ann. Mus. Genov. (2) x; O. densepunctatus, cribricollis, foveic Syria, FAIRMAIRE, p. 153, Ann. Ent. Belg. xxxvi; O. will Margelan, Reitter, p. 136, Wien. ent. Z. xi: n. spp.

- Tedinus, n. g., p. 153, for T. angustus, Georgia, n. sp., p. 154, Casey, Ann. N. York Ac. vi.
- Xystropus californicus, = (Prostenus fulgidus, Mäkl.); Casey, p. 74, Ann. N. York Ac. vi.
- X. fuscicularis, p. 94, alborittatus, gossypiatus, p. 95, Venezuela, FAIR-MAIRE, Ann. Soc. Ent. Fr. lxi, n. spp.

## MELANDRYIDÆ, LAGRIIDÆ.

- [Cf. FAIRMAIRE (261), GUILLEBEAU (Échange, 1887-88), PÉRINGUEY (672, 673), XAMBEU (1023).]
- Abdera triguttata, metamorphoses; XAMBEU, Mœurs, ii, p. 19.
- Eutrapela concolor, n. n. for unicolor, Pér.; Péringuey, p. 134, Tr. S. African Phil. Soc. vi (2).
- Hysterarthron collare, Thoms., referred to Lagriidæ; RITSEMA, p. 54, Notes Leyd. Mus. xiv.
- Lagria corrugata, Zambesi, PÉRINGUEY, Tr. S. African Phil. Soc. vi (2), p. 61, n. sp.
- Orchesia australis, reyi, p. 3, Échange, 1887, No. 31, subimpressa, France, No. 32, p. 4, corsica, Corsica, p. 13, abeillei, Marseilles, p. 14, No. 40, 1888, GUILLEBRAU, n. spp.
- Statira simonis, p. 95, variicollis, p. 96, Venezuela, FAIRMAIRE, Ann. Soc. Ent. Fr. lxi, n. spp.

# PEDILIDE, XYLOPHILIDE.

[Cf. FAIRMAIRE (261), HARRINGTON (389), Pic (678).]

Bactrocerus cribripennis, Venezuela, FAIRMAIRE, p. 97, Ann. Soc. Ent. Fr. lxi, n. sp.

Ischalia vancouverensis, N. America, HARBINGTON, p. 132, Canad. Ent. xxiv, n. sp.

Macratria comosella, Venezuela, FAIRMAIRE, p. 97, Ann. Soc. Ent. Fr. lxi, n. sp.

Scraptia trotommoides, Syria, Pic, Échange, 1892, p. 4, n. sp.

Xylophilus brericornis, Perris, habitat and characters; Champion, Ent. M. M. xxviii, p. 68.

X.-Euglenes (Anidorus, M.) testaceipes, Algeria, Ptc, p. 139, Échange, 1892, n. sp.?

#### ANTHICIDÆ.

- [Cf. FAIRMAIRE (261, 262), HORN (418), PIC (678, 680, 682, 684, 685, 687, 689, 690), REITTER (725).]
- Anthicus: synonymical notes, with A. brevipilis, n. n. for pilosus, Pic, marseuli, n. n. for scoticus, Mars., and reyi, n. n. for puberulus, Rey; Pic, pp. cexi & cexii, Bull. Soc. Ent. Fr. lxi.

- 1. delagrangei, Syria, pilosus, deslogesi, Algeria, Pic, p. 43, Échange, 1. henoni, Algeria, Pic, p. 103, t. c.; A. rubricollis, Marocco, t. c. p. 139 (? n. sp.); A. roberti, p. 213, testaceipes, p. 214, Alg Pic, Rev. Sci. Bourb. v; A. chobauti, Algeria, Pic, p. lxxv, Bull Ent. Fr. lxi; A. bedeli, carthaginiensis, Algeria, p. cxcvii, bl Spain, p. excviii, Pic, t. c.; A. demaisoni, lepaumi, Egypt, p. ccxxi, t. c.; A. kraatzi, Syria, p. cclxviii, nitidior, Turke p. cclxix, siciliæ, Sicily, angulifer, Syria, p. cclxx, croissane Corsica, p. cclxxi, Pic, t. c.: n. spp.
- Formicomus pegusus, Transcaspian Region, Reitter, p. 63, Wien. en xi; F. strigicollis, spinicrus, Obock, Fairmaire, p. 116, Rev. d xi: n. spp.
- Leptaleus truncatulus, Obock, FAIRMAIRE, p. 116, Rev. d'Ent. xi, n. s. Notoxus jucqueti, Wallachia, Pic, p. 4, Échange, 1892; N. schwarzi, 1 HORN, p. 47, Tr. Am. Ent. Soc. xix: n. spp.
- Tomo lerus bosnicus, Bosnia, Pic, p. 241, Rev. Sci. Bourb. v, n. sp.

## MORDELLIDÆ, RHIPIDOPHORIDÆ, STYLOPIDÆ.

- [Cf. Blackburn (95), Champion (163), Chobaut (169), Fairy (261, 264), Horn (418).]
- Conalia baudii, habits noticed; Fleischer, p. 209, Wien. ent. Z. xi. Mordella exilis, varr.? described, Blackburn, p. 46, Tr. R. Soc. S. A xvi.
- Mordellistena floridensis, larva and larval habits; RILEY, P. E. Soc. V ii, p. 321, figs. M. nigricans, habits; BRODIE, p. 139, Canad. xxiv.
- M. tenuis, Venezuela, FAIRMAIRE, p. 98, Ann. Soc. Ent. Fr. lxi, n. sp. Pentaria oberthuri, Pyrenees or., Champion, p. 109, Ent. M. M. x n. sp.
- Myodites: note on N. American; HORN, p. 48, Tr. Am. Ent. Soc. xi Rhipidius abeillei, Mt. Ventoux, Споваст, Mém. Ac. Vaucluse, xi, p. pl., n. sp.
- Rhipiphorus basicollis, Caracas, FAIRMAIRE, p. 98, Ann. Soc. Ent. Fr. R. argentinus, Argentine Rep., FAIRMAIRE, p. 252, Ann. Ent. xxxvi: n. spp.
- Stylopida: localities and dates of British; DALE, Ent. M. M. xxviii, Elenchus tenuicornis, parasitic on Homopteron (Liburnia), SAUNDERS, M. M. xxxviii, pp. 249 & 297, fig.; details of structure of the 1 EATON, t. c. p. 250, woodcut.
- Xenos sp.: life-history; Hubbard, pp. 257-261, Canad. Ent. xxiv.

#### Cantharidæ.

[G. Bedel (51), Blackburn (94, 95), Casey (161), Champion ( Escherich (258), Fairmaire (263, 264, 271), Frivaldszky ( Péringuey (672, 673), Prochazka (705), Reitter (725), Wi (989).]

- List of what is known as to habits and early stages of European Meloidæ; Bedel (51).
- Calospasta mirabilis, figured, Biol. Centr. Am. Col. iv (2) pl. xviii, fig. 12. C. sulcifrons, Mexico, id. t. c. p. 394, pl. xviii, fig. 14, n. sp.
- Cantharis: CHAMPION, Biol. Centr. Am. Col. iv (2), notices and figures the following—C. cardinalis, pl. xx, fig. 12, p. 438, erythrothorax, fig. 13, p. 439, quadrimaculuta, fig. 14, p. 440, corallifera, fig. 17, p. 441, sanguinea, fig. 19, p. 442, bipuncticollis, p. 443, fig. 21, variabilis, p. 447, proteus, figs. 22 & 23, p. 448.
- C. (Lytta) optata, S. W. Africa, Péringuey, p. 65, Tr. S. African Phil. Soc. vi (2); C. intricata, p. 441, pl. xx, fig. 18, erebea, p. 442, fig. 20, subviolacea, p. 445, scitula, fig. 25, signaticallis, fig. 24, p. 446, michoacanæ, p. 447, Mexico, Champion, Biol. Centr. Am. Col. iv (2): n. spp.
- Cephaloon: notes on the variation of the species; Heyden, pp. 165-169, Deutsche c. Z. 1892.
- Cysteodemus wizlizeni, figured; Biol. Centr. Am. Col. iv (2), pl. xvii, fig. 10.
- Epicauta: the following species are noticed, with much synonymy and many figures, in Biol. Centr. Am. Col. iv (2), by Champion: E. curricornis, pl. xviii, fig. 25, basimacula, pl. xix, fig. 1, p. 406, rufipedes, fig. 2, p. 407, rufipennis, fig. 4, p. 408, terminata, fig. 5, p. 409, funesta, fig. 6, punctum, fig. 7, p. 410, maculata, p. 413, ocellata, fig. 9, pardalis, fig. 10, ritticollis, fig. 11, p. 414, grammica, fig. 12, p. 415, niveolineata, figs. 14 & 15, albolineata, fig. 16, p. 416, carmelita, fig. 17, p. 417, cinctipennis, fig. 18, croceicincta, fig. 19, p. 420, cinerca, p. 421, stigmata, p. 422, neglecta, p. 423, pl. xix, fig. 21, obesa, p. 424. E. griseo-nigra, n. var. riojana; Fairmaire, p. 252, Ann. Ent. Belg. xxxvi.
- E. desgodinsi, Tibet, FRIVALDSZKY, p. 115, Term. füzetek, xv; E. levettei, Colorado, duplicata, Arizona, p. 172, Casey, Ann. N. York Ac. vi; E. atripilis, p. 410, cinereiventris, p. 411, pl. xix, fig. 8, evanescens, unicalcarata, p. 412, unilineata, fig. 13, p. 415, atricolor, p. 419, tripartita, p. 421, fig. 20, auricomans, p. 424, fig. 22, leucocoma, p. 425, fig. 23, candidata, emarginata, fig. 24, p. 426, singularis, fig. 25, compressicollis, fig. 26, p. 427, Champion, Biol. Centr. Am. Col. iv (2): n. spp.
- Eupompha fissiceps, figured, Biol. Centr. Am. Col. iv (2), pl. xviii, fig. 13. Gnathium minimum, figured, Biol. Centr. Am. Col. iv (2), pl. xvii, fig. 17. Gnathospasta: Cantharis labialis referred to this; Champion, p. 403, Biol. Centr. Am. Col. iv (2).
- Halosimus araxis, Araxes Valley, REITTER, p. 63, Wien. ent. Z. xi, n. sp. Hapalus and Stenoria, revision of the palearctic species; PROCHAZKA, pp. 263-270, Wien. ent. Z. xi.
- H. binævus, reitteri, Turkestan, PROCHAZKA, Wien. ent. Z. xi, p. 270, n. spp.
- Henous mexicanus, figured and noticed; Champion, Biol. Centr. Am. Col. iv (2) p. 369, pl. xvii, fig. 6.

- Hoplozonitis, n. g., p. 228, for H. mira, n. sp., Queensland, p. 229; Bi BURN, Tr. R. Soc. S. Austr. xv.
- Horia auriculata, N. and Centr. America, Champion, p. 372, pl. xvii 9-9d, Biol. Centr. Am. Col. iv (2) n. sp.
- Lydus pilicollis, rulneratus, p. 155, impressicollis, p. 156, Syria, FAIRM Ann. Ent. Belg. xxxvi, n. spp.
- Lytta mæsta, n. n. for lugubris, Pér., designata, n. n. for albolineata, Péringuey, p. 135, Tr. S. African Phil. Soc. vi (2).
- L. carneola, p. 125, amana, p. 126, S. Africa, Péringuey, Tr. S. Af Phil. Soc. vi (2), n. spp.
- Macrobasis fumosa, p. 395, pl. xviii, fig. 15, purpurea, p. 397, fig. diversicornis, p. 399, fig. 19, beckeri, p. 400, fig. 20, ochrea, p. fig. 23, segmentata, fig. 22, noticed and figured; Champion, Centr. Am. Col. iv (2).
- M. distorta, p. 396, pl. xviii, fig. 16, disparilis, p. 398, fig. 18, tenuic p. 400, fig. 21, megacephala, p. 401, fig. 24, Centr. America, Cham Biol. Centr. Am. Col. iv (2): n. spp.
- Megetra cancelluta and var. described and figured; Champion, p pl. xvii, fig. 8.
- Meloë: notes on young larva; Weber, Ber. Ver. Cassel, 1891, 1892, pp. 1-5, pl.
- M. meridianus, S. W. Africa, PERINGUEY, p. 62, Tr. S. African Phil vi (2), n. sp.
- Mylabris tenuepicta, posticuta, Syria, FAIRMAIRE, p. 154, Ann. Ent. xxxvi; M. hilaris, p. 63, M. (Ceroctis) blanda, aliena, p. 64, Africa, Péringuey, Tr. S. African Phil. Soc. vi (2): n. spp.
- Negalius, n. g., near Phodaga, for N. marmoratus, n. sp., Texas; C p. 175, Ann. N. York Ac. vi.
- Nemognatha caruleipennis, varr. figured and noticed, p. 374, pl. xvii 7, &c., tarasca, Dug., figured, fig. 4, redescribed, p. 378, zonit fig. 15, flara, fig. 16, figured; Champion, Biol. Centr. Col. iv (2).
- N. brevicollis, pl. xvii, fig. 11, foveifrons, fig. 12, p. 376, nigripes, e piata, fig. 13, p. 377, Mexico, Champion, Biol. Centr. Am. Col. in n. spp.
- Pyrota terminata, p. 429, pl. xx, fig. 1, punctata, fig. 2, mylabrina, p. 430, postica, fig. 5, p. 431, tenuicostalis, p. 432, fig. 6, dive fig. 7, decorata, fig. 8, p. 433, nobilis, fig. 10, quadrinervata, fig. 436, noticed and figured; Champion, Biol. Centr. Am. iv (2).
- P. rugulipennis, p. 431, pl. xx, fig. 4, hirticollis, fig. 9, mariarum, p. Mexico, Champion, Biol. Centr. Am. Col. iv (2); P. pun p. 173, concinna, p. 174, Texas, Casey, Ann. N. York Ac. p. spp.
- Tetraonyx sexguttatus, variation noticed, p. 388, figured, pl. xviii, figs batesi, fig. 7, bipartitus, fig. 8, sallai, fig. 9, maculatus, fig. 10, from fig. 11, figured; Biol. Centr. Am. Col. iv (2).

- Zonabris: regularity in variation of; ESCHERICH, Deutsche e. Z. 1892, pp. 113-130, pl. i. Z. beckeri, note on; BECKER, p. 69, Bull. Mosc. 1892.
- Z. chinensis, parvula, China, FRIVALDSZKY, p. 114, Term. füzetek, xv, n. spp.
- Zonitis analis, Ab., thoracica, Cast., abeillei, Bed., noticed, ruficollis, distinctive, and & characters, p. 174, and of allies, p. 176; ESCHERICH, Wien. ent. Z. xi. Z. posticalis, n. n. for apicalis, Pér.; Péringuey, p. 135, Tr. S. African Phil. Soc. vi (2). Z. flohri, fig. 18, nigromaculata, fig. 23, fulva, fig. 24, figured, pl. xvii; Biol. Centr. Am. Col. iv (2). Z. thoracica, habits; Vachal, p. xxiii, Bull. Soc. Ent. Fr. lxi.
- Z. gracilenta, p. 156, cæruleiceps, p. 157, Syria, FAIRMAIRE, Ann. Ent. Belg. xxxvi; Z. quadricollis, p. cliii, scutellaris, sinuatipennis, p. cliv, Turkestan, FAIRMAIRE, Bull. Soc. Ent. Fr. lxi; Z. pulchella, proxima, S. Africa, Péringuey, Tr. S. African Phil. Soc. vi (2); Z. helmsi, Australia, BLACKBURN, Tr. R. Soc. S. Austr. xvi; Z. dunniana, perforata, Texas, Casey, p. 170, Ann. N. York Ac. vi; Z. tenebrosa, p. 383, pl. xvii, fig. 19, sanguinicollis, p. 384, figs. 20 & 21, lateritia, p. 385, fig. 22, megalops, p. 386, fig. 26, Centr. America, Champion, Biol. Centr. Am. Col. iv (2): n. spp.

#### EDEMERIDÆ.

[Cf. Abeille (1), Fairmaire (263, 264), Gestro (334).]

Ananca pallida, Somaliland, Gestro, p. 777, Ann. Mus. Genov. (2) xii;
A. sellata, luridimembris, Argentine Rep., FAIRMAIRE, p. 253, Ann.
Ent. Belg. xxxvi: n. spp.

Asclera partitipennis, Syria, FAIRMAIRE, p. 158, Ann. Ent. Belg. xxxvi, n. sp.

Chrysanthia oralis, p. 158, chalcochroa, p. 159, Syris, FAIRMAIRE, Aun. Ent. Belg. xxxvi, n. sp.

Edemera basipes, Cyprus, p. 63, lutescens, Syria, p. 64, ABEILLE, Rev. d'Ent. xi; E. apicipennis, Syria, FAIRMAIRE, p. 157, Ann. Ent. Belg. xxxvi: n. spp.

Mycstochares grandicollis, longior, Syria, FAIRMAIRE, p. 149, Ann. Ent. Belg. xxxvi, n. spp.

Nacerdes melanura, distribution of; Champion, Ent. M. M. xxviii, p. 23. Stenaxis parallela, Syria, Fairmaire, p. 158, Ann. Ent. Belg. xxxvi, n. sp.

Xanthochroa foveata, Syria, FAIRMAIRE, p. 157, Ann. Ent. Belg. xxxvi n. sp.

#### CURCULIONIDÆ.

[Cf. Aurivillius (25), Becker (44), Blackburn (91, 93, 94, 95), Casev (161, 162), Desbrochers (200, 201, 202, 205-209), Duvivier (230, 231), Fairmaire (262, 270), Faust (272-278, 402), Frivaldszky (309, 310), Gestro (334), Grilat (364), Guillebeau (375), 1892. [vol. xxix.]

- HELLER (394), PERINGUEY (672, 673), REITTER (725), REY ROELOFS (772-776), ROST (780), SCHWARZ (805), SCUDDER STIERLIN (898, 899, 900), TRYON (953), WEBSTER (991).]
- Summary of and remarks on N. American Tertiary Rhynche Scudder (814).
- For many synonymical remarks on Palssarctic Curculionic Desprochers, Frelon, i, pp. 20-32.
- Abrachius, n. g., near Rhynchophorus, p. 210, for A. insularis, Aru Is., p. 211; ROELOFS, Notes Leyd. Mus. xiv.
- Acamptus echinus, New York, Casey, p. 445, Ann. N. York n. sp.
- Acalles edoughensis, subglobatus, p. 97, 7-costatus, p. 98, longus Algeria, DESEROCHERS, Frelon, ii; A. profusus, Texas, Casey Ann. N. York Ac. vi: n. spp.
- Acallestes: characters; FAUST, p. 39, S. E. Z. liii.
- Agnesiotis blanda, Australia, FAUST, p. 180, S. E. Z. liii, n. sp.
- Alaccyba: systematic position is in Erirhinini, Casey, p. 709, 1 York Ac. vi.
- Alcides angulus, hypocritus, &c., synonymical note; AURIVILLIUS, N. Arch. Mus. Paris (3) iii.
- A. eruditus, p. 312, josephus, p. 314, Congo, Duvivier, Ann. Er xxxvi; A. affaber, p. 218, clathratus, p. 219, Laos, decem Cambodia, p. 220, Aurivillius, N. Arch. Mus. Paris (3) richteri, pervicax, p. 207, reductus, N. Guinea, sellatus, p. 208, p. 209, Sumatra, FAUST, S. E. Z. liii: n. spp.
- Allomimus politus, Texas, Casey, p. 692, Ann. N. York n. sp.
- Alophus, = (Seidlitzia, Desb.); FAUST, p. 51, S. E. Z. liii.
- Alycodes dubius, Dietz, referred to Dorytomus; CASEY, p. 363, York Ac. vi.
- Ambates nobilis, p. 40, bipartitus, buqueti, p. 41, Venezuela, S. E. Z. liii, n. spp.
- Ampeloglypter longipennis, N. America, Casey, p. 549, Ann. N. Y vi, n. sp.
- Anchoning, new group of Molytides, defined, p. 19, genera ta pp. 19 & 20, species revised, pp. 21-60; FAUST, Deutsel 1892.
- Anchonus panamensis, Panama, pudens, Guadeloupe, p. 43, l Venezuela, p. 44, fraterculus, p. 45, murcidus, Costa Rica, lentus, Mexico, p. 46, intricatus, Panama, brevis, Venezuela mirus, Guatemala, p. 48, kirschi, Colombia, pedestris, Mexic tardus, Fonteboa, p. 50, morulus, femoralis, Venezuela, p. 51, Honduras, junix, p. 52, affaber, p. 53, Colombia, confidens, Ve p. 54, Faust, Deutsche e. Z. 1892: n. spp.
- Anephilus, n. g., Anchoninæ, p. 34, for A. simoni, p. 56, longulu zuela, guadulpianus, Guadeloupe, p. 57, claviger, Venezuel n. spp., FAUST, Deutsche e. Z. 1892.

- Anthonomus: characters of the European species; DESBROCHERS, Frelon, ii, pp. 106, &c. A. ulmi, pedicularius, synonymy; DESBROCHERS, Échange, 1891, No. 76, p. 31. A. musculus and signatus, characters noticed; HAMILTON, Canad. Ent. xxiv, p. 41.
- A. curtulus, Mediterranean Region, DESBROCHERS, p. 112, Frelon, ii, n. sp.
- Anypotactus: systematic position is in Sitoninæ; FAUST, p. 21, S. E. Z. liii.
- A. sulcicollis, Venezuela, FAUST, p. 21, S. E. Z. liii, n. sp.
- Apion semirufum, Lyons, Rey, Échange, 1888, No. 45, p. 4; A. fallaciosum, Algeria, turkestanicum, p. 107, reitteri, soricinum, p. 108,
  Turkestan, Desbrochers, Frelon, i; A. terræ-reginæ, Queensland,
  Blackburn, p. 151, P. Linn. Soc. N.S.W. (2) vii: n. spp.
- Apirocalus cornutus, n. var. ebrius; FAUST, p. 189, S. E. Z. liii.
- Apoderus cardinalis, n. n. for miniatus, Pér.; Péringuey, p. 135, Tr. S. African Phil. Soc. vi (2).
- A. balteatus, Transvaal, Péringuey, p. 129, Tr. S. African Phil. Soc. vi (2), n. sp
- Apotrepus, n. g., Cossonini, near Caulophilus, for A. densicollis, n. sp., Arizona; Casey, p. 695, Ann. N. York Ac. vi.
- Arachnopus fossulatus, stigmatus, p. 223, fenestratus, p. 224, New Guinea, FAUST, S. E. Z. liii, n. spp.
- Aterpus raucus, abruptus, Australia, BLACKBURN, p. 132, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Attactagenus cordicollis, Portugal, p. 75, Desbrochers, Frelon, i, n. sp.
- Attelabus (Lamprolabus) bihastatus, Yunnan, FRIVALDSZKY, p. 117, Term. füzetek, xv, n. sp.
- Auletobius glaber, tibialis, Venezuela, FAUST, p. 43, S. E. Z. liii, n. spp.
- Aulobaris dux, Nebraska, Casey, p. 546, Ann. N. York Ac. vi.
- Bagous frit, longitarsis, &c., synonymy and characters; FOWLER, Ent. M. M. xviii, p. 82.
- Balaninus: characters of the European species; DESBROCHERS, Frelon, ii, pp. 102-105.
- B. nivosus, p. 205, pluto, p. 206, New Guinea, FAUST, S. E. Z. liii, n. spp. Barini: the N. American genera tabulated and their classification dis-
- cussed; Casey, pp. 460-469, Ann. N. York Ac. vi.
- Baridius: revision of the European; Deserochers, Frelon, ii, pp. 19-69.
  B. phlomidis, Sarepta, Becker, p. 70, Bull. Mosc. 1892 (? n. sp.); B. abeillei, Syria, p. 44, litigiosus, Sarepta, p. 45, insularis, Corsica, p. 48, triplagiatus, p. 49, stricticollis, p. 54, Syria, edoughensis, p. 55, corinthius (Fairm.), p. 57, Algeria, siciliensis, Sicily, p. 59, mecinoides, Tunis, p. 62, demaisoni, Egypt, p. 64, Deserochers, Frelon, ii: p. spp.
- Barilepton famelicum, p. 677, fulciger, p. 678, N. America, Casey, Ann. N. York Ac. vi, n. spp.
- Barinus suffusus, p. 670, difficilis, p. 671, curticollis, p. 673, N. America, Casey, Ann. N. York Ac. vi, n. spp.

- Baris rectirostris, Alexander Mts., FAUST, p. 63, Deutsche e. Z. B. striolala, p. 222, strigosa, p. 223, Cambodia, AURIVILLI Arch. Mus. Paris (3) iii; B. ingens, p. 476, arizonica, p. 478, his gravida, p. 479, callida, p. 481, lubrica, p. 482, soluta, p. 484, ensis, p. 485, vespertina, p. 486, oblongula, p. 487, dilatata, p. cuneipennis, p. 489, aprica, dolosa, p. 490, zuniana, p. 491, o porosicollis, p. 492, punctiventris, p. 493, brunneipes, p. 495, de p. 496, futilis, p. 497, inconspicua, p. 498, subsimilis, socialis, aperta, p. 500, abrupta, tenuestriata, p. 501, discipula, p. 503, lans, p. 504, ansomicans, hyperion, p. 505, vitreola, p. 506, splendens, p. 507, exigua, p. 508, N. America, Casey, Ann. N. Ac. vi: n. spp.
- Barynotus scutatus, Finland, DESBROCHERS, p. 99, Frelon, i, n. sp. Barypeithes longicornis, Abchasia, STIERLIN, p. 410, MT. Schw. er viii, n. sp.
- Blosyrus obliquatus, Congo, DUVIVIER, p. 164, Ann. Ent. Belg. n. sp.
- Brachycerus incommodus, adustus, p. 72, viduatus, p. 73, S. W. PÉRINGUEY, Tr. S. African Phil. Soc. vi (2), n. spp.
- Brachyderes rosti, Caucasus, STIERLIN, p. 411, MT. Schw. ent. Gn. sp.
- Brachyomus histrio, bicostatus, p. 14, sallei, p. 15, Venezuela, S. E. Z. liii, n. spp.
- Brachysomus ornatus, Bohemia, STIERLIN, r. 366, MT. Schw. en viii, n. spp.
- Buttikoferia, n. g., Ulomascides, p. 136, for B. liberiensis, n. sp., W. p. 137, Roelofs, Notes Leyd. Mus. xiv.
- Byctiscus pariei, Laos, Aurivillius, p. 207, N. Arch. Mus. Paris n. sp.
- Cactophagus subnitens, Arizona, Casey, p. 685, Ann. N. York n. sp.
- Calandra rugicollis, Florida, CASEY, p. 687, Ann. N. York Ac. vi, : Calandrinus insignis, p. 619, obsoletus, p. 620, N. America, CASEY, I York Ac. vi, n. spp.
- Camptorhinus cineritius, Obock, FAIRMAIRE, p. 118, Rev. d'Ent. xi Canistes, n. g., near Acalles, for C. schusteri, n. sp., Missouri; p. 447, Ann. N. York Ac. vi.
- Canoixus nigroclavatus, Siam, Aurivillius, p. 214, N. Arch. Mus (3) iii, p. sp.
- Carphonotus, n. g., Cossonini, for C. testaceus, n. sp., Minnesota; p. 694, Ann. N. York Ac. vi.
- Catamonus robustus, Kolbe, referred to Heterostylus, Faust; FAUS1 S. E. Z. liii.
- Catapastus, n. g., Burini, p. 664, for C. diffusus, n. sp., N. America, and Zygobaris conspersus, Lec.
- Cutaponus, n. g., p. 48, for C. curtulus, Senegal, C. (Grypnus) ap. 49, bengalensis, p. 50, E. India, n. spp.; Desurochers, Fre

- Centrinites, n. g., near Centrinus, p. 615, for C. strigicollis, n. sp., N. America, p. 616; Casey, Ann. N. York Ac. vi.
- Centrinogyna, n. g., p. 620, for Centrinus strigata, Lec., and C. procera, n. sp., California, p. 622; Cassy, Ann. N. York Ac. vi.
- Centrinopus, n. g., near Centrinus, p. 601, for C. helvinus, alternatus, n. spp., N. America, p. 602, Casey, Ann. N. York Ac. vi.
- Centrinus tortuosus, p. 580, albotectus, p. 582, grisescens, p. 583, finitimus, hospes, p. 585, clarescens, p. 586, exulans, p. 588, acuminatus, p. 590, globifer, p. 591, nubecula, clientulus, p. 594, denticornis, p. 597, salebrosus, p. 598, pinguescens, p. 599, pulverulentus, p. 600, N. America, Casey, Ann. N. York Ac. vi, n. spp.
- Centyres sinuatus, Victoria, BLACKBURN, p. 129, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Cestophorus, n. g., Anchonina, p. 23, for C. idoneus, Colombia, p. 23, simoni, guerini, mendosus, granulicollis, necessarius, p. 24, Venezuela, tribulus, infidus, p. 25, Colombia: n. spp., and pp. 39-42; Faust, Deutsche e. Z. 1892.
- Ceutorrhynchus balsaminæ, Berne, Guillebeau, Échange i, No. 7, p. 3; C. conformis, Algeria, Grilat, Échange, 1887, No. 32, p. 1: n. spp.
- Charorrhinus squalidus, metamorphoses, XAMBEU, p. 38, Mœurs, ii.
- Chalcocybebus richteri, New Guinea, FAUST, p. 205, S. E. Z. liii, n. sp.
- Chaleponotus, n. g., near Chalcodermus, for C. elusus, n. sp., Indiana; CASEY, p. 444, Ann. N. York Ac. vi.
- Cherrus inconspicuus, p. 53, longulus, maurulus, p. 54, Australia, BLACKBURN, Tr. R. Soc. S. Austr. xvi: n. spp.
- Chiloneus ottomanus, Turkey, DESBROCHERS, p. 113, Frelon, i, n. sp.
- Chirozetus elaphus, Sumatra, FAUST, p. 221, S. E. Z. liii, n. sp.
- Cimbus ovampoensis, S. W. Africa, Péringuey, p. 67, Tr. S. African Phil. Soc. vi (2), n. sp.
- Cionus albopunctatus, Laos, Aurivillius, p. 218, N. Arch. Mus. Paris (3) iii, n. sp.
- Cleonidius, n. subg. of Cleonus, p. 186, including the following n. spp. from N. America, grandirostris, p. 186, bicarinatus, lecontei, lobigerinus, p. 187, kirbyi, = (vittatus, Lec.), circumductus, subcylindricus, graniferus, p. 188; CASEY, Ann. N. York Ac. vi.
- Cleonus perlatus and allies, synonymy; Desbrochers, pp. 18-20, Frelon, i.
- C. (Bothynoderes) dissimilis, S. W. Africa, Péringuey, p. 75, Tr. S. African Phil. Soc. vi (2), n. sp.
- Cneorhinus of Europe and allies, revision of; DESBROCHERS, Frelon, i, pp. 52, &c.
- Colobodes turbatus, New Guines, FAUST, p. 210, S. E. Z. liii, n. sp.
- Compsus virginalis, biimpressus, p. 9, simoni, p. 10, cræsus, p. 11, spectabilis, p. 12, gemmeus, p. 13, Venezuela, Faust, S. E. Z. liii, n. spp.
- Conoproctus: distinctions of sexes in; Casey, p. 540, Ann. N. York Ac. vi.

- Conorhynchus dissimilatus, Mén., = (pistor, Chev.); FAUST, p. 5 E. Z. liii.
- Conotrachelus compositus, p. 439, carinifer, p. 440, integer, duplex, p. rotundus, p. 442, N. America, Casey, Ann. N. York Ac. vi, n. s
- Corigetus humilis, Somaliland, Gestro, p. 783, Ann. Mus. Genov. (2
  C. pavici, Siam, p. 210, lineatus, Laos, p. 211, Aurivillius, N. Mus. Paris (3) iii; C. discolor, neglectus, Sumatra, p. 190, albot p. 191, Borneo, Faust, S. E. Z. liii: n. spp.
- Cossonides: table of the European genera; Deserochers, Frek pp. 72-82.
- Cotaster referred to Hylobiides; DESBROCHERS, p. 70, Frelon, ii.
- Cratopus aurostriatus, Seychelles, FAIRMAIRE, p. cli, Bull. Soc. Ellxi, n. sp.
- Ctenolobus, n. g., near Lacordaireus, for C. squamiger, n. sp., Ma: Desbrochers, Frelon, i (cf. Faust, S. E. Z. liii, p. 52).
- Cubicorhynchus mussoni, N. S. Wales; BLACKBURN, p. 124, P. Lini N.S.W (2) vii.
- Cyclomaurus moraguesi, p. 117, boghariensis, p. 118, Algeria, vau p. 119, Tunis, Desbrochers, Frelon, i, n. spp.
- Cyclomias, n. g. (no affinities given, probably Otiorhynchides) 1 mylacoides, n. sp., Asia Minor; Desbrochers, Frelon, i, p. 12
- Cylindromus, n. g., Cyphicerinarum, for C. plumbeus, n. sp., Caml AURIVILLIUS, p. 213, N. Arch. Mus. Paris (3) iii.
- Cylloderus josephus, Congo, DUVIVIER, p. 58, Ann. Ent. Belg. xxxvi. Cylloproctus, n. g., Sitoninarum, p. 22, for C. murinus, Venezuela, mo Colombia, p. 23, pyriformis, vehemens, p. 24, simoni, p. 25, Venn. spp.; FAUST, S. E. Z. liii.
- Cyphicerus erinaceus, Sumatra, p. 191, præcanus, New Guinea, 1 FAUST, S. E. Z. liii; C.? nigrofasciatus, Laos; AURIVILLIUS, N. Arch. Mus. Paris (3) iii: n. spp.
- Dereodus cylindricollis, p. 781, elongatus, p. 782, Somaliland, G. Ann. Mus. Genov. (2) xii, n. spp.
- Dermatodes albarius, Engano, p. 185, succinctus, aptus, Sumatra, FAUST, S. E. Z. liii, n. spp.
- Desmidophorus fasciatus, Laos, AURIVILLIUS, p. 221, N. Arch. Mus (3) iii; D. dohrni, Sumatra, p. 211, prætor, Almahera, p. 212, S. E. Z. liii: n. spp.
- Desmoglyptus, n. g. (Barini), for Ampeloglypter crenatus, Lec.; p. 550, Ann. N. York Ac. vi.
- Desmoris, Lec., merged in Smicronyx; CASEY, p. 382, Ann. N Ac. vi.
- Dialeptopus approximatus, W. Australia, BLACKBURN, p. 124, P Soc. N.S.W. (2) vii, n. sp.
- Diaphorus, n. g, Cyphinarum, for D. morosus, n. sp., Venezuela; p. 7, S. E. Z. liii.
- Diastrophilus, n. g., Hyperinarum, p. 31, for D. astutus, n. sp., Ven p. 32, FAUST, S. E. Z. liii.

- Dinocleus, n. n. for Centrocleonus, Lec., nec Chev., p. 176, with D. jacobinus, p. 179, denticollis, p. 180, farctus, p. 181, saginatus, p. 182, alborestitus, p. 183, wickhami, p. 184, densus, p. 185, n. spp., N. America; Casey, Ann. N. York Ac. vi.
- Dioptrophorus, n. g., Anchoninæ, p. 35, for D. sharpi, Guatemala, p. 58, simplex, Mexico, fahræi, Guatemala, p. 59, n. spp.; FAUST, Deutsche e. Z. 1892.
- Dorytomus inæqualis, p. 364, brevisetosus, p. 365, amplus, parvicollis, p. 367, hystricula, filiolus, p. 370, nubeculinus, p. 371, cuneatulus, alaskanus, p. 374, marginatus, indifferens, p. 375, vagenotatus, p. 376, fusciceps, p. 377, marmoreus, p. 381, N. America, Casey, Ann. N. York Ac. vi, n. spp.
- Dyscerus, n. g., for D. virgatus, Sumatra, p. 198, cruciatus, Java, p. 199, linnei, Sumatra, p. 200, n. spp.; FAUST, S. E. Z. liii.
- Dyschænium, position is Erirhinides; BLACKBURN, p. 56, Tr. R. Soc. S. Austr. xv.
- Dysostines glaber, p. 127, advena, p. 128, Victoria, BLACKBURN, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Edmundia sericea, Somaliland, GESTRO, p. 780, Ann. Mus. Genov. (2) xii, n. sp.
- Elleschus angustatus, Dietz, referred to Dorytomus; CASEY, p. 363, Ann. N. York Ac. vi.
- Ellimenistes ostentatus, S. W. Africa, Péringuey, p. 70, Tr. S. African Phil. Soc. vi (2), n. sp.
- Elytrodon baudii, Sicily, STIERLIN, p. 363, MT. Schw. ent. Ges. viii; E. baudii, Sicily, STIERLIN, p. 156, Nat. Sicil. xi; E. impressifrons, p. 115, carinifrons, p. 116, Algeria, DESBROCHERS, Frelon, i: n. spp.
- Emplesis monticola, p. 142, ocellata, p. 143, Victoria, BLACKBURN, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Enicoderus, n. g., near Siderodactylus, for E. thoracicus, n. sp., Namaqualand; Péringuey, p. 127, Tr. S. African Phil. Soc. vi.
- Epistrophus cristulatus, Venezuela, FAUST, p. 32, S. E. Z. liii, n. sp.
- Eremnini: notes on Australian; BLACKBURN, p. 229, Tr. R. Soc. S. Austr.
- Eremotes heydeni, Turkestan, FAUST, p. 104, Deutsche e. Z. 1892, n. sp.
- Eristus, n. g. (aberrant Erirhinidæ), p. 148, for E. setosus, bicolor, n. spp., Victoria, p. 149, BLACKBURN, P. Linn. Soc. N.S.W. (2) vii.
- Esmelina australis, N. S. Wales, BLACKBURN, p. 123, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Essolithna soriata, Australia, BLACKBURN, p. 50, Tr. R. Soc. S. Austr. xvi, n. sp.
- Eugithopus nobilis, Borneo, Roelofs, p. 7, Notes Leyd. Mus. xiv, n. sp. Eugnamptus bifenestratus, apicalis, Venezuela, Faust, S. E. Z. liii, p. 42, n. spp.
- Eunyssobia, n. n. for Euchates, Lec., with discussion of characters; Casey, p. 679, Ann. N. York Ac. vi.
- Eupholus compositus, New Guinea, FAUST, p. 188, S. E. Z. liii, n. sp.

- Evas everardensis, Australia, BLACKBURN, p. 47, Tr. R. Soc. S. . xvi, n. sp.
- Exophthalmida, n. g., Cyphinarum, p. 187, for E. glauca, n. sp., Adm and Solomon Is., p. 188; FAUST, S. E. Z. liii.
- Foucartia karamani, Dalmatia, STIERLIN, p. 364, MT. Schw. ent. Ge F. conicicollis, Ordubad, REITTER, p. 153, Wien. ent. Z. xi: n.
- Geosomus, n. g., Pantopeinarum, for G. macleayi, n. sp., Gawler; F p. 179, S. E. Z. liii.
- Gerynassa affinis, p. 144, minor, p. 145, Victoria, BLACKBURN, P. Soc. N.S.W. (2) vii, n. spp.
- Glyptobaris, n. g. (Barini), for Onychobaris rugicollis, Lec.; CASEY, | Ann. N. York Ac. vi.
- Gonipterus rufus, Victoria, BLACKBURN, p. 130, P. Linn. Soc. N.S.V vii, n. sp.
- Grypnus, n. subg. of Cataponus (q.v.); DESBROCHERS, p. 49, Frelon, Gymnetron nigritarse, p. 93, hircinum, p. 95, canescens, p. 96, A. furcatum, p. 94, aper, p. 95, hirsutum, p. 96, Syria, DESBROC Frelon, ii, n. spp.
- Hadromerus scabricollis, Venezuela, FAUST, p. 1, S. E. Z. liii, n. sp. Haplorhynchus valdani, Roel., = (Cyrtopisthen rubicundum, At ROELOFS, p. 36, Notes Leyd. Mus. xiv.
- Heilipus? mediator, Venezuela, FAUST, p. 38, S. E. Z. liii, n. sp.
- Hesperobaris, n. g., Barini, p. 559, for H. suavis, n. sp., Texas, p. Casey, Ann. N. York Ac. vi.
- Himatium nigritulum, Florida, CASEY, p. 691, Ann. N. York. Ac. vi, Hipporrhinus ovampoensis, S. W. Africa, Péringuey, p. 75, Tr. S. A Phil. Soc. vi (2), n. sp.
- Hylobiini: tabulation of genera; FAUST, pp. 201-204, S. E. Z. liii. Hyphantus maculifer, p. 16, distinguendus, p. 17, uncinatus, dehiscens, Brazil, DESBROCHERS, Frelon, ii, n. spp.
- Hypomeces atomarius, Java, p. 47, laniger, Malay Arch., p. 48, BROCHERS, Frelon, i, n. spp.
- Hypsonotus umbilicatus, Venezuela, FAUST, p. 28, S. E. Z. liii, n. sp Idiostethus, n. g. Barini, p. 649, for I. ellipsoideus, dispersus, p. 652, 1 N. America, Camptorhinus tubulatus, Say, and Zygobaris sub-Lec.; Casey, Ann. N. York Ac. vi.
- Iphthimorhinus, n. g., near Rhynchophorus, p. 207, for I. australasia, Queensland, p. 208; Roelofs, Notes Leyd. Mus. xiv.
- Ithyporus nigrosignatus, Congo, Duvivier, p. 316, Ann. Ent. xxxvi, n. sp.
- Larinus numidicus, p. 89, villosicollis, p. 91, Algeria, villosiventris, p. 90, distinguendus, Syria, p. 92, Desbrochers, Frelon, ii; 1 viusculus, Asia Minor, Desbrochers, p. 12, t. c.: n. spp.
- Leprosomus sulcicollis, p. 21, schonherri, contractus, vicarius, p. 22, (bia, FAUST, Deutsche e. Z. 1892, n. spp.
- Leptopsini: value of generic characters discussed; Blackburn, p. 4 R. Soc. S. Austr. xvi.

- Leptops biordinatus, p. 55, raucus, p. 56, gravis, p. 57, areolatus, planicollis, p. 58, interioris, p. 59, contrarius, sulcicollis, p. 60, Australia, BLACKBURN, Tr. R. Soc. S. Austr. xvi, n. spp.
- Leptostylus, n. g., near Megalostylus, for Cyphus juvencus, Ol., and L. suturalis, n. sp., Venezuela, FAUST, p. 8, S. E. Z. liii.
- Leucodrusus maria, characters of; FAUST, p. 45, S. E. Z. liii.
- Linonotus, n. g., for Centrinus distinctus, Boh.; CASEY, p. 603, Ann. N. York Ac. vi.
- Limnobaris bracata, p. 627, limbifer, blandita, p. 628, tabidu, p. 629, deplanata, p. 630, denudata, p. 632, planiuscula, p. 633, oblita, p. 634, seclusa, p. 635, ebena, p. 638, puteifer, p. 639, concurrens, p. 640, fratercula, seminitens, p. 642, nitidissima, p. 644, N. America, Casey, Aun. N. York Ac. vi, n. spp.
- Liophlaus paulinoi, Portugal, DESBROCHERS, p. 88, Frelon, i, n. sp.
- Lixus: table of the food-plants of N. American; WEBSTER, p. 340, P. E. Soc. Wash. ii. L. fascicularis, defloratus, sturmi, synonymical note, p. 44; turkestanus, acicularis, cinerascens, noticed, p. 46, FAUST, S. E. Z. liii.
- L. fecundus, p. 61, imitator, p. 62, Alexander Mts., christophi, Caucasus, p. 63, Faust, Deutsche e. Z. 1892; L. albopictus, farinifer, Araxes Valley, Reitter, p. 64, Wien. ent. Z. xi; L. itimbirensis, Congo, Duvivier, p. 164, Ann. Ent. Belg. xxxvi; L. amplexus, p. 199, oregonus, p. 200, eximius, tenellus, p. 201, pygmaus, p. 203, soror, laramiensis, p. 204, sobrinus, p. 205, semivitatus, p. 207, jülichi, p. 208, luculentus, p. 209, nitidulus, p. 210, obesulus, p. 211, sexualis, p. 212, N. America, Casey, Ann. N. York Ac. vi: n. spp.
- Lobotrachelus, = (Colosternus, Motsch.); FAUST, p. 47, S. E. Z. liii.
- Madarellus, n. g., for Rhynchænus undulatus, Say; CASEY, p. 540, Ann. N. York Ac. vi.
- Madarus, composition of, and sexual distinctions in; Casey, p. 540, Ann. N. York Ac. vi.
- Mandalotus, Er., = (Dysostines, Pasc.); BLACKBURN, p. 229, Tr. R. Soc. S. Austr. xv.
- Mechistocerus, Fauv., = (Berosiris, Pasc.); FAUST, p. 46, S. E. Z. liii.
- M. pascoei, Sumatra, FAUST, p. 213, S. E. Z. liii, n. sp.
- Mecysmoderes, = (Caliosomus, Motsch.); FAUST, p. 47, S. E. Z. liii.
- M. rhomboidalis, Cambodia, Aurivillius, p. 221, N. Arch. Mus. Paris (3) iii, n. sp.
- Medicasta lugubris, Victoria, BLACKBURN, p. 131, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Meira baudii, Rome, STIERLIN, p. 362, MT. Schw. ent. Ges. viii; M. subplana, Algeria, sicula, Sicily, p. 4, subconiceps, Algeria, p. 5, tunicensis, Tunis, p. 6, DESBROCHERS, Frelon, ii: n. spp.
- Menetypus opalescens, kirschi, p. 2, modestus, p. 3, robustus, p. 4, Venezuela, FAUST, S. E. Z. liii, n. spp.
- Meriphus humeralis, N. S. Wales, BLACKBURN, p. 146, P. Linn. Soc. N.S.W. (2) vii, n. sp.

- Metacinops calabrus, Sicily, STIERLIN, p. 361, MT. Schw. ent. Ges. n. sp.
- Metallites damryi, Sardinia, DESBROCHERS, p. 2, Frelon, ii, n. sp.
- Metopotoma, n. g., Cossonini, near Gononotus, p. 689, for M. repens, n. California, p. 690; Casey, Ann. N. York Ac. vi.
- Microbaris, n. g., Barini, p. 560, for M. galvestonica, n. sp., Texas, p. & Casey, Ann. N. York Ac. vi.
- Micromastus, Lec., characters amended; Casey, p. 446, Ann. N. York vi.
- Minyops bertolinii, Rome, STIERLIN, p. 364, MT. Schw. ent. Ges. viii, r Mitophorus globosus, p. 778, humerosus, p. 779, Somaliland, GESTRO, 1 Mus. Genov. (2) xii, n. sp.
- Myllocerus benignus, Askhabad, FAUST, p. 60, Deutsche e. Z. 1892; debilis, sparsutus, Obock, FAIRMAIRE, p. 117, Rev. d'Ent. xi: n.
- Myossita munda, crucigera, Australia, BLACKBURN, P. Linn. Soc. N.8 (2) vii, n. spp.
- Nauphæus simius, Philippines, FAUST, p. 222, S. E. Z. liii, n. sp.
- Nemosinus, n. g., near Styphloderes, for Anchonus angulicollis, S. FAUST, p. 18, Deutsche e. Z. 1892.
- Neotocerus, n. g., near Sophrorhinus, for N. fausti, n. sp., Congo; D. vier, Ann. Ent. Belg. xxxvi, p. 165.
- Nicentrus, n. g., p. 608, for some species of Contrinus and N. inger p. 610, scitulus, p. 611, effetus, contractus, p. 613, n. spp., N. Ame Casey, Ann. N. York Ac. vi.
- Notaris dalmatina, Dalmatia, STIERLIN, p. 365, MT. Schw. ent. Ges. n. sp.
- Nyssonotus, n. g., Cossonini, for N. seriatus, n. sp., Texas; Casey, p. Ann. N. York Ac. vi.
- Oligolochus, n. g., for Zygobaris? convexus, Lec.; Casey, p. 648, Anı York Ac. vi.
- Omoionotus: generic characters; FAUST, p. 29, S. E. Z. liii.
- O. jekeli, p. 29, lacordairei, p. 30, Venezuela, FAUST, S. E. Z. liii, n. sp. Omotemnus rhinoceros, Q noticed; Aurivillius, p. 223, N. Arch. Paris (3) iii.
- Onychobaris corrosa, depressa, p. 525, millepora, p. 526, austera, p. insidiosa, p. 528, arguta, p. 529, audax, stictica, p. 530, mystica, p. egena, p. 532, ambigua, pauperella, p. 533, remota, p. 535, mo p. 536, illex, p. 537, diluta, p. 538, porcata, p. 539, N. America, C. Ann. N. York Ac. vi, n. spp.
- Onychogymnus ursulus, Madagascar, p. 50, O.? ocellatus, Gaboon, p Roelofs, Notes Leyd. Mus. xiv, n. spp.
- Oomorphidius, n. g., Barini, for Microcholus erasus and lævicollis, I CASEY, p. 659, Ann. N. York. Ac. vi.
- Oopterinus, n. g., for Otidocephalus perforatus, Horn; Casey, p. 438, . N. York Ac. vi.
- Opetiopteryx, n. g., Amycterinæ, p. 125, for O. frigida, n. sp., Vict p. 126, Blackburn, P. Linn. Soc. N.S.W. (2) vii.

- Orchestes fagi, metamorphoses; Pissot, p. 91, Le Nat. 1892.
- Orthorhinus albosparsus, New Guinea, FAUST, p. 204, S. E. Z. liii; O. bicolor, Victoria, BLACKBURN, p. 139, P. Linn. Soc. N.S.W. (2) vii: n. spp.
- Orthoris cylindrifer, Arizona, Casey, p. 571, Ann. N. York Ac. vi, n. sp.
- Otidocephalus nivosus, p. 429, insignis, estriatus, p. 430, egregius, p. 431, floridanus, p. 433, speculator, p. 434, ruficornis, p. 435, cavirostris, p. 437, N. America, Casey, Ann. N. York Ac. vi, n. spp.
- Otiorrhynchus proreus belongs to subg. Arammichus; REITTER, p. 26, Wien. ent. Z. xi. O. valachiæ, distinctive characters; FRIVALDSZKY, p. 132, Term. füzetek, xv.
- O. dilatipes, France, Guillebeau, Échange, i, No. 8, p. 2; O. nevadensis, Spain, p. 357, latifrons, Abruzzi, p. 358, O. (Tournieria) brevipennis, Greece, thoracicus, Pennine Alps, p. 359, Stierlin, MT. Schw. ent. Ges. viii; O. kelecsenyii, Hungary, Frivaldszky, p. 132, Term. füzetek, xv; O. abchasicus, Caucasus, Rost, p. 402, Deutsche e. Z. 1892; O. (Tournieria) hystericus, Alexander Mts., Faust, p. 61, Deutsche e. Z. 1992; O. caroli, gracilicornis, p. 408, O. (Tournieria) subparallelus, p. 409, Caucasus, bugnioni, Tunis, p. 410, Stierlin, MT. Schw. ent. Ges. viii: n. spp.
- Ozyopisthen rittatum, and linea-alba, Thoms., noticed, referred to Ichthyopisthen, p. 34, also clavatum, Roel., = (rufo-clavatum, Auriv.), p. 35; Roelofs, Notes Leyd. Mus. xiv.
- Pachyrhynchus rubro-verrucatus, N. Guinea, TRYON (953), n. sp.
- Pachytychius simillimus, Lalla-Marghnia, smyrnensis, Smyrna, letourneuxi, Ramleh, abeillei, Jerusalem, siculus, Sicily, Desbrochers, p. 30, Frelon, i, n. spp.
- Pagiophloeus, n. g., Hylobiinorum, p. 195, type Aclees pacca, Fab., with P. javanicus, p. 196, rühli, p. 197, n. spp., Java; FAUST, S. E. Z. liii.
- Pandeleteius angustirostris, Venezuela, FAUST, p. 5, S. E. Z. liii, n. sp. Pantorhytes, n. g., Pachyrhynchinarum, for Pachyrhynchus chrysomelas,
- Montr., and P. batesi, n. sp., New Guinea; FAUST, p. 193, S. E. Z. liii.
- Parameleus, n. g., Hylobiinorum, p. 33, for P. stevensi, p. 34, insignatus, neglectus, p. 35, n. spp., Venezuela; FAUST, S. E. Z. liii.
- Paraphilernus, n. g., for Philernus bilunulatus, Db.; Desbrochers, p. 71, Frelon, ii.
- Paraplinthus, n. g., for Plinthus carinatus, Bob.; FAUST, p. 49, 8. E. Z. liii.
- Parascytopus baudii, Calabria, STIERLIN, p. 361, MT. Schw. ent. Ges. viii, n. sp.
- Paratasis viridiænea, Java, Heller, p. 269, Notes Leyd. Mus. xiv, n. sp. Pelororhinus proximus, p. 134, crassus, p. 135, Australia, Blackburn, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Pentarthrinus, n. g., Cossonini, p. 698, for Amaurorhinus nitens, Horn, and P. parvicollis, p. 699, piceus, atrolucens, p. 700, n. spp., N. America; Casey, Ann. N. York Ac. vi.

- Pephricus umbratus, W. Australia, p. 231, squalidus, S. Australia, p. Blackburn, Tr. R. Soc. S. Austr. xv, n. spp.
- Peritelus lostiæ, Sardinia, DESBROCHERS, Frelon, ii, p. 88; P. vit Sicily, DESBROCHERS, p. 7, t. c.: n. spp.
- Phacepholis albaria, Venezuela, FAUST, p. 6, S. E. Z. liii, n. sp.
- Phænoderus, n. g., near Siderodactylus, for P. distinctus, n. sp., S. Africa; Péringuey, Tr. S. African Phil. Soc. vi (2), p. 69.
- Phyllobius brevis, Gyll., = (emgei, Stierl.); Reitter, p. 239, Wien. Z. xi.
- P. siculus, Sicily, STIERLIN, p. 360, MT. Schw. ent. Ges. viii; siculus, Sicily, BAUDI, p. 156, Nat. Sicil. xi; P. delagrangei locality), DESBROCHERS, p. 3, Frelon, ii: n. spp.
- Phyrdenus bullatus, Arizona, Casey, p. 458, Ann. N. York Ac. n. sp.
- Phytobius griseomicans, N. America, Schwarz, p. 165, P. E. Soc. W ii, n. sp.
- Phytoscaphus, = (Rhypochromus, Motsch.); FAUST, p. 47, S. E. Z. lii P. interstitialis, p. 215, P. ? setosus, p. 216, Laos, AURIVILLIUS, N. A Mus. Paris (3) iii; P. leporinus, Sumatra, FAUST, p. 193, S. I liii: n. spp.
- Piasomias limbatellus, p. 66, sumptuosus, p. 67, S. W. Africa, Péringi Tr. S. African Phil. Soc. vi (2); P. desgodinsi, p. 115, Tibet, fa p. 116, trapezicollis, p. 117, China, Frivaldszky, Term. füze xv: n. spp.
- Platyopisthen, n. g., for Stenophida (sub. Oxyopisthen) suturale, R. Roeloff, p. 134, Notes Leyd. Mus. xiv.
- Plesiobaris, n. g., p. 509, for P. signatipes, p. 510, æmula, p. 512, disju: p. 513, n. spp., N. America, and including Baridius T-signum, I and Pseudobaris albilatus, Lec.; CASEY, Ann. N. York Ac. vi.
- Polydrusus pulchellus, Dalmatia, STIERLIN, p. 363, MT. Schw. ent. viii; P. delagrangei, Asia Minor, Desbrochers, p. 1, Frelon, n. spp.
- Polymicrus, n. g., Hylobiinorum, for P. tessellatus, p. 36, tuitus, p. 37, n. 1 Venezuela; Faust, S. E. Z. liii.
- Polyphrades tumidulus, p. 51, rugulosus, p. 52, Australia, BLACKBURN, R. Soc. S. Austr. xvi, n. spp.
- Promecops luteus, p. 25, jucundus, p. 26, olivieri, p. 27, stimulans, p. Venezuela, FAUST, S. E. Z. liii, n. spp.
- Promecotarsus, n. g., near Smicronyx, p. 408, for P. maritimus, p. densus, fumatus, p. 410, n. spp., N. America; Casey, 1 N. York Ac. vi.
- Proxyrodes, n. g. (Otiorhynchini, near Proxyrus), for P. maculatus, n. Australia; BLACKBURN, p. 48, Tr. R. Soc. S. Austr. xvi.
- Prypnus (?) squamosus, N. S. Wales, BLACKBURN, p. 115, P. Linn. N.S.W. (2) vii, n. sp.
- Pseudaclees, n. g., for Hylobius fasciatus, Pasc.; FAUST, S. E. Z. p. 50.

- Pseudanchonus, n. g., Heilipinarum, for P. æquatorius, n. sp., Ecuador; FAUST, p. 203, S. E. Z. liii.
- Pseudobaris luctuosa, Florida, discretu, Texas, p. 554, fausta, Arizona, p. 555, lugubris, p. 557, cælata, p. 559, New Mexico, Casey, Ann. N. York Ac. vi, n. spp.
- Pseudobarynotus, n. g., for laticeps, Desb.; Desbrochers, p. 102, Frelon, i (cf. Faust, S. E. Z. liii, p. 52).
- Pseudocholus bivittatus, p. 224, lacordairei, p. 225, New Guinea, viridimicans, Amboina, p. 226, FAUST, S. E. Z. liii, n. spp.
- Pseudopentarthrum robustum, simplex, N. America, Casky, Ann. N. York Ac. vi, n. spp.
- Psomus, n. g., Zygopini, near Acoptus, for P. politus, n. sp., Indiana; Casey, p. 458, Ann. N. York Ac. vi.
- Ptochus interruptus, Ephesus, p. 122, vagepictus, Peloponnesus, p. 123, Desbrochers, Frelon, i, n. spp.
- Pycnobaris, n. g., p. 514, for Baris pruinosa, Lec., and P. squamotecta, Texas, n. sp., p. 515; Casey, Ann. N. York Ac. vi.
- Raymondia referred to Erirhinides; DESBROCHERS, p. 70, Frelon, ii.
- Rhadinomerus, n. g., for Mechistocerus (? Berosiris) mastersi, Pasc.; FAUST, p. 46, S. E. Z. liii.
- R. vigilans, p. 215, mæstus, solidus, p. 216, simplex, p. 217, Sumatra, FAUST, S. E. Z. liii, n. spp.
- Rhamphocolus, n. g., near Rhyncolus, p. 702, for R. tenuis, n. sp., Texas, p. 703; Casey, Ann. N. York Ac. vi.
- Rhinaria tibialis, p. 136, debilis, p. 138, Victoria, BLACKBURN, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Rhinochenus stigma, metamorphoses; DECAUX, p. 202, Le Nat. 1892.
- Rhinochrosis, n. g., near Mylacus, p. 121, for R. subglobatus, n. sp., Tangier, p. 122, Desbrochers, Frelon, i.
- Rhynchites interpunctatus, Steph., distinguished from alliariæ; Thomson, p. 1773, Opusc. ent. xvi. R. seminiger, characters of; Reitter, p. 186, Wien. ent. Z. xi.
- Rhyncholus syriacus, Syria, p. 80, dalmatinus, Dalmatia, p. 81, Des-BROCHERS, Frelon, ii; R. pallens, p. 703, spretus, dilatatus, p. 704, relictus, p. 705, nimius, p. 706, discors, p. 707, N. America, CASEY, Ann. N. York Ac. vi: n. spp.
- Rhyparonotus, n. g., Anchoninæ, p. 33, for R. mendax, p. 54, jekeli, p. 55, Venezuela, n. spp.; FAUST, Deutsche e. Z. 1892.
- Rhytirhinus confusus, Greece, p. 8, carthagenensis, Spain, p. 9, varipes, p. 10, monstruosus, p. 11, Algeria, Desbrochers, Frelon, ii, n. spp.
- Ryssematus ovalis, Texas, CASEY, p. 443, Ann. N. York Ac. vi, n. sp.
- Schizonotus, n. g., Erirhinini, p. 708, for S. cacus, n. sp., California, p. 709; CASEY, Ann. N. York Ac. vi.
- Sciaphilus subnudus, Greece, DESBROCHERS, p. 114, Frelon, i, n. sp.
- Seidlitzia, n. g., for Liophlaus maroccanus, Fairm.; Desnochers, p. 93, Frelon, i (cf. Alophus, suprà).

- Sepionus, n. g., Cyphicerinarum, p. 211, type Episomus parallelus, Chwith S. frontalis, Cambodia, p. 212, tuberculatus, Battambang, p. 2 Aurivillius, N. Arch. Mus. Paris (3) iii.
- Siderodactylus elegans, egregius (simplicipes, olim.), p. 68, opalinus, p. S. W. Africa; Péringuey, Tr. S. African Phil. Soc. vi (2).
- Simophorus, n. g., Phyllobiinarum, for S. pustulosus, n. sp., Venezu Faust, S. E. Z. liii, p. 19.
- Sipalus aurivilli, Congo, DUVIVIER, p. 166, Ann. Ent. Belg. xxxvi, n. Smicronyx lineolatus, p. 385, quadrifer, p. 388, profusus, p. intricatus, p. 390, imbricatus, p. 391, silaceus, p. 392, spurcus, p. sparsus, p. 394, pleuralis, p. 395, connivens, p. 398, fiducialis, p. congestus, p. 401, eagittatus, p. 402, eculpticollis, instabilis, p. apionides, perpusillus, p. 405, defricans, p. 406, gibbirostris, squal p. 407, N. America, Casey, Ann. N. York Ac. vi, n. spp.
- Sophronopterus, n. g., near Mechistocerus, for S. humeralis, n. sp., Sum: FAUST, p. 214, S. E. Z. liii.
- Sphenocorynus scutellatus, Sumatra, FAUST, p. 227, S. E. Z. liii, n. sp. Sphenophorus cinerascens, Motsch., referred to Trochorhopalus; F2 p. 50, S. E. Z. liii.
- Stenancylus, n. g., Cossonini, near Rhyncolus, for S. colomboi, n. Florida; Casey, p. 694, Ann. N. York Ac. vi.
- Stenophida (sub Oxyopisthen) suturale, Roel., = (trilineata, Aur Roelors, p. 36, Notes Leyd. Mus. xiv.
- Stethobaris incompta, p. 655, congermana, egregia, p. 657, N. Ame Casey, Ann. N. York Ac. vi, n. spp.
- Stictobaris, n. g. (Barini), p. 516, for S. pimalis, p. 517, subacuta, p. n. spp., N. America, and Onychobaris cribrata, Lec.; Casey, An York Ac. vi.
- Stigmatotrachelus placidus, S. W. Africa, Péringuey, Tr. S. African Soc. vi (2) p. 70, n. sp.
- Strophomorphus uniformis, Syria, DESBROCHERS, p. 117, Fredon, i, n. Styphlus ursus, Greece, DESBROCHERS, p. 89, Fredon, ii, n. sp.
- Sympiezomias basalis, p. 207, Siam, setosus, Laos, p. 208, AURIVIL N. Arch. Mus. Paris (3) iii, n. spp.
- Synthlibonotus: systematic position discussed, FAUST, p. 17, S. E. Z. S. rufipes, p. 17, scapha, mucronotus, p. 18, Venezuela, FAUST, S. E. Z. n. spp.
- Tanymecus and allies, characters tabulated; FAUST, S. E. Z. liii, p. 5 T. (Esamus) piliscapus, Cambodia, Aurivillius, p. 208, N. Arch. Paris (3) iii, n. sp.
- Temnoschoita eruditus, basipennis, Congo, Duvivier, Ann. Ent. xxxvi, n. spp.
- Tentegia, Pasc., characters and position; FAUST, p. 181, S. E. Z. liii. T. basalis, p. 181, ingrata, p. 182, sana, p. 183, Australia, FAUST, S. liii, n. spp.
- Thylacites subvittatus, Spain, p. 110, cretaceus, p. 111, subahenus, p. Algeria, Desbrochers, Frelon, i, n. spp.

- Thysanocnemis graphica, p. 425, squamiger, horridula, p. 426, N. America, Casey, Ann. N. York Ac. vi, n. spp.
- Timus, n. g., near Siderodactylus, for T. simplex, p. 128, planiceps, p. 129, n. spp., S. Africa; Péringuey, Tr. S. African Phil. Soc. vi (2).
- Titinia and Idaspora discussed; BLACKBURN, p. 117, P. Linn. Soc. N.S.W. (2) vii.
- T. brevicollis, p. 118, læta, T. (Idaspora?) eremita, p. 119, bicolor, p. 120, Australia, Blackburn, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Trachyphlaus scabriculus and laticollis are the sexes of one species; Guillebeau, p. celv, Bull. Soc. Ent. Fr. lxi.
- Tragopus fossulatus, Oelebes, brevis, p. 218, sulcifrons, p. 219, Java, FAUST, S. E. Z. liii, n. spp.
- Trepobaris, n. g., for T. elongata, n. sp., Texas; Casey, p. 519, Ann. N. York Ac. vi.
- Trichobaris insolita, p. 565, compacta, p. 566, cylindrica, p. 567, Casey, Ann. N. York Ac. vi, n. spp.
- Trigonosomus, n. g., near Rhinochenus, for T. draco, n. sp., New Guinea; FAUST, p. 220, S. E. Z. liii.
- Trochorhopalus sumatranus, Sumatra, FAUST, p. 227, S. E. Z. liii, n. sp.
- Tychius longierus, p. 99, parallelogrammus, p. 100, Algeria, DESBROCHERS, Frelon, ii; T. soltaui, p. 416, lamellosus, p. 418, prolixus, p. 419, variegatus, p. 420, simplex, sibinoides, p. 421, mica, p. 422, subfasciatus, p. 423, hispidus, p. 424, N. America, Casey, Ann. N. York Ac. vi: n. spp.
- Tyloderma angustula, p. 451, contusa, p. 452, rufescens, p. 454, subpubescens, p. 455, N. America, Casey, Ann. N. York Ac. vi, n. spp.
- Xanthochelus miscellaneus, Lower India, FAUST, S. E. Z. liii, p. 48, n. sp. Yuccaborus sharpi, p. 688, Mexico, grossus, Texas, p. 689, Casey, Ann. N. York Ac. vi, n. spp.
- Zygops suffusus, Texas, CASEY, p. 459, Ann. N. York Ac. vi.

#### CERAMBYCIDE.

- [Cf. Balbi (31), Bates (36), Blackburn (90, 93, 94), Brongniart (133), Casey (161), Distant (210, 216), Duvivier (230, 231), Fairmaire (262, 266, 268, 269), Frivaldszey (309, 310), Gahan (315, 316), Gestro (324, 334), Lameere (534), Nonfried (644, 645, 647, 648), Olliff (655), Péringuey (672, 673), Pic (677, 678, 679, 681, 683, 686, 688, 691), Poll (697), Reitter (725), Bitsema (760, 763, 764), Théry (921), Villard (976).]
- Acanthoderes nigritarsis, White, = (sylvanus, Bates); GAHAN, p. 264, Tr. E. Soc. 1892.
- A. piperatus, p. 262, pl. xii, fig. 4, signatus, p. 263, fig. 3, Mexico, Gahan, Tr. E. Soc. 1892, n. spp.
- Acmaops: synonymy of the N. American species; Casey, p. 38, Ann. N. York Ac. vi.
- A. variipes, California, CASEY, p. 38, Ann. N. York Ac. vi, n. sp.

- Acyphoderes cribricollis, Mexico, Bates, p. 160, pl. vi, fig. 6, Tr. E. 1892, n. sp.
- Elaru severa, Congo. DUVIVIER, p. 360, Ann. Ent. Belg. xxxvi, n.: Ayada, n. g., near Callidium, FAIRMAIRE, p. clxxi, for A. clavico n. sp., Madagascar; FAIRMAIRE, Bull. Soc. Ent. Fr. lxi.
- Alphitopola robecchii, Somaliland, Gestro, p. 786, Ann. Mus. Genov xii; A. lameerei, Gaboon, Duvivier, p. 351, Ann. Ent. Belg. xx: n. spp.
- Anatinomma, n. g., near the Piezocrides, p. 150, for A. alreolatum, n. Mexico, p. 151, pl. v, fig. 8; BATES, Tr. E. Soc. 1892.
- Ancylocera rubella, Mexico, BATES, p. 170, pl. vii, fig. 8, Tr. E. Soc. 1 n. sp.
- Aneflus cylindricollis, p. 147, pl. v, fig. 1, A. ? fulvipennis, fig. 2, p. Mexico, Bates, Tr. E. Soc. 1892, n. spp.
- Anisopodus brevis, Mexico, Gahan, p. 266, pl. xii, fig. 12, Tr. E. Soc. 1 n. sp.
- Anthophilax subcittata, Colorado, Casey, p. 37, Ann. N. York Ac. vi, 1 Anybostetha quedenfeldti, Congo, DUVIVIER, Ann. Ent. Belg. xxxvi, p. n. sp.
- Aphneope quadrimaculata, Queensland, Poll, p. 220, pl. xiii, fi Tijdschr. Ent. xxxiv, n. sp.
- Apilocera breviformis, pl. vi, fig. 15, yucateca, fig. 16, Mexico, B. p. 165, Tr. E. Soc. 1892, n. spp.
- Aposites lanaticollis, S. Australia, p. 61, niger, W. Australia, p. 62, BL BURN, Tr. R. Soc. S. Austr. zv, n. spp.
- Arctolamia fasciata, Burma, GESTRO, p. 222, Ann. Mus. Genov. (2) x, 1
  A ristobia pulcherrima, S. China, Nonfried, p. 94, Ent. Nachr. xviii, 1
  A semum striatum, minute structure of elytra; HOFFBAUER (405).
- A. glubrellum, Mexico, BATES, Tr. E. Soc. 1892, p. 146, pl. v, fig. 6, r Athetesis convergens, Mexico, BATES, p. 171, pl. vii, fig. 10, Tr. E. 1892, n. sp.
- Axestoleus, n. g., near Batyle, p. 180, for Batyle meridionalis, Bates, Axestoleus quinquepunctatus, n. sp., Mexico, pl. vii, fig. 16, p. Bates, Tr. E. Soc. 1892.
- Batyle lævicollis, Mexico, Bates, p. 181, pl. vii, fig. 15, Tr. E. Soc. 1 n. sp.
- Bebius variegatus, Victoria, BLACKBURN, p. 64, Tr. R. Soc. S. Austr n. sp.
- Bethelium tricolor, S. Australia, p. 56, mundum, N. S. Wales, p. 57, Bi. Burn, Tr. R. Soc. S. Austr. xv, n. spp.
- Brachyta bifasciata, n. var. caucasica; Rost, Ent. Nachr. xviii, p. 81. Callichroma splendidum, notes on; Hamilton, p. 38, Canad. Ent. xxi Callimus akbesianus, Syria, Pic, p. 416, Ann. Soc. Ent. Fr. lxi, n. sp. Callipogon: monograph of; Nonfried, pp. 17-24, pl. iii, Berl. Zeit. xxxvii.
- C. hauseri, p. 20, friedluenderi, p. 22, Honduras, Nonfried, Berl. ]
  Zeit. xxxvii, n. spp.

- Cumira: was made from a composite insect, and the name should be deleted; RITSEMA, p. 54, Notes Leyd. Mus. xiv.
- Centrodera nevadica, male noticed; CASEY, p. 36, Ann. N. York Ac. vi.
- Cerambyz: characters of the European species; Belon, pp. 70 & 71, Echange, 1892.
- C. lucasi, Siam, Brongniart, p. 238, pl. x, fig. 4, N. Arch. Mus. Paris (3) iii, n. sp.
- Cerosterna ocellata, Siam, Nonfried, p. 379, Berl. Ent. Zeit. xxxvi, n. sp. Championa badeni, Mexico, Bares, p. 170, Tr. E. Soc. 1892, n. sp.
- Charisia, new name to replace Charis, Newm.; CHAMPION, p. 161, Tr. E. Soc. 1892.
- Charisia nigerrima, Mexico, BATES, p. 160, pl. vi, fig. 8, Tr. E. Soc. 1892,
- Chreostes cinerascens, S. W. Africa, Péringuey, p. 80, Tr. S. African Phil. Soc. vi (2), n. sp.
- Chrotoma, n. g., near Hesperophanes, for C. dunniana n. sp., Texas; Casey, p. 27, Ann. N. York Ac. vi.
- Chrysoprasis guerrerensis, Mexico, BATES, p. 167, pl. vii, fig. 3, Tr. E. Soc. 1892, n. sp.
- Cirrhicera conspicua, pl. xii, fig. 13, basalis, fig. 5, Mexico, GAHAN, p. 269, Tr. E. Soc. 1892, n. spp.
- Closteromerus amabilis, S. W. Africa, Péringuey, p. 78, Tr. S. African Phil. Soc. vi (2), n. sp.
- Clytanthus hololeucus, Mexico, BATES, p. 163, Tr. E. Soc. 1892, n. sp.
- Clutus bifarius, Heyd., = (reitteri, Théry); HEYDEN, p. cxxxi, Bull. Soc. Ent. Fr. lxi. C. ciliciensis, Chev., = (bifarius, Heyd.); Pic, p. cxlvii, Bull. Soc. Ent. Fr. lxi. C. arietis, metamorphoses; XAMBEU, Mœurs, ii, p. 23.
- C. (Sphegestes) reitteri, Syria, mayeti, Siberia, Théry, p. xcvi, Bull. Soc. Ent. Fr. lxi; C. (Xylotrechus) villioni, Japan, VILLARD, p. li, Bull. Soc. Ent. Fr. lxi: n. spp.
- Conisonia leprieuri, Algeria, Pic, p. lii, Bull. Soc. Ent. Fr. lxi, n. sp.
- Coptops hypocrita, Gaboon, LAMBERE, p. 506, Ann. Ent. Belg. xxxvi,
- Cordylomera annulicornis, Obock, FAIRMAIRE, p. 122, Rev. d'Ent. xi, n. sp.
- Cortodera umbripennis, & described; Reitter, p. 239, Wien. ent. Z. xi.
- C. reitteri, Sarepta, Pic, p. 43, Echange, 1891, n. sp.
- Cosmisoma nudicorne, Panama, BATES, p. 166, pl. vii, fig. 2, Tr. E. Soc. 1892, n. sp.
- Criodion fulvopilosum, Brazil, p. 23, antennatum, Venezuela, p. 24, dejeani, Brasil, tuberculatum, Peru, p. 25, testaceum, p. 26, quadrimaculatum, sommeri, p. 27, Brazil, GAHAN, Ann. N. H. (6) ix: n. spp.
- Crioprosopus gaumeri, Mexico, BATES, p. 172, Tr. E. Soc. 1892, n. sp.
- Crossidius longipennis, New Mexico, p. 31, crassipes, Washington, p. 32, nitidicollis, Arizona, p. 33, CASEY, Ann. N. York Ac. vi; C. militaris, p. 177, pl. vii, fig. 14, agrotus, p. 178, Mexico, Bates, Tr. E. Soc. 1892 : n. spp.
  - 1892. [VOL. XXIX.]

- Crossotus sublineatus, Somaliland, Gestro, p. 788, Ann Mus. Geno xii; C. klugii, Transvaal, Distant, Naturalist Transvaal, p. 205 fig. 8: n. spp.
- Cyclocranium, n. g. (Ametrocephalina), p. 219, for C. swierstra, Victoria, pl. xiii, fig. 1, p. 230; Poll., Tijdschr. Ent. xxxiv.
- Cyrtognathus siamensis, Siam, Nonfried, p. 375, Berl. Ent. Zeit.: n. sp.
- Dulterus, n. g., Anauxesides, p. 125, for D. auberti, n. sp., Obock, p. FAIRMAIRE, Rev. d'Ent. xi.
- Daramus, n. g., near Tetropium, for D. serricornis, n. sp., Obock; MAIRE, p. 121, Rev. d'Ent. xi.
- Delagrangeus, n. g., near Deilus, p. xeiii, for D. angustissimus, Syria, p. xeiv; Pic, Bull. Soc. Ent. Fr. lxi.
- Deliathis diluta, Mexico, GAHAN, p. 257, pl. xii, fig. 11, Tr. E. Soc n. sp.
- Deltaspis rufostigma, p. 173, fulva, p. 174, tuberculicollis, p. 175, M Bates, Tr. E. Soc. 1892, n. spp.
- Derobrachus smithi, Mexico, BATES, Tr. E. Soc. 1892, p. 144, n. sp. Dichostates molossus, Congo, DUVIVIER, p. 359, Ann. Ent. Belg. n. sp.
- Dichostethes nebulosus, Obock; FAIRMAIRE, p. 124, Rev. d'Ent. xi. Didymocantha novica, Victoria, BLAUKBURN, p. 233, Tr. R. & Austr. xv, n. sp.
- Distenia lineatopora and hægei, variation noticed; Bates, pp. 156 Tr. E. Soc. 1892.
- D. trifasciata, Mexico, BATES, p. 157, pl. vi, fig. 1, Tr. E. Soc. n. sp.
- Dorcadion equestre, variation described and figured; KRAATZ, ppl. iii, figs. 3-22, Deutsche e. Z. 1892. D. uhagoni, n. var. p. Théry, p. xevi, Bull. Soc. Ent. Fr. lxi.
- D. cervæ, Hungary, FRIVALDSZKY, p. 133, Term. füzetek, xv, n. sp. Eburia baroni, p. 148, pl. v, fig. 3, porulosa, fig. 5, p. 149, Mexico; 1
  Tr. E. Soc. 1892, n. spp.
- Ectosticta ornuta, Victoria, Blackburn, p. 66, Tr. R. Soc. S. Aus n. sp.
- Ecyrus and Œbaceres: notes on the composition of; GAHAN, p. 2 E. Soc. 1892.
- E. arcuatus, Mexico, GAHAN, p. 259, Tr. E. Soc. 1892, n. sp.
- Elaphidion arizonense, p. 28, levettei, p. 29, Arizona, Casey, Ann. N Ac. vi, n. spp.
- Elytroleptus scabricollis, Mexico, BATES, p. 172, pl. vii, fig. 7, Soc. 1892, n. sp.
- Entomosterna proliza, Mexico, BATES, p. 180, Tr. E. Soc. 1892, n. Ergates spiculatus and neomecicanus, distinctions of; CASEY, p. 20 N. York Ac. vi.
- Euclea nodicornis, Amboyna, RITSEMA, p. 38, Notes Leyd. M. n. sp.

- Euderces cribripennis, Mexico, BATES, p. 165, pl. vi, fig. 14, Tr. E. Soc. 1892, n. sp.
- Eupogonius tomentosus, habits; HAMILTON, p. 39, Canad. Ent. xxiv.
- Euryptera unicolor, pl. vi, fig. 3, Mexico, planicoxis, Panama, fig. 5, Bates, p. 159, Tr. E. Soc. 1892, n. spp.
- Exocentrus ruficollis, Congo, LAMEERE, p. 508, Ann. Ent. Belg. xxxvi, n. sp.
- Gasponia, n. g., near Gnathænia, for G. gaurani, n. sp., Obock; FAIR-MAIRE, p. 123, Rev. d'Ent. xi.
- Gaurotes multiguttatus, Mexico, BATES, p. 158, pl. vi, fig. 2, Tr. E. Soc. 1892, n. sp.
- Glenea florensis, p. 221, Flores, oberthuri, p. 222, Java, Ritsema, Notes Leyd. Mus. xiv, n. spp.
- Gnatholea (sub Hesperophanes) liturifer, Wlk., = (denticollis and picicornis), Fairm.; FAIRMAIRE, p. 120, Rev. d'Ent. xi.
- Gnomidolon denticorne, Panama, BATES, p. 155, Tr. E. Soc. 1892, n. sp.
- Gracilia pygmaa, habits; VERHOEFF, p. 298, Ent. Nachr. xviii.
- Grammoptera auricollis, n. var. lucidipes; PIC, p. 290, Abeille, xxvii.
- G. angustata, Algeria, Pic, p. 289, Abeille, xxvii, n. sp.
- Hammaticherus consobrinus, Cayenne, p. 19, lacordairei, Buenos Aires, p. 20, punctulatus, p. 21, lasiocerus, Brazil, murinus, Corrientes, luridipennis, French Guiana, p. 22, GAHAN, Ann. N. H. (6) ix, n. spp.
- Hecyrida lutulenta, Somaliland, Gestro, p. 789, Ann. Mus. Genov. (2)
- Herpetophygas fasciatus, S. Africa, Péringuey, p. 79, Tr. S. African Phil. Soc. vi (2).
- Hexoplon smithi, pl. v, fig. 12, p. 154, sylvarum, fig. 14, p. 155, Mexico, Bates, Tr. E. Soc. 1892, n. spp.
- Hippopsis imitans, Congo, DUVIVIER, p. 364, Ann. Ent. Belg. xxxvi, n. sp.
- Homzophlaus, n. g, near Onychocsrus, for H. licheneus, n. sp., Mexico, pl. xii, fig. 6; Gahan, p. 261, Tr. E. Soc. 1892.
- Hylotrupes litigiosus, N. America, CASEY, p. 25, Ann. N. York Ac. vi, n. sp. Hyperplatys californica, Sta. Cruz, CASEY, p. 51, Ann. N. York Ac. vi, n. sp.
- Hysterarthron belongs to Lagriida; RITSEMA, p. 54, Notes Leyd. Mus. xiv.
- Ibidion ruatanum, Honduras, pl. v, fig. 15, p. 155, griseolum, fig. 13, gaumeri, fig. 16, p. 156, Mexico, Bates, Tr. E. Soc. 1892, n. spp.
- Ipochus subnitidus, pubescens, California, Casey, p. 45, Ann. N. York Ac. vi, n. spp.
- Ischnocnemis cyaneus, BATES, p. 178, Tr. E. Soc. 1892, n. sp.
- Ispaterus, n. g., near Exocentrus, p. 122, for I. longipilis, n. sp., Obock, p. 123; FAIRMAIRE, Rev. d'Ent. xi.
- Lasiogaster, n. g., Monodesmina, p. 255, for L. costipennis, n. sp., British Honduras, p. 256; Gahan, Tr. E. Soc. 1892.

- Leiopus bedeli, Algeria, Pic, p. 289, Abeille, xxvii; L. setipes, min Texas, Casey, p. 48, Ann. N. York Ac. vi; L. batesi, Mexico, G p. 265, pl. xii, fig. 9, Tr. E. Soc. 1892: n. spp.
- Leontium binotatum, Siam, Brongniart, p. 245, pl. x, fig. 9, N. Mus. Paris (3) iii, n. sp.
- Leptostylus arciferus, Mexico, Gahan, p. 265, pl. xii, fig. 8, Tr. 1 1892, n. sp.
- Leptura attenuata, v. brunnescens, Caucasus, Balbi, Riv. Ital. Sc xii, p. 49, n. sp. (sic); L. serpentina, p. 41, haldemani, p. 42, la p. 43, N. America, Casey, Ann. N. York Ac. vi : n. spp.
- Litopus suturalis, S. Africa, Périnquer, p. 131, Tr. S. African Ph vi (2), n. sp.
- Lycidola levipennis, Panama, Gahan, p. 272, pl. xii, fig. 1, Tr. 1892, n. sp.
- Lygesis? ornata, Queensland, BLACKBURN, p. 63, Tr. R. Soc. S. Aun. sp.
- Macrotoma tenuelimbata, Madagascar, Nonfried, p. 139, Ent. Nachi M. alteni, Siam, p. 376, difformis, E. India, p. 377, Nonfried Ent. Zeit. xxxvi: n. spp.
- Malacosrylus humilis, n. varr. grisescens, fulvescens, pl. xii, fig. 16; (p. 271, Tr. E. Soc. 1892.
- M. bivittatus, Mexico, Gahan, Tr. E. Soc. 1892, p. 272, pl. xii, n. sp.
- Mallonia granulato, Transvaal, Distant, p. 407, Ann. N. H. (6) x Maulia, n. g. of doubtful position, ? near Phalota, p. 65, for M cornis, n. sp., Victoria, p. 66; BLACKBURN, Tr. R. Soc. S. Aus
- Mecas marmorata, Mexico, GAHAN, p. 268, pl. xii, fig. 7, Tr. E. Soc
- Mesolita inermis, p. 226, pl. xiii, fig. 8, pascosi, p. 227, fig. 6, Quee Poll, Tijdschr. Ent. xxxiv, n. spp.
- Metaleptus comis, Mexico, BATES, p. 175, pl. vii, fig. 9, Tr. E. Soc n. sp.
- Microtragus quadrimaculatus, Queensland, BLACKBURN, p. 235, Soc. S. Austr. xv, n. sp.
- Monocladum, n. subg. of Polyarthron; Pic, p. celx, Bull. Soc. E
- Monohammus: the Siberian species noticed; Reitter, p. 187, ent. Z. xi.
- M. senerini, S. China, Nonfried, p. 94, Ent. Nachr. zviii; M. J. S. Australia, Blackburn, P. Linn. Soc. N.S.W. (2) vi, p. 5-obtusus, California, Casey, p. 47, Ann. N. York Ac. vi : n. spi
- Moneilema (Collapteryx) spinicollis, Arizona, Casey, p. 45, Ann. N Ac. vi, n. sp.
- Mylothris bimaculatus, Siam, Brongniart, p. 252, N. Arch. Mus. (3) iii, n. sp.
- Mystacophorus recharacterised; Duvivier, p. 368, Ann. Ent.

- Nocydalis major and panzeri, distinctive characters; Chobaut, Échange, No. 61, 1890, p. 100.
- Nemophas malachiticus, Sumatra, Nonfried, Berl. Ent. Zeit. xxxvi, n. sp. Neoclytus smithi, Mexico, Bates, p. 164, pl. vi, fig. 9, Tr. E. Soc. 1892, n. sp.
- Neodorcadion flachneri, n. var., merkli, Pic, p. exevi, Bull. Soc. Ent. Fr. lxi.
- Nitocris gigantea, Zambesi, NONFRIED, p. 140, Ent. Nachr. xviii, n. sp.
- Nyoma, n. g., Acanthocinidarum, p. 366, for N. parallela, n. sp., Congo, p. 367; DUVIVIER, Ann. Ent. Belg. xxxvi.
- Nyphasia maculata, Saïgon, BRONGNIART, p. 240, pl. x, fig. 3, N. Arch. Mus Paris (3) iii, n. sp.
- Oberea schaumii, peculiar larva described and figured; RILEY, p. 323, P. E. Soc. Wash, ii.
- Ochresthes nigritus, pl. vi, fig. 10, clerinus, fig. 13, p. 162, obscuricornis, tulensis, fig. 12, p. 163, Mexico, Bates, Tr. E. Soc. 1892: n. spp.
- Ochyra nana, N. S. Wales, Poll, p. 225, pl. xiii, fig. 7, Tijdschr. Ent. xxxiv, n. sp.
- Odontocera yucateca, Mexico, BATES, p. 159, pl. vi, fig. 7, Tr. E. Soc. 1892, n. sp.
- Olenecamptus macari, Trop. Africa, Lameere, p. 507, Ann. Ent. Belg. xxxvi, n. sp.
- Ophistomis zanthotelus, Mexico, BATES, p. 158, pl. vi, fig. 4, Tr. E. Soc. 1892, n. sp.
- Opsidota guttata, S. Australia, BLACKBURN, Tr. R. Soc. S. Austr. xv, n. sp.
- Oreodera affinis, Guatemala, GAHAN, p. 262, pl. xii, fig. 14, Tr. E. Soc. 1892, n. sp.
- Pachylocerus sulcatus, Siam, BRONGNIART, p. 240, pl. x, fig. 4, N. Arch. Mus. Paris (3) iii, n. sp.
- Pachymerola, n. g., near Coremia for P. vitticollis, n. sp., Mexico, pl. vi, fig. 11; Bates, Tr. E. Soc. 1892, p. 161.
- Pachyteria pasteuri, p. 213, lugubris, p. 215, Nias Is., borneoensis, Sarawak, p. 218; RITSEMA, Notes Leyd. Mus. xiv.
- Paroeme gahani, Transvaal, DISTANT, Naturalist Transvaal, p. 202, pl. i, fig. 7, n. sp.
- Paryeatis nigritareis, Thoms., = (flavescens, Bates); GAHAN, p. 258, Tr. E. Soc. 1892.
- P. perplexa, Mexico, GAHAN, p. 258, Tr. E. Soc. 1892, n. sp.
- Pavieia, n. g., near Crioprosopus, p. 241, for P. superba, n. sp., Siam, p. 243, pl. x, fig. 6; Brongniart, N. Arch. Mus. Paris. (3) iii.
- Phasa unicolor, Mexico, GAHAN, p. 267, Tr. E. Soc. 1892, n. sp.
- Phryneta supera, Limpopo Riv., Peringuey, p. 133, Tr. S. African Phil. Soc. vi (2), n. sp.
- Phyllocnema speciosa, S. W. Africa, Péringuey, p. 78, Tr. S. African Phil. Soc. vi (2); P. pretiosa, Limpopo Riv., Péringuey, p. 130, t. c.: n. spp.

- Phymatodes obliques, California, CASEY, p. 26, Ann. N. York Ac. vi, Phytacia (Coptosia) trilinea; Pic, p. cxii, Bull. Soc. Ent. Fr. lxi; F. Araxes Valley, Reitter, p. 64, Wien. ent. Z. xi, n. spp.
- Plocederus incertus, Somaliland, GESTRO, p. 784, Ann. Mus. Geno xii, n. sp.
- Pacilomallus, n. g., Elaphidiina, for P. palpalis, n. sp., Mexico, fig. 6; Bates, p. 151, Tr. E. Soc. 1892.
- Polyarthron unipectinatum, White, sexes described and figured; HE p. 169, pl. iii, figs. 1 & 2, Deutsche e. Z. 1892.
- P. desvauxi, p. celix, moissoni, p. celx, Algeria, Pic, Bull. Soc. Ent. F n. spp.
- Praonetha lichenea, Congo, DUVIVIER, p. 363, Ann. Ent. Belg. xxxvi, Prionus ruficornis, nadari, Turkestan, FAIRMAIRE, p. cxxiii, Bull Ent. Fr. lxi; P. debilis, N. America, Casey, p. 21, Ann. N. Ac. vi: n. spp.
- Promeces crassicorais, Somaliland, GESTRO, p. 785, Ann. Mus. 6
  (2) xii, n. sp.
- Proteinidium, n. g., near Azestinus, p. 149, for P. brevicorne, 1 Mexico, p. 150, pl. v, fig. 4; Bates, Tr. E. Soc. 1892.
- Psenocerus supernotatus and tristis, variation of; Hamilton, p Canad. Ent. xxiv.
- P. tristis, New Mexico, Casey, p. 46, Ann. N. York Ac. vi, n. sp.
- Psyrassa sthenias, punctulata, cribellata, pl. v, fig. 9, p. 152, pil fig. 11, nigricornis, fig. 10, nigroænea, p. 153, Mexico, Bates, Soc. 1892, n. spp.
- Purpuricenus fasciatus, Siam, BRONGNIART, p. 241, pl. x, fig. 5, N. Mus. Paris (3) iii, n. sp.
- Rhopalophora longipes and messkei, note on; Hamilton, p. 159, ( Ent. xxiv.
- R. meeskei, New Mexico, Casey, p. 30, Ann. N. York Ac. vi; R. e Mexico, Bates, p. 166, pl. vii, fig. 1, Tr. E. Soc. 1892: n. spp.
- Rhytidodera siamica, Siam, Nonfried, p. 378, Berl. Ent. Zeit.: n. sp.
- Rosalia lameerei, Siam, Brongniart, p. 243, pl. x, figs. 7 & 8, N. Mus. Paris (3) iii.
- Saperda phoca, habits; Delagrange, Bull. Soc. Ent. Fr. lxi, p. ec. Sisyrium: note on characters, p. 58, and table of species, p. 61; B Burn, Tr. R. Soc. S. Austr. xv.
- S. fraternum, p. 58, ventrale, p. 59, sparsum, S. Australia, lævigatu:
  Australia, p. 60, Blackburn, Tr. R. Soc. S. Austr. xv, n. spp.
- Spalacopsis texana, Texas, Casey, p. 51, Ann. N. York Ac. vi; S. s. fusca, Mexico, Gahan, p. 260, Tr. E. Soc. 1892: n. spp.
- Sphallenum literatum, p. 28, spadiceum, p. 29, Brazil, Gahan, Ann. (6) ix, n. spp.
- Sphenothecus rubens, Arizona, Casey, p. 34, Ann. N. York Ac., quadrivittatus, cribricollis, cribellatus, Mexico, Bates, p. 179, Soc. 1892 : n. spp.

- Spondylis sinensis, S. China, Nonfried, p. 92, Ent. Nachr. xviii, n. sp. Stenaspis pilosella, Mexico, Bates, p. 173, pl. vii, fig. 11, Tr. E. Soc. 1892, n. sp.
- Stenopterus kraatzi, Smyrna, Pic, p. 22, Échange, 1892, n. sp.
- Stenosphenus longicollis, Texas, Casey, p. 34, Ann. N. York Ac. vi; S. sublævicollis, p. 168, sexlineatus, pl. vii, fig. 5, gaumeri, comus, vitticollis, fig. 6, p. 169, Mexico, Bates, Tr. E. Soc. 1892: n. spp.
- Sternoplistes schaiblei, S. China, Nonfried, p. 92, Eut. Nachr. xviii, n. sp. Strangulia: sexual distinctions; Casey, p. 39, Ann. N. York Ac. vi.
- S. montana, New Mexico, Casey, Ann. N. York Ac. vi, p. 40, n. sp.
- Stratioceros princeps, figured and noticed; BRONGNIART, p. 248, pl. x, fig. 10. N. Arch. Mus. Paris (3) iii.
- Strongylaspis lobulifer, Mexico, BATES, p. 145, Tr. E. Soc. 1892, n. sp. Sympleurotis armatus, Mexico, GAHAN, p. 266, pl. xii, fig. 10, Tr. E. Soc. 1892, n. sp.
- Taurotagus auberti, Obock, FAIRMAIRE, p. 118, Rev. d'Ent. xi, n. sp. Teispes frenchi, Queensland, BLACKBURN, p. 234, Tr. R. Soc. S. Austr. xv, n. sp.
- Tetropium luridum, minute structure of elytra; HOFFBAUER (405).
- T. parallelum, p. 23, schwarzianum, parvulum, p. 24, N. America, CASEY, Ann. N. York Ac. vi; T. guatemalanum, Tepan, BATES, p. 147, Tr. E. Soc. 1892: n. spp.
- Thesalia, n. g., for Acmaops lisa, Leng; CASEY, p. 36, Ann. N. York Ac. vi.
- Tibetobia, n. g., near Monochammus, p. 118, for T. szechenyiana, n. sp., Tibet; FRIVALDSZKY, Term. füzetek, xv.
- Tilloclytus conradti, Guatemala, BATES, p. 164, Tr. E. Soc. 1892, n. sp.
- Toxotus lateralis, California, Casey, p. 37, Ann. N. York Ac. vi, n. sp. Trachustola armata. S. China. Nonfried. p. 93. Ent. Nachr. xvi
- Trachystola armata, S. China, Nonfried, p. 93, Ent. Nachr. xviii, n. sp.
- Tragidion fulvipenne, oviposition; POPENOE, Tr. Kansas Ac. xii, p. 15.
- Tragocephala carbonaria, Trop. Africa, LAMEERE, p. 507, Ann. Ent. Belg. xxxvi; T. timida, Congo, DUVIVIER, p. 167, Ann. Ent. Belg. xxxvi; T. sulphurata, Transvaal, DISTANT, Naturalist Transvaal, p. 202, pl. i, fig. 9: n. spp.
- Tragosoma depsarium and harrisii are one species; Hamilton, p. 296, Canad. Ent. xxiv.
- T. nigripenne, Mexico, BATES, p. 146, Tr. E. Soc. 1892, n. sp.
- Triacetelus, n. g., near Metaleptus, for T. sericatus, n. sp., Mexico; Bates, p. 176, Tr. E. Soc. 1892.
- Trichoxys cinereolus, Mexico, BATES, p. 162, Tr. E. Soc. 1892, n. sp.
- Trichromia, n. g., near Anthribola, for T. oberthuri, n. sp., Madagascar; FAIRMAIRE, Bull. Soc. Ent. Fr. lxi, p. xxi.
- Tritomicrus, n. g., near Hoplosia for T. marmoreus, n. sp., Obock; FAIRMAIRE, Rev. d'Ent. xi, p. 125.
- Tryphocharia mitchelli, note on, BLACKBURN, p. 56, Tr. R. Soc. S. Austr. xv; T. frenchi, N. S. Wales, BLACKBURN, p. 233, t. c.: n. spp.

- Tylosis puncticollis, var., = (angusticollis, Bates); BATES, p. 17 E. Soc. 1892.
- T. dimidiata, Mexico, BATES, Tr. E. Soc. 1892, pl. vii, fig. 13, n. sp. Typhocesis floccosa, Poll, figured; Poll, pl. xiii, fig. 10, Tijdschr. Ent. Uracanthus cryptophagus, Australia, Olliff, p. 896, pl. lvii, Agrin N.S.W. iii, n. sp.
- Valenus, n. g., neur Lepturges, p. 49, for V. inornatus, n. sp., Texas, Casey, Ann. N. York Ac. vi.
- Volumnia transversalis, Trop. Africa, LAMEERE, p. 508, Ann. Ent. xxxvi, n. sp.
- Xestia denticornis, p. 29, longipennis, globulicollis, p. 31, Brazil, G Ann. N. H. (6) ix: n. spp.
- Xoanodera pascoei, Siam, BRONGNIART, p. 239, N. Arch. Mus. Pa iii, n. sp.
- Xylotrechus insignis, sexual distinctions, Casey, p. 35, Ann. N. Yo vi, n. sp.
- Xystrocera parvicollis, Obock, FAIRMAIRE, p. 120, Rev. d'Ent. x læta, S. W. Africa, Péringuey, p. 77, Tr. S. African Phil. S (2): n. spp.
- Zalates pulchra, Transvaal, Péringuey, p. 132, Tr. S. African Phi vi (2), n. sp.
- Zamium rusticum, S. W. Africa, Péringuey, p. 77, Tr. S. African Soc. vi (2), n. sp.
- Zatrephus nebulosus, Saïgon, BRONGNIART, p. 240, pl. x, fig. 2, N. Mus. Paris (3) iii, n. sp.
- Zenochloris barbicauda, Mexico, BATES, p. 168, pl. vii, fig. 4, Tr. F 1892, n. sp.
- Zočdia v-album, figured, pl. xiii, fig. 3; Z. dirisa, Pasc., = (el Waterh.), p. 222, Poll, Tidjschr. Ent. xxxiv.
- Z. longipes, p. 222, p. 13, fig. 4, gracilipes, p. 223, fig. 5, Australia, p. 224, fig. 6, Poll, Tijdschr. Ent. xxxiv, n. spp.

## SCOLYTIDÆ.

- [Cf. Blandford (100), Desbrochers (199), Knotek (500), XA (1022).]
- Hylesinus henscheli, Herzegovina, KNOTEK, p. 234, Wien. ent. Z. zi, Hypoborus ficus, metamorphoses; XAMBEU, Soc. Agricole Pyrén.-c xxxii, p. 473.
- Phlæotribus oles, metamorphoses; XAMBEU, Soc. Agricole Pyrén.-c xxxii, pp. 477-484.
- Pityophthorus annectens and consimilis, note on; Schwarz, p. P. E. Soc. Wash. ii.
- Scolytus: table of characters of the European; Desbrochers, Frei pp. 10-17. S. carpini, habits, Decaux, p. cexli, Bull. Soc. Ent. F S. aceris, Herzegovina, Knotek, p. 235, pl. ii, Wien. ent. Z. xi, n. sp Xyleborus perforans, in the West Indies; Blandford (100).

### BRENTHIDE.

- [Cf. Desbrochers (203), Kolbe (506), Semenow (831), Senna (837, 838, 839).]
- List of additions to, and notes on the Catalogue of, Brenthidæ; Senna, pp. 38-63.
- Observations on the dimorphic form of male rostrum; Senna, pp 30-37, Bull. Ent. Ital. xxiv.
- Agriorrhynchus quadrituberculatus, Burma, SENNA, p. 470, Ann. Mus. Genov. (2) xii, n. sp.
- Autarcus, n. g., for Ceocophalus laticollis, Perroud; SENNA, p. 59, Bull. Ent. Ital. xxiv.
- Basenius, n. g., for B. laticornis, E. Africa, n. sp.; Kolbe, p. 170, S. E. Z. liii.
- Callipareius, n. g., near Cyphagogus, p. 444, for C. fea, n. sp., Burma, p. 445; Senna, Ann. Mus. Genov. (2) xii.
- Carcinopisthius, n. g., for C. fruhstorferi, Java, p. 174, felschei, Burma, p. 175, n. spp.; Kolbe, S. E. Z. liii.
- Cediocera tristis, Java, SENNA, p. 183, Notes Leyd. Mus. xiv, n. sp.
- Cerobates seesulcatus, variation, pp. 449, 450, canaliculatus, n. var., carinensis, p. 450; Senna, Ann. Mus. Genov. (2) xii.
- Coptorhynchus francoisi, New Hebrides, Desbrochers, p. 109, Frelon, i, n. sp.
- Cordus peguanus, Burma, Senna, p. 463, Ann. Mus. Genov. (2) xii, n. sp. Cormopus, n. g., p. 166, for C. penicillifer, n. sp., Cameroons, p. 167; Kolbe, S. E. Z. liii.
- Diurus ominosus, Burma, Senna, p. 492, Ann. Mus. Genov. (2) xii, n. sp. Epicoinoneus, n. g., Belophorinæ, p. 476, for E. femoralis, n. sp., Burma, p. 478; Senna, Ann. Mus. Genov. (2) xii.
- Eremozenus, n. g., p. 438, for E. chan, n. sp., Transcaspian Region, p. 440, SÉMENOW, Hor. Ent. Ross, xxvi.
- Hopliterrhynchus, n. g., near Ectocemus, p. 26, for H. emmæ, n. sp., Malasia, p. 27; SENNA, Bull. Ent. Ital. xxiv.
- Hoplopisthius, n. g., Trachelisinæ, p. 451, for H. trichemerus, n. sp., Burma, p. 452; Senna, Ann. Mus. Genov. (2) xii.
- H. javanus, Java, p. 173, celebensis, Celebes, p. 174, Kolbe, S. E. Z. liii, n. spp.
- Hyperephanus, n. g., near Jonthocerus, p. 153, for E. (sic) hireutus, n. sp., Chili, p. 155; SENNA, Bull. Ent. Ital. xxiv.
- Ithystenus, = (Coptorhynchus, Desb.); SENNA, p. 152, Bull. Ent. Ital. xxiv.
- Microsebus, n. g., p. 168, for M. pusio, E. Africa, adelphus, Ceylon, n. spp., p. 169; Kolbe, S. E. Z. liii.
- Miolispa javanica, p. 167, metallica, p. 169, nupta, p. 171, conformis, p. 175, Java, Senna, Notes Leyd. Mus. xiv, n. spp.
- Prophthalmus versicolor, Burma, Senna, p. 466, Ann. Mus. Genov. (2) xii, n. sp.

- Oncodemerus, n. g., Taphroderini, p. 157, for O. costipennis, n. sp., Gap. 160; Senna, Bull. Ent. Ital. xxiv.
- Orychodes degener, Burma, Senna, p. 474, Ann. Mus. Genov. (2)
  O. piliferus, Java, Senna, Notes Leyd. Mus. xiv, p. 177: n. spj
- Schisotrachelus intermedius, p. 481, carinensis, p. 483, consanguineus manicus, p. 485, sculptiventris, p. 488, carinirostris, p. 489, B. Senna, Ann. Mus. Genov. (2) xii, n. spp.
- Sebasius: note on the components of; Kolbe, p. 171, S. E. Z. liii.
- Trachelizus brevitibia, p. 454, bhamoensis, p. 456, ghecuanus, p. rufomaculatus, p. 460, politus, p. 462, Burma, Senna, Ann. Genov. (2) xii; T. insularis, p. 163, modestus, mæstus, p. 165, Senna, Notes Leyd. Mus. xiv: n. spp.
- Usambius, n. g., p. 167, for U. conrudti, n. sp., E. Africa, p. 168; K. S. E. Z. liii.
- Zemioses pubens, p. 439, lostus, p. 441, Burma, Senna, Ann. Mus. G (2) xii, n. spp.

#### ANTHRIBIDA.

- [Cf. DUVIVIER (230), FAIRMAIRE (269), LESNE (547).]
- Cenchromorphus, n. g., Corrhécérides, for C. burbicornis, n. sp., Madag FAIRMAIRE, p. clxx, Bull. Soc. Ent. Fr. lxi, n. sp.
- Eugigas harmandi, Cochin China, LESNE, p. 227, N. Arch. Mus. 1 (3) iii, n. sp.
- Litocerus pariei, Siam, LESNE, p. 226, N. Arch. Mus. Paris (3) iii, n. Phlæotragus prasinus, Congo, DUVIVIER, p. 59, Ann. Ent. Belg. x n. sp.
- Vitalis, n. g., near Platyrhinus, for V. rubricollis, n. sp., Madaga FAIRMAIRE, Bull. Soc. Ent. Fr. lxi, p. clxxi.

## BRUCHIDE.

- [Cf. Heyden (402), RILEY (751, 752, 754).]
- Bruchus: temporary feet in early larves; RILEY, pp. 185 & 186, C
  Ent. xxiv. B. fabæ and pisi, metamorphoses; RILEY & Hov
  Ins. Life, iv, pp. 297-302. B. obsoletus, Say, note on; Hami
  p. 162, Canad. Ent. xxiv. B. obtectus, Say, nomenclature
  habits; RILEY, Ins. Life, v, pp. 27-33, and note on, with synon
  RILEY, p. 291, Canad. Ent. xxiv. B. alboscutellatus, habits; H.
  Ton, Ent. News, iii, p. 253.
- B. ruthenicus, Sarepta, BECKER, p. 69, Bull. Mosc. 1892, n. sp. ?

  Mylabris (Bruchus) orchesioides, Margelan, HEYDEN, p. 109, Deutsc
  Z. 1892, n. sp.

# CHRYSOMELIDÆ.

[Cf. Balbi (31), Blackburn (90, 93, 94), Chittenden (168), Dbi chers (204), Duvivier (230, 232), Everts (260), Fairmaire (Frivaldszky (309), Gahan (210), Gestro (325, 331, 335), Goi

- (350), GORHAM & GAHAN (353), GUILLEBEAU (375), HORN (416, 418, 419), JACOBSOHN (444, 445), JACOBY (210, 347, 446, 447, 448), LEFÈVRE (543, 544), PÉRINGUEY (672, 673), REITTER (725), SCHWARZ (806), WEISE (992, 994, 996, 997, 998, 999).]
- Food-habits of some N. American Chrysomelidæ; CHITTENDEN (168).
- Food-plants of various N. American *Halticids*; Schwarz, pp. 182-184, P. E. Soc. Wash. ii.
- Halticidæ: protest against generic names used by SEIDLITZ in Fauna transsylvanica; Weise, p. 414, Deutsche e. Z. 1892.
- Abirus antennatus, Burma, JACOBY, p. 912, Ann. Mus. Genov. (2) xii, n. sp.
- Achænops facialis, Transvaal, JACOBY in DISTANT, Naturalist Transvaal, p. 204, pl. i, fig. 4, n. sp.
- Acrocrypta discoidalis, Annam, ALLARD, p. 233, N. Arch. Mus. Paris (3) iii, n. sp.
- Ælianus, n. g., near Malacosoma, for Æ. scutellatus, n. sp., Madagascar, pl. xxxix, fig. 9; JACOBY, p. 577, P. Z. S. 1892.
- Enidea? pretoriæ, Transvaal, GAHAN in DISTANT, Naturalist Transvaal, p. 209, pl. i, fig. 10; Æ. crassipalpis, Burma, JACOBY, p. 986, Ann. Mus. Genov. (2) xii: n. spp.
- Æthonea, Baly, is same as Bonesia; GAHAN, p. 92, P. Z. S. 1892.
- Agelastica? flava, Burma, JACOBY, p. 967, Ann. Mus. Genov. (2) xii;
  A. viridis, Mexico, JACOBY, p. 334, Biol. Centr. Am. Col. vi
  (1 supp.): n. spp.
- Allochroma teapense, pl. xliii, fig. 2, flohri, fig. 5, Mexico, Jacoby, p. 319, Biol. Centr. Am. Col. vi (1 supp.), n. spp.
- Allomorpha, n. g., Halticidæ, for A. sericea, u. sp., Burma; Jacoby, Ann. Mus. Genov. (2) xii.
- Amphimela cyanea, Bengal, DUVIVIER, p. 420, Ann. Ent. Belg. xxxvi, n. sp.
- Anisodera nasuellii, Burma, Gestro, p. 233, Ann. Mus. Genov. (2) x, n. sp.
- Anthipha pubescens, p. 971, inornata, p. 972, antennata, fee, p. 973, lavicollis, p. 974, JACOBY, Ann. Mus. Genov. (2) xii; A. blanchardi, Indo-China, Allard, p. 234, N. Arch. Mus. Paris (3) iii: n. spp.
- Anthiphula, n. g., between Galerucina and Halticina, provisionally placed near Anthipha, for A. semifulva, n. sp., Burma; JACOBY, p. 991, Ann. Mus. Genov. (2) xii.
- Antsianaka viridis, elegantula, pl. xxxix, fig. 10, Madagascar, JACOBY, p. 576, P. Z. S. 1892, n. spp.
- Aoria nigrita, p. 904, fulvifrons, p. 905, semicostata, p. 906, Burma, Jacoby, Ann. Mus. Genov. (2) xii : n. spp.
- Aphthona atrovirens, var. described; Weise, Deutsche e. Z. 1892, p. 411.
- A. nigrilabris, Bengal, DUVIVIER, p. 426, Ann. Ent. Belg. xxxvi;
  A. birmanica, Pegu, JACOBY, p. 920, Ann. Mus. Genov. (2) xii:
  n. spp.
- Arcastes sanguinea, Perak, JACOBY, p. 88, Ent. xxv, Supp., n. sp.

- Argopistes bistripunctata, Bengal, DUVIVIER, p. 425, Ann. Ent. B xxxvi; A. nigromarginatus, Burma, JACOBY, p. 931, Ann. I Genov. (2) xii: n. spp.
- Argopistoides, n. g., p. 931, for A. septempunctata, n. sp., Burma, p. 9

  JACOBY, Ann. Mus. Genov. (2) xii.
- Arispoda pulcher, N. Guinea, TRYON (953), n. sp.
- Asphæra madagascariensis, brevicornis, Madagascar, Jacoby, p. P. Z. S. 1892, n. spp.
- Aspidispa albertisii, p. 265, nigritarsis, p. 266, New Guinea, Gestro, A Mus. Genov. (2) x, n. spp.
- Aspidolopha thoracica, p. 879, costata, p. 881, carulea, p. 882, Bur Jacoby, Ann. Mus. Genov. (2) xii, n. spp.
- Aspidomorpha aruwimiensis, Centr. Africa, GAHAN & GORHAM, p. P. Z. S. 1892, n. sp.
- Atyea? albofasciata, Burma, JACOBY, p. 977, Ann. Mus. Genov. (2) n. sp.
- Aulacophora, auct., divided into genera (two presumably new, cf. Papalpa, Orthaulaca); Weise, p. 392, Deutsche e. Z. 1892.
- A. melanocephala, p. 941, intermedia, semifusca, p. 942, gestroi, p. bhamoensis, p. 944, Burma, Jacoby, Ann. Mus. Genov. (2) n. spp.
- Aulacoscelis purpurea, Arizona, HORN, p. 46, Tr. Am. Ent. Soc. xix, n Blepharida singularis, Mexico, JACOBY, p. 347, pl. xliii, fig. 23, Biol. Ce Am. Col. vi (1 supp.), n. sp.
- Bonesia serricornis, var. described; GAHAN, p. 92, P. Z. S. 1892.
- Bufonida pallida, p. 159, piceo-limbata, p. 960, Burma, Jacoby, A. Mus. Genov. (2) xii; B. rubroviridis, intermedia, New Gui Tryon (953): n. spp.
- Cacoscelis caruleipennis, Mexico, Jacoby, p. 346, Biol. Centr. Am. vi (1 supp.), n. sp.
- Callispa arcana, Bengal, DUVIVIER, p. 446, Ann. Ent. Belg. xxxvi, n. Candezea longicornis, marginipennis, p. 979, apicalis, p. 980, orn p. 981, Burma, JACOBY, Ann. Mus. Genov. (2) xii, n. spp.
- Cassena, n. g., near Nisotra, for C. celebensis, p. 389, ribbei, p. 1. n. spp., Celebes, Weise, Deutsche e. Z. 1892.
- Cassida: monograph of the French species; DESEROCHERS, Frelor Supp., & ii, pp. 82-87.
- C. thoracica, p. 19, sanguinosa, p. 20, France, corrosa, oranensis, Alge p. 39, cerata, Syria, p. 40, salsolæ, Russia, p. 43, nitidula, Alge p. 44, Desbrochers, Frelon, i, Supp.; C. reitteri, Caucasus, We p. 238, Wien. ent. Z. xi; C. bonnyana, p. 94, strigosa, fuscopunct p. 95, Centr. Africa, Gorham, P. Z. S. 1892; C. (Odontionycha) i: cola, Bengal, Duvivier, p. 448, Ann. Ent. Belg. xxxvi: n. spp.
- Cassidula turcmenica, Turcomania, Weise, p. 238, Wien. ent. Z. xi, n. Cerania, subg. (n.?) of Orthaulaca; Weise, p. 396, Deuteche e. Z. 189 Ceratia cattigarensis, Shanghai, p. 397, tricolora, Sumatra, fune Celebea, p. 398, Weise, Deuteche e. Z. 1892, n. spp.

- Cerophysa fulricollis, p. 947, monstrosa, p. 948, Burma, Jacoby, Ann. Mus. Genov. (2) xii : n. spp.
- Cerotrus nigromarginatus, Burma, JACOBY, p. 940, Ann. Mus. Genov. (2) Xii, n. sp.
- Chatocnema nagpurensis, E. India, DUVIVIER, p. 422, Ann. Ent. Belg. xxxvi; C. birmanica, duvivieri, Burma, JACORY, p. 939, Ann. Mus. Genov. (2) xii; C. smithi, fulvilabris, interstitialis, p. 313, horni, balyi, costatipennis, p. 314, Mexico, JACOBY, Biol. Centr. Am. Col. vi (1 supp.): n. spp.
- Charaa? alboplagiata, Burma, JACOBY, Ann. Mus. Genov. (2) xii, p. 975, n. sp.
- Chlamys fea, Burma, Jacoby, p. 887, Ann. Mus. Genov. (2) xii, n. sp.
- Charidiona fea, Burma, Gestro, p. 239, Ann. Mus. Genov. (2) x, n. sp.
- Chrysochus pretiosus, metamorphoses; XAMBEU, Le Nat. 1892, p. 117.
- Chrysolampra varicolor, p. 893, minuta, p. 894, Burma, Jacoby, Ann. Mus. Genov. (2) xii : n. spp.
- Chryomela: minute structure of elytra; HOFFBAUER (405). C. cærulans, n. var. olivaceonigra; FLEISCHER, p. 141, Wien. ent. Z. xi; C. songorica, cedeagus of; Weise, p. 139, Deutsche e. Z. 1892.
- C. abchasica, p. 405, rosti, fuscicornis, p. 406, Caucasus, Weise, Deutsche
  e. Z. 1892; C. sogdiorum, p. 138, cyrtonoides, p. 139 (this altered to cyrtonastes, p. 413, t. c.), Turkestan, Weise, t. c.; C. corrugata,
  S. W. Africa, Péringuey, p. 87, Tr. S. African Phil. Soc. vi (2): n. spp.
- Cladocera spectabilis, p. 87, simplex, p. 88, S. W. Africa, Péringury, Tr. S. African Phil. Soc. vi (2): n. spp.
- Cleoporus lefevrei, Bengal, DUVIVIER, p. 414, Ann. Ent. Belg. xxxvi; C. birmanicus, p. 914, plugiatus, p. 915, Burma, JACOBY, Ann. Mus. Genov. (2) xii: n. spp.
- Cleorina jacobyi, Bengal, DUVIVIER, p. 415, Ann. Ent. Belg. xxxvi, n. sp. Cneorane braeti, Bengal, DUVIVIER, p. 435, Ann. Ent. Belg. xxxvi; C. subænea, p. 945, orientalis, p. 946, feæ, p. 947, Burma, JACOBY, Ann. Mus. Genov. (2) xii: n. spp.
- Canobius birmanicus, Burma, JACOBY, p. 889, Ann. Mus. Genov. (2) xii, n. sp.
- Colaspoides sublavicollis, p. 417, bengalensis, p. 418, E. India, DUVIVIER, Ann. Ent. Belg. xxxvi; C. violaceipennis, opacicollis, p. 229, viridimicans, p. 230, N. America, Horn, Tr. Am. Ent. Soc. xix: n. spp.
- Colasposoma aruwimiense, Centr. Africa, Gahan, p. 91, P. Z. S. 1892; C. parculum, Transvaal, Lefèvre, p. 42, Tr. S. African Phil. Soc. vi (2): n. spp.
- Copa, subg. (n.?) of Orthaulaca, with O. (C.) kunowi, n. sp.?, E. Africa, Weise, Deutsche e. Z. 1892, pp. 397, 398.
- Coptocyrla catenata, Boh., note on; Weise, p. 352, Deutsche e. Z. 1892. Coraia apicicornis, Mexico, Jacoby, p. 324, pl. xliii, fig. 6, Biol. Centr. Am. Col. vi (1 supp.), n. sp.

- Corynodes bonnyi, Centr. Africa, Gahan, p. 91, P. Z. S. 1892; C. insig Zambezi, Lepèvre, p. 43, Tr. S. African Phil. Soc. vi; C. dilatice p. 916, birmanicus, p. 917, Burma, Jacoby, Ann. Mus. Genov. xii: n. spp.
- Coscinoptera seminuda, Arizona, HORN, p. 13, Tr. Am. Ent. Soc. xix, n Crepidodera orientalis, obscurofasciata, Burma, JACOBY, p. 933, Ann. I Genov. (2) xii, n. spp.
- Crioceris koltsei, Turkestan, Weise, p. 131, Deutsche e. Z. 1892; C. giosa, S. W. Africa, Péringuey, Tr. S. African Phil. Soc. vi p. 82; C. humeralis, Burma, Jacoby, p. 874, Ann. Mus. Genov. xii; C. aterrima, Mexico, Jacoby, p. 342, Biol. Centr. Am. Co (1 supp.): n. spp.
- Cryptocephalus canguinolentus, Scop., invalidity of the species; W1 p. 414, Deutsche e. Z. 1892.
- C. duvivieri, p. 134, placidus, p. 135, personatus, p. 136, negligens, p. Turkestan, Weise, Deutsche e. Z. 1892; C. scutellatus, doi pl. xxxix, fig. 1, Madagascar, Jacoby, p. 566, P. Z. S. 1892 emphractus, p. 84, strenuus, p. 85, S. Africa, Péringury, Tr African Phil. Soc. vi (2); C. nagpurensis, p. 400, suavis, p. laterimaculatus, p. 403, fraternus, p. 404, Bengal, Duvivier, I Ent. Belg. xxxvi; C. nigriceps, semimarginatus, Indo-China, All. p. 232, N. Arch. Mus. Paris (3) iii; C. angulatofasciatus, gei p. 890, fem, p. 891, flavicinctus, p. 892, Burma, Jacoby, Ann. I Genov. (2) xii: n. spp.
- Cynorta violacea, p. 970, subænea, p. 971, Burma, JACOBY, Ann. ] Genov. (2) xii, n. spp.
- Cyphotarsis, n. g., near Scelida, for C. niger, n. sp., Mexico, JACOBY, p. Biol. Centr. Am. Col. vi (1 supp.).
- Demotina semifusciata, Burma, JACOBY, p. 906, Ann. Mus. Genov. (2) n. sp.
- Derectis, Clark, = (Antipha, Baly); JACOBY, Ent. xxv, p. 162.
- Derocrepis pubipennis, Araxes Valley, REITTER, p. 154, Wien. ent zi, n. sp.
- Diabrotica smithi, peregrina, pl. xliii, fig. 12, p. 330, guerreroensis, fig blattoides, p. 331, quatuor-decimpunctata, fig. 14, semiopaca, ri p. 332, D.? antennata, fig. 15, p. 333, Centr. America, Jacoby, Centr. Am. Col. vi (1 supp.), n. spp.
- Diacantha lugubris, opulenta, S.W. Africa, Péringuey, Tr. S. African ]
  Soc. vi (2); D. balteata, Transvaal, Péringuey, p. 133, t. c. : n.
- Diapromorpha curulea, p. 877, melanocephala, gigantea, p. 878, Bu Jacoby, Ann. Mus. Genov. (2) xii, n. spp.
- Dioryctus castaneus, Burma, JACOBY, p. 888, Ann. Mus. Genov. (2) n. sp.
- Diphaulacosoma, n. g., Halticinæ, for D. lævipenne, n. sp., Madaga: Jacoby, p. 574, pl. xxxix, fig. 8, P. Z. S. 1892.
- Donaciini: tabulation of the Netherlands species; Everts, Tijds Ent. xxxv, pp. 31-58.

- Donacia and Plateumaris: characters of the groups and species of the Old World; Jacobson, pp. 412-437, Hor. Ent. Ross. xxvi. D. sericea: abnormal coloration in; Paulcke, Deutsche e. Z. 1892, p. 416.
- D. australasia, Queensland, BLACKBURN, p. 235, Tr. R. Soc. S. Austr. xv, n. sp.
- Doryida and D. mouhoti, redefined; JACOBY, pp. 989 & 990, Ann. Mus. Genov. (2) xii.
- Dononesia grandis, p. 241, elegans, p. 242, longipennis, p. 243, Burma, Gestro, Ann. Mus. Genov. (2) x, n. spp.
- Echtrusia, n. g., near Eurysthenes, for E. rhomboidea, n. sp., Cape Town; Lepèvre, p. 41, Tr. S. African Phil. Soc. vi.
- Enneamera birmanica, limbatipennis, Burma, JACOBY, p. 935, Ann. Mus. Genov. (2) xii, n. spp.
- Erystus podagroides, Ceram, p. 390, villicus, Bangkei, p. 391, Weise, Deutsche e. Z. 1892, n. spp.
- Eulychius nigritarsis, Madagascar, JACOBY, p. 567, P. Z. S. 1892, n. sp.
- Eumelepta, n. g., near Ergana, for E. biplagiata, n. sp., Burma, JACOBY, p. 988, Ann. Mus. Genov. (2) xii.
- Eumolpini, of N. America, revised and tabulated; HORN, Tr. Am. Ent. Soc. xix, pp. 195-234.
- Eurispa loria, New Guinea, GESTRO, p. 1016, Ann. Mus. Genov. (2) x;
  E. nigripes, S. Australia, simplex, Victoria, fraterna, S. Australia,
  BLACKBURN, p. 68, Tr. R. Soc. S. Austr. xv, n. spp.
- Eurydemus metallicus, Madagascar, JACOBY, p. 570, P. Z. S. 1892; E. depressus, Zambezi, LEFÈVRE, p. 42, Tr. S. African Phil. Soc. vi : n. spp.
- Euryope notabilis, S. W. Africa, Péringuey, p. 86, Tr. S. African Phil. Soc. vi (2), n. sp.
- Euryscopa pusilla, Texas, p. 16, subtilis, California, p. 17, Horn, Tr. Am. Ent. Soc. xix, n. spp.
- Eustetha varians, Indo-China, Allard, p. 233, N. Arch. Mus. Paris (3) iii, n. sp.
- Fidia cana, Texas, Horn, p. 199, Tr. Am. Ent. Soc. xvi, n. sp.
- Galeruca, Geoff., = (Adimonia, Laich.); Weise, p. 414, Deutsche e. Z. 1892. G. xanthomelæna, habits, &c.; SMITH, Canad. Ent. xxiv, pp. 246-250: habits and life-history; RILEY, pp. 282 & 286, Canad. Ent. xxiv.
- Galerucella albopilosa, Burma, JACOBY, p. 969, Ann. Mus. Genov. (2) xii; G. celebensis, Celebes, Weise, p. 404, Deutsche e. Z. 1892; G. amabilis, Mexico, pl. xliii, fig. 11, terminalis, Panama, JACOBY, p. 327, Biol. Centr. Am. Col. vi (1 supp.): n. spp.
- Galerusoma, n. g., near Mimastra, for G. apicicorne, n. sp., Madagascar; JACOBY, p. 578, P. Z. S. 1892.
- Gonophora parvula, p. 237, rugicollis, p. 238, Burma, Gestro, Ann. Mus. Genov. (2) x; G. modiglianii, Engano, Gestro, p. 793, op. cit. xii; n. spp.

- Gynandrophthalma, 11-notata, p. 132, punctatissima, riridiceps, p. Turkestan, Weise, Deutsche e. Z. 1892; G. laticollis, Be Duv.vier, p. 399, Ann. Ent. Belg. xxxvi; G. mandarina, C. Lefèvre, p. xciv, Bull. Soc. Ent. Fr. lxi; G. pictipennis, p. decemnotata, octomaculata, p. 884, birmanica, p. 885, Burma, Jac Ann. Mus. Genov. (2) xii: n. spp.
- Haltica semenosoi, p. 462, weissi, = (laviuscula, Weiss), balassogloi, p. Central Asia, Jacobsohn, Hor. Ent. Ross. xxvi, n. spp.
- Haplosonyx ruspennis, Pedong, DUVIVIER, p. 439, Ann. Ent. Belg. xi H. orientalis, p. 962, ornata, p. 963, inornatus, varipes, p. 964, Bt JACOBY, Ann. Mus. Genov. (2) xii: n. spp.
- Hemyloticus, n. g., Typophorina, p. 571, for H. geniculatus, n. sp., h gascar, p. 572, pl. xxxix, fig. 6; Jacoby, P. Z. S. 1892.
- Heteraspis viridis, Burma, JACOBY, p. 86, Ent. xxv, Supp.; H. par nitida, p. 909, bhamoensis, p. 910, Burma, JACOBY, Ann. Mus. G (2) xii : n. spp.
- Hispida of Burma, with many notes; Gestro, Ann. Mus. Genov. pp. 225-268.
- Hispa elegantula, Bengal, DUVIVIER, p. 447, Ann. Ent. Belg. xxxvi megacantha, p. 249, cariana, p. 251, maculata, p. 252, brevic p. 253, discicollis, p. 254, doriæ, p. 256, monticola, p. 257, m p. 259, paronæ, p. 260, xunthospila, p. 261, platyprioides, p. 262, fida, p. 263, Gestro, Ann. Mus. Genov. (2) x; H. rubus, Guines, Gestro, p. 1019, Ann. Mus. Genov. (2) x: n. spp.
- Homophæta simulans, pl. xlii, figs. 24 & 25, violacea, Mexico, Jap. 316, Biol. Centr. Am. Col. vi (1 supp.), n. spp.
- Hoplacerus, n. g., near Diphaulaca, for H. tibialis, n. sp., Mexico, pl fig. 21, JACOBY, p. 344, Biol. Centr. Am. Col. vi (1 supp.).
- Hylaspoides, n. g., near Hylaspes, p. 443, for H. magnifica, n. sp., Si p. 444; Duvivier, Ann. Ent. Belg. xxxvi.
- Hyperacantha flavonigra, var. described; GAHAN, p. 92, P. 1892.
- Hyperaxis distincta, Bengal, DUVIVIER, p. 411, Ann. Ent. Belg.: n. sp.
- Hyphamia submetallica, Burma, JACOBY, p. 985, Ann. Mus. Genov. (n. sp.
- Hyphasis signata, Bengal, DUVIVIER, p. 429, Ann. Ent. Belg. xxxv distincta, p. 936, intermedia, inornata, p. 937, balyi, p. 938, J. Ann. Mus. Genov. (2) xii : n. spp.
- Hypolampsis inornata, Mexico, JACOBY, p. 323, Biol. Centr. Am. ( (1 supp.), n. sp.
- Ivongius nigromaculatus, Madagascar, JACOBY, p. 569, pl. xxxix, P. Z. S. 1892, n. sp.
- Konbirella, n. g., Galerucinæ, for K. cardoni, n. sp., Bengal; Duv p. 438, Ann. Ent. Belg. xxxvi.
- Labidostomis glycyrrhizæ, note on; BECKER, p. 70, Bull. Mosc. 189 L. urticarum, China, FRIVALDSZKY, p. 119, Term. füzetek, xv. n. si

Lactica bipustulata, Burma, JACOBY, p. 919, Ann. Mus. Genov. (2) xii;
L. vittatipennis, p. 345, pl. xliii, fig. 22, hidalgoensis, L.? varicornis,
fig. 24, p. 346, Mexico, JACOBY, Biol. Centr. Am. Col. vi (1 supp.) :
n. spp.

Lema melanopa, habits; VERHOEFF, p. 298, Ent. Nachr. xviii; L. sayi,

habits; RICHARDSON, p. 240, P. E. Soc. Wash. ii.

L. rugicollis, madagascariensis, Madagascar, JACOBY, p. 565, P. Z. S. 1892;
L. pulchella, S. W. Africa, Péringuey, p. 82, Tr. S. African Phil.
Soc. vî (2);
L. mashuana, S. Africa, Péringuey, p. 133, t. c.;
L. robusta, p. 869, mandibularis, p. 870, dimidiatipennis, p. 871, obscuritarsis, birmanica, p. 872, feæ, palonensis, p. 873, Burma, JACOBY, Ann.
Mus. Genov. (2) xii;
L. centromaculata, Nias I., Weise, p. 386,
Deutsche e. Z. 1892;
L. smithi, fulvipes, p. 341, quinquelineata, sponsa, p. 342, Mexico, JACOBY, Biol. Centr. Am. Col. vi (1 supp.):
n. spp.

Lina: minute structure of elytra; HOFFBAUER (405).

Longitarsus rangoonensis, p. 920, birmanicus, p. 921, Burma, JACOBY, Ann.

Mus. Genov. (2) xii, n. spp.

Luperodes braeti, Bengal, Duvivier, p. 437, Ann. Ent. Belg. xxxvi; L. impressipennis, p. 950, subrugosus, tarsalis, p. 951, pygidialis, p. 952, Burma, Jacoby, Ann. Mus. Genov. (2) xii; L. salcini, Guatemala, Jacoby, p. 336, Biol. Centr. Am. Col. vi (1 supp.): n. spp.

Luperomorpha vittata, p. 427, nigripennis, albofasciata, p. 428, Bengal,

DUVIVIER, Ann. Ent. Belg. xxxvi : n. spp.

Luperus tenuelimbatus, Obock, FAIRMAIRE, p. 126, Rev. d'Ent. xi; L. constricticollis, Burma, JACOBY, p. 949, Ann. Mus. Genov. (2) xii: n. spp.

Mucetes peringueyi, S. Africa, LEFÈVRE, p. 41, Tr. S. African Phil. Soc. vi, n. sp.

Macrima abdominalis, varietal note; JACOBY, p. 88, Ent. xxv (Supp.).

Malacorhinus cobanensis, pl. xliii, fig. 18, exclamationis, fig. 19, Central America, Jacoby, p. 336, Biol. Centr. Am. Col. vi (1 supp.), n. spp.

Malacosoma sikoræ, aterrimum, p. 575, flavicorne, p. 576, pl. xxxix, fig. 12, Madagascar, Jacoby, P. Z. S. 1892; M. nigricolle, Burma, Jacoby, Ann. Mus. Genov. (2) xii: n. spp.

Malaxia pallipes, Burma, JACOBY, p. 969, Ann. Mus. Genov. (2) xii,

n. sp.

Mandarella, n. g., Agelasticites, p. 433, for M. nagpurensis, n. sp., Bengal, p. 434; DUVIVIER, Ann. Ent. Belg. xxxvi.

Mastostethus angustovittatus, Mexico, JACOBY, p. 343, Biol. Cent. Am. Col. vi (1 supp.), n. sp.

Megalognatha bohemanni, = (Cneorane foveicollis, Jac.), M. elegans, = (Apophylia elegantula, Jac.); Jacoby, Ent. xxv, p. 162.

Megalopus basalis, Mexico, Jacoby, p. 344, Biol. Centr. Am. Col. vi (1 supp.), n. sp.

Megalostomis: characters of, N. American; HORN, p. 11, Tr. Am. Ent. Soc. xix.

- Megamerus mandibukuris, N. W. Australia, BLACKBURN, p. 67, Tr. B. S. Austr. xv, n. sp.
- Melasoma populi, n. var., janaceki; REITTER, p. 240, Wien. ent. Z. x Melitonoma decorata, S. W. Africa, Péringuey, p. 84, Tr. S. Af Phil. Soc. vi (2), n. sp.
- Menippus ænsipennis, Celebes, WEISE, p. 404, Deutsche e. Z. 1892, n. Menius distunti, Transvaal, JACOBY in Distant, Naturalist Transp. 205, pl. i, flg. 6, n. sp.
- Merists cardoni, E. India, DUVIVIER, p. 440, Ann. Ent. Belg. x n. sp.
- Mesodera brevicollis, Mexico, JACOBY, p. 347, Biol. Centr. Am. C (1 supp.), n. sp.
- Metachroma aneicolle, p. 211, longulum, p. 212, aterrimum, p. 214, anale, p. 215, parallelum, p. 217, N. America, Horn, Tr. Am. Ent xix, p. spp.
- Metacoryna laripennis. Guatemala, pretiosa, Mexico, pl. xliii, fi Jacoby, p. 338, Biol. Centr. Am. Col. vi (1 supp.), n. sp.
- Metacycla rugipennis, Mexico, Jacoby, p. 337, Biol. Centr. Am. (1 supp.), n. sp.
- Metasyonycha circumcincta, N. America, Horn, p. 228, Tr. Am. Ent xvi, n. sp.
- Mimastra longicornis, p. 944, hirsuta, p. 945, Burma, Jacoby, Ann. Genov. (2) xii, n. spp.
- Mimastroides, n. g., Galerucines, p. 577, for M. madagascariensis, : Madagascar, pl. xxxix, fig. 11, p. 578; Jacoby, P. Z. S. 1892.
- Monolepta marginata, p. 981, birmanensis, p. 982, alboplagiata, g p. 983, fea, p. 984, Burma, Jacoby, Ann. Mus. Genov. (2 M. benalla, frogratti, Victoria, Blackburn, p. 549, P. Line N.S.W. (2) vi; M. morosa, Mexico, Jacoby, p. 340, Biol. Centr Col. vi (1 supp.): n. spp.
- Monoplatus puncticollis, Mexico, JACOBY, p. 320, Biol. Centr. Au vi (1 supp.), n. sp.
- Mouhotina birmanica, Burma, JACOBY, p. 913, Ann. Mus. Genov. (n. sp.
- Neobrotica pallida, ruatanæ, pl. xliii, fig. 17, Mexico, JACOBY, p. 335 Centr. Am. Col. vi (1 supp.), n. spp.
- Neolepta ruficollis, Burma, JACOBY, p. 992, Ann. Mus. Genov. (n. sp.
- Nestinus longicornis, Mexico, JACOBY, p. 324, Biol. Centr. Am. ( (1 supp.), n. sp.
- Nirina, n. g., near Pachytoma, for N. jacobyi, n. sp., E. Africa; 1 p. 403, Deutsche e. Z. 1892.
- Nisotra nigritarsis, klugii, Madagascar, JACOBY, p. 572, P. Z. S. N. papuensis, New Guinea, TRYON (953): n. spp.
- Nodina tarsalis, Bengal, DUVIVIER, p. 409, Ann. Ent. Belg. xxxvi; 1 vula, p. 902, robusta, p. 903, birmanica, p. 904, Burma, JACOBY Mus. Genov. (2) xii: n. spp.

- Nodonota clypealis, N. America, Horn, p. 232, Tr. Am. Ent. Soc. xix, n. sp. Nodostoma variabile, p. 406, angulicolle, p. 408, Bengal, Duvivier, Ann. Ent. Belg xxxvi; N. bellum, p. 895, bhamoense, p. 896, semicæruleum, feæ, p. 897, birmanicum, p. 898, capitatum, bimaculicolle, p. 899, denticolle, p. 900, violaceofasciatum, p. 901, semipurpureum, p. 902, Burma, Jacoby, Ann. Mus. Genov. (2) xii; N. indicum, Burma, multicostata, p. 86, apicipes, p. 87, Sumatra, Jacoby, Ent. xxv (Supp.): n. spp.
- Nonarthra variabilis, varr. described; DUVIVIER, p. 423, Ann. Ent. Belg.
- N. albofasciata, Duvivier, p. 424, Ann. Ent. Belg. xxxvi, n. sp.
- Ochralea rufobasalis, p. 977, pallida, p. 978, Burma, JACOBY, Ann. Mus. Genov. (2) xii, n. spp.
- Odontionopa discolor, Cape Town, LEFRVRE, p. 39, Tr. S. African Phil. Soc. vi, n. sp.
- Edionychis clypeata, Madagascar, Jacoby, p. 574, P. Z. S. 1892; E. clypeata, p. 317, haroldi, durangoensis, p. 318, pavonina, p. 319, pl. xliii, fig. 1, Mexico, Jacoby, Biol. Centr. Am. Col. vi (1 supp.): n. spp.
- Oides basalis, flavolimbatus, New Guinea, TRYON (953), n. spp.
- Omotyphus suturalis, pl. xliii, fig. 3, p. 322, carinatus, fig. 4, p. 323, Mexico, Jacoby, Biol. Centr. Am. Col. vi (1 supp.), n. spp.
- Ootheca modesta, Transvaal, GAHAN in Distant, Naturalist Transvaal, p. 206, pl. i, fig. 11, n. sp.
- Ophraa opaca, Mexico, Jacoby, p. 327, Biol. Centr. Am. Col. vi (1 supp.), n. sp.
- Orsodacna atra, variation and synonyms; Horn, p. 6, Tr. Am. Ent. Soc. xix: variation, Hamilton, p. 160, Canad. Ent. xxiv.
- Orthaulaca, n. g. (with 5 subgg.), type, Galeruca similis, Ol.; Weise, p. 393, Deutsche e. Z. 1892.
- Omycephala longipennis, New Guinea, GESTRO, p. 1017, Ann. Mus. Genov. (2) x, n. sp.
- Pachnephoptrus, n. g., near Pachnephorus, for P. weisei, n. sp., Ordubad; Reitter, p. 153, Wien. ent. Z. xi.
- Pachnephorus plagiatus, Burma, JACOBY, p. 913, Ann. Mus. Genov. (2) xii, n. sp.
- Pachybrachys nigropunctatus, glycyrrhizæ, distinctive characters; Weise, p. 413, Deutsche e. Z. 1892. P. lecontei, n. n. for brevicollis, Jac., nec Lec.; Jacoby, p. 348, Biol. Centr. Am. Col. vi (1 supp.).
- Puchygnatha, n. g., for Orina dolene, Kr.; Weise, p. 352, Deutsche e. Z. 1892.
- Pachypalpa, n. g., for Galeruca luteicornis, Fab.; Weise, p. 392, Deutsche e. Z. 1892.
- Pachytoma dives, Karsch, = (Mesodonta submetallica, Jac.); JACONY, Ent. xxv, p. 162.
- P. obscura, Centr. Africa, GAHAN, p. 93, P. Z. S. 1892, n. sp.
- Pallena chapuisi, Cape Town, LEFEVRE, p. 41, Tr. S. African Phil. Soc. vi, n. sp.

- Parvelrylos. n. g., Arajuolites. p. 420, for P. apicipennia, n. sp., Bengal, p. 421. D. vivier. Ann. East. Belg. xxxvi.
- Peris merged in Typephorus: Hous., p. 207, Tr. Am. Ent. Soc. xix.
- Perides approximate. p. 43), Eride, p. 432, DUVIVIER, Ann. Ent. Belg. XXVI; P. bifurcute, reficellie, p. 953, cornute, p. 954, nigripensis, p. 956, proipensis, p. 956, mifacciate. p. 957, Burma, JACOBY. Ann. Mm. Genov. '2 xii: n. app.
- Perspeides perdelle, aigrepunctates. Burms, Jacost, p. 918, Ann. Mus. Genov. (2) xii, n. app.
- Personie gilgarnessia. p. 545. latipes, p. 546, regularia, p. 547. Australia, Blacketen, P. Linn. Soc. N.S.W. (2) vi; P. refopieta, W. Australia, Blacketen, p. 237. Tr. R. Soc. S. Anstr. xv; n. spp.
- Penniris stripics Vis., Cape Colomy, LEFRVEZ, p. 41, Tr. S. African Phil. Soc. vi. z. sp.
- Photocus brumens, p. 557, pl. xxxix, fig. 2. P.? eneicolles, fig. 4, bifasciena, fig. 3, p. 562, Madagascar, Jacoby, P. Z. S. 1892, n. spp.
- Phylletrete dentifera, S. France, Guillebeau, Échange, i, No. 7; P. chotenica, Bengal, Duvivier, p. 426, Ann. Ent. Belg. xxxvi: n. spp.
- Physinerus elongatulus, p. 321, flavopilosus, p. 322, Mexico, JACOBY, Biol. Centr. Am. Col. vi (1 supp.), n. spp.
- Phytodorta linneana, Schrank, synonymical note; WEISE, p. 414, Deutsche e. Z. 1492
- Piasso, n. g., near Lactica, for P. fulrus, n. sp., Mexico, pl. xliii, fig. 25; JACOBY, p. 345, Biol. Centr. Am. Col. vi (1 supp.).
- Pilemostoma, n. subg. of Cassida: Deserochers, p. 14, Supp., Frelon, i. Platypria acanthion, p. 245, ericulus, p. 247, Burma, Gestro, Ann. Mus. Genov. (2) x, n. spp.
- Platycantha africana, Congo, Duvivier, p. 167, Ann. Ent. Belg. xxxvi; P. indica, Bengal, Duvivier, p. 445, Ann. Ent. Belg. xxxvi: n. spp. Plectroscelis ollifi. N. S. Wales, Blackburn, p. 548, P. Linn. Soc. N.S.W. (2) vi. n. sp.
- Podagrica decolorata, Congo, DUVIVIER, p. 60, Ann. Ent. Belg. xxxvi, n. sp.
- Pacilomorpha binotata, S. W. Africa, Phringuey, Tr. S. African Phil. Soc. vi (2), p. 83, n. sp.
- Prionispa pulchra, Borneo, Gorham, p. 84, pl. iv, fig. 2, P. Z. S. 1892, n. sp.
- Pseudocassida, n. subg. of Cassida, Desbrochers, p. 15 (Supp.) Frelon, i.
- Pseudocolaspis hottentota, rariegata, brachiata, p. 39, gracilipes, viridana, p. 40, S. Africa, Lefèvre, Tr. S. African Phil. Soc. vi; P. bengalensis, India, Duvivier, p. 412, Ann. Ent. Belg. xxxvi: n. spp.
- Pseudocrania, n. g., near Monolepta, for P. latifrons, n. sp., Quango; Weise, p. 400, Deutsche e. Z. 1892.
- Pseudodiabrotica, n.g., near Microbrotica, for P. metallica, n. sp., Mexico, pl. xliii, fig. 16; Jacoby, p. 334, Biol. Centr. Am. Col. vi (1 supp.).

- Pseudoides, n. g., Agelastina, for P. bivittata, n. sp., Burma; Jacoby, p. 966, Ann. Mus. Genov. (2) xii.
- Pseudopioméra, n. g., Leprotina, p. 907, for P. pallidicornis, n. sp., Burma, p. 908; Jacoby, Ann. Mus. Genov. (2) xii.
- Pteleon pubescens, Mexico, JACOBY, p. 337, Biol. Centr. Am. Col. vi (1 supp.), n. sp.
- Rhaphidopalpa bengalensis, Calcutta, aruensis, N. Guinea, ceramensis, Borneo, &c., niasensis, Sumatra, p. 394, chinensis, Shanghai, p. 395, Weise, Deutsche e. Z. 1892, n. spp.
- Rhembastus dimidiaticornis, p. 569, antennatus, p. 570, Madagascar, Jacoby, P. Z. S. 1892, n. spp.
- Rhinotmetus modestus, p. 320, minutus, p. 321, Mexico, Jacoby, Biol. Centr. Am. Col. vi (1 supp.), n. spp.
- Rhyparida striaticollis, Madagascar, Jacoby, p. 571, pl. xxxix, fig. 7, P. Z. S. 1892; R. aterrima, Burma, Jacoby, p. 911, Ann. Mus. Genov. (2) xii; R. mastersi, Queensland, Blackburn, p. 236, Tr. R. Soc. S. Austr. xv: n. spp.
- Rupilia rugulosa, Queensland, BLACKBURN, p. 238, Tr. R. Soc. S. Austr. xv, n. spp.
- Rybakowia, n. g., near Haltica, p. 465, for R. pyriformis, = (globicollis, Byb.), n. sp., Amdo, p. 466; Jacobsohn, Hor. Ent. Ross. xxvi.
- Sastra fulvicornis, Burma, JACOBY, p. 958, Ann. Mus. Genov. (2) xii, n. sp. Sastroides crassipalpis, referred to Sastra; JACOBY, p. 958, Ann. Mus. Genov. (2) xii.
- S. parvula, Burma, JACOBY, p. 959, Ann. Mus. Genov. (2) xii, n. sp.
- Saxinis politula, speculifera, California, HORN, p. 10, Tr. Am. Ent. Soc. xix, n. spp.
- Scelidopeis (?) violacea, Mexico, JACOBY, p. 339, Biol. Centr. Am. Col. vi, (1 supp.), n. sp.
- Sebaethe quadrimaculata, elongata, p. 922, immaculata, p. 923, S. Precticollis, p. 924, Burma, JACOBY, Ann. Mus. Genov. (2) xii, n. spp.
- Sermyloides dilaticornis, Sumatra, JACOBY, p. 88, Ent. xxv (Supp.), n. sp. Solenia abdominalis, p. 986, integricollis, p. 987, Burma, JACOBY, Ann. Mus. Genov. (2) xii, n. spp.
- Sphenoraia imitans, Burma, JACOBY, p. 961, Ann. Mus. Genov. (2) xii, n. sp.
- Sphærarthra, subg. (n.?) of Orthaulaca; Weise, p. 396, Deutsche e. Z. 1892.
- Sphæroderma antennata, p. 925, nigrita, p. 926, birmanica, acutangula, p. 927, varipennis, p. 928, discicollis, terminata, p. 929, pallidicornis, p. 930, Burma, Jacoby, Ann. Mus. Genov. (2) xii, n. spp.
- Spharometopa intermedia, Burma, JACOBY, p. 925, Ann. Mus. Genov. (2) xii, n. sp.
- Spilocephalus viridipennis, var. described and figured; GAHAN in Distant, Naturalist Transvaal, p. 207, pl. i, fig. 12.
- S. distanti, Transvaal, Gahan in Distant, Naturalist Transvaal, p. 208, n. sp.

- Spilopyra olliff, N. S. Wales, BLACKBURN, p. 544, P. Linn. Soc. N.5 (2) vi, n. sp.
- Sternoplatys picsus, Fab., = (pascosi, Baly); DUVIVIER, p. 374, .
  Ent. Belg. xxxvi.
- Stethomela caudata, Queensland, BLACKBURN, p. 237, Tr. R. Soc. S. A xv, n. sp.
- Syagrus rufo-brunneus, rugiceps, Zambezi, Leftvre, p. 43, Tr. S. Af Phil. Soc. vi, n. spp.
- Syneta: characters and synonymy of the N. American species; H pp. 1-5, Tr. Am. Ent. Soc. xix.
- Temnaspis flavicornis, p. 875, nigroplagiata, p. 876, Burma, JACOBY, Mus. Genov. (2) xii; T. squalidus, Indo-Chins, ALLARD, p. 23
  Arch. Mus. Paris (3) iii: n. spp.
- Theopea nigricollis, Borneo, JACOBY, p. 87, Ent. xxv (Supp.), n. sp. Timarcha arragonica, Spain, BALBI, Riv. Ital. Sci. Nat. xii, p. 50, n. Trichostola capensis, S. Africa, LEFEVRE, p. 42, Tr. S. African Phil. vi. n. sp.
- Trichotheca basifemorata, Bengal, Duvivier, p. 410, Ann. Ent. xxxvi, n. sp.
- Trirrhabda vicina, p. 324, hogsi, pl. xliii, fig. 8, sublavicollis, fig. 7, p. semiviridis, fig. 9, rugosa, fig. 10, guatemalensis, p. 326, Ce America, Jacoby, Biol. Centr. Am. Col. vi (1 supp.), n. spp.
- Tymnes violaceus, Pennsylvania, HORN, p. 222, Tr. Am. Ent. Soc n. sp.
- Wallacea inornata, Engano, Gestro, p. 792, Ann. Mus. Genov. (2 n. sp.
- Zeugophora: characters of N. American; HORN, p. 7, Tr. Am. Ent.

#### EROTYLIDÆ.

- [Cf. Gorham (350), Reitter (725), Ritsema (757).]
- Asmonax, n. g., near Encaustes, for A. whiteheadi, n. sp., Borneo, j. fig. 5; Gorham, p. 85, P. Z. S. 1892.
- Helota feæ, p. 886, serratipennis, p. 890, rotundata, p. 891, gestroi, p immaculata, p. 895, difficilis, p. 896, dohertyi, p. 898, notata, p dubia, p. 901, Burma, BITSEMA, Ann. Mus. Genov. (2) x, n. spp
- Triplax (Platychna) analis, Araxes Valley, p. 133, swanetica, Cau p. 134, Reitter, Wien. ent. Z. zi, n. spp.

## ENDOMYCHIDÆ.

- [Cf. Gahan (210), Gorham (350)]
- Amphisternus armatus, Borneo, Gorham, p. 86, pl. iv, fig. 3, P. Z. S. n. sp.
- Ancylopus fuscipennis, Trausvaal, Gahan in Distant, Naturalist T vaal, p. 210, pl. iv, fig. 10, n. sp..

- Epipocus cinctus, metamorphoses; Wickham, Bull. Lab. Iowa, ii, p. 198, pl. i.
- Eumorphus costatus, & described; Gorham, p. 87, P. Z. S. 1892.
- E. tumescens, p. 86, pl. iv, fig. 4, lucidus, p. 87, fig. 1, Borneo, Gorham, P. Z. S. 1892, n. spp.

#### COCCINELLIDAE.

- [Cf. Blackburn (93, 91), Blaisdell (97), Frivaldszky (309), Gorham (347, 348, 351), Sicard (846), Weise (402, 993, 995, 997, 1001)]
- Habit of congregating of Coccinellida, discussion; P. E. Soc. Wash. ii, pp. 169 & 170.
- Weise's "Bestimmungs-Tabellen" of European Coccinellide, translated into French, with notes by SICARD & BEDEL; SICARD, Abeille, xxviii, pp. 1–80.
- Adonia weisei, China, FRIVALDSZKY, p. 120, Term. füzetek, xv., n. sp.
- Bucolinus, n. g., near Bucolus, for B. longicornis, n. sp., Victoria; Black-Burn, p. 252, Tr. R. Soc. S. Austr. xv.
- Bacolus convexus, New S. Wales, Blackburn, p. 72, Tr. B. Soc. S. Austr. xv, n. sp.
- Chilocorides: characters of the group noticed; GORHAM, p. 174, Biol. Centr. Am. Col. vii.
- Chilocorus bipustulatus and renipustulatus, destroying Coccida; BEDEL, p. 2, Abeille, xxviii.
- C. chalybeatus, Kiu-Kiang, GORHAM, Ent. xxv (Supp.), p. 84; C. flavidus, Queensland, BLACKBURN, p. 239, Tr. R. Soc. S. Austr. xv: n. spp.
- Chmootriba antinorii, Scioa, Gorham, p. 911, Ann. Mus. Genov. (2) x, n. sp.
- Clanis soror, Bengal, WEISE, p. 25, Ann. Ent. Belg. xxxvi, n. sp.
- Cleis lynx, figured, pl. ix, fig. 19; GORHAM, Biol. Centr. Am. Col. vii.
- Coccinella 12-maculata, n. var., jakolewi; Weise, p. 140, Deutsche e. Z. 1892. C. septempunctata, temporary colonies of; Plateau, p. 393, Ann. Ent. Belg. xxxvi.
- Calophora unicolor, great variation and synonymy; Weise, pp. 20-24, Ann. Ent. Belg. xxxvi.
- C. mastersi, p. 238, guttata, p. 239, Queensland, Blackburn, Tr. R. Soc. 8. Austr. xv, n. spp.
- Cycloneda sallai, pl. ix, fig. 21, retrospiciens, pl. x, fig. 2, figured, rubida, variation and synonymy noticed, p. 171, abdominalis, varr. figured, pl. ix, figs. 22-24, gilardini, figured, pl. ix, fig. 25; GORHAM, Biol. Centr. Am. Col. vii.
- C. electra, Guatemala, GORHAM, p. 173, pl. x, fig. 3, Biol. Centr. Am. Col. vii., n. sp.
- Cycloscymnus, n. g., Bucolites, for C. minutus, n. sp., New S. Wales; Blackburn, p. 251, Tr. R. Soc. S. Austr. xv.
- Egleis kingi, Macl., == (varicolor, Muls.); BLACKBURN, p. 238, Tr. R. Soc. S. Austr. xv.

- Epilachna scioensis, p. 910, agreta, p. 911, Scion, Gorham, Ann Genov. (2) x; E. fairmairei, China, Frivaldezky, p. 121, T füsetek, xv; E. insignis, Kiu-Kiang, Gorham, Ent. xxv (Su p. 84: n. spp.
- Erithionyz, n. g., for Rhinobius lanosus, BLACKBURN, p. 259, Tr. R. Sc Austr. xv.
- Gymnoscymnus, n. g., p. 241, for G. quadrimaculatus, n. sp., Aust. p. 242; Blackburn, Tr. R. Soc. S. Austr. xv.
- Halysia oblongoguttata, minute structure of elytra; HOFFBAUER (40. H. smaciata, pl. ix, figs. 12 & 13, spistictica, figs. 14 & 15, p. 164, c pioni, pl. x, fig. 1, p. 165, Panama, Gorham, Biol. Centr. Am. vii, n. spp.
- Harmonia impustulata, larva and pupa; Planet, Le Nat. 1892, p. 17 Hyperaspis salaamensis, E. Africa, Weise, Deutsche e. Z. 1892, p. 15, Hypocyrema, n. g., near Cyrema, p. 250, for H. pauxilium, n. sp., tralia, p. 251; Blackburn, Tr. R. Soc. S. Austr. xv.
- Leis azyridis, n. var., duodecimsignasa; FRIVALDEZKY, p. 121, T füzetek, xv.
- Lioadalia lenticula, Kiu-Kiang, Gorham, Ent. xxv (Supp.), p. 82, n. Lipernes subviridis, New S. Wales, BLACKBURN, p. 72, Tr. R. Sc Austr. xv, n. sp.
- Megilla maculata, habits; RILEY, P. E. Soc. Wash. ii, p. 168.
- Midus pygmaus, Australia, BLACKBURN, p. 261, Tr. B. Soc. S. A xv, n. sp.
- Neda bicolor, Queensland, BLACKBURN, p. 238, Tr. R. Soc. S. Aust n. sp.
- Neocalvia arcolata, Panama; GORHAM, p. 169, pl. ix, fig. 20, Biol. (Am. Col. vii.
- Neohalysia perroudi, figured; Gobham, pl. ix, fig. 11, Biol. Centr. Col. vii.
- Orcus carulous, figured; GORHAM, pl. x, fig. 4, Biol. Centr. Am. Co. C. chalybeus, early stages, Ins. Life, v, p. 42.
- O. punctulatus, splendens, p. 240, ovalis, obscurus, p. 241, Aust Blackburn, Tr. R. Soc. S. Austr. xv, n. spp.
- Ortalia discoidea, E. India, Weise, p. 27, Ann. Ent. Belg. xxxvi, n. Oryssomus: Clypeaster variegatus, Phil., referred to this; Ger p. 261, Act. Soc. Chili, ii.
- Platynaspis, = (Microrrhymbus, Gerst.); Weise, p. 413, Deutsche 1892. P. abdominalis, n. n. for rufipennis, Weise; Weise, l.c.
- Psyllobora confuens, fig. 18, luctuosa, fig. 16, roci, fig. 17, figured, noticed, pp. 165, 168; GORHAM, Biol. Centr. Am. Col. vii.
- Ptychanatis axyridis, variation; GORHAM, Ent. xxv (Sapp.), p. 82.
- Rhizobius fugux, dorsalis, p. 70, coccus, lanosus, pulcher, p. 71, Aus BLACKBURN, Tr. B. Soc. S. Austr. xv; R. nigronotatus, ornatiz p. 253, speculifer, toowoomba, p. 254, satelles, aurantii, p. fasciculatus, apicalis corruleus, p. 256, plebeius, p. 257, Aus BLACKBURN, t. c.: n. spp.

- Rodolia: characters of; Weise, p. 25, Ann. Ent. Belg. xxxvi.
- R. breviuscula, E. India, pumila, Hong-Kong, Weise, p. 26, Ann. Ent. Belg., n. spp.
- Scymnodes koebeli, New S. Wales, BLACKBURN, Tr. R. Soc. S. Austr. xv, p. 69, n. sp.
- Scymnomorpha, n. g., near Pullus, for S. duplopunctulata, n. sp., Victoria; BLACKBURN, Tr. B. Soc. S. Austr. xv.
- Scymnus koebeli, variation; BLACKBURN, p. 242, Tr. R. Soc. S. Austr. xv.
  S. (Nephus) jakolewi, Irkutsk, Weise, p. 141, Deutsche e. Z. 1892; S. (Nephus) severini, p. 27, S. (Pullus) dorsualis, p. 28, E. India, Weise, p. 28, Ann. Ent. Belg. xxxvi; S. sydneyensis, australasia, p. 243, jocosus, cucullifer, p. 244, victoriensis, subelongatulus, p. 245, sublatus, pretiosus, inaffectatus, p. 246, whittonensis, S. (n. g.?) queenslandicus, p. 247, S. vagans, p. 248, Australia, Blackburn, Tr. R. Soc. S. Austr. xv: n. spp.
- S. lophantha, California, BLAISDELL, Ent. News, iii, p. 51.
- Semichnoodes, n. g., near Bucolus, p. 15, for S. kunowi, n. sp., E. Africa, p. 16; Weise, Deutsche e. Z. 1892.
- Semiadalia heydeni, Margelan, Weise, p. 110, Deutsche e. Z. 1892, n. sp.
  Serangium maculigerum, Queensland, hirtuosum, New S. Wales,
  BLACKBURN, p. 73, Tr. R. Soc. S. Austr. xv, n. spp.
- Summius, n. g., near Rhizobius for S. renardi, p. 29, cardoni, p. 30, n. spp., E. India; Weise, Ann. Ent. Belg. xxxvi.
- Trichorcus, n. g., near Orcus, for T. cinctus, n. sp., New S. Wales, BLACKBURN, Tr. R. Soc. S. Austr. xv, p. 73.
- Verania cardoni, E. India, Weise, p. 19, Ann. Ent. Belg. xxxvi, n. sp.

#### CORYLOPHIDA.

Clypeaster variegatus, Phil., referred to Oryssomus in Coccinellidæ; Germain, p. 261, Act. Soc. Chili, ii.

# (B.) HYMENOPT.

Structure of ovipositor; MARLATT (581).

Retardation of appearance of Hymenoptera summit of Mont Ventoux; NICOLAS, I XX. pt. 2.

Early stages of embryology; HERKING (396). Mode of union of abdominal segments; CARL Characters for the determination of the fi FREY-GESSNER, Bull. Soc. Murith. xiii, pp

### ACULEATA.

[Cf. André (12, 13), Benton (54), Benzi (55), & Marshall (149), Cameron (151), Cobi (197), EMERY (249-254), FOREL (287-2 FREY-GES-NER (307, 308), GRAPPE (358) KATURIC (473, 474), KELLER (475), KOI M (GRETTI (575, 576), MARCHAL (580), (615), MORAWITZ (624), NEREN (631), N (671), PERKINS (674), REED (719), BILEY SAUSSURE (210, 793). SCHULTHESS-RECH STADELMANN (883), THEOBALD (920), VAC (961, 962, 963, 964, 966, 967, 968, 969), \ (984, 986, 987). WILLISTON (1012), WROUG The use of the asperities on pupse; VERHOEFI Position of the sexes in pupal cells; relative: time of their appearance; VERHOEFF (96 Catalogue of Australian Acukata, pt. ii; FROG In Saussure's great work recorded below, by: the novelties are not indicated in a suf the letter "n" being all that is used, a species or genera described by the author previously. The Recorder begs to point any errors he may have fallen into.

#### APIDE.

Mode of secretion of wax; MAYER, MT. z. 8 pl. xxxii, figs. 4, 4a.

Revision of the bees described in Gray's Chili; Andrena mandibularis, Illinois, ROBERTSON, p.

- Anthidium manicatum, habits; VERHOEFF, Zool. Jahrb. Syst. vi, p. 710.
- A. buyssoni (Perez), Caracas, Buysson, p. 55, Ann. Soc. Ent. Fr. lxi, n. sp.
- Anthophora parietina, personata, pilipes, nests and habits, Verhoeff, Zool. Jahrb. Syst. vi, pp. 699-703; A. pilipes, habits of 3, Verhoeff, Ent. Nachr. xviii, p. 244.
- A. locsyi, Tibet, Mocsary, p. 130, Term. füzetek, xv, n. sp.
- Apis mellifica: workers feeding drones; Cook, Bull. Dep. Agric. Ent. xxvi, p. 89. Koschewnikoff on male organs, translated in Ann. N. H. (6) ix, p. 185. Distance it flies, Benton, Ins. Life, iv, p. 319.
- Bombus, the species of the Valais, with note on economy; FREY-GESSNER, Bull. Soc. Murith, xiii, pp. 32-36.
- B. silantjewi, p. 132, suworzewi, p. 135, Russia; Morawitz, Hor. Ent. Ross. xxvi, n. spp.
- Calliopsis parvus, Illinois, ROBERTSON, p. 273, Am. Nat. xxvi, n. sp.
- Camptopæum altaicum, Russia, Morawitz, p. 142. Hor. Eut. Ross. xxvi, n. sp.
- Ceratina carulea, habits; VERHOEFF, Zool. Jahrb. Syst. vi, p. 736.
- C. sitidula, Transcaspian region, Morawitz, p. 141, Hor. Ent. Ross. xxvi, n. sp.
- Chalicodoma muraria, parasites; CARPENTIER, Bull. Soc. L. Nord France, x, p. 328.
- Chelostoma florisomme, habits; ALFKEN, p. 209, Ent. Nachr. xviii.
- Colioxys dentigera, China, Mocsáry, p. 131, Term. füzetek, xv. n. sp.
- Colletes and Prosopis to form one family called Colletides; VERHOEFF, p. 734, Zool. Jahrb. Syst. vi.
- Epeclus productus and rusipes, distinctions of \$; SAUNDERS, Ent. M. M. xxviii, p. 161. E. pusillus, \$ described; Fox, Ent. News, iii, p. 29.
- Ecomulopsis neglectus, Caracas, Buysson, p. 55, Ann. Soc. Ent. Fr. lxi, n. sp.
- Halictus cylindricus, hermaphrodite; KRIEGER, pp. 137-140, Z. Naturw. lxv. H. sezcinctus, habits; VERHOEFF, pp. 711-717, Zool. Jahrb. Syst. vi.
- H. sphecodimorphus, Spain, Oran, angustifrons, Algeria, Vachal, p. xxii, Bull. Soc. Ent. Fr. lxi; H. denticollis, E. Siberia, p. 145, dmitrijewi, p. 146, Russia, monstrificus, E. Siberia, p. 147, Jakowlew, Hor. Ent. Ross. xxvi; H. luridipes, gibber, p. cxxxvi, magrettii, p. cxxxvii, Soudan, Vachal, Bull. Soc. Ent. Fr. lxi; H. reticulatus, Florida, hartii, Illinois, p. 268, floridanus, Florida, pruinosus, p. 269, cephalicus, obscurus, p. 270, Illinois, ashmeadii, Florida, illinoensis, p. 271, anomalus, Illinois, apophensis, longiceps, Florida, p. 272, Robertson, Am. Nat. xxvi; H. clavatus, Venezuela, Buysson, p. 55, Ann. Soc. Ent. Fr. lxi: n. spp.
- Heriades copetica, Transcaspian Region, Morawitz, p. 149, Hor. Ent. Ross. xxvi, n. sp.
- Megachile: nests of N. American species noticed; PACKARD, Psyche, vi, p. 340. M. dufourii, nest; André, Échange, i, No. 7, p. 1.

- M. rhinoceros, China, Mocsáry, p. 131, Term. füsetek, xv, n. sp. Meliseodes palustris, Illinois, Robertson, p. 273, Am. Nat. xxvi, n. sp. Meliturga pictipes, spinosa, Russia, Morawitz, Hor. Ent. Ross, n. spp.
- Nomada baldiniana, Italy, Benzi, Atti Soc. Mod. (3) xi, p. 220, n. sp. Osmia smarginata, habits, architecture; Verhoeff, Zool. Jahrb. Syr p. 703. A. leucomelana, habits; Verhoeff, t. c. p. 736: account the mode in which its larva is devoured by Stelis minuta; Verhop. 41, Zool. Ans. xv. O. tridentata, nidification and habits; hoeff, Ent. Nachr. xviii, pp. 225-227. O. simillima, pacifica, noticed; Packard, Psyche, vi, p. 340.
- O. (Acanthosmia) gräffei, Trieste, SCHMIEDEKNECHT, p. 137, Atti Trieste, viii; O. friesei, Dalmatia, VERHOEFF, p. 289, Ent. N xviii; O. globicola, Cape Good Hope, STADELMANE, p. 235, i fig. 3, Berl. Ent. Zeit. xxxvii: n, spp.
- Panurginus hersi, E. Siberia, MORAWITZ, p. 144, Hor. Ent. Ross. n. sp.
- Procopis annulatus, habits of 3; VERHOEFF, p. 247, Ent. Nachr.: P. brevicornis, habits discussed; VERHOEFF, p. 732, Zool. J Syst. vi.
- P. magrettii, Soudan, Vachal, p. exxxv, Bull. Soc. Ent. Fr. lxi, n.: Rhophites mandibularis, p. 138, dispar, p. 139, Russia, Morawitz Ent. Ross. xxvi, n. spp.
- Spheoodes: Sabulicola is a subg. S. cirsii, characters; VERHOEFF, Berl. Ent. Zeit. xxxvii.
- Stelis cognata, Araxes Valley, Kohl, p. 230, Ann. Hofmuseum Wie n. sp.
- Tetralomia pygialis, Venezuela, Buysson, p. 54, Ann. Soc. Ent. F. n. sp.
- Trigona apiformis, ochrotrichus, Venezuela, Buysson, p. 54, Ann. Soc Fr. lxi, n. spp.
- Xylocopa virginica, hibernation of &, egg noticed; HOWARD, p P. E. Soc. Wash. ii.
- X. gribodoi, Somaliland, MAGRETTI, p. 959, Ann. Mus. Genov. (n. sp.

# VESPIDÆ (DIPLOPTERA).

- Eumenes fraterna, habits; Southwick, Ins. Life, v, p. 107.
- E. kohli, Nossibé, Brancsik, p. 159, Jahresh. Ver. Trencsin. xiii, n Hoplopus spinipes, habits; Verhoeff, p. 710, Zool. Jahrb. Sy H. lævipes, habits; id. t. c. p. 739.
- Odynerus murarius, habits; ALFKEN, p. 210, Ent. Nachr. xviii: morphoses; King, Sci. Goss. 1892, pp. 196-198, woodcuts spinipes, metamorphoses and parasites; CARPENTIER, Bull. S. Nord. France, x, p. 326.
- O. pauli, Cadiz, Medina, p. 142, Act. Soc. Esp. xxi; O. aldric. America, Fox, p. 197, Ent. News, iii: n. spp.

- Polistes gallicus, color differences in sexes; Medina, Act. Soc. Esp. xxi p. 177.
- P. puncticollis, E. Siberia, Morawitz, p. 150, Hor. Ent. Ross. xxvi, n. sp. Pterocheilus hispanicus, Ciudad-Real, Medina, p. 144, Act. Soc. Esp. xxi; P. bezzii, Pavia, Cobelli, p. 67, Verh. z.-b. Wien, xlii: n. spp.

Zethus slossonæ, Florida, Fox, p. 29, Ent. News, iii, n. sp.

# Hymenoptera Fossoria. (Sphegidæ, Crabbonidæ, Mutillidæ, Etc.)

- Systematic arrangement and classification of the Heterogyna, consisting of Scoliens, Thynniens, Bradynobæniens, Mutilliens, Fedischenkiens, Sapygiens; SAUSSURE in Grandidier, Madagascar Hym. Ac. xx, pt. 1, pp. 176, &c.
- Many of the species described by SAUSSURE last year in MT. Schw. ent. Ges. viii, are now figured and described more fully in GRANDIDIER, Madagascar, xx, pt. 1.
- General observations on the dwellers in Rubus stems; Verhoeff, Zool. Jahrb. Syst. vi, pp. 746-751.
- Agenia tæniata, p. 347, pl. xxiv, fig. 8, melæna, p. 348, pl. xxv, fig. 11, nitidula, p. 349, pl. xxii, fig. 10, Madagascar, Saussure in Grandidier, Madagascar, xx, pt. 1, n. spp.
- Ammophila affinis, account of its instinct and mode of stinging; Marchal, Arch. Z. exper. (2), pp. 23-36. A. yarrowi, habits; Williston, Ent. News, iii, p. 85. A. gryphus, habits; Pergande, P. E. Soc. Wash. ii, p. 256.
- A. imerinæ, Madagascar, SAUSSURE, p. 435, Grandidier, Madagascar, xx, pt. 1, n. sp.
- Ampulex nigrocerulea, Transvaal, SAUSSURE, p. 212, pl. iv, fig. 6, in Distant, Naturalist Transvaal, n. sp.
- Aporus compressus, p. 309, pl. xxiv, fig. 1, prismaticus, p. 311, fig. 2, Madagascar, Saussure in Grandidier Madagascar, xx, pt. 1, n. spp.
- A status: synopsis of the N. American species; Fox, p. 232, Canad. Ent. xxiv.
- A. pygidialis, N. America, Fox, Canad. Ent. p. 234; A. glaber, Rio grande do Sul, Kohl, p. 230, Ann. Hofmuseum Wien, vii; A. blandus, Madagascar, Saussure, p. 519, pl. xxvii, fig. 4, Grandidier, Madagascar, xx, pt. 1: n. spp.
- Bembex fischeroides, Somaliland, MAGRETTI, p. 958, Ann. Mus. Genov. (2)
  x; B. hova, Madagascar, SAUSSURE, p. 457, Grandidier, Madagascar,
  xx, pt. 1: n. spp.
- Bothynostethus, Pisonitus, Silaon, are one; PATTON, p. 90. Ent. News, jii. Bradynobænus, to form a distinct group; SAUSSURE, p. 250, Grandidier, Madagascar, xx, pt. 1.
- Callosila, n. subg. of Cosila, p. 232, with C. flavicornis, minuta, inornata, n. spp., S. Australia, p. 233, SAUSSURE in Grandidier, Madagascar, xx, pt. 1.

- Corcoris sibirica, E. Siberia, MORAWITZ, p. 159, Hor. Ent. Ross. xx:
  C. carnaria, p. 545, pl. xxvii, fig. 13, spirans, fig. 12, p. 5
  Madagascar, SAUSSURE in Grandidier, Madagascar, xx, pt.
  n. spp.
- Ceropales: revision of the N. American species; Fox, Tr. Am. Ent. 5 xix, pp. 49-63.
- C. stretchii, p. 52, cressoni, p. 58, N. America, Fox, Tr. Am. Ent. 8 xix, n. spp.
- Chevrieria unicolor, habits; VERHOEFF, Zool. Jahrb. Syst. vi, p. 730.
- Chlorampulex, n. g., p. 443, near Ampulex, for A. sibirica and compre Fab., with C. striolata, n. sp., Zanzibar, p. 446, Saussune in Gradier, Madagascar, xx, pt. 1.
- Chlorion forficula, Madagascar, p. 422, SAUSSURE in Grandidier, Magascar, xx, pt. 1, n. sp.
- Cosila jheringi, S. Brazil, SAUSSURE in Grandidier, Madagascar, xx, p p. 234, n. sp.
- Crabronidæ, Pemphredonidæ, Trypozylidæ, to form three distinct fami characters given; Verhoeff, p. 728, Zool. Jahrb. Syst. vi.
- Crabro capitosus and spp. in Sambucus twigs, habits noticed; VERHO Zool. Jahrb. Syst. vi, pp. 717-724.
- C. (Clytochrysus) longipalpis, Heidelberg, Verhoeff, p. 70, Ent. Na xviii; C. femoralis, p. 161, biguttatus, p. 163, E. Siberia, jaroschewi p. 164, altaicus, p. 165, Russia, jakolewi, E. Siberia, p. 170, flavici Transcaspian, p. 172, ruthenicus, p. 174, flagellarius, p. 175, Ru martjanowi, E. Siberia, p. 178, Morawitz, Hor. Ent. Roes, xxvi chrysites, Sikkim? p. 197, C. (Calocrabro) styrius, Admont, p. amurensis, Amurland, p. 199, C. (Crossocerus) acanthophorus, Swi land, p. 201, C. circumscriptus, Bahia, p. 201, C. (Lindenius) al Algeria, p. 202, columbianus, British Columbia, p. 203, Kohl., . Hofmuseum Wien, vii; C. nitidirentris, New Jersey, p. 9, excav Florida, p. 10, Fox, Ent. News, iii: n. spp.
- Ctenagenia, n. subg. of Agenia, for Heteronyx madecasea, Priocn viduus and A. vespiformis; SAUSSURE, p. 343, in Grandidier, M gascar, xx, pt. 1.
- Cyphononyx muelleri, Africa, p. 402, SAUSSURE in Grandidier, Madaga xx, pt. 1; C. antennatu, Transvaal, SAUSSURE, p. 217, pl. v, fig. Distant, Naturalist Transvaal: n. spp.
- Dasyproctus crudelis, pl. xxvi, fig. 5, sævus, p. 579, ferox, p. 580, imm Madagascar, immitis, S. Africa, p. 581, indicus, Calcutta, p. Saussure in Grandidier, Madagascar, xx, pt. 1, n. spp.
- Diodontus hyalipennis, Caucasus, Kohl, p. 204, Ann. Hofmuseum V vii; D. occidentalis, rugosus, p. 315, flaviturnis, gillettei, p. 316, nig p. 317, N. America, Fox, Tr. Am. Ent. Soc. xix: n. spp.
- Discolia præcana, pl. iv, fig. 11, præstabilis, fig. 9, Transvaal, SAUSS p. 222, in Distant, Naturalist Transvaal, n. spp.
- Dolichurus secundus, p. 450, pl. xxvi, fig. 1, tertius, p. 452, Madaga SAUSSURE in Grandidier, Madagascar, xx, pt. 1, n. spp.

- Elis barbata, S. Africa, p. 217, pilosella, Madagascar, p. 220, hisurta, Ceylon, p. 221, grandidieri, Madagascar, p. 222, SAUSSURE in Grandidier, Madagascar, xx, pt. 1; E. reticulata, Poona, p. 109, E. (Trielis) orientalis, Ceylon, p. 112, CAMERON, Mem. Soc. Manch. (4) v:n. spp.
- Fedtschenkiens: characters and affiuities; SAUSSURE, p. 301, in Grandidier, Madagascar, xx, pt. 1.
- Gorytes cribratus, Transcaspian Region, Morawitz, p. 157, Hor. Ent. Ross. xxvi; G. albosignatus, mirandus, p. 152, rufocinctus, p. 153, Fox, Canad. Ent. xxiv: n. spp.
- Harpactes histrio, Madagascar, SAUSSURE, p. 530, in Grandidier, Madagascar, xx, pt. 1, n. sp.
- Heliocausus, n. g., near Tachysphex, p. 210, for H. fairmairei, n. sp., Chili, p. 211; Kohl, Ann. Hofmuseum Wien, vii.
- Hemimeria, n. g. (Scoliidæ), for M. semirufa, Guer., and H. savignyi, n. sp., Egypt; SAUSSURE, p. 249, in Grandidier, Madagascar, xx, pt. 1.
- Hemipogonius, n. subg. of Pogonius, for Priocnemis venustipennis, Saus., figured, pls. ix & x, fig. 28; SAUSSURE, p. 334, in Grandidier, Hymenoptera, xx, pt. 1.
- Hemisalius, n. subg. of Salius, q. v.; SAUSSURE, infrd.
- Heteronyx madecassus, = (Agenia marginipennis); SAUSSURE, p. 387, in Grandidier, Madagascar, xx, pt. 1.
- Homonotus hirundo, Madagascar, SAUSSURE, p. 331, pl. xxiv, fig. 4, in Grandidier, Madagascar, xx, pt. 1; H. cærulans, p. 213, pl. v, fig. 1, pedestris, p. 214, fig. 6, Transvaal, SAUSSURE in Distant, Naturalist Transvaal: n. spp.
- Isodontia leonina, Madagascar, SAUSSURE in Grandidier, Madagascar, xx, pt. 1, p. 431, pl. xix, fig. 6, n. ep.
- Lurra divisa, Pttn., is 2 of athiops, Cresson; Patton, p. 90, Ent. News, iii.
- L. carbonaria, p. 491, rariipes. p. 493, pl. xxvii, fig. 1, heydenii, p. 494, pls. x & xii, fig. 18, Madagascar, SAUSSURE in Grandidier, Madagascar, xx, pt. 1; L. angustifrons, Java, saussurei, Madagascar, Kohl, p. 219, Ann. Hofmuseum Wien, vii: n. spp.
- Larrada similis, p. 127, sinensis, p. 128, China, Mocsáry, Term. füzetek, xv. n. sp.
- Larrazena duz, Æthiopian Region, Kohl, p. 220, Ann. Hofmuseum Wien, vii, n. sp.
- Larropeis, n. g., type Larrada tenuicornis, Smith; PATTON, p. 90, Ent. News, iii.
- Liacos fulro-picta, E. India, CAMERON, p. 98, Mem. Soc. Manch. (4) v, n. sp.
  Liris: table of characters of the species; Kohl, pp. 227-229, Ann. Hofmuseum Wien, vii.
- L. pedestris, Madagascar, SAUSSURE, p. 517, in Grandidier, Madagascar, xx, pt. 1; L. gibbosa, Arabia, p. 222, mordax, Borneo, p. 223, anthracina, p. 224, sagax, morio, p. 225, Sikkim?, Kohl, Ann. Hofmuseum Wien, vii; L. cozalis, Florida, Patron, p. 90, Ent. News, iii: n. spp.

Lyreda subita, Say, generic position queried; PATTON, p. 90, Eut. Nev iii, and Pox, p. 138, t. c.

Mellines arrensis: statement as to habits corrected; VERHORFF, p. 69 Zool. Jahrb. Syst. vi.

Meria: table of characters of the 2, p. 248; SAUSSURE in Grandidi Madagascar, xx, pt. 1.

M. hottentota, S. Africa, Saussunz in Grandidier, Madagascar, xx, pt. p. 249, n. sp.

Mesa, n. g.?, for a part of Plesia, with the following n. spp.:—hete gamia, Mozambique, atopogamia, Zanzibar, peringueyi, pl. vi, fig.: hottentotta, S. Africa; SAUSSURE, pp. 244 & 245, in Grandidier, Magascar, xx, pt. 1.

M. diapherogamia, Transvaal, SAUSSURE, p. 225, pl. iv, fig. 8, in Dista Naturalist Transvaal, n. sp.

Mesocrabro. n. subg., for a division of the subg. Solenius; VERHOEI p. 70, Ent. Nachr. xviii.

Methocs orientalis, n. n. for orientalis, Smith (ex parts); MAGRETTI, AI Mus. Genov. (2) xii.

M. rufa, Mexico, Saussure in Grandidier, Madagascar, xx, pt. 1, n. sp. Microcrabro, n. subg. of Crabro, p. 574, for M. micromegas, n. sp., Madgascar, p. 575, pl. xxvi, fig. 6; Saussure in Grandidier, Madgascar, xx, pt. 1.

Mimesa picicornis, Irkutsk, Morawitz, p. 155, Hor. Ent. Ross. xx n. sp.

Miscophus kandlirschui, Algeria, Kohl., p. 214, Ann. Hofmuseum Wie vii, n. sp.

Mutilla representans, & described, p. 205, dimidiata, Q, p. 20
MAGRETTI, Ann. Mus. Genov. (2) xii. M. crythrocephala, a parase
of Crabro rubicola; BORRIES, p. 247, Ent. Tidskr. xiii.

M. somalica, p. 951, robecchii, p. 952, gestroi, cuneatiformis, p. 953, decke p. 954, doriæ, teleki, p. 955, rufo-guttata, p. 956, conjunctoides, p. 9 Somaliland, MAGRETTI, Apn. Mus. Genov. (2) x; M. albisty Transvaal, SAUSSURE, p. 225, pl. iv, fig. 7, in Distant, Natural Transvaal; M. sikoræ, p. 291, divorta, p. 296, Madagascar, SAUSSUI in Grandidier, Madagascar, xx, pt. 1; M. freyi, Nossibé, BRANCS p. 160, pl. vii, fig. 7, Jahresh. Ver. Trencein, xiii; M. kauthel p. 124, taprobana, p. 125, trimaculata, p. 126, wroughtoni, pulch rentris, p. 127, metallica, p. 128, poonaensis, p. 129, veda, pulchrice p. 130, erythrocera, p. 131, bhudda, p. 132, serratula, insularis, p. 13 kanara, constancea, p. 134, yerburghi, p. 135, pulcherina, p. 136, India, CAMERON, Mem. Soc. Manch. (4) v; M. ruficrus, p. 2 tridungulata, p. 209, fee, p. 210, ninnii, p. 211, pectinospina pulchrinella, p. 212, tenasserimica, conjungenda, p. 214, laminel p. 215, kirbyi, p. 216, kohli, p. 217, maximina, p. 218, dissimilan p. 219, pilosella, p. 220, emergenda, p. 222, subanalis, p. 223, emer p. 225, kandlirechi, stephani, p. 227, josephi, p. 228, gribod tornatorei, schlettereri, p. 229, cariana, p. 231, exiloides, p. 2;

- circumscribenda, p. 233, M. (Dasylabris) sejugoides, p. 234, Burma, MAGRETTI, Ann. Mus. Genov. (2) xii; M. stephana, cincta, signata, lamellifera, Venezuels, Buysson, p. 58, Ann. Soc. Ent. Fr. lxi: n. spp.
- Mygnimia muelleri, regina, W. Africa, SAUSSURE, p. 412, in Grandidier, Madagascar, xx, pt. 1; M. belzebuth, p. 218, pl. v, fig. 8, depressa, p. 219, distanti, p. 220, fig. 7, fallax, p. 221, fig. 5, Transvaal, SAUSSURE, in Distant, Naturalist Transvaal: n. spp.
- Mysine clavata, Transvaal, SAUSSURE, p. 242, pl. xxii, fig. 7; M. nitida, Poona, Cameron, p. 114, pl. iv, fig. 2, Mem. Soc. Manch. (4) v: n. spp.
- Notogonia sculpturata, Egypt, Kohl, p. 221, Ann. Hofmuseum Wien, vii; N. alaris, p. 503, pl. xxvi, fig. 2, dyscheira, p. 510, rufipes, p. 511, pl. xxvi, fig. 3, reticulata, p. 512, Madagascar, Saussure, in Grandidier, Madagascar, xx, pt. 1: n. spp.
- Nysson curtulus, Turkestan, Morawitz, p. 156, Hor. Ent. Ross. xxvi, n. sp. Oxybelus rufopictus, minutissimus, Transcaspian Region, Morawitz, p. 180, Hor. Ent. Ross. xxvi; O. cristatus pl. xxvii, fig. 5, p. 559, subcristatus, p. 561, Madagascar, pinnatus, p. 562, pl. xxvii, figs. 6 & 7, angustus, fig. 8, peringueyi, fig. 9, p. 563, S. Africa, Saussure, in Grandidier, Madagascar, xx, pt. 1; O. (Oxybeloides) odontophorus, p. 205, modestus, p. 206, Araxes Valley, obscurus, Corfu, p. 207, columbianus, British Columbia, p. 208, O. latifrons, Araxes Valley, p. 209, Kohl, Ann. Hofmuseum Wien, vii: n. spp.
- Passalacus relativus, distinctus, p. 319, dispar, p. 320, N. America, Fox, Tr. Am. Ent. Soc. xix, n. spp.
- Pemphredonidæ: revision of the N. American species; Fox, pp. 307-326, Tr. Am. Ent. Soc. xix.
- Pemphredon angularis, rileyi, p. 310, confertim, p. 311, bipartior, tenax, p. 313, N. America, Fox, Tr. Am. Ent. Soc. xix, n. spp.
- Philanthus triangulum, oviposition; Verhoeff, Zool. Jahrb. Syst. vi, p. 696.
- P. immitis, Madagascar, SAUSSURE, p. 537, pl. xxiii, fig. 24, in Grandidier, Madagascar, xx, pt. 1, n. sp.
- Planiceps planatus, California, Fox, p. 171, Ent. News, iii, n. sp.
- Plesia, tabulation of the females; SAUSSURE, p. 243, in Grandidier, Madagascar, xx, pt. 1.
- P. argentina, New World, SAUSSURE, p. 243, in Grandidier, Madagascar, xx, pt. 1; P. mandalensis, Burma, Magrerri, p. 257, Ann. Mus. Genov. (2) xii: n. spp.
- Pogonius macilentus, Madagascar, SAUSSURE, p. 334, pl. xxiv, fig. 5, in Grandidier, Madagascar, xx, pt. 1, n. sp.
- Polemistus, n. g., Pemphredoniens, p. 565, for P. macilentus, Madagascar, p. 567, pl. xxvii, fig. 11, pusillus, Mexico, p. 567, n. spp., SAUSSURE, in Grandidier, Madagascar, xx, pt. 1.
- Pompiliens: characters and definition, with table of genera; SAUSSURE, pp. 303-308, in Grandidier, Madagascar, xx, pt. 1.

- Pompilus nigerrimus, n. var. kohli, p. 71, P. speciosus, characters of (Planiceps) latreillei, = (Aporus lugubris, Verh.), p. 72; Verho Ent. Nachr. xviii.
- Priocnemis (Salius) vitripennis, p. 65, P. fuscopennis, p. 66, fallax, relip. 69, Germany, Verhoeff, Ent. Nachr. xviii; P. hirsutus, Tivaal, Saussurs, p. 216, pl. v, fig. 3, in Distant, Naturalist Transv P. muelleri, W. Africa, p. 372, deplanatus, p. 376, pls. viii & x. fig. sikora, p. 386, pl. xxv, fig. 13, Madagascar, Saussure in Grandi Madagascar, xx, pt. 1: n. spp.
- Psen dilectus, Madagascar, SAUSSURE, p. 571, pl. xxvi, fig. 7, in Grand Madagascar, xx, pt. 1, n. sp.
- Pseudagenia blaisdelli, California, Fox, p. 171, Ent. News n. sp.
- Rhopalum clavipes, habits; Verhoeff, Zool. Jahrb. Syst. vi, pp. 731.
- Salius spectabilis, p. 151, scarlatinosus, p. 153, Russia, MORAWITZ, Ent. Ross. xxvi; S. (Hemisalius) albistylus, p. 315, pl. xxv,
  - S. (Schistosalius) atristylus, p. 318, pl. xxiv, fig. 3, Madag
  - S. cyanirentria, Zanzibar, p. 822, bidens, Madagascar, p. pl. xxiii, fig. 25, SAUSSURE in Grandidier, Madagascar, xx, r
- n. spp.

  Schistonyx gryps, pl. xxv, fig. 15. albiculus, fig. 14, Madagascar, Saus
- p. 393, in Grandidier, Madagascar, xx, pt. 1, n. spp.
- Schistosalius, n. subg. of Salius, q.r. supra, BAUSSURE.
- Scolia bengalensis, Poons, CAMERON, p. 104, Mem. Soc. Manch. S. bhamoensis, p. 239, burmanica, p. 242, kirbyi, p. 244, Burma, RETTI, Ann. Mus. Genov. (2) xii: n. spp.
- Solierella spinola, p. 212, chilensis, p. 213, Chili, Kohl, Ann. Hofmi Wien, vii, n. spp.
- Sphærophthalma anthracina, California, Fox, p. 172, Ent. News, i simoni, p. 56, uncifera, fraterna, crocata, p. 57, Venezuela, Bu Ann. Soc. Ent. Fr. lxi, n. spp.
- Sphecius speciosus, habits; RILEY, Ins. Life, iv, pp. 248-252, figg.: and cocoon, peculiarities; RILEY, p. 170, P. E. Soc. Wash. ii.
- S. freyi, Madagascar, HANDLIRSCH, p. 27, pl. vii, fig. 5, Jahreal Trencsin, xiii, n. sp.
- Sphex malagassus, SAUSSURE, p. 427, pl. xviii, fig. 38, in Gran Madagascar, xx, pt. 1; S. (Priononyx) ferrugineus, Californis p. 170, Ent. News, iii: n. spp.
- Stenagenia, n. g. (Pompilides), p. 338, for S. pedunculata, p. 339, pl fig. 6, consobrina, p. 341, fig. 7, n. spp., Madagascar; Sai in Grandidier, Madagascar, xx, pt. 1.
- Stigmus inordinatus, p. 322, fulvipes, p. 324, N. America, Fox, T Ent. Soc. xix; S. temporalis, Guatemala, Kohl, p. 204, Ann museum Wien, vii: n. spp.
- Stizolarra chichimeca, Mexico, histrio, Zanzibar, p. 466, Saussi Grandidier, Madagascar, xx, pt. 1, n. spp.

- Stizus: monograph; HANDLIRSCH, SB. Ak. Wien, ci, pp. 25-205, pls. i-iii.
- S. magrettii, Oriental Region, p. 43, socius, Amboyna, p. 44, proximus, Oriental Region, p. 45, pusillus, Indo-Malay, semperi, Philippines, p. 46, insularis, Celebes, p. 47, facialis, Aru, versicolor, Oriental Region, p. 48, javanus, Java, p. 50, egens, p. 51, an'ipodum, p. 52, signatus, p. 53, Australia, anthracinus, New Guinea, p. 54, excisus, Brazil, p. 59, bolivari, Amazons, p. 60, nanus, Georgia, consobrinus, Rio Grande do Sul, p. 61, mexicanus, p. 66, guttulatus, p. 67, Mexico, zanthochrous, Texas, p. 69, kotschyi, E. Africa, p. 71, discolor, Algeria, p. 78, hoplites, Caffraria, p. 84, mayri, p. 85, gazagnairei, p. 86, Algeria, gynandromorphus, Syria, p. 89, inermis, Fiji, p. 91, gracilicornis, Caucasus, p. 93, clavicornis, Caffraria, p. 95, calopteryz, E. India, p. 110, pacilopterus, Africa, p. 111, chryssorrhaus, hab.?, p. 122, gracilipes, Caucasus, p. 131, melanurus, Senegal, p. 132, tricolor, Syria, p. 133, niloticus, Egypt, p. 134, hugelii, E. India, p. 139, pygidialis, Bhodes, p. 140, imperialis, Caffraria, p. 141, hyalipennis, Sinai, p. 142, marthæ, Biskra, p. 143, marnonis, Centr. Africa, p. 145, suchromus, Caucasus, p. 153, arenarum, N. Africa, p. 154, dewitzii, S. Africa, p. 156, rapax, Egypt, p. 157, HANDLIRSCH, SB. Ak. Wien, ci, n. spp.
- Tachyrhostus claviger, Australia, Kohl, p. 229, Ann. Hofmuseum Wien, vii, n. sp.
- Tachysphex algira, Oran, austriacus, Vienna, p. 215, T.? integer, Araxes Valley, p. 216, vestitus, Algeria, consocius, Caucasus, p. 217, simonyi, Canary Is., p. 218, Kohl, Ann. Hofmuseum Wien, vii; T. micromegas, p. 481, pl. xxvii, fig. 2, dimidiatus, fig. 3, p. 483, Madagascar, Saussure in Grandidier, Madagascar, xx, pt. 1: n. spp.
- Tachytes: monograph of the N. American; Fox, Tr. Am. Ent. Soc. xix, pp. 234-252. T. cœlebs is & T. abdominalis, T. mandibularis, habits; PATTON, p. 90, Ent. News, iii. T. aurulenta, = (Liris coxalis, Patt.); Fox, p. 138, Ent. News, iii.
- T. oviventris, Madagascar, SAUSSURE, p. 478, in Grandidier, Madagascar, xx, pt. 1, n. sp.
- T. calcaratus, p. 239, prædutor, p. 240, columbiæ, p. 241, spatulatus, p. 243, contractus, p. 245, minimus, p. 248, parvus, p. 249, obductus, mergus, p. 250, N. America, Fox, Tr. Am. Ent. Soc. xix; T. callosa, Brazil, Kohl, p. 218, Ann. Hofmuseum Wien, vii: n. spp.
- Trypoxylon figulus, habits; Verhoeff, Zool. Jahrb. Syst. vi, pp. 709 & 725-730.
- T. scutifrons, p. 523, pl. xxv, fig. 18, hova, fig. 17, p. 525, Madagascar, SAUSSURE in Grandidier, Madagascar, xx, pt. 1, n. spp.
- Tiphia fumipennis, sexes described, MAGRETTI, p. 248, Ann. Mus. Genov. (2) xii; T. picta, Bulgaris, Schulthess-Rechberg, p. 384, MT. Schw. ent. Ges. viii; T. policarinata, p. 250, vicina, p. 251, lyrata, p. 252, rostrata, p. 253, lævigata, cordata, p. 254, nodata, trapesata, p. 255, Burms, Magretti, Ann. Mus. Genov. (2) xii: n. spp.

### FORMICIDE.

- Account of the nests of ants; FOREL, Neujahrabl. Naturf. Ges. xcv. Organs of hearing in ants; JANET, p. ccxlvii, Bull. Soc. Ent. Fr. lxi. Classification of *Myrmicida*; FOREL, p. 143, in Grandidier, Madagas xx, pt 2.
- Classification of the Attini and Cryptocerini discussed, with suggesti as to their grouping; FOREL, pp. 344 & 345, MT. Schw. Ges. viii.
- WROUGHTON gives a list of the ants of India, amounting to 212 speand including a great deal of information as to habits and int gence; J. Bomb. Soc. vii, pp. 13, etc.
- Complete systematic treatise on the ants of Madagascar; FOREL (287 Sense organs in antenns; Sergi (840).
- New species of ants in amber; EMERY (249), and of. Palssontology, p Speculation on the origin of the present ant fauna of EUR EMERY, Mem. Acc. Bologn. (5) i, pp. 156-162, and Bull. Soc. V xxvii, p. 258.
- Symbiosis of ants and plants; WARBURG (982).
- Symbiosis of ants and plants in Somaliland; Keller (475).
- The behaviour of various ants in connection with Lomechusa; MANN (987).
- The ants of Belgium, with brief characters; Lameere, Ann. Ent. xxxvi, p. 61.
- Notes on the habits and sociology of ants; KATURIC (473, 474).
- New names proposed for numerous known species; Dalla Te Wien. ent. Z. xi, pp. 89-92.
- Acantholepis: synopsis of the species and varr.; Forel, pp. 41-43, Ent. Belg. xxxvi.
- A. carbonaria, Somaliland, EMERY, p. 119, Ann. Mus. Genov. (2) x gracilicornis, Aden, simplex, Somaliland, p. 42 (more fully desc MT. Schw. ent. Ges. viii, pp. 350-352), opaca, with pulchella, E. India, p. 43, FOREL, Ann. Ent. Belg. xxxvi: n. spp.
- Acanthomyrmex, n. g., for part of Pheidole; EMERY, p. cclxxvi, Soc. Ent. Fr. lxi.
- Acanthoponera brownii, New Zealand, FOREL, p. 335, MT. Schw Ges. viii, n. sp.
- Aeromyrma, n. g., near Pheidologeton, p. 198, for A. nosindambo, fig. 1, p. 199; FOREL in Grandidier, Madagascar, xx, pt. 2.
- Amblyopone gheorghieffi, Bulgaria, Forel, p. 309, Verh. z.-b. Wien A. saundersi, New Zealand, Forel, p. 336, MT. Schw. ent viii: n. spp.
- Ancuretus, n. g., between Ponerides and Dolichoderides (for A. n. sp., from Ceylon, not described); EMERY, p. cclxxv, Bul Ent. Fr. lxi.
- Anochetus grandidieri, Madagascar, FOREL, p. 108, pl. iii, fig. Grandidier, Madagascar, xx, pt. 2, n. sp.

- Anomma arcens, large worker from Gaboon, EMERY, p. liv, Bull. Soc Ent. Fr. lxi.
- Aphænogaster (Ischnomyrmex) swammerdami, redescribed and figured, p. 167, pl. iv, fig. 14; FOREL in Grandidier, Madagascar, xx, pt. 2.

  A. barbara, n. st. sordida, p. 452, striola, n. var., lævior, hispanica worker, p. 453; FOREL, Ann. Ent. Belg. xxxvi.
- Apterostigma soasmannii, p. 345, molleri, p. 348, S. Brazil, FOREL, MT. Schw. ent. Ges. viii, n. spp.
- Asteca instabilis, hermaphrodite described; Forel, pp. 268-270, pl. xvi, Bull. Soc. Vaud, xxviii.
- Bothroponera wasmannii, figured, pl. iv, fig. 4, comorensis, p. iv, fig. 5; Forel in Grandidier, Madagascar, xx, pt. 2.
- B. perroti, p. 131, pl. iv, fig. 6, cambouei, p. 133, pl. iv, fig. 7, Madagascar, FOREL in Grandidier, Madagascar, xx, pt. 2; B. gabonensis, W. Africa, Andre, p. 50, Rev. d'Ent. xi: n. spp.
- Camponotus is dealt with in Part I of "Systematic Revision of the Ants of India;" FOREL, J. Bomb. Soc. vii, pp. 219-245; C. lateralis, n. var. rectus; FOREL, p. 306, Verh. z.-b. Wien, xlii. C. micans, n. st. albisectus; EMERY, p. 120, Anu. Mus. Genov. (2) xii. C. foraminosus, C. robecchii, infrà, is a form of this, C. ruspolii, n. st. described; FOREL, p. 350, MT. Schw. ent. Ges, viii. C. maculatus, variation discussed, pp. 23-36, radamæ, n. st., p. 31, pl. i, fig. 4, boivini, p. 34, hova, figured, fig. 5, C. grandidieri, figured, fig. 6, described, p. 39, darwinii, p. 43, pl. ii, fig. 1, ursus, p. 46, pl. ii, fig. 6, edmondi, p. 49, pl. i, fig. 7, robustus, p. 54, pl. i, fig. 10, quadrimaculatus, p. 58, pl. ii, fig. 3, lubbocki, p. 60, pl. i, fig. 11, with n. st. christoides, p. 61, kelleri, pl. ii, fig. 4, christi, pl. i, fig. 12, p. 63, with n. sp. foersteri, p. 64, noticed and figured; FOREL in Grandidier, Madagascar, xx, pt. 2. C. mistura, n. st. fornaronis, p. 232, rufo-glaucus, n. st. dolendus, p. 238, maculatus, n. stt. taylori, p. 241, infuscus, p. 242, junctus, p. 243; Forel, J. Bomb. N. H. Soc. vii. C. cameroni, n. n. for mayri, Cam., nec. For.; Forel, p. 256, Ann. Ent. Belg. xxxvi. C. buchneri, worker noticed; André, p. 45, Rev. d'Ent. xi.
- C. robecchii, Somaliland, EMERY, p. 120, Ann. Mus. Genov. (2) xii; C. dufouri, p. 16, pl. i, fig. 2, ellioti, p. 37, echinoploides, p. 51, pl. i, fig. 8, sibreei, p. 53, pl. i, fig. 9, dromedarius, p. 65, pl. ii, fig. 5, nasica, p. 67, cambousi, p. 68, pl. i, fig. 13, & pl. ii, fig. 6, imitator, p. 209, pl. iv, fig. 15, concolor, p. 214, pl. vi, fig. 12, gibber, p. 215, fig. 13, pictipes, p. 217, Madagascar, Forel in Grandidier, Madagascar, xx, pt. 2; C. radiatus, p. 233, invidus, p. 234, lamarckii, p. 236, buddhæ, p. 238, Forel, J. Bomb. N. H. Soc. vii; C. ectatomnoides, New Zealand, Forel, p. 333, MT. Schw. ent. Ges. viii: n. spp.
- Cardiocondyla elegans, n. var. bulgarica; Forel, p. 312, Verh. z.-b. Wien, xlii. C. ergatoid, male of C. stambuloff; Forel, p. 461, Ann. Ent. Belg. xxxvi.

- C. stambulofi, Bulgaria, Forel, p. 810, Verh. s.-b. Wien, xlii; shuckardi, Madagascar, Forel in Grandidier, Madagascar, xx, pt. n. spp.
- Cataulacus oberthuri, Madagascar, EMERY in Grandidier, Madagascar, pt. 2, p. 146, pl. iv, fig. 9, n. sp.
- Cremastogaster scutellaris, n. var. christossitchii; Forel, p. 316, V z.-b. Wien, zlii. C. ranavalona, p. 184, pl. vi, figs. 3, 4, tric p. 187, fig. 5, degesri, p. 189, fig. 6, madagascariensis, p. 191, p fig. 7, redescribed and figured, C. inermis, n. st. secollii, p. pl. vi, fig. 9; Forel in Grandidier, Madagascar, xx, pt. 2. chiarinii, variation, 3 and 2 described; Forel, p. 353, MT. 8 ent. Ges. viii.
- C. acacia, p. 140, ruspolii, p. 142, Somaliland, FOREL, Zool. Ans. xv ferruginea, Somaliland, FOREL, p. 353, MT. Schw. ent. Ges. vii: striatulus, W. Africa, EMERY, p. liii, Bull. Soc. Ent. Fr. lxi schenki, p. 182, pl. vi, fig. 2, grevei, p. 183, fig. 3, rasoherina, p. kelleri, p. 197, emma, p. 227, pl. vi, fig. 11, Madagascar, FORI Grandidier, Madagascar, xx, pt. 2; C. biformis, Borneo, André, Mem. Soc. Zool. Fr. v: n. spp.
- Cryptopone, n. g., Ponerides, for Amblyopone testacea, Motsch.; El p. colxxv, Bull. Soc. Ent. Fr. lxi.
- Cylindromyrmex, Mayr, =(Holcoponera, Cam.); FOREL, p. 256, Ann. Belg. xxxvi.
- C. longiceps, Brazil, ANDRE, p. 47, Rev. d'Ent. xi, n. sp.
- Cyphomyrmes foxi, Jamaica, André, p. 56, Rev. d'Ent. xi, n. sp.
- Dimorphomyrmex, n. g., near Gesomyrmex, p. 49, for D. janeti, r Borneo, p. 51; André, Mem. Soc. Zool, v.
- Dorylus nigricans, agyptiacus, noticed; EMERY, p. liv, Bull. Soc. Ent. F Drepanognathus rugosus, worker described; MAYE, Term. füzete p. 127.
- Echinopla rugoza, Borneo, Andrá, p. 47, Mem. Soc. Zool. v, n. sp. Emeryia: this is the ergatold male of Cardiocondyla wroughtonii; F p. 461, Ann. Ent. Belg. xxxvi.
- Epoecus, n. g., near Anergates; EMERY, p. cclxxvi, Bull. Soc. Ent. F Euponera, n. subg. of Ponera; FOREL, p. 126, in Grandidier, Madaş xx, pt. 2.
- Eusphinctus, n. g., near Sphinctomyrmex, for E. furcatus, n. sp. described); Emery, p. colxxv, Bull. Soc. Ent. Fr. lxi.
- Forelius, n. g., for Iridomyrmex, mccookii; Emery, Z. wiss. Zool p. 389 (omitted in Zool. Rec. xxv).
- Formica nasuta, habits; FOREL, p. 307, Verb. z.-b. Wien, xlii.
- Gesomyrmex chaperi, Borneo, André, p. 47, Mem. Soc. Zon. sp.
- Lasius: embryology of normal and parthenogenetic eggs; Henki wiss. Zool. liv, pp. 137-140.
- Leptogenys falcigera, described and figured, p. 109, pl. iii, fig. 10; in Grandidier, Madagascar, xx, pt. 2.

- L. pavesii, Somaliland, EMERY, p. 111, Ann. Mus. Genov. (2) xii; L. incisa, Madagascar, p. 113, pl. iv, fig. 1, Forel in Grandidier, Madagascar, xx, pt. 2: n. spp.
- Leptothorax bulgaricus, Bulgaria, Forel, p. 314, Verh. s.-b. Wien, xlii; L. grisii, Canary Is., Forel, p. 456, Ann. Ent. Belg. xxxvi: n. spp.
- Liometopum microcephalum, habits; Forel, p. 308, Verh. z.-b. Wien, xlii: structure of nest; MAYE, t. c. p. 317.
- Lobopelta sulcinoda, Gaboon, André, p. 48, Rev. d'Ent. xi; L. oswaldi, p. 119, pl. iv, fig. 2, saussurei, pp. 121 & 218, jonesii, p. 219, Madagascar, Forel in Grandidier, Madagascar, xx, pt. 2: n. spp.
- Mayria madagascariensis, figured and noticed, p. 75, pl. ii, fig. 7; FOREL in Grandidier, Madagascar, xx, pt. 2.
- Meranoplus nanus, Gaboon, André, p. 55, Rev. d'Ent. xi; M. radama, Madagascar, Forel in Grandidier, Madagascar, xx, pt. 2, p. 148, pl. iv, fig. 10: n. spp.
- Monomorium gracillimum, n. st. robustior; FOREL, p. 352, MT. Schw. ent. Ges. viii.
- M. medinæ, Canaries, FOREL, p. 454, Ann. Ent. Belg. xxxvi; M. suteri, p. 340, smithii, p. 342, New Zealand, FOREL, MT. Schw. ent. Ges. viii: n. spp.
- Myrmicaria subcarinata, habits; ROTHNEY, P. E. Soc. 1892, p. v.
- Mystrium: characters of, pp. 114-116; M. mysticum, figured, pl. iii, fig. 11, Q described, p. 117; FOREL in Grandidier, Madagascar, xx, pt. 2.
- Ocymyrmex robecchii, Somaliland, p. 114, weitzeckeri, Basutoland, p. 116, EMERY, Ann. Mus. Genov. (2) xii, n. spp.
- Odontomachida: notes on the springing powers of; WASMANN, Wien. ent. Z. xi, pp. 316 & 317.
- Oligomyrmez grandidieri, Madagascar, FOREL in Grandidier, Madagascar, xx, pt. 2, p. 201, n. sp.
- Otomyrmez, n. subg. of Cataulacus; FOREL in Grandidier, Madagascar, xx, pt. 2, p. 147.
- Parasyscia imerinensis, Madagascar, FOREL in Grandidier, Madagascar, xx, pt. 2, p. 138, pl. iii, fig. 12, n. sp.
- Pheidole cameroni, Mayr., (= monticola, Cam.); Cameron, Ent. M. M. xxviii, p. 67.
- Pheidole longispinosa, p. 170, pl. v, fig. 1, ostoaldi, pp. 173 & 227, fig. 2, sikora, p. 223, veteratria, p. 225, Madagascar, Forel in Grandidier, Madagascar, xx, pt. 2; P.? variolosa, Somaliland, Emery, p. 113, Ann. Mus. Genov. (2) xii; P. risii, Buenos Ayres, Forel, p. 38, Ann. Ent. Belg. xxxvi: n. spp.
- Plagiolepis custodiens, n. varr. pilipes, hireuta, detrita; EMERY, p. 118, Ann. Mus. Genov. (2) xii.
- Polyrhacis concava, 2 described; ANDRÉ, p. 45, Rev. d'Ent. xi.
- P. paradoza, Gaboon, André, p. 46, Rev. d'Ent. xi; P. jerdonii, Ceylon, Forel, p. 17, Neujahrsbl. Naturf. Ges. xcv: n. spp.

- P. sakulava, pp. 124 & 222, pl. iv, fig. 3, P. (Euponera) sikoræ, p. 1 johannæ, p. 220, pl. v, fig. 11, elisæ, p. 221, pl. v, fig. 10, Madagas Forel in Grandidier, Madagascar, xx, pt. 2; P. annamita, FANDRÉ, p. 49, Rev. d'Ent. xi: n. spp.
- Prenolepis: characters of species from Madagascar and elsewhere, pp & 93, pl. ii, detail of & characters, &c., P. bourbonica, For., is a g species, p. 82, P. braueri, n. st. humbloti, p. 90, pl. ii, fig. 11; Fo in Grandidier, Madagascar, xx, pt. 2. P. imparis, Say, habits synonymy; PATTER, Am. Nat. xxvi, p. 871.
- P. ellisii, p. 87, pl. ii, fig. 10, glabra, p. 92, Madagascar, FOREL in Grandic Madagascar, xx, pt. 2: n. spp.
- Prolasius, n. subg. of Lasius, for Formica advena, Sm.; Forel, p. MT. Schw. ent. Ges. viii.
- Sima sahlbergii, redescribed and figured, p. 204, pl. v, fig. 4, n morondaviensis, p. 206; FOREL in Grandidier, Madagascar, pt. 2.
- S. spininoda, Gaboon, André, p. 51, Rev. d'Ent. xi; S. grandid pp. 203 & 229, pl. v, fig. 3, rakotonis, p. 206, pl. v, fig. 5, Madaga: Forel in Grandidier, Madagascar, xx, pt. 2: n. spp.
- Simopone, n. g., near Oylindromyrmez, p. 139, for S. grandidieri, n. Madagascar, p. 141, pl. iv, fig. 8, Forel in Grandidier, M gascar, xx, pt. 2.
- Stenomyrmex africanus, female described; André, p. 47, Rev. d' xi.
- Strumigenys antarctica, New Zealand, Forel, p. 338, MT. Schw. ent. viii, n. sp.
- Tapinoma flavidum, Borneo, André, p. 51, Mém. Soc. Zool n. sp.
- Technomyrmex mayri, Madagascar, FOREL, p. 99, pl. iii, fig. 6, in Gradier, Madagascar, xx, pt. 2, n. sp.
- Tetramorium cæspitum, n. st. depressum; Forel, p. 455, Ann. Ent. l xxxvi: and n. st. exasperatum; Emery, Expl. Sci. Tunis. Hym. r T. striatum, habits, formation of nests, &c.; Smith, Ent. M xxviii, p. 60. T. blochmannii, figured, pl. v, fig. 6, n. var., tanum, described, p. 153; Forel in Grandidier, Madaga xx, pt. 2.
- T. (Xiphomyrmax) kelleri, redescribed and figured, p. 153, pl. iv, fig.
  FOREL in Grandidier, Madagascar, xx, pt. 2.
- T. (X.) humbloti, p. 154, pl. iv, fig. 12, bessonii, p. 156, fig. 13, schauj p. 158, Madagascar, FOREL in Grandidier, Madagascar, xx, p n. spp.
- Tomognathus sublavis, note on; MEINERT, Ent. Med. iii, p. 205.
- Triglyphothrix: & described; Andre, p. 53, Rev. d'Ent. xi.
- T. gabonensis, W. Africa, ANDRE, p. 53, Rev. d'Ent. xi, n. sp.

#### TEREBRANTIA.

[Gf. Ashmead (17, 18, 19, 20, 21, 22), Berthoumieu (74), Bignell (82), Brischke (126, 127), Brodie (129), Buysson (11, 148), Buysson & Marshall (149), Carpentier (160), Cunningham (195), Fox (298), Froggatt (311), Fyles (314), Gillette (344), Harrington (388), Howard (432, 433, 434, 435, 436), Kohl (501), Kriechbaumer (523, 524, 525, 526), Kulagin (528), Lesne (548), Marshall (583), Mocsáry (615), Nerén (631), Riley (747), Saussure (210), Semenow (827, 828, 829), Stadelmann (883), Thomson (927), Townsend (949), Trail (950), Verhoeff (964).]

Hymenopterous parasites of spiders; Howard (435).

Spider parasites, list of; HASSELT, Tijdschr. Ent. xxxv, pp. xxxviii-xli.

#### CHRYSIDIDÆ.

Catalogue of the family; DALLA TORRE (198).

Chrysididæ of Valais described, with figures of terminal segments; FREY-GESNER, Bull. Soc. Murith. xvi, pp. 44-112, pl. iii.

Achrysis, n. subg. of Pseudochrysis; SEMENOW, p. 486, Hor. Ent. Ross.

Chrysis scutellaris, synonymy; BECKERS, p. 318, Wien. ent. Z. xi.

- C. (Gonochrysis) tekensis, p. 75, C. (Dichrysis) bispina, p. 77, C. (Tetrachrysis) gracilicornis, p. 78, Transcaspian Region, schalfewi, Chinese Turkestan, p. 80, chrysophora, Transcaspian Region, p. 81, buchneri, E. Siberia, p. 83, strauchi, p. 85, pleskei, p. 87, Chinese Turkestan, leptoposcila, Montenegro, p. 88, bianchii, Transcaspian Region, p. 90, grumorum, Amdo, p. 92, C. (Hexachrysis) herzensteini, Persia, p. 94, Semenow, Bull. Pétersb. (n.s.) iii, n. spp.
- Cleptes flammifer, S. Europe, p. 498, obsoletus, Sarepta, p. 500, buyssonis, Montenegro, p. 501, mocsarii, Hungary, p. 502, Semenow, Bull. Pétersb. (n.s.) ii, n. spp.
- Ellampus scutellaris, Panz., = (panzeri, Fab.); Dalla Torre, p. 205, Wien. ent. Z. xi. E. tournieri, n. n. for (Notozus viridis, Tourn.); Dalla Torre, p. 93, t. c. E. auratus, habits; Verhoeff, Zool. Jahrb. Syst. vi, p. 740.
- E. testaceicornis, E. Russia, Buysson in André, Spec. Hym. vi; E. heros, China, Semenow, p. 71, Bull. Pétersb. (n.s.) iii: n. spp.
- Hedychridium jakolewi, Transcaspian Region, Senenow, Bull. Pétersb. (n.s.) iii, n. sp.
- Hedychrum collare, Sarepta, Semenow, p. 73, Bull. Pétersb. (n.s.) iii, n. sp. Notosus konosoi, p. 102, particeps, p. 105, Mecklenburg, putoni, France, p. 108, Buysson in André, Spec. Hym. vi, n. spp.
- Pseudochrysis and allies, characters discussed and tabulated; SEMENOW, pp. 480-491, Hor. Ent. Ross. xxvi.
- Spintharina, n. subg. of Pseudochrysis; Semenow, p. 485, Hor. Ent. Ross. xxvi.

#### ICHNEUMONIDAL

- On the relations of primary and secondary parasites; CARPENTIER (16 On the existence of poison-glands in *Ichneumonides*; BUYSSON, R d'Ent. xi, pp. 257 & 258, pl. i.
- Agrypon stenostigma, Scandinavia, Thomson, Opusc. ent. xvi, p. 17 n. sp.
- Amblyteles speciosus and I. apricus, note on; KRIECHBAUMER, p. 2 Ent. Nachr. xviii.
- A. flavolatus, Caucasus, p. 41, inermis, Switzerland, bicuspis, Sav frustrator, Basses Alpes, p. 42, angustus, radosskowskii, capita Askhabad, adventor, Tunis, p. 43, Berthoumieu, Rev. d'Ent. n. spp.
- Anomalon: Thomson discusses the allied forms, Opusc. ent. xvi, 1757-1772.
- A. lapponicum, p. 1763, orbitale, annulitarse, clavipennis, p. 1764, Scalnavia, A. (Barylypa) laticeps, p. 1766, genalis, p. 1767, Scandina Thomson, Opusc, ent. xvi, n. spp.
- Astiphromma pectoralis, Texas, Ashmead, p. 107, Ent. News, n. sp.
- Bassus clypearis, Prussia, BRISCHKE, p. 43, Schr. Ges. Danz. (n.s.) n. sp.
- Blaptocampus, n. subg., for Anomalon nigricorns, and perspicus Thomson, Opusc. ent. xvi, p. 1765.
- Coencryptus bimaculatus, oviposition and larvas; Verhoeff, Zool. Ja Syst. vi, pp. 692-696.
- Campoplex lateralis, Prussia, BRISCHEE, p. 44, Schr. Ges. Danz. (n.s.) n. sp.
- Catadelphus pestrei, France, BERTHOUMIEU, Rev. d'Ent. xi, pp. 45 n. sp.
- Distantella, n. g., Cryptinæ, p. 229, for D. trinotata, n. sp., Transp. 230; SAUSSURE in Distant, Naturalist Transvaal.
- Ephialtes divinator, mediator, habits; VERHORFF, pp. 741-744, 2 Jahrb. Syst. vi.
- Erromenus defectivus, Prussia, BRISCHKE, p. 41, Schr. Ges. Danz. (viii, n. sp.
- Gnathowys tristis, Prussia, BRISCHKE, p. 31, Schr. Ges. Dans. (viii, n. sp.
- Hemipimpla, n. g., for H. oaffra, p. 227, calliptera, p. 228, n. app., Ti vaal; SAUSSURE in Distant, Naturalist Transvaal.
- Hemiteles subimpressus, p. 48, coxalis, rufus, p. 49, Prussia, BRISCHKE, S. Ges. Danz. (n.s.) viii; H. prosthesimæ, p. 294, micariroru, p. drassi, p. 296, N. America, Howard, P. E. Soc. Wash. ii; H. mius, p. 63, pothinus, p. 64, Venezuela, Marshall, Ann. Soc. Fr. lxi: n. spp.
- Hepiopelmus leucostigma, n. var., 4-guttata; Kriechbaumer, p. 202, Nachr. xviii.

- H. apicalis, Prussia, BRISCHKE, p. 30, Schr. Ges. Danz. (n.s.) viii, n. sp.
- Ichneumon molitorius and deliratorius, synonymy discussed; KRIECH-BAUMER, p. 39, Ent. Nachr. xviii. I. rubens, & described, p. 196, Q var., p. 198, alpicola, var. &, p. 198; KRIECHBAUMER, t. c. I. gerstaeckeri, erythromerus, croceipes, dissimilis, præstigiator, deletus, notes on; KRIECHBAUMER, pp. 292-295, t. c. I. firmipes, Wesm., note on; NEBÉN, Ent. Tidskr. xiii, p. 67.
- I. singularis, Transcaspian, scopulator, Oural Mts., evanidus, p. 37, paganus, erraticus, France, cintranus, Estremadura, p. 38, delphinas, buyssoni, bifossatus, p. 39, trifarius, France, levicoxa, Transcaspian, canescens, p. 40, bifarius, p. 41, France, Berthoumieu, Rev. d'Ent. xi; I. szechenyii, China, Frivaldezky, p. 126, Term. füzetek, xv; I. discoensis, W. Greenland, Fox, P. Ac. Philad. 1892, p. 134: n. spp.
- Ischnocryptus, n. subg. of Phygadenon, q.v.; KRIECHBAUMER, p. 351, Ent. Nachr. xviii.
- Ischyrocnemis ottawaensis, Canada, Harrington, p. 98, Canad. Ent. xxiv, n. sp.
- Leptocryptus albomarginatus, p. 371, bellulus, p. 372, rubens, p. 373, Germany, Kriechbaumen, Ent. Nachr. xviii, n. spp.
- Limneria consumtor, & described; BRISCHKE, p. 21, Schr. Ges. Danz. (n.s.) viii.
- Lochetica (Frst.), n. g., for Phygadenon pimplarius; KRIECHBAUMER, p. 340. Ent. Nachr. xviii.
- Mesochorus splendidulus, larval peculiarities and development; Kulagin (528).
- Mesoleius (? Spudæa) similis, Prussia, BRISCHKE, p. 20, Schr. Ges. Danz. (n.s.) viii; M. bipunctatus, albopictus, p. 34, affinis, similis, p. 35, longicornis, p. 36, Prussia, BRISCHKE, Schr. Ges. Danz. (n.s.) viii: n. spp.
- Mesoleptus nigricornis, Prov., referred to Ctenopelma; HARRINGTON, p. 99, Canad. Ent. xxiv.
- M. (Hypocryptus) testaceicornis, Prussia, BRISCHKE, p. 33, Schr. Ges. Danz. (n.s.) viii, n. sp.
- Mesostenus simonis, Venezuela, Marshall, p. 61, Ann. Soc. Ent. Fr. lxi, n. sp.
- Microcryptus amænus, Worms, KRIECHBAUMEB, p. 362, Ent. Nachr. zviii, n. sp.
- Nemeritis raphidiæ, Munich, KRIECHBAUMBB, p. 234, Ent. Nachr. xviii, n. sp.
- Notopygus americana, Canada, HARRINGTON, p. 98, Canad. Ent. xxiv, n. sp.
- Oneista bohemani, redescribed; KRIECHBAUMER, p. 41, Ent. Nachr. xviii.
- Ophion wustneii, Sonderburg, p. 232, slaviceki, Moravia, p. 233, KRIECH-BAUMER, Ent. Nachr. xviii, n. spp.

- Orthocontrus palpalis, Prussia, BRISCHKE, Schr. Ges. Danz. (n.s.) v p. 43, n. sp.
- Periliseus (Rhæstes) testaceipes, Prussia, BRISCHKE, p. 19, Schr. G Danz. (n.s.) viii; P. nigropunetatus, Prussia, BRISCHKE, p. 37, Sc Ges. Danz. (n.s.) viii; n. spp.
- Perosis albopicta, p. 214, gracilis, p. 216, Germany, KRIECHBAUMER, E Nachr. xviii, n. spp.
- Pezomachus micariæ, Massachusetts, Howard, p. 294, P. E. Soc. Wii, n. sp.
- Phæstus? sericeus, Prussia, BRISCHKE, p. 39, Schr. Ges. Danz. (n.s.) v n. sp.
- Phygadeuon cretatus, p. 363, diaphanus, p. 364, notes on; Krie Baumer, Ent. Nachr. xviii.
- P. (Ischnocryptus) geniculatus, p. 343, forticornis, clotho, p. 344, lach. p. 345, atropos, p. 346, Bavaria, Kriechbaumer, Ent. Naxviii; P. caracacensis, Venezuela, Marshall, Ann. Soc. Ent. Fr. p. 62: n. spp.
- Pimpla instigator, bred from Trichiosoma; BRISCHKE, p. 57, Schr. (Dans. (n.s.) viii.
- P. ellopiæ, Brit. Columbia, HARRINGTON, p. 99, Canad. Ent. xxiv argiopes, p. 65, simonis, p. 67, Caracas, Marshall, Ann. Soc. Ent. lxi: n. spp.
- Platylabus uranius, & described; Thomson, p. 1773, Opusc. xvi.
- Plesiopthalmus paniscoides, Massachusetts, Ashmead, p. 107, Ent. N iii, n. sp.
- Polyblastus (Scopiorus) lucidus, p. 40, albiventris, p. 41, Prussia, BRISCI Schr. Ges. Danz. (n.s.) viii, n. spp.
- Polysphincta (Zatytopa) strigis, p. 291, P. theridii, p. 292, P. (Zaglyzkoebelei, p. 293, N. America, Howard, P. E. Soc. Wash. ii dichroa, Venezuela, Marshall, p. 67, Ann. Soc. Ent. Fr. n spp.
- Prionopoda canadensis, Ottawa, HARRINGTON, p. 98, Canad. Ent. 3 n. sp.
- Pseudacænites, n. g., Pimplidarum, for P. moravicus, n. sp., Mäh Kriechbaumer, p. 219, Ent. Nachr. xviii.
- Psilomastax cyaneus, Sardinia (from pupa of Papilio hospiton), KRI BAUMER, p. 101, Ent. Nachr. xviii; P. violaceus, = (cyaneus, Kriec Mocsáry, Ent. Nachr. xviii, p. 208: n. spp.
- Rhimphalia? pilosa, Prussia, BRISCHKE, p. 39, Schr. Ges. Danz. (n.s.)
  n. sp.
- Tryphon mesoxanthus and punctus, note on; KRIECHBAUMER, p. 203, Nachr. xviii. T. punctus, habits, synonymy, characters; S1 pp. 102-105, Ent. Nachr. xviii.
- T. grossus, Prussia, BRISCHKE, p. 38, Schr. Ges. Danz. (n.s.) viii, n. Udenia, Förster, characters given; KRIECHBAUMER, p. 40, Ent. N xviii.

# BRACONIDE.

- List of Virginian hosts of Braconidæ; HOPKINS, Ins. Life, iv, pp. 256-259. Aleiodes ungularis, p. 1677, punctipes, p. 1678, arcticus, borealis, p. 1679, crassipes, p. 1681, Scandinavia, Thomson, Opusc. ent. xvi, n. spp.
- Apanteles: habits and distinction by secondary parasites; CARPENTIER, Bull. Soc. L. Nord France, x, pp. 35-40: synonymy of the American species; PATTON, Psyche, vi, p. 260.
- Arrhaphis americana, minuta, Florida, Ashmead, Psyche, vi, p. 289, n. spp. Aspicolpus borealis, Scandinavia, Thomson, p. 1726, Opusc. ent. xvi, n. sp. Ascogaster fuscipennis, leptopus, p. 1718, clypealis, nigricornis, p. 1719,

cynipum, sternalis, cavifrons, gibbiscuta, p. 1720, graniger, p. 1721, Scandinavia, Thomson, Opusc. ent. xvi, n. spp.

- Blacus spinifer, Scandinavia, THOMSON, p. 1735, Opusc. ent. xvi, n. sp.
- Bracon furtivus, Canada, Fyles, p. 34, Canad. Ent. xxiv; B. hamobaphes, Venezuela, Marshall, p. 68, Ann. Soc. Ent. Fr. lxi: n. spp.
- Calyptus mucronatus, p. 1703, strigator, truncatus, p. 1704, lapponicus, macrurus, p. 1705, Scandinavia, Thomson, Opusc. ent. xvi, n. spp.
- Chelonus seticornis, Scandinavia, THOMSON, p. 1714, Opusc. ent. xvi, n. sp. Clinocentrus tenuicornis, p. 1687, petiolaris, striolatus, p. 1688, gracilipes,
- Clinocentrus tenuscornis, p. 1687, petiolaris, striolatus, p. 1688, graculipes, breviculcar, p. 1689, Scandinavia, Thomson, Opusc. ent. xvi, n. spp.
- Colastes caudatus, N. Europe, Thomson, p. 1691, Opusc. ent. xvi, n. sp. Euphorus grandiceps, brevispina, facialis, p. 1747, microcerus, p. 1748, rubricollis, obscuripes, p. 1749, fulviceps, arenicola, p. 1751, Scandinavia, Thomson, Opusc. ent. xvi, n. spp.
- Exothecus foveolator, p. 1698, lævis, flavitarsis, lapponicus, pubicornis, p. 1699, laticarpus, flaviventris, p. 1700, Scandinavia, Thomson, Opusc. ent. xvi, n. spp.
- Helcon femoralis, Scandinavia, Thomson, p. 1725, Opusc. ent. xvi, n. sp. Microgaster glomeratus, particulars as to development of embryo and larva; Kulagin (528). M. sp. parasitic on imago of Eleodes; Riley (747).
- Pambolus bifusciatus, California, ASHMEAD, Psyche, vi, p. 290, n. sp.
- Perilitus omophli, Algeria, with metamorphoses, LESNE, Ann. Soc. Ent. Fr. lxi, pp. 305 & 308, pl. v; P. gracilipes, p. 1742, caudatus, flaviventris, areolatus, p. 1743, brevipetiolatus, p. 1744, facialis, borealis, p. 1745, Scandinavia, Thomson, Opusc. ent. xvi: n. spp.
- Phanomeris glabricollis, Scandinavia, Thomson, p. 1696, Opusc. ent. xvi, n. sp.
- Rhogas medianus, p. 1668, rufipes, p. 1669, alpinus, p. 1671, flavipalpis, hirtus, p. 1672, Scandinavia, THOMSON, Opusc. ent. xvi; R. xanthus, Caracas, Marshall, p. 69, Ann. Soc. Ent. Fr. lxi: n. spp.
- Rhysipolis obscuripes, varicoxa, Scandinavia, Thomson, p. 1692, Opusc. ent. xvi, n. spp.
- Sigalphus antennalis, p. 1706, parvus, p. 1707, breviventris, rimulosus, p. 1708, ambiguus, opacus, nigripes, p. 1709, crussiceps, p. 1710, Scandinavia, Thomson, Opusc. ent. xvi : n. spp.

# EVANIIDE, STEPHANIIDE, MONONACHIDE, TRIGONALIDE.

- Aulacus sibiricola, Centr. Siberia, p. 27, morawitzi, Transcaspian Rep. 29, Semenow, Bull. Pétersb. (n.s.) iii, n. spp.
- Evania schlettereri, Araxes Valley, Kohl., p. 231, Ann. Hofmus Wien, vii; E. cribrata, hab. ?, Semenow, p. 10, Bull. Pétersb. (iii: n. spp.
- Gasteryption, intermedium, p. 13, fallaciosum, p. 14, dubiosum, obsolinsidiosum, p. 15, foveiceps, p. 17, schenoyrevoi, p. 19, Russia, con Turkestan, p. 20, dimidiatum, E. Siberia, p. 22, dilutum, Ch Turkestan, p. 23, sibiricum, Central Siberia, p. 24, fortic Andalusia, p. 26, Semenow, Bull. Pétersb. (u.s.) iii: n. spp.
- Monomachus viridis, Brazil, Stadelmann, p. 237, Berl. Ent. Zeit. xz pl. vi, fig. 5, n. sp.
- Stephanus martini, Sumatra, Stadelmann, Berl. Ent. Zeit. xxxvii, p pl. vi, fig. 4, n. sp.
- Trigonalis simoni, Caracas, Buysson, p. 59, Ann. Soc. Ent. Fr. lxi, n

## CHALCIDIDE.

- The biology of Chalcidida, Howard, pp. 567-588, P. U. S. Nat.
- Mode of destruction of Coecidæ by parasites; Howard, p. 351, Soc. Wash. ii.
- Odoriferous Chalcidida, HOWARD, P. E. Soc. Wash. ii, p. 215.
- Blastophaga: fertilisation of Ficus roxburghii by; CUNNINGHAM

  B. psenes: its value as caprifying agent; RILEY, P. Biol.

  Washington, vii, p. 99.
- Calocerinus, n. g., for Tetracnemus floridanus, Ashm.; HOWARD, | P. U. S. Nat. Mus. xv.
- Chalcura californica, California, Ashmead, p. 356, P. E. Soc. Win. sp.
- Cratæpus fletcherii, Canada, ASHMEAD, p. 309, Canad. Ent. xxiv, n. Elasmus: habits, E. varius. figured; Howard, Ins. Life, iv, p. 253 Eunotus lividus, N. America, ASHMEAD, P. E. Soc. Wash. ii, p. 288, Eupelmus piceus, p. 296, drassi, p. 297, N. America, Howard, P. I. Wash. ii, n. spp.
- Eurytoma rubicola, habits; VERHOEFF, Zool. Jahrb. Syst. vi, p. 73
  Holcopelte nitens, Washington, HOWARD, p. 298, P. E. Soc. Wash. ii
  Kapala terminalis, Cuba. ASHMEAD, p. 358, P. E. Soc. Wash. ii, n. :
  Lælups callisto, Venezuela, MARSHALL, p. 73, pl. iv, fig. 3, Ann. Soc.
  Fr. lxi, n. sp.
- Larradomorpha, n. g., near Polistomorpha, p. 238, for L. insignis, W. Africa, p. 239, pl. vi, fig. 7, STADELMANN, Berl. Ent. Zeit., Lophyrocera apicalis, California, ASHMEAD, p. 357, P. E. Soc. W. n. sp.

- Melittobia: the hosts and habits of the species discussed; HOWARD, P. E. Soc. Wash. ii, pp. 244-248.
- M. chalybii, N. America, ASHMEAD, p. 231, P. E. Soc. Wash. ii, n. sp.
- Mestocharis wilderi, N. America, Howard, p. 298, P. E. Soc. Wash. ii, n. sp.
- Metagea schwarzii, N. America, Ashmead, p. 356, P. E. Soc. Wash. ii, n. sp.
- Orasema occidentalis, California, Ashmead, p. 355, P. E. Soc. Wash. ii, n. sp.
- Pentacnemus, n. g., Encyrtinæ, for P. bucculatricis, n. sp., Missouri; Howard, p. 366, pl. xlvii, fig. 1, P. U. S. Nat. Mus. xv.
- Podagrion, notes on, with fig.; Ins. Life, iv, p. 243.
- Prionopelma basilica, Venezuela, MARSHALL, p. 71, pl. iv, fig. 2, Ann. Soc. Ent. Fr. lxi, n. sp.
- Development of Smicra clavipes in Stratiomys; Henneguy, C.R. exiv, p. 133.
- Tetracladia, n. g., Encyrtines, for T. texana, Texas, p. 367, pl. xlvi, fig. 1, gracilis, Florida, p. 368, n. spp., N. America; Howard, P. U. S. Nat. Mus. xv.
- Tetrastichus banksii, Washington, Howard, p. 299, P. E. Soc. Wash. ii, n. sp.

### PROCTOTRUPIDE.

- Pristocera rosmarus, Cameroons, Stadelmann, p. 237, pl. vi, fig. 6, Berl. Ent. Zeit. xxxvii, n. sp.
- Procacantha bæiformis, Caracas, MARSHALL, p. 75, pl. iv, fig. 5, Ann. Soc. Ent. Fr. lxi, n. sp.
- Scelio venezuelensis, Caracas, MARSHALL, p. 74, pl. iv, fig. 4, Ann. Soc. Ent. Fr. lxi, n. sp.
- Tanyzonus, n. g. (Belytidæ), for T. bolitophilæ, n. sp., New Zealand; MARSHALL, p. 276, Ent. M. M. xxviii: note on habits by Hudson, p. 277. P. 308, t. c., this is said to be the same as Betyla fulva, Cameron.

#### CYNIPIDE.

- Embryology in parthenogenetic eggs; Henking (396).
- Notes on synonymy, proposing new names; Dalla Torre, pp. 131 & 132, Wien. ent. Z. xi.
- Andricus hartigii, Foorst.: this name is valid; Cameron, Ent. M. M. xxviii, p. 67.
- A. cellularius, frequens, Colorado, GILLETTE, p. 247, Ent. News, iii, p. spp.
- Aulaz nabali, Toronto, BRODIE, p. 13, Canad. Ent. xxiv, n. sp.
- Cynips acaciondiscoloris, p. 153, acacialongifolia, p. 154, maideni, p. 155, Sydney, Froggatt, P. Linn. Soc. N.S.W. (2) vii, n. spp.

- Pediaspis acerina, Bremi., ==(sorbi, Tischb., and aceris, Foerst.); CAMERIEL. M. M. xxviii, p. 67.
- Rhodites rose, embryology of parthenogenetic eggs; Henking, Z. wi Zool. liv, pp. 142, &c., and oviposition, pp. 145 & 146.
- Spathegaster punctatus, England, BIGNELL, Ent. M. M. xxviii, p. 1 n. sp.

#### SESSILIVENTRES.

[Cf. Borries (111), Brischke (127), Buysson & Marshall (149), F (298), Gasperini (320), Jakowlew (453), Konow (510), Krief Baumer (522), Krieger (527), Nerén (631), Biley (74 Saussure (210), Semenow (828).]

# SIRICIDE and CEPHIDE.

- Cephus niger, Prussia, BRISCHKE, p. 53, Schr. Ges. Danz. (n.s.) vi C. carbonarius, p. 12, grombczewskii, p. 13, Turkestan, JAKOWL Hor. Ent. Ross. xxxvi: n. spp.
- Everemanella, n. g., for Cephus cruentatus, Everem.; JAKOWLEW, p. Hor, Ent. Ross, xxvi.
- Phyllacus sibiricola, p. 13, cylindrus, p. 14, Irkutsk, JAKOWLEW, Hor. I Ross. xxvi, n. spp.
- Xiphydria potanini, Gan-ssu, JAKOWLEW, p. 15, Hor. Ent. Ross. x n. sp.

# TENTHREDINIDE.

- Abia: table of characters of the subgenera and species; SEMENOW, 513-517, Bull. Pétersb. (n.s.) ii. A. nigricornis, larva noti-BRISCHKE, p. 51, Schr. Ges. Danz. (n.s.) viii.
- A. symballophthalma, W. Europe, Semenow, p. 510, Bull. Pétersb. (ii, n. sp.
- Allantus monozonus, & described; KRIECHBAUMER, p. 99, Ent. Na
- Amauronematus morawitsi, p. 24, jaroslawensis, Russia, nigridorsis, p. bicolor, Irkutsk, glacialis, Novaja Zemlja, p. 26, JAKOWLEW, Ent. Ross. xxvi, n. spp.
- Allantus potanini, p. 45, andreas, p. 47, dioctroides, p. 48, Gar superbus, p. 49, capucinus, p. 50, grombesewskii, p. 51, fi p. 52, heros, Turkestan, maculiger, p. 53, xylota, p. 54, Sil semenowi, Centr. Russia, mongolicus, Mongolia, p. 55, omiss. Siberia, p. 56, jakowlewi, Turkestan, p. 57, Jakowlew, Hor. Ross. xxvi, n. spp.
- Arge cingulata, Turkestan, p. 16, forficula, p. 17, potanini, p. 18, zowskii, subtilis, p. 19, zonata, p. 20, coriacea, p. 21, Gan-ssu, JALEW, Hor. Ent. Ross. xxvi, n. spp.

- Athalia bicolor, Transvaal, SAUSSURE, p. 226, in Distant, Naturalist Transvaal, n. sp.
- Blennocampa sibirica, Irkutsk, Jakowlew, p. 29, Hor. Ent. Ross. xxvi, n. sp.
- Cimber americana, dipterous parasites; Townsend, p. 126, Canad. Ent. xxiv, and Aldrich, t. c. pp. 144 & 145.
- Cladius pectinicornis, habits; RILEY, Ins. Life, v, pp. 6 & 7, figg.
- C. major, N. Italy, Cobelli, p. 70, Verh. z.-b. Wien, zlii (cf. SB. p. 69, t. c., where this is said to be C. crassicornis), n. sp.
- Dolerus grombczewskii, Turkestan, p. 31, pusillus, p. 32, variegatus, p. 33, Irkutsk, purus, Gan-ssu, p. 34, asceta, Thibet, p. 35, Jakowlew, Hor. Ent. Ross. xxvi, n. spp.
- Emphytus arcticus, Sweden, Neren, Ent. Tidskr. xiii, p. 57, n. sp.
- Hylotoma graeffei, pallipes, Triest, KRIECHBAUMBR, p. 38, Ent. Nachr. xviii, n. spp.
- Jermakia, n. g., for Allantus cephalotes, Jak.; JAKOWLEW, p. 58, Hor. Ent. Ross. xxvi.
- Lophyrus: observations on the Danish species and their metamorphoses; Borres, Ent. Med. iii, pp. 97-124, pl. ii. L. rufus, ravages and metamorphoses; LAMPA, Ent. Tidskr. xiii, pp. 41-44.
- Lyda carpini, Prussia, BRISCHKE, p. 53, Schr. Ges. Danz. (n.s.) viii; L. konowi, Centr. Russia, JAKOWLEW, p. 9, Hor. Ent. Ross. xxvi: n. spp.
- Lygeonemutus B. jakowlewi, Irkutsk, A. Jakowlew, p. 28, Hor. Ent. Ross. xxvi, n. sp.
- Macrophya rustica, hermaphrodite; KRIEGER, Z. Naturw. lxv, p. 138.
- M. dalmatina, Dalmatia, GASPERINI (320); M. soror, p. 43, potanini, p. 45, Gan-ssu, JAKOWLEW, Hor. Ent. Ross. xxvi: n. spp.
- Monophadnus athalioides, Gan-ssu, JAKOWLEW, p. 30, Hor. Ent. Ross. xxvi, n. sp.
- Nematus: remarks on species of and their parasites; Neren, Ent. Tidskr. xiii, pp. 61-67.
- N. insubricus, Milan, Cobelli, p. 70, Verh. z.-b. Wien. xlii (cf. p. 69, SB. t. c., where this is said to be N. cæruleocarpa, Htg.); N. borealis, West Greenland, Fox, P. Ac. Philad. 1892, p. 133: n. spp.
- Pachynematus quinquemontanus, Caucasus, gracilis, Turkestan, Jakowlew, p. 27, Hor. Ent. Ross. xxvi, n. spp.
- Pachyprotasis antennata, n. var. chinensis; JAKOWLEW, p. 43, Hor. Ent. Ross. xxvi.
- P. semenowi, p. 39, macrophyoides, p. 40, obscura, misera, p. 41, longicornis, p. 42, Gan-ssu, Jakowlew, Hor. Ent. Ross. xxvi, n. spp.
- Paccilosoma plana, Irkutsk, Jakowlew, p. 31, Hor. Ent. Ross. xxvi, n. sp.
- Priophorus hyalopterus, Irkutsk, JAKOWLEW, Hor. Ent. Ross. xxvi, n. sp.
- Pteronus schewyresoi, Caucasus, irkutensis, Siberia, p. 22, konosoi, Caucasus, balassagloi, Turkestan, p. 23, Jakowlew, Hor. Ent. Ross. xxvi, n. spp. 1892. [vol. xxix.]

- Rhogogustera robusta, Gan-ssu, Jakowlew, p. 38, Hor. Ent. Ross. x n. sp.
- Schizocera: analytic table of the European species; Konow, Wien. ent xi, pp. 11-18. S. privatus, metamorphoses; Marlatt, Ins. Life pp. 24-26, figg.
- S ballioni, Tiflis, p. 19, similis, p. 20, austriaca, Austria, humerata, iniolia, p. 21, Konow, Wien. ent. Z. xi, n. spp.
- Sciopteryx pusilla, Gan-ssu, JAKOWLEW, p. 35, Hor. Ent. Rosa z
- Sciandria bimaculata, N. Italy, Cobelli, p. 71, Verh. s.-b. Wien, slii p. 69, SB. t. c., where this is said to be S. coronata, Kl.), n. sp.
- Strongylogaster konowi, Centr. Russia, Jakowlew, p. 30, Hor. Ent. I xxvi, n. sp.
- Synairema simoni, Venezuela, Buysson, p. 59, Ann. Soc. Ent. Fr. n. sp.
- Tarpa borealis, Irkutsk, p. 10, victoriosa, Turkestan, p. 11, Jakow Hor. Ent. Ross. xxvi, n. spp.
- Tenthredo maura, F., synonymical note; KRIECHBAUMER, p. 100, Nachr. xviii.
- T. ruthena, Ural Mountains, pediculus, p. 59, stulta, p. 60, pulchra, p. Gan-ssu, JAKOWLEW, Hor. Ent. Ross. xxvi, n. spp.
- Tenthredopsis kokuswi, Centr. Russia, p. 36, ganssuensis, China, gra Irkutsk, p. 37, Jakowlew, Hor. Ent. Ross. xxvi, n. spp.
- Trichiosoma pubescens, E. Siberia, JAKOWLEW, p. 16, Hor. Ent. : xxvi, n. sp.

# (c.) LEPIDOPTERA.

Wing neuration; SPULER (881).

Lepidoptera to be divided into two suborders: Frenatz and Jugatæ; Comstock, P. Am. Ass. xli, p. 200.

Mode of development of wings after leaving the pupal skin; Petersen (675).

The peculiarities of the scale-pigments in numerous Lepidoptera; URECH (958).

Histology of metamorphosis; Bugnion (140).

Histology of alimentary canal of larvæ; HOLMGREN (411).

Value of appearance of sexes at different times; Petersen (676).

Lepidoptera reared in Zoological Gardens, London; Thomson, P. Z. S. 1892, pp. 193-195.

#### RHOPALOCERA.

[Cf. Alphebaky (779), Aurivillius (23), Baker (30), Brancsik (115), Butler (141), Butler & Rothschild (147), Cosmovici (191), Druce (227), Eckstein (245), Edwards (246, 247), Elwes (248), Hagen (383), Holland (406, 407, 408, 409, 410), Honrath (412), Hormuzaki (415), Karsch (460, 461, 462, 463), Leech (542), Mabille & Vuillot (573), Manders (579), Moore (622), Nicéville (638), Oberthur (650), Postans (701), Poujade (702), Rippon (755, 756), Robbe (766), Röber (767), Rogenhoper (777, 778), Rothschild (782, 783), Rühl (789), Scudder (813), Semper (835), Skinner (847, 848, 849), Smith (857), Smith & Kirby (858), Snellen (869, 870, 871, 874, 876), Spuler (882), Standfuss (884), Staudinger (779, 885, 886, 888, 889, 890, 892, 893, 894), Tryon (954), Vuillot (979), Weymer (1003), Wright (1016, 1017).]

Remarks on upwards of 100 species of Rhopalocera found in Southern Siberia; STAUDINGER, Deutsche e. Z. Lep. v, pp. 301, etc.

The Bhopalocera of Amurland discussed; STAUDINGER, pp. 83-219, Romanoff Mem. vi.

Remarks on Haase's system of Rhopalocera, Settz, S. E. Z. liii, pp. 157-159.

Obtaining eggs in captivity; WRIGHT, Canad. Ent. xxiv, pp. 70-74.

#### PAPILIONIDE.

Phylogeny of Papilionida; SPULER (882).

Cosmodesmus, n. subg. of Papilio; HAASE, Bibl. Zool. Heft. viii, p. 15.

Doritis apollinus, varr. described; STAUDINGER, p. 225, Deutsche e. Lep. iv.

Druryia antimachus, 9 noticed; WATKINS, Ent. xxv, p 143: 9 descrit STAUDINGER, p. 270, Deutsche e. Z. Lep. v.

Ornithoptera arruana, n. var. valentina; Vuillot, p. cxxiv, Bull. 1 Ent. Fr lxi. O. flavicollis, var., = (olympia, Honr.), figured, pl fig. 1; Honrath, p. 429, Berl. Ent. Zeit. xxxvi. O. (Priamopt lydius, O. (Pompeoptera) nereis, figured; Rippon, Ornithopt. Ico O. riedeli figured; Smith & Kirby, Rhop. ex. ii, Papilionida, Polionina, Ornithoptera, ii. O. zalmoxis, Q described; Stauding p. 268, Deutsche e. Z. Lep. v. O. priamus, sexes of var. from Ké described; Röber, pp. 263-268, Tijdschr. Ent. xxxiv. O. posein variation; Tryon (954).

O. andromache, N. E. Borneo, STAUDINGER, p. 393, Deutsche e. Z. I. v; O. sumæus, Aru Is., Rippon, p. 193, Ann. N. H. (6) x: also figu and described in Rippon, Ornithopt. Icones: n. spp.

Papilio: the following figured and noticed—P. (Menelaides) anna, pl. x fig. 2, phlegon, fig. 3, p. 271, P. (Atrophaneura) semperi, pl. xlvii, fi 1-3, p. 272, P. (Araminta) antonio, fig. 4, p. 274, P. (Pathysa) at phates, pl. xlviii; Semper, Reisen Philipp. ii, v. P machaon, varieti AUSTAUT, Le Nat. 1892. pp. 13, &c. P. machaon and xuthus, note red markings of hind wings of; Sanson, Ent. xxv, p. 43. P. and geos, n. var. depelchini; ROBBE, p. 125, Ann. Ent. Belg. xxxvi. alexanor, n. var. maccabæus; STAUDINGER, p. 224, Deutsche e. Z. Le iv. P. macareus, var. noticed and figured; NICEVILLE, p. 345, pl. fig. 1, J. Bomb. N. H. Soc. vii. P. neumoegeni, figured, pl. xv, fig. neptunus, n. var. fehri, p. 432; HONRATH, Berl. Ent. Zeit. xxxvi. schönbergianus, 2 described; Honrath, p. 441, Berl. Ent. Zeit. xxx P. anactus, figured, pl. xvi; Scott, Austral. Lep. ii, sarpedon, lyca pl. xvii. P. sisenna, figured; MABILLE & VUILLOT, pl. x, fig. 4, No Lep. P. gelon and var. described and figured; ROTHSCHILD, Tr. Soc. 1892, p. 141, pl. iv. P. antimachus, Q noticed, p. 162, figur pl. v, p. 189; WATKINS, Ent. M. M. xxviii. P. cresphontes, ea. stages; BEUTENMÜLLER, p. 75, Bull. Am. Mus. iv. P. asterias, ii of mixed sexes; EDWARDS, Canad. Ent. xxiv, p. 49.

P richelmanni, E. Africa, WEYMER, p. 98, S. E. Z. liii; P. policenoic W. Africa, Holland, p. 287, Ann. N. H. (6) x; P. catoris, N. Born Smith, p. 426, Ann. N. H. (6) x; P. (Achillides) discordia, Suma Nicéville, p. 343, pl. i, fig. 2, J. Bomb. N. H. Soc. vii; P. godma Alor, Röber, p. 271, Tijdschr. Ent. xxxiv (fig. in xxxv); P. hollan Colorado, Edwards, p. 50, Canad. Ent. xxiv; P. mathani, Up Amazons, Oberthur, Études d'Ent. xvi, p. 1, pl. ii, figs. 8 & 12; garleppi, p. 427, lamarchei, p. 428, Bolivia, Staudinger, Deuts e. Z. Lep. v: n. spp.

Paraussius: mode of formation of pouch, with new organ cal peraplast; Scudder, pp. 249-253, Tr. E. Soc. 1892; and discussi P. E. Soc. 1892, p. xxxiii. P. apollo, var. described; Bossan

Jahresh. Ver. Trencsin. xiii, p. 79. P. delphius, n. var., cæcus; Dönitz, Ent. Nachr. xviii, p. 193, fig. P. delphius, var. stenosemus, figured; Honrath, Berl. Ent. Zeit. xxxvi, pl. xv, fig. 3. P. delius, larva noticed and described; Sellon, Ent. xxv, p. 234: n. var., leonhardi; Rühl, p. 105, Soc. Ent. vii. P. orleans, pl. ii, fig. 14, orleans-groumi, fig. 10, p. 3, szechenyi, pl. ii, figs. 11 & 13, p. 3, noticed and figured; Oberthur, Études d'Ent. xvi.

P. poeta, Thibet, OBERTHUR, p. 2, pl. ii, fig. 9, Études d'Ent. xvi, n. sp. Pharmacophagus, n. subg. of Papilio; HAASE, Bibl. Zool. Heft. viii, p. 15.

Sericinus telamon, variation and larva described; STAUDINGER, pp. 130-136, Romanoff Mem. vi.

Thais polyzena, n. var., bipunctata; Cosmovici, Le Nat. 1892, p. 254.

#### PIERIDE.

- Anthocharis charlonia, varr. described; STAUDINGER, pp. 227-229, Deutsche e. Z. Lep. iv. A. sara and reakirtii are dimorphic forms; EDWARDS, pp. 52 & 109, Canad. Ent. xxiv.
- A. flora, Washington, WRIGHT, p. 154, Canad. Ent, xxiv, n. sp.
- Appias lalassis, Smith, = (Pieris indroides, Honr.); WEYMER, p. 121, S. E. Z. liii.
- A. flavius, N. Borneo, SMITH, p. 427, Ann. N. H. (6) x; A. dohertyi, Celebes, ROTHSCHILD, p. 441, pl. v, fig. 1, Deutschee. Z. Lep. v: n. spp.
- Aporia cratugi, notes on, irregular appearance, odour, metamorphoses, &c; Eckstein, Zool. Jahrb. Syst. vi, pp. 230-240. A. potanini, distinct from martineti; Alphéraky, p. 1, Romanoff Mem. vi.
- Belenois antsianaka, u. varr, drurii, figs. 4 & 5, persimilis, figs. 6 & 7, B. ramona, figs. 1, 2, & 3, figured; Smith & Kirby, Rhop. ex., Papilionidæ, Pierinæ, Belenois, i. B. lasti, nagare, majungana, isokani, figured; Smith & Kirby, Rhop. ex. ii, Papilionidæ, Pierinæ, Belenois, pl. ii.
- Callidryas eubule, life history of; DYAR, Canad. Ent. xxiv, p. 7. C. sennæ, habits; PANTON, J. Inst. Jamaica, i, p. 106.
- Callosune bacchus and mrogoroana, distinctions of; Vuillot, p. viii, Bull. Soc. Ent. Fr. lxi.
- Colias: characters of the palsearctic species; Bramson, Le Nat. 1892, pp. 83, 91, 104. C. rautieri, Guer., = (flaviola, Blanch., minuscula, Butl.); Beed, p. 68, Act. Soc. Chili, i. C. wiskotti, varr. described; Staudinger, pp. 230-232, Deutsche e. Z. Lep. iv. C. werdandi, and hecla, varietal note; Lampa, Ent. Tidskr. xiii, p. 249. C. viluiensis, note on; Graeser, p. 210, Berl. Ent. Zeit. xxxvii. C. edusa, earlier stages; Frohawk, Ent. xxv, p. 201. C. hyale, earlier stages; Frohawk, t. c. pp. 271-274. C. christina, larva and pupa noticed; Edwards, p. 110, Canad. Ent. xxiv. C. scudderi and nastes, food-plants; Edwards, p. 54, meudii and elis, characters noticed, p. 55, Canad. Ent. xxiv. C. philodice, albino; Edwards, p. 49, t. c.

. :

- Delias abnormis, n. var. euryxantha; HONRATH, p. 435, Berl. Ent. Ze xxxvi. D. peribæa, n. var. wallacei; ROTHSCHILD, p. 441, pl. v, fig. Deutsche e. Z. Lep. v.
- D. parthenia, Borneo, STAUDINGER, p. 449, Deutsche e. Z. Lep. v; tobahana, p. 571, hageni, p. 572, Sumatra, ROGENHOFEE, Verh. z. Wien, zlii: n. spp.
- Eronia cleodora, n. var. latimarginata; WEYMER, p. 96, S. E. Z. liii. Huphina celebensis, p. 439, pl. iv, figs. 1 & 2, dissimilis, p. 440, pl. v, fig 5 & 6, ROTHSCHILD, Deutsche e. Z. Lep. v, n. spp.
- Itias flavipennis, Smith, = (pyritis, Weym.); WEYMER, p. 12 S. E. Z. liii.
- I. kühni, Wetter, Röber, p. 287, Tijdschr. Ent. xxxiv (fig. in xxxv n. sp.
- Leuconea cratægi, n. var. alepica; Cosmovici, p. 254, Le Nat. 1892.
- Leucophasia amurensis, n. var. vernalis; GRAESER, p. 209, Berl. Em Zeit. xxxvii.
- Metaporia harrietæ, Bhutan, Nicéville, p. 341, pl. i, fig. 1, J. Bomb. N. H. Soc. vii, n. sp.
- Pieris (Mylothris) rubricosta, Mab., = (mackenziana, Sharpe), p. 61, figured, pl. x, fig. 3; Mabille & Vuillot, Nov. Lep. P. rape, var.; Swierstra, p. 337, pl. xvii, figs. 1 & 2, Tijdschr. Ent. xxiv. P. oberthuri, figured, pl. i, fig. 2, P. larraldei, n. var. melania, p. 5, pl. i, fig. 5, nutans, p. 6, fig. 3; Oberthur, Études d'Ent. xvi; P. zeuxippe, & var. figured; Poujade, pl. xi, figs. 1 & 1a, N. Arch. Mus. Paris (3) iii. P. brassicæ, immigration noticed; Kerry, Essex Nat. vi, p. 205. P. brassicæ and rapæ, pupal variation in colour; Poulton, Tr. E. Soc. 1892, pp. 437-446. P. oleracea and Pontia protodice are disappearing in N. America where P. brassicæ is introduced, Soc. Ent. vii, p. 94; but cf. Smith, t. c. p. 131.
- P. hilara, W. Africa, Karsch, p. 169, Ent. Nachr. xviii; P. hastata Yunnan, Oberthur, p. 5, pl. i, fig. 6, Études d'Ent. xvi; P. fruk storferi, Java, Honrath, p. 434, pl. xv, figs. 4 & 4a, Berl. Ent. Zeit xxxvi; P. oberthüri, p. 277, synchroma, p. 278, Flores, Röber Tijdschr. Ent. xxxiv (figg. in xxxv): n. spp.
- Tachyris adelpha, Alor, p. 281, paula, Wetter, p. 282 (with fig. in xxxv) Röber, Tijdschr. Ent. xxxiv, n. spp.
- Terias mandarinula, E. Africa, Holland, Ent. xxv (Supp.) p. 91, n. sp. Thestias reinwardtii, figured, pl. xv, fig. 2, noticed, p. 249; Snellen Tijdschr. Ent. xxxiv. T. flavipennis, Q figured; Swierstrp pl. 16: described; Snellen, p. 335, t. c.

#### DANAIDÆ.

- Anosia plexippus, note on early larva; SCUDDER, Psyche, vi, p. 250. Chlorochropsis, n. g., for C. dohertyi, n. sp., Celebes, pl. iv, fig. 3; ROTHE CHILD, p. 430, Deutsche e. Z. Lep. v.
- Elsa, n. g., near Danais, for E. morgeni, n. sp., Cameroons; HONRATE p. 436, pl. xv, fig. 5, Berl. Ent. Zeit. xxxvi.

- Euplea schreiberi, Butl., = (maasseni, Weym.); WEYMER, p. 119, S. E. Z. liii. E. alcathæ, p. 1, doubledayi, durrsteini, = (mesocala), p. 2, synonymy; SNELLEN, Tijdschr. Ent. xxxv. E. perdita, n. var., swierstræ; SNELLEN, p. 339, pl. xvii, fig. 4, Tijdschr. Ent. xxxiv. E. callithoë, Q described; TRYON (954). E. kansemanni, Q described and figured, p. 442, pl. xv, fig. 6, callithoë, Q described, p. 443; HONRATH, Berl. Ent. Zeit. xxxvi. E. corinna, figured; SCOTT, Austral. Lep. ii, pl. xvi. E. harrisii and allies, great variability; NICÉVILLE, p. 247, Tr. E. Soc. 1892.
- E. kuhni, Flores, p. 294 (fig. in vol. xxxv), oropina, Wetter, p. 296, sobrina, Goram, neptis, Flores, p. 297, nepos, Goram, p. 298 (fig. in xxxv), jacobseni, Wetter, p. 299 (fig. in xxxv), compta, Timor Laut, p. 300 (fig. in xxxv), Röber, Tijdschr. Eut. xxxiv, n. spp.
- E. (Calliplasa) parvior, N. Guines; TRYON (954).
- Glinama suctemon, Q = (configurata, Feld.); ROTHSCHILD, p. 432, Deutsche e. Z. Lep. v.
- Nasuma celebensis, Celebes, STAUDINGER, p. 431, pl. iv, fig. 4, Deutsche e. Z. Lep. v. n. sp.

# ACREIDE and HELICONIIDE.

- Acrea (Telchinia) anemosa, buettneri, abdominal pouch of Q figured;
  ROGENHOFER, pp. 574, 575, & 579-581, Verh. s.-b. Wien, xlii. A.
  vesperalis, figs. 1 & 2, crystallina, figs. 3 & 4, bomba, figs. 5 & 6,
  nelsoni, figs. 9 & 10, figured; SMITH & KIRBY, Rhop. ex. Nymphalida, Acreina, Acrea, pl. iii. A. cerasa, iturina, igola, andromba,
  § Q, mombasa, figured, SMITH & KIRBY, t. c., Nymphalida,
  Acreina, Acrea, iv. A. dohertyi, Q described and figured;
  ROTHSCHILD, p. 435, Deutsche e. Z. Lep. v.
- A. ombria, p. 82, albomaculata, p. 83, E. Africa, adrasta, W. Africa, p. 85, Weymer, S. E. Z. liii; A. balina, W. Africa, Karsch, p. 170, Ent. Nachr. xviii; A. (Telchinia) welwitschii, Angola, Rogenhofer, p. 573, Verh. z.-b. Wien, xlii; A. pharsaloides, minima, abbottii, E. Africa, Holland, Ent. xxv (Supp.) p. 89; A. marmorata, Madagascar, Smith & Kirby, Rhop. ex. Nymphalidæ, Acræinæ, Acræa, p. 9, pl. iii: n. spp.
- Heliconius rubellius, figs. 1 & 2, Columbia, amandus, fig. 3, p. 1, phyllidis, fig. 4, p. 2, anacreon, fig. 5, p. 3, SMITH & KIRBY, Rhop. ex. Nymphalidæ, Heliconinæ, Heliconius, i, n. spp.

# NYMPHALIDÆ.

Abrota pratti, figured; LEECH, pl. xiv, figs. 7 & 8, Butt. China.

Ageronia: sound-apparatus described; Hampson, P. Z. S. 1892, p. 190.

Amnosia martini, Sumatra, Honrath, p. 439, Berl. Ent. Zeit. xxxvi, n. sp.

- Apatura: Chinese species figured, and varr. described; LEE pp. 153-164, pls. xv & xvi, Butt. China. A. fasciola, subcarufulva, Leech, figured; Smith & Kirby, Rhop. ex. Nymphali Nymphalina, Apatura, i.
- Apatura laverna, China, LEECH, p. 164, pl. xv, fig. 6, Butt. China, n. : Araschnia: Chinese species figured, varr. described; LEECH, pp. 268-: pl. xxvi, Butt. China.
- A. doris, China, LERCH, p. 272, pl. xxvi, figs. 4 & 5, Butt. China, n. s. Argynnis: Chinese species figured, and varr. described; Leipp. 222-244, pls. xxii-xxiv, Butt. China. A. zenobia, figured Oberthur, p. 7, pl. i, fig. 1, Études d'Ent. xvi. A. lathonio = (anna, Blanchd.); Reed, p. 68, Act. Soc. Chili, i. A. pales, n. killiasi; Rühl, Soc. Ent. vii, p. 113. A. pandora, n. var. dac Hormuzaki, Ent. Nachr. xviii, p. 1. A. paphia, n. var. nigric Cosmovici, p. 256, Le Nat. 1892. A. epithore, early sta Edwards, p. 108, Canad. Ent. xxiv. A. paphia, colour of larva surroundings; Poulton, Tr. E. Soc. 1892, p. 437. Oviposition species of, Skinner, p. 128, Canad. Ent. xxiv.
- A. rosea, Roumania, Cosmovici, p. 256, Le Nat. 1892; A. pene Amurland, Staudinger, p. 195, Romanoff Mem. vi, and p. Deutsche e. Z. Lep. iv; A. ornatissima, China, Leech, p. pl. xxii, figs. 1 & 2, Butt. China; A. oweni, Mt. Shasta, p. 105, nelia, Colorado, p. 106, Edwards, Canad. Ent. xxiv: n. spp.
- Atella sinha, specific validity; SNELLEN, p. 5, Tijdschr. Ent. xxxv. Athyma: Chinese species figured, varr. described; Leech, pp. 167-pls, xvi & xvii, Butt. China.
- A. recurra, W. China, LEECH, p. 176, pl. xvii, fig. 9, Butt. China; A. a. Borneo, STAUDINGER, p. 452, Deutsche e. Z. Lep. v: n. spp.
- Cutuna duodecimpunctata. note on; SNELLEN, p. 4, Tijdschr. Ent. xi Charaxes agoba, Dist., = (homerus, Stdgr.), porthos, Smith, = (n Stdgr.); STAUDINGER, p. 223, Deutsche e. Z. Lep. iv. C. nic Smith, = (ogovensis, Holl., = hamatus, Dew.), C. hildebrandti, = (talagugæ, Holl.); WEYMER, p. 120, S. E. Z. liii. C. rothen n. n. for ganymedes, Leech, nec Westw., p. 128, figured, pl. xiv, : Leech, Butt. China. C. pelias, n. var. tanyanika; Robbe, p Ann. Ent. Belg. xxxvi. C. nitebis, figured, pl. vi, fig. 3, C. n. var. dohertyi, p. 437, pl. vi, fig. 2, Deutsche e. Z. Lep. v.
- C. lactetinctus, W. Africa, KARSCH, p. 113, Eut. Nachr. xviii; C. od: W. Africa, p. 260, georgius, Miudoro, p. 262, STAUDINGER, Det e. Z. Lep. v.: n. spp.
- Cirrechroa niassica, Nias I., Honrath, p. 437, Berl. Ent. Zeit. : n. sp.
- Crenis umbrina, W. Africa, KARSCH, p. 114, Ent. Nachr. xviii; C. g W. Africa, KARSCH, p. 173, t. c.: n. spp.
- Cymothoë blassi, E. Africa, WEYMER, p. 90, S. E. Z. liii, n. sp.
- Doleschallia pfeili, New Ireland, HONRATH, p. 438, pl. xv, fig. 7, Ent. Zeit. xxxvi, n. sp.

- Euphædra edwardsii, Hoeven, = (pralinas, Dbl.), inanum, Butl., = (vespasia, Moesch.); Weymer, p. 120, S. E. Z. liii.
- E. judith, Sierra Leone, WEYMER, p. 88, S. E. Z. liii, n. sp.
- Euptoista hortensia, Blanch., =(Argynnis valdiviana, Phil.); REED, p. 68, Act. Soc. Chili, i.
- Eurytela velleda, figured, pl. x, fig. 1; MABILLE & VUILLOT, Nov. Lep.
- Euthalia: Chinese species figured and varr. described; LEECH, pp. 135-140, pl. xxi, Butt. China.
- E. pyrrha, China, LEECH, p. 137, pl. xxi, fig. 4, Butt. China, n. sp.
- Godartia: generic characters and components; Snellen, p. 8, Tijdschr. Ent. xxxv.
- Grapta gigantea, figured, pl. xxv, fig. 6, c-album, varr. described; Leech, pp. 263-265, Butt. China.
- Harma canis, migration of, larva and chrysalis figured; Holland, Psyche, vi, pl. v, figs. 4 & 5, p. 215.
- Harmilla, n. g., near Cymothoë, for H. elegans, n. sp., Cameroons; Aurivillius, p. 200, Ent. Tidskr. xiii.
- Hestina mena, variation; LEECH, p. 143, Butt China.
- Hypolimnas angustolimbata, E. Africa, WEYMER, p. 87, S. E. Z. liii, n. sp. Junonia touhilimasa, Tanganyika, VUILLOT, p. exlviii, Bull. Soc. Ent. Fr. lxi, n. sp.
- Kullima spiridion, knyvetti, figured; Smith & Kirby, Rhop. ex. Nymphalidæ, Kallima, i.
- Limenitis: Chinese species figured, varr. described, pp. 178-188, pls. xvi & xvii; Leech, Butt. China. L. sibylla, n. var. stenotænia; Honrath, p. 440, Berl. Ent. Zeit. xxxvi. L. albomaculata, figured and described; Oberthur, p. 7, pl. ii, fig. 15, Études d'Ent. xvi. L. weidemeyerii, early stages; Edwards, p. 107, Canad. Ent. xxiv.
- L. dosrriesi, Amurland, Staudinger, p. 173, pl. xiv, fig. 1, Romanoff Mem. vi; L. hageni, Sumatra, Staudinger, p. 452, Deutsche e. Z. Lep. v: n. spp.
- Melitza: Chinese species figured; LEECH, pl. xxiv, Butt. China. M. maculata, Brem., generic position; SNELLEN, p. 6, Tijdschr. Ent. xxxv. M. anicia, larva going over two years; EDWARDS, p. 53, Canad. Ent. xxiv. M. chalcedon, oviposition; KUNZE, p. 190, Canad. Ent. xxiv.
- M. bellona, W. China, LEECH, p. 219, pl. xxiv, figs. 1-5, Butt. China, n. sp. Neptis: Chinese species figured, varr. described; LEECH, pp. 190-207, pls. xviii & xix, Butt. China. N. biafra, n. var. continuata; Holland, Ent. News, iii, p. 249.
- Neptis nicobule, metanira, mixophyes, Ogowe Valley, Holland, Ent. News, iii, p. 249, pl. ix; N. thestias, W. China, Leech, p. 196, pl. xviii, fig. 3, Butt. China; N. patalina, Philippine Is., Semper, p. 347, Reisen Philipp. ii, v; N. florensis, Flores, Snellen, p. 238, Tijdschr. Ent. xxxiv: n. spp.
- Nymphalis freyi, Nossibé, Brancsik, p. 161, pl. vii, fig. 8, Jahresh. Ver. Trencsin, xiii, n. sp.

- Palla lichas, Dbl., p. 93, falcata, Butl., p. 94, Qs described; WEYNE S. E. Z. liii.
- P. publius, Sierra Leone, STAUDINGER, p. 267, Deutsche e. Z. Lep. v; rectifascia, W. Africa, WEYMER, p. 91, S. E. Z. liii: n. spp.
- Phyciodes gorgone, Hub., = (ismeria, Boisd., and carlota, Reak.); WE MER, p. 119, S. E. Z. liii.
- Precis serena, Sierra Leone, WEYMER, p. 86, S. E. Z. liii, n. sp.
- Prothoë calydonia, = (belisama, Crowley); WEYMER, p. 120, S. E. Z. I Pseudacrae uhelda, Mab., figured; Mabille & Vuillot, pl. x, fig. Nov. Lep. P. poggei, & described and figured; Butler & Roti Child, p. 202, pl. x, fig. 2, Tr. E. Soc. 1892.
- P. warburgi, Cameroona, AURIVILLIUS, Ent. Tidskr. xiii, p. 200; clarkii, Congo River, BUTLER, p. 201, pl. x, fig. 1, Tr. E. Soc. 189: n. spp.
- Pseudargynnis, n. g., for Catuna duodecimpunctata, auct.; KARSCH, p. 1 Ent. Nachr. xviii.
- Pyrameis cardui, life-history; DYAR, Canad. Ent. xxiv, p. 6. P. ii phylogenetic note; SMITH, p. 27, Ent. xxv.
- Rahinda sandaka, Borneo, BUTLER, p. 120, P. Z. S. 1892, n. sp.
- Stibochiona persephone, Borneo, STAUDINGER, p. 451, Deutsche e. Z. L v, n. sp.
- Symbrenthia lucina, hippoclus, distinctive characters; SEMPER, p. 3 Reisen Philipp. ii, v.
- Terinos novæ-guinesnais, New Guinea; TRYON (954), n. sp.
- Thaleropis trigona, W. Africa, Holland, Ann. N. H. (6) x, p. 284, n. t
- Timelwa nana, W. China, Leecii, p. 246, pl. xxiii, fig. 8, Butt. China, n. Vanessa fenestra, Leech, referred to Dilipa, = (Apatura chrysus, Obert
  - p. 165, figured, pl. xiv, fig. 2; LEECH, Butt. China. V. urtivariation in China; LEECH, p. 258, pl. xxv, fig. 1, Butt. Chin variation produced by exposure of pupes to different temperature MERRIFIELD, p. 36, Tr. E. Soc. 1892. V. urtica and io, colour pupa and surroundings; Poulton, pp. 391-402, polychloros: atalanta, pp. 432-437. V. urtica, gilded appearance of pupes cussed; Bateson, p. 210-214.
- Vanessula: characters and position discussed, V. milca, fig., p. 171, (sub Liptena) milca, Hew., = (buchneri, Dew.), p. 172; KARSCH, F. Nachr. xviii.

#### MORPHIDÆ.

- Clerome ærope, figured, pl. i, fig. 4; LEECH, Butt. China.
- Discophora dis, Sumatra, &c., Nicáville, p. 325, pl. H, fig. 3, J. Bol N. H. Soc, vii, n. sp.
- Enispe lunatus, figured; LEECH, Butt. China, pl. i, figs. 1 & 2.
- Morpho sulkowskyi, n. var. psyche; STAUDINGER, Deutsche e. Z. Lep. p. 197.
- Stichophthalma hovoqua, var. figured, pl. i, fig. 3; LEECH, Butt. China. S. neumogeni, W. China, LEECH, p. 114, pl. i, fig. 5, Butt. China, n. sp

# SATYRIDE.

- MOORE continues his descriptions and figures of E. Indian Satyridæ in Lep. Ind. i, pp. 177-310. pls. lxv-xciv.
- Acrophthalmia thalia, figured; LEECH, pl. x, fig. 1, Butt. China.
- A. misarte, Philippines, SEMPER, p. 327, Reisen Philipp. ii, v, n. sp.
- Amecera majuscula, W. China, LEECH, Butt. China, p. 67, pl. xii, fig. 6, n. sp.
- Archondesa, n. g., type, Lethe lanaris, Butl.; Moore, p. 270, Lep. Ind. i.
- Bicyclus ephorus, W. Africa, WEYMER, p. 79, S. E. Z. liii, n. sp.
- Blanaida pulahoides, Naga Hills, Moore, p. 304, pl. 94, fig. 2, Lep. Ind. i, n. sp.
- Callarge, n. g. for Zethera sagitta, Leech, which is figured, pl. xii, fig. 1; Leech, p. 57, Butt. China.
- Callerebia phyllis, albipuncta, orixa, figured; Leech, pls. ix & xii, Butt. China.
- Calysisme persecides, Burma, p. 179, pls. lx, fig. 2, & lxi, fig. 1, nicobarica, Nicobar Is., intermedia, p. 187, subdita, p. 194, pl. lxv, figs. 1, a-h, E. India, rama, pl. 57, fig. 3, Ceylon, andamana, pl. 66, fig. 1, Andaman Is., p. 196, Moore, Lep. Ind. i, n. spp.
- Catargynnis, n. n. for Pseudomaniola (infrå); RÖBER, p. 284, Exot. Schmett. Theil ii.
- Chionobas chryxus, calais, figured, Chionobas ii, jutta, Chionobas v, crambis, brucci, Chionobas vi; Edwards, Butt. N. Am. (3).
- Choranesa, n. g., type Lethe trimacula, Leech; Moore, p. 270, Lep. Ind. i.
- Cwlites adamsoni, p. 229, pl. 75, fig. 1, binghami, p. 230, fig. 3, E. India, Moore, Lep. Ind. iii, n. spp.
- Comonympha pamphilus, n. var. bi-pupillata; Cosmovici, p. 264, Le Nat. 1892.
- Debis (sub Satyrus) hyrania, Koll., = (goalpara, Moore), p. 3, samio, Dbl., = (manthara, Feld.), p. 4; Snellen, Tijdschr. Ent. xxxv.
- Dionana, n. g., type Lethe margaritæ, Elw.; Moore, p. 271, Lep. Ind. i.
- Epinephele monarchus, Bl., = (valdiviæ, Feld.); REED, p. 68, Act. Soc. Chili, i. E. tithonus, n. var. mincki; SEEBOLD, p. 467, Berl. Ent. Zeit. xxxvi. E. janira, var. cinerascens; Fuchs, JB. nass. Ver. xlv, p. 85.
- E. cinerea, erebiformis, Roumania, Cosmovici, p. 264, Le Nat. 1892, n. spp. Erebia: Chinese species figured, varr. described, Leech, pl. ix, pp. 97-101, Butt. China. E. medusa, n. var. procopiani; Hormuzaki, Ent. Nachr. xviii, p. 2. E. sofia, Streck., = (ethela, Edw.); Strecker, Ent. News, iii, p. 255. E. melas, metamorphoses; Dognin, p. 211, Le Nat. 1892.
- E. nigra, Roumanis, Cosmovici, p. 264, Le Nat. 1892, n. sp.
- Harima, n. g., type Neope callipteris, Butl.; Moore, p. 299, Lep. Ind. i.
- Idiomorphus vala, larva described; Holland, Psyche, vi, p. 214.
- Kerrata, n. g., for Lethe tristigmata, Elw.; Moore, p. 285, Lep. Ind. i.

- Kirrodesa, n. g., type Debis sicelis, Hew.; MOORE, p. 237, Lep. Ind. i. Lethe: Chinese species figured, and varr. described; LEECH, pp. 18-4 pls. iii-vi, Butt. China.
- L. prozima, W. China, LEECH, p. 32, pl. vi, fig. 8, Butt. China, n. sp. Mandarinia, n. g., for Mycalesis regalis, Leech; LEECH, Butt. China, p. Malanassia lada, Leech (Assaurance, Oberth), LEECH p. 60, Bu
- Melanargia leda, Leech, = (Arge yunana, Oberth.); LEECH, p. 60, Bu China. M. titea, n. var. titania; STANDFUSS, p. 659, pl. xv, fig. Romanoff Mem. vi.
- M. grumi, Kurdistan, Standfuss, p. 661, pl. xv, fig. 2, Romanoff Me vi, n. sp.
- Mycalesis: Chinese species figured and n. varr. described; LEECH, pp. 1 15, pl. ii, Butt. China. M. dinon, Hew., = (Messaras ophthalmic Westw.); WEYMER, p. 119, S. E. Z. liii.
- M. anisops, W. Africa, Karsch, p. 176, Ent. Nachr. xviii; M. uni Moupin, Leech, Butt. China, p. 15, pl. ii, fig. 9; M. kina, Born Staudinger, p. 451, Deutsche e. Z. Lep. v; M. (Calysisme) pernota N. Guinea, Tryon (954): n. spp.
- Nemetis, n. g. (for Lethe group, 1 pt., Nioéville); Moore, p. 237, L. Ind. i.
- Neomænas servilia, Wllgr., = (Satyrus decorata, Butl.); REED, p. 68, A Soc. Chili, i.
- Neope: Chinese species figured, pls. vii & viii, varr. described, pp. 48 54; Leech, Butt. China.
- Neorina patria, Leech, figured, pl. xii, fig. 2; LEECH, Butt. China.
- N. westwoodii, p. 226, pl. lxxiv, fig. 2, E. India, Moore, Lep. Ind. i, n. Neosatyrus ambiorix, Wllgr., = (minimus, Butl.), violaceus, Butl., (ochreivittatus, Butl.); Reed, p. 68, Act. Soc. Chili, i.
- Ninguta, n. g., type Pronophila schrenkii, Men.; Moore, p. 310, Lep. Inc Pararge præusta, catena, deidamia, figured, pls. xi & xii; Leech, Bt China. P. megæra, var. mediolugens; Fuchs, p. 87, JB. nass. V xlv.
- Patala, n. g. (type Zophoessa yama, Moore), p. 305, with P. yamoin n. sp., N. India, p. 307, pl. xciv, fig. 4; Moore, Lep. Ind. i.
- Pegada, n. g., for Mycalesis oculatissima, Pouj.; MOORE, pp. 224, L. Ind. i.
- Placilla, n. g. (type Lethe christophi, Leech); Moore, p. 253, L. Ind. i.
- Pseudomaniola, n. g., p. 222 (replaced by Caturgynnis, p. 284), for part Dædalma, Hew.; Röber, Exot. Schmett., Theil ii.
- Ptychandra mindorana (Stdgr.), Philippines, Semper, p. 329, Rei Philipp. ii, v, n. sp.
- Putlia, n. n. for Charma, Doh. (preocc.); Moore, p. 287, Lep. Ind. i. Rugadia critolaus, Burma, Nickville, p. 322, J. Bomb. N. H. Soc. pl. H, fig. 1; R. melita, Borneo, Staudinger, p. 450, Deutsche e. Lep. v: n. spp.
- Ranghia, n. g. (for Lethe group, 1 pt., Nicéville), p. 232, with R. pegua n. sp., Burma, p. 235; Moore, Lep. Ind. i.

- Samanta bethami, p. 205, pl. lxviii, fig. 4, davisoni, p. 206, fig. 3, E. India, Moore, Lep. Ind. i, n. spp.
- Satyrus autonoë, dryas, n. varr. described and figured; Leech, pp. 69 & 70, pls. xi & xiii, Butt. China. S. autonoë, var. extrema, figured, pl. i, fig. 1, Romanoff Mem. vi. S. bischoffi, varr. described; STAUDINGER, p. 237, Deutsche e. Z. Lep. iv.
- S. favonius, Malatia, STAUDINGER, p. 239, pl. iii, fig. 1, Deutsche e. Z. Lep. iv, n. sp.
- Sinchula, n. g. (for Lethe group, 3 pt., Nicéville); MOORE, p. 275, Lep. Ind. i.
- Trophonina, n. g., for Lymanopoda acræida, Butl.; Röber, p. 222, Exot. Schmett., Theil ii (omission from 1889 Record).
- Ypthima: Chinese species figured, and varr. described; Leech, pp. 81-91, pls. ix & x, Butt. China. Y. baldus, figured; Snellen, pl. xv, fig. 1, p. 236, Tijdschr. Ent. xxxiv.
- Y. argillosa, p. 133, nigricans, p. 135, Java, Snellen, Tijdschr. Ent. xxxv; Y. florensis, Flores, Snellen, p. 235, pl. xiv, fig. 3, Tijdschr. Ent. xxxiv; Y. celebensis, Celebes, Rothschild, p. 433, pl. v, figs. 3 & 4, Deutsche e. Z. Lep. v; Y. subarctous, New Guinea, Tryon (954): n. spp.
- Zophoessa helle, procee, argentata, dora, figured; Lebch, pl. vii, Butt. China.
- Z. gummiei, p. 294, pl. xei, fig. 3, elwesi, p. 298, pl. xeii, fig. 3, Bhutau; Moore, Lep. Ind. i, n. spp.

# ELYMNIDE and ERYCINIDE.

- Drepanula, n. g., for Lemonias calous, Stdgr.; Röber, p. 246, Exot. Schmett., Theil ii (omitted in Zool. Rec. of last year).
- Dyctis esacoides, Perak, Niceville, p. 323, pl. H, fig. 2, J. Bomb. N. H-Soc. vii, n. sp.
- Elymnias borneensis, Borneo, SMITH, p. 428, Ann. N. H. (6) x, n. sp.

#### LYCENIDE

- Lycanida of the South Pacific Is., with numerous synonymical and critical remarks; DRUCE, P. Z. S. 1892, pp. 434-446, pl. xxvii.
- Acesina zephyretta, fig. 8, ariel, fig. 9, figured; ELWES, pl. xliv, P. Z. S. 1892.
- Allotinus panormis, Karen Hills, ELWES, p. 619, P. Z. S. 1892, n. sp.
- Aphneus vulcanus, n. var. maximus, p. 637, pl. xliii, fig. 5, sani, var. ?, p. 638, fig. 6; ELWES, P. Z. S. 1892.
- Argyrocheila, n. g., for A. undifera, n. sp., W. Africa; STAUDINGER, p. 215, Deutsche e. Z. Lep. iv: figured; SMITH & KIRBY, Rhop. ex. ii, African Lycanida, pl. xviii, figs. 11 & 12.
- Arhopala ace, p. 329, pl. H, fig. 13, acestes, p. 330. fig. 14, Perak, arca, Celebes, p. 331, fig. 15, asia, Malay Penins., fig. 16, acta, Burma, fig. 17, p. 333, Niceville, J. Bomb. N. H. Soc. vii, n. spp.

- Aslauga marginalis, figured; Smith & Kirby, Rhop. ex. ii, Africa Lycanida, pl. xix, figs. 15 & 16.
- Camena carmentalis, Khasi Hills, NICEVILLE, p. 335, pl. H, fig. 10, Bomb. N. H. Soc. vii; C. icetoides, p. 636, pl. xliv, fig. 3, cleoboid p. 637, figs. 4 & 5, Karen Hills, Elwes, P. Z. S. 1892: n. spp.
- Catapacilma subochrea, Karen Hills, ELWES, p. 640, pl. xliv, fig. 1 P. Z. S. 1892.
- Chrysophanus abbottii, E. Africa, Holland, Ent. xxv (Supp.), p. 90, n. : Curetis stigmata (Moore), Röber, p. 318, Tijdschr. Ent. xxxiv, n. sp.
- Cyaniris ceyx, Java, Niceville, p. 328, pl. H, figs. 6 & 7, J. Bomb. N. Soc. vii, n. sp.
- Deudorix carulea, Druce, = (obscurata, Trim.), the species referred Rapala; DRUCE, Ent. M. M. xxviii, p. 65.
- D. gatulia, Khasi Hills, Nickville, p. 338, pl. H, fig. 12, J. Bomb. N. Soc. vii; D. verriculata, Celebes, Snellen, p. 257, pl. xv, fig. Tijdschr. Ent. xxxiv; D. mathewi, New Hebrides, DRUCE, p. 4 P. Z. S. 1892, pl. xxvii, fig 9: n. spp.
- Drupadia casarea, Weym., = (Sithon ravindra, var. niasica, Stdg: Weymer, p. 121, S. E. Z. liii.
- Durbania gerda, figured; SMITH & KIRBY, Rhop. ex. i, African Lycami pl. xvi, figs. 9 & 10.
- D. cornucopiæ, W. Africa, Holland, p. 286, Ann. N. H. (6) x, n. sp. Eupsychellus, n. g., near Plebeius, for dionysius, Boisd.; Röber, p. ? Tijdschr. Ent. xxxiv, and p. 273, Exot. Schmett., Theil ii.
- Epitola doleta, henleyi, goodii, barombiensis, catuna, figured; Smite Kirby, Rhop. ex. i, African Lycanida, pl. xvii.
- E. ciconia, Sierra Leone, Smith & Kirby, Rhop. ex. i, African Lycon pl. xvii, p. 58, n. sp.
- Hewitsonia kirbyi, Dew., = (preussi, Stdgr.); STAUDINGER, p. Deutsche e. Z. Lep. iv.
- Hypochlorosis, n. g., for Sithon antipha, Hew.; RÖBER, p. 267, E Schmett., Theil ii.
- Hypolycena rava, W. Africa, Holland, p. 286, Ann. N. H. (6) x, n. Iolaus julianus, Staud., and alcibiades, Don., noticed, I. julius, Staud (lekanion, Druce); Druce, p. 185, Ent. M. M. xxviii.
- Jamides pulcherrima, pl. xxvii, fig. 16, carissima, fig. 17, figured; Dr. P. Z. S. 1892.
- J. kava, pl. xxvii, fig. 15, walkeri, figs. 13 & 14, South Pacific Is., DR p. 443, P. Z. S. 1892, n. spp.
- Kopelatus virgata, Druce, = (Hypolycana gracilis, Stdgr.); STAUDINp. 223, Deutsche e. Z. Lep., and Druce, p. 66, Ent. M. M. xxviii
- Lachnocnema reutlingeri, Benita, Holland, p. 286, Ann. N. H. (6) x, r. Læosopis roboris, n. var. lusitanica; Staudinger, p. 232, Deutsche dep. iv.
- Larinopoda sylpha, opaca, figured; SMITH & KIRBY, Rhop. ex., Afi
  Lycanida, pl. xvi. L. lara, sylphida, lybia, campinus, figured; Si
  & KIRBY, Rhop. ex. ii, African Lycanida, xviii.

- L. sylphida, lybia, p. 217, lara, p. 218, W. Africa, STAUDINGER, Deutsche e. Z. Lep. iv, n. spp.
- Liptena mnestra, Möschl., = (Teriomina dispar, Kirby); DRUCE, Ent. M. M. xxviii, p. 65. L. rubrica, Druce, = (Pseuderesia soraida, Smith & Kirby); DRUCE, t. c.
- Lycana rhymnus, tengstramii, pretiosa, referred to Thecla, their & characters fully described and figured; Baker, pp. 27-31, pl. ii, Tr. E. Soc. 1892. L. arion, var. unicolor; Hormuzaki, Ent. Nachr. xviii, p. 1. L. amanda, n. var. lydia; Krulikowsky, Eut. Nachr. xviii, p. 97. L. arion, n. var. ruhli; Krulikowsky, Soc. Ent. vii, p. 1. L. dolus, damon and rippertii, note on; Oberthur, p. ix, Bull. Soc. Ent. Fr. lxi. L. lucifera, note on, with figure; Staudinger, p. 316, pl. iii, fig. 1, Dentsche e. Z. Lep. v. L. comyntas, metamorphoses; Brutenmüller, p. 76, Bull. Am. Mus. iv. L. lygdamas, larva noticed, Edwards, p. 109, Canad. Ent. xxiv. L. icarus, number of yearly generations in E. Russia; Krulikowsky, p. 370, Ent. Nachr. xviii.
- L. dama, Malatia, p. 234, triphysina, Kashgar, p. 235, pl. iii, fig. 2, STAUDINGER, Deutsche e. Z. Lep. iv; L. stormsi, Tanganyka, Robbe, p. 132, Ann. Ent. Belg. xxxvi; L. perpulchra. E. Africa, Holland, Ent. xxv (Supp.), p. 90; L. saturata, p. 137, deliana, p. 139, datarica, p. 140, glauca, p. 142, quadriplaga, p. 143, Java, SNELLEN, Tijdschr. Ent. xxxv: n. spp.
- Nacaduba vitiensis, pl. xxvii, figs. 3 & 4, florinda, fig. 12, figured; DRUCE, P. Z. S. 1892.
- N. aberrans, E. Pegu, Elwes, p. 626, pl. xliv, fig. 6, P. Z. S. 1892; N. samoensis, p. 437, pl. xxvii, figs. 5 & 6, Samoa Is., novahebridensis, figs. 7 & 8, p. 438, mallicollo, fig. 18, p. 439, New Hebrides, nebulosa, Australasia, figs. 10 & 11, p. 440, Druce, P. Z. S. 1892: n. spp.
- Narathura pryeri, Borneo, BUTLER, p. 121, P. Z. S. 1892, n. sp.
- Papua, n. g., for P. lucifer, Röber; Röber, p. 273, Exot. Schmett., Theil ii.
- Pentila subundularis, figured; Smith & Kirby, Rhop. ex. ii, African Lycænidæ, xviii.
- P. umbra, W. Africa, Holland, p. 285, Ann. N. H. (6) x; P. subundularis, W. Africa, Staudinger, p. 215, Deutsche e. Z. Lep. iv: n. spp.
- Philiris, n. g., for Plebeius ilias, Feld., Röher, p. 317, Tijdschr. Ent. xxxiv, and p. 273, Exot. Schmett., Theil ii.
- Pilodeudorix barbatus, Druce, = (Hypolycana nobilis, Stdgr.); Stau-DINGER, p. 223, Deutsche e. Z. Lep., and Druce, Ent. M. M. XXVIII, p. 66.
- Pithecops bassaris, Ké Is., Nicéville, p. 327, pl. H, figs. 4 & 5, J. Bomb. N. H. Soc. vii, n. sp.
- Plebeius cyaniris, p. 315, P. (Theclinesthes) eremicola, p. 316, Flores, Röber, Tijdschr. Ent. xxxiv (figg. in xxxv), n. spp.
- Polyommatus agestis, var. noticed; BARRETT, Ent. M. M. xxviii, p. 245.

  P. dispar, var. rutilus, taken in England; BAKER, p. xxx, P. E. Soc.
  1892.

- P. cupreus, Roumania, Cosmovici, p. 255, Le Nat. 1892, n. sp.
- Pseudaletis agrippina, Druce, & = (tricolor, Stand.); DRUCE, Ent. M. M. xxviii, p. 65.
- P. nigra, W. Africa, Holland, p. 286, Ann. N. H. (6) x, n. sp.
- Pseuderesia tullia, W. Africa, STAUDINGER, p. 221, Deutsche e. Z. Lep. iv, n. sp.
- Rapala (sub Deudorix) diyllus, Hew., = (Hypolycana debilis, Stand.); DRUCE, p. 185, Ent. M. M. xxviii.
- B. abnormis, p. 642, pl. xliv, fig. 2, hypargyria, p. 643, fig. 7, subguttata, fig. 1, p. 644, Karen Hills, ELWES, P. Z. S. 1892, n. spp.
- Spalgis lemolea, Druce, = (latimarginata, Sharpe, and s-signata, Holland); DRUCE, Ent. M. M. xxviii, p. 66. S. s-signata, habits of larva, peculiar chrysalis resembling old man's head; Holland, Psyche, vi, p. 201, pl. iv.
- Suasa suessa, Perak, Nicéville, p. 337, pl. H, figs. 8 & 9, J. Bomb. N. H. Soc. vii, n. sp.
- Tajuria thyia, Khasi Hills, Nickville, p. 336, pl. H, fig. 11, J. Bomb. N. H. Soc. vii, n. sp.
- Teriomima delicatula, figs. 5 & 6, decipiens, figs. 7 & 8, serena, figs. 11 & 12, figured; SMITH & KIRBY, Rhop. ex. i, African Lycanida, pl. xvi. T. adelgunda, adelgitha, modesta, cordelia, alberta, figured; SMITH & KIRBY, op. cit. ii, pl. xix.
- T. adelgunda, p. 219, alberta, p. 220, W. Africa, STAUDINGER, Deutsche e. Z. Lep. iv, n. spp.
- Thecla rumania, Roumania, Cosmovici, p. 255, Le Nat. 1892; T. leechii, Khasi Hills, Niceville, p. 335, J. Bomb. N. H. Soc. vii: n. spp.
- Theclinesthes, n. subg. of Plebeius, q.v.; RÖBER, p. 316, Tijdschr. Ent.
- Thestor romanovi, n. var. maculifera, STAUDINGER, p. 233, Deutsche e. Z. Lep. iv.
- Tingra lunaris, Sierra Leone, WEYMER, p. 95, S. E. S. liii, n. sp.
- Zizera labradus, pl. xxvii, fig. 1, lulu, = (conjungens, Luc.), fig. 2, figured; DRUCE, P. Z. S. 1892.

#### HESPERIIDE.

- Many of the Philippine Is. species described by Mabille are referred to other genera by SEMPER, in Reisen Philipp. ii, v, pp. 287-319.
- Notes on N. American Hesperiidæ (genera); AARON, pp. 119 & 120, Canad. Ent. xxiv.
- Abaratha syricthus, var. figured; ELWES, p. 656, pl. xliii, fig. 2, P. Z. S. 1892.
- Abantis bismarcki, W. Africa, KARSCH, p. 228, Ent. Nachr. zviii, n. sp.
- Astictopterus ladana, fig. 4, sindu, fig. 3, figured, pl. xliii; ELWES, P. Z. S.
- Calpodes ethius, life-history; COCKERELL, Ent. News, iii, p. 78.

- Capila jayadeva and Pisola zennara, correction as to characters and sexes of; Nickville, p. 347, J. Bomb. N. H. Soc. vii.
- Caprona adelica, W. Africa, KARSCH, p. 242, Ent. Nachr. xviii, n. sp.
- Carterocephalus palæmon, life-history; FROHAWK, Ent. xxv, pp. 225-228 & 254-256.
- Carystus duris, Mab., referred to Lotongus and figured; SEMPER, p. 289, pl. xlix, fig. 7, Reisen Philipp. ii, v.
- Crossiura, n. g., near Capila, p. 350, for C. pennicillatum, n. sp., Khasi Hills, p. 351, pl. J, figs. 1 & 2, NICEVILLE, J. Bomb. N. H. Soc. vii,
- Cyclopides romi, Congo; ROBBE, p. 133, Ann. Ent. Belg. xxxvi, n. sp.
- Erionota alexandra, Philippine Is., SEMPER, Reisen Philipp. ii, v, p. 312, n. sp.
- Eudamus coyote, Texas, Skinner, p. 164, Canad. Ent. xxiv, n. sp.
- Halps fusca, E. India, ELWES, p. 653, pl. xliii, fig. 1, P. Z. S. 1892, n. sp.
- Hesperia gemma, Plötz., = (Pamphila ravola, Godm. & Salv.); WEYMER, p. 122, S. E. Z. liii. H. lineola, early stages described; HAWES, Ent. xxv, pp. 177-180.
- H. mucorea, sintgraffi, W. Africa, Karsch, p. 178, Ent. Nachr. xviii, n. spp. Ismene fergusonii, S. India, Nickville, p. 345, pl. J, fig. 6, J. Bomb. N. H. Soc. vii; I. antigone, Flores, Röber, p. 320, Tijdschr. Ent. xxxiv (fig. in xxxv), n. spp.
- Lotongus parthenope, Weymer, sexes figured, & described; NICÉVILLE, p. 354, pl. J, figs. 4 & 5, J. Bomb. N. H. Soc. vii.
- Osmodes lux, W. Africa, Holland, p. 291, Ann. N. H. (6) x, n. sp.
- Oxypalpus annulifer, W. Africa, Holland, Ann. N. H. (6) x, p. 293, n. sp.
- Padraona procles, Ké Is., Nicéville, p. 353, pl. J, figs. 7 & 8, J. Bomb. N. H. Soc. vii, n. sp.
- Pamphila dimila, Q figured; NICEVILLE, pl. J, fig. 9, J. Bomb. N. H. Soc. vii. P. manitoba, variation; LYMAN, Canad. Ent. xxiv, p. 57.
- P. errans, N. America, Skinner, p. 174, Ent. News, iii (metamorphoses Dyar, t. c., p. 175); P. carolina, N. America, Skinner, p. 222, t. c.: n. spp.
- Pardaleodes cedipus, Cr., n. var. diluta; Robbe, p. 134, Ann. Ent. Belg.
- P. zanthopeplus, p. 289, astrape, zunthioides, p. 290, W. Africa, Holland, Ann. N. H. (6) x, n. spp.
- Plastingia viburnia (Stdgr.), cuneiformis, pl. xlix, fig. 11, Philippines, Semper, p. 314, Reisen Philipp. ii, v, n. spp.
- Procampta, n. g., for P. rara, n. sp., W. Africa; Holland, p. 294, Ann. N. H. (6) x.
- Sarangesa, Moore, = (Erites, Hyda, and Sape, Mab.); Holland, p. 288, Ann. N. H. (6) x.
- S. motosioides, W. Africa, Holland, p. 288, Ann. N. H. (6) x, n. sp.
- Satarupa corona (Stdgr.), Philippines, SEMPER, p. 306, Reisen Philipp. ii, v, n. sp.
- Suastus albinus, pl. xlix, fig. 8, p. 299, migreus, fig. 9, p. 300, Philippines, Semper, Reisen Philipp. ii, v. n. spp.
  - 1892. [VOL. XXIX.]

Tagiades princeps (Stdgr.), Philippines, Sempre, p. 307, pl. xlix, fig. Reisen Philipp. ii, v, n. spp.

Tapena igna, Philippines, Semper, p. 316, Reisen Philipp. ii, v, n. sp. Teniorhinus watsoni, Gaboon, Holland, p. 292, Ann. N. H. (6) x, n. : Tricosemeia, n. g., for T. subolivescens, n. sp., W. Africa, Holland, p. 2 Ann. N. H. (6) x.

#### HETEROCERA.

- Catalogue of Lepidoptera Heterocera; vol. i, Sphinges, Bombyces; KII (496).
- The classification, phylogeny, neuration, &c., of Heterocera discuss Hampson, Moths of India, i, pp. 1-12.
- 707 species of Amurland Heterocera of the groups Sphinges, Bombs Noctues, catalogued and discussed seriatim; STAUDINGER in Roma: Mem. vi, pp. 219-658.
- Notes on many species, with named varieties, of *Heterocera* from Sc Siberia; STAUDINGER, Deutsche e. Z. Lep. v, pp. 342-393.
- Brief notes on some of the early stages of numerous Australian Hecera, by the late Henry Edwards; Brutenhüller, Bull. Mus. iv, pp. 71-74.
- List of 12 genera established by Philippi, and of the genera of w they are synonyms; Reed, pp. 68 & 69, Act. Soc. Chili, i.
- Experiments as to colour variation in Lepidopterous larvæ, and p and in occoons; BATESON (40, 41), GOULD (354), POULTON (702 Walker's types and their labelling noticed; GROTE, p. 189, Canad. xxiv.

#### SPHINGIDA.

- [Cf. Aurivillius (24), Austaut (29), Cosmovici (191), Graeseb ( Karsch (460, 461), Lucas (568), Miskin (611, 612), Staudi (779), Swinhoe (908).]
- Revision of the Australian Sphingida; MISKIN (611, 612).
- Synonymy of the species described by Lucas in "Queenalander, Miskin at end of "Further note" (612).
- Brief note on phylogeny; GROTE, p. 17, Canad. Ent. xxiv.
- Abrisa, n. n. for Microlophia, Feld.; KIRBY, p. 641, Cat. Lep. Het. Acanthosphinx, n. g. for Ambulyx gussfeldtii, Dew., with n. var. g. Aurivillius, Ent. Tidskr. xiii, pp. 181 & 182.
- Acherontia atropos, stridulation of; Roth, p. 250, Ent. Tidskr. xiii.

  Acosmeryx mixtura, figured, pl. i, fig. 7; SWINHOE, Cat. Lep. Oxfo.

  Ambulyx wilder, Queensland, Miskin, P. R. Soc. Queensl. viii, pt.

  n. sp.
- Anceryx fasciata, larva described; Cockerell, Ent. xxiv, p. 19. Charocampa rosetta, p. 16, insularis, p. 18, Ceram, amara, Mysol, pl. i, fig. 9, Swinhoe, Cat. Lep. Oxford; C. cleopatra, cloacina, pallida, Queensland, Miskin, P. R. Soc. Queensl. viii, pt. 3, pp. 18, sep. copy): n. spp.

- Deilophila japonica, var. ? suifuna, figured, pl. iv, fig. 2, described, p. 228; STAUDINGER, Romanoff Mem. vi.
- .D. proxima, Japan, Austaut, p. 69, Le Nat. 1892; D. euphorbioides, Japan, Swinhoe, p. 22, Cat. Lep. Oxford: n. spp.
- Dolbina exacta, Amurland, STAUDINGER, p. 222, pl. iv, fig. 1, Romanoff Mem. vi, n. sp.
- Gargantua, n. n. for Lintneria, Butl.; KIRBY, p. 692, Cat. Lep. Het. i.
- Hemaris rubra, N. India, Hampson, Moths of India, i, p. 120; H. janus, Queensland, Miskin, P. R. Soc. Queensl. viii, pt. 3, p. ?: n. spp.
- Kentochrysalis (sub Sphinx) streckeri, Stgr., = (davidis, Oberth.); STAUDINGER, p. 220, Romanoff Mem. vi.
- Lophura hyas, figured, pl. i, fig. 2, Romanoff Mem. vi.
- Macrosila severina, Cape York, MISKIN, P. R. Soc. Queensl. viii, pt. 3, p.? n. sp.
- Macroglossa nigra, Roumania, Cosmovici, p. 280, Le Nat. 1892; M. semifasciata, Pegu, Hampson, Moths of India, i, p. 115; M. limata, Java, pl. i, fig. 1, p. 4, kanita, fig. 2, Sumatra, labrosa, Bouru, p. 5, inusitata, Dorey, p. 6, Swinhoe, Cat. Lep. Oxford: n. spp.
- Nephele pachyderma, W. Africa, Karsch, p. 180, Ent. Nachr. xviii, n. sp. Ocyton radiata, W. Africa, Karsch, p. 116, Ent. Nachr. xviii, n. sp.
- Pachylia ficus, notes on larva; COCKERELL, Ent. News, iii, p. 4.
- Panacra variolosa, figured, pl. i, fig. 4; SWINHOE, Cat. Lep. Oxford.

  P. turneri, Lucas, redescribed; Miskin, P. R. Soc. Queenal. viii, pt. 3, p.?
- P. frena, Silhet, pl. i, fig. 5, p. 48, mira, Cape York, p. 13, pl. i, fig. 6, Swinhoe, Cat. Lep. Oxford, n. spp.
- Protoparce cæstri, Blanch., = (Sphinx surylochus, Phil.); REED, p. 68, Act. Soc. Chili, i.
- P. minimus, Queensland, MISKIN, P. R. Soc. Queensl. viii, pt. 3, p.?, n. sp. Pseudosphinz grisea, p. 104, concolor, p. 106, India, Hampson, Moths of India, i, n. spp.
- Pterogon kuldjaensis, Centr. Asia, GRAESER, p. 299, Berl. Eut. Zeit. xxxvii, n. sp.
- Smerinthus populi, anatomy of hermaphrodite; TETENS (918). S. tiliæ and populi, note on markings of larvæ; Gould, pp. 236-243, Tr. Ent. Soc. 1892.
- S. heynei, Japan, Austaut, p. 68, Le Nat. 1892, n. sp.
- Sphinz eremophilæ, redescribed; Lucas, p. 277, P. Linn. Soc. N.S.W. (2) vi.
- S. marmorata, Queensland, Lucas, P. Linn. Soc. N.S.W. (2) vi, p. 278, n. sp.

### ÆGERIIDE.

[Gf. Druce (226), Hampson (384), Kellicott (476, 477), Mibrin (613), Staudinger (886), Swinhoe (908).]

Adizoa, n. g. for A. auricollum, n. sp., Moths of India, i.

Egeria corni, habits and characters, K xxiv, and metamorphoses, Ina. Life on; Kellicott, Canad. Ent. xxiv, p

Æ. pensilis, Sarawak, SWINHOE, p. 36, stigma, KELLICOTT, p. 211, Canad. E. Æ. armasata, mardia, Mexico, DRUG. n. spp.

Albuna modesta. Ohio, KELLICOTT, p. 46, Anthrenoptera, n. g., p. 35, for Sphecia fig. 11; SWINHOE, Cat. Lep. Oxford.

Aschistophleps, n. g. for A. lumpropoda, p India: HAMPSON, Moths of India, i.

Ceratocorema, n. g. for C. postcristatum, n. India, i, p. 200.

Ichneumenoptera, n. g. for I. auripes, | flaripalpus, ignifera, p. 195, n. spp., India, i.

Macrotarsipus, n. g. for M. albipunctus, n. Moths of India, i.

Melittia ceto, notes on; Smith, p. 129, Ca M. grandia, Sikkim, Hampson, Moths of N. China, Swinhoe, p. 38, Cat. Queensland, Miskin, P. R. Soc. beckeri, Mexico, Druce, p. 276, Ann.

Sciapteron ferridum, varr. described; Sta Lep. iv.

S. splendidum, Taurus, p. 242, S.? dispa Deutsche e. Z. Lep. iv; S. flamman metallicum, cupreivitta, tenuimarginat Moths of India, i: n. spp.

Sesia proximata, Amasia, p. 244, regule Deutsche e. Z. Lep. iv; S. qu xanthosticta, p. 197, E. India, HAMPS

Sphecodoptera, n. g., p. 189, type repand n. sp., Kashmir, p. 190: Hampson, 1

Trichocerota, n. g. for T. ruficineta, Peg India, i.

Trilochana ignicauda, Burma, HAMPSON, Trochilium apiforme, n. var. cafischii; Ent. Ges. viii.

T. ommatiæforme, ignicolle, India, HAMF T. cupreifascia, Queensland, MISKIN, p. ?: n. spp.

# CASTNUDE.

[Cf. Lower (566), Swinhoe (908).]

Synemon collecta, p. 151, pl. vii, fig. 16, adelaida, fig. 10, p. 154, Australia, Swinhoe, Cat. Lep. Oxford; S. pyrrhoptera, Australia, Lower, p. 13, Tr. R. Soc. S. Austr. xvi: n. spp.

## BOMBYCES.\*

- [Cf. Alpheraky (779), Aurivillius (24), Butler (141, 146), Carpentier (160), Christoph (175), Distant (210, 216), Dognin (220), Druce (226), Dyar (238, 239, 240), Fenk (279), Graeser (356, 357), Heylaerts (403), Holland (408, 409), Honrath (413), Karsch (460, 464), Kirby (497), Lower (565, 566), Lucas (568, 569), Mabille (571, 572), Mabille & Vuillot (573), Moore (623), Neumoegen (633, 634), Oberthur (650), Packard (665, 666), Poujade (702), Püngeler (706), Röber (767), Schaus (798, 799), Slosson (854, 855), Sonthonnax (878), Standfuss (779), Staudinger (779, 886, 890, 891), Swinhoe (908), Vuillot (978, 980), Weymer (1003).]
- The following new or revived names of families are recognized in the first volume of Hampson's Moths of India, viz., Brahmæidæ, Eupterotidæ, Syntomididæ, Arbelidæ, Pterothysanidæ.
- KIRBY, in his catalogue, makes use of the names Cyllopodidæ, Cymbidæ, and Pinaridæ, to represent families of Bombyces.
- Achelura, n. n. for Chelura, Hope; KIRBY, p. 56, Cat. Lep. Het. i.
- Aclytia petra, Peru, p. 276, hecale, p. 277, Brazil, Schaus, P. Z. S. 1892 (figured, Schaus, American Lep. pl. i), n. spp.
- Adelocephala invalida, Brazil, Schaus, p. 289, P. Z. S. 1891 (figured, Schaus, pl. iii, American Lep.); A. columbia, Columbia, Schaus, p. 21, pl. iii, fig. 13, American Lep.: n. spp.
- Adlullia lunifera, pl. vi, fig. 3, innotata, fig. 1, xanthomela, fig. 8, divisa, fig. 14, figured; SWINHOE, Cat. Lep. Oxford.
- A. bollora, Borneo, SWINHOE, p. 186, pl. vi, fig. 2, Cat. Lep. Oxford, n. sp. Egocera and Hecatesia, stridulation in; Hampson, pp. 188 & 189, P. Z. S. 1892. E. triphænoides, synonymical note; Weymer, p. 124, S. E. Z. liii.
- Æ. confluens, Zanzibar, WEYMER, p. 100, S. E. Z. liii, n. sp.
- Æmilia, n. n. for Ameles, Walk.; KIRBY, p. 218, Cat. Lep. Het. i.
- Agarea minuta, Brazil, SCHAUS, p. 288, P. Z. S. 1892, n. sp.
- Agaristida, definition of; AURIVILLIUS, Ent. Tidskr. xiii, p. 183.

Owing to increasing divergence of opinion as to the limits and positions of the families hitherto placed between the Sphingid and the Noctuid Moths, and to the large number of new genera that have been established of late years, we are compelled to unite the groups under the comprehensive term Bombyess.

- Agarista albamedia, simplex, p. 301, tropica, p. 302, caruleotincta, p. 30 Queensland, Lucas, P. Linn. Soc. N.S.W. (2) vi, n. spp.
- Aglia tau, life-history; Andersson, Ent. Tidskr. xiii, pp. 93-96.
- Agnidra carnea, Borneo, BUTLER, p. 125, pl. vi, fig. 3, P. Z. S. 1892, n.
- Agonis lutensis, Ternate, SWINHOE, p. 136, Cat. Lep. Oxford, n. sp.
- Albara reversaria, figured, pl. vii, fig. 9; SWINHOE, Cat. Lep. Oxford Allodonta? leucodera, Amurland, STAUDINGER, p. 357, Romanoff Me vi, n. sp.
- Aloa gangara, pl. iv, fig. 2, corsima, fig. 1, SWINHOE, p. 171, Cat. L Oxford, n. spp.
- Alophogaster, n. g., Phaudinæ, for A. rubribasis, Nága Hills, n. sp., HAI SON, p. 287, Moths of India, i.
- Alpenus trifasciata, E. Africa, Holland, Ent. xxv (Supp.), p. 92, n. sp
- Alypia crescens, Wlk., = (grutei, Bdv.); SMITH, p. 133, Canad. Ent. x7
- Amaxia hebe, p. 278, pyga, p. 279, Brazil, Schaus, P. Z. S. 1892 (figur Schaus, American Lep. pl. i): n. spp.
- Ameles byblis, Brazil, Schaus, p. 279, P. Z. S. 1892 (figured, Schauerican Lep. pl. ii, fig. 2), n. sp.
- Amerila nivaria, p. 105, phædra, p. 106, E. Africa, WEYMER, S. E. Z. n. spp.
- Ammalo thrailkilli, Mexico, Schaus, p. 280, P. Z. S. 1892, n. sp.
- Amydona sericea, Peru, Schaus, p. 324, P. Z. S. 1892, n. sp.
- Anaphe: characters and position noticed; Butler, p. 398, Ann. N. H. x. A. carteri, Walsingh., = (parra, Aur); Aurivillius, p. Ent. Tidskr. xiii.
- A. aurea, Madagascar, BUTLER, p. 398, Ann. N. H. (6) x, n. sp.
- Anaxidia, n. g., Limacodida, for A. luctea, n. sp., W. Australia; Swini p. 231, Cat. Lep. Oxford.
- Anisota stigma, metamorphoses; MARLATT, pp. 111 & 112, Tr. Ka Ac. xi, figs. A. stigma, p. 65, senatoria, p. 66, metamorpho Beutenmüller, Bull. Am. Mus. iv.
- Antarctia rubra, var., = (walsinghami, Butl.); Smith, p. 135, Canad.
- Antheræa anthina, Cameroons, Karsch, p. 353, Ent. Nachr. xviii knyretti, Sikkim, Hampson, Moths of India, i, p. 19; A. pull fasciata, p. 448, olivescens, ochripicta, p. 449, Bombay, versie Bengal, hartii, N. China, p. 450, borneensis, p. 451, ridlyi, p. Borneo, surakarta, Java, p. 453, Moore, Ann. N. H. (6) ix alephostra, Waigiou, Swinhoe, p. 252, Cat. Lep. Oxford; A. lore Queensland, Lucas, p. 292, P. Linn. Soc. N.S.W. (2) vi: n. spp.
- Antipha biplagata, Java, HEYLAERTS, p. 10, Ann. Ent. Belg. xxxvi, Aon, n. g., Cossidæ, for A. noctuiformis, n. sp., Texas; NEUMOEGEN,
- News, iii, p. 258.

  Aræocera, n. g., Zygænidæ, for A. cyanescens, posthyalina, n. spp., II

  HAMPSON, Moths of India, i, p. 244.
- Aræogyia, n. g., Limacodida, p. 394, for A. spatulata, castanea, n. India, p. 395; Hampson, Moths of India, i.

- Archylus mexicana, Jalapa, p. 288, vesta, Brazil, p. 289, Schaus, P. Z. S. 1892, n. spp.
- Arctia fortunata and glauca, notes on, with figg.; STAUDINGER, pp. 250 & 251, pl. iii, figs. 3 & 5, Deutsche e. Z. Lep. iv. A. mongolica, figured, pl. i, fig. 6, Romanoff Mem. vi. A. nais; rhoda and ochreata, Butl., and radians, Wlk., are probably varr. of this: A. phyllira, = (dodgei, Butl.); SMITH, pp. 134 & 135, Canad. Ent. xxiv. A. caju, var.; LADDIMAN, Ent. xxv, p. 1.
- A. flava, n. var. campestris; GRAESER, p. 213, Berl. Ent. Zeit. xxxvii.
- A. turbans, Irkutsh, Christoph, p. 460, Hor. Ent. Ross. xxvi; A. minea, N. America, Slosson, p. 257, Ent. News, iii: n. spp.
- Ardices garida, Australia, SWINHOE, p. 179, Cat. Lep. Oxford, n. sp.
- Ardonea metallica, Peru, Schaus, p. 284, P. Z. S. 1892 (figured, pl. ii, fig. 8; Schaus, American Lep.), n. sp.
- Areas sparsalis, conflictalis, rudis, Wlk., figured; SWINHOE, pl. iv, figg. 4-6, Cat. Lep. Oxford.
- Argina pulchra, Philippines, SWINHOE, p. 117, pl. iii, fig. 5, Cat. Lep. Oxford, n. sp.
- Arguda rectilinea, Burma, HAMPSON, p. 414, Moths of India, i, n. sp.
- Argyroeides magon, Brazil, Schaus, p. 276, P. Z. S. 1892, n. sp.
- Aroa atrella, p. 439, aurantifascia, p. 440, India, Hampson, Moths of India, i; A. exoleta. Queensland, SWINHOE, p. 197, pl. vi, fig. 18, Cat. Lep. Oxford, n. spp.
- Arotros, n. g., Bombycidæ, p. 325, for A. striata, n. sp., Brazil, p. 326; SCHAUS, P. Z. S. 1892.
- Artace anula, meridionalis, Brazil, Schaus, p. 319, P. Z. S. 1892, n. spp.
- Artaxa chrysophea, n. var. iobrota; Lucas, p. 251, P. Linn. Soc. N.S.W.

  (2) vii. A. signiplaga, figured; SWINHOE, pl. vi, fig. 9, Cat. Lep.
  Oxford. A. charmetanti, figured; MABILLE & VUILLOT, pl. xi, fig. 2,
  Nov. Lep.
- Artaza albinotata, p. 9, bicolor, p. 10, Sumatra, HEYLAERTS, Ann. Ent. Belg. xxxvi; A. singapura, Singapore, p. 189, pl. vi, fig. 19, enunciativa, Philippines, p. 190, SWINHOE, Cat. Lep. Oxford; A. lucifuga, Brisbane, Lucas, p. 250, P. Linn. Soc. N.S.W. (2) vii: n. spp.
- Artona sieversi, Mongolia, Alpheraky, p. 5, pl. i, fig. 4, Romanoff Mem. vi, n. sp.
- Asterocopus nodosus, Tasmania, SWINHOE, p. 299, pl. viii, fig. 13, Cat. Lep. Oxford, n. sp.
- Asura? bisecta, Queensland, Lucas, P. Linn. Soc. N.S.W. (2) vi, p. 279, n. sp.
- Atasca, n. g. for pt. of Nyctemera, p. 139, A. simplex and quadriplaga, Wlk., figured, pl. v, figs. 13 & 9; SWINHOB, Cat. Lep.
- Atossa palæarctica, Stdgr., systematic position; SNELLEN, p. 11, Tijdschr. Ent. xxxv.
- Attacus atlas and allies, synonymy; KIRBY, Ent. xxv, pp. 186-188.

- Automeris macareis, rubicunda, Braxil, nopalisin, Mexico, p. 290, tamphilus, Braxil, p. 291, Schaus, P. Z. S. 1892 (A. macareis, rubicunda, figured, pl. iii, Schaus, American Lep.); A. quadridentata, Braxil Kirby, p. 175, pl. xi, fig. 3, Ann. N. H. (6) x: n. spp.
- Automolidina, = (Hexaneurida, Wallengr.), composition of ; Aurivillius p. 189, Ent. Tidskr. xiii.
- Automolis elissa, Brazil, SCHAUS, p. 277, P. Z. S. 1892 (figured and referred to Euplesia, SCHAUS, p. 7, pl. i, fig. 21, American Lep.) n. sp.
- Asygophleps, n. g., Cossidæ, type scalaris, Fab.; Hampson, p. 309, Moth of India, i.
- Balacra elegans, p. 190, pulchra, B.? croceipes, p. 200, Cameroom Aurivillius, Ent. Tidskr. xiii, n. spp.
- Barsine hypoprepoides, fig. 22, placens, fig. 15, figured, pl. iii, SWINHOI Cat. Lep. Oxford.
- B. lucibilis, Borneo, pl. iii, fig. 6, p. 107, intrita, Ceram, fig. 23, p. 104 Swinhoe, Cat. Lep. Oxford, n. spp.
- Bharetta flammans, Manipur, Hampson, p. 416, Moths of India, i, n. sp. Bijugis sp., larva and female noticed; Püngeler, p. 139, Deutsche e. Lep. v.
- Birthama obliqua, Wlk., figured, pl. viii, fig. 9; SWINHOE, Cat. Le Oxford.
- B. bilineata, Philippines, SWINHOE, p. 240, Cat. Lep. Oxford, n. sp.
- Bizone dubenskii, China, Alpheraky, p. 11, pl. i, fig. 5, Romanoff Mer vi, n. sp.
- Blera causia, p. 332, bianca, apella, sida, Brazil, p. 333, SCHAUS, P. Z. 1892, n. spp.
- Bombyx trifolii, n. var. bathseba; STAUDINGER, p. 260, Deutsche e. Z. Le iv. B. quercus and callunæ, effects of temperature in produci variation; MERRIFIELD, p. 38, Tr. E. Soc. 1892. B. mori, existen of clinging papillæ in abdominal legs of larva; VERSON, Zool. At xv, p. 279: embryology; HENKING (396): food-plants; Hu (442).
- B. salomonis, Beyrout, STAUDINGER, p. 259, Deutsche e. Z. Lep. iv; rueckbeili, Central Asia, GRAESER, p. 301, Berl. Ent. Zeit. xxxv B. (Lebeda) rotundata, Java, HEYLAERTS, Ann. Ent. Belg. xxx p. 8: n. spp.
- Borocera madagascarieneis, notes on; SIKORA, p. 179, Soc. Ent. vi.
- B. esteban, Lojn, Dognin, p. 33, Le Nat. 1892, n. sp.
- Brachartona catoxantha, Tenasserim, Hampson, p. 233, Moths of India n. sp.
- Bremeria, n. g., near Rhaphidognatha, for B. manza, n. sp., China, p fig. 3; Alpheraky, p. 7, Romanoff Mem. vi.
- Brycea peruviana, Peru, Schaus, p. 284, P. Z. S. 1892 (figured, pl. fig. 16, Schaus, American Lep.), n. sp.
- Bunæa mitfordi, Sierra Leone, KIRBY, p. 173, pl. xi, fig. 1, Ann. N. H. x, n. sp.

- Caligula castanea, Japan, Swinhoe, p. 249, Cat. Lep. Oxford, n. sp.
- Callartona, n. g., Zygænidæ, type purpurascens, Hmpn.; Hampson, p. 233, Moths of India, i.
- Calligenia pilcheri, Rockhampton, Lucas, P. Linn. Soc. N.S.W. (2) vi, p. 279, n. sp.
- Callimorpha suffusa, metamorphoses; MARLATT, p. 113, figg., Tr. Kansas Ac. xi. C. lecontei, localisation of variation; MASON, Ent. News, iii, p. 52.
- Callitomis multifasciata, India, Hampson, Moths of India, i, p. 225, n. sp. Callisygæna auricincta, Nilgiri Hills, Swinhoe, p. 62, Cat. Lep. Oxford, n. sp.
- Calybia, n. n. for Phryns, Grote; KIRBY, p. 446, Cat. Lep. Het. i.
- Camptochilus, n. g., Drepanulidæ, type reticulatum, Moore; HAMPSON, p. 351, Moths of India, i.
- Caprima gelida, fig. 3, C. mutilata, Wlk. (sub Soritia), fig. 15, figured, pl. ii; SWINHOE, Cat. Lep. Oxford.
- Carales celer, Brazil, SCHAUS, p. 282, P. Z. S. 1892, n. sp.
- Cascera inconcisa, Mysol, SWINHOE, p. 299, pl. vii, fig. 6, Cat. Lep. Oxford, n. sp.
- Ceratonema, n. g., Limacodidæ, p. 393, type retractatum, Wlk., with C. albifusum, fasciatum, n. spp., India, p. 394; Hampson, Moths of India, i.
- Cerura: list of the N. American species; Dyar, Psyche, vi, p. 290: mode of emergence from cocoon; Chapman, Ent. xxv, pp. 302-304.
- C. meridionalis, Texas, Dyar, Psyche, vi, p. 291, n. sp.
- Chalcocelis, n. g., Limacodidæ, type fumifera, Swinh.; Hampson, p. 392, Moths of India, i.
- Chalcosia fragilis, Wlk., figured, pl. ii, fig. 7; SWINHOE, Cat. Lep. Oxford. C. griseifusa, Pegu, Hampson, p. 264, Moths of India, i; C. paviei, Laos, Poujade, p. 268, pl. xi, fig. 5, N. Arch. Mus. Paris (3) iii; C. indistincta,
- Borneo, SWINHOE, p. 74, Cat. Lap. Oxford: n. spp. Chalina, n. subfam. of Psychida; Hampson, Moths of India, i, p. 300.
- Chalioides, n. g., p. 227, for C. vitrea, n. sp., Ceylon, p. 228; Swinhoe, Cat. Lep. Oxford.
- Charagia pomalis, Queensland, pl. viii, fig. 3, dulcis, Swan River, SWINHOE, p. 288, Cat. Lep. Oxford, n. spp.
- Charidea inachia, Brazil, Schaus, p. 277, P. Z. S. 1892 (figured and referred to Cyanopepla, Schaus, American Lep. i, p. 6, pl. i, fig. 19), n. sp.
- Chelonia mirifica, Ta-Tsien-Lou, OBERTHUR, p. 8, pl. i, fig. 7, Études d'Ent. xvi, n. sp.
- Chenuala, n. g., Liparidæ, for C. rufa, n. sp., Queensland; SWINHOE, p. 212, Cat. Lep. Oxford.
- Chondrostega pastrana, n. var. palæstrana; STAUDINGER, p. 257, Deutsche e. Z. Lep. iv.
- C. fasciana, Jerusalem, Staudinger, Deutsche e. Z. Lep. iv, p. 258,

Chrysartona, n. g., Zyganida, for Procris st. Cat. Lep. Oxford.

C. refulgens, Burma, Hampson, p. 232, Moth Chrysopsyche mirifica, coccon figured; Hot p. 216, Psyche, vi.

Cispia flavipes, Sikkim, Hampson, p. 493, Mc Cisthene petrovna, Brazil, Schaus, p. 283, fig. 4, Schaus, American Lep.), n. sp.

Citheronia vogleri, = (Eacles leona, Druce), fi American Lep.

Clelea plumbeola, Burma, Hampson, p. 240, Clisiocampa constricta, var. = (strigosa, Strevi. C. erosa, metamorphoses; Dyar, t.

Cochliotheca: habits, systematic position; P. v, p. 133.

Codiosoma, n. n. for Kodiosoma, Stretch; K. Calodasys tonac, p. 330, pegasis, p. 331, B. n. spp.

Collita leucanioides, Wlk. (sub Lithosia), figu: Cat. Lep. Oxford.

Comacla simplex, Wlk., = (murina, Wlk., and SMITH, p. 134, Canad. Ent. xxiv.

Contheyla melanosticta, India, HAMPSON, p. 3 Corcura torta, fig. 10, discalis, fig. 12, figured Oxford.

C. mysolica, Mysol, Swinhor, p. 119, pl. ii n. sp.

Corma maculata, Burma, Hampson, p. 268, Cosmosoma harpulyce, Petropolis, Schaus, p. Schaus, American Lep. pl. i, fig. 11), n.

Cossa vagivitta, Wlk. (sub Lithosia), figured, Lep. Oxford.

Cossidæ and Hepialidæ, wing-neuration disc Zool. liii, pp. 635-639.

Cossula notodontoides, præclara, Brazil, Schin. spp.

Cossulinus, n. n. for Cossulus, Stdgr.; KIRBY, Cossus ussuriensis, E. Siberia, GRAESER, p. 2
C. nigromaculatus, p. 305, parvipunctus
HAMPSON, Moths of India, i; C. chloratu
pl. viii, fig. 8, Cat. Lep. Oxford; C. pari
p. 327, P. Z. S. 1892: n. spp.

Costria abnoba, Brazil; SCHAUS, p. 327, P. Z. Cosuma marginata, E. Africa, HOLLAND, Ent. Cothocida nigrifera, Wlk., referred to Crocol Ent. xxiv.

Crambidia pallida, egg and young larva; Dy

- Crambomorpha auraria, Loja, Dognin, p. 169, Le Nat. 1892, n. sp.
- Crinocraspeda, n. g., Lasiocampida, type'torrida, Moore; Hampson, p. 420, Moths of India, i.
- Crocomela, n. n. for Durna, Wlk.; KIRBY, p. 405, Cat. Lep. Het. i.
- Crothama decorata, Transvaal, DISTANT, p. 237, pl. ii, fig. 6, Naturalist Transvaal, n. sp.
- Ctenucha azteca, Mexico, SCHAUS, p. 278, P. Z. S. 1892, n. sp.
- Cyclosia ficta, Wlk., referred to Milleria, p. 70, and figured, pl. ii, fig. 4; Swinhoe, Cat. Lep. Oxford.
- C. chuna, Menado, p. 68, enodis, Sumatra, pl. ii, fig. 2, chartacea, Manilla, fig. 3, p. 69, SWINHOE, Cat. Lep. Oxford, n. spp.
- Cyllopodidæ, n. n. for a part of the family Chrysaugidæ; KIRBY, Cat. Lep. Het. i, p. 398.
- Cyllopoda (Flaviniu) dubia, durno, janeira, quicha, chibcha, Schaus, figured; SCHAUS, American Lep. pl. ii.
- Cyphanta chortochlora, Himalaya, Hampson, Moths of India, p. 175, n. sp.
- Dactylorhyncha, n. g., Liparidæ, type pallida, Hmpson.; Hampson, p. 470, Moths of India, i.
- Dalcera tijucana, Brazil, SCHAUS, p. 322, P. Z. S. 1892, n. sp.
- Dalaca prytanes, p. 329, mummea, oreas, Brazil, terea, Mexico, p. 330, Schaus, P. Z. S. 1892, n. spp.
- Damata microsticta, E. India, SWINHOE, p. 302, Cat. Lep. Oxford; D. nigrosticta, E. India, HAMPSON, Moths of India, i, p. 157, n. spp.
- Damias angustifasciata, Ké Is., Röber, p. 324, Tijdschr. Ent. xxxiv (fg. in xxxv), n. sp.
- Danima banksia, larva noticed; ASH, Ent. M. M. xxviii, p. 232, woodcut.
- Darala complens, p. 209, uniformis, minuta, p. 210, Australia; SWINHOE, Cat. Lep. Oxford; D. ochropteru, Centr. Australia, LOWER, p. 14, Tr. R. Soc. S. Austr. xvi; D. expansa, magnifica, p. 286, asciscens,
  - p. 287, linearis, p. 289, succinea, scortea, p. 290, rubriscripta, roseu, p. 291, Australia, Lucas, P. Linn. Soc. N.S.W. (2) vi : n. spp.
- Darna metaleuca, Wlk. (sub Artaxa), pl. vii, fig. 14, basalis, Wlk. (sub Susica), pl. viii, fig. 11, figured; SWINHOE, Cat. Lep. Oxford.
- D. inca, Peru, Schaus, p. 286, P. Z. S. 1892 (figured, Schaus, American Lep. pl. ii, fig. 10), n. sp.
- Dasychira chloroptera, India, HAMPSON, p. 450, Moths of India, i; D. tristis, Sumatra, p. 12, D. (Mardara) preangerensis, Java, p. 13, HEYLAERTS, Ann. Ent. Belg. xxxvi: n. spp.
- Dasylophia puntagorda, Florida, Slosson, p. 139, Canad. Ent. xxiv, n. sp. Datana major, metamorphoses; Beutenmuller, p. 76, Bull. Am. Mus. iv.
- Demas coryli, habits noticed; STILL, Ent. M. M. xxviii, p. 148.
- Dendrophleps, n. g., Liparidæ, for D. semihyalina, n. sp., Khasi Hills; Hampson, p. 491, Moths of India, i.
- Diabasis nilgirensis, n. n. for Bijugis sikkimensis, Heyl.; HAMPSON, p. 303, Moths of India, i.

Dicranura vinula, escape from cocoon by mea LATTER, Tr. E. Soc. 1892, p. 287.

Digama burmana, Myingan, Hampson, p. 505 Dirphia aculea, Braxil, Vuillot, p. exci, Bull Dixoa, n. g., Thyridida, type albatalis, Swinh of India, i.

Dolecta juturna, macrochir, invenusta, Brazi 1892, n. spp.

Dolgoma nebulosa, Wlk. (sub Lithosia), figure Cat. Lep. Oxford.

Dreata pomona, E. Africa, WEYMER, p. 113, S. Drepana curvaria, Wlk. (sub Canucha), pl. vii. figured; SWINHOE, Cat. Lep. Oxford.

D. grisearia, pl. iv, fig. 7, muscula, pl. xii, fig. p. 335, Romanoff Mem. vi; D. rufofi innotata, fasciata, p. 335, perhamata, p. i p. 338, pulcherrima, p. 339, xanthopteru, Hampson, Moths of India, i: n. spp.

Drucea, n. n. for pt. of Mastigocera; KIRBY, Drymæa, Butl.: characters and systematic pc pp. 20-23, Tijdschr. Ent. xxxv.

Duga rana, Goram, SWINHOE, p. 96, pl. iii, fig Duomitus kilimanjarensis, E. Africa, HOLLAI

D. fuscipars, Sikhim, Hampson, p. 309, I Dycladia catherina, rogenhoferi, epimethew Brazil, Schaus, P. Z. S. 1892 (figure pl. i), n. spp.

Dysodia albifurca, Sikkim, Hampson, p. 369, Earias chlorophyllana, Mardin, Staudinger iv, n. sp.

Echedorus argentina, Brazil, Schaus, p. 321, l Ecpantheria pellucida, Brazil, Schaus, p. 2 Schaus, American Lep. pl. iii, figs. 11 &

Ectrocta, n. g., Bombycidæ, for E. diaphana, Moths of India, i, p. 33.

Eligma duplicata, Trop. Africa; AURIVILLIU Ellida (sub Edema) transversata, Wlk., = (ge Canad. Ent. xxiv.

Encaumaptera, n. g., Arbelidæ, p. 315, for i p. 316, Hampson, Moths of India, i.

Endagria fuscula, Tunis, STAUDINGER, p. 2 e. Z. Lep. v; E. cossoides, Centr. Asia, G Zeit. xxxvii, n. sp.

Ephemeroidea, n. g., Zygænidæ, for E. ariel, Hampson, p. 242, Moths of India, i.

Epimydia, n. g. for E. dialampa, n. sp., Siberis p. 346, Deutsche e. Z. Lep. v.



- Eremocossus, n. g., Cossidæ, type fædus, Swinh.; Hampson, p. 313, Moths of India, i.
- Eressa politula, India, p. 52, detola, olinda, Australia, p. 53, SWINHOE, Cat. Lep. Oxford, n. spp.
- Eriogaster lanestris, variation in colour of cocoons discussed; BATESON (40, 41).
- Eriphioides, n. n. for Eriphia, Feld.; KIRBY, p. 159, Cat. Lep. Het. i. Eterusia submarginalis, Cambodia, SWINHOE, p. 81, Cat. Lep. Oxford, n. sp.
- Etobesa tizoc, Peru, SCHAUS, p. 339, P. Z. S. 1892, n. sp.
- Eucereon ladas, Brazil, Schaus, p. 278, P. Z. S. 1892 (figured, Schaus, American Lep. pl. i, fig. 26); E. clementsi, St. Lucia, Schaus, p. 10, pl. i, fig. 25, t. c. : n. spp.
- Eudelia venusta, Wlk., = (Bombyx aristotelia, rufescens, Phil.); REED, p. 68, Act. Soc. Chili, i.
- Eudoxyla bosschæ, Borneo, HEYLAERTS, p. 45, Ann. Ent. Belg. xxxvi, n. sp.
- Eudryas grata, var., = (stos. johannis, Wlk.), note on; SMITH, p. 134, Canad. Ent. xxiv. E. stos. johannis, note on; GROTE, p. 236, Canad. Ent. xxiv.
- E. ? liturata, Cameroons, AURIVILLIUS, Ent. Tidskr. xiii, p. 187, n. sp.
- Eudule venata, Pern, aurata, Brazil, Schaus, p. 284, P. Z. S. 1892 (figured, Schaus, pl. ii. American Lep.), n. spp.
- Eulimacodes möschleri, Mexico, Schaus, p. 324, P. Z. S. 1892, n. sp.
- Eumeta javanica, Java, wallacei, Borneo, Swinhor, p. 226, Cat. Lep. Oxford, n. spp.
- Euproctis fulviceps, variation; MEYRICK, p ii, P. E. Soc. 1892.
- E. negrita, p. 471, arenacea, p. 476, antiphates, p. 478, flavimaculata, p. 480, albopunctata, nigripennis, p. 481, India, Hampson, Moths of India, i; E. (Chærotricha) sabulosa, p. 8, cinerea, p. 10, Java, Heylaerts, Ann. Ent. Belg. xxxvi: n spp.
- Eupyra psittacus, consors, aurata, p. 273, bacchans, p. 274, Peru, Schaus, P. Z. S. 1892 (figured, Schaus, American Lep. i), n. spp.
- Euschema: vide Geometridæ.
- Eusemia communis, fasciatrix, communicans, vetula, variation noticed, E. hesperioides, pl. v, fig. 3, saturata, fig. 1, figured; Swinhoe, p. 160, Cat. Lep. Oxford. E. opheltes, & figured; Poujade, pl. xi, figs. 4 & 4a, N. Arch. Mus. Paris (3) iii. E. xanthopyga, fig. 1, mosambica, fig. 2, longipennis, fig. 3, deficiens, fig. 4, pl. viii, novemmaculata, pl. ix, fig. 1, nigridorea, fig. 2, eoa, pl. xi, fig. 1, figured; Mabille & Vuillot, Nov. Lep. E. vilthoroides, figured; Oberthur, pl. i, fig. 4, p. 8, Études d'Ent. xvi.
- E. millionata, Philippines, pl. v, fig. 2, p. 162, doreana, Dorey, fig. 4, p. 164, SWINHOE, Cat. Lep. Oxford, n. spp.
- Eutane brevis, Borneo, SWINHOE, p. 126, pl. iii, fig. 13, Cat. Lep. Oxford, n. sp.
- Felderia, n. n. for Acreagris, Feld.; KIRBY, p. 163, Cat. Lep. Het. i.

- Flavinia dubia, darna, janeira, Brasil, quicha, p. 286, chibcha, p. 287, Peru, Schaus, P. Z. S. 1892 (cf. Cyllopoda), n. spp.
- Galethalea peruviana, Peru, Schaus, p. 277, P. Z. S. 1892 (figured, Schaus, pl. i, fig. 22, American Lep.), n. sp.
- Gangamela aymara, Peru, Schaus, p. 285, P. Z. S. 1892 (figured, Schaus, American Lep. pl. ii, fig. 9), n. sp.
- Gangarides vardena, Borneo, SWINHOE, p. 270, pl. vii, fig. 3, Cat. Lep. Oxford, n. sp.
- Gardinia, n. n. for Chrysocale, Butl.; KIRBY, p. 286, Cat. Lep. Het. i. Gargetta nagaensis, p. 135, curvaria, albimacula, p. 136, India, Hampsox.
- Moths of India, i, n. spp.
- Gartha dalsa, Brazil, Schaus, American Lep. i, pl. i, fig. 15, n. sp.
- Gastropacha quercifolia, scale-like hairs; PACKARD, Ann. N. H. (6) ix, p. 373.
- G. gabunica, W. Africa, Aurivillius, p. 199, Ent. Tidskr. xiii, n. sp. Givira tecmessa, philomela, Brazil, Schaus, p. 329, P. Z. S. 1892,
- Glaucopis iridea, Sierra Leone, Mabille, p. exxxviii, Bull. Soc. Ent. Fr. lxi, n. sp.
- Gluphisia trilineata, full-grown larva; Beutenmuller, p. 67, Bull. Am. Mus. iv.
- G. quinquelinea, Oregon, DYAR, p 168, Ent. News, iii, n. sp.
- Goë, n. g., Zygænidæ, for G. diaphana, n. sp., Burma: Hampson, p. 242, Moths of India, i.
- Gonimbrasia rubricostalis, Sierra Leone, Kirby, p. 174, pl. xi, fig. 2, Ann. N. H. (6) ix, n. sp.
- Gonometa sjostedti, Cameroons, Aurivillius, p. 199, Ent. Tidskr. xiii, n. sp.
- Gorgopis abbottii, E. Africa, Holland, Ent. xxv (Supp.), p. 94, n. sp.
- Gurna, n. g. for Dysauxes indica, Moore; SWINHOE, p. 123, Cat. Lep. Oxford.
- Gymnelia serra, Brazil, Schaus, p. 274, P. Z. S. 1892 (figured, Schaus, American Lep. pl. i, fig. 10), n. sp.
- Gynanisa isis, Westw., Q described; KIRBY, Ent. xxv, p. 267.
- Halias chlorana, variation in colour of cocoons; Tutt, Ent. Rec. iii, p. 9.
- Halisidota, notes on species of; DYAR, Canad. Ent. pp. 304-306. H. alwi and agassizii, larvæ noticed; DYAR, Psyche, vi, p. 323.
- H. sanguivenosa, Vancouver I., Neumoegen, p. 228, Canad. Ent. xxiv;
  H. oruba, Brazil, Schaus, p. 280, P. Z. S. 1892: n. spp.
- Hallicarnia, n. n. for Polymona, Wlk.; KIRBY, p. 822, Cat. Lep. Het. i. Haloplia, n. g., Drepanulidæ, for Nelcynda orciferaria, Wlk.; SNELLEN, p. 14, the insect figured, pl. i, fig. 1, Tijdschr. Ent. xxxv.
- Hanisa subnotata, figured, pl. viii, fig. 16; SWINHOE, Cat. Lep. Oxford.
- Hapigia xolotl, Mexico, p. 339, accipiter, phocus, p. 340, Brazil, Schaus, P. Z. S. 1892, n. spp.
- Harpyia? tenuis, Brazil, Schaus, p. 336, P. Z. S. 1892, p. sp.

- Harrisina janeira, dantasi, eminens, Brazil, Schaus, p. 272, P. Z. S. 1892 (figured, pl. i, figs. 1 & 2, Schaus, American Lep. i): n. spp.
- Hectomones fusca, crocea, Australia, LUCAS, p. 283, P. Linn. Soc. N.S.W. (2) vi; H. pteromela, S. Australia, LOWER, p. 5, Tr. R. Soc. S. Austr. xv: n. spp.
- Heliura venata, Butl., = (ælia, Schaus), figured, pl. i, fig. 20; Schaus, American Lep.
- Hepialiscus, n. g., Hepialidæ, type nepalensis, Wlk.; Hampson, p. 317, Moths of India, i.
- Hepialus (Charagia) blackburnii, S. Australia, LOWER, p. 5, Tr. R. Soc. S. Austr. xv; H. daphnandræ, hilaris, Australia; LUCAS, p. 284, P. Linn. Soc. N.S.W. (2) vi; H. keniæ, E. Africa, Holland, Ent. xxv (Supp.), p. 94: n. spp.
- Heterocampa læca, p. 333, hertha, limosa, virgea, Brazil, p. 334, epona, Pera, atrax, perilleus, Brazil, p. 335, Schaus, P. Z. S. 1892, n. spp.
- Heteropan = (Birtina, Wlk.) lycænoides, Wlk., figured, pl. ii, fig. 12; Swinhoe, Cat. Lep. Oxford.
- H. flavimacula, Burma, HAMPSON, p. 256, Moths of India, i, n. sp.
- Heterusia luteoradiata, Costa Rica, THIERRY-MIEG, p. 236, Le Nat. 1892, n. sp.
- Heylaertsia, n. subg. of Psyche for griseata, fusca, nudilineata, Hmpsn.; Hampson, p. 298, Moths of India, i.
- Hildala, n. g., Limacodida, p. 231, for H. miniacea, uniformis, n. spp., Port Essington, p. 232; Swinhoe, Cat. Lep. Oxford.
- Histia albimacula, Burma, Hampson, p. 280, Moths of India, i, n. sp.
- Hydrius bochica, Peru, theresa, braganza, lucrimosa, p. 320, sordida, p. 321, Brazil, Schaus, P. Z. S. 1892; H. victorio, Loja, pompilus (sic), Palandra, Dognin, p. 181, Le Nat. 1892; n. spp.
- Hydrusa chroma, Australia, p. 50, basirufa, Cambodia, p. 51, SWINHOE, Cat. Lep. Oxford; H. recedens, Queensland, Lucas, p. 281, P. Linn. Soc. N.S.W. (2) vi : n. spp.
- Hygrochroa limosa, Brazil, Schaus, p. 325, P. Z. S. 1892, n. sp.
- Hylesia muscula, Brazil, VUILLOT, p. cxcii, Bull. Soc. Ent. Fr. lxi, n. sp. Hyparpax venus, Colorado, NEUMOEGEN, p. 226, Canad. Ent. xxiv, n. sp.
- Hypatia, n. n. for Hysia, Wlk.; KIRBY, p. 147, Cat. Lep. Het. i.
- Hyperæschra nigribasis, Sikkim, Hampson, p. 165, Moths of India, i, n. sp.
- Hypolamprus, n. g., Thyridida, p. 364, type striatalis, Swinhoe, and including H. obscuralis, atrostriatus, n. spp., India, p. 365; Hampson, Moths of India, i.
- Hypoprepia miniata, egg and young larva; BLAISDELL, p. 246, Ent. News, iii.
- Hypsa perimele, Weym., characters and position noticed, p. 88, isthmia. Wlk., figured, pl. iii, fig. 2; SWINHOE, Cat. Lep. Oxford.
- Hypsa nicobarica, Nicobar Is., malisa, Halmaheira, p. 87, ghara, Ké I., p. 89, albiformis, Borneo, p. 90, ochrealis, Goram, avacta, Batchian, p. 92, SWINHOE, Cat. Lep. Oxford, n. spp.

- Ichyosoma cassiope, Brazil, Schaus, p. 341, P. Z. S. 1892, n. sp.
- Ichythyara bifiria, early stages; DYAR, Ent. News, iii, p. 5. Lran, larval stages; BEUTENMÜLLER, p. 67, Bull. Am. Mus. iv; I. albesigna, n. var. specifics; DYAR, Canad. Ent. xxiv, p. 180.
- multnoma. Oregon, Dyar, Canad. Ent. xxiv, p. 179; I. costiconna, India, Hampson, Moths of India, i, p. 173: Nilgiri Hills, Swinhor, p. 301, Cat. Lep. Oxford: n. app.
- Idalus ortus, Brazil, SCHAUS, p. 279, P. Z. S. 1892 (figured, SCHAUS, American Lep., pl. ii, fig. 1), n. sp.
- Idonauton, n. g. for Limacodes apicalis, Wik.; SWINHOE, p. 238, Cat. Lep. Oxford.
- Isa textula, notes on; MORTON, Ent. News, iii, p. 1.
- Isanthrene gaza. Peru, SCHAUS, p. 274, P. Z. S. 1892 (figured, pl. i, fg. 9, American Lep. i), n. sp.
- Jana gabunica, W. Africa, AURIVILLIUS, p. 195, Ent. Tidskr. xiii, n. sp.
- Janassa cerigoides, figured; SWINHOR, pl. vii, fig. 5, Cat. Lep. Oxford. Katha apicalis, Wlk. (sub Lithesia), figured, pl. iv, fig. 9; SWINHOR, Cat. Lep. Oxford.
- Kosala rufa, Nága Hills, HAMPSON. p. 419, Moths of India, i, n. sp.
- Lalia prolata, India, SWINHOE, p. 195, Cat. Lep. Oxford; L. heterogyas, Kashmir, Hampson, p. 443, Moths of India, i: n. spp.
- Laganda picaria, Wlk., figured, pl. vii, fig. 2, SWINHOR, Cat. Lep. Oxford.
- Lagoa to be the type of a new family, Lagoida; PACKARD, Psyche, vi, p. 281: on the seven pairs of abdominal legs and metameric glandular processes of larva; PACKARD, Zool. Ann. xv, pp. 229-234.
- Langedorfia polybia, Brazil, SCHAUR, p. 329, P. Z. S. 1892, n. sp.
- Lasiocampa dieckmanni, figured, pl. iv, fig. 5; Romanoff Mem. vi. L. leta, Wlk., figured, pl. v, fig. 1, larva noticed, p. 315; Staudinger, t. c.
- L (Chrysopoloma) theorini, Gaboon, AURIVILLIUS, p. 198, Ent. Tidskr. xiii; L. salammbo, Delagoa Bay, Vuillot, p. exc, Bull. Soc. Ent. Fr. lxi: n. spp.
- Lebeda doeri, Brazil, SCHAUS, p. 319, P. Z. S. 1892, n. sp.
- Leptosoma mundipicta, figured, pl. v, fig. 14; SWINHOE, Cat. Lep. Oxford.
- L. sonticum, Philippines, p. 142, absurdum, Salawatti, acceptum, Celebes, pl. v, fig. 5, kala, Ké I., fig. 8, p. 143, confusum, kondekum, Java, proprium, Manilla, fig. 12, p. 144, agrotum, New S. Wales, fig. 15, p. 145, galbanum, Philippines, p. 146, SWINHOE, Cat. Lep. Oxford, n. spp.
- Leptothrix, n. g., Zygænidæ, for L. tettigonioides, n. sp., Sumatra; Hev-LAERTS, p. 47, Ann. Ent. Belg. xxxvi.
- Leucodrepana, n. g., Drepanulidæ, for L. idæoides, nivea, obliquilinea, n. spp., Sikkim, Hampson, p. 333, Moths of India, i.
- Leucoma salicis, embryology; HENKING, Z. wiss. Zool. liv, p. 166.

- L. tavetensis, E. Africa, Holland, Ent. xxv (Supp.), p. 93; L. thyridophora, India, Намрзон, р. 488, Moths of India, i; L. lobipennis, Dorey, Swinhoe, p. 203, Cat. Lep. Oxford: n. spp.
- Liparis monacha, habits of larva, &c.; FENK (279). L. salicis, destruction by parasites; CARPENTIER, Bull. Soc. L. Nord France, x, p. 35.
- L. preussi, W. Africa, Mabille & Vuillot, Nov. Lep. p. 57, pl. ix, fig. 5 (referred by Kirby in Cat. Lep. Het. to Phægorista in Nyctemerida), n. sp.
- Lipuropsis, n. g., Notodontidæ, for L. postalbida, n. sp., Naga Hills; Hampson, Moths of India, i, p. 154.
- Lirimiris mechanica, Loja, Dognin, p. 169, Le Nat. 1892, n. sp.
- Lithosia (sub Tinea) entella, Cr., = (Gnophria ceramensis, Voll.); WEYMER, S. E. Z. liii, p. 122. L. caniola, var. complanoides; Fuchs, JB. nass. Ver. xlv, p. 88. L. tosola, sexmaculata, figured, pl. ix, figs. 2 & 3, Mabille & Vuillot, Nov. Lep.
- L. pusa, Loja, Dognin, p. 185, Le Nat. 1892, n. sp.
- Lobeza favilla, Loja, Dognin, p. 33, Le Nat. 1892, n. sp.
- Laspa dogninia, China, Sonthonnax, Échange, 1892, p. 20, n. sp.
- Lomera albifrons, Australia, SWINHOE, p. 225, Cat. Lep. Oxford, n. sp.
- Lonomia carnica, Brazil, SCHAUS, p. 318, P. Z. S. 1892, n. sp.
- Lophopteryz atrofusa, Sikkim, Hampson, p. 166, Moths of India, i, n. sp. Lophosoma, n. g. for Syntomis cuprea and quadricolor, Wlk.; SWINHOE, p. 58, Cat. Lep. Oxford.
- Lyces mæra, Brazil, Schaus, p. 285, P. Z. S. 1892 (figured, pl. ii, fig. 17, Schaus, American Lep.), n. sp.
- Lyclene vagilinea, figured, pl. iv, fig. 11; SWINHOE, Cat. Lep. Oxford.
- L. ruenca, Sula, SWINHOE, p. 101, pl. iv, fig. 15, Cat. Lep. Oxford, n. sp.
- Lymantria manicata, Cameroons, Aurivillius, p. 193, Ent. Tidskr. xiii,
- Macrocrambus, n. n. for Crambomorpha, Feld.; KIRBY, p. 337, Cat. Lep. Het. i.
- Macromphalia dedecora, Feist., = (ancilla, Phil.), rubrogrisea, Phil., = (rustica, Phil.); REED, p. 68, Act. Soc. Chili, i.
- Macronola, n. n. for Cyana, Wlk.; KIRBY, p. 299, Cat. Lep. Het. i.
- Macroplectra, n. g., Limacodidæ, type minutissima, Swinh.; Hampson, p. 376, Moths of India, i.
- Mardara albostriata, Manipur, Hampson, p. 455, Moths of India, i, n. sp.
- Marthula nora, pleione, Brazil, Schaus, p. 341, P. Z. S. 1892, n. spp.
- Marsypophora dissimilipennis, Loja, Dognin, Le Nat. 1892, p. 123, n. sp.
- Massaga delicia, Butl., Q described; Aurivillius, p. 186, Ent. Tidskr.
- Maxia, n. n. for Phryganopteryx, Saalm.; Kirby, p. 206, Cat. Lep. Het. i. Megaceramis, n. g., Notodontidæ, for M. lamprolepis, n. sp., Sikkim; Hampson, p. 167, Moths of India, i.
- Megalopyge radiata, acca, Brasil, Schaus, p. 322, P. Z. S. 1892, n. spp.

1892. [vol. xxix.]

- Mekla, n. g., Cossidæ, for Degia deficiena, Wlk.; SWINHOE, p. 283, Ca Lep. Oxford.
- Melanchroia braganza, palmeira, Brazil, SCHAUS, p. 288, P. Z. S. 189 (figured and referred to Leucopeumis, SCHAUS, p. 8, pl. ii, figs. 24 & 26, American Lep.), n. spp.
- Melanothriz lencotrigona, Burma, Hampson, Moths of India, i, p. 44, n. sp. Melia, n. g., near Ptilophora, for M. danbyi, n. sp., British Columbia Neumorgen, p. 225, Canad. Ent. xxiv.
- Mennis sceata, cytherea, Peru, una, Brazil, SCHAUS, p. 287, P. Z. S. 189. (figured, SCHAUS, American Lep. pl. ii), n. spp.
- Metagarista mænas, synonymical note; WEYMER, p. 125, S. E. Z. liii M. triphænoides and mænas, synonymy; AURIVILLIUS, Ent. Tidskr xiii, p. 184.
- M. aziyade, Sierra Leone, Vuillot, p. cxl, Bull. Soc. Ent. Fr. lxi, n. sp. Metanastria obliquifascia, Darjiling, p. 263, lucifuga. Singapore, p. 264 pl. vi, fig. 5, SWINHOE, Cat. Lep. Oxford, n. spp.
- Metarctia inconspicua, Trop. Africa, Holland, Ent. xxv (Supp.), p. 93 n. sp.
- Metaschalis, n. g., Notodontidæ, type disrupta, Moore; Hampson, Moth of India, i, p. 158.
- Metosamia, n. g., Saturniidæ, p. 276, for M. godmani, n. sp., Mexico, p. 277 DRUCE, Ann. N. H. (6) ix.
- Metriophyla lena, Peru, SCHAUS, p. 277, P. Z. S. 1892 (figured, pl. i fig. 24, SCHAUS, American Lep.), n. sp.
- Mila, n. g., Ayaristida, for Hypsa concinnula, Mab.; AURIVILLIUS, Ent Tidskr. xiii, p. 187.
- Milleria cyanivena, Burma, HAMPSON, p. 263, Moths of India, i, n. sp. Mimeusemia tara, Borneo, SWINHOE, p. 167, pl. vi, fig. 11, Cat. Lep Oxford, n. sp.
- Monotropia, n. n. for Licnoptera, Meyr.; Kirby, p. 307, Cat. Lep. Het. i. Mydrothauma, n. g., near Eupyra, p. 121, for M. ada, n. sp., Borneo, p. 121 pl. vi, fig. 1, Butler, P. Z. S. 1892, n. sp.
- Neærasa, n. n., to replace Neæra, H.-Sch.; STAUDINGER, p. 299, Romano Mem. vi.
- Naclia syntomida, Mardin, Staudinger, p. 248, Deutsche e. Z. Lep. is N.? erythrogaster, Freetown, Mabille, p. cxxxix, Bull. Soc. Ent. F lxi; N. aītcha, p. cxl, msila, p. cxli, Sierra Leone, Vuillot, t. c. n. spp.
- Nadata oregonensis, number of larval stages discussed; DYAR, Psyche, p. 337.
- Narosa velutina, figured; SWINHOE, pl. vii, fig. 8, Cat. Lep. Oxford.
- Nelo lippa, p. 287, caullama, p. 288, Pern, Schaus, P. Z. S. 1892 (figur Schaus, American Lep. pl. ii, fig. 18); N. solimara, Colomb Thierry-Mieg, p. 262, Le Nat. 1892: n. spp.
- Nemeta: characters and affinities; SNELLEN, p. 9, Tijdschr. Ent. xxx Neochera butleri, Assam, p. 84, zaria, Manilla, p. 85, SWINHOB, Cat. L. Oxford, n. spp.

- Neomiresa copac, Peru, Schaus, p. 324, P. Z.S. 1892, n. sp.
- Nerice bidentata, early stages; Soule, Psyche, vi, p. 276: metamorphoses; Marlatt, Tr. Kansas Ac. xi, p. 110, figs.
- N.? leechi, Amurland, Staudinger, p. 348, pl. v, fig. 2, Romanoff Mem. vi; N. upina, China, Alphéraky, p. 17, pl. i, fig. 7, t. c.; N. disjuncta, Loja, Dognin, p. 85, Le Nat. 1892: n. spp.
- Nesace, n. n. for part of Euchromia, Walk.; KIRBY, p. 112, Cat. Lep. Het. i.
- Neurosymploca agria, Transvaal, DISTANT, p. 236, pl. ii, fig. 9, Naturalist Transvaal, n. sp.
- Neumania, n. g., Liparidæ, for Teara guenei, Newm.; SWINHOE, p. 199, Cat. Lep. Oxford.
- Nishada fusciculosa figured, pl. iv, fig. 16; SWINHOE, Cat. Lep. Oxford.
- Nola fumosa, Butl. = (strigulosa, Stgr.); STAUDINGER, p. 256, Romanoff Mem. vi. N. minuscula, larva; Dyar, Psyche, vi, p. 248.
- N. candidalis, Amurland, Staudinger, p. 258, Romanoff Mem. vi; N. lactaria, Amurland, Graeser, p. 211, Berl. Ent. Zeit. xxxvii; N. quadrimaculata, p. 43, ranhassellii, p. 44, Java, Heylaerts, Ann. Ent. Belg. xxxvi: n. spp.
- Nossa, n. n. for Atossa, Moore; KIRBY, p. 56, Cat. Lep. Het. i.
- Notodonta streckfussi, Carinthia, Honrath, p. 445, Berl. Ent. Zeit. xxxvi; N. vittata, Anti-Taurus, Staudinger, p. 262, Deutsche e. Z. Lep. iv; N. graeseri, p. 351, pl. v, fig. 3, aliena, fig. 4, p. 352, oberthuri, fig. 5, p. 354, Amurland, Staudinger, Romanoff Mem. vi; N. P argenteo-maculata, Cameroons, Aurivillius, p. 196, Ent. Tidskr. xiii; N. rufa, Nága Hills, Hampson, Moths of India, i, p. 163; N. descherei, British Columbia, Neumoegen, p. 227, Canad. Ent. xxiv: n. spp.
- Nudaria murina, varr. described; STAUDINGER, p. 249, Deutsche e. Z. Lep. iv.
- Numeres siletti, neuration outlined; SWINHOE, p. 206, Cat. Lep. Oxford. Nyctemera amica, metamorphoses; WHITE, p. 68, Vict. Nat. ix.
- N. secundiana, Queensland, Lucas, p. 280, P. Linn. Soc. N.S.W. (2) vi, n. sp.
- Nyssia vetusta, pl. viii, fig. 15, cruda, fig. 12, figured and referred to Thosea, p. 237; SWINHOE, Cat. Lep. Oxford.
- Nyssia sulla, Brazil, Schaus, p. 324, P. Z. S. 1892, n. sp.
- Oberthueria, n. n. for Euphraor, Oberth.; KIRBY, p. 720, Cat. Lep. Het. i. Oberthuria, n. g. for Euphranor caca, Oberth.; STAUDINGER, p. 337,
- Romanoff Mem. vi.

  Ocinara albiceps, Wlk. (sub Naprepa), figured, pl. vii, fig. 7; Swinhoe,
- Cat. Lep. Oxford.

  \*\*Ocneria lapidicola\*, varr. described; STAUDINGER, p. 337, Deutsche e. Z. Lep. iv.
- O. destituta, Taurus, STAUDINGER, p. 256, Deutsche e. Z. Lep. iv, n. sp.
- Ocnsrogyia, n. g., Liparidæ, for O. amanda, n. sp., Taurus; STAUDINGER, p. 254, Deutsche e. Z. Lep. iv.

- Odonestis inobtrusa, Wlk., figured; SWINHOE, pl. vi, fig. 15, Cat. Lep. Oxford.
- O. directa, Japan, p. 261, pl. vi, fig. 4, SWINHOE, Cat. Lep. Oxford; O. castanea, p. 426, isocymu, p. 427, E. India, Hampson, Moths of India, i: n. spp.
- Edema matheis, Brazil, tlotzin, Mexico, Schaus, p. 332, P. Z. S. 1892, n. spp.
- Edenasia salicis, = (Dryocampa riversii, Behr.); DYAR, Psyche, vi, p. 222.
- E. terrena, maxtla, Mexico, guarana, p. 331, inca, Peru, p. 332, Schats, P. Z. S. 1892: n. spp.
- Œlonia, n. g. for Pseudopsyche, Edw.; KIRBY, p. 524, Cat. Lep. Het. i. Oiketicus abbotii, prolongation of larval life; Slosson, Ent. News, v. p. 49.
- Olceclosteria maya, Mexico, mutusca, Brazil, Schaus, p. 326, P. Z. S. 1892, n. spp.
- Olene basinigra, Sumatra, HEYLAERTS, p. 11, Ann. Ent. Belg. xxivi, n. sp.
- Opharus lugubris, morosus, Brazil, SCHAUS, p. 282, P. Z. S. 1892, n. spp. Oreta obliquilinea, p. 349, grissotineta, p. 350, India, HAMPSON, Moths of
  - India, i; O. singapura, Singapore, SWINHOE, p. 243, pl. vii, fig. 15, Cat. Lep. Oxford: n. spp.
- Orgyia cana, metamorphoses; DYAR, Psyche, vi, p. 203. O. leucostigma, metamorphoses; DYAR, t. c., p. 324.
- Ormiscodes eumedidoides, Amazons, aurora, Bogota, Vuillot, p. cxciii, Bull. Soc. Ent. Fr. lxi; O. opis, Brazil, Schaus, p. 318, P. Z. S. 1892: n. spp.
- Orna, n. n. for Euctenia, Feld.; KIRBY, p. 81, Cat. Lep. Het. i.
- Orthocraspeda, n. g., Limacodidæ, type trima, Moore; Hampson, p. 393, Moths of India, i.
- Orthorene arpi, janeira, Brazil, Schaus, p. 289, P. Z. S. 1892 (figured, pl. iii, Schaus, American Lep.), n. spp.
- Othria meridionalis, Brazil, Schaus, p. 272, P. Z. S. 1892, n. sp.
- Oxyplax, n. g., Limacodidæ, type ochracea, Moore; Hampson, p. 376, Moths of India, i.
- Pachyma, n. g., near Pachypasa, p. 115, for P. trapezina, n sp., Cameroons, p. 116; WEYMER, S. E. Z. liii.
- Pachypasa richelmanni, E. Africa, WEYMER, S. E. Z. liii, p. 113, n. sp.
- Paidina obtusa, var. sutschana, described; STAUDINGER, p. 260, Romanoff Mem. vi.
- Palpifer, n. g., Hepialidæ, type sexnotatus, Moore; Hampson, p. 316, Moths of India, i.
- Parasa argentilinea, p. 389, albipuncta, p. 390, dentata, p. 391, India, HAMPSON, Moths of India, i; P. minima, Mexico, SCHAUS, p. 322, P. Z. S. 1892: n. spp.
- Pelobrochis, n. g., near Nudaria, for Sceodora rava, Luc.; Lucas, p. 250, P. Linn. Soc. N.S.W. (2) vii.

- Pericopis cubana, H. Sch., = (Phaloë gaumeri, Druce); WEYMER, p. 122, S. E. Z. liii.
- P. montesuma, Mexico, SCHAUS, p. 283, P. Z. S. 1892, n. sp.
- Peronetis naringa, Philippines, pl. iii, fig. 16, p. 99, SWINHOE, Cat. Lep. Oxford, n. sp.
- Perophora corcovada, p. 326, acuta, p. 327, Brazil, Schaus, P. Z. S. 1892, n. spp.
- Phacusa cyanocera, Burma, Hampson, p. 241, Moths of India, i; P. variata, Dorey, Swinhoe, p. 58, Cat. Lep. Oxford: n. spp.
- Phagoptera schäfferi, Brazil, p. 280, proba, Mexico, notata, granifera, ursina, Brazil, p. 281, SCHAUS, P. Z. S. 1892 (notata, ursina, granifera, figured, SCHAUS, American Lep. pl. iii), n. spp.
- Phalacra excisa, Sikkim, Hampson, p. 345, Moths of India, i, n. sp.
- Phalænoides butleri, pl. v, fig. 6, macleayi, fig. 11, p. 157, Australia, Swinhoe, Cat. Lep. Oxford, n. spp.
- Phassus punctimargo, viridis, India, SWINHOE, p. 291, Cat. Lep. Oxford; P. punctimargo, p. 319, auratus, viridis, p. 321, India, HAMPSON, Moths of India, i; P. marcius, Mexico, DRUCE, p. 273, Ann. N. H. (6) ix; P. absyrtus, Brazil, SCHAUS, p. 330, P. Z. S. 1892: n. spp.
- Phauda fuscalis, Assam, SWINHOE, p. 60, Cat. Lep. Oxford, n. sp.
- I'heosia portlandia, early stages; DYAR, Psyche, vi, p. 351.
- P. excurvata, Sikkim, Hampson, Moths of India, i, p. 161; P. grisea, India, Swinhoe, p. 298, Cat. Lep. Oxford: n. spp.
- Philagria, n. n. for Æonistis, Hb.; KIRBY, p. 336, Cat. Lep. Het. i.
- Philania, n. n. for Pharmacis, Möschl.; KIRBY, p. 885, Cat. Lep. Het. i.
- Philudoria, n. n. for Odonestis, Steph.; KIRBY, p. 820, Cat. Lep. Het. i.
- Phlebohecta, n. g., Chalcosiidæ, type fuscescens, Moore; Hamison, p. 251, Moths of India, i.
- Phragmatæcia gummata, Java, SWINHOE, p. 285, pl. viii, fig. 14, Cat. Lep. Oxford, n. sp.
- Phragmatobia lymphasea, Loja, Dognin, p. 16, Le Nat. 1892, n. sp.
- Phya psamathe, Brazil, Schaus, p. 339, P. Z. S. 1892, n. sp.
- Piarosoma, n. g., Zygænidæ, for P. albicinctum, n. sp., Burma; Hampson, p. 243, Moths of India, i.
- Pielus albofasciatus, Swan River, SWINHOE, p. 289, Cat. Lep. Oxford, n. sp.
- Pinconia coa, Mexico, SCHAUS, p. 322, P. Z. S. 1892, n. sp.
- Pintia insularis, Singapore, SWINHOE, p. 76, Cat. Lep. Oxford, n. sp.
- Pirga, n. g. for P. mirabilis, Auriv.; Aurivillius, p. 193, Ent. Tidskr. xiii: figured; op. cit. xii, pl. ii, fig. 2.
- Pisara opalina, fig. 14, acontioides, fig. 19, lucidalis, fig. 12, figured, pl. iv; Swinhoe, Cat. Lep. Oxford.
- Platysygæna, n. g., for Soritia? moëlleri, Elw.; Swinhoe, p. 57, Cat. Lep. Oxford.
- Paccilocampa simplex, Australia, SWINHOE, p. 267, Cat. Lep. Oxford, n. sp. Polythysana albescens, Phil., = (cinerescens, Phil.); Reed, p. 68, Act. Soc. Chili, i.

Pompelon subcyanea, Singapore, p. 71, rotundata, Java, p. 72, SWINHO Cat. Lep. Oxford, n. spp.

Porina kershawi, Australia, Lucas, p. 282, P. Linn. Soc. N.S.W. (2)

Porthesia melania, Asia Minor, STAUDINGER, p. 255, Dentsche e. Z. Le iv; P. isabellina, Atjeh, HEYLAERTS, p. 9, Ann. Ent. Belg. xxxv. P. aurantiaca, Sikkim, HAMPSON, p. 485, Moths of India, i: n. sp. Preptos, n. g., Lasiocampida, p. 318, for P. oropus, n. sp., Mexico, p. 318 SCHAUS, P. Z. S. 1892.

Pseudantherwa, n. g., Saturniidæ, p. 109, for P. daphne, n. sp., Gold Comp. 110; WEYMER, S. E. Z. liii.

Pseudantiora, n. n. for Antiora, Möschl.; KIRBY, p. 566, Cat. Lep. Het. Pseudaphelia, n. n. for Aphelia, Westw.; KIRBY, p. 771, Cat. Lep. Het. Pseudasbolia, n. n. for Asbolia, Möschl.; KIRBY, p. 877, Cat. Lep. Het. Pseuderchia, n. n. for Erchia, Wlk.; KIRBY, p. 410, Cat. Lep. Het. i. Pseudhypsa, n. n. for Aganais, Boisd. pt.; KIRBY, p. 384, Cat. Lep. Het. Pseudojana, n. g., Eupterotidæ, type incandescens, Wlk.; Hampson, Motl of India, i, p. 48.

Pseudomya pellucida, p. 274, tijuca, Brasil, musca, p. 275, Mexico, Schau P. Z. S. 1892 (figured, Schaus, American Lep. pl. i), n. spp.

Pseudopsyche dembowskii, notes on; Staudinger, p. 294, Romanoff Mer

Psilura rubripunctata, E. Africa, WEYMER, p. 108, S. E. Z. liii, n. sp. Psychida: wing-neuration discussed; SPULER, Z. wiss. Zool. liii, p. 639
Psyche (Eurycyttarus) nigra, p. 299, decemvena, p. 300, India, Hampso Moths of India, i, n. spp.

Pterostoma palpina, n. var. gigantina; STAUDINGER, p. 363, Romano Mem. vi.

Pydna eupatagia, p. 141, nigropuncta, nigrofasciata, p. 142, India, Han son, Moths of India, i; P. calyptis, Dorey, Swinhor, p. 297, Ca Lep. Oxford: n. spp.

Pygara curtula, n. var. canescens; Graeser, p. 303, Berl. Ent. Ze xxxvii. P. troglodyta, figured, pl. v, fig. 7, Romanoff Mem. vi.

P. sieversi, Amurland, STAUDINGER, p. 370, pl. v., figg. 6 a, b, Romand Mem. vi; P. moderata, Central Asia, Graeser, p. 303, Berl. Es Zeit. xxxvii: n. spp.

Pygetera, n. n. for Cutaphractes, Feld.; Kirby, p. 446, Cat. Lep. Het. i. Pyrosis idiota, & Q figured, pl. iv, figg. 6, a, b, Romanoff Mem. vi. Rachia striata, Sikkim, Hampson, p. 132, Moths of India, i, n. sp. Rumesa fuscipennis, Burma, Hampson, Moths of India, i, p. 143, n. sp. Rezia, n. n. for Lagaria, Walk.; Kirby, p. 149, Cat. Lep. Het. i.

Rhodareas, n. g., for part of Areas; KIRBY, p. 254, Cat. Lep. Het. i. Rhodoneura obliqualis, p. 355, multipunctata, p. 356, hypargyra, p. 3 glaphyralis, p. 359, bipuncta, p. 360, cuprealis, p. 361, oligostic subcostalis, ferrofusa, uniformis, p. 362, bracteata, rufareta, p. 3 hypoxantha, tristis, munda, p. 364, India, Hampson, Moths of India (cf. also Pyralidæ, &c., p. 277), n. spp.

Rhuda endymion, Brazil, SCHAUS, p. 340, P. Z. S. 1892, n. sp.

Rifargia cloelia, Brazil, p. 338, gelduba, Mexico, p. 339, Schaus, P. Z. S. 1892, n. spp.

Rosema emavata, luna, Brazil, luppa, p. 337, incita, Peru, languida, unda, Brazil, p. 338, Schaus, P. Z. S. 1892, n. spp.

Sampatissa pretoriæ, Transvaal, DISTANT, p. 408, Ann. N. H. (6) x, n. sp. Supinella, n. n. for Eutheca, Grote; KIRBY, p. 524, Cat. Lep. Het. i.

Sarmalia alba, Philippines, Swinhoe, p. 275, Cat. Lep. Oxford, n. sp.

Sarothroceras rhomboidea, Cameroons, WEYMER, p. 104, S. E. Z. liii, n. sp. Sarrothripa reveyana, larva; Dyar, Psyche, vi, p. 259.

Saturniida, notes on various; KIRBY, Ent. xxv, pp. 267-269.

Saturnia pavonia, nature, and use of its spines; Hrim, p. xlviii, Bull.
Soc. Ent. Fr. lxi. S. arnobia, peculiar chrysalis; Holland, Psyche,
vi, p. 213, pl. v. S. carpini, variation in colour of cocoons discussed;
BATESON (40, 41).

Scaptius juno, Brazil, SCHAUS, p. 279, P. Z. S. 1892 (figured, SCHAUS, American Lep. pl. ii, fig. 3), n. sp.

Seea solaris, Peru, Schaus, p. 285, P. Z. S. 1892 (figured, Schaus, pl. ii, fig. 14, American Lep.), n. sp.

Scepsis nigricollum, Loja, Dognin, p. 123, Le Nat. 1892, n. sp.

Schizuru con\*pecta, egg and young larva; Dyar, Psyche, vi, p. 325.

Sebastea, n. n. for Grotea, Moore; KIRBY, p. 383, Cat. Lep. Het. i.

Selca cuneifera, Wlk. (sub Melia), figured, pl. iv, fig. 22; SWINHOE, Cat. Lep. Oxford.

8. obscura, Queensland, Swinhoe, p. 133, Cat. Lep. Oxford, n. sp.

Semyra struminea, Mexico, SCHAUS, p. 324, P. Z. S. 1892, n. sp.

Sermyla morta, Brazil, Schaus, p. 282, P. Z. S. 1892, n. sp.

Secapa divisa, figured, pl. iv, fig. 21; SWINHOB, Cat. Lep. Oxford.

Siglophora, n. g., near Chandica, p. 123, for S. bella, n. sp., Borneo, p. 124, pl. vi, fig. 2; BUTLER, P. Z. S. 1892.

Sitina albicans, W. Australia, SWINHOE, p. 268, pl. vi, fig. 10, Cat. Lep. Oxford, n. sp.

Somabrackys, n. n. for Brackysoma, Aust.; KIRBY, p. 930, Cat. Lep. Het. i.

Somatrichia, n. n. for Trichosoma, Ramb.; KIRBY, p. 274, Cat. Lep. Het i.

Soritia bipartita, Wlk., figured, pl. ii, fig. 9; SWINHOE, Cat. Lep. Oxford. S. viridivena, nigribasalis, HAMPSON, p. 253, Moths of India, i, n. spp.

Sozusa stevensii, E. Africa, Holland, Ent. xxv (Supp.), p. 91, n. sp.

Spatalia albonotata, Wlk. (sub Allata), pl. viii, fig. 2, punctifera, Wlk. (sub Caschara), fig. 4, lativitta, Wlk. (sub Xylina), fig. 1, figured; SWINHOE, Cat. Lep. Oxford.

S. albifasciata, Nilgiris, Hampson, Moths of India, i, p. 170, n. sp.

Spilarctia abbotti, E. Africa, Holland, Ent. xxv (Supp.), p. 92, n. sp.

Spilosoma urtica, n. var. quadripunctata; Cosmovici, p. 280, Le Nat. 1892; S. urtica, var. paucipuncta; Fuchs, p. 89, JB. nass. Ver. xlv.

Stauropus fagi, habits noticed; Holland, Ent. M. M. xxviii, p. 206.

- pallidifascia, p. 151, albivertex, orbifer, p. 152, India, Hampson, Moths of India, i, n. spp.
- Stenophylloides, n. g., Lasiocampida, type sikkima, Moore; Hampson, p. 429, Moths of India, i.
- Stibolepis atomaria, E. Africa, Holland, Ent. xxv (Supp.), p. 93, n. sp.
- Stichobasis, n. n. for Diabasis, Heyl.; KIRBY, p. 519, Cat. Lep. Het. i.
  Striggides fuscus, p. 280. decoratus, p. 281. Australia, SWINHOR, Cat. Lep.
- Strigoides fuscus, p. 280, decoratus, p. 281, Australia, SWINHOE, Cat. Lep. Oxford, n. spp.
- Susica congrua, Wlk., referred to Natada, p. 233, figured, pl. viii, fig. 6; SWINHOE, Cat. Lep. Oxford.
- Sychesia hartmanni, janeira, Brazil, Schaus, p. 280, P. Z. S. 1892 (figured, Schaus, American Lep. pl. iii), n. spp.
- Symmerista dentata, p. 335, tethys, proces, myconos, p. 336, Brazil, Schaus, P. Z. S. 1892, n. spp.
- Syntomis gracilis, n. var. lugens; Röber, p. 325, Tijdschr. Ent. xxiv. S. caspia and annetta are varr. of maracandina; Alpheraky, p. 4, Romanoff Mem. vi. S. longipennis, Wlk., figured, pl. ii, fig. 13, S. annosa, Wlk., figured and referred to Hydrusa, p. 51, pl. ii, fig. 17; Swinhoe, Cat. Lep. Oxford.
- S. sintenisi, Kurdistan, STANDFUSS, p. 664, pl. xv, fig. 3, Romanoff Mem. vi; S. rubicunda, pterophorina, Natal, Mabille, p. cxxxix, Bull. Soc. Ent. Fr. lxi; S. unifuscia, p. 212, flavifrons, quadrifascia, p. 215, chlorocera, luteifascia, p. 218, eressoides, p. 220, nigra, serrata. p. 221, ochreipuncta, p. 222, erythrosoma, p. 223, India, Hampson, Moths of India, i; S. cymatilis, Philippines, p. 40, insueta, E. India, pl. ii, fig. 16, p. 41, aurofasciata, New Guinea, pl. ii, fig. 19, p. 43, frustulenta, Lizard I., p. 44, deflocca, Philippines, p. 45, ypleta, Waigiou, formicina, New Guinea, p. 47, exapta, Flores, fata, Borneo, p. 48, Swinhoe, Cat. Lep. Oxford; S. trifenestra, Letti, Röber, p. 324, Tijdschr. Ent. xxxiv (fig. in xxxv): n. spp.
- Syntomoides, n. g., p. 209, type imaon, Cram., and including S. microchilu, n. sp., Tenasserim, p. 212; Hampson, Moths of India, i.
- Syntrichura doeri, Brazil, Schaus, p. 276, P. Z. S. 1892 (figured, pl. i, fig. 16, American Lep.), n. sp.
- Syssphinz basirei, Brazil, Schaus, p. 289, P. Z. S. 1892 (figured, Schaus, American Lep. pl. iii, fig. 5), n. sp.
- Tamphana, n. g., Bombyciidæ, for T. marmorea, n. sp., Brazil; Schats, p. 325, P. Z. S. 1892.
- Tantura, n. n. for Penthetria, Edw.; KIRBY, p. 86, Cat. Lep. Het. i.
- Tarchon minois, Brazil, SCHAUS, p. 325, P. Z. S. 1892, n. sp.
- Tasema longipennis, p. 236, nigra, p. 237, Burma, Hampson, Moths of India, i, n. spp.
- Teara erebodes, Centr. Australia, Lower, p. 14, Tr. R. Soc. S. Austr. xvi; T. togata, fimbriata, Queensland, Lucas, p. 285, P. Linn. Soc. N.S.W. (2) vi: n. spp.
- Telea aurelia, Mexico, Druce, p. 278, Ann. N. H. (6) ix, n. sp.
  Teracotona clara, E. Africa, Holland, Ent. xxv (Supp.), p. 92, n. sp.

- Termessa xanthomelas, S. Australia, Lower, p. 6, Tr. R. Soc. S. Austr. xv, n. sp.
- Tetraphleps, n. g., type brevilinea, Wlk.; HAMPSON, p. 383, Moths of India, i.
- Teucer, n. n. Telioneura, Feld.; KIRBY, p. 165, Cat. Lep. Het. i.
- Teulisna plagiata, pl. iii, fig. 11, chiloides, pl. iv, fig. 10, figured; Swin-Hoe, Cat. Lep. Oxford.
- Theages vestalis, Peru, Schaus, p. 278, P. Z. S. 1892 (figured, Schaus, American Lep. pl. i, fig. 23); T. nubilosa, mamona, Loja, Dognin, Le Nat. 1892, p. 144: n. spp.
- Thyrassia inconcinna, Lizard I., SWINHOE, p. 55, Cat. Lep. Oxford, n. sp. Ticera, n. g, Lasiocampidæ, for T. castanea, n. sp., Philippines; SWINHOE, p. 269, Cat. Lep. Oxford.
- Ticilia argentilinea, figured, pl. vii, fig. 13, Cat. Lep. Oxford.
- Titya fumida, p. 321, argentata, p. 322, Brazil, Schaus, P. Z. S. 1892, n. spp.
- Trabala cicur, brumalis, Brazil, drucei, cebrenis, Mexico, Schaus, p. 323, P. Z. S. 1892, n. spp.
- Trabaloides, n. n. for Protogenes, Saalm.; KIRBY, p. 818, Cat. Lep. Het.i. Trichata, n. g., for Syntomis teneiformis, Wlk.; SWINHOE, p. 52, Cat. Lep. Oxford.
- Trichomelia celenna, Brazil, Schaus, p. 283, P. Z. S. 1892 (figured, Schaus, American Lep. pl. ii, fig. 5), n. sp.
- Trichura cyanea, Brazil, Schaus, p. 276, P. Z. S. 1892, n. sp.
- Triplophleps, n. g., type inferma, Swinh.; HAMPSON, p. 393, Moths of India, i.
- Tristania, n. n. for Zonosoma, Butl.; KIRBY, p. 423, Cat. Lep. Het. i.
- Tropæa mandschurica, Ussuri, Staudinger, p. 331, Romanoff Mem. vi, n. sp.
- Trypanophora flavalis, Burma; HAMPSON, p. 250, Moths of India, i, n. sp. Urocoma baliolalis, N. S. Wales, SWINHOE, p. 215, pl. vi, fig. 7, Cat. Lep.
- Urocoma baliolalis, N. S. Wales, SWINHOE, p. 215, pl. vi, fig. 7, Cat. Lep Oxford, n. sp.
- Uropyia meticulodina, larva figured, pl. iv, fig. 8, noticed, p. 314, STAU-DINGER, Romanoff Mem. vi.
- Vipsania frigida, Mexico, Schaus, p. 323, P. Z. S. 1892, n. sp.
- Virbia varians, p. 284, parva, p. 285, Peru, Schaus, P. Z. S. 1892 (figured, Schaus, American Lep. pl. ii), n. spp.
- Kanthospilopteryx abacata, p. 181, lomata, p. 182, W. Africa, Karsch, Ent. Nachr. zviii; X. athene, p. 101, minerva, p. 103, E. Africa, Weymer, S. E. Z. liii: n. spp.
- Zatrephes biseriata, Plötz, systematic position queried; Aurivillius, p. 196, Ent. Tidskr. xiii.
- Zebeeba, n. n. for Nycteola, H.-S.; KIRBY, p. 279, Cat. Lep. Het. i.
- Zeuzera pyrina, as N. American; SMITH, p. 136, Canad. Ent. xxiv: habits in N. America; Pike, Ins. Life, iv, p. 317.
- Z. ? regia, Taurus, Staudinger, p. 253, pl. iii, fig. 6, Deutsche e. Z. Lep.; Z. congerens, Australia, Swinhoe, p. 282, pl. viii, fig. 5, Cat. Lep. Oxford; Z. ramosa, Brazil, Schaus, p. 329, P. Z. S. 1892: n. spp.

Zonosoma: of. Tristania, p. 249, and Geometridæ, p. 273.

Zutulba, n. n. for Anteris, Wallengr.; KIRBY, p. 80, Cat. Lep. Het. i.

Zygana niphona, Butl., = (christophi, Stgr.); STAUDINGER, p. 251. Romanoff Mem, vi. Z. cynarce, n. var. turatii; Standfuss, p. 368, MT. Schw. ent. Ges. viii. Z. rhadamanthus, n. var. roederi; Stat-DINGER, p. 247, Deutsche e. Z. Lep. iv.

Z. sardoa, Sardinia, MABILLE, p. cl, Bull. Soc. Ent. Fr. lxi; Z. octo, Transvaal, holozanthina, Kitui, MABILLE, p. cxxxviii, Bull. Soc. Ent. Fr. lxi; Z. erythrosoma, E. India, HAMPSON, p. 231, Moths of

India, i : n. spp.

Zygenopsis, n. g. for Z. fuscimarginalis, Dorey, p. 61, flavibasis, Ké I., p. 62, n. spp.; Swinhor, Cat. Lep. Oxford.

## CYMATOPHORIDAL.

[Cf. Hampson (384), Staudinger (779).]

Gaurena aurofasciata, p. 181, fuscescens, lichenea, p. 182, India, HAMPSON, Moths of India, i, n. spp.

Habrosyne argenteipunctata, Sikkim, Hampson, Moths of Iudia, i, p. 179,

Keralu: generic characters; STAUDINGER, p. 381, Romanoff Mem. vi. Nemacerota, n. g., type cinerea, Warren; HAMPSON, p. 185, Moths of India, i.

Polyploca anguligera, p. 183, albibasis, p. 184, India, HAMPSON, Moths of India, i, n. spp.

Tozoides, n. g., p. 185, for T. longipennis, n. sp., Sikkim, p. 186, HAMPSON, Moths of India, i.

### NOCTUIDÆ.

[Cf. Alphéraky (4, 779), Aurivillius (23, 26), Butler (141, 142, 144, 145), DISTANT (216), DOGNIN (220), GRAESER (356, 357), HOLLAND (408), LUCAS (568, 569), MALLY (577), OLLIFF (654), POUJADE (702), RÜHL (790), SKINNER & MENGEL (850), SMITH (861, 862, 863, 867), Standfuss (779), Staudinger (779, 886, 890, 891), Tutt (955).]

Criticism of Butler's synonymy of N. American Noctuida; Smith, Ent. xxv, p. 2.

Note on the points available for generic characters; SMITH, Ent. xxv, pp. 101-105.

Number of subdivisions; GROTE, Canad. Ent. xxiv, p. 17.

TUTT has published the second vol. of his work on the variation of British Noctuida.

Acantholipes (sub Hydrelia) circumdata, Wlk., = (Docela vetustalis, Wlk.); Butler, Ent. xxv, p. 141.

Acherdoa ferraria, Wik., = (Varina ornata, Neum.); SMITH, p. 135. Canad, Ent. xxiv.

- Acontia delecta, full-grown larva; Beutenmüller, p. 68, Bull. Am. Mus. iv. A. gratiosa, Wallengr., = (luteola, Saalm.); Snellen, p. 77, Tijdschr. Ent. xxxv.
- Ædia, Hb., = (Anophia, Gn.); BUTLER, p. 213, Ent. xxv.
- Acronycta: structural details of larvæ and pupæ of British species, continued by Chapman, in Ent. Rec. iii, pp. 1, &c., pls. iv & vi. A. dactylina, full-grown larva; Beutenmüller, p. 77, Bull. Am. Mus. iv. A. alni, use of hairs; Fitz-Gibbon, Ent. xxv, p. 39.
- terrigena, Amurland, GRAESER, p. 214, Berl. Ent. Zeit. xxxvii, n. sp.
- Agrophila sulphuralis, = (Erastria pardalina, Wlk.); Butler, p. 91, Ent. xxv.
- Agrotis sepulcralis, pl. ii, fig. 2, degeniata, var. aucta, fig. 3, pæcila, fig. 4, characteristica, pl. i, fig. 9, figured; Romanoff Mem. vi. A. festiva, n. var. thulei, glareosa, n. var. edda, depuncta, p. 266, renigera, p. 268, varr. described; Staudinger, Deutsche e. Z. Lep. iv. A. pachnoboides, pl. vi, fig. 3, militaris, fig. 5, nigricostatu, fig. 6, præcurrens, fig. 7, figured; Staudinger, Romanoff Mem. vi.
- A. succica, Sweden, Aurivillius, Nordens Fjärilar, pp. 113 & 121; A. amæna, Amasia, p. 267, hadjina, Taurus, p. 270, mollis, Margelan, p. 271, Staudinger, Deutsche e. Z. Lep. iv; A. honesta, p. 354, pl. iii, fig. 3, albivenata, fig. 4, p. 359, præcipua, fig. 5, p. 360, trifurcula, fig. 6, p. 362, Siberia, Staudinger, op. cit. v; A. vidua, p. 409, pl. vii, fig. 3, umbra, p. 411, Amurland, Staudinger, Romanoff Mem. vi; A. nigrita, p. 217, cicatricosa, p. 218, coturnicola, p. 219, lapidosa, p. 220, E. Siberia, Graeser, Berl. Ent. Zeit. xxxvii; A. dulcis, Centr. Asia, Alpheraky, Hor. Ent. Ross. xxvi, p. 444: n. spp.
- Ala pretiosa, Central Asia, Alphéraky, p. 454, Hor. Ent. Ross. xxvi, n. sp.
- Alamis subcinerea, Snell., = (albangula, Saalm.); SNELLEN, p. 79, Tijdschr. Ent. xxxv.
- Allotria amica, Hb., = (linsella, Gr.); BUTLER, p. 214, Ent. xxv.
- Ammoconia vetula, var. mediorhenana; Fuchs, p. 91, JB, nass. Ver. xlv.
- Amyna selenampha, Gn., = (paradoxa, Saalm.); SNELLEN, p. 75, Tijdschr. Ent. xxxv.
- Anarta besta, Greenland, SKINNER, P. Ac. Philad. 1892, p. 158, n. sp.
- Anartomorpha, n. g., near Anarta, for A. potanini, n. sp. (Chiua?), pl. ii, fig. 8; Аlphéraky, p. 39, Romanoff Mem. vi.
- Anchocelis litura, food-plant; JEFFREY, Ent. M. M. xxviii, p. 47.
- Anophia leucomelas, L., = (discistriga and trispilosa, Saalm.); SNELLEN, p. 80, Tijdschr. Ent. xxxv.
- Antha, n. g., near Chloantha, for A. pretiosa, n. sp., Amurland, pl. vii, fig. 6; STAUDINGER, p. 448, Romanoff Mem. vi.
- Anthophila: note on use of the name; SNELLEN, p. 78, Tijdschr. Ent. xxxv.
- Apamea dumerilii, varr. described; STAUDINGER, p. 284, Deutsche e. Z. Lep. iv.

- Aplectoides caliginea, figured, pl. vi, fig. 9, Romanoff Mem. vi.
- Aporophyla haasi, Spain, STAUDINGER, p. 280, Deutsche e. Z. Lep. iv, n. sp.
- Argiva and Patula, peculiarity of neuration and glandular organs; Hampson, P. Z. S. 1892, p. 193.
- Arzama: the name is more recent than Bellura, Wlk.; Smith, p. 135, Canad. Ent. xxiv.
- Audea: Phoberia catocala, Feld., referred to, p. 214, A. bipunctata, Wlk, = (P. fatua, Feld.), p. 213; Butler, Ent. xxv.
- Aventina, n. g., p. 605 (? same as Aventiola), for A. costimacula, pl. xiv, fig. 4, p. 604, solituria, fig. 5, p. 605, n. spp., Amurland; STAUDINGEE, Romanoff Mem. vi.
- Aventiola, n. g., near Aventia, for A. maculifera, n. sp., Amurland, pl. xiv, fig. 3; STAUDINGER, p. 602, Romanoff Mem. vi.
- Bagada (sub Acontia) plumbata, Butl., = (Marimatha freda, Sw.). B. spicea, Gn. = (Acontia firina, Sw.); BUTLER, p. 90, Ent. xxv.
- Bellura gortynoides, Wlk., = (Arzama densa); SMITH, p. 135, Canad. Ent. xxiv.
- Berresa, Wik., = (Pteratholix); Butler, p. 11, Ent. xxv.
- Boletobia cognata, Amurland, STAUDINGER, p. 606, Romanoff Mem. vi. n. sp.
- Brada curvilinea, Amurland; STAUDINGER, p. 553, pl. ix, fig. 12, Romanoff Mem. v.
- Bryophila tabora, Aintab, p. 264, paulina, Jerusalem, p. 265, STAUDINGER, Deutsche e. Z. Lep. iv; B. roderi, Kurdistan, STANDFUSS, p. 665, pl. xv, fig. 4, Romanoff Mem. vi; B. glaucula, p. 394, pl. v, fig. 8, albonotata, fig. 9, p. 396, B. P (Bryomoia) melachlora, fig. 10, p. 397. Amurland, STAUDINGER, t. c.; B. confucii, p. 19, pl. i, fig. 8, chloromixta, p. 21, pl. ii, fig. 1, China, Alphéraky, t. c.: n. spp.
- Bryophilina, n. g., near Bryophila, for B. blandula, n. sp., Amurland, pl. v, fig. 11; STAUDINGER, p. 398, Romanoff Mem. vi.
- Culliodes pretiosissima, E. Africa, Holland, Ent. xxv (Supp.), p. 94, n. sp. Culocampa brucci, Colorado, Smith, p. 252, Ent. News, iii, n. sp.
- Culymnia, Hb., = (Elyptron, Saalm.); SNELLEN, p. 76, Tijdschr. Ent. xxxv. C. picta, pl. x, fig. 2, moderata, fig. 1, figured, Romanoff Mem. vi.
- Capnodes trifusciata, Moore, = (alboguttata, Heyd.); Snellen, Tijdschr. Ent. xxxv, p. 82.
- Caradrina vicina, varr. described; STAUDINGER, p. 294, Deutsche e. Z. Lep. iv. C. funesta, fig. 10, maurella, fig. 9, illustrata, fig. 11, figured, pl. viii, Romanoff Mem. vi.
- C. (Amphidrina) agrotina, Amasia, p. 293, pl. iii, fig. 11, STAUDINGER, Deutsche e. Z. Lep. iv, n. sp.
- Catocala: synonymic notes; Butler, Ent. xxv, pp. 283-285: synonymy of some N. American; Hulst, p. 74, Canad. Ent. xxiv, and French, p. 19, t. c. C. lineella and alabama, synonymical note; Grote, p. 155, Canad. Ent. xxiv. C. deuteronympha, pl. xii, fig. 2, agitatrix, fig. 3,

- serenides, pl. xi, fig. 6, dærriesi, fig. 8, greyi, fig. 7, streckeri, fig. 9, figured, Romanoff Mem. vi. C. elocata, n. var. locata; STAUDINGER, p. 327, Deutsche e. Z. Lep. iv. C. illecta, preparatory stages; FRENCH, p. 307, Canad. Ent. xxiv. C. stretchii, early stages; FRENCH, p. 229, t. c. C. electa, sponsa, colour variation of larvæ; Poulton, Tr. E. Soc. 1892, p. 300. C. nupta and fraxini, variation in colour of larvæ; Gould, pp. 219-234, Tr. E. Soc. 1892.
- C. remissa, p. 328, pl. iv, fig. 10, contemnenda, p. 329, fig. 11, æstimabilis, p. 330, fig. 12, Turkestan, STAUDINGER, Deutsche e. Z. Lep. iv; C. hetæra, p. 592, pl. xii, fig. 4, eminens, fig. 5, p. 594, Amurland, STAUDINGER, Romanoff Mem. vi: n. spp.
- Chamyris cerintha, full-grown larva; Beutenmüller, p. 78, Bull. Am. Mus. iv.
- Chara, n. g., near Thalpochares, for C. albosignata, pl. xiv, fig. 2, p. 559, bimaculata, p. 561, n. spp., Amurland; STAUDINGER, Romanoff Mem. vi.
- Chariclea erubescens, Amurland, GRAESER, p. 229, Berl. Ent. Zeit. xxxvii; C. hedemanni, Amurland, STAUDINGER, p. 552, Romanoff Mem. vi; C. sanguinata, Brisbane, Lucas, p. 254, P. Linn. Soc. N.S.W. (2) vii: n. spp.
- Charidea v-brunneum, Gn., = (Appana rosacea, Saalm.); SNELLEN, p. 76, Tijdschr. Ent. xxxv.
- Chobata discalis, Wlk., = (Erastria concludens, Wlk.); BUTLER, p. 65, Ent. xxv.
- Cirradia xerampelina, varr. described; Staudinger, p. 299, Deutsche e. Z. Lep. 1v.
- Clavipalpula, n. g. for Taniocampa auraria, Oberth.; STAUDINGER, p. 499, Romanoff Mem. vi.
- Cleophana pauli, Jerusalem, STAUDINGER, p. 306, Deutsche e. Z. Lep. iv, n. sp.
- Cocytodes modesta and polygrapha, synonymy; Butler, Ent. xxv, p. 212. Copipanolis borealis, fasciata, N. America, Smith, p. 62, P. U. S. Nat. Mus. xv, n. spp.
- Cosmia trapezoides, figured, pl. ix, fig. 7, Romanoff Mem. vi.
- C. regina, Amasia, STAUDINGER, p. 297, Deutsche e. Z. Lep. iv, n. sp.
- Cucullia maculosa, figured, pl. x, fig. 4, Romanoff Mem. vi. C. cyrtana, figured; MABILLE & VUILLOT, pl. xi, fig. 3, Nov. Lep. Monograph of the N. American species; SMITH, P. U. S. Nat. Mus. xv, pp. 33-52, pl. ii.
- C. similaris, obscurior, p. 41, dorsalis, p. 47, bistriga, p. 48, cinderella, p. 49, N. America, Smith, P. U. S. Nat. Mus. xv; C. umbristriga, Central Asia, Alphéraky, p. 452, Hor. Ent. Ross. xxvi; C. sublutea, p. 312, hemidiaphana, p. 313, Central Asia, Graeser, Berl. Ent. Zeit. xxxvii; C. distinguenda, Siberia, Staudinger, p. 368, pl. iii, fig. 7, Deutsche e. Z. Lep. v: n. spp.
- Curvatula, n. g., near Aventia, for C. pallicostata, n. sp., Amurland, pl. xii, fig. 6; STAUDINGER, p. 601, Romanoff Mem. vi.

- Dasygaster hollandia, Gn., (leucanioides, Gn., and Grophiphora facili Wlk.), D. epundoides, Gn., (punctivena, Wlk.); BUTLER, p. 1 Ent. xxv.
- Dasypolia templi, n. var. caftischli; ROHL, p. 170, Soc. Ent. vi.
- D. ferdinandi, Stilfser Joch, RÜHL, Soc. Ent. vi, p. 169; D. paged gerbillus, Central Asia, Alpheraky, p. 452, Hor. Ent. Ross. xxv D. ? fani, Amurland, Staudinger, p. 522, pl. ix, fig. 10, Roman Mem. vi : n. spp.
- Deva, Polychrysia, and Plusia, note on the confusion as to these gener their members mentioned; BUTLER, pp. 407 & 408, Ann. N. H. ( ix.
- Dianthocia filigrama, n. var. melanochroa; Staudinger, p. 273, Deutse e. Z. Lep. iv.
- D. musculina, Malatia, STAUDINGER, p. 272, Deutsche e. Z. Lep iv; deliciosa, lurida, Central Asia, ALPHERAKY, p. 446, Hor. Ent. Roxxvi: n. spp.
- Dicopis: revision of the N. American; SMITH, P. U. S. Nat. Mus.: pp. 53 & 68. D. viridescens, Wlk., = (muralis, Grt.): Psaphic resumens, Wlk., is the same; SMITH, p. 135, Canad. Ent. xxiv.
- Drasteria erichto, note on ; SLINGERLAND, Ent. News, iii, p. 63.
- Dryobota contermina, Central Asia, Graeser, p. 308, Berl. Ent. Ze xxxvii; D.? amurensis, Ussuri, Staudinger, p. 432, pl. vii, fig Romanoff Mem. vi: n. spp.
- Dyschorista (sub Cerastis) prodeuns, Wlk., = (Taniocampa pueri Grote); Butler, p. 10, Ent. xxv.
- D. contecta, Central Asia, Graeser, p. 310, Berl. Ent. Zeit. xxxvii, n. Eccopteroma, n. g., for Calymnia fulcata, Graeser; Staudinger, p. 5
  Romanoff Mem. vi.
- Eicomorpha? argillacea, Amdo, Alpheraky, p. 445, Hor. Ent. B. xxvi, n. sp.
- Eliochræa chrysochlora, Wlk., = (Amphipyra laportei, Feld.); BUTL Ent. xxv, pp. 285 & 286.
- Epimecia subtilis, pl. xi, fig. 4, quadrivirgula, fig. 5, figured; Mabilli Vuillot, Nov. Lep.
- Episema lederi, varr. described; STAUDINGER, pp. 274-276, Deutsch Z. Lep. iv.
- Epunda muscosa, Amasia, STAUDINGER, p. 281, Deutsche e. Z. Lep. n. sp.
- Erastria wiskotti, pl. xi, fig. 4, numisma, fig. 3, distinguenda, fig. 4, figu: Romanoff Mem. vi.
- E. staudingeri, Kurdistan, Standfuss, p. 667, pl. xv, fig. 7, Roma: Mem. vi, n. sp.
- Erebus odorata, = (marquesii, Phil.); REED, p. 68, Act. Soc. Chili, i. Eriopus venus, figured, pl. viii, fig. 6, Romanoff Mem. vi.
- Erygia apicalis, numerous synonyms of; BUTLER, Ent. xxv, p. 213.
- Erythrophaia, n. g. for Chariclea? sudoxia, n. sp., Kashgar, pl. iv, fig. Staudinger, p. 316, Deutsche e. Z. Lep. iv.

- Eublemma (sub Micra) hemirhoda, Wlk., = (Anthophila roseifascia, Wlk.);
  BUTLER, Ent. xxv, p. 189.
- Eumestla (sub Thalpochares) patula, Mor., = (Tarache patruelis, Gr., and Anthophila flammicineta, Wlk.); BUTLER, p. 189, Ent. xxv.
- Euplexia græseri, Amurland, Staudinger, p. 462, Romanoff Mem. vi, n. sp.
- Eurhipia bowkeri, Feld., = (Eutelia exquisita, Saalm.); Snellen, p. 78, Tijdschr. Ent. xxxv.
- E. adoratriz, STAUDINGER, p. 308, Deutsche e. Z. Lep. iv, n. sp.
- Euromoia. n. g. of doubtful position, for E. mixta, n. sp., Amurland, pl. xii, fig. 10; STAUDINGER, p. 632, Bomanoff Mem. vi.
- Eutolype bombyciformis, Ohio, SMITH, p. 59, P. U. S. Nat. Mus. xv, n. sp. Fodina ostorius, figured; SCOTT, Austral. Lep. ii, pl. xviii.
- F. gloriosa, Queensland, Lucas, p. 304, P. Linn. Soc. N.S.W. (2) vi, n. sp. Fruva (sub Xanthodes) parvulu, Wlk., = (georgica, Gr.); Butler, p. 91, Ent. xxv.
- Gonitis editrix, Gn., = (fractifera and cosmioides, Wlk.), G. sabulifera, Gn., = (involuta, var., G., and busalis, Wlk., Tiridata colligata, Wlk., and G. propinqua, Butl.); Butler, p. 11, Ent. xxv. G. fulvida, Gn., complex synonymy; Butler, t. c., p. 12.
- Grammesia bifasciata, figured, pl. viii, fig. 8, Romanoff Mem. vi.
- Grammodes excellens, p. 257, pulcherrima, p. 258, divaricata, p. 259, pullens, p. 260, Brisbane; Lucas, P. Linn. Soc. N.S.W. (2) vii, n. spp.
- Graphiphora lichenodes, E. Siberia, GRAESER, p. 216, Berl. Ent. Zeit.
- Grumia, n. g., near Heliothis for G. flora, n. sp., Amdo; Alphéraky, p. 453, Hor. Ent. Ross. xxvi.
- Hadena monoglypha, n. var. ? polyglypha; STAUDINGER, p. 286, Deutsche e. Z. Lep. iv. H. fraudulenta, pl. viii, fig. 5, dorriesi, fig. 4, figured; Romanoff Mem. vi.
- H. subaquila, p. 224, radicosa, p. 225, E. Siberia, GRAESER, Berl. Eut. Zeit. xxxvii, n. spp.
- Hadjina, n. g. (Segetia) for H. lutosa, n. sp., Asia Minor; STAUDINGER, p. 286, Deutsche e. Z. Lep. iv.
- Hamodes simplicia, W. Africa, WEYMER, p. 116, S. E. Z. liii, n. sp.
- Heliocheilus foveolatus, figured, pl. x, fig. 3, Romanoff Mem. vi.
- Heliophobus scillæ, 2 noticed; Chrétien, Le Nat. 1892, p. 7.
- H. hirsutus, Asia Minor, p. 277. H. ? discrepans, Jerusalem, p. 278, STAUDINGER, Deutsche e. Z. Lep. iv; H. grumi, anachoretoides, Central Asia, Alphéraky, p. 447, Hor. Ent. Ross. xxvi: n. spp.
- Heliothis armigera, notes on; Mally, Bull. Dep. Agric. Ent. xxvi, pp. 45-56.
- Hepatica, n. g. of doubtful position, for H. anceps, n. sp., Amurland, pl. xii, fig. 11; STAUDINGER, p. 634, Romanoff Mem. vi.
- Herama, n. g. for Puchnobia mandschurica, Graes.; STAUDINGER, p. 501, Romanoff Mem. vi.

Hiptelia lorezi, note on; Lorez, p. 370, M'l Holocryptis, n. g., Plusiides, p. 256, for H p. 257; Lucas, P. Linn. Soc. N.S.W. (Hydrilla? ornata, Margelan, Staudinger, n. sp.

Hydracia burkhana, Amdo, Alphtraky, p H. medialis, senilis, unimoda, Colorado iii: n. spp.

Hypertra: characters and composition of, H. (6) x, pp. 297-300.

Hypocala: revision of the genus; BUTLES

H. deflorata, figured, pl. xii, fig. 1, Ron

H. moorei, clarissima, Ceylon, BUTLER, Ans

Hypsophila grumi, Central Asia, Alphér

xxvi; H. daishi, China, Alphéraky,

Mem. vi: n. spp.

Isochlora albiritta, grumi, Central Asia, A.
Rosa, xxvi, n. spp.

Janthinea divalis, Malatia, STAUDINGER, p. n. sp.

Kuldecha oberthuri, Amdo, ALPHERAKY, 1
n. sp.

Lasionycia, n. g. for Phlogophora skrælir p. 285, Ent. Tidskr. xiii.

Leocyma cilia, pl. x, fig. 10, borussica, Mem. vi.

Leucania impuncta. Amurland, STAUDINGEI n. sp.

Leucanitis: note on structure of legs of;
Mem. vi; L. hedemanni, Stgr., figured,

L. scolopax, Central Asia, ALPHERAKY, p

Leucocosmia ceres, = (Caradrina euthusa, I

Lithacodia bellicula, Hb., = (Hydrelia a p. 65, Ent. xxv.

Lithocampa ramulosa, figure; Romanoff M Lithophasia, n. g. (Lithocampa), for L. ver fig. 3; STAUDINGER, p. 305, Deutsche

Luperina ditaiu, Brisbane, Lucas, p. 254, L.? malitiosa, Centr. Asia, Alphéa. xxvi: n. spp.

Mamestra: synonymical notes on Smith's 1
Canad. Ent. xxiv. M. legitima, larval s
Bull. Am. Mus. iv. M. adjuncta, pl. v
declinans, fig. 2, incommoda, fig. 3, fig.
Mem. vi.

- M. suavis, Amurland, STAUDINGER, p. 428, pl. vii, fig. 4, Romanoff Mem. vi; M. tancrei, p. 305, longicornis, p. 306, Central Asia, GRAESER, Berl. Ent. Zeit. xxxvii; M. satanella, p. 445, texturata, p. 446, Centr. Asia, Alphéraky, Hor. Ent. Ross. xxvi; M. fuscolutea, Colorado, Smith, p. 250, Ent. News, iii: n. spp.
- felipotis: revision of; BUTLER, Ann. N. H. (6) x, pp. 315-327.
- walkeri, Callao, p. 325, yerburyi, Aden, p. 327, BUTLER, Ann. N. H.
   x, n. spp.
- fesotrosta? incerta, Amurland, STAUDINGER, p. 572, Romanoff Mem. vi, n. sp.
- 1etalopha, n. g. for M. plusina, N. Persia, p. 313, pl. iv, fig. 8, gloriosa, Syria, p. 314, fig. 7, n. spp.; STAUDINGER, Deutsche e. Z. Lep. iv.
- Vetecia cornifrons, figured, pl. i, fig. 2: described, pp. 15-17; SNELLEN, Tijdschr. Ent. xxxv.
- Vetopoceras? beata, Taurus, STAUDINGER, p. 274, Deutsche e. Z. Lep. iv, n. sp.
- Vetoponia pallida, p. 323, gratiosa, p. 325, nubila, p. 326, Asia Minor, STAUDINGER, Deutsche e. Z. Lep. iv, n. spp.
- ficra rosita, Gn., = (derogata, Wlk.); BUTLER, p. 141, Ent. xxv.
- fiselia oxyucantha, varr. described; STAUDINGER, p. 283, Deutsche e. Z. Lep. iv.
- f. sabulosa, E. Siberia, GRAESER, p. 222, Berl. Ent. Zeit. xxxvii, n. sp.
- foma virens, Butl., figured, pl. vii, fig. 1; Romanoff Mem. vi.
- fonostola, n. g., near Gortyna, for M. asiatica, n. sp., Ganssu, pl. ii, fig. 7; Alphéraky, p. 37, Romanoff Mem. vi.
- forrisonia: the N. American species revised; Smith, pp. 78-86, P. U. S. Nat. Mus. xv.
- !ythimna pectinata, Central Asia, ALPHÉRAKY, p. 450, Hor. Ent. Ross. xxvi, n. sp.
- 'ipista (sub Diastema) tigris, Gn., = (lineata, Wlk.); BUTLER, p. 11, Ent. xxv.
- octua variicollis, figured; MABILLE & VUILLOT, pl. xi, fig. 6, Nov. Lep. . favotincta, N. America, Smith, p. 252, Ent. News, iii, n. sp.
- 'onagria phragmiticola, pl. ix, fig. 2, p. 469, impudica, fig. 3, p. 470, Amurland, Staudinger, Romanoff Mem. vi, n. spp.
- govia, n. g., Hypopyrides, for O. tavetensis, n. sp., E. Africa; Holland, Ent. xxv (Supp.), p. 95.
- morphina, n. g., near Heliaca, for O. aurantiaca, n. sp., Amdo, Alphé-RAKY, p. 452, Hor. Ent. Ross. xxvi.
- neocnemis nigrocaput, p. 250, extranea, p. 251, N. America, Smith, Ent. News, iii, n. spp.
- phideres princeps, Gn., is not princeps, Boisd., and should be called Halastus intricatus; BUTLER, p. 375, Ann. N. H. (6) ix.
- phiodes parcemacula, Queensland, Lucas, p. 304, P. Linn. Soc. N.S.W. (2) vi, n. sp.
- phisma pretorice, Transvaal, Distant, p. 239, pl. ii, fig. 7, Naturalist Transvaal, n. sp.
  - 1892. [vol. xxix.]

- Ophiusa myope, frontina, senez, figured; Scott, Austral. Lep. ii, pl. xvi Orresia hartmanni, Möschl., == (cuprea, Saalm.); SNELLEN, p. 79, Tijdsci Ent. xxxv.
- Orrhodia acutula, Jerusalem, STAUDINGER, p. 302, Deutsche e. Z. Le iv; O. ciliata, Amurland, STAUDINGER, p. 516, Romanoff Mem. v n. spp.
- Orthosia kindermanni, n. var., pauli; STAUDINGER, p. 301, Deutsche s. Lep. iv. O. (sub Apamea) reciproca, Wlk., = (breviuscula, Wlk BUTLER, p. 10, Ent. xxv.
- O. scabra, Jerusalem, STAUDINGER, p. 299, Deutsche e. Z. Lep. iv, n. s Osarba punctigera, Wlk., = (Grammodes excavata, Wlk.); BUTLER, p. 1 Ent. xxv.
- Pachetra leucophaa, notes on rearing; JEFFREY, Ent. M. M. xxviii, p. Pachydasys, n. g., Callopistriida, p. 125, for Eutelia consentanea, Wlpl. vi, fig. 5; BUTLER, P. Z. S. 1892.
- Palindia: synonymy of species of; Butler, Ent. xxv, pp. 189-191.
- Panopoda carneicosta and rufimargo, larval stages; Beutenmüller, 68 & 69, Bull. Am. Mus. iv.
- Panthauma, n. g., near Panthea, for P. egregia, n. sp., Amurland, pl. fig. 2; STAUDINGER, p. 404, Romanoff Mem. vi.
- Paragona. n. g., for Acidalia multisignata, Chr., figured, pl. xii, fig. Staudinger, p. 557, Romanoff Mem. vi.
- Perigea conducta, = (Caradrina consocia, Wlk.); Butler, p. 11, Exxv. P. xanthioides, larval stages; Beutenmüller, p. 69, Bull. Mus. iv.
- Phalacra, n. g., for Perigea gemella, Leech; STAUDINGER, p. 568, Romar Mem. vi.
- Phoebophilus? lenis, Jerusalem, STAUDINGER, p. 279, Deutsche e. Z. I iv. n. sp.
- Platydasys, n. g., Callopistriidæ, for P. pryeri, n. sp., Borneo, pl. vi, fig BUTLER, p. 126, P. Z. S. 1892.
- Pleroma apposita, N. America, Smith, p. 252, Ent. News, iii, n. sp. Plotheia: notes on its species; Butler, Ent. xxv, p. 285.
- Plusia gammoides, Bl., = (chilensis, Butl.); REED, p. 68, Act. Soc. Ci. P. oxygramma, Hb., note on; SNELLEN, Tijdschr. Ent. x: p. 18. P. emichi, n. var. paulina; STAUDINGER, p. 310, Deutsel Z. Lep. iv. P. emichi, n. var. dorujtava; STANDFUSS, p. 666, pl. fig. 6, Romanoff Mem. vi. P. gamma, var. of larva; PORRITT, M. M. xxviii, p. 255. P. moneta, habits; FALLOU, p. clxv, Bull. Ent. Fr. lxi. P. aurata, pl. x, fig. 6, splendida, fig. 8, adscripta, fi figured, Romanoff Mem. vi.
- P. augusta, Aintab, p. 309, maria, Jerusalem, p. 311, STAUDIN Deutsche e. Z. Lep. iv; P. inconspicua, Central Asia, GRAE p. 315, Berl. Ent. Zeit. xxxvii; P. agnata, Amurland, STAUDIN p. 547, Romanoff Mem. vi: n. spp.
- Plusilla, n. g., near Telesilla, for P. rosalia, n. sp., Amurland, pl. ix, fig Staudinger, p. 533, Romanoff Mem. vi.

- Pacilogramma, n. g., near Gnamptocera, for P. picatum, n. sp., Borneo, pl. vi, fig. 4; BUTLER, p. 126, P. Z. S. 1892.
- Polia rebecca, Jerusalem, Staudinger, p. 282, Deutsche e. Z. Lep. iv; P. mirafloresa, Loja, Dognin, p. 185, Le Nat. 1892: n. spp.
- Præstilbia, n. g. for P. armeniaca, n. sp, Amasia; Staudinger, p. 288, Deutsche e. Z. Lep. iv.
- Premusia intrahens, Wlk., = (Anophia smaragdaria, Wlk., and Dysedia sibellina, Feld.); Butler, Ent. xxv, p. 213.
- Prionoptera serraoides, Loja, Dognin, p. 85, Le Nat. 1892, n. sp.
- Pseudathyrma, n. g. for part of Hypætra; BUTLER, p. 299, Ann. N. H. (6) x.
- Pseudohadena armata, figured, pl. ii, fig. 6, description amended, p. 35; ALPHÉRAKY, Romanoff Mem. vi.
- P. pugnax, Central Asia, Alpheraky, p. 449, Hor. Ent. Ross. xxvi, n. sp. Pseudomicra, n. g. for Anthophila marginalis and semipurpurea, Wlk.;
  BUTLER, p. 91, Ent. xxv.
- Raddea, u. g., near Mesogona, for R. digna, n. sp., Amdo; Alpheraky, p. 450, Hor. Ent. Ross. xxvi.
- Rancora, n. g., for R. strigata, n. sp., N. America; SMITH, p. 253, Ent. News, iii.
- Rusidrina, n. g., near Caradrina, for R. rasdolnia, n. sp., Amurland, pl. vii, fig. 9; STAUDINGER, p. 491, Romanoff Mem. vi.
- Segetia albopicta, Amurland, Graeser, p. 226, Berl. Ent. Zeit. xxxvii; S.? mandarina, Amurland, Staudinger, p. 492, pl. ix, fig. 4, Romanoff Mem. vi: n. spp.
- Sidemia, n. g., near Trigonophora, for S. snelleni, n. sp., Amurland, pl. vii, fig. 8; STAUDINGER, p. 459, Romanoff Mem. vi.
- Sonagara scitaria, Wlk., = (Timandra cancellata, Wlk.); STAUDINGER, p. 636, Romanoff Mem. vi.
- Sphragifera, n. g., ? near Leocyma, p. 555, for Anthecia sigillata, Mén.; Staudinger, Romanoff Mem. vi.
- Spinipalpa, n. g., near Agrotis, p. 444, for S. maculata, n. sp., Amdo, p. 445; Alphéraky, Hor. Ent. Ross. xxvi.
- Stilbia anomala, varr. described; STAUDINGER, p. 289, Deutsche e. Z. Lep. iv.
- Stilbina, n. g., for S. hypænides, n. sp., Beyrout, pl. iii, fig. 12; STAU-DINGER, Deutsche e. Z. Lep. iv.
- Taniocampa lata, pl. ix, fig. 5, p. 496, stenoptera, p. 497, fig. 6, Amurland, Staudinger, Romanoff Mem. vi, n. spp.
- Tapinostola? (Canagria) nana, Amurland, Staudinger, p. 473, Romanoff Mem. vi, n. sp.
- Turache, synonymy of various species; Butler, Ent. xxv, pp. 61-63.
- Thalpochares: notes on the composition of the genus and the species belonging to allied forms of Eublemmidæ; Butler, Ent. xxv, pp. 91-93. T. albida, varr. described; STAUDINGER, p. 317, Deutsche e. Z. Lep. iv. T. bella, figured, pl. xi, fig. 1; STAUDINGER, Romanoff Mem. vi.

- T. suppuncta, p. 319, suppura, p. 320, substrigula, p. 321, straminea, p. 325
  Asia Minor, Staudinger, Deutsche e. Z. Lep. iv; T. subvessate
  Tunis, Staudinger, p. 288, op. cit. v; T. pulvinariæ, Australia
  Olliff, p. 179, Agric. Gas. N.S.W. iii: n. spp.
- Toxocampa emaculata, E. Siberia, GRAESER, p. 231, Berl. Ent. Zei xxxvii, n. sp.
- Traches similis, Amurland; STAUDINGER, p. 456, pl. ix, fig. 1, Romano Mem. iv, n. sp.
- Trigonophora grumi, Central Asia, Alphéraky, p. 449, Hor. Eut. Ros xxvi, n. sp.
- Ulochlæna superba, Central Asia, Alphéraky, p. 447, Hor. Ent. Ros xxvi, n. sp.
- Valeria? sauberi, E. Siberia, Graeser, p. 223, Berl. Ent. Zeit. xxxv n. sp.
- Varnia fenestrata, Lace, POUJADE, p. 270, pl. xi, fig. 6, N. Arch. M. Paris (3) iii, n. sp.
- Virgo, n. g. for Nonagria amana, Stgr., figured, pl. viii, fig. STAUDINGER, p. 467, Romanoff Mem. vi.
- Xanthia divergens, Butl., figured, pl. ix, fig. 8, Romanoff Mem. vi.
- Xylina ledereri, Rhodus, STAUDINGER, p. 304, Deutsche e. Z. Lep. iv; brachyptera, Amurland, STAUDINGER, p. 520, pl. ix, fig. 9, Roman Mem. vi: n. spp.
- Xylomiges: the N. American species revised; SMITH, pp. 65-78, pl. P. U. S. Nat. Mus. xii.
- X. peritalis, p. 69, ochracea, p. 75, N. America; Smith, P. U. S. N. Mus, xv, n. spp.
- Xylomoia, n. g., near Chloantha, for X. separata, n. sp., Amurland, pl. fig. 8; Staudinger, p. 451, Romanoff Mem. viii.
- Zalissa, Wlk, referred to Agaristida, = (Saudyra, Stretch.); Butl Ent. xxv, p. 214.

### DELTOIDIDE.

- [Cf. Alphéraky (779), Butler (143), Distant (216), Lucas (5 Poujade (702), Staudinger (779, 886).]
- Bocana flavopunctalis, Laos, Poujade, p. 275, pl. xi, fig. 10, N. A. Mus. Paris (3) iii, n. sp.
- Bomolocha rhombalis, figured, pl. ii, fig. 10, Romanoff Mem. vi.
- B. bipartita, Amurland, STAUDINGER, p. 625, pl. xiv, fig. 12, Roma Mem. vi, n. sp.
- Episeuxis athiops, Transvaal, DISTANT, p. 239, pl. ii, fig. 2, Natur Transvaal, n. sp.
- Gynaphila, n. g., near Hypenodes, for G. maculifera, n. sp., Amurl pl. xiv, fig. 13; STAUDINGER, p. 629, Romanoff Mem. vi.
- Helia curripalpalis, Butl., = (quadra, Graes.); STAUDINGER, p. Romanoff Mem. vi.
- Herminia semicircularis, Brisbane, Lucas, p. 260, P. Linn. Soc. N.:
  (2) vii, n. sp.

- Hollandia, n. g., p. 295, for H. sigillata, n. sp., W. Africa, p. 296; BUTLER, Ann. N. H. (6) x.
- Hypena conspersalis, figured, pl. xiii, fig. 10, Romanoff Mem. vi; H. ravalis, varr. described; STAUDINGER, p. 334, Deutsche e. Z. Lep. iv.
- H. mardinalis, Asia Minor, STAUDINGER, p. 336, Deutsche e. Z. Lep. iv, n. sp.
- Nodaria amurensis, figured, pl. xiii, fig. 1, Romanoff Mem. vi.
- Pangrapta marmorata, pl. xiii, fig. 5, incisa, fig. 6, flavomacula, fig. 7, suaveola, fig. 8, costæmacula, fig. 9, figured, Romanoff Mem. vi.
- Pangrapta griscola, Amurland, STAUDINGER, p. 620, pl. xiv, fig. 11, Romanoff Mem. vi, n. sp.
- Pechipogon? grandis, China, Alphéraky, p. 49, pl. ii, fig. 11, Romanoff Mem. vi. n. sp.
- Rivula unctalis, Amurland, STAUDINGER, p. 631, Romanoff Mem. vi, n. sp.
- Zanclognatha fumosa, pl. xiii, fig. 2, robiginosa, fig. 3, bistrigata, fig. 4, figured, Romanoff Mem. vi.
- Z. cristulalis, p. 610, pl. xiv, fig. 7, violacealis, fig. 6, p. 612, umbrosalis, fig. 8, p. 613, Z. P helialis, fig. 9, p. 615, obliqua, fig. 10, p. 616, Amurland, STAUDINGER, Romanoff Mem. vi, n. spp.

#### GEOMETRIDÆ.

- [Cf. Alphéraky (4, 779), Butler (141), Distant (216), Dognin (220), Druce (225, 347), Fuchs (313), Graeser (356, 357), Gumppenberg (380), Heylaerts (403), Holland (408), Klemensiewicz (499), Lower (565), Lucas (568, 569), Meyrick (603, 604, 605), Poujade (702), Röber (767), Skinner & Mengel (850), Standfuss (779), Staudinger, (886, 887, 890, 891), Steinert (897), Swinhoe (909), Thierry-Mieg (923), Weymer (1003).]
- Classification of the Geometrinæ of Europe; they are divided into six families and 117 genera, of which the characters are given; MEYRICK, Tr. E. Soc. 1892, pp. 53-140.
- The classifications of Meyrick and Gumppenberg discussed and criticised, many remarks on generic nomenclature; Rebell, S. E. Z. liii, pp. 247-268.
- Colour variation in larvæ; Poulton (703).
- Abraxas khasiana, Khasia Hills, SWINHOE, p. 17, Tr. E. Soc. 1892; A. subhyalinata, Flores, RÖBER, p. 333, Tijdschr. Ent. xxxiv (fig. in xxxv): n. spp.
- Achlora gaujoniaria, Ecuador, Dognin, p. 186, Le Nat. 1892, n. sp.
- Acidalia lumenaria, Hb., = (Zanclopteryx zincaria, Gn.); SNELLEN, p. 20, Tijdschr. Ent. xxxv. A. privata, fig. 23, calidata, fig. 24, figured, pl. lii, Biol. Centr. Am. Heter. ii. A. pallidata, larva described; HABICH, p. 161, S. E. Z. liii. A. luridata, metamorphoses; REBEL, p. 518, Verh. z.-b. Wien, xlii.

- A. bicertaria, Tunis, STAUDINGER, p. 290, Deutsche e. Z. Lep. v; A. subochraria, Central Asia, p. 144, fasciata, Syria, p. 145, striolata, Beyrout, p. 146, pl. ii, fig. 5, incisaria, fig. 6, p. 147, Spain, nocturna, p. 148, sartharia, p. 149, tristriata, p. 150, effeminata, p. 151, Central Asia, completa, Algeria, p. 152, lutulentaria, Spain, p. 153, grisescens, p. 155, annubiata, p. 156, Central Asia, STAUDINGER, Deutsche e. Z. Lep. v. A. latelineata, Central Asia, GRAESER, p. 316, Berl. Ent. Zeit. xxxvii ..., A.? obrimo, pl. lii, fig. 13, A. cavonu, fig. 14, p. 117, onca, fig. 15, celenna, fig. 16, A.? enona, fig. 17, A. carna, fig. 18, p. 118, nursica. ..., fig. 19, olmia, fig. 20, p. 119, caseca, fig. 21, catenes, fig. 22, p. 120. ..., A.? numitoria, fig. 25, carmenta, fig. 26, p. 121, A.? ocrisia, fig. 27, A. nycteis, fig. 28, odrussa, fig. 29, p. 122, Central America, DRUCE E., Biol. Centr. Am. Heter. ii; A. minuscula, nigroapicata, Chiriqui ..., tirescens, Colombia, rubricolor, rubellula, nubicolor, Cayennee, THIERRY-MIEG, p. 235, Le Nat. 1892: n. spp.
- Acropteris luteopictata, Laos, POUJADE, p. 274, pl. xi, fig. 9, N. Arch. Muz. Raris (3) iii, n. sp.
- Acrotomia muta, pl. xlv, figs. 15 & 16, polla, figg. 17 & 18, A? mucia, figs. —g.
  19 & 20, p. 48, A. minucia, figg. 21 & 22, A.? bola, fig. 23, nis— a,
  fig. 24, p. 49, Central America; DRUCE, Biol. Centr. Am. Heter.
- Agathia codina, pl. i, fig. 3, p. 7, gemma, p. 8, Khasia Hills, SWINHOE, T. E. Soc. 1892; A. iodioides, distributa, Queensland, Lucas, p. 296, F. Linn. Soc. N.S.W. (2) vi : n. spp.
- Amelora, n. g., Selidosemidæ, p. 646, for A. goniota, p. 647, demistis, p. 648, amblopa, p. 649, catacris, p. 650, arotras, p. 651, n. spp., Australia; MEYRICK, P. Linn. Soc. N.S.W. (2) vi.
- Amphidasys betularia, variation in colour of larva; BATESON, p. 213, Tr. E. Soc. 1892. A. superans, Butl., referred to Boarmia; GRAESER, p. 232, Berl. Ent. Zeit. xxxvii. A. arnobia, figured, pl. xlviii, fig. 4, Biol. Centr. Am. Heter. ii.
- Amphidasis matharma, Central America, DRUCE, p. 71, pl. xlviii, figg. 5 & 6, Biol. Centr. Am. Heter. ii, n. sp.
- Anaitis mundata, Beyrout, p. 216, submundulata, Taurus, p. 217, musculata, Asia Minor, pl. ii, fig. 19, p. 218, STAUDINGER, Deutsche e. Z. Lep. v, n. spp.
- Anaplodes pistaciaria, figured, pl. xlix, fig. 20, Biol. Centr. Am. Heter. ii. Anhibernia, n. g. for Hybernia (A.) orientalis, n. sp., Syria; STAUDINGER p. 170, Deutsche e. Z. Lep. v.
- A. burætica, Siberia, Staudinger, p. 376, pl. iii, fig. 8, Deutsche e. Z. Lep v, n. sp.
- Anisodes lichenaria, intermixtaria, Khasia Hills, SWINHOE, p. 8, Tr. E. Soc 1892; A. melitia, pl. li, fig. 3, caletra, fig. 4, carbina, fig. 5, Centra America, DRUCE, p. 104, Biol. Centr. Am. Heter. ii; A. melano chrysus, Loja, Dognin, p. 237, Le Nat. 1892: n. spp.
- Anteia canescens, Rockhampton, Lucas, p. 300, P. Linn. Soc. N.S.W. (2 vi, n. sp.

- Anthyria, n. g., p. 2, type Hyria gruturia, Wlk., with A. iole, n. sp., Khasia Hills, p. 3, pl. i, fig. 7; Swinhoe, Tr. E. Soc. 1892.
- Anticypella, n. g. for gigantaria, Stgr.; MEYRICK, p. 108, Tr. E. Soc. 1892.
- Aphrogeneia, n. subg. of Acidalia; Gumppenberg, p. 257, N. Acta Ac. L.-C. Nat. cur. lviii.
- Apicia medusa, pl. xliv, fig. 20, atillu, fig. 21, mesada, fig. 22, melenda, figg. 23 & 24, p. 39, aurana, fig. 25, micca, fig. 26, atia, fig. 27, mera, figg. 28 & 29, p. 40, aurelia, p. 41, pl. xlv, fig. 1, megania, p. 42, fig. 2, Central America, DRUCE, Biol. Centr. Am. Heter. ii, n. spp.
- Aporoctena, n. g., Selidosemidæ, for A. scierodes, n. sp., Queensland; MEYRICK, p. 622, P. Linn. Soc. N.S.W. (2) vi.
- Artemidora, n. g. for maracandaria, Ersch.; MEYRICK, p. 112, Tr. E. Soc. 1892.
- Artiora, n. g. for evonymaria, Schiff.; MEYRICK, p. 114, Tr. E. Soc. 1892.
- Asellodes laternaria, figured, pl. li, fig. 8, Biol. Centr. Am. Heter. ii.
- Aspilates curvaria, Evers, figured, pl. iii, fig. 5, Q described, p. 70; Alphé-Baky, Romanoff Mem. vi.
- A. revocaria, Beyrout, STAUDINGER, p. 203, pl. ii, fig. 16, Deutsche e. Z. Lep. v, n. sp.
- Asthena prasina, Khasia Hills, SWINHOR, p. 13, pl. i, fig. 10, Tr. E. Soc. 1892; A. pellucida, Brisbane, Lucas, p. 253, P. Liun. Soc. N.S.W. (2) vii; A. carnana, Mexico, pl. lii, figg. 10 & 11, neclas, Guatemala, p. 12, DRUCE, p. 115, Biol. Centr. Am. Heter. ii: n. spp.
- Astrapephora, n. g., near Ortholithu, for A. romanovi, n. sp., Amdo, Alphéraky, p. 457, Hor. Ent. Ross. xxvi.
- Atomophora, n. g., near Lithostege, p. 65, for A. falsaria, n. sp., Ganssu, Alpheraky, p. 66, Romanoff Mem. vi.
- Azelina terrenaria, Loja, Dognin, p. 106, Le Nat. 1892; A. attagena, barbarata, fortunata, Ecuador, Dognin, p. 186, Le Nat. 1892; A. contorta, Peru, Thierry-Mieg, p. 217, Le Nat. 1892; A. asterodia, pl. xlvii, figg. 8 & 9, astapa, figg. 10 & 11, p. 62, melissa, fig. 12, metella, fig. 13, aurunca, fig. 14, p. 63, micca, figg. 15 & 16, rumina, figg. 17 & 18, p. 64, Central America, Druce, Biol. Centr. Am. Heter. ii : n. spp.
- Bessophora, n. g., = (Ptychoptera, Christoph.); MEYBICK, p. 63, Tr. E. Soc. 1892.
- Biston hirtarius, n. var. hanoviensis; Heymons, p. 446, pl. xv, fig. 9, Berl. Ent. Zeit. xxxvi.
- Boarmia repandata, n. var. nobilis; ALPHERAKY, p. 61, Romanoff Mem. vi.
- B. exustaria, Siberia, STAUDINGER, p. 378, pl. iii, fig. 9, Deutsche e. Z. Lep. v; B. nobilitaria, p. 173, variolaria, p. 174, subrepandata, p. 175, Central Asia, STAUDINGER, Deutsche e. Z. Lep. v; B. conjungens, Amdo, Alphéraky, p. 456, Hor. Ent. Ross. xxvi; B. rybakowi, Ganssu, Alphéraky, p. 61, pl. iii, fig. 4, Romanoff Mem. vi; B.

leucopterata, Laos, POUJADE, p. 273, pl. xi, fig. 7, N. Arch. Mas. Paris (3) iii; B. noctiluca, pl. xlviii, fig. 7, B.? cubira, fig. 8, p. 73, B. nephalia, fig. 10, p. 75, brotes, fig. 11, nepia, fig. 12, p. 76, Datcz, Biol. Centr. Am. Heter. ii: n. spp.

Boletobia robiginosa, p. 332, pl. iv, fig. 13, detersa, p. 333, fig. 14, Am. Minor, STAUDINGER, Deutsche e. Z. Lep. iv, n. spp.

Bonatea, n. g., Boarminæ, for Oxydia duciata, Maassen, figured, pl. xlix, figg. 6 & 7; DRUCE, p. 82, Biol. Centr. Am. Heter. ii.

Bronchelia subroraria, p. xlix, fig. 1, conjugaria, fig. 4, detexta, fig. 3, figured, Biol. Centr. Am. Heter. ii.

B. masica, Mexico, DRUCE, p. 81, pl. xlix, fig. 4, Biol. Centr. Am. Heter. ii, n. sp.

Bryoptera pruna, Loja, Dognin, p. 59, Le Nat. 1892; B. (?) candara, Panama, Druce, p. 80, pl. xlviii, fig. 26, Biol. Centr. Am. Heter. ii : n. spp.

Byssodes appropriata, figured, pl. l, fig. 25, Biol. Centr. Am. Heter. ii.

B. miletia, fig. 27, navia, fig. 26, p. 99, nasada, fig. 28, p. 100, Central America, DRUCE, Biol. Centr. Am. Heter. ii, u. spp.

Caberodes azona, pl. xlvi, fig. 6, messens, fig. 7, Mexico, Decce, p. 53, Biol. Centr. Am. Heter. ii, n. spp.

Culothyeanis amaturaria, preparatory stages; Kellicott, Canad. Ent. xxiv, p. 242.

Cambogia obada, pl. lii, figg. 1 & 2, numida, fig. 3, p. 112, occia, fig. 4, odatis, fig. 5, numeria, fig. 6, p. 113, nucula, fig. 7, cedon, fig. 8, C.? necula, fig. 9, p. 114, Central America, Druce, Biol. Centr. Am. Heter. ii, n. spp.

Curphoxera ptelearia, further notes on; RILEY, Ins. Life, v, p. 40.

Chlenias melanoxysta, p. 633, sonæa, p. 665, Australia, MEYRICK, P. Linn. Soc. N.S.W. (2) vi, n. spp.

Charodes clemitaria, full-grown larva; BEUTENMÜLLER, p. 79, Bull. Am. Mus. iv.

Chrysoctenis, n. g., for filacearia, H.-S.; MEYRICK, p. 86, Tr. E. Soc. 1892. Cidaria minorata, larva described; Habich, S. E. Z. liii, p. 160. C. tetrica, figured, pl. iii, fig. 11, Romanoff Mem. vi.

C. inconsiderata, Amasia, p. 226, apiciata, p. 227, fidonaria, p. 229, alexandraria, p. 230, interpositaria, p. 231, prapositaria, p. 233, postpositaria, p. 234, Central Asia, ignorata, Caucasus, p. 235, desperata, p. 236, antelataria, p. 237, pl. ii, fig. 21, Central Asia, numidiata, Algeria, fig. 22, p. 239, viduata, Central Asia, p. 241, sintenisi, Asia Minor, p. 242, mærens, p. 245, rogata, p. 248, rejectaria, p. 249, Central Asia, STAUDINGER, Deutsche e. Z. Lep. v; C. semenovi, p. 458, excentricata, with var. virginea, p. 459, Amdo, Alphéraky, Hor. Ent. Ross. xxvi; C. tima, Peru, Thierry-Mieg, p. 217, Le Nat. 1892; C. cinereolimitata, Colombia, Thierry-Mieg, p. 235, t.c.; C. apiciflava, alboscripta, Loja, Dognin, p. 274, Le Nat. 1892; n. spp.

Cimelia raulogeri, Tunis, STAUDINGER, p. 291, pl. iii, fig. 11, Deutsche e. Z. Lep. v, n. sp.

- Cimicodes portentosa, Peru, THIERRY-MIEG, p. 235, Le Nat. 1892, n. sp. Cleora godmani, Mexico, DRUCE, p. 72, pl. xlviii, fig. 9, Biol. Centr. Am. Heter. ii, n. sp.
- Cleta? minuta, Central America, DRUCE, p. 111, pl. li, fig. 30, Biol. Centr. Am. Heter. ii, n. sp.
- Cnidia, n. subg. of Acidalia; Gumppenberg, p. 223, N. Acta Ac. L.-C. Nat. cur. lviii.
- Comibæna venezuelata, figs. 14 & 15, invasata, fig. 18, pl. xlix, figured, Biol. Centr. Am. Heter. ii.
- C. belisama, p. 87, pl. xlix, figg. 16 & 17, callicula, fig. 19, p. 88, Panama, Deuce, Biol. Centr. Am. Heter. ii; C. ecuadorata, Zamora, Dognin, p. 186, Le Nat. 1892; C. lafayaria, Loja, Dognin, p. 206, t. c.: n. spp.
- Conosara, n. g., Selidosemidæ, for C castanea, n. sp., Tasmania; MEYRICK, p. 660, P. Linn. Soc. N.S.W. (2) vi.
- Coricia bicincta, Loja, Dognin, Le Nat. 1892, p. 144, n. sp.
- Cratoptera arvina, p. 34, pl. xliv, fig. 10, mena, fig. 11, atina, figs. 12 & 13, p. 35, Central America, DRUCE, Biol. Centr. Am. Heter. ii, n. spp.
- Criomacha, n. g. for Chlenias belidearia, Feld.; MEYRICK, p. 659, P. Linn. Soc. N.S.W. (2) vi.
- Crocallis? eduxaria, Loja, Dognin, Le Nat. 1892, p. 123, n. sp.
- Dasydia tarpeia, Colombia, THIERRY-MIEG, p. 236, Le Nat. 1892, n. sp.
- Dasypteroma, n. g., near Hibernia, for D. thaumasia, n. sp., Spain, pl. ii, fig. 10; STAUDINGER, p. 168, Deutsche e. Z. Lep. v.
- Deilinia eccentritis, Australia, p. 635, ænias, Tasmania, lithodora, p. 637, cremnias, p. 638, ochthadia, p. 639, Australia, MEYRICK, P. Linn. Soc. N.S.W. (2) vi, n. spp.
- Descoreba rosalinda, Brazil, THIERRY-MIEG, p. 236, Le Nat. 1892, n. sp. Diastictis margaritis, Queensland, MEYRICK, p. 588, P. Linn. Soc. N.S.W. (2) vi. n. sp.
- Dichromodes ptilomacra, S. Australia, Lower, p. 8, Tr. R. Soc. S. Austr. xv; D. petrina, New Zealand, Meyrick, p. 216, Tr. N. Z. Inst. xxiv: n. spp.
- Discalma, n. g., for Tephrina normata, Wlk.; MEYRICK, p. 590, P. Linn. Soc. N.S.W. (2) vi.
- Doryodes acutaria, full-grown larva; BEUTENMÜLLER, p. 78, Bull. Am. Mus. iv.
- Drepanodes valeria, Peru, THIEBRY-MIEG, p. 217, Le Nat. 1892; D. moneta, pl. xliv, fig. 2, astanda, fig. 3, p. 30, asinu, fig. 4, D.? melina, fig. 5, p. 31, Central America, DRUCE, Biol. Centr. Am. Heter. ii : n. spp.
- Drymoptila, n. g., Selidosemidæ, for D. temenitis, n. sp., Australia; Meybick, P. Linn. Soc. N.S.W. (2) vi, p. 670.
- Ectropis pristis, p. 626, argalea, isombra, p. 627, Australia, MEYRICK, P. Linn. Soc. N.S.W. (2) vi, n. spp.
- Egea temeraria, Central Asia, STAUDINGER, p. 196, Deutsche e. Z. Lep. vi, n. sp.

- Ellopia cinereostrigaria, Galicia, KLEMENSIEWICZ, p. 67, SB. z.-b. Wien, xlii, n. sp.
- Ennomos bitias, Panama, DRUCE, p. 69, pl. xlviii, figs. 1 & 2, Biol. Centr. Am. Heter. ii, n. sp.
- Eois subtectata, figured, pl. li, figg. 18 & 19, Biol. Centr. Am. Heter. ii.
- E. ascantia, pl. li, fig. 20, batea, fig. 21, camma, fig. 22, p. 109, nundine, figg. 23 & 24, ops, figg. 25 & 26, cassandra, fig. 27, E.? ceno, fig. 28, p. 110, catana, fig. 29, p. 111, Central America, DRUCE, Biol. Centr. Am. Heter. ii, n. spp.
- Ephoria, n. g. for arenosa, Butl.; MEYRICK, p. 109, Tr. E. Soc. 1892.
- Ephyra megista, p. 101, pl. li, fig. 1, citrina, p. 102, fig. 2, Central America, DRUCE, Biol. Centr. Am. Heter. ii, n. spp.
- Epicompsa, n. g., Selidosemidæ, for E. xanthocrossa, n. sp., S. Australia; MEYRICK, P. Linn. Soc. N.S.W. (2) vi.
- Epifidonia fractistriga, Ganssu, Alpheraky, p. 65, pl. iii, fig. 7, Romanoff Mem. vi, n. sp.
- Epione? polydora, Peru, THIERRY-MIEG, p. 235, Le Nat. 1892; E. (?) asteria, p. 43, pl. xlv, figg. 3 & 4, megamede, p. 44, figg. 5 & 6, Drice, Biol. Centr. Am. Heter. ii : n. spp.
- Erateina semiluctuata, Colombia, THIERRY-MIEG, p. 236, Le Nat. 1892; E. tibicina, Loja, Dognin, p. 238, Le Nat. 1892; E. artemisia p. 97, mecyra, rhesa, p. 98, medama, aroma, p. 99, hewitsomi, artemis, p. 100, buckleyi, arocha, p. 101, meduthina, bossora, p. 102, N. W. South America, capua, St. Catharina, masura, media, p. 103, peloria, wheeleri, p. 104, artabates, faventia, p. 105, whitelyi, necysia, p. 106, hermaa, tryphosa, p. 107, N. W. South America, DRUCE, Ann. N. H. (6) ix: n. spp.
- Eremia calestinaria, Central Asia, Alphebaky, p. 456, Hor. Ent. Ross. xxvi, n. sp.
- Ergavia bogotaria, figured, pl. xlviii, fig. 20, Biol. Centr. Am. Heter. ii.
- E. calvina, figg. 21 & 22, caletra, fig. 23, assecoma, figg. 24 & 25, Mexico, Druce, p. 79, pl. xlviii, Biol. Centr. Am. Heter. ii, n. spp.
- Erosiidæ, systematic position discussed; Heylaerts, p. 15, Ann. Ent. Belg. xxxvi.
- Erosia incisaria, figured, pl. liii, figg. 3 & 4, Biol. Centr. Am. Heter. ii.
- E. (Dirades) tridentata, p. 14, bidentata, p. 15, Java, HEYLAERTS, Ann. Ent. Belg. xxxvi; E. opigena, pl. liii, figg. 1 & 2, p. 126, omana, fig. 5, cassera, figg. 6 & 7, E. P carilla, fig. 8, E. ocalea, fig. 9, p. 127, ora, fig. 10, p. 128, DRUCE, Biol. Centr. Am. Heter. ii: n. spp.
- Erosina meropia, Costa Rica, DRUCE, p. 55, pl. xlvi, fig. 10, Biol. Centr. Am. Heter. ii, n. sp.
- Erythrolophus, n. g., Ephyrides, p. 10, for E. fascicorpus, n. sp., Khasia Hills, p. 11; Swinhoe, Tr. E. Soc. 1892.
- Eubolia? flavescens, Ordoss, Alphéraky, p. 68, pl. iii, fig. 6, Romanoff Mem. vi, n. sp.
- Eucosmia montivagata, larva; Püngeler, S. E. Z. liii, p. 75.

- Eumera, n. g., near Eugonia for E. regina, n. sp., Amasia, pl. ii, fig. 8; STAUDINGER, p. 162, Deutsche e. Z. Lep. v.
- Eupithecia innotata and frazinata, note on; Habich, p. 159, S. E. Z. liii, E. castigata, note on melanic var. of; Tugwell, Ent. xxv, p. 41. E. extensaria, larval habits; Porritt, Ent. M. M. xxviii, p. 122.
- E. horticolaria, Rhineland, Fuchs, JB. nass. Ver. xlv, p. 102; E. accurata, Central Asia, p. 251, irritaria, Armenia, p. 253, albidulata, Caucasus, p. 254, kuldschaensis, p. 255, æquata, p. 256, marginata, p. 257, Central Asia, subfenestrata, Trans-Caucasus, p. 259, STAUDINGER, Deutsche e. Z. Lep. v; E. filicata, p. 1, rigida, p. 2, pl. 1, fig. 6, Khasia Hills, SWINHOE, Tr. E. Soc. 1892: n. spp.
- Eusarca jacularia, n. var., minor; ALPHÉRAKY, p. 71, Romanoff Mem. vi. E. acidaria, p. 204, dispar, p. 205, Central Asia, STAUDINGER, Deutsche e. Z. Lep. v, n. spp.
- Euschema ares, characters and synonymy; WEYMER, p. 122, S. E. Z. liii.
- Fergana, n. g., near Chemerina, for F. oreophila, n. sp., Alai Mts., pl. ii, fig. 9; STAUDINGER, p. 166, Deutsche e. Z. Lep. v.
- Fidonia cretacea, Central Asia, STAUDINGER, p. 197, Deutsche e. Z. Lep. v; F. opulenta, Peru, THIERRY-MIEG, p. 216, Le Nat. 1892; F. incandescens, Chili, THIERRY-MIEG, p. 236, t. c.: n. spp.
- Geometra iridaria, figured, pl. xlix, fig. 5, Biol. Centr. Am. Heter. ii.
- G. muscipunctata, Loja, Dognin, p. 186, Le Nat. 1892; G. carbina, pl. xlix, fig. 8, aturia, fig. 9, Mexico, Druce, p. 84, Biol. Centr. Am. Heter. ii: n. spp.
- Glaucopteryx immaculata, Greenland, SKINNER, p. 159, P. Ac. Phil. 1892, n. sp.
- Gnopharmia, n. g., p. 181, for Gnophos colchidaria, Led., with G. rubraria, Syria, pl. ii, fig. 12, p. 184, maculifera, Samarcand, fig. 13, p. 187; STAUDINGER, Deutsche e. Z. Lep. v.
- Gnophos vastaria, Central Asia, p. 189, dubitaria, Persia, p. 192, STAU-DINGER, Deutsche e. Z. Lep. v, n. spp.
- Gonanticlea, n. g., p. 3, for G. aversa, n. sp., Khasia Hills, p. 4; SWINHOE, Tr. E. Soc. 1892.
- Gonodela kilimanjarensis, rhabdophoru, E. Africa, Holland, Ent. xxv (Supp.), p. 95, n. spp.
- Gynopteryx fundaria, multiple synonymy; DRUCE, p. 36, Biol. Centr. Am. Heter. ii.
- G. asopia, pl. xliv, fig. 14, mynes, fig. 15, megasa, fig. 16, minoa, fig. 17, p. 37, asteria, figg. 18 & 19, p. 38, Central America, Biol. Centr. Am. Heter. ii, n. spp.
- Halia berytaria, Beyrout, STAUDINGER, p. 200, Deutsche e. Z. Lep. v, n. sp.
- Halthia nigripars, Khasia Hills, SWINHOE, p. 16, pl. i, fig. 1, Tr. E. Soc. 1892; H. aurantiaca, Ganssu, Alphéraky, p. 56, pl. iii, fig. 2, Romanoff Mem. vi : n. spp.
- Hemithea idaa, Khasia Hills, SWINHOE, p. 6, Tr. E. Soc. 1892, n. sp.

Heriote tenebrics. Loja, Dogwin, p. 185, Le Nat. 1892; H.? a
Persama, Deuce, p. 25, pl. xliv, fig. 1, Biol. Centr. Am. He
H.? capacina. THERRY-MIEG, p. 217, Le Nat. 1892: 1

Huners nester. Gustemala, DRUCE, p. 70, pl. xlviii, fig. 3, Biol. Cen. Heter. ii. n. sp.

Hydromens gypeonels, S. Australia, Lower, p. 11, Tr. R. Soc. S. xv. z. sp.

Byprodreme rieda, Colombia, THIRREY-MIRG, p. 237, Le Nat. 18 asseina, pl. xlvi. figg. 24-26, bubona, figg. 27 & 28, p. 58, pl. xlvii. figg. 1 & 2 asseova, figg 4-7, p. 59, Central America, I Biol. Centr. Am. Heter. ii : n. spp.

Hyperycire refo-fescieta, Leon, POUJADE, p. 274, pl. xi, fig. 8, N Mas. Paris 3' iii, n. sp.

Hyperirums surentiacon, p. 297, difundens, p. 298, Australia, l. P. Linn. Soc. N.S.W. (2) vi; H. engramma, S. Australia, l. p. 14. Tr. R. Soc. S. Austr. xv : n. spp.

Idea Lampooni, p. 14. albivertez, p. 15. Khasia Hills, SWINHOE, Soc. 1892. n. app.

Inis impirente, bernarde, p. 293, cresulata, p. 294, multitineta, ni p. 255. Australia, Lucas, P. Linn. Soc. N.S.W. (2) vi; I. con marie. Lucas, p. 252, op. cit. vii: I. ipomopsis, S. Australia, L. p. 14. Tr. R. Soc. S. Austr. xv: n. spp.

Kyratiche, n. g. for Orthelithe (K.) obstinate, n. sp., Central Asia, fig. 18; Statusger, p. 213, Deutsche e. Z. Lop. v.

Leurspan, n. g., = (Trichopleure, Stgr.): MEYRICK, p. 70, Tr. E. Soc Leurs is series, pl. xlvi, fig. 3, magazire, fig. 5, DRUCE, p. 52, Biol. Am. Heter. ii, n. app.

Limerou, E. g. mear Egen, for L. mecreria, n. sp., Sarepta, pl. ii, fi STATINGER, p. 195, Deutsche e. Z. Lep. v.

Linescope evincensenii, Kurdistan, Standeuss, p. 668, pl. xv, Romanoff Mem. vi. n. sp.

Lober in ex comin. Loja, Dognin, p. 274, Le Nat. 1892; L. ebriok georgeoise. Loja, Dognin, p. 295, t. c.: n. spp.

Li mogeophia speciosa, p. 633, isocyma, p. 634, Australia, Meyrick, P. See, N.S.W. (2) vi. n. spp.

Longuis, n. g., near Acidelia, for L. extraordinaria, n. sp., Beyrout, fig. 7; STATDINGER, p. 157, Deutsche e. Z. Lep. v.

Lycusges donocusi, Transvaal, Distant, p. 240, pl. ii, fig. 4, Nat Transvaal, n. sp.

Lucinas cajamasa, Loja, Dogwin, p. 185, Lo Nat. 1892, n. sp.

Macuria suglista, Gn., = (arata, Saalm.): SNELLEX, p. 83, Tijdsch xxxv.

M. m. mata, anomalata, Amdo, Alpheraky, p. 455, Hor. Ent. Ross M. drepanita, Flores, goramata, Goram, Röber, p. 331, Ti Ent. xxxiv (figg. in xxxv): n. spp.

Marceia? undifusciata. Borneo, BUTLER, p. 132, pl. vi, fig. 10, P 1892, n. sp.

- Mecoceras aulis, Mexico, DRUCE, p. 95, pl. l, fig. 15, Biol. Centr. Am. Heter. ii, n. sp.
- Megalochlora, n. g., near Euchloris; MEYRICK, p. 95, Tr. E. Soc. 1892.
- Megametoron, n. g., Boarmiidæ, for M. piperaium, n. sp., Ganssu, pl. iii, fig. 3; Alpheraky, p. 58, Romanoff Mem. vi.
- Metanema cabiria, p. 67, pl. xlvii, figg. 20 & 21, bonadea, figg. 22 & 23, neon, figg. 24 & 25, brea, fig. 26, p. 68, Central America, DRUCE, Biol. Centr. Am. Heter. ii, n. spp.
- Metrocampa glaucias, Australia, MEYRICK, p. 644, P. Linn. Soc. N.S.W. (2) vi, n. sp.
- Micronia: systematic position; SNELLEN, p. 19, Tijdschr. Ent. xxxv.
  M. archilis, Obert., referred to Pseudomicronia, and figured, pl. iii, fig. 1; Alphéraky, p. 53, Romanoff Mem. vi.
- M. simpliciata, Goram, Röber, p. 331, Tijdschr. Ent. xxxiv (fig. in xxxv), n. sp.
- Mictodoca, n. g., Selidosemidæ, for M. tozeuta, n. sp., Australia; MEYRICK, p. 661, P. Linn. Soc. N.S.W. (2) vi.
- Mnesampela, n. g., with M. comarcha, p. 656, lenæa, p. 657, n spp., Australia, and including Idiodes privata, Gn., and Chlonias fucata, Feld.; MEYRICK, P. Linn. Soc. N.S.W. (2) vi.
- Mnesampela privata, metamorphoses; LIDGETT, Vict. Nat. ix, p. 67.
- Mochlotona, n. g., Selidosemidæ, p. 672, for M. phasmatias, n. sp., Australia, p. 673; MEYRICK, P. Linn. Soc. N.S.W. (2) vi.
- Monoctenia iximia, calladelpha, S. Australia, Lower, p. 7, Tr. R. Soc. S. Austr. xv, n. spp.
- Mychonia? bityla, pl. xlvii, fig. 27, noela, fig. 28, Central America, DBUCE, p. 69, Biol. Centr. Am. Heter. ii, n. spp.
- Myinodes, n. g., for interpunctaria, H.-S.; MEYRICK, p. 82, Tr. E. Soc. 1892.
- Mysticoptera, n. g., for sexalisata, Hb.; MEYRICK, p. 62, Tr. E. Soc. 1892. Nelo, see Bombyces, p. 242.
- Nemoria pretiosaria, n. var. gigantaria; STAUDINGER, p. 143, Deutsche e. Z. Lep. v. N. iris, figured, pl. xlix, fig. 10, Biol. Centr. Am. Heter. ii.
- N. mollissima, N.? vitiosaria, Ecuador, Dognin, p. 186, Le Nat. 1892;
  N. assa, Costa Rica, Druce, p. 85, pl. xlix, figg. 11 & 12, Biol. Centr.
  Am. Heter. ii: n. spp.
- Neolythria, n. g., near Lythria, p. 71, for N. abraxaria, n. sp., China, p. 72, pl. iii, fig. 8; Alphéraky, Romanoff Mem. vi.
- Neoteristis, n. g., Selidosemida, for N. paraphanes, n. sp., Tasmania; MEYRICK, p. 672, P. Linn. Soc. N.S.W. (2) vi.
- Neurophana amatonga, E. Africa, Vuillot, p. exciv, Bull. Soc. Ent. Fr. lxi, n. sp.
- Nipteria impunctata, panacea, valens, panopea, Peru, THIERRY-MIEG, p. 217, Le Nat. 1892; N. semigrisea, minor, erigone, Peru, tapponia, Brazil, THIERRY-MIEG, p. 236, t. c.; N. viatrix, THIERRY-MIEG, p. 262, t. c.; N. athiopissa, sororcula, tironaria, pellucenta, costistigmata, secturata, dividua, Loja, Dognin, p. 238, t. c.: n. spp.

Nychiodes lividaria, varr. described; STAUI Lep. v.

Odontopera muscularia, Central Asia, STAUI Lep. v; O. bubula, Mexico, DRUCE, p. ( Am. Heter. ii: n. spp.

Enospila, n. g., p. 5, Thalera flavifusata, Khasia Hills, p. 6; SWINHOE, Tr. E. S Omphax nænia, Mexico, DRUCE, p. 86, pl.

Heter. ii, n. sp.

Onychora, n. g. for agaritharia, Dard.; M 1892.

Ophthalmophora melæ, pl. 1, figg. 16 & 17, a 19 & 20, aspledon, fig. 22, p. 97, molpadia, Druce, Biol. Centr. Am. Heter. ii, n. s<sub>1</sub>

Ortholitha appropinquaria, p. 208, integraria, Central Asia, subvicinaria, Caucasus, Caucasus, pl. ii, fig. 17, p. 212, STAUDI n. spp.

Oulobophora, n. g. for Lobophora (O.) maced fig. 19, STAUDINGER, p. 219, Deutsche e Oxydia gastropachata, figured, pl. xliii, fig. 1

O. dognini, Peru, THIERRY-MIEG, Le Nat Mexico, p. 27, pl. xliii, fig. 11, masth Guatemala, figg. 15 & 16, artaxa, Centi angusta, Panama, figg. 17 & 18, p. 29, Di

ii: n. spp.

Palæoctenis, n. g., = (Heteropsis, Gn.); 1 1892. Panagra fixseni, Amdo, Alphéraky, p. 456,

Paralæa, n. g., Selidosemidæ, for P. promach RICK, p. 671, P. Linn. Soc. N.S.W. (2) v

Pelagia, n. subg. of Acidalia, Gumppenbel Nat. cur. lviii.

Pellonia tabidaria, Z., and sieversi, Chr., di TOPH, S. E. Z. liii, p. 245.

P. adauctata, Margelan, STAUDINGER, p. 1
P. grumaria, Amdo, Alphéraky, p. 4
n. spp.

Perigramma religiosa, P.? theodora, Peru, Ti 1892, n. sp.

Perixera pulverulenta, Khasia Hills, SWINHC fig. 8, n. sp.

Phasiane boarmiata, Amurland, GRAESEE xxxvii; P. tancrearia, Kuldja, STAUD Lep. v: n. spp.

Phorodesma smaragdaria, varr. described; S e. Z. Lep. v.



- P. smaragdularia, Margelan, STAUDINGER, p. 142, Deutsche e. Z. Lep. v. n. sp.
- Photoscotosia, Warr., is same as Trichopleura, Stgr.; T. amplicata, pl. iii, fig. 10, T. undulosa, fig. 9, figured; Alphéraky in Romanoff Mem. vi.
- Phthorarcha, n. g. for primigena, Stgr.; MEYRICK, p. 80, Tr. E. Soc. 1892.

  Phyletis prasonaria, Khasia Hills, SWINHOE, p. 12, Tr. E. Soc. 1892,
  n. sp.
- Phyllia triangularia, Blanch., = (cinerescens, Butl.); REED, p. 68, Act. Soc. Chili, i.
- Planolocha, n. g., Selidosemidæ, p. 645, for P. autoptis, n. sp., Australia, p. 646; MEYRICK, P. Linn. Soc. N.S.W. (2) vi.
- Polla avellana, Peru, THIERRY-MIEG, p. 235, Le Nat. 1892, n. sp.
- Polynesia, n. g., p. 4, type Pomasia sunandeva, Wlk., with P. truncapex, n. sp., Khasia Hills, p. 5; Swinhoe, Tr. E. Soc. 1892.
- Polythrena croceitincta, Loja, Dognin, p. 107, Le Nat. 1892, n. sp.
- Proboloptera, n. g., Selidosemida, for P. embolias, n. sp., Australia; MEY-RICK, p. 641, P. Linn. Soc. N.S.W. (2) vi.
- Pseudotagma, n. g., near Chemerina, for Eusarca interpunctaria, H.-Sch.; STAUDINGER, p. 168, Deutsche e. Z. Lep. v.
- Psilosticha, n. g., for Tephrosia mactaria, Gn.; Meyrick, p. 624, P. Linn. Soc. N.S.W. (2) vi.
- Pyrinia arxata, p. 33, pl. xliv, figg. 6 & 7, megara, p. 34, figg. 8 & 9, Panama, Druce, Biol. Centr. Am. Heter. ii, n. spp.
- Racheospila roseilinearia, pelluciduria, Ecuador, Dognin, p. 206, Le Nat. 1892; R. arycauda, pl. xlix, fig. 21, athena, fig. 22, p. 89, calliops, pl. 1, figg. 1 & 2, mustela, fig. 3, capys, fig. 4, astrœu, fig. 5, p. 90, atrapes, fig. 6, p. 91, busa, fig. 8, R. P batina, figg. 9 & 10, asmura, fig. 11, p. 92, Central America, Druce, Biol. Centr. Am. Heter. ii: n. spp.
- Rhodometra, n. g., for the Sterrha of other authors than Hubner; MEYBICK, p. 75, Tr. E. Soc. 1892.
- Rhoptria, n. subg. of Boarmia; Gumppenberg, p. 299, N. Acta Ac. L.-C. Nat. cur. lviii.
- Rhyparia hamiltonia, Khasia Hills, SWINHOE, p. 17, pl. i, fig. 5, Tr. E. Soc. 1892, n. sp.
- Rumia cratægata, variation in colour of larvæ; Gould, Tr. E. Soc. 1892, p. 216.
- Rumina lateolata, n. var. emaculuta; GRAESER, p. 317, Berl. Ent. Zeit.
- Sabulodes miliaria, Loja, Dognin, Le Nat. 1892, p. 170; S. lojanata, sticta, franciscata, xylinochroma, Dognin, p. 185, t. c.; S. ornatissima, Peru, Thierry-Mieg, p. 262, t. c. : n. spp.
- Scardamia chrysolina, Australia, MEYRICK, p. 640, P. Linn. Soc. N.S.W. (2) vi. n. sp.
- Scioglyptis, n. g., Selidosemidæ, for S. lithinopa, p. 593, hemeropa, p. 594, n. spp., Australia; MEYRICK, P. Linn. Soc. N.S.W. (2) vi.

Scodiomima, n. g. for Scodiona (S.) crncallaria, n. sp., Central Asi fig. 15; STAUDINGER, p. 202, Deutsche e. Z. Lep. v.

Scordylia valbum, Peru, THIERRY-MIEG, p. 216, Le Nat. 1892, n. 1 Scotosia senescens, N. Persia, p. 223, neglectata, Central Asia, STAUDINGER, Deutsche e. Z. Lep. v; S. confirmata, Loja, I p. 59, Le Nat. 1892: n. spp.

Selenia illunaria and lunaria, colour influenced by temperature in stage; MERRIFIELD, Tr. E. Soc. 1892, p. 34.

S. monima, pl. xlvi, figg. 11 & 12, asteria, fig. 13, myrina, fig. 14, figg. 15 & 16, p. 56, nabdalsa, fig. 21, narcæa, figg. 17 & 18, b figg. 19 & 20, azamora, figg. 22 & 23, p. 57, Mexico and Gui Druce, Biol. Centr. Am. Heter. ii, n. spp.

Selidosemida, tabulation of the Australian genera of; M

pp. 582-585, P. Linn. Soc. N.S.W. (2) vi.

Selidosema aristarcha, N. Zealand, MEYRICK, Tr. N. Z. Inst. xi cheleuta, p. 598, amphiclina, p. 599, eremias, thermea, p. 600, c p. 605, leptodesma, p. 606, leucoplecta, p. 607, agoraea, p. 608, a p. 610, adelphodes, p. 611, sascia, p. 613, argoplaca, p. 615, At MEYRICK, P. Linn. Soc. N.S.W. (2) vi, n. spp.

Sicya aurunca, pl. xlv, fig. 7, mesapia, fig. 8, bala, fig. 9, meda, figg. 1 p. 46, myron, figg. 12 & 13, S.? ascolia, fig. 14, p 47, Central A Drucce, Biol. Central Am. Heter, ii p. spp.

DRCCE, Biol. Centr. Am. Heter. ii, n. spp.

Spargania longipalpata, Peru, THIERRY-MIEG, p. 262, Le Nat n. sp.

Stathmorrhopa, n. g. for Chlenias beggaria, Gn.; MEYRICK, p. Linn. Soc. N.S.W. (2) vi.

Stegania crina, Khasia Hills, SWINHOE, p. 14, pl. i, fig. 14, Tr. 1892, n. sp.

Sterrha virgenpamba, Loja, Dognin, p. 237, Le Nat. 1892, n. sp.

Stibaroma, n. g., Selidosemidæ, for S. melanotoxa, n. sp., Au Meyrick, p. 669, P. Linn. Soc. N.S.W. (2) vi.

Stibaroma trigramma, S. Australia, LOWER, p. 9, Tr. R. Soc. S. Au n. sp.

Stigma isthmensis, Panama, THIERRY-MIEG, p. 236, Le Nat, 1892, 1 Streptopteron, n. g. (Ephyrides), p. 9, for S. posticamplum, n. sp., Hills, p. 10, pl. i, fig. 12; SWINHOE, Tr. E. Soc. 1892.

Strophidia falcifera, E. Africa, WEYMER, p. 118, S. E. Z. liii, n. sp. Syllexis extendata, chartularia, Loja, DOGNIN, p. 144, Le Nat. 189; Synchlora? nortia, Mexico, DRUCE, p. 93, pl. l, fig. 12, Biol. Cen Heter. ii, n. sp.

Synegiodes, n. g., type Anisodes diffusaria, Moore, with S. diffus p. 11, pl. i, fig. 9, histrionaria, p. 12, n. spp., Khasia Hills; Sw Tr. E. Soc. 1892.

Syngonorthus, n. g., Urapterygidæ, for S. subpunctatus, n. sp., Borned fig 9; Butler, p. 130, P. Z. S. 1892.

Tachyphyle? neis, Central America, DRUCE, p. 93, pl. l, figs. 13
Biol. Centr. Am. Heter. ii, n. spp.

- \*ephronia oranaria, Algeria, p. 179, castilliaria, Spain, p. 180, pl. ii, fig. 11, STAUDINGER, Deutsche e. Z. Lep. v, n. spp.
- "ephrosia bostar, p. 76, pl. xlviii, fig. 13, bura, fig. 14, T.? azenia, fig. 15, balsa, fig. 16, p. 77, T. muræna, figg. 17 & 18, axica, fig. 19, p. 78, Central America, DRUCE, Biol. Centr. Am. Heter. ii, n. spp.
- "erpnomicta lala, Khasia Hills, SWINHOE, p. 13, Tr. E. Soc. 1892, n. sp. 'etracis azonax, pl. xlvi, fig. 8, belides, fig. 9, Central America, DRUCE, p. 54, Biol. Centr. Am. Heter. ii, n. spp.
- 'halassodes liliana, Khasia Hills, SWINHOE, p. 7, pl. i, fig. 2, Tr. E. Soc. 1892, n. sp.
- \*\*Thalera acte, Khasia Hills, SWINHOE, p. 6, pl. i, fig. 13, Tr. E. Soc. 1892; T. spumosaria, eximia, Peru, Dognin, p. 237, Le Nat. 1892: n. spp. \*\*Thamnonoma wavaria, aberration; VRIES & SNELLEN, Tijdschr. Ent. xxxv, pp. 24 & 25.
- "herina azion, pl. xlv, figg. 25 & 26, betala, fig. 27, munda, fig. 28, p. 50, bada, pl. xlvi, figg. 1 & 2, T. ? nomia, fig. 4, p. 51, Central America, DRUCE, Biol. Centr. Am. Heter. ii, n. spp.
- Posaura, n. g., p. 15, type Metabrazas falcipennis, Moore, with T. picaria, n. sp., Khasia Hills, p. 16, pl. i, fig. 16; SWINHOE, Tr. E. Soc. 1892.
   Prichoclada opsinaria, Khasia Hills, SWINHOE, p. 15, Tr. E. Soc. 1892, n. sp.
   Prichopleura leechi, Amdo, Alpheraky, p. 458, Hor. Ent. Ross. xxvi, n. sp.
   Priphosia ravulata, Central Asia, STAUDINGER, p. 221, Deutsche e. Z. Lep. v, n. sp.
- Prygodes musivaria, figured, pl. li, figg. 6 & 7, Biol. Centr. Am. Heter. ii.
  P. niobe, pl. li, figg. 9 & 10, merta, fig. 11, p. 106, ategua, fig. 12, melia, fig. 13, callas, fig. 14, carbania, fig. 15, p. 107, T. ? auxesia, figg. 16 & 17, p. 108, Central America, DRUCE, Biol. Centr. Am. Heter. ii, n. spp.
- Trapteryz hilaris, Peru, THIERRY-MIEG, p. 217, Le Nat. 1892, n. sp.
- Kanthorhos paradelpha, p. 11, hyperythra, p. 12, xanthospila, p. 13, S. Australia, Lower, Tr. B. Soc. S. Austr. xv, n. spp.
- Kenomusa metallica, Brisbane, Lucas, p. 299, P. Linn. Soc. N.S.W. (2)
- Zamacra, n. g. for flabellaria, Heeg.; MEYRICK, p. 121, Tr. E. Soc. 1892. Zonosoma albilineata, Central Asia, Staudinger, p. 159, Deutsche e. Z. Lep. v; Z. alodia, Loja, Dognin, p. 16, Le Nat. 1892: n. spp.

# Pyralidæ (including Cranbidæ, Phycitidæ, Siculodidæ).

- Cf. Cockerell (177), Distant (216), Fernald (280), Hulst (441), Kellogg (479), Lucas (568, 569), Müller (629), Pagenstecher (667, 668, 669), Ragonot (712), Rebel (715), Röber (767), Snellen (872, 873), Staudinger (890, 891), Swinhoe (909), Warren (983).]
- 4charana rudis, Japan, minoralis, Accra, maledicta, Pitcairn I., p. 435, descripta, S. Paolo, simplex, Bombay, olivescens, Ecuador, indistincta, Japan, p. 436, elongalis, Formosa, fuscescens, Japan, p. 437, WARREN, Ann. N. H. (6) ix, n. spp.

i

- Acrobasis cirreferella, Texas, Hulst, p. 60, Canad. Ent. xxiv, n. sp. Actenicides, n. g., type Nephopteryx creperalis, Swinhoe; WARREN, 1
  Ann. N. H. (6) ix.
- Adeloides, n. g., type Hoterodes cinerealis, Moore; WARREN, p 299, N. H. (6) ix.
- Aglaope, n. g., type Mecyna furnacalis, Gn.; Warren, p. 298, Ann. (6) ix.
- Agrammia cervinalis, lutealis, S. Paolo, WARREN, p. 439, Ann. N. ix, n. spp.
- Anomostictis, n. n. (to replace Diastictis, Hb. Verz., p. 355), wi coccelis, n. sp., California; Warren, p. 434, Ann. N. H. (6) ix
- Anthocrypta, n. g. for Ebulea subinquinalis, Gn.; WARBEN, p. 296 N. H. (6) ix.
- Aphomia? progritella, Canary Is., REBEL, p. 262, Ann. Hofmuseum vii, n. sp.
- Aphytoceros nigrolinealis, Goya, p. 390, longipalpis, Transvaal, p. WARREN, Ann. N. H. (6) ix, n. spp.
- Aplographe, n. g., type Scopula bisignata, Butl, with A. fulvalis, Good Hope, umbrosalis, N. China, n. spp.; Warren, p. 301, A. H. (6) ix.
- Aporodes obscura, Sarepta, versicolor, Washington Terr.; WARREN, Ann. N. H. (6) ix, n. spp.
- Archernis pubescens, E. Ania, WARREN, p. 300, Ann. N. H. (6) ix, n. Autocosmia, n. g., near Cosmocreon, p. 432, for A. concinna, n. sp., America, WARREN, p. 433, Ann. N. H. (6) ix.
- Basotarcha eunealis, Port Darwin, WARREN, p. 396, Ann. N. H. (6) ix Blepharucha, n. g., type Phalana naide, Stoll.; WARREN, p. 177, A H. (6) ix.
- Botys commellalis, early stages; Christien, Bull. Soc. Ent. Fr. lxi, p
  B. gracesri, Siber.a, Staudinger, p. 384, pl. iii, fig. 24, Deutsche e. 2
  v: B. catasemalis, Ké I., Röber, p. 333, Tijdschr. Ent. xxxiv (
  xxxv): n. spp.
- Bradyrrhoa ochrospilella, Canary Is., REBEL, p. 258, pl. xvii, fig. 13 Hofmuseum Wien, vii, n. sp.
- Camptomastiz, n. g., type Botys pacalis, Leech; WARREN, p. 439, N. H. (6) ix.
- Cataclysta lemna, respiration of larva; MÜLLER, Zool. Jahrb. Sy p. 623: Cataclysta spp.? subaquatic oviposition; MÜLLES p. 625.
- C. hapilista, Khasia Hills, Swinhor, p. 20, pl. i, fig. 11, Tr. F 1892, n. sp.
- Cephis, n. g., Crambina, p. 295, for C. galleriellus, n. sp., Tunis, pl. iii, fig. 12; RAGONOT, Deutsche e. Z. Lep. v.
- Chilo infuscatellus, Java, Snellen, p. 347, Tijdschr. Ent.:
- Chipeta, n. g., near Tampa, for C. perlepidella, n. sp., Florida; F p. 62, Canad. Ent. xxiv.

- Choristostigma, n. g., type Botys plumbosignata, Fern., with O. elegantalis, n. sp., California; WARREN, Ann. N. H. (6) ix.
- Cirrochrista margarita, S. Paolo, WARREN, p. 430, Ann. N. H. (6) ix, n. sp.
- Chephidia, n. g., near Selagia, for C. kenteriella, n. sp., Siberia; RAGONOT, p. ccxxxv, Bull. Soc. Eut. Fr. lxi.
- Conogethes nubifera, Brisbane, Lucas, p. 264, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Coptobasis ridopalis, Khasia Hills, SWINHOE, p. 18, Tr. E. Soc. 1892, n. sp. Cornifrons pulveralis, N. America, Warren, p. 435, Ann. N. H. (6) ix, n. sp.
- Cosmocreon, n. g., type B. albiceralis, Grote; Warren, p. 433, Ann. N. H. (6) ix.
- Crambus atlanticus, n. var. canariensis; REBEL, p. 254, pl. xvii, figg. 11 & 12, Ann. Hofmuseum Wien, vii.
- Cremnophila, n. g., near Adelosemia, for C. auranticiliella, n. sp., Siberia; RAGONOT, p. ccxxxvi, Bull. Soc. Ent. Fr. lxi.
- Cybolomia extorris, N. America, Warren, p. 395, Ann. N. H. (6) ix, n. sp. Diacme, n. g., type Samea phyllisalis, Wlk.; Warren, p. 389, Ann. N. H. (6) ix,
- Diatron saccharalis, notes on in Jamaica, with metamorphoses; Cock-ERELL (177).
- D. striatalis, Java, Snellen, p. 349, pl. xix, figs. 1-4, Tijdschr. Ent. xxxiv. n. sp.
- Dichotis, n. g., type Botys teneralis, Led.; WARREN, p. 392, Ann. N. H. (6) ix.
- Didymostoma, n. g., type Botys emphranoralis, Wlk.; WARBEN, p. 392, Ann. N. H. (6) ix.
- Dioryctria nivaliensis (Rag. i. l.), Canary Is., REBEL, p. 256, pl. xvii, fig. 10, Ann. Hofmuseum Wien, vii, n. sp.
- Diviana nymphæella, Florida, Hulst, p. 62, Canad. Ent. xxiv, n. sp.
- Durdara ovifera, Borneo, Butler, p. 129, pl. vi, fig. 7, P. Z. S. 1892, n. sp. Ebulea fumipennis, p. 392, E.? straminea, California, pulverulenta,
- Dharmsala, p. 393, Warren, Ann. N. H. (6) ix, n. spp.
- Ematheudes tunesiella, Tunis, RAGONOT, p. 298, Deutsche e. Z. Lep. v, n. sp.
- Endotricha rogenhofers, Canary Is., Rebel, p. 249, pl. xvii, fig. 1, Ann. Hofmuseum Wien, vii; E. crobulus, p. 305, dispergens, p. 306, Queensland, Lucas, P. Linn. Soc. N.S.W. (2) vi: n. spp.
- Ennychia intrudens, New Zealand, WARREN, p. 175, Ann. N. H. (6) ix, n. sp.
- Ephestia sericaria, Scott, queried as identical with kuhniella; RAGONOT, p. oclxxiv, Bull. Soc. Ent. Fr. lxi.
- Euclasta warreni, Transvaal, Distant, Naturalist Transvaal, p. 241, pl. ii, fig. 5, n. sp.
- Euctenospila, n. g., near Blepharucha, for E. castalis, n. sp., Abyssinia; WARREN, Ann. N. H. (6) ix.

- Energestis consimilis (no locality), WARREN, p. 433, Ann. N. H. (6) ix Energhers concolorella, Tunia, RAGONOT, p. 297, Deutsche e. Z. I n. sp.
- Glauconer, n. g., type Botys deductalis, Wlk., with G. subj. Madagascar, fuscessess, Sumatra, n. spp.; WARREN, pp. 296
  Ann. N. H. (6) ix.
- Glyphodes prothymedia, Khasia Hills, SWINHOR, p. 19, pl. i, fig. 15, Soc. 1892, n. sp.
- Hemiscopis cineres, Japan, expense, Dharmsala, WARREN, p. 396 N. H. (6) ix, n. app.
- Homeosoma caneriella, Canary Ia., REBEL, p. 260, Ann. Hofm Wien, vii; H. delinesta, Brisbane, Lucas, p. 266, P. Lim N.S.W. (2) vii: n. spp.
- Hyaloplagia, n. g., type Hydrocampa pulchralis, Moore; WARREN, Ann. N. H. (6) ix.
- Hyslorista, n. g., type Rhederia teniolalis, Gn., with H. imitans, S. Paolo: WARREN, Ann. N. H. (6) ix, p. 299.
- Hydrocamps symphesis, structure of larva, skin, breathing, habi discussed; MCLLER, Zool, Jahrb. Syst. vi, pp. 617-622.
- Hypotomorpha, n. g., near Ancylolomia, p. 252, for H. lance n. sp., Canary Ia., p. 253, pl. xvii, figs. 4 & 5; Rebell, Ann museum Wien, vii.
- Lipographis subosseella, Bahamaa, Hulst, p. 62, Canad. Ent. xxiv, Larocreon, n. g., type Salbis continustalis, Wilng.; Warren, p. 431 N. H. (6) ix.
- Loxestoge baccestalis, Texas, HULST, p. 63, Canad. Ent. xxiv, n. sp. Melitera dentata, metamorphoses; Kelloge, Kanasa Quart. i, pl. viii.
- Mesographe juncialis. Japan, Warren, p. 434, Ann. N. H. (6) ix, r Mesothyris, n. g., type Botys aluensis, Butl.; Warren, p. 300 N. H. (6) ix.
- Metasia excavatalis, Tunis, RAGONOT, p. 294, pl. iii, fig. 14, Deutsel Lep. v, n. sp.
- Micractis, n. g., type Botys subilalis, Hb., with M. sanguinealis, Japan: WARREN, p. 294, Ann. N. H. (6) ix.
- Mimochinia, n. g., type Botys thalialis, Wlk.; WARREN, p. 174 N. H. (6) ix.
- Minudea, n. g., p. 440, for M. olivelis, subrosea, S. Paolo, flavinotai Janeiro, n. spp., p. 441; WARREN, Ann. N. H. (6) ix.
- Monocona, n. g., near Tegostoma, p. 173, for M. rubralis, n. sp., Calif WARREN, Ann. N. H. (6) ix.
- Myelois sarcotineta, Brisbane, LUCAS, p. 265, P. Linn. Soc. N.S. vii; M. fructetella, elegantella, p. 59, texanella, leucophæella, p. America, HULST, Canad. Ent. xxiv: n. spp.
- Nuscia citrinalis, Dharmmla, WARREN, p. 302, Ann. N. H. (6) ix, t Niphograpta, n. g., type Epichronistis albiguttalis, War.; WARREN, Ann. N. H. (6) ix.

- Noctuelia flavifimbrialis, California, WARREN, p. 174, Ann. N. H. (6) ix, n. sp.
- Notarcha exculta, Brisbane, Lucas, p. 262, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Notaspis, n. g., type Botys tranquillalis, Led.; WARREN, p. 297, Ann. N. H. (6) ix.
- Nymphula sinuosa, Brisbane, Lucas, p. 261, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Ocala, n. g., near Dolichorrinia, for O. dryadella, n. sp., Florida; HULST, p. 61, Canad. Ent. xxiv.
- Œdematophora cacaalis, Brisbane, Lucas, p. 306, P. Linn. Soc. N.S.W. (2) vi, n. sp.
- Opeibotys latipennis, ocellalis, Japan, Warren, p. 295, Ann. N. H. (6) ix, n. spp.
- Pachynoa limitata, Borneo, WARREN, p. 395, Ann. N. H. (6) ix, n. sp.
- Pachyzancia dissimilis, Acora, WARREN, p. 442, Ann. N. H. (6) ix, n. sp.
- Paralipsa decorella, N. America, HULST, p. 63, Canad. Ent. xxiv, n. sp.
- Paraponyx, sp. (Brazil), respiration of larva; Müller, Zool. Jahrb. Syst. vi, p. 625.
- Pessocosma suffusalis, Goya, WARREN, Ann. N. H. (6) ix, p. 390, n. sp.
- Pharambara sphoraria, Khasia Hills, SWINHOE, p. 18, pl. i, fig. 17, Tr. E. Soc. 1892; P. vinosa, Borneo, Butler, p. 130, pl. vi, fig. 8, P. Z. S. 1892: n. spp.
- Phlyctenia ferruginealis, paolinalis, arenacea, Rio de Janeiro, WARREN, p. 431, Ann. N. H. (6) ix, n. spp.
- Pindicitora flavifrons, Accra, WARREN, p. 172, Ann. N. H. (6) ix, n. sp.
- Prionopaltis, n. g., p. 437, for P. sericea, Dharmsala, consocia, P. ? suffusalis, Japan, p. 438, n. spp.; WARREN, Ann. N. H. (6) ix.
- Prionophora torquesauria, Brisbane, LUCAS, p. 255, P. Linn. Soc. N.S.W. (2) vii, n. sp.
- Prodasycnemis, n. g., type Botys inornata, Butl.; WARREN, p. 301, Ann. N. H. (6) ix.
- Pygospila tyres, n. var., cuprealis; SWINHOE, p. 19, pl. i, fig. 4, Tr. E. Soc. 1892.
- Pyrausta dorsipunctalis, Canary Ia., Rebel, p. 245, Ann. Hofmuseum Wien, vii; P. straminea, violacea, Queensland, Lucas, p. 263, P. Linn. Soc. N.S.W. (2) vii; P. coccinea, California, Warren, p. 176, Ann. N. H. (6) ix; P. magdalena, Florida, Fernald, p. 122, Canad. Ent. xxiv; P. nigralis, Florida, Fernald, p. 178, t. c.: n. spp.
- Ramila angustifimbrialis, Burma, WARREN, p. 430, Ann. N. H. (6) ix, n. sp. Rhectocraspeda, n. g., type Botys periusalis, Wlk.; WARREN, p. 439, Ann. N. H. (6) ix.
- Rhodoneura notula, hab. ?, R. ? tessellatula, Philippines, PAGENSTECHER, p. 58, Deutsche e. Z. Lep. ▼ (for other n. spp. of Rhodoneura see Bombyces, p. 246): n. spp.
- Salebria levigatella, purpurella, N. America, Hulst, p. 61, Canad. Ent. xxiv, n. spp.

Scirpophaga intecta, Java, Shellen, p. 343, pl. xviii, figu 1-4, Ent. xxxiv, n. sp.

Semaiomima n. g. for Hypotrita flavicepe, Burm.; WARREN, p. 1 N. H. (6) ix.

Sericophera guttata, figured, Romanoff Mem. vi, pl. xiv, fig. 14. Sericoplaga n. g., near Opsibotys, p. 295, for S. externalis, n. sp. 296; Warren, Ann. N. H. (6) ix.

Sicul[od]ida: monographic revision of the group; Pagen Deutsche e. Z. Lep. v, pp. 5-131, and Supplement to the id. t. c. pp. 443-449.

Seculodes (?) ignotalis, Bonerate, Röber, p. 329, Tijdschr. E. (fig. in xxxv); S. pulchelloides, Peru, p. 64, pl. i, fig. 4, Chiriqui, p. 65, violalis, Ecuador, arcuata, Chiriqui, p. 69, Peru, p. 70, furcatula, Cameroons, p. 71, pl. i, fig. 6, er Brazil, pl. i, fig. 17, equamigera, p. 72, catenula, p. 7. anastomosalis, Chiriqui, albipunctula, Peru, p. 76, tristriate video, p. 77, levigata, p. 78, ferruginea, p. 79, S. (Z. leuconotula, p. 80, pl. i, fig. 15, fumatilis, p. 81, Chiriqui, p. 82, selenioides, p. 83, pl. i, fig. 14, Peru, maculata, p. 91, st p. 92, patercula, p. 94, Brazil, fratercula, California, p. 95, Peru, p. 96, pl. i, fig. 10, denticulata, Chiriqui, p. 98, pl. politula, Philippines, p. 101, rotundula, Amasons, p. 107, pl. polychloralis, Chiriqui, p. 108, nigropunctula, Natal, p. 109 punetula, Sumatra, p. 110, flavula, Gaboon, p. 111, guttula, 1 pl. i, fig. 11, p. 112, rosacea, Malacca, p. 119, pl. i, fig. 3, Borneo, p. 121, triangularis, Philippines, p. 122, globulifere PAGENSTECHER, Deutsche e. Z. Lep. v : n. spp.

Spilodes bicoloralis, Bagdad, WARREN, p. 178, Ann. N. H. (6) ix, Stenochora, n. g., type Rhodaria lancinalis, Gn.; WARREN, p. 2: N. H. (6) ix.

Stemomeles, n. g., type Botys agavealis, Wlk.; WARREN, p. 437, H. (6) ix.

Striglina radiata, Portorico, p. 41, sordida, Borneo, p. 47, PAGENi Deutsche e. Z. Lep. v, n. spp.

Syllythria? rubrivena, Madagascar, p. 176, subnicalis, Californis Warren, Ann. N. H. (6) ix, n. spp.

Tanaophysa, n. g. for T. adornatalis, n. sp., S. Paolo; WARREN Ann. N. H. (6) ix.

Terastiodes, n. g., type Terastia ochraceulis, Wlk.; WARREN, p. 2 N. H. (6) ix,

Tetraphana alipes, Amazons, p. 3, lilacina, Bolivia, p. 4, pl. i, fig p. 131 said to be Euclea fenestrata, Mén.); PAGENSTECHER, le. Z. Lep. v, n. spp.

Tholoria stigmosalis, S. Paolo, WARREN, p. 429, Ann. N. H. n. sp.

Trichauchenia, n. g., type Euclita dharmsalæ, Butl.; WARREN, Ann. N. H. (6) ix.

- Trischistognatha, n. g., type Spilodes palindialis, Gn.; WARREN, p. 429, Ann. N. H. (6) ix.
- Tritæa protealis, Callao, ferruginea, Ooquimbo, WARREN, p. 179, Ann. N. H. (6) ix, n. spp.
- Udea sabulosalis, Coquimbo, p. 393, indistinctalis, N. America, inhospitalis, California, nigripunctata, Callao, p. 394, WARREN, Ann. N. H. (6) ix, n. spp.
- Varnia aurea, Chiriqui, p. 32, pl. i, fig. 13, incudigera, p. 33, ypsiloides, p. 35, Java, Pagenstecher, Deutsche e. Z. Lep. v, n. spp.
- Zophodia bella, Massachusetts, HULST, p. 61, Canad. Ent. xxiv, n. sp.
- Zeuzerodes, n. subg. of Siculodes, q.v.; PAGENSTECHER, p. 80, Deutsche e. Z. Lep. v.

### TORTRICIDÆ.

- [G. Chretien (174), Constant (182), Fernald (280), Meyrick (605), Rebel (715), Riley (748, 749), Snellen (872), Staudinger (891).]
- Bactra lanceolana, = (Chiloides straminea, Butl.); MEYRICK, p. 217, Tr. N. Z. Inst. xxiv.
- Caccecia semiferana, egg-clusters noticed; GILLETTE, Canad. Ent. xxiv, p. 36. C. rosana and parallela, full-grown larvæ; BEUTENMÜLLER, p. 80, Bull. Am. Mus. iv.
- C. magnoliana, New York, FERNALD, p. 121, Canad. Ent. xxiv, n. sp.
- Carpocapea saltitass, determination of the plant; RILEY, P. E. Soc. Wash. ii, pp. 178 & 213.
- Ditula woodiana, habits; Wood, Ent. M. M. xxviii, p. 225.
- Doloploca buratica, Siberia, STAUDINGER, p. 388, Deutsche e. Z. Lep. ▼, n. sp.
- Eupacilia geyeriana and larva figured; RICHARDSON, P. Dorset Club, ziii, pl.
- Grapholitha carduana, n. var. subvittana; STAUDINGER, p. 299, Deutsche e. Z. Lep. v.
- G. kenteana, Siberia, STAUDINGER, p. 390, Deutsche e. Z. Lep. v; G. schistaceana, Java, SNELLEN, p. 353, pl. xix, figs. 5-7, Tijdschr. Ent. xxxiv; G. sebastiana, N. Mexico, RILEY, p. 214, P. E. Soc. Wash. ii: n. spp.
- Noteraula, n. g. (Grapholithidæ), for Chiloides straminea, Meyr., nec Butler; Meyrick, p. 217, Tr. N. Z. Inst. xxiv.
- Pandemis simonyi, Canary Is., REBEL, p. 263, pl. xvii, figs. 8 & 9, Ann. Hofmuseum Wien, vii, n. sp.
- Penthina oblongana, n. var. adelana; REBEL, p. 526, Verh. s.-b. Wien, zlii.
- Pyrgotis plinthoglypta, New Zealand, MEYRICK, p. 218, Tr. N. Z. Inst. xxiv, n. sp.
- Sciaphila cinareana, Carcassonne, Chrétien, Le Nat. 1892, p. 132, n. sp. Steganoptycha lindana, Ontario, Fernald, p. 178, Canad. Ent. xxiv, n. sp.

Teras ferrugana, metamorphoses, CHRÉTIEN, p. 240, Le Nat. 1892, T. comandrana. N. America, FERNALD, p. 121, Canad. Ent. xxiv angusana, N. America, FERNALD, p. 178, t. c. : n. spp.

### TIMBIDE.

[Cf. Chapman (164), Constant (182), Durrant (210), Ebschoff Fernald (280), Lower (565), Meyrick (605), Ragonot Rebel (715, 716), Riley (741, 742, 743), Staudinger (891), (1014, 1015).]

Wing-neuration discussed; SPULER, Z. wiss. Zool. liii, pp. 626-635 Tineid larva mining leaves of *Populus fremonti* described; Town Zoe, iii, pp. 234-236.

Adela infantella, figure, pl. xvi, fig. 3, Romanoff Mem. vi. A. vir oviposition of; Chapman, Ent. M. M. xxviii, p. 42.

Blastobasis mamorosella, Woll., characters discussed; Rebel, ppl. xvii, fig. 6, Ann. Hofmuseum Wien, vii.

Bucculatrix canadensisella, metamorphoses; PACKARD, Ins. Life, v, fig.

Butalis sagittatella, figure, pl. xvi, fig. 12, Romanoff Mem. vi. Cerostoma falculella, figure, pl. xvi, fig. 4, Romanoff Mem. vi.

C. affinitella, Siberia, STAUDINGER, p. 392, Deutsche e. Z. Lep. v, n. Choreutis solaris, figure, Romanoff Mem. vi, pl. xvi, fig. 1.

Coleophora lativittella, figure, pl. xvi, fig. 11, Romanoff Mem. vi.

C. sylvaticella, p. 118, glaucicolella, p. 172, England, Wood, Ent. 1 xxviii; C. agrammella, England, Wood, t. c. p. 283; C. comi Canary Ia, Rebel, p. 278, pl. xvii, fig. 15, Ann. Hofmuseum vii : n. spp.

Cryptophaga blackburnii, ochroleuca, S. Australia, Lower, p. 15, 7 Soc. S. Austr. xv, n. spp.

Depressaria multiplicella, pl. xvi, fig. 5, anticella, fig. 6, figured, Ron Mem. vi. D. doronicella, metamorphoses; Constant, Bull. Autun, iv, pp. 277-283.

D. conciliatella, Canary Is., REBEL, p. 272, pl. xvii, fig. 14, Ann. museum Wien, vii, n. sp.

Erechthias erebistis, New Zealand, MEYRICK, p. 220, Tr. N. Z. Inst. n. sp.

Etiella zinckenella, Tr., = (madagascariensis, Saalm., and chrysopo Meyr.); Snellen, p. 84, Tijdschr. Ent. xxxv.

Gelechia unthyllidella, = (sparsiciliella, Barrett); BARRETT, Ent. I xxviii, p. 80. G. gallædiplopappi, larva noticed; Fyles, I Canad. Ent. xxiv. G. cinctipunctella, pl. xvi, fig. 7, G. (Lita): notephrella, fig. 8, G. (Teleia) trijugella, fig. 9, figured, Rom Mem. vi.

Glyphipteryx magnatella, figure, pl. xvi, fig. 10, Romanoff Mem. vi. Hyponomeuta gigas, Canary Is.; REBEL, p. 271, pl. xvii, fig. 17, Hofmuseum Wien, vii, n. sp.

- Lampronia capitella, life-history; CHAPMAN, Ent. M. M. xxviii, p. 297.
- Lichenaula selenophora, S. Australia, LOWER, p. 16, Tr. R. Soc. S. Austr. xv, n. sp.
- Lita solanella, metamorphoses figured, Ins. Life, iv, p. 239.
- Nemophora dorsiguttella, figure, pl. xvi, fig. 2, Romanoff Mem. vi.
- Palparia leucosta, S. Australia, Lower, p. 10, Tr. R. Soc. S. Austr. xv, n. sp.
- Phylomictis monochroma, S. Australia, Lower, p. 17, Tr. R. Soc. S. Austr. xv, n. sp.
- Prodoxus: variation, habits, &c.; RILEY, P. Biol. Soc. Washington, vii, pp. 96-99.
- P. pulverulentus, p. 315, y-inversus, p. 316, reticulatus, p. 317, coloradensis, p. 318, sordidus, p. 319, N. America, RILEY, P. E. Soc. Wash. ii, n. spp.
- Prosuba, and fertilisation of Yucca; RILEY, P. Biol. Soc. Washington, vii, pp. 81, &c., woodcuts. P. (and Prodoxus), some anatomical peculiarities; RILEY, P. E. Soc. Wash. ii, pp. 305-311: maxillary tentacles; SMITH, P. Am. Ass. xli, p. 198: problematical larva; TOWNSEND, Zoe, iii, p. 115.
- Pronuba synthetica, Mojave Desert, RILEY, P. E. Soc. Wash. ii, p. 312, metamorphoses, with figg. : n. sp.
- Psecadia libyella, Tunis, RAGONOT, p. lxxxii, Bull. Soc. Ent. Fr. lxi, n.sp. Ptocheusa multistrigella, Tunis, RAGONOT, p. lxxxii, Bull. Soc. Ent. Fr. lxi, n. sp.
- Sematocera, n. g., near Lasioctena, p. 242, for S. fuliginipuncta, n. sp., Transvaal, p. 243, pl. iv, fig. 4; DURRANT in Distant, Naturalist Transvaal.
- Semiocosma caustopa, paraneura, New Zealand, MEYRICK, p. 219, Tr. N. Z. Inst. xxiv, n. spp.
- Setomorpha discipunctella, Canary Is., Rebell, p. 267, pl. xvii, fig. 16, Ann. Hofmuseum Wien, vii, n. sp.
- Staintonia fulgens, figured, pl. xvi, fig. 13, generic position queried, p. 671, Bomanoff Mem. vi.
- Swammerdamia apicella, larva described; ELISHA, Ent. M. M. xxviii, p. 257.
- Tinea merdella, n. var.? immaculatella, REBEL, p. 269, Ann. Hofmuseum Wien, vii, n. sp.; T. propulsatella, Tyrol, REBEL, p. 527, Verh. s.-b. Wien, xlii; T. bipartitella, Tunis, RAGONOT, p. lxxxii, Bull. Soc. Ent. Fr. lxi: n. spp.
- Tineola alutella, Canary Is., REBEL, p. 270, pl. xvii, fig. 3, Ann. Hof-museum Wien, vii, n. spp.
- Tineomima, n. g. for Tineola? (T.) kenteella, n. sp., Siberia; STAUDINGER, p. 391, Deutsche e. Z. Lep. v.
- Trachypepla hieropis, New Zealand, MEYRICK, p. 218, Tr. N. Z. Inst. xxiv, n. sp.
- Xylorycta leucophanes, S. Australia, Lower, p. 17, Tr. R. Soc. S. Austr. xv, n. sp.

## Agdistidz,

Cf. Herisc (401), Mabille & Vi Acipolis timeta, Tunia, Staudings Attroptorys doori, figured; Man Lep.

Gilbertia eques, figured; MASILLE Minascoptiles pinaredactylus, figur Ochyretics fuscista, figured; Mai Lep.

Platypolia terminalia, figure, pl.: ochrodactyla, dichrodactyla, opinious as to distinctions of;

Sockehore donatella, figured; Mai Lep.

Stepenedactyla concursa, pl. xii, fig. & VUILLOY, Nov. Lop.

Uroloba fuscicostatu, figured; Maing. 8.

Ctace ocirucealis, figured; MABILI



# (D.) DIPTERA.

[Cf. Aldrich (5, 6), Arribalzaga (15), Becker (45), Bergroth (61), Bezzi (79, 80), Bidenkap (81), Bigot (83-87), Blanchard (101, 102), Brauer (116, 117, 118), Brauer & Bergenstamm (119), Cholodkovsky (171), Cockebell (180), Coquillett (185-190), Giglio-Tos (339-343), Guthrie (381), Henneguy & Binet (399), Hudson (440), Jenkinson (456), Kieffer (492-495), Kowarz (512), Lampa (537, 538), Liebel (562), Meade (595, 596, 597), Meunier (602), Mik (606-610), Mortensen (627), Muggenburg (628), Osten-Sacken (659-663), Portschinsky (698, 699), Biley (753), Reed (718), Röder (769, 770, 771), Rübsaamen (784-787), Sangagli (791), Sawtschenko (796), Snow (877), Stein (895, 896), Strobl (902-906), Theobald (919), Thomas (924, 925), Tömösváry (929), Townsend (931-944, 946, 947), Trybom (951), Verrall (971), Webster (990), Williston (347, 1010, 1011), Wulp (1019).]

Table of the N. American genera of Nemocerous Diptera; Townsend, Tr. Am. Ent. Soc. xix, pp. 144-160.

Catalogue of Diptera of the Oriental Region; BIGOT (84).

A new spinning Hilara, cf. H. aëronetha, infrà.

### CECIDOMYIIDÆ.

Structure and mode of preparation of larvæ; Thomas, pp. 356-358, Verh. z.-b. Wien, xlii.

General notes on the structure of larvee of Cecidomyiidæ; Rübsaamen, Berl. Ent. Zeit. xxxvi, pp. 381-391, pl. xiv.

New Cecidomyiid galls; BÜBBAAMEN, pp. 403-406, Berl. Ent. Zeit. xxxvi. Account of various Cecidomyiid galls; Thomas (925).

Mode of preparing for collections, classification; RÜBSAAMEN, Berl. Ent. Zeit, xxxvii, pp. 319-341: Amended descriptions of many species; id. t. c. pp. 341-404.

Asphondylia, notes on the characters and species, KIEFFER, pp. 220-222, Wien. ent. Z. xi; A. sarothamni, Lw., and bitensis, Kieff., noticed, Mik, p. 184, Wien. ent. Z. xi; A. cytisi, sarothamni, genista, notes on, Rübsaamen, pp. 59-61, Verh. z.-b. Wien. zlii.

Asynapta citrina, description amplified; Kieffer, p. 264, Berl. Ent. Zeit. xxxvi.

- Cecidomyia rosaria, descriptive note, p. 246, salicina, Schrank, syr p. 247, saliciperda, Duf., descriptive note, p. 249, KIEFFEI Ent. Zeit. xxxvi; C. viciæ gall, noticed and figured, MIK, Wien. ent. Z. xi, pl. iii, figg. 1-3: sp. ? gall on Thalictrum fig. 4.
- C. praticola, Bitsch, Kieffer, p. 216, Wien. ent. Z. xi; C. op. 241, pulvini, p. 244, karschi, p. 251, medullaris, p. 253 p. 255, Bitsch, Kieffer, Berl. Ent. Zeit. xxxvi; C. sa. Bitsch, Kieffer, p. 262, & c.; C. dulcamaræ, p. 397, kleini, symphyti, p. 399, tetensi, p. 400, gemmæ, p. 401, Germany, Rüßl. c.: n. spp.
- Choristoneura, n. g., for obtusa, Lw.; Rübsaamen, p. 342, Bei Zeit. xxxvii.
- Clisorrhyncha, characters noticed; KIEFFER, p. 223, Wien. ent. Z Colomyia, n. g. near Epidosis, for C. clavata, n. sp., Bitsch, K. p. 259, Berl. Ent. Zeit. xxxvi.
- Cystiphora, n. g. for a part of Cocidomyia, p. 213, including C. p. n. sp., Bitech, p. 213, KIEFFER, Wien ent. Z. xi.
- Dichelomyia, n. g. for a part of Cocidomyia; Rübsaamen, p. 34/ Ent. Zeit. xxxvii.
- Diplosis tritici, ravages in Sweden; LAMPA (536).
- D. loescii, p. 49, rosiperda, p. 54, Weidenau, rhamni, Germany, Rübsaamen, Verh. z.-b. Wien, zlii; D. voraz, p. 393, necans, pastinaca, p. 395, solani, p. 396, Germany, Rübsaamen, Ber Zeit. xxxvi; D. coccidarum, Jamaica, Cockerell, Ent. xxv, p. n. spp.
- Hormomyia hartigi, Lorraine, LIEBRL, p. 285, Ent. Nachr. xviii; H. haueri. p. 394, brunnea, p. 396, Europe, RUBSAAMEN, Berl. En xxxvii: n. spp.
- Lasioptera carophila, & and larva described; KIEFFER, Wien. ent Macrolabis, n. g. for a part of Cocidomyia, p. 214, including M. s n. sp., Elbeuf, p. 215; KIEFFER, Wien. ent. Z. xi.
- Monarthropalpus, n. g. for flavus, Schrank; RÜBSAAMEN, p. 381. Ent. Zeit. xxxvii.
- Rhopalomyia, n. g. for species previously referred to Cecidomyia o momyia; RÜBSAAMEN, pp. 370 & 375, Berl. Ent. Zeit. xxxvii.
- Schicomyki, n. sp., larva noticed, p. 220, pl. i, fig. 8; S. (sub E<sub>1</sub> sigripes, Lw., larva described, Kieffer, Wien. ent. Z. xi.

# Mycetophilida, Simulida, Bibionida, Blepharoceri

- Blepharocerida: synopsis of the described genera and species; ( SACKEN, Berl. Ent. Zeit. xxxvi, pp. 407-411.
- Bolitophila luminosu, noticed, metamorphoses figured; Hudson, M p. 49, frontispiece.
- Colosia fusca, Pavia, BEZZI, Bull. Ent. Ital. xxiv, p. 68, n. sp.

- Mycetophila antarctica, New Zealand, Hudson, Manual, p. 46, pl. iv, fig. 5, n. sp.
- Simulia columbacensis, metamorphoses; Tömösváry, Glasnik Naravosl. Druzt. vi, p. 187, &c., pl. ii.
- Plecia concolor, New Grenada, WULP, p. 197, Tijdschr. Ent. xxxiv, n. sp.

## CULICIDE, CHIRONOMIDE.

- Ceratopogon antipodum, New Zealand, Hudson, Manual, p. 45, pl. iv, fig. 4, n. sp.
- Chironomus: Mermis larva in larva of; TRYBOM, Ent. Tidskr. xiii, pp. 81-92.
- C. zealandicus, New Zealand, Hudson, Manual, p. 43, pl. iv, fig. 2, n. sp. Corethra antarctica, New Zealand, Hudson, Manual, p. 43, pl. iv, fig. 3, n. sp.

#### TIPULIDÆ.

- Adelphomyia, n. g. near Limnophila, for A. helvetica, n. sp., Switzerland; Векскогн, р. 134, МТ. Ges. Bern. 1890.
- Dicranota brevitarsis, p. 135, brevicornis, longitarsis, p. 136, Switzerland, Вевскотн, МТ. Ges. Bern. 1890, n. spp.
- Gnophomyia tripudians, Switzerland, BERGROTH, p. 131, MT. Ges. Bern. 1890, n. sp.
- Limnophila denticulata, Switzerland, Bergroth, p. 132, MT. Ges. Bern. 1890, n. sp.
- Orimargula, Mik, is the same as Antocha, O.-S., the species discussed; OSTEN SACKEN, pp. 313 [in error for 413]-416, Berl. Ent. Zeit. xxxvi.
- Pachyrhina immaculata, Java, Wulp, p. 196, Tijdschr. Ent. xxxiv, n. sp. Rhypholophus egenus, Switzerland, Bergroth, p. 131, MT. Gea. Bern. 1890, n. sp.
- Tipula bicornis: habits, oviposition, WEBSTER, Bull. Dep. Agric. Ent. xxvi, pp. 66-70; T. dispar and pagana, distinctions of, Dale, Ent. M. M. xxviii, p. 310.
- T. saginata, Switzerland, BERGROTH, p. 137, MT. Ges. Bern. 1890;
  T. ornaticornis, New Grenada, Wulp, p. 195, pl. xii, fig. 1, Tijdschr. Ent. xxxiv;
  T. fumiponnis, New Zealand, Hudson, Manual, p. 48, pl. v, fig. 2: n. spp.

#### XYLOPHAGIDA and STRATIOMYLIDA.

Beris vallata, mimics Athalia; GIARD, p. 43, CR. Soc. Biol. (n.s.) iv. Dialysis disparilis, spurs of tibia; BERGROTH, p. 162, Wien. eut. Z. xi. Platyna hastata, characters of; RÖDER, p. 271, Wien. ent. Z. xi. Stratiomys longicornis, histology of nervous system of larva; Henneguy & Binet, Ann. Soc. Ent. Fr. lxi, pp. 309-316, pl. vi.

### TABANIDA.

- Atylotus letourneusi, Batna, p. 643, camaronensis, Madeira, cereolus, E. Africa, p. 644, niveipalpis, rusiceps, p. 645, namaquinus, p. 646, soulousneis, S. Africa, alazinus, Caucasus, p. 647, polyzonatus, p. 648, canipalpis, Persia, macer, albocostatus, p. 649, conicus, p. 650, melanopygatus, leucopogon, p. 651, angustus, oxyceratus, p. 652, assamensis, birmanicus, p. 653, pagodinus, monilifer, p. 654, monotaniatus, p. 655, nephodes, leucocnematus, p. 656, joidus, flaviventris, p. 657, E. India, fuscicauda, Ceylon, pallidepectoratus, Saigon, p. 658, bituberculatus, China ?, p. 659, tenessensis, acutus, N. America, obesus, p. 660, erythraus, p. 661, fuscicrura, Mexico, picticornis, Haiti, p. 662, unipunctatus, S. America, rubescens, Uruguay, p. 663, eutemiatus, maletectus, p. 664, Brazil, pulverulentus, Cayenne, aurisquammatus, Brazil, p. 665, indecisus, p. 666, simplex, Amazons, polytænia, Buenos Ayres, p. 667, erythrocephalus, Panama, andamanicus, Andaman Is., p. 668, cinerascens, lacrymans, p. 669, abbreviatus, Java, albonotatus, New Caledonia, p. 670, A.? picticornis, p. 671, sonnerati, alfourensis, p. 672, New Guinea, laglaisei, Waigiou, avidus, rufinotatus, p. 673, hyperythreus, p. 674, sanguinarius, claripennis, p. 675, erraticus, rubrinotatus, hab. ? p. 676, BIGOT, Mém. Soc. Zool.Iv, n. spp.
- Bellardia sinica, China, p. 629, annamita, Salgon, rubribarbis, S. America, p. 630, furcata, hab. ?, p. 631, Bigor, Mém. Soc. Zool. v, n. spp.
- Bolbodimyia, n. g. for B. bicolor, n. sp., Venezuela, Bigor, p. 162, Wienent. Z. xi; see also Röder, t. c. p. 237.
- Chrysops iranensis, Persia, cinctus, Philippines, p. 602, atricornis, p. 603, nigriventris, p. 604, coloradensis, p. 605, N. America, oculatus, Brazil, p. 606, trimaculatus, Senegal, p. 607, Bigot, Mém. Soc. Zool. v, n. spp.
- Corizoneura annulata, S. Europe ?, p. 612, albifrons, p. 613, brachyrrhyncha, Cape Good Hope, nigricornis, Australia, p. 614, ruficornis, velutina, p. 615, California, leucopicta, New Caledonia, trichocera, p. 616, rubiginosa, C. ? angusta, p. 617, Australia, Bigor, Mém. Soc. Zool, v. n. spp.
- Dasybasis tristis, Chili, BIGOT, p. 621, Mém. Soc. Zool. v, n. sp.
- Diachlorus barbatus, Piedmont, p. 622, maroccanus, Marocco, notatus, California, p. 623, D. ? hamatopotides, N. America, p. 624, D. melas, Sydney, p. 625, Bigot, Mém. Soc. Zool. v. n. spp.
- Diatomineura californica, California, p. 618, hirtipalpis, Chili, carnea, p. 619, D. ? gagantina, p. 620, Australia, Bigot, Mém. Soc. Zool. v, n. spp.
- Dichalacera pachypalpus, Mexico, p. 631, japonica, Japan, satanica, p. 632, castanea, albopicta, p. 633, marmorata, p. 634, Brazil, peruviana, p. 635, Peru, BIGOT, Mém. Soc. Zool. v. n. spp.
- Erephosis vertebrata, Australia, BIGOT, p. 618, Mém. Soc. Zool. v, n. sp.
- Goniops, n. g., p. 236, for G. hippoboscoides, n. sp., Dakota, p. 237, Aldrich, Payche, vi.

- Hamatopota: the species tabulated last year now described; Bigot, pp. 626-629, Mém. Soc. Zool. v.
- Mesomyia? maoriorum, New Zealand, Bigot, p. 621, Mém. Soc. Zool. v, n. sp.
- Mycteromyia nigrifacies, India, p. 607, ensata, elegans, Cape Good Hope, p. 608, nitens, p. 609, penicillata, Brazil, cinerascens, Chili, p. 610, albipectus, p. 611, erythronotata, p. 612, Brazil, Bigot, Mém. Soc. Zool. v, n. spp.
- Pangonia bricchettii, Somaliland, Bezzi, p. 181, Ann. Mus. Genov. (2) xii; P. bahiuna, Brazil, Bigot, p. 612, Mém. Soc. Zool. v: n. spp.
- Selasoma tibialis, note on; RÖDER, p. 237, Wien. ent. Z. xi.
- Silvius bicolor, Marocco, Bigot, p. 625, Mém. Soc. Zool. v, n. sp.
- Stibasoma pachycephalum, Mexico, bicolor, Brazil, Bigot, p. 636, Mém. Soc. Zool. vi, n. spp.
- Tabanus swiridowi, figured, pl. i, fig. 2, Hor. Ent. Ross. xxvi.
- T. lama, Mongolia, Portschinsky, p. 201, Hor. Ent. Ross. xxvi; T. yulensis, New Guinea, Röder, p. 244, S. E. Z. liii; T. oritensis, Caucasus, p. 677, glaber, India, yokoamensis, Japan, p. 678, nitidulus, rufocallosus, Java, p. 679, leonum, Sierra Leone, p. 680, fuscipalpis, tetropsis, p. 681, N. America, sexvittatus, Mexico, p. 682, atricornis, S. America, pruinosus, Mexico, p. 683, villosulus, California, discifer, Amazons, p. 684, venosus, p. 685, callicera, albidocinctus, p. 686, macroceratus, Brazil, erythræus, La Plata, p. 687, lunulatus, p. 688, Australia, lifuensis, Lifu Is., indistinctus, p. 689, raffreyi (sic), p. 690, New Guinea, Bigot, Mém. Soc. Zool. v: n. spp.
- Therioplectes athereus, Europe?, trichocerus, p. 637, fezianus, p. 638, Marocco, batnensis, calopis, p. 639, Batna, leucophorus, p. 640, maculifer, hirtulus, p. 641, N. America, limbatus, Buenos Ayres, T.? melanorhinus, N. America, p. 642, BIGOT, Mém. Soc. Zool. v, n. spp.
- Veprius rufopilosus, Chili, BIGOT, p. 620, Mém. Soc. Zool. v, n. sp.

### NEMESTRINIDE and BOMBYLIIDE.

- Classification of Bombyliidæ, with table of genera; Bigor, pp. 321-342, Ann. Soc. Ent. Fr. lxi.
- Acreotrichus inappendiculatus, Australia, BIGOT, p. 366, Ann. Soc. Ent. Fr. lxi, n. sp.
- Amictus auripilus, N. America, p. 372, cinerascens, Chili, p. 373, Bigor, Ann. Soc. Ent. Fr. lxi, n. spp.
- Anthrax: revision of the N. American species, with notes, synonymy, and tables; Coquillett, Tr. Am. Ent. Soc. xix, pp. 168-187.
- A. tangerinus, Marocco, p. 353, aurocinctus. Senegal, fissus, N. America, p. 354, eurhinatus, Mexico, ruficollis, Venezuela, p. 355, bifenestratus, California, unicinctus, p. 356, bipenicillatus, obscuripes, p. 357, melanogaster, pallipes, micromelas, p. 358, Chili, Bigot, Ann. Soc. Ent. Fr. lxi; A. nitidofasciatus, Central Asia, Portschinsky, p. 208, pl. i, fig. 5, Hor. Ent. Ross. xxvi; A. muscaria, p. 178, pretiosa, vacans,

- p. 179, cinefacta, p. 180, inculta, terrena, squamigera, p. 181, gemella, hircina, telluris, p. 182, vulpina, crocina, p. 183, tantilla, variata, rasta, p. 184, lacunaris, fenestratoides, p. 185, sabulosa, p. 186, arenosa, p. 187, N. America, Coquillett, Tr. Am. Ent. Soc. xix: n. spp.
- Apiomidas, n. g. near Bhaphiomidas, for A. trochilus, n. sp., California; Coquillett, p. 314, Canad. Ent. xxiv.
- Argyromæba appendiculata, Pondichery, varicolor, p. 347, melanopogon, albosparsa, N. America, p. 348, aterrima, no locality, acroleuca, China ?, crinita, p. 349, plurinota, p. 350, Chili, Bigot, Ann. Soc. Ent. Fr. lxi, n. spp.
- Bombylius nigropenicillatus, Asia Minor, fulvipes, albosparsus, Pondichery, p. 302, albopenicillatus, Mexico, laticeps, p. 363, cinereus, California, ochraceus, Monte Video, australianus, p. 364, rubriventris, p. 365. Sydney, Bigot, Ann. Soc. Ent. Fr. lxi, n. spp.
- Comptosia fulvipes, Australia, BIGOT, p. 359, Aun. Soc. Ent. Fr. lxi; C. virida, New Zealand, HUDSON, Manual, p. 55, pl. vi, fig. 3: n. spp. Conophorus mauritanicus, p. 360, Algeria, melanoceratus, California,

p. 361, Bigot, Ann. Soc. Ent. Fr. lxi, n. sp.

- Dischistus cycnus, leucophys, Cape Good Hope, p. 368, melanurus, Natal, fuscipes, N. America, p. 369, Bigor, Ann. Soc. Ent. Fr. lxi, n. spp.
- Eclimus venosus, Europe ?, Bigot, p. 371, Ann. Soc. Ent. Fr. lxi, n. sp.
- Epacmus rufolimbatus, California, BIGOT, p. 359, Ann. Soc. Ent. Fr. lxi; E. concinnus, pellucidus, fumosus, California, LYMAN, pp. 10 & 11, Canad. Ent. xxiv: n. spp.
- Eurycarenus? pachyceratus, Cape Good Hope, Bigor, Ann. Soc. Ent. Fr. p. 371, n. sp.
- Exoprosopa: characters of some N. American; Coquillett, p. 123
  Canad. Ent. xxiv.
- E. herzi, Korea, Portschinsky, p. 207, pl. i, fig. 4, Hor. Ent. Ross. xxvi E. vitrea, Pondichery, melanura, N. America, p. 344, pallens, Cali fornia, p. 345, bipartita, Colombia, latelimbata, Australia, p. 346 Bigot, Ann. Soc. Ent. Fr. lxi; E. grata, California, Coquillett p. 124, Canad. Ent. xxiv: n. spp.
- Geron cothurnatus, Tasmania, dicroma, Cape Good Hope?, BIGOT, Ant Soc. Ent. Fr. lxi; G. fasciola, p. 125, capax, p. 126, California Coquillett, Canad. Ent. xxiv: n. spp.
- Glossista costata, Amazons, multicolor, Chili, p. 352, lipposa, Sydne p. 353, Bigot, Ann. Soc. Ent. Fr. lxi, n. spp.
- Hemipenthes castanipes, p. 350, latelimbatus, p. 351, N. America, Bigo Ann. Soc. Ent. Fr. lxi, n. spp.
- Heterostylum pallipes, Haīti, BIGOT, p. 361, Ann. Soc. Ent. Fr. lxi, n. s Hyperalonia flavosparsa, Brazil, argenticincta, Australia, BIGOT, p. 34 Ann. Soc. Ent. Fr. lxi, n. spp.
- Nemestrina bombiformis, Caucasus, Portschinsky, p. 203, pl. i, fig. Hor. Ent. Ross. xxvi, n. sp.
- Phthiria tristis, pallipes, Chili, Bigot, p. 367, Ann. Soc. Ent. Fr. l. n. spp.

Sparnopolius limbatus, Australia, BIGOT, p. 370, Ann. Soc. Ent. Fr. lxi, n. sp.

Spogostylum? inappendiculatum, Chili, description, Bigor, p. 351, Ann. Soc. Ent. Fr. lxi,? n. sp.

Systechus eupogonatus, E. India, p. 365, eulabiatus, Tasmania, p. 366, Bigor, Ann. Soc. Ent. Fr. lxi, n. spp.

Systropus calopus, Chili, BIGOT, p. 373, Ann. Soc. Ent. Fr. lxi, n. sp.

Thevenetimya melanopogon, N. America, Bigot, p. 370, Ann. Soc. Ent. Fr. lxi, n. sp.

Usia gagathea, Algeria, Bigor, p. 374, Ann. Soc. Ent. Fr. lxi, n. sp.

### Asilida.

Additions and corrections to Williston's Catalogue of S. American; OSTEN-SACKEN, Berl. Ent. Zeit. xxxvi, pp. 417-428.

Apiocerina, aberrant group of Asilidæ discussed and defined; OSTEN-SACKEN, Berl. Ent. Zeit. xxxvi, pp. 311-315.

Apogon dufourii, Perr., synonymical note; Röder, p. 248, Ent. Nachr. xviii.

Brachyrrhopala victoriæ, Australia, Röder, p. 241, S. E. Z. liii, n. sp.

Chrysopogon mulleri, Victoria, RÖDER, p. 243, S. E. Z. liii, n. sp.

Lophonotus dubius, Somaliland, Bezzi, p. 187, Ann. Mus. Genov. (2) xii, n. sp.

Mochtherus striatus, Java, Wulp, p. 198, pl. xii, figg. 3-6, Tijdschr. Ent. xxxiv, n. sp.

Philodicus pavesii, Somaliland, Bezzi, p. 185, Ann. Mus. Genov. (2) xii, n. sp.

Promachus leucotrichodes, p. 210, tristis, p. 211, Mahé, Bigot, Bull. Soc. Z. Fr. 1892, n. spp.

Saropogon luctuosum, n. var. ticinense; BEZZI, p. 107, Bull. Ent. Ital. xxiv.

### LEPTIDE, THEREVIDE, EMPIDE.

Chamedipsia, diagnosis corrected; Mik, p. 55, Wien. ent. Z. xi. Hilara, monographic revision of the Central European species; STROBL, Verh. s.-b. Wien. xlii, pp. 85–182.

H. aëronetha, p. 81, novakii, p. 83, S.E. Europe, Mik, Wien. ent. Z. xi;
H. pesudochorica, p. 109, lasiochira, p. 110, hystrix, p. 112, bivittata,
p. 113, qudrifaria, pectinipes, p. 116, andermattensis (subsp. of longevittata), simplicipes, p. 119, tyrolensis, p. 121, sulcitarsis, p. 123, diversipes, p. 125, dimidiata, p. 128, angustifrons, p. 133, braueri,
p. 135, argyrosoma, p. 136, lacteipennis, p. 137, hirta, p. 141, beckeri,
p. 143, carinthiaca, p. 144, tiefii, p. 150, pseudosartrix, p. 152, cinereomicans, p. 156, mikii, p. 158, discolor, p. 165, pilosopectinata,
p. 169, Central Europe, Strobl, Verh. z.-b. Wien. xlii: n. spp.

Leptis, attacking man; Leprevost & Gazagnaire, pp. clv & clvi, Bull. Soc. Ent. Fr. lxi.

- L. subpilosa, Switzerland, BECKER, p. 23, Wien. ent. Z. xi, n. sp.
- Ptiolina, Schin. noc Zett., to be replaced by Symphoromyia, Francef.; Strobl., p. 123, Wien. ent. Z. xi.
- Spania: notes on the genus and its species; Ptiolina, Zett., merged in it; Strobl, pp. 121-125, Wien. ent. Z. xi.
- Steleocheta meridionalis, n. n. for stiriensis, Beck.; BECKER, p. 126, Wien. ent. Z. xi.

#### DOLICHOPODIDE.

- Achalcus scutellaris, Java, Wulp, p. 202, Tijdschr. Ent. xxxiv, n. sp. Bathycranium, n. g. for Dolichopus bicolorellus, Zett.; Strobl., p. 102, Wien. ent. Z. xi.
- Campsionemus thalhammeri, Austria, STROBL, p. 106, Wien. ent. Z. xi. n. sp.
- Chrysotus nigricilius, Loew., referred to Melanostolus; STROBL, p. 105 Wien, ent. Z. xi.
- Coracocephalus, n. g. near Eucoryphus, p. 279, for C. stroblii, n. sp., Alp of Steiermark, p. 281; Mik, Wien. ent. Z. xi.
- Dolichopus bicingulatus, Zett., referred to Gymnopternus, and characterised STROBL, p. 103, Wien. ent. Z. xi.
- Lyroneurus chalybæus, Ecuador, Böder, p. 81, Soc. Ent. vii, n. sp.
- Psilopus fenestratus, p. 200, pl. xii, fig. 6, bifilum, p. 201, setosus, p. 202 Java, Wulp, Tijdschr. Ent. xxxiv, n. spp.
- Sphyrotarsus hygrophilus, characters amended; BECKER, p. 126, Wiel ent. Z. xi.
- Thrypticus divisus, & described; Strobl, p. 105, Wien. ent. Z. xi.

#### SYBPHIDE.

- Synonymical and critical notes on the Syrphidæ of Biol. Centr. At Dipt.
- Argentinomyia grandis, Argentine Rep., ARRIBALZAGA, p. 255, An. 80 Arg. xxxiv, n. sp.
- Asemosyrphus olivaceus, griseus, impurus, Mexico, Giglio-Tos, p. 6, Bo Mus. Zool. Tor. vii, No. 123, n. spp.
- Baccha clavata, synonyms of; WILLISTON, p. 145, Ent. News, iii.
- B. spatulata, Mexico, Giglio-Tos, p. 4, Boll. Mus. Zool. Tor. vii, No. 15 n. sp.
- Brachyopa cynops, Colorado, Snow, Kansas Quart. i, p. 37, pl. vii, fig. n. sp.
- Callicera montensis, Colorado, Snow, Kansas Quart. i, p. 34, pl. vii, fig. n. sp.
- Camerania, n. g. for Temnocera megacephala, Loew.; Giglio-Tos, B. Mus. Zool. Tor. vii, No. 117.
- Ceria verralli, Panama, p. 75, meadei, pl. ii, fig. 12, schnablei, p. 76, be rothi, p. 77, Mexico, Williston, Biol. Centr. Am. Dipt. iii.
- Ceriogaster auricaudata, Mexico, Williston, p. 73, pl. ii, fig. 10, Bi Centr. Am. Dipt. iii, n. sp.

- Chilosia aurotecta, Mexico, Giglio-Tos, p. 4, Boll. Mus. Zool. Tor. vii, No. 132, n. sp.
- Chrysotoxum derivatum, variation; Snow, Kansas Quart. i, p. 34.
- Copestylum parvum, simile, Mexico, Giglio-Tos, p. 2, Boll. Mus. Zool. Tor. vii, No. 123, n. spp.
- Crioprora arctophiloides, Mexico, Giglio-Tos, p. 7, Boll. Mus. Zool. Tor. vii, No. 123, n. sp.
- Criorhinia coquilletti, California, Williston, p. 145, Ent. News, iii, n. sp. Eristalis alpinus, n. var. caucasicus; Portschinsky, p. 209, pl. i, fig 6, Hor. Ent. Ross. xxvi. E. vinetorum, albifrons, scutellaris, complex synonymy; Williston, p. 146, Ent. News, iii.
- E. bombusoides, p. 4, trilimbatus, præclarus, clarissimus, sallei, p. 5, triangularis, p. 6, Mexico, Giglio-Tos, Boll. Mus. Zool. Tor. vii, No. 123; E. sumichrasti, atropos, Mexico, Giglio-Tos, p. 1, 4. c. No. 132; E. persa, p. 58, circe, p. 59, pl. ii, fig. 3, trigonus, p. 61, fig. 4, cemulus, pl. ii, fig. 5, minutalis, fig. 6, p. 64, Central America, Williston, Biol. Centr. Am. Dipt. iii : n. spp.
- Helophilus mexicanus, variation noticed, and synonymy; WILLISTON, p. 68, Biol. Centr. Am. Dipt. iii.
- Lycastrirhyncha nitens, Big., redescribed; WILLISTON, p. 66, Biol. Centr. Am. Dipt. iii.
- Mallota? championi, p. 69, margarita, pl. ii, fig. 7, smithi, fig. 8, p. 70, Mexico, Williston, Biol. Centr. Am. Dipt. iii, n. spp.
- Megametopon, Giglio-Tos, = (Ophromyia, Will.); Williston, p. 79, Biol. Centr. Am. Dipt. iii.
- Melanostoma stegnum, Say, = (tigrina, O. Sack.); WILLISTON, p. 145, Ent. News, iii.
- M. elegans, p. 2, bellum, p. 3, Mexico, Giglio-Tos, Boll. Mus. Zool. Tor. vii, No. 132, n. spp.
- Mesogramma rombicum, diversum, ciliatum, p. 3, comma, p. 4, Mexico, Giglio-Tos, Boll. Mus. Zool. Tor. vii, No. 132, n. spp.
- Mesograpta willistonii, Argentine Rep., ARRIBALZAGA, An. Soc. Arg. xxxiii, p. 54, n. sp.
- Microdon: oviposition in ants' nest; VERHOEFF, p. 13, Ent. Nachr. xviii.
- M. megalogaster, Colorado, Snow, Kansas Quart. i, p. 34, pl. vii, fig. 1;
  M. aquilinus, Mexico, Giglio-Tos, Boll. Mus. Zool. Tor. vii, No. 123,
  p. 2: n. spp.
- Milesia pulchra, Gnatemala, WILLISTON, p. 74, pl. ii, fig. 11, Biol. Centr. Am. Dipt. iii, n. sp.
- Mixogaster dimidiata, Mexico, Giglio-Tos, Boll. Mus. Zool. Tor. No. 123, p. 1, n. sp.
- Nausigaster bonariensis, S. America, ARRIBALZAGA, p. 254, An. Soc. Arg. xxxiv, n. sp.
- Ocyptamus trigonus, Wied., = (Baccha torva, Willist.); WILLISTON, p. 146, Ent. News, iii.
- Omegasyrphus, distinctions from Microdon; Giglio-Tos, Boll. Mus. Zool. Tor. No. 118.

- Phalacromyia albitarsis, Argentine Rep., ARRIBALZAGA, p. 244. An. So Arg. xxxiv, n. sp.
- Platymochatus niger, Mexico, Giglio-Toa, Boll. Mus. Zool. Tor. v No. 123, p. 6, n. sp.
- Promilesia, n. g., Eristalini, p. 241, for P. nectarinoides, n. sp., Argenti Bep. p. 243, Arribalzaga, An. Soc. Arg. xxxiii.
- Filota, habits and systematic position; MIK, p. 183, Wien. ent. Z. xi. Rhopalosyrphus. = (Holmbergia, Arribalk.); Giglio-Tos, Boll. Mi Zool. Tor. vii, No. 118.
- Sulpingogaster nova, Mexico, Giglio-Tos, p. 7, Boll. Mus. Zool. Tor. v No. 123, n. sp.
- Spherophoria syrphica, Mexico, Giolio-Tos, p. 2, Boll. Mus. Zo. Tor. vii, No. 132, n. sp.
- Syrphus pauxillus, 2 described; Snow, Kansas Quart. i, p. 37.
- S. rusicauda, Colorado, Snow, Kansas Quart. i, p. 36, pl. vii, fig. ; S. lautus, saussurii, Mexico, Giglio-Tos, p. 2, Boll. Mus. Zool. To vii, No. 132; S. patagonus, p. 115, walkeri, p. 117, Patagonia, Arr Balzaga, An. Soc. Arg. xxxiii: n. spp.
- Temnocera spinithorax, Argentine Rep., BERG, p. 190, An. Soc. Ar xxxiii, n. sp.
- Tropidia insularis, Parana, Arribalzaga, p. 238, An. Soc. Arg. xxxi
- Ubristes chrysopyga, Mexico, Giglio-Tos, Boll. Mus. Zool. Tor. No. 12 p. 1. p. sp.
- Volucella pallens, Wied., = (serpunctata, Loew.); WILLISTON, p. 14 Ent. News, iii.
- V. omochroma, furens, craverii, p. 2, flavissima, minima, dichroica, trigos hyaloptera, hirsuta, p. 3, hispida, volucris, hystrix, brevis, obesoid p. 4, Mexico, Giglio-Tos, Boll. Mus. Zool. Tor. vii, No. 123; V. t. tacea. Curaços, Wulp, p. 203, Tijdschr. Ent. xxxiv; V. ambroset p. 179, obscuripennis, p. 180, missionera, picturata, p. 184, Argenti Rep., Berg, An. Soc. Arg. xxxiv: n. spp.
- Xiphopheromyia, n. g., Bigot, for X. glossata, n. sp., E. India, Bigot, p. 161, Wien. ent. Z. xi.
- Xylota angustiventris, Lw., = (slongata, Willist.); WILLISTON, p. 1-Ent. News, iii.
- X. ruspes, paurilla, pl. ii, fig. 9, p. 71, brachygaster, stenogaster, p. Mexico, Williston, Biol. Centr. Am. Dipt. iii, n. spp.

#### Œstrida.

- Historical, critical, and descriptive review of the larvæ of *Œstrides* th are known to attack man in America; Blanchard, Ann. Soc. El Fr. lxi, pp. 109-154, and Bull. Soc. Ent. Fr. lxi, p. ccix.
- Ciphenomyia and Hypoderma: note on the unknown stage in the law history: Brayer, pp. 79-84. Verh. z.-b. Wien, zlii.

- Cuterebra, larva from Lepus artemisia? described; Townsend, Pysche, vi, p. 299.
- Dermatobia sp. from Lepus callotis described; Townsend, Psyche, vi, p. 299.
- Dermatæstrus, n. g., with strepsicerontis, n. sp. from Strepsiceros capensis, S. Africa; Brauer, p. 12, pl. i, fig. 1, SB. Ak. Wien, ci.
- Gastrophilus nasalis: habits; Townsend, p. 227, Ent. News, iii.
- Hypoderma lineata: habits and larva described; RILEY, Ins. Life, iv, pp. 302-317.
- Œstrus attacking man, Sangagli (791); Œ. ovis attacking Capra hircus, Blanchard, p. cexlvi, Bull. Soc. Ent. Fr. lxi.
- **E.** perplexus, New Zealand, Hudson, Manual, p. 63, pl. vii, fig. 12, n. sp. Strobilæstrus, n. g., with S. antilopinus, n. sp., from Antelopes in Transvaal; Brauer, p. 10, pl. i, fig. 2, SB. Ak. Wien, ci.
- Tachinæstrus semenowi, note on; Portschinsky, p. 209, Hor. Ent. Ross.

## CONOPIDÆ, PIPUNCULIDÆ, PLATYPEZIDÆ.

- Brachyceraa, n. g., for Conops brevicornis, Loew.; Röder, p. 366, Ent. Nachr. xviii.
- Callomyia bella, Mexico, Williston, p. 89, Biol. Centr. Am. Dipt. iii, n. sp.
- Conops curbonarius, nigrifacies, ochreiceps, Bigot, redescribed and referred to Physocephala; WILLISTON, pp. 82 & 83, Biol. Centr. Am. Dipt. iii.
- C. ornatus, Mahé, BIGOT, p. 211, Bull. Soc. Z. Fr. 1892; C. magnus, p. 43, grandis, rufus, augustifrons, p. 44, nobilis, inornatus, p. 45, ornatus, parvus, p. 46, Chapada, Brazil, WILLISTON, Kansas Quart. i; C. ocellatus, Mexico, GIGLIO-TOS, Boll. Mus. Zool. Tor. vii, No. 132; C. discalis, anthreas, sequax, Mexico, WILLISTON, Biol. Centr. Am. Dipt. iii: n. spp.
- Dalmannia vitiosa, California, Coquillett, p. 150, Ent. News, iii, n. sp.
   Physocephala sororcula, xanthops, Mexico, Williston, p. 83, Biol. Centr.
   Am. Dipt. iii; P. maxima, Mexico, Giglio-Tos, p. 5, Boll. Mus.
   Zool. Tor. vii, No. 132: n. spp.
- Pipunculus elegantulus, flavitarsis, xanthopodus, p. 87, aculeatus, flavicornis, p. 88, Mexico, WILLISTON, Biol. Centr. Am. Dipt. iii, n. spp.
- Stylogaster: notes on and characters of the species; RÖDER, p. 286, Wien. ent. Z. xi.
- Zodion auricaudatum, Mexico, WILLISTON, p. 85, pl. ii, fig. 14, Biol. Centr. Am. Dipt. iii, n. sp.

## MUSCIDÆ CALYPTRATÆ.

- Biology of coprophagous and necrophagous Diptera (in Russian); PORT-CHINSKY, Hor. Ent. Ross. xxvi, pp. 63-131.
- Table of the genera of N. American Calyptrate Muscidæ; Townsend pp. 133-144, Tr. Am. Ent. Soc. xix.

- BRAUER & BENGENSTAMM, Denk. Ak. Wien, lviii, pp. 3.5-445, have vised their work, published two years ago, on Maccaria echinam making a large number of alterations and additions: these late give on pp. 298-301, in the order followed by the authors; taxonomical and synonymical changes involved by this memoir be sought in the pages of the work itself. It concludes will very useful lists of the whole of the genera and groups, and a species, 1520 in number, examined.
- Acyglosa diversa, Rond., = (Anthonyia corvina, Loew.): Röder, Nachr. xviii, p. 366.
- Anisia randerwulpi, Jamaica, Towesend, p. 81, Ent. News, iii, n. s. Anthomyiida: study of the Anthomyiid types in the Fallen-Zette collection at Lund, Stein, Ent. Nachr. xviii, pp. 321-333; Sylof the N. American genera, Towesend, pp. 290-293, Tr. Am. Soc. xix.
- Aricia parasitic in Carabus; GIGLIO-Tos (339, 340).
- Aphria angustifrons, England, MEADE, p. 114, Wien. ent. Z. xi, n. sp. Aporia limacodis, New York, Townsend, Psyche, vi, p. 275, n. sp.
- Atropharista, n. g. Hystriciina, for A. jurinoides, n. sp., Dakota; T. END, p. 92, Tr. Am. Ent. Soc. xix.
- Atrotophopalpus, n. g., Phytoinæ, p. 130, for A. angusticornis, 1 Florida, p. 131; TOWESEND, Ent. News, iii.
- Blepharipeza ezul, New Hampshire, Townsend, p. 64, Canad. Ent. B. rufescens, Maryland, Townsend, p. 90, Tr. Am. Ent. Soc B. nigrisquamis, Jamaica, Townsend, p. 80, Ent. News. iii: n
- Brachycalia, n. g., for Scopolia ocypterina, Zett.; Meade, p. 26 M. M. xxviii.
- Brachycoma smerinthi, England, MEADE, Ent. M. M. xxviii, p. chihuahuaensis, Mexico, Townsend, p. 165, Canad. Ent. xxiv:
- Calliphora sp. larve killing toads; Guthrie, Ent. M. M. xxviii, p Chatoglossa, n. g., Phytoina, p. 125, for C. picticornis, viola, nigri
- n. spp., Florida, p. 126; TOWNSEND, Tr. Am. Ent. Soc. xix.

  Clista americana, Illinois, Townsend, p. 78, Canad. Ent. xxiv, n. s
- Clistomorpha, n. g. (Phytoinæ). p. 79, for C. hyalomoides, n. sp., N. TOWNSEND, p. 80, Canad. Ent. xxiv.
- Cænosia brachialis, Rond., = (Caricea stroblii, Mik); Mik, p. 186, ent. Z. xi.
- C. submaculata, p. 208, ferruginea, p. 209, Java; WULP, Tijdsch xxxiv, n. spp.
- Coronimyia, n. g., Phytoinæ, p. 128, for C. geniculata, n. sp., I p. 129; Townsend, Tr. Am. Ent. Soc. xix.
- Crossocosmia: affinities of, and validity of name; Mik, p. 116, ent. Z. xi.
- Daochata, n. g., Tachinina, p. 97, for D. harveyi, n. sp., Maine, Townsend, Tr. Am. Ent. Soc. xix.
- Desvoidia n. g. (Tachinida), near Metopia, for D. fusca, n. sp., En MEADE, p. 179, Ent. M. M. xxviii.

- Deviides: synoptic table of the N. American genera; Townsend, pp. 273-284, Tr. Am. Ent. Soc. xix.
- Deponchelia, B. Desv., = (Parasetigena, Br. & Berg.); Mik, p. 117, Wien. ent. Z. xi.
- Echinomyia dakotensis, N. America, Townsend, p. 94, Tr. Am. Ent. Soc. xix, n. sp.
- Emphanopteryx, n. g., Thryptoceratina, p. 120, for E. eumyothyroides, n. sp., New York, p. 121; TOWNSEND, Tr. Am. Ent. Soc. xix.
- Euceromyia, n. g., p. 115, near Raselia, for E. robertsoni, n. sp., Illinois, p. 116; Townsend, Tr. Am. Ent. Soc. xix.
- Eucnephalia, n. g., Phoroceratinæ, p. 166, for E. gonoides, n. sp., New Mexico, p. 167; TOWNSEND, Canad. Ent. xxiv.
- Eulasiona, n. g., Thryptoceratina, p. 119, for E. comstocki, n. sp., New York, p. 120; Townsend, Tr. Am. Ent. Soc. xix.
- Eumacronychia, n. g., Tachinina, p. 98, for E. decens, n. sp., N. America, p. 99; Townsend, Tr. Am. Ent. Soc. xix.
- Eumyothyria, n. g., Phytoinæ, p. 121, for E. illinoiensis, n. sp., N. America, p. 122; TOWNSEND, Tr. Am. Ent. Soc. xix.
- Euphorocera, n. g., Phoroceratinæ, for E. tuchinomoides, n. sp., New Mexico; Townsend, p. 112, Tr. Am. Ent. Soc. xix.
- Euscopolia, n. g., Phytoina, p. 123, for E. dakotensis, n. sp., N. America, p. 124; TOWNSEND, Tr. Am. Ent. Soc. xix.
- Euthyprosopa, n. g., Tachininæ, p. 106, for E. petiolata, n. sp., New Mexico, p. 107; Townsend, Tr. Am. Ent. Soc. xix.
- Exorista endryæ, p. 287, platysamiæ, datanæ, p. 288, N. America, .
  TOWNSEND, Tr. Am. Ent. Soc. xix, n. spp.
- Ginglymia, n. g., Thryptoceratinæ, p. 118, for G. acrirostris, n. sp., Michigan, p. 119; TOWNSEND, Tr. Am. Ent. Soc. xix.
- Glaucophana, Br. & Berg., queried as identical with Neara, Rond.; Mik, p. 185, Wien. ent. Z. xl.
- Gonia sagax, Iowa, Townsend, p. 65, Canad. Ent. xxiv, n. sp.
- Gymnoprosopa, n. g., Tachinina, p. 108, for G. polita, argentifrons, clarifrons, n. spp., N. America, p. 109; Townsend, Tr. Am. Ent. Soc. xix.
- Helocera: note on its presumed identity with Stauferia. H. delecta, Meig., = (S. diaphana, Br. & Berg.); Mik, p. 55, Wien. ent. Z. xi, also Brauer & Bergenstamm, t. c. p. 108, and Mik, t. c. pp. 110-113.
- Homalomyia corrina, Wales, kowarsi, England, VERRALL, p. 149, Ent. M. M. xxviii; H. hirticeps, p. 70, barbata, p. 73, latipalpis, p. 75, Centr. Europe, STEIN, Wien. ent. Z. xi: n. spp.
- Hyadesimyia clausa, referred to Tachinida, and H. sarcophagoides to the genus Zelleria; BIGOT, p. 54, Wien. ent. Z. xi.
- Hylemyia coronata, & described; BIDENKAP, Ent. Tidakr. xiii, p. 244.
- Hyphantrophaga, n. g. for Meigenia hyphantria, Towns.; Townsend, p. 247, Psyche, vi.
- Hystricia aldrichi, Dakota: Townsend, p. 91, Tr. Am. Ent. Soc. xix, n. sp.

- Idia fasciata, destroying locusts in Algeria; BRONGNIART, Le Nat. 1892, p. 140.
- Lachnomma, n. g., p. 103, Tachinina, for L. magnicornis, n. sp., New Mexico, p. 104; Townsend, Tr. Am. Ent. Soc. xix.
- Leptoda elegans, Java, Wulp, p. 207, pl. xii, figg. 11 & 12, Tijdschr. Ent xxxiv, n. sp.
- Leucostoma neomezicana, New Mexico, Townsend, p. 169, Canad Ent xxiv, n. sp.
- Limnophora zetterstedti, Norway, BIDENKAP, p. 245, Ent. Tidskr. xiii n. sp.
- Lispa, monograph of the European species; Kowarz, Wien. ent. Z. zi pp. 33-54.
- L. leucomelæna, Sarepta, candicans, Ægina, p. 36, convexiuscula, Sicily p. 48, Kowarz, Wien. ent. Z. xi, n. spp.
- Locuoia globosa, Florida, Townsend, p. 129, Ent. News, iii; L. restcorni nigrifrons, N. America, Townsend, Canad. Ent. xxiv, p. 77: n. spi
- Lucilia sylvarum attacking toads; Mortensen, Zool. Anz. xv, p. 193. Lyperosia minuta, Somaliland; Bezzi, p. 192, Ann. Mus. Genov. (2) xi
- Macquartia johnsoni, N. America, Townsend, p. 81, Canad. Ent. xxiv
- Macropalpus, n. g. for Exerista villica, Zett.; MEUNIER, Bull. Soc. En Fr. lxi, p. ceviii.
- Masicera tenthredinidarum, p. 285, sphingivora, eufitchia, p. 286, datamum p. 287, N. America, Townsend, Tr. Am. Ent. Soc. xix; M. prote parcis, Jamaica, Townsend, p. 70, J. Inst. Jamaica, i : n. spp.
- Masistylum, n. g., Pyrrhosiides, for Pachystylum arcuatum, Mik; BRAUEI p. 595, SB. Ak. Wien, ci.
- Medoria and Morinia, note on; RÖDER, p. 374, Ent. Nachr. xviii.
- Megaprosopus michiganensis, N. America; Townsend, p. 111, Tr. At Ent. Soc. xix.
- Meigenia websteri, referred to Prospherysa, Wulp; TOWNSEND, p. 25 Pysche, vi.
- Metopia luggeri, Minnesota, Townsend, p. 69, Canad. Ent. xxiv, n. sp. Microtachina, n. g. for Tachina nympharum and erucarum, Rond.: Mil p. 116, Wien. ent. Z. xi.
- Miltogramma decisa, Florida, Townsend, p. 81, Ent. News, iii; M. kasensis, Kansas, Townsend, p. 68, Canad. Ent. xxiv: n. spp.
- Musca corvina, n. var., somalorum; Bezzi, p. 190, Ann. Mus. Genov. (1xii. M. vomitoria, embryology; Henking, Z. wiss. Zool. liv, p. 171
- Muscopteryx, n. g., Phytoinæ, p. 170, for M. chætosula, n. sp., Chihuaha p. 171; Townsend, Canad. Ent. xxiv.
- Myothyria vanderwulpi, Florida, Townsend, p. 131, Ent. News, iii, n. sp. Neotractocera, n. g, Tachinina, p. 105, for N. anomala, n. sp., Ne Mexico, p. 106; Townsend, Tr. Am. Ent. Soc. xix.
- Olenochata, n. g., Phoroceratine, p. 114, for O. kansensis, n. sp, 1 America; Townsend, p. 115, Tr. Am. Ent. Soc. xix.

- Pachyophthalmus floridensis, Florida, Townsend, Ent. News, iii, p. 80, n. sp.
- Pachystylum and Pseudopachystylum, and their species discussed, BRAUER, pp. 593-607, SB. Ak. Wien, ci; Pachystylum, note on the characters and composition of, and on Brauer's views thereon, Mik, pp. 245-248, Wien. ent. Z. xi; P. arcuatum, systematic position, Mik, p. 182, t. c.
- Phorbia seneciella, Centr. Europe; MEADE, p. cxvi, Bull. Soc. Ent. Fr. lxi. Phorocera lophyri, Canada, Townsend, p. 289, Tr. Am. Ent. Soc. xix, n. sp.
- Phyto senilis, New York, TOWNSEND, p. 81, Canad. Ent. xxiv; P. nigricornis, New Mexico, Townsend, p. 170, t. c.: n. spp.
- Phytomyptera unicolor, abdominal macrochætæ; MIK, p. 182, Wien. ent. Z. xi.
- Plagia aurifrons, Pennsylvania, Townsend, p. 67, Canad. Ent. xxiv, n. sp.
- Plagiprospherysa, n. g. near Phorocera, for P. valida, p. 113, New Mexico, floridensis, Florida, p. 114, n. spp.; Townsend, Tr. Am. Ent. Soc. xix.
- Polidea americana, Michigan, Townsend, p. 82, Canad. Ent. xxiv, n. sp. Pseudatractocera, n. g., Tuchinina, p. 107, for P. neomexicana, n. sp., New Mexico; Townsend, Tr. Am. Ent. Soc. xix.
- Pseudogonia ruficauda, Dakota, Townsend, p. 66, Canad. Ent. xxiv, n. sp.
- Pseudohystricia exilis, Jamaica, Townsend, p. 146, Ent. News, iii, n. sp. Pseudomyothyria, n. g., Phytoinæ, p. 131, for P. indecisa, n. sp., Illinois, p. 132; Townsend, Tr. Am. Ent. Soc. xix.
- Rhinophora valida, p. 167, mexicana, p. 168, New Mexico, Townsend, Canad. Ent. xxiv, n. spp.
- Sarcoclista, n. g., Phytoinæ, p. 122, for S. dakotensis, n. sp., N. America, p. 123; TOWNSEND, Tr. Am. Ent. Soc. xix.
- Sarcodexia, n. g., Dexiidæ, p. 105, for S. sternodontis, n. sp., Jamaica, p. 106; TOWNSEND, J. Inst. Jamaics, i.
- Sarcomacronychia, n. g., Tachinina, p. 100, for S. unica, n. sp., New Mexico, p. 101; TOWNSEND, Tr. Am. Ent. Soc. xix.
  - S. sarcophagoides, New Mexico, Townsend, p. 165, Canad. Ent. xxiv, n. sp.
  - Sarcophagida: synopsis of the N. American genera; Townsend, pp. 279-283, Tr. Am. Ent. Soc. xix.
  - Sarcophaga cimbicis, Dakota, TOWNSEND, p. 126, Canad. Ent. xxiv; S. helicis, Ohio, TOWNSEND, Psyche, vi, p. 220; S. opifera, N. America (destroying Melanoplus devastator), Coquillett, Ins. Life, v, p. 22, n. spp.
  - Sarcotachinella, n. g., Tachininæ, p. 110, for S. intermedia, n. sp., Illinois, p. 111; TOWNSEND, Tr. Am. Ent. Soc. xix.
  - Saundersia rubicunda, New Grenada, Wulp, p. 206, Tijdschr. Ent. xxxiv, n. sp.

Siphoclytia, n. g., Thryptoceratina, p. 116, for S. robertsoni, n. sp., Florida, p. 117; TOWNSEND, Tr. Am. Ent. Soc. xix.

Siphophyto, n. g., Phytoine, p. 127, for S. foridensis, neomezicanus, n. spp. N. America, p. 128; Townsend, Tr. Am. Ent. Soc. xix.

Synamphoneura africana, Somaliland, Bezzi, p. 190, Ann. Mus. Genov. (2) xii, n. sp.

Tuchina: application of the name and synonymy; BRAUER, pp. 602-606, SB. Ak. Wien, ci.

T. orgyiæ, p. 284, tenthredinivora, p. 285, N. America, Townsend, T. Am. Ent. Soc. xix, n. spp.

Tachinomyia, n. g., Tachinine, for T. robusta, Michigan, p. 96, foridensis, Florida, p. 97, n. spp.; TOWNSEND, Tr. Am. Ent. Soc. xix.

Tachinophyto, n. g., Phytoinas, p. 130, for T. floridensis, n. sp., Floride, p. 131; TOWNSEND, Tr. Am. Ent. Soc. xix.

Thryptocera americana, Washington, Townsend, p. 69, Canad. Ent. xxiv, n. sp.

Trixa gillettei, Colorado, Townsend, p. 68, Canad. Ent. xxiv, n. sp.

Trixoclista, n. g., Tachinina, p. 102, for T. distincta, n. sp., Illinois, p. 103;
TOWNSEND, Tr. Am. Ent. Soc. xix.

Tryphera americana, p. 78, polidoides, p. 79, N. America; Townsend, Canad. Ent. xxiv, n. spp.

Vanderwulpia sequens, New Mexico, Townsend, p. 172, Canad. Ent. xxiv, n. sp.

The following are the new forms described by Brauer & Bergenstamm in Denk. Ak. Wien, lviii, as mentioned on p. 294.

Macromeigenia, n. g. for Chrysoprocta, Wd.; Brauer & Bergenstamm, p. 311, Denk. Ak. Wien, lviii.

Cryptomeigenia, n. g. for C. setifacies, n. sp., Brazil, p. 311.

Pseudoviviania, n. g., p. 311, for P. platypoda, n. sp., Venezuela, p. 312.

Masiphya, n. g. for M. brasiliana, n. sp., Ypanema, p. 313.

Alsopsyche, n. g. for A. nemoralis, n. sp., Venezuela, p. 313.

Conogaster, n. g. for Viviania nubilis, Rond, p. 314.

Megalochata brachystoma, Austria, p. 317, n. sp.

Chatomyia, n. g. for Exorista crassiseta, Rond, p. 317.

Dexodes machairopsis, eggeri, auripilus, Europe, p. 316, n. spp.

Parexorista raiblensis, tultschensis, intermedia, setosa, antennata, acrochata, p. 318, clavellaria, magnicornis, p. 320, brevifrons, p. 321, gnava, dubia, blepharipoda, p. 322, grossa, trifasciata, irregularis, p. 323 latifrons, p. 324, Europe, n. spp.

Alsomyia, n. g. for A. gymnodiscus, n. sp., Austria, p. 328.

Paratryphera, n. g. for P. handlirschii, n. sp., Tyrol, p. 328.

Catachæta, n. g. for C. depressariæ, n. sp., Austria, p. 329.

Blepharidopsis, n. g. for B. nemea, Meig., p. 329.

Bactromyia, n. g. for Masicera scutelligera, Stein, p. 329.

Pexomyia, n. g. for Ræselia aberrans, Egg., p. 329.

Ceromasia ruspes, wulpii, Europe, p. 330, n. spp.

Leptotachina, n. g. for Ceromasia gratiosa, Stein., p. 330.

Thelymyia, n. g. for T. lowii, n. sp., Bohemia, p. 330.

Myxexorista, n. g. for M. pexops, flavipalpis, roseanæ, n. spp., Europe, and others, p. 332.

Achatoneura, n. g. for A. hesperus, n. sp., N. America, and others from Mexico, p. 334.

Tachinoptera, n. g. for T. eggeri, n. sp., Sicily, p. 336.

Chatogadia, n. g. for part of Prospherysa, Wulp, pp. 336.

Gadiopsis, n. subg. of Chatogadia, for G. mexicana, n. sp., Mexico, p. 336.

Parahypochata, n. g. for P. heteroneura, n. sp., N. America, p. 337.

Paralispe, n. g. for P. brasiliana, n. sp., Brazil, p. 337.

Paradoria, n. g. for P. nigra, n. sp., Venezuela, p. 339.

Neomintho, n. g. for macilenta, Wd., &c, p. 339.

Parasetigena, n. g., p. 339 (no species).

Chætoprocta, n. g., p. 341, for C. tarsalis, n. sp., Venezuela, p. 342.

Ctenophorocera, n. g., pp. 339 & 342, for C. blepharipus, n. sp., Cape or Brazil?, p. 42.

Xylotachina, n. g., p. 342, for X. ligniperdæ, n. sp., Europe, p. 343.

Argyrophylaz, n. g., p. 343, for A. galii, pelmatoprocta, Europe, idonea, Cape Good Hope, n. spp., p. 344, and others.

Sisyropa augusta, Austria, p. 345, ingens, Prachatitz, cinerea, Rockhampton, rufiventris, p. 346, prosopinu, leptotrichopa, p. 347, Brazil, n. spp.

Bolomyia, n. g. for Mystacella violacea, Wulp, p. 347.

Catagonia, n. g. for Exorista nemestrina, Egg., p. 348.

Anagonia, n. g. for A. spylosioides, n. sp., Tasmania, p. 348.

Thysanomyia, n. g. for Brachycoma fimbriata, Wulp, p. 340.

Mesochæta, n. g. for Didyma connexa, Wulp, p. 341.

Paramesochæta, n. g. for Mystacella fuscicostalis, Wulp, p. 341.

Masipoda, n. g. for M. geminata, n. sp., Brazil, p. 341.

Anamastax, n. g. for Blepharipesa goniæformis, Mcq., p. 349.

Paragædia, n. g. for P. hedemanni, n. sp., Mexico, p. 350.

Ptilotachina, n. g. for Tachina civilis, Rond., p. 350.

Podotachina, n. g. for P. americana and vibriseata, n. spp., New York, p. 350.

Tetragrapha, n. g. for T. tessellata, n. sp., Cuba, p. 351.

Hypotachina, n. g. for H. disparata, n. sp., Brazil, p. 352.

Pseudogermaria, n. g. for P. georgiæ, n. sp., N. America, p. 352.

Chephaliodes, n. g. for C. perversus, n. sp., Russia, p. 353.

Cnephalia bisetosa, n. n. for C. bucephala, Schin. pars, p. 353.

Paraplagia, n. g. for part of Plagia, p. 354.

Goniocera, n. g. for G. schistacea, n. sp., Europe, p. 354.

Glaucophana, n. g. for G. amasiæ, n. sp., Asia Minor, p. 354.

Paraneara, n. g. for P. longicornis, n. sp., Askold I., p. 355.

Microphana, n. g. for M. minuta, n. sp., Central Europe, p. 355.

Nemorilloides, n.g., p. 355, for N. flaviventris, n. sp., Cape of Good Hope, p. 356.

Petinops, n. g. for P. schnablii, n. sp., Poland, p. 356.



bifasciata, p. 359, stelviana, E. Europe, fulva, Bordeau n. spp.

Erythronychia, n. g. for Demoticus australensis, S., p. 360. Arrenopus, n. g. for A. americanus, n. sp., Georgia, p. 361.

Selenomyia, n. g. for S. brevicornis (Phil.), n. sp., Chili, p. 361.

Nemore a tropidobothra, Java, p. 361, n. sp.

Myiophasia, n. g. for M. anea, Wd., p. 362.

Thorarites, n. g. for Musca abdominalis, Wd., p. 363. Myzodezia, n. n. for (Tropidomyia, Br. Berg.), p. 363.

Myxodexia, n. n. for (Tropidomyia, Br. Berg.), p. 36. Myiomima brasiliana, Brazil, p. 364, n. sp.

Gymnodexia, n.g. for Sirostoma trianguliferum, Zett., and G. ste Hungary, p. 364.

Prorhynchops, n. g. for P. bilimeki, n. sp., Mexico, p. 364. Styloneuria, n. g. for S. manni, n. sp., Ragusa, p. 365.

Sarothromyia, n. g. for Sarcophila femoralis, S., p. 365. Tephromyia, n. g. for Sarcophila grisea, Mg., p. 366.

Paramorinia, n. g. for P. cincta, n. sp., Tyrol, p. 367.

Leucomyia, n. g. for Sarcophila alba, Schin., p. 368.

Ptilozeuxia brachycera, n. n. for (P. brevicornis, Egg.), p. 368.

Erythrandra, n. g. for E. picipes, n. sp., N. America, p. 368.

Oxytachina, n. g. for Heteronychia vittata, Wd., p. 369.

Mochlosoma simonyi, Canary Is., n. sp., p. 370.

Prosenoides, n. g. for Prosena papilio, S., p. 370.

Dezionhuna, n. y. for Prosenherusa amulans. Wnl

Dexiophuna, n. g. for Prospherysa æmulans, Wulp, p. 374.

Gymnostylia ornata, Venezuela, p. 374, n. sp.

Eufischeria, n. g. for E. ceylanica, n. sp., Ceylon, p. 374.

Ptilodegeeria, n. g. for Hypostena obumbrata, Wulp, p. 375.

Prosheliomyia, n. g. for P. nietneri, n. sp., Ceylon, p. 375.

Minthodexia, n. g. for M. gravipes, flavicornis, n. spp., Venezueli

Zosteromyia, n. g. for Myobia cingulata, Moq., p. 376.

Preudodexia, n. g. for Dexia eques, Wd., p. 378.

Paragymnomma, n. g. for P. hystrix and diaphana, n. spp., Brazil, p. 384. Plagiomima, n. g. for P. disparata, n. sp., Mexico, p. 384.

Seciophaga, n. g. for Myobia glirina, Rdi., p. 384.

Micromyobia, n. g. for Myobia montana, Schin., p. 385.

Chatodemoticus, n. g. for Demoticus chilensis, Schin., p. 385.

Aneogmena, n. g. for A. fischeri, n. sp., Agra, p. 385.

Peteina stylata, Greenland, n. sp., p. 387.

Stylogymnomyia, n. g. for Gymnosoma nitens, Mg., p. 387.

Mormonomyia, n. g. for laniventris, Wd., ? n. sp., p. 388.

Epineura, n. g. for Phasia helva and taniata, Wd., p. 388.

Elioseta americana, Georgia, n. sp., p. 388.

Paramphibolia, n. g. for Rutilia assimilis, Mcq., p. 389.

Microcerella steinduchneri, Galapagos Is., n. sp., p. 389.

Thelychæta, n. g. for T. chalybea, n. sp., Borneo, p. 390.

Arrhinidia, n. g. for Rhynchomyia aberrans, Sch., p. 390.

Phasiophana, n. g. for Musca obsoleta, Wd., p. 390.

Pararicia, n. g. for a part of Cyrtoneura, p. 391.

Neocalliphora, n. g. for Call. dasyophthalma, Mcq., and ochracea, S., p. 391.

Paralucilia, n. g. for Calliph. fulvipes, Blanch., p. 391.

Zonochroa, n. g. for Ochromyia euzona, Löw, p. 391.

Aucheromyia, n. g. for Ochromyia luteola, Fbr., p. 391, Brauer & Bergenstamm, Denk. Ak. Wien, lviii.

#### MUSCIDE ACALYPTRATE.

Agromyza lappæ, metamorphoses; LAMPA, Ent. Tidskr. xiii, pp. 117-120.

Cacoxenus indagator, characters of; Mik, p. 182, Wien. ent. Z. xi: chartotaxy; Osten-Sacken, Berl. Ent. Zeit. xxxvi, p. 311.

Chlorops pumilionis, life-history; LAMPA, Ent. Tidskr. xiii, p. 257, pl. vii. Calopa aquatorialis, Somaliland, Bezzi, p. 194, Ann. Mus. Genov. (2) xii, n. sp.

Drosophila lacteo-guttata, W. Russia, Portschinsky, p. 226, pl. i, fig. 20, Hor. Ent. Ross. xxvi; D. nigropunctata, Java, Wulp, p. 216, Tijdschr. Ent. xxxiv: n. spp.

Ephydra dorsopunctata, Wied., referred to Notiphila; WULP, p. 215, Tijdschr. Ent. xxxiv.

Eurosta solidaginis: galls and parasites noticed; Brodle, p. 137, Canad. Ent. xxiv.

Homoneura, n. g., Sapromyzarum, p. 213, for H. picea, n. sp., Java, p. 214; Wulp, Tijdschr. Ent. xxxiv.

Hypenidium gracum: note on, with fig. of wing; PORTSCHINSKY, p. 215, pl. i, fig. 13, Hor. Ent. Ross. xxvi.

H. sibiricum, Amurland, Portschinsky, p. 216, pl. i, fig. 14, Hor. Ent. Ross. xxvi, n. sp.

Lagarosia, n. g., Ortalinarum, p. 210, for L. lacteata, p. 211, striatella, p. 213, n. spp., Java; Wulp, Tijdschr. Ent. xxxiv.

Lamprophthalma, n. g., near Lazocera, p. 225, for L. metallica, n. sp.. Persia, p. 226, pl. i, fig. 7; PORTSCHINSKY, Hor. Ent. Rosa xxvi.

Macrotrypeta, n. g., p. 223, for M. ortalidina, n. sp., Siberia, p. 224, pl. i, fig. 11; PORTSCHINSKY, Hor. Ent. Ross. xxvi.

Myennie sibirica, Siberia, Portechinery, p. 213, pl. i, fig. 10, Hor. Ent. Ross. xxvi, n. sp.

Neottiophilum: characters and systematic position discussed; RODER, p. 204, Ent. Nachr. xviii.

Opomyza punctella, Fall.: variation and characters noticed; SINTENIS, p. 177. Wien, ent. Z. xi.

Paralimna javana, Java, WULP, p. 215, pl. xii, fig. 17, Tijdschr. Ent. xxxiv, n. sp.

Platyparea? terebratula, E. Siberia, PORTSCHINSKY, p. 216, pl. i, fig. 9, Hor. Ent. Ross. xxvi, n. sp.

Ptilona amurensis, E. Siberia, Portschinsky, p. 214, pl. i, fig. 12, Hor. Ent. Ross. xxvi, n. sp.

Pyrgota microcera, Amur, Portschinsky, p. 212, pl. i, fig. 8, Hor. Ent. Ross. xxvi, n. sp.

Sapromysa: characters of the N. American; Townsend, pp. 301-303, Canad. Ent. xxiv.

S. ocellaris, New Mexico, Townsend, p. 303, Canad. Ent. xxiv; S. mikii, p. 155, styriaca, p. 156, Austrian Alps, Strobl, Wien. ent. Z. xi: n. spp.

Spilographa armifrons, p. 221, pl. i, figs. 16 & 17, bipunctata, p. 222, fig. 15.
E. Siberia, Portschinsky, Hor. Ent. Ross. xxvi, n. spp.

Tephritis tatarica, Turkestan, p. 218, pl. i, fig. 19, angulato fasciuta. Persia, p. 220, fig. 18, PORTSCHINSKY, Hor. Ent. Ross. xxvi.

Timia testacea, Central Asia, Portschinsky, p. 211, Hor. Ent. Ross xvi, n. sp.

## PHORIDE.

Phora: remarks on characters, habits, and variation of the Austrian species; STROBL, pp. 193-204, Wien. ent. Z. xi.

Phora crassicosta, p. 194, srythronota, p. 195, spinosissima, p. 196, pseudo concinna, p. 199, nitidifrons, p. 200, xanthosona, p. 203, Austria Strobl, Wien. ent. Z. xi; P. cimbicis, p. 143, setacea, p. 144, luggeri p. 145, minuta, p. 146, Dakota, Aldrich, Canad. Ent. xxiv P. omnivora, New Zealand, Hudson, Manual, p. 62, pl. vii, fig. 15 n. spp.

#### PUPIPARA.

The mouth parts of various *Pupipara*, including *Braula* and *Nycteribia* described; Müggenburg, Arch. f. Nat. lviii, i, pp. 287-332 pls. xv & xvi.

Trichobius dugesii, note on ; TOWNSEND, p. 177, Ent. News, iii.

# (E.) RHYNCHOTA.

# HEMIPTERA-HETEROPTERA.

[Cf. Ashmead (18a), Berg (56), Bergroth (62-72), Carlini (158), Carpenter (159), Distant (210, 217, 218, 347), Dugès (229), Febrari (281), Forbes (286), Gerstaecker (323), Hridemann (392), Horváth (428-431), Hudson (440), Jakowleff (452), Karsch (460, 465), Léon (545), Moberg (614), Montandon (617-620), Puton (707, 708), Reuter (730-733), Schlechtendal (800), Sharp (842), Tryon (952).]

Homologies of the Hemipterous mouth, SMITH (866); and Homology of the trophi, Léon (545).

Silurian fossil Hemipterous insect, Moberg (614).

Transvaal: n. spp.

BERGROTH, Rev. d'Ent. xi, pp. 262-264, records many synonyms in view of the promised general Catalogue of *Hemiptera*; these it appears unnecessary for us to give in detail.

Hemidiptera haeckeli, Léon, note on ; BERGROTH, p. 169, Wien. ent Z. xi.

#### PENTATOMIDÆ.

- Alcimocoris potanini, China, Horváth, p. 134, Term füzetek, xv. n. sp. Antestia guttifera, E. Africa, Gerstaecker, p. 46, JB. Hamb. Anst. ix; A. transvaalia, Transvaal, DISTANT, p. 251, pl. iii, fig. 6, Naturalist
- Aphanopneuma stali, Africa, = (biloba, Sign., Stal); Montandon, p. 297, Rev. d'Ent. xi, n. sp.
- Apotomogonius, n. g., Plataspidinæ, p. 301, for A. exornatus, n. sp., Centr. Africa, p. 302; MONTANDON, Rev. d'Ent. xi.
- Arma prasina, New Guinea, TRYON, p. 15, Ann. Queensland Mus. ii, n. sp. Asopus figuratus, var. described and figured; DISTANT, p. 248, pl. iii, fig. 1, Naturalist Transvaal.
- A. glaucolimbata, New Guinea, TRYON, p. 15, Ann. Queensland Mus. ii, n. sp.
- Aspongopus balinus, W. Africa, Karsch, p. 166, Ent. Nachr. xviii, n. sp. Basicryptus elongatus, Transvaal, antennatus, E. Africa, DISTANT, p. 239, Ent. M. M. xxviii, n. spp.
- Brachyplatys flavosparsus, Borneo, BERGROTH, p. 171, Wien. ent. Z. zi; B. simplex, pusilla, N. Guinea, TRYON, p. 13, Ann. Queenaland Mus. ii : n. spp.

- Cintharodes nubilosus, Cougo, Montandon, p. 299, Rev. d'Ent. xi, n. sp. Cirpocoris alienus, E. Siberia, Reuter, p. 176, Œfv. Finska. Förh. xxxiii, n. sp.
- Caura superba, W. Africa, KARSCH, p. 164, Eut. Nachr. xviii, n. sp.
- Cermatulus pulcher, New Guinea, TRYON, p. 15, Ann. Queensland Mus. ii, n. sp.
- Chrysocoris nicobarensis, Nicobar Is., DISTANT, Ent. xxv (Supp.), p. 96, n. sp.
- Coptosoma atenes, W. Africa, KARSCH, p. 129, Ent. Nachr. xviii. C. sculpturatum, Madagascar, p. 278, nebulosum, p. 279, confusum, p. 280, E. Africa, parvipictum, p. 281, sordidulum, p. 283, China, lethierryi, E. India, p. 284, Montandon, Rev. d'Ent. xi. C. respersum, Java, p. 171, mundum, p. 172, lascicum, p. 173, China, BERGROTH, Wien. ent. Z. xi: n. spp.
- Derula delagrangei, Syria, Puton, p. 25, Rev. d'Ent. xi, n. sp.
- Dictyotus transversus, Murray I., CARPENTER, p. 138, pl. xii, fig. 1, P. B. Dublin Soc. (n.s.) vii, n. sp.
- Dolycoris rutherfordi, Old Calabar, DISTANT, p. 187, Ent. M. M. xxviii, n. sp.
- Edessa herrichi, p. 45, rugiventris, p. 48, Argentine Rep., BERG, An. 80c. Arg. xxxiii, n. spp.
- Ennius monteironis, Delagoa Bay, DISTANT, p. 188, Ent. M. M. xxviii, n. sp.
- Eysarcoris megaspiloides, New Guinea, TRYON, p. 17, Ann. Queensland Mus. ii, n. sp.
- Gonocerus juniperi, n. var., obtusangulus, PUTON, p. 26, Rev. d'Ent. xi.
- Halyomorpha bimaculata, Sierre Leone, BERGROTH, p. 160, Ann. Ent. Belg. xxxvi. H. capitata, p. 249, pl. iii, fig. 3, pretoriæ, p. 250, fig. 4, Transvaal, DISTANT, Naturalist Transvaal: n. spp.
- Handhirschiella, n. g., Plataspidina, p. 294, for H. anea, p. 295, emarginata, p. 296, n. spp., Africa; Montandon, Rev. d'Ent. xi.
- Holcostethus distanti, S. Africa, BERGROTH, p. 160, Ann. Ent. Belg. xxxvi. H. obscuratus, Transvaal, Distant, p. 249, pl. iii, fig. 2, Naturalist Transvaal: n. spp.
- Hymenomaga, n. g., near Tropicorypha, for T. formosa, Dist., KARSCH, p. 130, Ent. Nachr. xviii.
- Isoplatys, n. g., Plataspidinα, for I. flavonotatus, n. sp., Gaboon; Montandon, p. 303, Rev. d'Ent. xi.
- Lyramorpha maculifer, New Guinea, TRYON, p. 19, Ann. Queensland Mus. ii, n. sp.
- Megymenum severini, Bengal, BERGROTH, p. 504, Ann. Ent. Belg. xxxvi, n. sp.
- Menida quadrimaculata, Mongolia, Horvath, p. 134, Term. füzetek, xv, n. sp.
- Nezara abnormis, Buenos Aires, p. 7, vicina, p. 8, fucosa, p. 9, Argentine Rep., Berg, An. Soc. Arg. xxxiii, n. spp.
- Odontotarsus horvathi, N. Africa, Puton, p. 24, Rev. d'Ent. xi, n. sp.

- Orthoschizops plagosa, S. Africa, DISTANT, p. 238, Ent. M. M. xxviii, n. sp. Patanocnema, n. g., near Aspongopus, for P. ovata, n. sp., W. Africa; KARSCH, p. 130, Ent. Nachr. xviii.
- Pelioderma, n. g. Plataspidinæ, for P. æthiops, n. sp., W. Africa, Венскотн, p. celv, Bull. Soc. Ent. Fr. lxi.
- Pentatoma (Cappea) pretoria, E. Africa, GERSTAECKER, JB. Hamb. Anst. ix, p. 46; P. submarginalis, p. 16, maculifer, prasina, p. 17, New Guinea, TRYON, Ann. Queensland Mus. ii: n. spp.
  - Peromatus sulcifer, Argentine Rep., BERG, p. 43, An. Soc. Arg. xxxiii, n. sp.
  - Philia area, Australia, DISTANT, Ent. xxv (Supp.), p. 96, n. sp.
  - Phimodera mongolica, E. Siberia, REUTER, p. 173, Œfv. Finska Förh. xxxiv, n. sp.
  - Piezosternum subulatum, egg figured; SHARP, pl. viii, fig. 9, Tr. E. Soc. 1892. Plataspis gambeyi, Gaboon, p. 273, plagifera, Guinea, p. 275, guttulata, Gaboon, p. 276, conspersa, Madagascar, p. 277, Montandon, Rev. d'Ent. xi, n. spp.
  - Platytatus, n. g. (Eustheninæ), p. 158, for P. ambiguus, n. sp., Madagascar, p. 159; Векскотн, Ent. M. M. xxviii.
  - Pododus striatus, S. Africa, DISTANT, p. 238, Ent. M. M. xxviii, n. sp.
- Pacilochroma balteata, E. India, DISTANT, Ent. xxv (Supp.), p. 96, n. sp.
- Ponsila severini, Centr. Africa, Montandon, p. 304, Rev. d'Ent. xi, n. sp. Schizops icterica, E. Africa, GERSTAECKER, p. 47, JB. Hamb. Anst. ix, n. sp.
- Schirus biguttatus, n. var. tibialis; Puton, p. 25, Rev. d'Ent. xi: n. var. concolor; Nickerl, p. 62, S. E. Z. liii.
- Sephela morosa, E. Africa, GERSTAECKER, p. 45, JB. Hamb. Aust. ix, n. sp.
- Solenosthedium madagascariensis, Madagascar, DISTANT, p. 187, Ent. M. M. xxviii, n. sp.
- Spudæus variabilis, New Guinea, TRYON, p. 18, Ann. Queensland Mus. ii, n. sp.
- Tesseratoma hornimani and allies, characters noticed; DISTANT, p. 188, Ent. M. M. xxviii.
- Tiarocoris luminatus, Malacca, Montandon, p. 307, Rev. d'Ent. xi, n. sp. Uhlunga, n. g. (Pentatominæ), for U. typicu, n. sp., Natal; DISTANT, p. 237, Ent. M. M. xxviii.
- Vaterna patula, Transvaal, DISTANT, p. 250, pl. iii, fig. 5, Naturalist Transvaal, n. sp.

#### COREIDÆ.

- Acanthocephala granulosa, Dall., = (luctuosa, Stål); DISTANT, p. 359, Biol. Centr. Am. Rhynch. Heter.
- Anasa montivaga, p. 366, pl. xxxiii, fig. 19, delibata, fig. 17, decretoria, fig. 16, versicolor, fig. 18, p. 367, Centr. America, DISTANT, Biol. Centr. Am. Rhynch. Heter.; A. limbata, p. 97, declivicollis, p. 153, Argentine Rep., Berg, An. Soc. Arg. xxxiii : n. spp.
  - 1892. [VOL. XXIX.]

- Archimerus chiriquinus, Panama, Distant, p. 355, pl. xxxiii, fig. 4, Biol. Centr. Am. Rhynch. Heter., n. sp.
- Athaumastus subfoveolatus, Argentine Bep., BERG, p. 66, An. Soc. Arg. xxxiii, n. sp.
- Burdistus serrulatus, figured, pl. xxxiii, fig. 7, Biol. Centr. Am. Rhynch. Heter.
- B. formidabilis. Panama, DISTANT, p. 357, pl. xxxiii, fig. 9, Biol. Centr. Am. Rhynch, Heter., n. sp
- Capaneus chontalensis, Nicaragua, pl. xxxiii, fig. 3, humerosus, fig. 5, Mexico, Distant, p. 354, Biol. Centr. Am. Rhynch. Heter., n. spp.
- Catorhintha sinuatipennis, Buenos Aires, Brag, p. 152, An. Soc. Arg. xxxiii, n. sp.
- Cebrenis rubro-conspersa, Argentine Rep., BERG, An. Soc. Arg. xxxiii, p. 99, n. sp.
- Chariesterus robustus, Mexico, pl. xxxiii, fig. 12, cuspidatus, Panama, fig 14, DISTANT, p. 364, Biol. Centr. Am. Rhynch. Heter., n. spp.
- Diactor bogotanus, figured, pl. xxxiii, fig. 8, Biol. Centr. Am. Rhynch. Heter.
- Dulichius, Stål, = (Formicoris, Kirb.); BERGROTH, p. 225, Wien. ent. Z.
- D. wroughtoni, E. India, BERGROTH, p. 107, Ent. M. M. xxviii [D. (sub Formicoris) inflatus, Kirby, = (wroughtoni), BERGROTH, t. c. p. 126], n. sp.
- Elusmornema, n. g., near Mictis, for E. limpidipensis, n. sp., W. Africa; KARSCH, p. 131, Ent. Nachr. xviii.
- Homeocerus vicarians, Cameroons, Karsch, p. 132, Ent. Nachr. xviii, n. sp.
- Hormambogaster, n. g, near Dalader, for H. expansus, n. sp., W. Africa; KARSCH, p. 131, Ent. Nachr. xviii.
- Hypselonotus bitrianguliger, Argentine Rep., BERG., p. 101, An. Soc. Arg. xxxiii, n. sp.
- Laminiceps quadrisignata, Central America, DISTANT, p. 360, pl. xxxiii, fig. 11, Biol. Centr. Am. Rhynch. Heter., n. sp.
- Leptoglossus dentatus, p. 68, impressicollis, p. 69, concaviusculus, p. 70, Argentine Rep., Berg, An. Soc. Arg. xxxiii, n. spp.
- Melucha quinquelineata, figured, pl. xxxiii, fig. 2, Biol. Centr. Am. Rhynch. Heter.
- M. bicolor, Guatemala, DISTANT, p. 353, pl. xxxiii, fig. 1, Biol. Centr. Am. Rhynch. Heter., n. sp.
- Mictis Iceta, E. Africa, GERSTARCKER, p. 48, JB. Hamb. Anst. ix,
- Myrmus calcaratus, E. Siberia, REUTER, p. 181, Œfv. Finska Förh. xxxiii. n. sp.
- Narnia inornata, Mexico, DISTANT, p. 361, pl. xxxiii, fig. 10, Biol. Centr. Am. Rhynch. Heter., n. sp.
- Ojedana, n. g., near Archimerus, p. 355, for O. loricata, n. sp., Panama p. 356, p. xxxiii, fig. 6, Distant, Biol. Centr. Am. Rhynch. Heter.

- Ovengua, n. g. (Daladeraria), for O. apertu, n. sp., Old Calabar: DISTANT, Ent. M. M. xxviii, p. 285.
- Plinachtus falcatus, Transvaal, Distant, p. 251, pl. iii, fig. 8, Naturalist Transvaal, n. sp.
- Prismatocerus (Philonus) hamatocerus, E. Africa, GERSTAECKER, p. 48, JB. Hamb. Anst. ix, n. sp.
- Rhopalus (Stictopleurus) nysioides, p. 178, R. (Æschynteles) robustus, p. 179, E. Siberia, Reuter, Œfv. Finska Förh. xxxiii, n. spp.
- Savius diagonalis, Corrientes, BERG, p. 102, An. Soc. Arg. xxxiii, n. sp. Sethenira uruguayensis, S. America, BERG, p. 71, An. Soc. Arg. xxxiii, n. sp.
- Spartocera gigantea, Costa Rica, DISTANT, p. 363, pl. xxxiii, fig. 13, Biol. Centr. Am. Rhynch. Heter., n. sp.
- Vilga mexicana, Omilteme, DISTANT, p. 368, pl. xxxiii, fig. 20, Biol. Centr. Am. Rhynch. Heter., n. sp.
- Zicca castanea, Argentine Rep., BERG, p. 98, An. Soc. Arg. xxxiii, n. sp.

# LYGÆIDÆ, PYBBHOCORIDÆ.

- Antilochus violaceus, Somaliland, Carlini, p. 532, Ann. Mus. Genov. (2) xii, n. sp.
- Beosus hoplites, E. Africa, GERSTAECKER, p. 49, JB. Hamb. Anst. ix, n. sp.
- Cenœus? exeanguis, Zanzibar, GERSTAECKER, p. 50, JB. Hamb. Anst. ix, n. sp.
- Dysdercus festivus, E. Africa, GERSTAECKER, p. 50, JB. Hamb. Anst. ix;
  D. melanoderes (Germ.), Cameroons, Karsch, p. 133, Ent. Nachr. xviii: n. spp.
- Geocoris putonianus, n. n. for G. picsus infrà; BERGROTH, Rev. d'Ent. xi, p. 264.
- G. henoni, Cairo, Puton, p. 26, Rev. d'Ent. xi; G. (Piocoris) piceus, Turkestan, Puton, p. 72, t. c.: n. spp.
- Henestaris irroratus, S. France, Horvath, p. 136, Rev. d'Ent. xi, n. sp. Ischnodemus atricolor, Uruguay, Berg, p. 158, An. Soc. Arg. xxxiii, n. sp.
- Ischnopeza pallipes, Biskra, Puton, p. 27, Rev. d'Ent. xi; I. flavogrisea, Somaliland, Carlini, p. 530, Ann. Mus. Genov. (2) xii: n. spp.
- Lamprodema rufipes, E. Siberia, REUTER, p. 182, Œfv. Finska Förh. xxxiii, n. sp.
- Lygaus (Melanocoryphus) heydeni, Turkestan, Puton, p. 71, Rev. d'Ent. xi; L. planitia, pl. iii, fig. 7, desertus, fig. 9, p. 252, campestris, p. 253, fig. 10, Transvaal, Distant, Naturalist Transvaal: n. spp.
- Myrmoplasta, n. g., Pyrrhocorides, for M. mira, n. sp., E. Africa, Gerstaecker, p. 51, JB. Hamb. Anst. ix.
- Nysius novitius, Transvaal, DISTANT, p. 254, pl. iii, fig. 11, Naturalist Transvaal; N. gibbifer, p. 155, ellipticus, p. 156, Arg. Rep., Berg, An. Soc. Arg. xxxiii: n. spp.

- Œ lancala (sub Pamera) dorsalis, Say, = (dorsilinea, Am. & Serv.); Berg, p. 160, An. Soc. Arg. xxxiii.
- Oncopeltus busanus, Cameroons, Karsch, p. 132, Ent. Nachr. xviii, n. sp. Pamera procerula, p. 162, sororcula, p. 163, Buenos Aires, Berg, An. Soc. Arg. xxxiii, n. spp.
- Peritrechus: characters of the Russian species; JAKOWLEFF, p. 228, Hor. Ent. Ross, xxvi.
- P. insignis, Derbent, JAEOWLEFF, p. 231, Hor. Ent. Ross. xxvi, n. sp. Phygadicus australis, Murray L., OARPENTER, p. 140, pl. xii, fig. 2, P. B. Dublin Soc. (n.s.) vii, n. sp.
- Pyrrhopeplus carduelis, n. var. posthumus, Horvátii, p. 135, Term. füzetek, xv.
- Rhyparochromus puncticollis, n. var. nigripes, Hobváth, p. 138, Rev. d'Ent. xi.
- Scolopostethus: characters of the palsarctic species, and synonymical list; HORVÁTH, pp. 253-256, Rev. d'Ent. xi.
- S. patruelis, S. France, HORVATH, p. 138, Rev. d'Ent. xi, n. sp.
- Sericocoris, n. g., p. 132 (Pyrrhocorides), for S. acromelanthes, n. sp., Cameroons, p. 133, Karsch, Ent. Nachr. xviii.
- Transvaalia, n. g., for T. lugens, n. sp., Transvaal, pl. iii, fig. 12; DISTANT, p. 253, Naturalist Transvaal.
- T. flarescens, Pretoria, DISTANT, Ent. M. M. xxviii, p. 285.

#### TINGITIDE.

- Campylostira: revision of species with characters and varr. described; HORVÁTH, pp. 309-313, Wien. ent. Z. xi.
- C. bornica, p. 309, Bosnia, eximia, Turkestan, p. 311, HORVÁTH, Wienent. Z. xi, n. spp.
- Cyclotynaspis, n. g., Cantacaderaria, for C. acalyptoides, n. sp., Singapore; Montandon, Rev. d'Ent. xi, p. 265.
- Dictyonota (Elina) henschi, Budapest, Puton, p. 72, Rev. d'Ent. zi, n. sp.
- Eurycera glabricornis, Mozambique, Montandon, p. 267, Rev. d'Ent. xi,
- Leptostyla carmelana, Uruguay, BERG, p. 200, An. Soc. Arg. xxxiv, n. sp.

#### ARADIDÆ.

- Aneurus indicus, Burma, BERGROTH, p. 717, Ann. Mus. Genov. (2) xii, n. sp.
- Aradus similis, var., = (centriguttatus), p. 332, list of the N. American Aradidæ, pp. 335 & 336; BERGROTH, P. E. Soc. Wash. ii.
- A. mirabilis, Burma, Bergroth, p. 710, Ann. Mus. Genov. (2) xii; A. duzeei, Ontario, p. 333, concinnus, California, p. 337, Bergroth, P. E. Soc. Wash. ii: n. spp.
- Artabanus excelsus, &, with stridulating area, described; BERGROTH, p. 806, Ann. Mus. Genov. (2) xii.

- A. excelsus, setulosus, p. 713, sexspinosus, p. 714, Burma, BERGROTH, Ann. Mus. Genov. (2) xii, n. spp.
- Brachyrrhynchus verruciger, Amurland, BERGROTH, p. 261, Rev. d'Ent. xi; B. undulatus, Bengal, BERGROTH, p. 504, Ann. Ent. Belg. xxxvi; B. tenericornis, Burma, BERGROTH, p. 716, Ann. Mus. Genov. (2) xii: n. spp.
- Carventus gestroi, Burma, BERGROTH, p. 712, Ann. Mus. Genov. (2) xii, n. sp.
- Lobocura, n. g. for L. oblonga, Argentine Rep., p. 259, orata, Mexico, p. 260, n. spp.; Bergroth, Rev. d'Ent. xi.
- Neuroctenus debilicornis, Gaboon, Montandon, p. 268, Rev. d'Ent. xi, n. sp.
- Pictinus asiaticus, Burma, BERGROTH, p. 716, Ann. Mus. Genov. (2) xii; P. modiglianii, pusio, Engano, BERGROTH, p. 807, t. c.: n. spp.

#### HYDROMETRIDÆ.

- Gerris (Tenagogonus) hypoleuca, E. Africa, GERSTAECKER, p. 55, JB. Hamb. Anst. ix, n. sp.
- Halobates regalis, Mabuiag, CARPENTER, p. 144, pl. xiii, figg. 1-8, P. R. Dublin Soc. (n.s.) vii, n. sp.
- Hermatobates, n. g., p. 142, for H. haddonii, n. sp., Mabuiag, p. 143, pl. xii, figg. 4-8; CARPENTER, P. B. Dublin Soc. (n s.) vii.
- Limnometra lineata, Mabuiag, CARPENTER, p. 141, pl. xii, fig. 3, P. R. Dublin Soc. (n.s.) vii, n. sp.
- Tenagogonus, subg. (n.?) of Gerris; GERSTAECKER, p. 56, JB. Hamb. Aust. ix.

#### REDUVIDE.

- Eggs of a Reduviid described and figured with micropyle (?) apparatus; Sharp, Tr. E. Soc. 1892, p. 191, pls. viii & ix.
- Acanthaspis armata, Somalilaud, CARLINI, 'p. 535, Ann. Mus. Genov. (2) xii, n. sp.
- Cleptria bicolor, S. Africa, BERGROTH, p. 162, Ann. Ent. Belg. xxxvi, n. sp.
- Coranus longiceps, S. Africa, BERGROTH, p. 161, Ann. Ent. Belg. xxxvi; C. hammarstroemii, Mongolia, REUTER, p. 184, Œfv Finska Förh. xxxiii; C. lóczyi, Calcutta, Horváth, p. 135, Term. füzetek. xv: n. spp.
- Ectrichodia problematica, E. Africa, GERSTAECKER, p. 54, JB. Hamb. Anst. ix, n. sp.
- Ectrychotes jakowleffi, China, Horváth, p. 136, Term. füzetek. xv. n. sp. Endochus africanus, W. Africa, BERGROTH, p. 161, Ann. Ent. Belg. xxxvi, n. sp.
- Enicocephalus: note on, and on Hymenodectes, Uhl.; ASHMEAD, p. 329, P. E. Soc. Wash. ii.

- Formicoris, Kirby, is the same as Dulichius; BERGROTH, Ent. M. M. xxviii, p. 126.
- Henicocephalus, = (Dicephalus, Kirb.); BERGROTH, p. 225, Wien. ent. Z. xi.
- Holotrichius henoni, Sues, Puton, p. 27, Rev. d'Ent. xi, n. sp.
- Ischnobæna preussi, W. Africa, KARSCH, p. 136, Ent. Nachr. xviii, n. sp. Laphyctes? brachialis, E. Africa, GERSTAECKER, p. 52, JB. Hamb. Aust. ix, n. sp.
- Leptodema, n. g. for L. kirtum, p. 533, acanthocephalum, p. 534, n. spp., Trop. Africa; Carlini, Ann. Mus. Genov. (2) xii.
- Oncocephalus insignis, E. Africa, GERSTAECKER, p. 55, JB. Hamb. Anst. ix; O. vaulogeri, Algeria, MONTANDON, p. 272, Rev. d'Ent. xi: n. spp.
- Maranaspis, n. g. Ectrichodiides, for M. typhlops, n. sp., Cameroons; Karsch, p. 134, Ent. Nachr. xviii.
- Miomerocerus, n. g. Ectrichodiides, for M. scopaceus, n. sp., Cameroons; Karsch, p. 135, Ent. Nachr. xviii.
- Nagusta junodi, Mosambique, Montandon, p. 269, Rev. d'Ent. xi, n. sp. Phonoctonus principalis, E. Africa, Gerstarcker, p. 52, JB. Hamb. Anst. xi, n. sp.
- Pirutes brachypterus, p. 54, morosus, collaris, p. 55, E. Africa, Gerstaecker, JB. Hamb. Anst. ix; P. conspurcatus, Transval, DISTANT, p. 255, pl. ii, fig. 10, Naturalist Transval: n. spp.
- Ploiaria melanacantha, Carcassonne, Horváth, p. 139, Rev. d'Eut. ri, n. sp.
- Polymazus, n. g., p. 52, for P. singularis, n. sp., E. Africa, p. 53; Gerstaecker, JB. Hamb. Anst. ix.
- Raphidosoma echinatum, E. Africa, GERSTAECKER, p. 54, JB. Hamb. Anst. ix, n. sp.
- Reduvius polystictus, E. Africa, GERSTAECKER, p. 53, JB. Hamb. Anst. ix; R. notabilipes, Obock, Montandon, p. 271, Rev. d'Ent. xi; R. pulvisculatus, pl. ii, fig. 3, sertus, fig. 8, p. 254, capitalis, p. 255, fig. 1, Transvaal, DISTANT, Naturalist Transvaal; n. spp.
- Rhochmogaster, n. g. Ectrichodiides, for R. dimerus, n. sp., Cameroons; KARSCH, p. 135, Ent. Nachr. xviii.
- Santosia semistriata, Cameroons, Karsch, p. 135, Eut. Nachr. xviii, n. sp. Sastrapada novæguinensis, Fergusson I., Tryon, p. 23, Ann. Queensland Mus. ii, n. sp.
- Sphedanolesthes fenestriculatus, W. Africa, KARSCH, p. 134, Ent. Nachr. xviii, n. sp.
- Vadimon bergrothi, Mozambique, Montandon, p. 270, Rev. d'Ent. xi, n. sp.

## SALDIDÆ, CIMICIDÆ.

- Acanthia: notes on the species of, in N. America; Osborn, p. 262, Canad. Ent. xxiv.
- A. inodora, Mexico, Duges, p. 169, pl. viii, n. sp.

- Anthocoris carinulatus, Madagascar, REUTER, p. 186, Ent. M. M. xxviii, n. sp.
- Xylocoris humeralis, Sign., referred to Lasiochilus, and redescribed; Reuter, p. 186, Eut. M. M. xxviii.

#### CAPSIDÆ.

Food-plants of Capsida, near Washington; Heidemann (392).

Antias lucidus, Argentine Rep., BERG, p. 95, An. Soc. Arg. xxxiii, n. sp.

Arsinotus, n. g. (Bryocoraria), for A. albipes, n. sp., Buenos Aires; Berg, p. 198, An. Soc. Arg. xxxiv.

Brachycoleus steini, n. var. thoracicus; Puton, p. 29, Rev. d'Ent. xi.

Calocoris humeralis, p. 89, lineolatus, p. 90, Argentine Rep., Berg, An. Soc. Arg. xxxiv, n. spp.

Camponotidea saundersi, macropterous form; Puton, p. 28, Rev. d'Ent. xi. Capsus delagrangei, Syria, Puton, p. 29, Rev. d'Ent. xi, n. sp.

Chrysodasia, n. g. near Lygus, p. 400, for C. strigifrons, n. sp., Venesuela, p. 401; REUTER, Ann. Soc. Ent. Fr. lxi.

Cyrtocapsus femoralis, Venezuela, REUTER, p. 394, Ann. Soc. Ent. Fr. lxi, n. sp.

Dolichomiris tibialis, Venezuela, Reuter, p. 392, Ann. Soc. Ent. Fr. lxi, n. sp.

Eccritotarsus luculentus, p. 195, nigriclavus, p. 196, Argentine Rep., Berg, An. Soc. Arg. xxxiv, n. spp.

Fulvius simoni, Venezuela, REUTER, p. 391, Ann. Soc. Ent. Fr. lxi, n. sp. Halticus minutus, injurious in Cochin-China; GIARD (337).

Helopeltis bergrothii, Gaboon, REUTER, p. 159, Ent. M. M. xxviii, n. sp.

IscAnias, n. g. (Capsaria), p. 193, for I. saltensis, n. sp., Argentine Rep., p. 194; Berg, An. Soc. Arg. xxxiv.

Linocerocoris, n. g. near Eucerocoris, for L. cariniventris, n. sp., Cameroons, Karsch, p. 133, Ent. Nachr. xviii.

Lopus flavomarginatus, n. var., luctuosus; Puton, p. 29, Rev. d'Ent. xi. Lygus æruginosus, p. 92, phaleratus, p. 93, Argentine Rep., Berg, An. Soc. Arg. xxxiv, n. spp.

Mecomma madagascariensis, Madagascar, Reuter, p. 185, Ent. M. M. xxviii, n. sp.

Oncognathus binotatus, habits; HOWARD, Ins. Life, v, pp. 90-92, figg. Orinonotus, n. subg., vide Physionotus infrà.

Orthotylus ochrotrichus, = (propinquus, Reut.); SAUNDERS, Ent. M. M. xxviii, p. 125.

Perissobasis, n. g., Perissobasaria (nov. divisio), p. 397, for P. aurora, n. sp., Venesuela, p. 398, Reuter, Ann. Soc. Ent. Fr. lxi.

Physetonotus, n. g., with n. subg. Orinonotus, and with P. (O.) simoni, n. sp., Venesuela; Reuter, p. 395, Ann. Soc. Ent. Fr. lxi.

Phytocoris riparum, N. Italy, FERRARI, Ann. Mus. Genov. (2) xii; P. bergrothi, p. 398, tripunctatus, p. 399, Venezuela, Reuter, Ann. Soc. Ent. Fr. lxi: n. spp.

- Paciloscytus rubricunsus, Argentine Rep., BERG, p. 94, An. Soc. Arg. xxxiv, n. sp.
- Pristoneura, n. g., Bryocoraria, for P. picea, n. sp., Venezuela; Reuter, Ann. Soc. Ent. Fr. lxi, p. 396.
- Resthenia (Callichila) occipitalis, p. 83, R. (R.) hirtula, p. 84, dissociata, p. 87, Argentine Rep., Berg, An. Soc. Arg. xxxiv, n. spp.
- Schizonotus, n. g., Cyllocoruria, p. 401, for S. dromedarius, n. sp., Caracas, p. 402; Reuter, Ann. Soc. Ent. Fr. lxi.
- Stenodema lateralis, E. Siberia, REUTER, p. 187, Œfv. Finska Förh. xxxiii, n. sp.

## HYDROCORISM.

- Aphelochirus sinensis, Ho-Chan, Montandon, p. 73, Rev. d'Ent. xì, n. sp. Coriza sealandica, New Zealand, Hudson, Manual, p. 120, pl. xx, fig. 5, n. sp.
- Enithares compacta, E. Africa, GERSTAECKER, p. 57, JB. Hamb. Anst. ix; E. bergrothi, New Caledonia, Montandon, p. 75, Rev. d'Ent. xi: n. spp.
- Luccotrephes ellipticus, E. Africa, GERSTAECKER, p. 56, JB. Hamb. Anst. ix, n. sp.
- Mononyx asiaticus, China, HORVÁTH, p. 136, Term. füzetek, xv, n. sp. Notonecta: fossil noticed; Schlechtendal (800).
- Ranatra: characters of the Palsearctic species; BERGROTH, p. 127, Rev. d'Ent. xi.
- R. nodiceps, E. Africa, GERSTAECKER, p. 57, JB. Hamb. Anst. ix, n. sp.

# HEMIPTERA-HOMOPTERA.

[Cf. CARLINI (158), DISTANT (210-215), DUZBE (233-237), FERRARI (281), GERSTAECKER (323), GODING (345, 346), HORVÁTH (428), HUDSON (440), LETHIERRY (549, 550), PUTON (707, 708), SLINGERLAND (851), TOWNSEND (945), and Coccidæ, p. 317.]

## CICADIDÆ.

- Baturia bicolorata, N. Guines, DISTANT, p. 316, Ann. N. H. (6) ix, n. sp. Callipsaltria bicolorata, S. Africa, DISTANT, p. 318, Ann. N. H. (6) ix, n. sp.
- Carineta tracta, p. 320, centralis, Ecuador, matura, Venezuela, p. 321, Distant, Ann. N. H. (6) ix, n. spp.
- Cicada viridis, Fab., = (D. mixta, Kirby); DISTANT, p. 325, Ann. N. H. (6) ix.
- C. flammata, Japan, Distant, p. 99, pl. xiii, fig. 15, Mon. Or. Cicad.;
  C. extrema, Swan River, p. 56, madagascariensis, Madagascar, p. 57,
  DISTANT, Ann. N. H. (6) x; C. timorensis, Timor, DISTANT, p. 406,
  Ann. N. H. (6) x; C. iolanthe, New Zealand, Hudson, Manual,
  p. 119, pl. xx, fig. 3: n. spp.

ł

- Cosmopsaltria feæ, p. 101, ficulnea, p. 102, Burma, DISTANT, Mon. Or. Cicad., n. spp.
- Dorachosa, n. g., p. 63, near Tibicen, for D. explicata, n. sp., Panama, p. 64; DISTANT, Ann. N. H. (6) x.
- Dundubia mixta, note on; KIRBY, p. 304, Wien. ent. Z. xi.
- Fidicina mülleri, Brazil, DISTANT, p. 319, Ann. N. H. (6) ix; F. amazona, Ega, bogotana, Bogota, p. 58, rubricata, Brazil, p. 59, DISTANT, Ann. N. H. (6) x: n. spp.
- Gazana and its species, characters of, with figures of numerous species; DISTANT, pp. 104-108, Mon. Or. Cicad.
- Graptotettix thoracica, Burma, DISTANT, p. 315, Ann. N. H. (6) ix, n. sp. Huechys: the species described, and most of them figured; DISTANT, pp. 111-116, Mon. Or. Cicad.
- H. fusca, p. 114, pl. iii, fig. 7, lutulenta, p. 115, pl. xiv, fig. 1, dohertyi, p. 116, fig. 2, Borneo, DISTANT, Mon. Or. Cicad., n. spp.
- Kanakia, n. g., near Malagasia, for K. typica, n. sp., New Caledonia; DISTANT, p. 62, Ann. N. H. (6) x.
- Lembeja, n. n., to replace Perissoneura; DISTANT, p. 103, Mon. Or. Cicad. Leptopsaltria japonica, Japan, Horváth, p. 136, Term. füzetek, xv; L. morrisi, E. India, DISTANT, p. 101, Mon. Or. Cicad., n. spp.
- Masupha, n. g., near Cicadatra, for M. ampliata, p. 317, delicata, p. 318, n. spp., S. Africa; DISTANT, Ann. N. H. (6) ix.
- Melampsalta: synonymy of New Zealand species, nervosa, Wlk., = (cassiope, Huds.); DISTANT, pp. 326 & 327, Ann. N. H. (6) ix. M. nervosa, Walk., = (mangu, White); DISTANT, p. 67, Ann. N. H. (6) x.
- M. exequata, Naga Hills, DISTANT, p. 144, pl. xv, fig. 11, Mon. Or. Cicad;
  M. labeculata, Australia, DISTANT, p. 66, Ann. N. H. (6) x; M. convicta, Norfolk I., rosacea, New Caledonia, p. 322, abdominalis, extrema, p. 323, Australia, rotundata, S. Africa, p. 324, DISTANT, op. cit. ix: n. spp.
- Mogannia: the species described and figured; DISTANT, Mon. Or. Cicad. pp. 119-122.
- M. effecta, E. India, DISTANT, Ann. N. H. (6) ix, p. 316, n. sp.
- Platypleura testacea, Somaliland, CARLINI, p. 537, Ann. Mus. Genov. (2) xii, n. sp.
- Pacilopsaltria trimeni, p. 313, peringueyi, p. 314, S. Africa, DISTANT, Ann. N. H. (6) ix, n. spp.
- Pomponia ransonneti, Dist., = (greeni, Kirby); DISTANT, p. 325, Ann. N. H. (6) ix.
- P. japonensis, Japan, DISTANT, p. 102, Mon. Or. Cicad., n. sp.
- Psaltoda flavescens, Australia, DISTANT, p. 55, Ann. N. H. (6) x, n. sp.
- Psilotympana infuscata, S. Africa, DISTANT, p. 319, Ann. N. H. (6) ix, n. sp.
- Talainga binghami, figured; DISTANT, pl. xiii, fig. 4, Mon. Or. Cicad.
- Terpnosia, n. g., near Pomponia, T. psecas, Wlk., = (P. elegans, Kirby); DISTANT, p. 325, Ann. N. H. (6) ix.

- T. pryeri, p. 139, pl. xv, fig. 5, Japan, andersoni, p. 141, pl. ix, fig. 13, China, Distant, Mon. Or. Cicad., n. spp.
- Tettigades parva, Argentine Rep., BERG, p. 65, Ann. N. H. (6) x, n. sp. Thopha sessiliba, Australia, Distant, p. 314, Ann. N. H. (6) ix, n. sp.
- Tibicen primitiva, Wlk., = (hæmatina, Stål), monilifera, Wlk., = (maculinervis, Stål); DISTANT, p. 325, Ann. N. H. (6) ix. T. nubifurca, Wlk., = (apicalis, Kirby); DISTANT, p. 325, t.e. T. putoni, n. n. for lacteipennis, Puton; DISTANT, p. 67, op. cit. x.
- T. (Quintilia) wealei, S. Africa, DISTANT, p. 316, Ann. N. H. (6) ix;
  T. egæ, Amazons, p. 64, pumilus, New Caledonia, p. 65, DISTANT,
  op. cit. x: n. spp.
- Tympanoterpes sodalis, Wlk., = (Fid. vultur, Wlk.); DISTANT, p. 67, Ann. N. H. (6) x.
- T. colombia, Bogota, p. 60, bergi, Argentine Rep. p. 61, DISTANT, Ann. N. H. (6) x, n. spp.

## FULGORIDE.

- Birdantis pallescens, Batchian, DISTANT, p. 276, Tr. E. Soc. 1892, n. sp. Copsyrna ochrucea, Perak, DISTANT, p. 286, Tr. E. Soc. 1892, n. sp.
- Cenestra ligata, Perak, pl. xiii, fig. 3, copulanda, Java, DISTANT, p. 285, Tr. E. Soc. 1892, u. spp.
- Cynthila viridimaculata, Perak, DISTANT, p. 275, Tr. E. Soc. 1892, n. sp. Dardus albomaculatus, p. 282, obscurus, p. 283, Australia, DISTANT, Tr. E. Soc. 1892, n. spp.
- Delphaz lugubris, Q noticed; LETHIERRY, p. 70, Rev. d'Ent. xi.
- Desudaba maculata, Australia, DISTANT, p. 277, Tr. E. Soc. 1892, n. sp.
- Dichoptera nubila, p. 277, hampsoni, E. India, nasuta, Celebes, p. 278, DISTANT, Tr. E. Soc. 1892, n. spp.
- Dictyophora unicolor, n. var. vittata, Puton, p. 30, Rev. d'Ent. xi.
- D. fuminervis, E. India, LETHIERRY, p. 208, Bull. Soc. Z. Fr. 1892; D. præferrata, bifasciata, insignis, Peak Downs, Australia, DISTANT, p. 279, Tr. E. Soc. 1892; n. spp.
- Eriphyle? paradoxa, E. Africa, Gerstaecker, p. 58, JB. Hamb. Anst. ix, n. sp.
- Eurybrachys apicata, India, DISTANT, p. 281, Tr. E. Soc. 1892, n. sp.
- Flata radiata, Borneo, labeculata, Celebes, p. 284, F. (Colobesthes?) semanga, Malay Penins., p. 285, pl. xiii, fig. 6, DISTANT, Tr. E. Soc. 1892, n. spp.
- Kandiana, n. g., near Messena, for K. lewisi, n. sp., Ceylon, pl. xiii, fig. 2, DISTANT, p. 280, Tr. E. Soc. 1892.
- Loxocephala custanea, E. India, DISTANT, p. 281, Tr. E. Soc. 1892, n. sp. Messena radiata, E. India, DISTANT, p. 280, pl. xiii, fig. 1, Tr. E. Soc. 1892, n. sp.
- Mycterodus orthocephalus, & described; FERRARI, p. 568, Ann. Mus. Genov. (2) xii.
- Ecleus decens, oviposition in Yucca; Townsend, Psyche, vi, p. 353.

- Oliarus angusticeps, Japan, Horváth, p. 137, Term. füzetek. xv, n. sp. Platybrachys signata, p. 281, insignis, Australia, ærata, Australia and Samoa, p. 282, DISTANT, Tr. E. Soc. 1892, n. spp.
- Phromnia parmata, Philippines, p. 283, montivaga, Borneo, p. 284, pl. xiii, fig. 5, Distant, Tr. E. Soc. 1892, n. spp.
- Ricania (Pochazia) flavocostata, Borneo, DISTANT, p. 283, Tr. E. Soc. 1892, n. sp.
- Scamandra diana, Sangir, DISTANT, p. 276, pl. xiii, fig. 4, Tr. E. Soc. 1892, n. sp.

# CERCOPIDÆ.

- Isthmia (?) elegantula, E. Africa, GERSTAECKER, p. 57, JB. Hamb. Anst. ix, n. sp.
- Phymatostetha deschampsi, E. India, LETHIERRY, p. 208, Bull. Soc. Z. Fr. 1892, n. sp.

## MEMBRACIDÆ.

- Synopsis of the N. American genera; Goding, Tr. Am. Ent. Soc. xix, pp. 253-260.
- Controdontus, n. g., Controtinæ, for Gargara atlus, Goding, n. sp., infrà; Goding, Ent. News, iii, p. 201.
- Gargara atlas, California, Goding, p. 110, Ent. News, iii, n. sp.
- Heliria scalaris, Colorado, Goding, p. 200, Ent. News, iii, n. sp.
- Platycotis minax, California, Goding, p. 109, Ent. News, iii, n. sp.
- Potnia asodalis, California, Goding, p. 110, Ent. News, iii, n. sp.
- Pubilia bicinctura, Colorado, Goding, p. 200, Ent. News, iii, n. sp.
- Stictocephala gillettei, Colorado, Goding, Ent. News, iii, pp. 108 & 200, n. sp.
- Stictopelta nova, California, p. 110, marmorata, New Mexico, p. 201, Goding, Ent. News, iii, n. spp.
- Telamona rileyi, California, Goding, p. 108, Ent. News, iii, n. sp.

#### JASSIDÆ.

- Tabulation of the N. American genera; Duzer, pp. 295-300, Tr. Am. Ent. Soc. xix.
- Acinopterus, n. g., near Athysanus, p. 307, for A. acuminatus, n. sp., N. America, p. 308, Duzee, Psyche, vi.
- Agallia sanguinolenta, oviposition and larvæ noticed; Osborn, Canad. Ent. xxiv, p. 35.
- Athysanus comma, bicolor, p. 114, obtusus (altered to obtutus, p. 156), p. 115, N. America, Duzee, Canad. Ent. xxiv, n. spp.
- Chlorotettix, n. g., near Thamnotettix, type Bythoscopus unicolor, Fitch, p. 306, with C. viridius, p. 309, galbanatus, p. 310, n. spp., N. America; Duzee, Psyche, vi.

- Deltocephalus ecchelii, Tyrol, FERRARI, p. 574, Ann. Mus. Genov. (2) xii;
  D. flavocostatus, Mississippi, Duzee, p. 116, Canad. Ent. xxiv;
  D. simplex, osborni, p. 304, cinerosus, signatifrons, p. 305, weedi,
  p. 306, N. America, Duzee, Tr. Am. Ent. Soc. xix: n. spp.
- Eupteryx zelleri, n. var. hipposideros, Honvath, p. 140, Rev. d'Ent. xi,
- Eutettix, n. g, type Thamnotettix lurida, Duzee; Duzee, Psyche, vi, p. 307.
- E. pictus, p. 301, marmoratus, p. 302, N. America, Duzer, Tr. Am. Ent. Soc. xix, n. spp.
- Gnathodus impictus, abdominalis, New Jersey, Duzee, p. 113, Canad. Ent. xxiv, n. spp.
- Goniagnathus elongatus, E. India, LETHIERRY, p. 209, Bull. Soc. Z. Fr. 1892, n. sp.
- Henschia, n. g., near Aconura, for H. seticauda, n. sp., Hungary; LETHIERRY, Rev. d'Ent. xi, p. 69.
- Macropsis indica, Mahé, Lethierry, p. 209, Bull. Soc. Z. Fr. 1892, n. sp. Phlepsius latifrons, p. 66, uhleri, p. 67, cinereus, p. 68, pallidus, p. 69, fuscipennis, p. 70, truncatus, p. 72, incisus, p. 73, punctiscriptus, p. 75, apertus, humidus, p. 76, nebulosus, p. 77, spatulatus, p. 78, ovatus, p. 79, superbus, p. 81, Duzee, Tr. Am. Ent. Soc. xix, n. spp.
- Tettigonia mitrata, E. Africa, GERSTAECKER, p. 58, JB. Hamb. Anst. ix, n. sp.
- Thamnotettix ghilianii, Turiu, FERRARI, p. 573, Ann. Mus. Genov. (2) xii; T. brevissimus, E. India, Lethierry, p. 209, Bull. Soc. Z. Fr. 1892; T. inornata, New York, Duzee, p. 303, Tr. Am. Ent. Soc. xix; T. smithi, longiseta, p. 266, gillettii, p. 267, montanus, p. 268, N. America, Duzee, Canad. Ent. xxiv: n. spp.
- Ulopa canadensis, Cauada, Duzee, p. 301, Tr. Am. Ent. Soc. xix, n. sp.

## PSYLLIDE.

Buctericera maritima, S. France, Horváth, p. 140, Rev. d'Ent. xi, n. sp. Diaphorina ægyptiaca, Cairo, Puton, p. 30, Rev. d'Ent. xi, n. sp. Psylla pyricola, habits and structure; SLINGERLAND (851). Trioza alacris, ravages on Laurus nobilis; Thomas (926).

### APHIDIDE.

- Pemphigiens: two forms of biological cycles in this group; Horvath (427).
- Lachnus pini and pineti, males described, pini, n. var. cembræ, p. 74; Cholodkovsky, Zool. Anz. zv, p. 67
- L. farinosus, St. Petersburg, Cholodkovsky, Zool. Ans. xv, p. 74, n. sp. Phylloxera vastatrix: abstract of life-history; Teich, KB. Ver. Riga, xxxiv, pp. 36-38: habits; Behr, Zoe, ii, pp. 305-310: sucking-apparatus, &c.; Krassilstschik (519).

- Siphonophora cucurbite, habits; Webster, p. 311, Canad. Ent. xxiv. S. tiliæ, mode of deposition of eggs; Howard, p. 249, P. E. Soc. Wash. ii.
- Toxoptera graminum, notes on; WEBSTER, Ins. Life, iv, pp. 245-248.

# Coccidæ, Aleurodidæ.

- [Gf. Cockerell (178, 179), Coquillett (184), Douglas (224, 625),
   Maskell (590, 591, 592), Mayer (594), Morgan (179, 625, 626),
   Newstead (637), Targioni-Tozzetti (912, 913), Townsend (949).
- Mode of preparing and mounting Coccidæ; MASKELL, Tr. N. Z. Inst. xxiv, pp. 3-5.
- General information about *Coccidæ*: their enemies, &c.; COCKERELL (178).
- For localities of a large number of species of Coccidæ, see MASKELL, Ent. M. M. xxviii, pp. 69-71.
- Coecidæ of California; Coquillett, Bull. Dep. Agric. Ent. xxvi, pp. 13-36.
- Note on the "spinnerets" of Diaspidinæ; MASKELL, Tr. N. Z. Inst. xxiv, pp. 5 & 6.
- Acanthococcidæ and Dactylopidæ, generic and group distinctions discussed; MASKELL, p. 30, pl. iv, figg. 11-17, Tr. N. Z. Inst. xxiv.
- Aleurodicus, n. g. for A. anonæ, n. sp., Demerara, and Aleurodes cocois, Curt.; Douglas & Morgan, Ent. M. M. xxviii, p. 32, pl. i.
- Aonidia blunchardi, Sahara, on date-palm, TARGIONI, p. 69, Mém. Soc. Zool., and Bull. Ent. Ital. xxiv, pp. 170-188, with developmental history: n. sp.
- Aspidiotus aurantii, synonymical note; MASKELL, Tr. N. Z. Inst. xxiv, p. 12.
- A. subrubescens, p. 9, pl. i, figg. 1 & 2, fodiens, figg. 3 & 4, bossieæ, figg. 5 & 6, p. 10, rossi, p. 11, figg. 7-9, Australia, Maskell, Tr. N. Z. Inst. xxiv, n. spp.
- Brachyscelidæ: notes on the group; MASKELL, p. 53, Tr. N. Z. Inst. xxiv.
- Carteria melaleucæ, p. 54, pl. xii, figg. 1-10, acaciæ, p. 56, figg. 11-15, Australia, Maskell, Tr. N. Z. Inst. xxiv, n. spp.
- Ceroplastes: spp. from Mexico and their parasites noticed; TOWNSEND, Zoe, iii, pp. 255-257.
- Chionaspis: note on species of; MORGAN, Ent. M. M. xxviii, pp. 14-16.
- C. sugenice, p. 14, pl. i, figg. 10-12, nitida, p. 15, figg. 13 & 14, Australia, Maskell, Tr. N. Z. Inst. xxiv, n. spp.
- Coccus cacti: observations on anatomy of, and production of wax; MAYER (594).
- Cielostoma compressum, New Zealand, p. 45, pl. xi, figg. 1-8, immane, p. 49, pl. xi, figg. 9-12, Australia, Maskell, Tr. N. Z. Inst. xxiv, n. spp.

- Ctenochiton eleocarpi, &, p. 17, depressum, small form, p. 19, described; MASKELL, Tr. N. Z. Inst. xxiv.
- C. dacrydii, N. Zealand; Maskell, p. 18, pl. ii, figg. 1-4, Tr. N. Z. Inst. xxiv, n. sp.
- Cylindrococcus, n. g. (position doubtful), p. 41, for C. casuarina, p. 41, pl. ix, spiniferus, p. 43, pl. x, figg. 1-7, sp. ?, p. 44, fig. 8, n. spp.; Australia, MASKELL, Tr. N. Z. Inst. xxiv.
- Dactylopius albizziæ, p. 31, pl. vi, figg. 3-10, hibbertiæ, p. 32, figg. 11-14, acaciæ, p. 33, figg. 15 & 16, Australia, iceryoides, New Zealand, p. 33, pl. vii, figg. 1-5, globosus, p. 34, figg. 6-8, sucalypti, p. 35, figg. 9-13, Australia, graminis, S. Africa, p. 36, pl. x, figg. 9-12, MASKELL, Tr. N. Z. Inst. xxiv, n. spp.
- Diuspis lanatus, Jamaica, Morgan, p. 136, J. Inst. Jamaica, i, n. sp.
- Eriococcus hoherice, apterous & noticed and figured, p. 26, pl. iv, figg. 9 & 10, Maskell, Tr. N. Z. Inst. xxiv; E. frazini, referred to Ripersia, Newstead, p. 147, Ent. M. M. xxviii.
- E. phyllocladi, New Zealand, p. 25, pl. iv, figg. 1-4, confusus, Australia, p. 26, figg. 5-8, fagicorticis, N. Zealand, pl. v, figg. 1-5, eucalypti, p. 27, figg. 6-14, tepperi, Australia, figg. 15-17, p. 29, MASKELL, Tr. N. Z. Inst. xxiv, n. spp.
- Fiorinia acacia, Australia, Maskell, p. 16, pl. i, figg. 15-17, Tr. N. Z. Inst. xxiv, n. sp.
- Frenchia, n. g. (Brachyscelidæ), p. 56, for F. casuarinæ, n. sp., Australia, p. 57, pl. xiii, Maskell, Tr. N. Z. Inst. xxiv.
- Gossyparia mannifera and manna in Algeria; GIARD, p. celxxiii, Bull. Soc. Ent. Fr. lxi.
- Icerya koebelei, Australia, Maskell, Ent. M. M. xxviii, p. 184, n. sp.
- Inglisia inconspicua, New Zealand, MASKELL, p. 19, pl. ii, figg. 5-7, Tr. N. Z. Inst. xxiv, n. sp.
- Lecanium capreæ, descriptive note; Douglas, Ent. M. M. xxviii, p. 278. L. rubi, Q described; Douglas, p. 105, t. c.
- L. minimum, assimile, England, NEWSTEAD, p. 141, Ent. M. M. xxviii; L. baccatum, Australia, MASKELL, p. 20, pl. ii, figg. 8-16, Tr. N. Z. Inst. xxiv; L. begoniæ, Demerara, Douglas, p. 209, pl. iii, fig. 4, Ent. M. M. xxviii: n. spp.
- Leucaspis pini, note on scales of; Morgan, Ent. M. M. xxviii, p. 13.
- Monophlebus crawfordi, notes on; Maskell, p. 51, Tr. N. Z. Inst. xxiv.
- Pinnaspis, n. g. or subg. of Mytilaspis, for part of Mytilaspis; Cockerell, J. Inst. Jamaica, i, p. 136.
- Planchonia stypheliæ, Australia, Maskell, p. 24, pl. iii, figg. 10-18, Tr. N. Z Inst. xxiv, n. sp.
- Poliuspis exocarpi, Australia, Maskell, p. 17, Tr. N. Z. Inst. xxiv, n. sp. Prosopophora, n. g. for P. dendrobii, n. sp., Demerara; Douglas, p. 207, pl. iii, fig. 1, Ent. M. M. xxviii.
- Pseudococcus socius, England (Pulvinaria ribesia, Doug., & is the & of this), NEWSTEAD, Ent. M. M. xxviii, p. 114, n. sp.

- Pseudoparlatoria, n. g. or subg. of Aspidiotus, for P. ostreala, n. sp., Jamaica; Cockerell, p. 136, J. Inst. Jamaica, i.
- Pulvinaria maskelli, life history; OLIFF, p. 176, pl. xi, Agric. Gas. N.S.W.
- P. persicæ, England, NEWSTEAD, p. 142, Ent. M. M. xxviii, n. sp.
- Rhizococcus grandis, Australia, Maskell, p. 29, pl. vi, figg. 1 & 2, Tr. N. Z. Inst. xxiv, n. sp.
- Ripersia tomlinii, habits noticed; DALE, Ent. M. M. xxviii, p. 219. R. formicicola, habitat in certain ants-nests; SMITH, Ent. M. M. xxviii, p. 307.
- R. pulveraria, England, p. 145, tomlinii, Guernsey, p. 146, NEWSTEAD, Ent. M. M. xxviii; R. rumicis, p. 37, pl. viii, figg. 1-3, formicicola, p. 38, pl. viii, figg. 4-7, New Zealand, Maskell, Tr. N. Z. Inst. xxiv: n. spp.
- Signoretia luzulæ, & described; NEWSTEAD, p. 142, Ent. M. M. xxviii.
- S. atriplicis, Australia, Maskell, p. 23, pl. iii, figg. 1-9, Tr. N. Z. Inst. xxiv, n. sp.
- Spharococcus, n. g. undefined, for S. casuarina, pl. viii, figg. 8-20, n. sp., Australia; Maskell, p. 39, Tr. N. Z. Inst. xxiv.
- Uhleria fioriniæ, synonymy and habitats; Morgan, Ent. M. M. xxviii, p. 12.

# (F.) NEUROPTERA.

Tabulation of the suborders and genera of N. American Neuropteroid insects, and Catalogue of the species; Banks, Tr. Am. Ent. Soc. xix, pp. 327-373.

#### TRICHOPTERA.

. [Cf. Linden (563), Martin (585).]

List of Trichoptera of the Department of the Indre, with notes on habits and dates; MARTIN, Rev. d'Ent. x, pp. 1-23 & 76.

On the earliest stages of a Phryganeid larva; Linden, Biol. Centralbl. xii, pp. 523-527.

Heteroplectron: systematic position queried; Banks, p. 373, Tr. Am. Ent. Soc. xix.

Hydroptila maclachlani, & characters; Morton, Ent. M. M. xxviii, p. 108.

Philopotamus montanus, var. chrysopterus, variation in neuration; MacLachlan, Ent. M. M. xxviii, p. 182.

## PLANIPENNIA.

[Cf. Banks (33), MacLachlan (574), Verhoeff (970)]

Bittacus tipularius, habits; Martin, Rev. Sci. Bourb. v, pp. 46-52.

Chrysopa ramburi, habits and localities; MacLachlan, Ent. M. M. xxviii, p. 50.

Chrysopa oculata, = (illepida, fulvibucca, chi, ypsilon, mississippiensis); BANKS, p. 373, Tr. Am. Ent. Soc. xix.

Formicaleo wilsoni, Hawaiian Is., MACLACHLAN, p. 178, Ann. N. H. (6) n. sp.

Hemerobius subnebulosus, development of colour in pupa; VERHOEF p. 297, Ent. Nachr. xviii.

Nothochrysa californica, N. America, BANKS, p. 373, Tr. Am. Ent. Soxix, n. sp.

# THYSANURA and COLLEMBOLA (APTERA).

[Cf. Harvey (390, 391), Reuter (734), Stummer-Traunfels (907).] The trophi of Japyx, Campodea, Collembola, Machilides, and Lepismide, discussed and figured; Stummer-Traunfels, SB. Ak. Wien, c, Abth. i, pp. 216-235, pls. i & ii.

Note on some Thysanura observed in connection with ants'-nests; Moniez (616).

Isotoma grandiceps, Siberia, REUTER, p. 229, Œfv. Finska Förh. xxxiii, n. sp.

Sminthurus viridis, n. var. tripunctatus, REUTER, p. 227, Œfv. Finska Förh. xxxiii, p. 227.

S. sex-maculatus, Maine, HARVEY, p. 169, Ent. News, iii, n. sp.

Templetonia americana, Maine, HARVEY, Ent. News, iii, p. 57, n. sp.

Tomocerus vulgaris, n. var. sibiricus; Reuter, p. 228, Œfv. Finska Förh. xxxiii.

# MALLOPHAGA, TERMITIDE, PSOCIDE.

[Cf. Grassi (359), Hudson (440), Kellogg (478), Martin (589).]

Mallophaga: notes on classification, and characters of genera; Kellogg, Tr. Kansas Ac. xii, pp. 46-48.

Habits and social relations of Termitidæ in Sicily; GRASSI (359).

Psocides of l'Indre, with notes on habits; MARTIN, pp. 285-288, Rev. d'Ent. xi.

Psocus zealandicus, New Zealand, Hudson, Manual, p. 107, pl. xvi, fig. 2, n. sp.

## PERLIDÆ,

[Cf. Martin (588), Nunney (649), Rostock (781).]

List of Perlide of l'Indre, with notes on habits; MARTIN, Rev. d'Ent. xi, p. 198.

Notes on and brief characters of British Perlidæ; NUNNEY, Sci. Goss. 1892, pp. 35, &c.

Capnodes, n. g. for C. schilleri, n. sp., Dresden; Rostock, p. 1, pl., Berl. Ent. Zeit. xxxvii.

#### EPHEMERIDÆ.

[Cf. Eaton, 241, 242, 243, 347).]

Baëtis salvini, & imago described; EATON, p. 13, Biol. Centr. Am. Neuropt.

Bleptus fasciatus, subimago described; EATON, p. 303, Ent. M. M. xxviii. Callibetis montanus, 2 subimago described; EATON, p. 15, Biol. Centr. Am. Neuropt.

Choroterpes exiguus, Tenasserim Valley, EATON, p. 189, Tr. E. Soc. 1892;
 C. inornata, Mexico, pl. i, fig. 5, nervosa, fig. 6, Guatemala, EATON,
 p. 6, Biol. Centr. Am. Neuropt. : n. spp.

Dipterominus tipuliformis, subimago noticed; EATON, p. 302, Ent. M. M. xxviii.

Euthyplocia hecuba, figured, pl. i, fig. 3, & imago described, p. 3; EATON, Biol. Centr. Am. Neuropt. E. sikorui, species of large size; VAYSSIÈRE, CR. Ass. Sci. Fr. xx, pt. l, p. 243.

Ephemera strigata, Japan, Eaton, p. 302, Ent. M. M. xxviii; E. remensa, p. 410, consors, p. 412, E. India, Eaton, J. A. S. B lx; E. pulcherrima, Tenasserim Valley, Eaton, p. 185, Tr. E. Soc. 1892: n. spp.

1892. [VOL. XXIX.] E 21

- Hagenulus monstratus, Tenasserim Valley, EATON, p. 189, Tr. E. Soc. 1892, n. sp.
- Hexagenia mexicana, & imago figured and described; EATON, p. 5, pl. i, fig. 4, Biol. Centr. Am. Neuropt.
- Homæoneuria salviniæ, figured, pl. i, fig. 2; EATON, Biol. Centr. Am. Neuropt.
- Lachlania lucida, figured, Biol. Centr. Am. Neuropt. pl. i, fig. 1.
- Leptohyphes brevissimus, Guatemala, Eaton, p. 12, pl. i, fig. 9, Biol. Centr. Am. Neuropt., n. sp.
- Palingenia robusta, p. 407, minor, p. 408, E. India, EATON, J. A. S. B. lx, n. spp.
- Potamanthus formosus, Tenasserim Valley, EATON, p. 186, Tr. E. Soc. 1892, n. sp.
- Rhoënanthus amabilis, Tenasserim Valley, EATON, p. 188, Tr. E. Soc. 1892, n. sp.
- Siphlurus binotatus, Japan, Eaton, p. 302, Ent. M. M. xxviii, n. sp.
- Thraulus mexicanus, 2 imago, lepidus, 3 2 described; EATON, p. 8, Biol. Centr. Am. Neuropt.
- T. primanus, Mexico, pl. i, fig. 7, versicolor, Costa Rica and Panama, valens, Panama, hilaris, Mexico, p. 8, EATON, Biol. Centr. Am. Neuropt.: n. spp.
- Tricorythus explicatus, Mexico, EATON, p. 11, pl. i, fig. 8, Biol. Centr. Am. Neuropt., n. sp.

#### ODONATA.

- [Cf. Calvert (150), Chatin (165, 166), Karsch (466, 467, 468), Martin (585), Selys (824, 825).]
- Odonata of Burma, with many notes; SELYS, Ann. Mus. Genov. (2) x, pp. 433 518.
- Revision of the African Gomphines; SELYS, pp. 86-107, Ann. Ent. Belg. xxxvi.
- Nature of respiration of larva of Eschna; MARTIN, Bull. Soc. Philom. (8) iv, p. 122.
- Epipharynx aud Hypopharynx in Odonata; Smith, p. 326, P. Am.
- Aciagrion, n. subg. for part of Pseudagrion, p. 509, with A. pallidum, n. sp., Burma. p. 512; Selys, Ann. Mus. Genov. (2) x.
- Eschna rileyi, Kilimanjaro, Calvert, p. 164, Tr. Am. Ent. Soc. xix, n. sp.
- Amphiæschna, Selys: composition and characters; KARSCH, p. 251, Ent. Nachr. xviii.
- Amphithemis, n. subg., p. 454, near Lyriothemis, for A. curristyla, p. 455, vacillans, p. 457, n. spp., Burma; Selys, Ann. Mus. Genov. (2) x.
- Anax pupuensis: characters; KARSCH, p. 250, Ent. Nachr. xviii.
- Anisopleura furcata, Burma, Selys, p. 488, Ann. Mus. Genov. (2) x, n. sp.

- Argiolestes melanothorar, Burma, Selys, p. 500, Ann. Mus. Genov. (2) x, n. sp.
- Brachydiplax gestroi, Burma, Selys, p. 451, Ann. Mus. Genov. (2) x, n. sp.
- Culicnemis chromothorax, p. 502, erythromelas, p. 505, Burma, Selys, Ann. Mus. Genov. (2) x, n. spp.
- Crenigomphus, n. g. for part of Onychogomphus, p. 97, with O. denticulatus, n. sp., Shoa; Selys, Ann. Ent. Belg. xxxvi.
- Deielia fasciata, Kirby: habitat and systematic position discussed; MacLachlan, Ann. N. H. (6) x, p. 177.
- Diplax dilatata, St. Helena, Calvert, p. 161, Tr. Am. Ent. Soc. xix, n. sp.
- Disparoneura abbotti, Kilimanjaro, Calvert, p. 164, Tr. Am. Ent. Soc. xix, n. sp.
- Echo incarnata, China, KARSCH, p. 455, Berl. Ent. Zeit. xxxvi, n. sp.
- Gynacantha bayadera, Burma, Gestro, p. 483, Ann. Mus. Genov. (2) x; G. limbalis, p. 252, musa, p. 254, Java, Karsch, Ent. Nachr. xviii: n. spp.
- Idionyx yolanda, optata, males described; Selys, pp. 470-473, Ann. Mus. Genov. (2) x.
- Isomma, n. g. (Legion, Lindenia), p. 104, for I. hieroglyphicum, n. sp.; Nossibé, p. 105; Selys, Ann. Ent. Belg. xxxvi.
- Lepthemis bluckburni: systematic position discussed; MacLachlan, Ann. N. H. (6) x, p. 177.
- Leptogomphus nietneri, description of &; Selys, p. 474, Ann. Mus. Genov. (2) x.
- L. ? maculivertex, Burms, SELYS, p. 478, Ann. Mus. Genov. (2) x, n. sp.
- Lestes nodalis, p. 496, umbrina, p. 497, L.? bilineata, p. 498, Burma, Selys, Ann. Mus. Genov. (2) x, n. spp.
- Libella testacea, n. race, chrysis, SELYS, p. 462, Ann. Mus. Genov. (2) x. Macromia paula, Cameroons, Karsch, p. 15, Berl. Ent. Zeit. xxxvii,
- Macromia paula, Cameroons, KARSCH, p. 15, Berl. Ent. Zeit. xxxvii, n. sp.
- Matrona kricheldorffi, China, Karsch, p. 456, Berl. Ent. Zeit. xxxvi, n. sp.
- Megalagrion bluckburni, variation in Lanai; MacLachlan, Ann. N. H. (6) x, p. 178.
- Mnais andersoni, description of sexes; Selys, pp. 485 & 486, Ann. Mus. Genov. (2) x.
- Onychogomphus æquistylus, p. 88, fritilarius, p. 89, obliteratus, p. 92, Madagascar, Selys, Ann. Ent. Belg. xxxvi, n. spp.
- Orogomphus speciosus, Burma, Selys, p. 481, Ann. Mus. Genov. (2) x, n. sp.
- Orthetrum truncatum, abbotti, Kilimanjaro, Calvert, p. 162, Tr. Am. Ent. Soc. xix, n. spp.
- Platygomphus few, Burma, Selys, p. 479, Ann. Mus. Genov. (2) x, n. sp. Pseudagrion asureum, Burma, Selys, p. 513, Ann. Mus. Genov. (2) x, n. sp.

- Rhinocypha iridea, Burma, Szlys, p. 492, Ann. Mus. Genov. (2) x, n. sp.
- Rhyothemis marcia and variegata are the sexes of one species, phyllis, distinguished therefrom; SELYS, pp. 440-444, Ann. Mus. Genov. (2) x.
- R. aterrima, Borneo, SELYS, p. 445, Ann. Mus. Genov. (2) x, n. sp.
- Thore concinna, MacLachlan, = (Sapho pulchella, Kirb.); MacLachlan, p. 456, Berl. Ent. Zeit. xxxvi.
- Trithemis aurora, intermedia, yerburii, synonymy discussed; Selys, p. 465, Ann. Mus. Genov. (2) x.
- T. limbata, Burma, p. 463, liturata, Celebes, p. 464, dryas, Burma, p. 465, Selys, Ann. Mus. Genov. (2) x, n. spp.
- Urothemis brevipennis, & described; SELYS, p. 469, Ann. Mus. Genov. (2) x.

# (G.) ORTHOPTERA.

[Cf. Adelung (3), Bateson & Brindley (42), Blatchley (104, 105), Blathwayt (107), Bolivar (110, 710), Brongniart (132, 134), Bruner (135), Bruner, Coquillett, & Osborn (136), Brunner (137), Brunner & Redtenbacher (138), Cholodkovsky (173), Distant (210), Dohrn (222), Eberli (244), Finot (282), Griffini (361), Heymons (404), Karsch (469-472), Krauss (520, 521), Künckel (529-532), Lataste (540), Pictet & Saussure (692), Rath (713), Redtenbacher (717), Saussure (210, 794, 795), Scudder (812, 816), Selys (826), Tepper (916, 917), Werner (1002).]

Notes on the Orthontera of Ceylon: Brunner pp. 337-339 Ent. Nachr

Notes on the Orthoptera of Ceylon; BRUNNER, pp. 337-339, Ent. Nachr. xviii.

Self-mutilation of carnivorous Orthoptera; WERNER (1002).

Hemimerus talpoides, systematic position queried; SHARP, Ent. M. M. xxviii, p. 212.

#### FORFICULIDE.

Dermaptera or Dermoptera, orthography discussed; KRAUSS & MIK, pp. 164-169, Wien. ent. Z. xi.

Forficula auricularia, oviposition, reluctance for flight; BLATHWAYT, Int. J. Micr. (3) ii, pp. 7-16, pls. i & ii: variation in & forceps; BATESON & BRINDLEY, P. Z. S. 1892, p. 586.

#### BLATTIDE.

Anaptycha, n. g., Ectobiida, for A. bipunctulata, n. sp., St. Vincent, pl. xv, fig. 1; Brunner, P. Z. S. 1892, p. 202.

Deropeltis bueana, Cameroons, KARSCH, p. 65, Berl. Ent. Zeit. xxxvii, n. sp.

Epilampra brevis, St. Vincent, BRUNNER, p. 203, pl. xv, fig. 3, P. Z. S. 1892, n. sp.

Holocampea simonyi, Tenerife, KRAUSS, p. 165, Zool. Anz. xv, n. sp.

Homalopteryz laminata, St. Vincent, BRUNNER, p. 204, pl. xv, fig. 4, P. Z. S. 1892, n. sp.

Loboptera fortunata, Canaries, KRAUSS, p. 165, Zool. Ans. xv, n. sp.

Nocticola, n. g., p. 29, for N. simoni, cæca, n. spp., found in caves in the Philippines, pp. 32 & 33, pl. i; BOLIVAR, Ann. Soc. Ent. Fr. lxi.

- Parasphæria nigra, St. Vincent, BRUNNER, p. 206, pl. xv, fig. 7, P. Z. S. 1892, n. sp.
- Periplaneta truncata, Canaries, &c., &c., KRAUSS, p. 165, Zool. Anz. xv, n. sp.
- Phyllodromia germanica, embryology; Cholodkovsky (173).
- Pseudophyllodromia semivitrea, St. Vincent, Brunner, p. 203, pl. xv, fig. 2, P. Z. S. 1892, n. sp.
- Rhyparobia, n. g. for Panchlora madera; Krauss, p. 165, Zool. Anz. xv. Stylopyga antillarum, St. Vincent, Brunner, p. 204, pl. xv, fig. 5, P. Z. S. 1892, n. sp.

# MANTIDAL

- Achlæna, n. g. (Mantides), p. 149, for A. crypsichroma, n. sp., Cameroons, p. 150; KARSCH, Ent. Nachr. xviii.
- Amorphoscelis annulipes, Cameroons, KARSCH, p. 145, Ent. Nachr. xviii, n. sp.
- Chlidonoptera, n. g., Harpagides, for C. vexillum, n. sp., Cameroons; Karsch, p. 68, Berl. Ent. Zeit. xxxvii, and p. 150, Ent. Nachr. xviii.
- Ductylopteryx, n. g. (Iris group), p. 8, for D. flexuosa, n. sp., Gaboon, p. 10; Karsch, Ent. Nachr. xviii.
- Hierodula bioculata, egg-case figured; KRAUSS, p. 146, Wien. ent. Z. xi. Hypsicorypha, n. g., near Idolomorpha, for H. juliæ, n. sp., Tenerife; KRAUSS, p. 166, Zool. Anz. xv.
- Leptocola tenuissima, Cameroons, Karsch, p. 146, Ent. Nachr. xviii; L. seriepunctata, p. 10, lignea, p. 12, W. Africa, Karsch, t. c.: n. spp.
- Litaneutria, n. g. for L. ocularis (n. sp., Mexico, not described); SAUSSURE, p. 124, Soc. Ent. vii.
- Melliera, n. g. for M. atopogamia, n. sp., Mexico; SAUSSURE, p. 123, Soc. Ent. vii.
- Miomantis armicollis, Cameroons, Karsch, p. 66, Berl. Ent. Zeit. xxvii, and Ent. Nachr. xviii, p. 148; M. preussi, p. 148, gracilis, p. 149, Cameroons, Karsch, t. c. : n. spp.
- Mionyx, n. g., Thespiens; SAUSSURE, p. 122, Soc. Ent. vii.
- Parastagmatoptera lobipes, St. Vincent, REDTENBACHER, p. 206, pl. xv, fig. 8, P. Z. S. 1892, n. sp.
- Phitrus, n. g., near Danuria, for P. lobulipes, n. sp., Cameroons; KARSCH, p. 149, Ent. Nachr. xviii.
- Phyllomantis, n. g., p. 124, for P. laurifolia, Panama, not described; SAUSSURE, p. 124, Soc. Ent. vii.
- Polyspilota laticollis, flavipennis, truncatipennis, Cameroons, Karsch, Ent. Nachr. xviii, p. 147, n. spp.
- Rhombodera scutata, W. Africa, KARSCH, p. 6, Ent. Nachr. xviii, n. sp.
- Stenopyga, n. g. (Euchomena, Gerst. nec Sauss.) for S. extera, n. sp., Cameroons, Karsch, p. 146, Ent. Nachr. xviii.
- Tenodera limbalicollis, W. Africa, KARSCH, p. 7, Ent. Nachr. xviii, n. sp. Thrinaconyx, n. g., Thespiens; SAUSSURE, p. 122, Soc. Ent. vii.

# PHASMATIDÆ.

- Leptynia kanigi, Tunis, KRAUSS, p. 147, Wien. ent. Z. xi, n. sp.
- Palophus: monograph of; Brongniart, N. Arch. Mus. Paris (3) iii, pp. 193-204, pls. viii & ix.
- Phanocles curvipes, St. Vincent, REDTENBACHER, p. 207, pl. xv, fig. 9, P. Z. S. 1892, n. sp.

### GRYLLIDÆ.

- Apithes mcneilli, Indiana, BLATCHLEY, p. 27, Canad. Ent. xxiv, n. sp.
- Ectatoderus antillarum, St. Vincent, REDTENBACHER, p. 218, pl. xvii, fig. 16, P. Z. S. 1892, n. sp.
- Endacustes dispar, St. Vincent, REDTENBACHER, p. 219, pl. xvii, fig. 18, P. Z. S. 1892, n. sp.
- Gryllodes rufipes, St. Vincent, REDTENBACHER, p. 217, pl. xvii, fig. 15, P. Z. S. 1892, n. sp.
- Gryllotalpa vulgaris, anatomy and histology of alimentary canal; EBERLI (244): phosphorescent; Ludwing, CB. Bakt. Parasit. ix, p. 561. G. borealis, notes on; FLETCHER, Canad. Ent. xxiv, p. 23, and DORAN, t. c. p. 147, life-history, id., t. c., p. 270.
- Gryllus guanchicus, Tenerife, KRAUSS, p. 171, Zool. Auz. xv, n. sp.
- Larandus marmoratus, St. Vincent, REDTENBACHER, p. 218, pl. xvii, fig. 18, P. Z. S. 1892, n. sp.
- Œcanthus niveus and allies distinguished; HART, Ent. News. iii, pp. 33 & 34.
- Phæophyllacris phalangium, Cameroons, KARSCH, p. 78, Berl. Ent. Zeit. xxxvii, n. sp.

#### LOCUSTIDÆ.

- Minute anatomy of tibial ears of Locustidæ; ADELUNG (3).
- Monographic revision of *Mecopodides*; REDTENBACHER, pp. 183-224, pl. iii, Verh. z.-b. Wien, xlii.
- Synopsis of generic characters of *Pseudophyllides*; PICTET & SAUSSURB, Icon. Saut. vertes.
- Acanthoprion, n. g., Pseudophyllides, p. 12, for A. aztecum, n. sp., Mexico, p. 26, pl. iii, fig. 21; Pictet & Saussure, Icon. Saut. vertes.
- Acilacris: systematic position; KARSCH, p. 327, Berl. Ent. Zeit. xxxvi.
- Alectoria superba, Brun., = (Hectoria pontoni, Tep.); Tepper, p. 87, Tr. B. Soc. S. Austr. xv.
- Ametrosomus, n. g. near Ametrus, p. 169, for A. helmsi, n. sp., Centr. Australia, p. 170; Tepper, Tr. R. Soc. S. Austr. xv.
- Amytta mutillata, & described; KARSCH, p. 328, Berl. Ent. Zeit. xxxvi.
- Anepitacta, systematic position; KARSCH, p. 327, Berl. Ent. Zeit. xxxvi.
- Anoedopoda, n. g., Mecopodides, p. 333, for A. erosa, n. sp., Cameroons, p. 334, Karsch, Berl. Ent. Zeit. xxxvi; & figured, op. cit. xxxvii, p. 77.

- Anonconotus, Cam., = (Analota, Brun.); GRIFFINI, p. 2, Boll. Mus. Zool. Tor. vii, No. 125.
- Aprion oculatum, Ceylon, maculifolia, Sumatra, pl. iii, fig. 19, PICTET & SAUSSURE, p. 21, Icon. Saut. vertes, n. spp.
- Apteronomus, n. g. near Apotrechus, for A. bordaensis, n. sp, Kangaroo I; TEPPER, p. 168, Tr. R. Soc. S. Austr. xv.
- Apteroscirtus, n. g., Mecopodides, for A. denudatus, n. sp., Cameroons; Karsch, p. 330, Berl. Ent. Zeit. xxxvi.
- Ariagona, n. g., Decticina, for A. margarita, n. sp., Tenerife; KRAUSS, p. 170, Zool. Anz. xv.
- Arota rosaura, noticed and figured; KARSCH, p. 316, Berl. Ent. Zeit.
- Bliastes superbus, p. 211, pl. xvi, fig. 12, striolatus, p. 212, fig. 13, St. Vincent, Redtenbacher, P. Z. S. 1892, n. spp.
- Brochopeplus, n. g., Pseudophyllides, p. 12, for B. reticulatus, n. sp., hab. ?, p. 25, pl. iii, fig. 20, Picter & Saussure, Icon. Sant. vertes.
- Cadicia longipennioides, p. 89, halmaturina, p. 91, hirsuta, p. 93, roseopennis, p. 94, Australia, Tepper, Tr. B. Soc. S. Austr. xv, n. spp.
- Calliphona, n. g. near Locusta, p. 169, for C. köniyi, n. sp., Tenerife, p. 170; KRAUSS, Zool. Anz. xv.
- Characta, n. g., Mecopodides, for C. bituberculata, n. sp., Borneo, pl. iii, fig. 9; Redtenbacher, p. 207, Verh. z.-b. Wien, xlii.
- Chloracris, n. g., p. 10, for C. prasina, India, pl. iii, fig. 14, brullei, Java, n. spp., p. 22; PICTET & SAUSSURE, Icon. Sant. vertes.
- Chlorotribonia, n. g., Pseudophyllides, p. 7, for C. acutipennis, n. sp., Java, p. 16, pl. ii, fig. 9; Pictet & Saussure, Icon. Saut. vertes.
- Clonia wahlbergi, habits; DISTANT, p. 83, Naturalist Transvaal.
- Condylodera tricondyloides, Westw., = (Trochalodera violascens, Brunner); DOHRN, p. 64, S. E. Z. liii.
- Corycus intermedius, W. Africa, REDTENBACHER, p. 221, pl. iii, fig. 12 Verh. z.-b. Wien, xlii, n. sp.
- Cyrtaspis scutata and variopicta, characters and synonymy; BOLIVAR, Rev. d'Ent. xi, pp. 289-293.
- Cyrtophyllus crepitans, St. Vincent, REDTENBACHER, p. 213, pl. xvii, fig. 14, P. Z. S. 1892, n. sp.
- Dasyphleps, n. g. for Locusta novæ-guineæ; KARSCH, p. 343, Berl. Ent. Zeit. xxxvi.
- Diaphlebus, n. g. for D. brevimarginatus, n. sp., Fiji Is.; KARSCH, p 343, Berl. Ent. Zeit. xxxvi.
- D. bivittatus, p. 193, marmoratus, p. 194, pl. iii, fig. 2, REDTENBACHER, Verh. z.-b. Wien, xlii, n. spp.
- Dicranopsyra, n. g., Phaneropterides, for D. multicolor, n. sp., Borneo; Dohrn, S. E. Z. liii, p. 71.
- Dictyota indivisa, Adelaide, p. 98, elderi, W. Australia, p. 99, Teppen, Tr. R. Soc. S. Austr. xv, n. spp.
- Elæoptera, n. g., Mecopodides, p. 195, for E. nitida, lineata, pl. iii, fig. 3, n. spp., Fiji Is., p. 196; REDTENBACHER, Verh. z.-b. Wien, xlii.

- Encentra, n.g., Mecopodides, for E. longipes, n. sp., Medellin, pl. iii, fig. 7, REDTENBACHER, p. 204, Verh. z.-b. Wien, xlii.
- Eonius, n. g. near Neanius, for E. tigrinus, p. 163, atrifrons, p. 164, fumatus, p. 165, n. spp., S. Australia; Tepper, Tr. R. Soc. S. Austr. xv.
- Ephippithyta quadrigesimaguttata, S. Australia, TEPPER, p. 84, Tr. R. Soc. S. Austr. xv, n. sp.
- Ephippithytoidea, n. g., Phaneropterides, for E. sparsa, n. sp., S. Australia; Tepper, p. 85, Tr. R. Soc. S. Austr. xv.
- Eucorypha mutica, Cameroons, Karsch, p. 327, Berl. Ent. Zeit. xxxvi; E. adicra, Cameroons, Karsch, p. 75, op. cit. xxxvii, n. spp.
- Eumegalodon, n. n. for Megalodon, Brullé nec Sowerby, p. 277; the species described and figured, pp. 279-285, pl. xii; a new tribe, Eumegalonidæ, proposed, p. 285; BRONGNIART, N. Arch. Mus. Paris (3) iii.
- Goëtia, n. g. (Psyres) for G. galbana, n. sp., Cameroons; Karsch, pp. 322 & 323, Berl. Ent. Zeit. xxxvi.
- Gravenreuthia, n. g., Phaneropterides, p. 73, for G. saturata, n. sp., Cameroons, p. 74; Karsch, Berl. Ent. Zeit. xxxvii.
- Gryllacridæ of Australia; TEPPER, pp. 139-170, Tr. R. Soc. S. Austr. xv.
- Gryllacris eximia, p. 339, genufusca, p. 340, nigriceps, p. 341, Cameroons,
  KARSCH, Berl. Ent. Zeit. xxxvi; G. atrogeniculata, Centr. Australia,
  p. 144, marmoriceps, p. 145, longicornis, p. 146, ferrotestacea, p. 147,
  S. Australia, lutescens, Centr. Australia, p. 148, subdebilis, p. 151,
  incerta, p. 154, paulula, molineusiana, p. 155, S. Australia, Tepper,
  Tr. B. Soc. S. Austr. xv: n. spp.
- Hemisaga, note on the genus, with corrections of errors; SAUSSURE, pp. 12-14, Ann. Soc. Ent. Fr. lxi.
- H. prædatoria, pp. 63 & 258, Transvaal, DISTANT, Naturalist Transvaal, n. sp.
- Ityocephala, n. g., Mecopodides, p. 219, for I. falcata, n. sp., Fiji Is., pl. iii, fig. 11, p. 220; REDTENBACHER, Verh. z.-b. Wien, xlii.
- Leproscirtus, n. g., Mecopodides, p. 328, for Mecopoda granulosa, Karsch; Karsch, Berl. Ent. Zeit. xxxvi.
- Leptodera ornatipennis, characters of; DOHRN, p. 66, S. E. Z. liii.
- Macroscirtus brevipennis, Sierra Leone, REDTENBACHER, p. 218, Verh. z.-b. Wien, xlii, n. sp.
- Mataus casamanca, Guinea, Pictet & Saussure, p. 23, pl. iii, fig. 13, Icon. Saut. vertes, n. sp.
- Mecopodides: tabulation of certain genera of; Karsch, Berl. Ent. Zeit. xxxvi, pp. 341-346.
- Mecopoda dilatata, Borneo, p. 213, divergens, hab. ?, REDTENBACHER, p. 213, Verh. z.-b. Wien, xlii, n. spp.
- Microprion, n. g., Pseudophyllides, p. 17, for M. philippinensis, Philippines, elliptifolia, Java, pl. ii, fig. 11, p. 18, ; PICTET & SAUSSURE, Icon. Saut. vertes.

- Microsaga, Sauss., referred to Conocephaliens; SAUSSURE, p. 15, Ann. Soc. Ent. Fr. 1xi.
- Mioacris, n. g., Pseudophyllides, p. 7, for M. javana, n. sp., Java, p. 16: Pictet & Saussure, Icon. Sant. vertes.
- Miosagu, n. g., Sagiens, p. 10, for M. orthoxipha, n. sp, New Holland, p. 11; SAUSSURE, Ann. Soc. Ent. Fr. lxi.
- Mirollia carinatu, & characters; DOHRN, p. 66, S. E. Z. liii.
- Onomarcus latipennis, p. 15, mandarinus, p. 16, China, PICTET & SAUSSURE, Icon. Saut. vertes, pl. i, figg. 6 & 7, n. spp.
- Orchelimum gracilis is not Xiphidium fasciatum; BRUNER, p. 264, Ent. News, iii.
- Orophila, n. g. (isolated), p. 168, for O. nubigena, n. sp., Tenerife, p. 169; Krauss, Zool. Anz. xv.
- Pachypodagrus, n. g., Stenopelmatidæ, p. 176, for P. magareyi, crassipes, n. spp., S. Australia, p. 177; Tepper, Tr. R. Soc. S. Austr. xv.
- Paragryllacris pallidolinea, p. 159, insignis, p. 161, deserta, p. 162, S. Australia, Tepper, Tr. R. Soc. S. Austr. xv, n. spp.
- Penthoplophora, n. g., Stenopelmatidæ, p. 175, for S. driffieldi, n. sp., Australia; Tepper, p. 176, Tr. B. Soc. S. Austr. xv.
- Phaneropteridæ of Australia; TEPPER, pp. 77-113, Tr. R. Soc. S. Austr. xv.
- Phaula trichopus, Deh., = (chlorotica, Brunner); DOHEN, p. 69, S. E. Z. liii.
- Phlaurocentrum mecopodoides, Cameroons, KARSCH, p. 321, Berl. Ent. Zeit. xxxvi, n. sp.
- Phricta, n. g., p. 191 (Mecopodides), for P. spinosa, n. sp., Queensland, p. 192, pl. iii, fig. 1; REDTENBACHER, Verh. z.-b. Wien, xlii.
- Phyllomimus truncatifolia, Moluccas, PICTET & SAUSSURE, p. 19, pl. ii, fig. 10, Icon. Saut. vertes, n. sp.
- Phyllotribonia, n. g., Pseudophyllides, p. 10, for P. verruculosa, n. sp., Centr. Africa, p. 23, pl. iii, fig. 17; Pictet & Saussure, Icon. Saut. vertes.
- Phyllozelus, n. g. for P. genicularis, n. sp., hab. ?, pl. iii, fig. 16, PICTET & SAUSSURE, p. 24, Icon. Saut. vertes.
- Plangiopsis foraminata, Cameroons, KARSCH, p. 324, Berl. Ent. Zeit. xxxvi, n. sp.
- Platycleis azami, S. France, Finot, Bull. Soc. Ent. Fr. lxi, p. xxxv, n. sp. Pæcilopsyra, n. g. for Phaneroptera octoseriata, Deh.; Dohrn, p. 69, S. E. Z. liii.
- Polichne ferruginea, S. Australia, TEPPER, p. 100, Tr. R. Soc. S. Austr. xv, n. sp.
- Preussia lobatipes, & described; KARSCH, p. 326, Berl. Ent. Zeit. xxxvi. Pyrgophylax emarginatus, Borneo, DOHEN, p. 68, S. E. Z. liii, n. sp.
- Rhammatcpoda, n. g., Mecopodides, for R. opilionoides, n. sp., Peru, pl. iii, fig. 6; Redtenbacher, p. 203, Verb. z.-b. Wien, xlii.
- Sagiens: synopsis of genera; SAUSSURE, pp. 5-16, Ann. Soc. Ent. Fr. lxi.

- Scutotribonia, n. g., Pseudophyllides, p. 11, for S. humbertiana, n. sp., Ceylon, pl. iii, fig. 18, p. 24; Picter & Saussure, Icon. Saut. vertes.
- Segestes punctipes, fuscus, Philippines, unicolor, Pelew Is., p. 199, decoratus, New Guines, p. 200, pl. iii, fig. 4, Redtenbacher, Verh. z.-b. Wienzlii, n. spp.
- Stenopelmatidæ, synopsis of Australian and Polynesian; Tepper, pp 170-178, Tr. R. Soc. S. Austr. xv.
- Sthenaropoda, n. g., Mecopodides, p. 231, for S. preussiana, n. sp., Cameroons, p. 332; Karsch, Berl. Ent. Zeit. xxxvi.
- Sympæstria longipes, Java, Dohrn, S. E. Z. liii, p. 73, n. sp.
- Taniomena sororoides, p. 106, S. Australia, fraserensis, W. Australia, p. 108, minor, S. Australia, p. 109, Tepper, Tr. R. Soc. S. Australia, v., n. spp.
- Terp audrus horridus, redescription and synonymy; SAUSSURE, Ann. Soc. Ent. Fr. 1xi, pp. 7 & 8.
- Trachyzulpha, n. g. near Zulpha, p. 67, for T. fruhstorferi, n. sp., Java, p. 68; Dohrn, S. E. Z. liii.
- Tympanoptera, n. g., p. 19, for T. grivleti, Moluccas, pl. iii, fig. 15, oceanica, Fiji Is., pl. ii, fig. 12, p. 20; Pictet & Saussure, Icon. Saut. vertes.
- Xiphidium scudderi, Indiana, BLATCHLEY, Canad. Ent. xxiv, p. 26, n. sp.

#### ACRIDIDAE.

- Acridiid ovipositing in Rubus stems; VERHOEFF, p. 298, Ent. Nach. xviii.
- Arminda, n. g. near Platyphyma, for A. brunneri, n. sp., Tenerife; KRAUSS. p. 168, Zool. Anz. xv.
- Caletes, n. g, p. 209, for C. opterus, n. sp., St. Vincent, p. 210, pl. xvi, fig 11; REDTENBACHER, P. Z. S. 1892.
- Caloptenus vulcanius, Canary Is., KRAUSS, p. 167, Zool. Anz. xv, n. sp.
- Chrotogonus meridionalis, Transvaal, SAUSSURE, p. 262, pl. iv, fig. 5, in Distant, Naturalist Transvaal, n. sp.
- Eremobia clareli, n. var., læviuscula; KRAUSS, p. 149, Wien. ent. Z. xi.
- Gemeneta, n. g., Acridiides, p. 71, for G. terrea, n. sp., Cameroons, p. 72; Karsch, Berl. Ent. Zeit. xxxvii.
- Hippiscus pantherinus, p. 285, haldemanii, p. 286, compactus, p. 288, variegatus, suturalis, p. 301, H. (Sticthippus) californicus, p. 317, marmoratus, p. 318, H. (xanthippus) conspicuus, p. 319, eremitus, p. 320, maculatus, p. 333, tigrinus, p. 334, affrictus, p. 336, altivolus, p. 347, cupidus, p. 348, pumilus, p. 349, albulus, p. 350, latefasciatus, obscurus, p. 359, griseus, aurilegulus, p. 362, stigmosus, p. 363, N. America, Scudder, Psyche, vi, n. spp.
- Melanoplus griseus, & described; BLATCHLEY, p. 30, Canad. Ent. xxiv. Leptysma marginicollis, habits; BLATCHLEY, p. 28, Canad. Ent. xxiv. Elipoda canariensis, Tenerife, KRAUSS, p. 167, Zool. Anz. xv, u. sp.
- Pamphagus foreli, 2 described; KRAUSS, p. 107, 2001. ALZ. 17, H. Sp.

Petusia spumans, var. ater; DISTANT, p. 259, pl. iv, fig. 3, Naturalist Transvaal.

Pezotettix hoosieri, Indiana; Blatchley, p. 31, Canad. Ent. xxiv, n. sp. Pygostolus inuncatus, Cameroons, Karsch, p. 72, Berl. Ent. Zeit. xxxvii, n. sp.

Schintocerca peregrina: Criquet pelerin, Sauterelle de la Bible: oviposition, KÜNCKEL, C.R. Ass. Fr. Sci. xx, pt. 1, p. 241, and pt. 2, p. 554; correction as to egg-cases, Lucas, p. lxxxi, Ball. Soc. Ent. Fr. lxi; note on the changes of colour that accompany its moults, KÜNCKEL, p. xxv, Bull. Soc. Ent. Fr. lxi, and Brongniart, t. c. p. lix; S. americana, the migratory locust of Chili, Bolivar, Act. Soc. Chili, ii, pp. 196-199: but it is not this species, Lataste, t. c. pp. 200-209.

Stauronotus maroccanus, structure of Ootheca; TARRAL, Bull. Soc. Bésiers, xii, p. 10.

Stenobothrus simonyi, Canary Is., Krauss, p. 166, Zool. Anz. xv, n. sp. Sticthippus, n. subg. of Hippiscus, Scudder, Psyche, vi, p. 317.

Tettix quadriundulatus, St. Vincent, REDTENBACHER, p. 208, pl. xvi, fig. 10, P. Z. S. 1892, n. sp.

Thalpomena picteti, Tenerife, Krauss, p. 167, Zool. Anz. xv, n. sp. Xiphocera distanti, pl. iv, fig. 1, picta, fig. 2, Saussure, p. 261, in Distant, Naturalist Transvaal, n. spp.

# XIV. ECHINODERMATA.\*

BY

# F. A. BATHER.

## I.—TITLES.

BARTHELS, P. [See LUDWIG & BARTHELS (32).]

- BATHER, F. A. Suggested Terms in Crinoid Morphology. Ann. N. H. (6) ix, pp. 51-66. [Crin.]
- British Fossil Crinoids. VI. Botryocrinus quinquelobus, n. sp., Wenlock Limestone; and note on B. pinnulatus. T. c. pp. 189-194, pl. xi. [Crin.]
- British Fossil Crinoids. VII. Mastigocrinus loreus, n. g. & sp., Wenlock Limestone, Dudley. T. c. pp. 194-202, pls. xi & xii. [Crin.]
- British Fossil Crinoids. VIII. Cyathocrinus: C. acinotubus, Ang., and C. vallatus, n. sp., Wenlock Limestone. T. c. pp. 202-226, pl. xiii. [Crin.]
- The Fossil Crinoidea in the British Museum. (An attempt to put into practice modern ideas of Museum arrangement.) Ann. Rep. Museums' Assoc. for 1891, pp. 78-110. [Crin.]
  - —. [See also Blake (13), Nortling (38), Woods (55).]

<sup>\*</sup>While Mr. Bather's Record was still incomplete he was compelled to take a voyage, of some months duration, for the benefit of his sight. He has not returned in time to finish his work, the completion of which must therefore be given in the next volume of our Record, in combination with the literature of 1893.—Editor Z. R.

<sup>1892. [</sup>VOL. XXIX.]

- BEACHLER, C. S. Rocks of the Niagara age in Indiana. Am. Geol. ix, pp. 408 & 409. [Crin.]
- Keokuk Group of the Mississippi Valley. Op. cit. x, pp. 88-96. [Crin.]
- 8. Bell, F. J. On the Echinoderms collected by the Ss. "Fingal" in 1890, and by the Ss. "Harlequin" in 1891, off the West Coast of Ireland. Sci. P. R. Dubl. Soc. (n.s.) vii, pp. 520-529, pls. xxiii-xxv. [Hol., Crin., Ast., Oph., Ech.]
  - 9. —. A contribution to the classification of *Ophiwoids*, with descriptions of some new and little-known forms. P. Z. S. 189 pp. 175-183, pls. xi & xii. [Oph.]
  - 10. —. Description of a new species of Antedon from Mauritius——. Ann. N. H. (6) ix, pp. 427 & 428, pl. xviii. [Orin.]
  - 11. Bigot, A. [Remarks on Cidaris cenomanensis, Cott.] Bull. Soc\_ \_\_\_\_\_ L. Norm. (4) v, pp. 54 & 55. [Ech.]
  - 12. BIZET, P. Considérations Géologiques and Paléontologiques sur le terrains des Environs de Bellême et de Mamers. Bull. Soc. Géol. Norm. xiii, pp. 95-132 (Havre, 1890). [Ech.]
    Distribution of Echinoids in Jurassic.
  - 13. Blake, J. F. Annals of British Geology, 1891, pp. 224-231. [Gives abstracts of the following papers in Record for 1891: GREGORY (3) and (8), ROBERTS, SLADEN (2), CARPENTER (1), BATHER (1) and (2).]
  - BOEHM, G. Ueber eine Anomalie im Kelche von Millericrinumespiliformis. Z. geol. Ges. zliii, pp. 741-743. [Crin.]
     CARPENTER, P. H. [See under BLAKE, J. F. (13).]
  - COLE, A. H. Palaeaster eucharis, Hall. Bull. Geol. Soc. Am. iii, pp. 512-514, pl. xv. [Ast.]
  - 16. CUNNINGHAM, J. T. The Preservation of Marine Animals for Zoological purposes. Ess. Nat. vi, p. 124.

  - 18. DYCHE, D. T. D. [Termination of the column of certain Crinoids.]

    Am. Geol. x, p. 130. [Crin.]
  - 19. Eck, H. Abnormer Kelchbau bei Encrinus gracilis. Z. geol. Ges\_\_\_\_ zliv, pp. 138 & 139. [Crin.]

- FABER, C. [See MILLER & FABER (34).]
- 21. Field, G. W. The Larva of Asterias vulgaris. Q. J. Micr. Sci. (n.s.) xxxiv, pp. 105-128, pls. xiii-xv. [Ast. and General.]
- 22. Gregory, J. W. The Relations of the American and European Echinoid Faunas. Bull. Geol. Soc. Am. iii, pp. 101-108. [Ech.]
- Archæopneustes abruptus, a new genus and species of Echinoid from the Oceanic Series in Barbados. Q. J. Geol. Soc. xlviii, pp. 163-169. [Ech.]
- 24. —. Further additions to Australian Fossil Echinoidea. Geol. Mag. (n.s.), (iii), ix, pp. 433-437, pl. xii. [Ech.]
  - —. [See also Blake, J. F. (13), & Woods (55).]

GUETTARD, J. E. [See LETACQ (29).]

HARRISON, J. B. [See JUKES-BROWNE & HARRISON (26).]

- HARTLAUB, C. Ueber die Arten und den Skelettbau von Culcita.
   Notes Leyd. Mus. xiv, pp. 65-118, pls. i & ii. [Ast.]
- JUKES-BROWNE, A. J., & HARRISON, J. B. The Geology of Barbados. Part 11. The Oceanic Deposits. Q. J. Geol. Soc. xlviii, pp. 170-226. [Ech.]
- KEYES, C. R. The Principal Mississippian Section. Bull. Geol. Soc. Am. iii, pp. 283-300, pl. ix. [Crin.]
  - KOKEN, E. [See WÖHRMANN & KOKEN (54).]
- 28. LENT, C. Die Fauna der Renggerithone von Kandern. MT. badischen Geol. Landesanst. ii, Heft 3, pp. 623-639. [Crin., Ast.]
- 29. LETACQ, A. L. Notice sur les Travaux Scientifiques de Guettard aux environs d'Alençon et de Laigle (Orne). Bull. Soc. L. Norm. (4) v, pp. 67-85. [Crin.]
- On p. 75 is a note "Sur les Encrines et les Pierres étoilées recuellies à Alençon," described by J. E. Guettard.
- LOVEN, S. Echinologica. Bih. Sv. Ak. Handl. xviii (4), No. 1, pp. 1-74, pls. i-xii. [Ech.]
- Ludwig, H. Ueber die Rädchen der Synaptiden. Z. wiss. Zool. liv, pp. 350-364. [Hol.]
- Ludwig, H., & Barthels, P. Beiträge zur Anatomie des Holothurien. T. c. pp. 631-654, pl. xxviii. [Hol.]
- 33. MACBRIDE, E. W. The Development of the Genital Organs, Ovoid Gland, Axial and Aboral Sinuses in Amphiura squamata; together with some remarks on Ludwig's Hæmal System in this Ophiurid. Q. J. Micr. Sci. (n.s.) xxxiv, pp. 129-153, pls. xvi-xviii. [Oph.]

- 34. MILLER, S. A., & FABER, C. Some new Species and new Structural parts of Fossils. J. Cincinn. Soc. xv, pp. 79-81, pl. i. [Cyst.] Critical note in Am. Geol. x, pp. 316 & 317, Nov.
- MILLS, J. E. Stratigraphy and Succession of the Rocks of the Sierra Nevada of California. Bull. Geol. Soc. Am. iii, pp. 413-444, pl. xiii. [Oris., stems in Lower Mesosoic.]
- Minchin, E. A. Notes on the Cuvierian Organs of Holothuria nigra.
   Ann. N. H. (6) x, pp. 273-284, pl. xvii. [Hol.]

Newton, E. T. [See Sharman & Newton (46).]

- 37. NICKLÈS, R. Recherches Géologiques sur les Terrains Secondaires et Tertiaires de la Province d'Alicante et du Sud de la Province de Valence. Ann. Hébert, i, pp. 1-219, pls. i-x.
  - Pp. 202-204, and pls. viii & ix (parts) refer to Echinoderms.
- NOETLING, F. Field Notes from the Shan Hills. [See Record for 1891.] Critical note by F. A. B.; Geol. Mag. (n.s.) ix, p. 521. [Cyst.]
- 39. ŒHLERT, D. P. Description de deux Crinoïdes nouveaux du Dévonien de la Manche. Bull. Soc. Géol. (3) xix, pp. 834-853, pl. xviii. [Crin.]
- 40. Ramsay, E. P. Catalogue of *Echinodermata* in the Australian Museum. Part I. 2nd Ed. Sydney: 1891, 8vo, vii & 60 pp., 4 pls.
- 41. Reed, F. R. C. Woodwardian Museum Notes. Geol. Mag. (n.s.), (iii), ix, pp. 548 & 549. [Crin.]
- RIGAUX, E. Notice géologique sur le Bas Boulonnais. Mém. Soc. Acad. Boulogne, xiv, p. 108, pls. i & ii. [Ech., Ast., Oph., Crin., localities and horizons; especially Jurassic.]
   ROBERTS, T. [See BLAKE, J. F. (13).]
- ROTHPLETZ, A. Die Perm-, Trias-, und Jura-Formation auf Timor und Rotti im indischen Archipel. Palseontogr. xxxix, pp. 57-106, pls. ix-xiv. [Crin., Ech.]
- SEELIGER, O. Die Embryonal-entwicklung der Comatula (Antedon rosacea). Zool. Anz. xv, pp. 391-393. [Crin.]
- The Embryonic Development of Comatula (Antedon rosacea).
   Ann. N. H. (6) x, pp. 481-483, Dec. [Crin.]
   A translation of the preceding.
- 46. SHARMAN, G., & NEWTON, E. T. On a New Form of Agelacrinites (Lepidodiscus milleri, n. sp.) from the Lower Carboniferous Lime stone of Cumberland. Q. J. Geol. Soc. xlviii, pp. 150-152, pl. ii [Cyst.]
  - SLADEN, W. P. [See BLAKE, J. F. (13).]

- SPRINGER, F. [See WACHSMUTH & SPRINGER (50).]
- 47. STOLLEY, E. Die Kreide Schleswig-Holsteins. MT. Min. Inst. Kiel, i, pp. 191-309, pls. vii-x. [Ech., Crin.]
- ULRICH, A. Palæozoische Versteinerungen aus Bolivien. Beitr. z. Geol. u. Paleont. v. Südamerika, i, pp. 5-116, pls. i-v. Stuttgart. [Crin., columnals only.]
- 49. Wachsmuth, C. [A Letter on *Pentacrinus* stems in Jurassic (?) Limestones of California; see Mills, J. E. (35), p. 428.]
- 50. Wachsmuth, C., & Springer, F. Description of Two new Genera and Eight Species of Camerate Crinoids from the Niagara Group. Am. Geol. x, pp. 135-144. [Crin.]
- WAGNER, R. Ueber einige Versteinerungen des unteren Muschelkalks von Jena. Z. geol. Ges. xliii, pp. 879-901, pl. xlix. [Crin.]
- 52. WHITEAVES, J. F. The Fossils of the Devonian Rocks of the islands, shores, or immediate vicinity of Lakes Manitoba and Winnepeg. Geol. Surv. Canada. Contrib. to Canadian Palæont. i, pt. iv, pp. 255-359, pls. xxxiii-xlvi. [Advance sheets to p. 318 and pl. xi, Sept.; the rest Dec.] [Crin., p. 276, fragments of Ctenocrinus sp.]
- WILLIAMSON, W. C. Exhibit and Remarks on Astropecter orion, Kellaways Rock, Pickering, Yorkshire. P. Geol. Soc. 1892, pp. 184 & 185. [Ast.]
- 54. WOHEMANN, S. VON, & KOKEN, E. Die Fauna der Raibler Schichten vom Schlernplateau. Z. geol. Ges. xliv, pp. 167-223, pls. vi-xvi. [Crin., p. 170, columnals of Encrinus cassianus.]
- 55. Woods, H. Catalogue of the Type Fossils in the Woodwardian Museum, Cambridge. With preface by T. McK. Hughes. Cambridge: 1891 (publ. Jan., 1892), 8vo, xvi & 180 pp. [Ech., pp. xvi & 30-34; Ast., pp. 35 & 36, revised by J. W. Gregory; Oph., pp. 36 & 37; Crin., pp. xvi & 37-42; Cyst., pp. 42 & 43, revised by F. A. Bather; Blast., pp. 43 & 44; Hol., p. xvi.]

### II.—BIOLOGY.

# (a.) ANATOMY.

Echinodermata: general account; BATHER, (5) p. 84.

Holothurioidea: Cuvierian organs of Holothuria and Muelleria; LUD-WIG & BARTHELS (32): of Holothuria nigra; MINCHIN (36).—The wheelspicules ("Bädchen") of Synaptida; LUDWIG (31).

Echinoidea: Interambulacra of Archæocidaris selwyni, and possible new genus; ETHERIDGE (20).—The buccal membrane, its origin and history; the auricles, their structure and development, pp. 36 et seq.; the dental apparatus, its structure and development, pp. 40 et seq.; Muscles of dental

apparatus, p. 42; LOVEN (30).—Corona of the Cidarida; LOVEN (p. 18.

Asteroidea: General structure and homologies of skeleton in Culci Habitaub, (25) p. 99.—Ambulacra of Palæaster, new subgenus Monter; Etheridge (20).—Detailed description of oral surface of Palæaeucharis; Cole (15).

Ophiuroidea: Large calycinal plates of young Pectinura, described a figured, p. 176; arm-ossicles of Ophioteresis elegans, and of Ophiuro generally as basis of classification, pp. 176-178; radial shields of Ophimaza obscura, p. 182; Bell (9).—Ovoid gland, p. 134; Hæmal systems, 140; genital organs (p. 142) of Amphiura squamata; MACBRI (33).

Crinoideu: New terms suggested: "Super-radials" and "Infer-radia for halves of transversely bisected radials; "Primibrachs, Secundibrac &c.," for series of non-pinnulate arm-ossicles, or brachials; "Mono chals, Distichals, Tetrastichals, &c.," for the same when pinnula corresponding names for intercalated and interradial plates, by pre "Inter"; "Deltoids," for certain tegminal plates; "Cirrals," "Radica and "Imperforate articulation"; BATHER (1).—Arms of Tribrack crinus; ETHERIDGE, (20) p. 87.—Axial folds of cup-plates as connec with dorsal nervous system, p. 840; radiate ridging of cup-plates tinguished from axial folds (q.v.), p. 841; interradial plates of c their morphological importance, p. 842; arms, their palsontologi development, p. 846; stem of Diamenocrinus compared with that Herpetrocrinus and others, p. 848; stem angles and cirri, Wachsmut Law of Orientation discussed, p. 849; base, dicyclic and monocycle types, p. 852; OEHLERT (39).—Tegmen of Encrinus wagneri careft described and compared with those of recent forms, p. 879; regenerat of arms in Encrinus wagneri, p. 887; arms of Encrinus aculeatus, p. 8 WAGNER (51).—Tegmen, orals of Idiocrinus ventricosus; WACHSMUTI SPRINGER, (50) p. 138.—Structure of dorsal cup in Hypocrinus; Ro PLETZ, (43) p. 75.—Roots of Glyptocrinus and Heterocrinus; DYC (18).—Arms of Cyathocrinus, the branching (p. 218), the axial ca (p. 223), the covering-plates (p. 224); anal tube of Cyathocrinus, p. 2 tegmen of Cyathocrinus, p. 225; BATHER (4).—Anal tube, detadescription of structure in Mastigocrinus, p. 198; stem, altern arrangement of pentameres in Mastigocrinus, p. 200; BATHER (3) Arms, peculiarities of branching, p. 193; anal tube, articulation plates, p. 194, and (3) p. 199; BATHER (2).—Abnormal structure dorsal cup in Millericrinus mespiliformis; BOEHM (14).—Abnor structure of dorsal cup in Encrinus gracilis; Eck (19).—Abnor dorsal cup, 4-rayed, Platycrinus pileatus; REED (41).—Abnormal sti ture of dorsal cup in Hypocrinus pyriformis; ROTHPLETZ, (43) p. 7 Nervous system, dorsal; OEHLERT, (39) p. 840.

Cystidea: Under-side and internal framework of Agelacrinus piles MILLER & FABER, (34) p. 85.

# (b.) Physiology.

Water-pore and madreporic plate, their functions; FIELD, (21) p. 120.—Enlarged *Crinoid* stems; ETHERIDGE, (20) p. 117.—Ejection of Cuvierian organs in *Holothuria*; Minchin (36).

# (c.) GENERAL.

Relation of Crinoid faunas to environment in N. American Palæozoic rocks; Keyes, (27) pp. 291-293.

Variation in species-characters, colour, calcification, &c.; Bell, (8) pp. 520, 521, 523, 527, & 528.

Echinoderm larva, ancestral or comogenetic?; FIELD, (21) p. 118.

Ancestral Echinoderm; FIELD, (21) p. 121.

Radial symmetry secondarily acquired; FIELD, (21) p. 121.

Phylogeny of Synaptida, based on wheel-spicules; LUDWIG, (31) p. 363.

# (d.) DEVELOPMENT.

General Echinoderm embryology: Gonads derived from peritoneal cells, as in other Calomata; MACBRIDE, (33) p. 149.

Corona of Cidaridæ and of Echinoidea Ectobranchiata, its gradual growth; descriptions of the young of Abatus, Goniocidaris, Echinus, Strongylocentrotus; provisional sucking-feet in larval Echinoids, pp. 4, 5, & 12; intestine of larval Echinoids, pp. 4 & 7; skeleton system of larval Echinoids; LOVEN (30).

Asterias vulgaris: Oogenesis, p. 106; spermatogenesis, p. 107; cleavage of ovum, p. 107; mesenchyme formation, pp. 108 & 116; bipinnaria of, p. 108; enteroccel formation, pp. 109 & 115; waterpores in larva of, p. 110; ciliated bands in larva of, p. 112; FIELD (21).

Embryonic development of genital organs, ovoid gland, axial and aboral sinuses in Amphiura squamata; MACBRIDE (33).

Growth of Cyathocrinus; BATHER, (4) p. 223.

Early stages of Antedon; abnormal bigastric embryo of Antedon; development of nervous system in Antedon; SEELIGER (44).

# III.—DISTRIBUTION.

#### A.—GEOGRAPHICAL.

British area, W. Coast of Ireland: Synapta, Psolus, Thyone, Holothuria, Antedon, Astropecten, Psilaster, Luidia, Porania, Asterina, Cribrella, Stichaster, Asterias, Ophiura, Amphiura, Ophiocoma, Ophiothrix, Ophiobyrea, Cidaris, Phormosoma, Asthenosoma, Echinus, Spatangus; BELL (8).

British area, S. W. Ireland: Ophiobyrsa hystricis; Bell (9).

Distribution of European and American Echinoid faunas; GREGORY (22).

Indian Ocean and Archipelago: species of Culcita; HARTLAUB, (25) p. 75.—Indian Ocean, Seychelles: Ophioteresis elegans; Bell (9).—Mauritius: Antedon emendatrix; Bell (10).

N. W. Australia, Bassett-Smith bank: Ophiomaza obscura; Bell (9). Pacific Ocean: species of Culcita; HARTLAUB, (25) p. 75.

General distribution of the genus Culcita; HARTLAUB, (25) p. 75.

#### B.—GEOLOGICAL.

Tertiary: of Australia, Laganum decagonale, Schizaster, Macropneutic decipiens; GREGORY (24).

Pliocene: Echinoid faunas of Europe and America; GREGORY, (22) p. 107.—Pliocene?: Barbados, Cystechinus crassus; JUKES-BROWNE & HARRISON, (26) p. 194.—Tertiary (Miocene or Pliocene), top of Oceanic series, Barbados, Archæopneustes abruptus; GREGORY, (23).

Miocene: Echinoid faunas of Europe and America; GREGORY, (22 p. 105.

Oligocene: Echinoid faunas of Europe and America; Gregory, (22 p. 104.

Eccene: Echinoid faunas of Europe and America; Gregory, (22 p. 104.—Mid. Eccene, Spain: Arachnopleurus reticulatus, D. & Sl. NICKLÈS (37).—Mid. Murravian, Australia, Cassidulus florescens and oth Echinoids; Gregory (24).

Cretaceous: Echinoid faunas of Europe and America; GREGORY, (2 p. 103.—Maëstrichian of Almaceres: Isopneustes heberti, n. sp.; NICKL (37).—(Senonian) Schleswig Holstein: Micraster glyphus, M. haasi, gottschei, M. schræderi, M. gibbus, Echinocorys vulgaris, Offaster pilul O. corculum, Cardiaster lehmanni, Stegaster facki, Echinoconus vulgar Pentacrinus carinatus, P. bicoronatus, Austinocrinus zitteli, A. rothplet A. meyni, Bourgueticrinus, Antedon; doubtful, Asteroidea; STOLLEY (4—(Cenomanien), Rouen: Cidaris cenomanensis; BIGOT (11).

Jurassic:—(Kellaways Rock), Yorkshire, Astropecten orion; WILLIA SON (53).—Normandy: distribution of Echinoids; Bizet (12).—(Ren gerithone), Baden: Balanocrinus pentagonalis and Goniaster impressu Lent (28).—(Oberer weisser Jura), Sontheim a. d. Brenz, Millericrin mespiliformis; Boehm (14).—(Bathonian) p. 42, (Oxfordian) p. 4 (Corallian) p. 59, various Echinoids and Crinoids; (Astartian and Kinmeridgian), Echinoids, Asteroids, Crinoids, and an Ophiurella, pp. 63-8 (Portlandian), various Echinoids, p. 85; (Bas-Boulonnais): Rigaux (4—Sierra Nevada, California: Pentacrinus stems; Mills, (35) p. 428.

Trias: (Lower Muschelkalk), Jena, Encrinus wagneri and E. aculeate Wagner (51).—(Chorzower Beds), Gogolin, Upper Silesia, E. gracile Eck (19).—(Dolomite, above Raibler Beds), Schlern, E. cassianus; WŒH MANN & KOKEN (54).

Permian-Jurassic: Echinoid faunas of Europe and America; Gu GORY, (22) p. 103.

Permian: Timor, Hypocrinus mulleri, H. (?) pyriformis; Timor and Rotti, columnals of Platycrinidæ and other Crinoids; Echinoid spine; ROTHPLETZ (43).

Carboniferous: Echinoid faunas of Europe and America; GREGORY, (22) p. 102.

Permo-Carboniferous: Australia, N.S.W., Archæocidaris, Palæaster, Poteriocrinus, Tribrachiocrinus, Phialocrinus; ETHERIDGE (20).

Carboniferous: Australia, N.S.W., Actinocrinus, Periechocrinus, Platycrinus, Symbathocrinus; Etheridge (20).—(Osage Limestone), Mississippi Basin: change in Crinoid faunas; Keyes, (27) p. 291.—(Keokuk), Mississippi Valley: numerous Crinoids recorded, with remarks on their palseontological development; Beachler, (7) pp. 94–96.—(Lower), Cumberland and Northumberland: Lepidodiscus milleri; Sharman & Newton (46).

Devonian: (Stringocepholus beds), Manitoba, Ctenocrinus; WHITEAVES (52).—(Hamilton Shales), Hamilton, N.Y.: Palæaster eucharis; Cole (15).—Lower Dev. la Manche, France: Ctenocrinus and Diamenocrinus; OEHLERT (39).—Bolivia: undeterminable fragments of Crinoids: ULRICH, (48) p. 83.

Ordovician: Burmah: Echinosphaera; Noetling (38). — Central Piedmont, Virginia: Crinoid remains apparently of Trenton age, and supposed to be allied to Schizocrinus, Heterocrinus, and Poteriocrinus; Darton. —(Hudson R. gr.), Cincinnati: Cyclocystoides and Agelacrinus; MILLER & FABER (34).

Silurian: (Niagara), Indiana: Holocystites affinis; MILLER & FABER (34).—Indiana: Idiocrinus elongatus, I. ventricosus, Hyptiocrinus typus, Callicrinus beachleri, Melocrinus oblongus, M. parvus; Tennessee, Eucalyptocrinus lindahli, Melocrinus rosmeri; Kentucky, Melocrinus oblongus, Wachsmuth & Springer (50).—(Niagara), Indiana: Pisocrinus, Zophocrinus, Stephanocrinus, Caryocrinus, Holocystites; Beachler (6).—(Wenlock), Dudley, England: Botryocrinus quinquelobus, Bather (2); Mastigocrinus loreus, Bather (3); Cyathocrinus acinotubus and C. vallatus, Bather (4).

### IV.—SYSTEMATIC.

#### 1. HOLOTHURIOIDEA.

A + prefixed indicates that the reference or the paragraph following is palseontological.

Acanthotrochus: wheel-spicules; Ludwig, Z. wiss. Zool. liv, p. 358. Bohadschia marmorata, Minchin, Ann. N. H. (6) x, p. 275.

Chiridota, wheel-spicules of, p. 352; C. lævis, Fabr., rotifera, Pourt., purpurea, Less., contorta, Ludw., p. 357; C. pisanii, Ludw., p. 352; C. rufescens, Br., variabilis, Semp., amboinensis, Ludw., liberata, Sluit., p. 354; Ludwig, Z. wiss. Zool. liv.

†Chirodota convexa, Whidb., C. ? gracillima, Whidb.; Woods, Cat. Typ. Foss. p. xvi.

- Holothuria catanensis, impatiens, p. 275, tubulosa, sanctori, p. 281, nigra, p. 273; MINCHIN, Ann. N. H. (6) x. H. nigra, auct., intestinalis, A. & R., tremula, Gunn.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 522. H. poli, Delle Chiaje, forskali, Delle Chiaje, catanensis, Grube (sub forskali), lamperti, Ludw., lagoena, Harcke, klunzingeri, Lampert, pervicax, Sel., marenzelleri, Ludw., marmorata, Jaeg., fusco-cinerea, Jaeg.; H. koellikeri, Semp., noticed: Ludwig & Barthels, Z. wiss. Zool. liv, p. 632.
- Muelleria mauritiana, Quoy. & Gaim., noticed; Ludwig & Barthels, Z. wiss. Zool. liv, p. 632.
- Myriotrochus, wheel-spicules of, M. rinki, Steenstrup; Ludwig, Z. wiss. Zool. liv, p. 359.
- Oligotrochus vitreus, wheel-spicules; Ludwig, Z. wiss. Zool. liv, p. 359.
- Psolus sp., juv.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 521.
- Synapla digitata, Mont.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 521.
- Thyone raphanus, Düb. & Kor.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 522.
- Trochoderma, wheel-spicules; Ludwig, Z. wiss. Zool. liv, p. 358.
- Trochodota venusta, Semon, p. 355, studeri, Theel, p. 358, wheel-spicules; Ludwig, Z. wiss. Zool. liv.

#### 2. ECHINOIDEA.

- †An Echinoid spine: Radiolus radiatus-tubulatus, Permian, Timor, described and figured; ROTHPLETZ, Palssontogr. xxxix, p. 74.
- Abatus cavernosus, Phil.; LOVEN, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 4.
- †Acrosalenia decorata, Haime, Corallian Boulonnais, p. 59, Astartian Boulonnais, p. 63, Kimmeridgian Boulonnais, p. 69, A. spinosa, Ag., lycetti, Wright, pustulata, Forbes, p. 42, hemicidaroides, Wright, legayi, R., pulchella, Cott., porifera, Cott., marioni, Cott., lamarcki, Phill., bradfordiensis, Phill., wastensis, R., lowei, Wright, Bathonian Boulonnais, p. 43, königi, Desm., Portlandian Boulonnais, pp. 80 & 81: RIGAUX. A. rarispina, McCoy; see Hemicidaris minor; A. sp.; Woods, Cat. Type Foss. p. 30.
- †Agassizia; Gregory, Bull. Geol. Soc. Am. iii, pp. 105-107.
- †Anorthopygus orbicularis, Cotteau; Loven, Bih. Sv. Ak. Handl. xviii, (4) No. 1, p. 51.
- Arachnoides placenta, Linn., p. 63, zelandia, Gray, p. 65; LOVEN, Bih. Sv. Ak. Handl. xviii, (4) No. 1.
- † Arachnopleurus reticulatus, Duncan & Sladen, Mid. Eccene, Spain; NICKLES, Ann. Hébert, i, p. 204, pl. viii, figg. 8 & 9.
- † Arbacia inflata, McCoy, see Cottaldia species.
- † Archæocidaris urei, Fleming; Woods, Cat. Type Foss. p. 30.
- †A. ? selwyni, n. sp., Permo-Carboniferous, New South Wales; ETHERIDGE, Pal. Mem. N.S.W. v. p. 67.

- †Archæopneustes, n. g. (type, Palæopneustes hystrix, A. Ag.), with A. abruptus, n. sp., Upper Tertiary, Barbados; Gregory, Q. J. Geol. Soc. xlviii, p. 163.
- †Archaopneustes; GREGORY, Bull. Geol. Soc. Am. iii, p. 107.
- Aspidodiadema antillarum, A. Ag.; Loven, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 32.
- †Asterostoma cubense, Cotteau, referred to Archæopneustes, and jimenoi, Cotteau, referred to Palæopneustes; GREGORY, Q. J. Geol. Soc. xlviii, p. 167.
- Asthenosoma hystrix, Wyv. Th., variation in colour and calcification, A. fenestratum, regarded as syn.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 526. A. varium, Grube; Loven, Bih. Sv. Ak. Handl. xviii (4), No. 1, pp. 26, 45, 52.
- †Australanthus, Bittn., adopted as a subgenus of Cassidulus, by GREGORY, Geol. Mag. (n.s.) Dec. iii, ix, p. 436.
- † Breynella: in Eccene of North America; Gregory, Bull. Geol. Soc. Am. iii, p. 104.
- † Cardiaster suborbicularis, 800 Holaster.
- †Cardiaster lehmanni, n. sp., Senonian, Schleswig-Holstein; Stolley, MT. Min. Inst. Kiel, i, p. 267.
- † Cassidulus florescens, n. sp., Upper Eccene; Gregory, Geol. Mag. (n.s.) Dec. iii, ix, p. 435.
- Centrostephanus coronatus, A. Ag.; LOVEN, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 32.
- Ciduris papillata, Leske; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 526, and Loven, Bih. Sv. Ak. Handl. xviii (4), No. 1, pp. 19 & 52: C. thouarsi, Ag.; Loven, t. c. p. 21.
- †; Cidaris: in Eocene of N. America; GREGORY, Bull. Geol. Soc. Am. iii, p. 104. C. blumenbachi, Münst., florigemma, Phill., houllefortensis, Cott., Oxfordian, Boulonnais, p. 52, C. species, Bathonian, Boulonnais, p. 42, C. coronata, Goldf., Astartian, Boulonnais, p. 67, C. florigemma, Phill., p. 73, beaugrandi, Cott., p. 75, Kimmeridgian, Boulonnais, C. legayi, R. S., p. 81, florigemma, Phill., p. 85, Portlandian, Boulonnais, C. florigemma, Phill., Corallian, Boulonnais, p. 59; RIGAUX (42). C. gaultina, Forbes, p. 30, sedgwicki, Seeley, C. species, p. 31; Woods, Cat. Type Foss. C. melitensis, Wright; GREGORY, Bull. Geol. Soc. Am. iii, p. 105. C. cenomanensis, Cott.; BIGOT, Bull. Soc. L. Norm. (4) v, p. 54.
- Clypeastridæ: dental system; Loven, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 55.
- Clypeaster reticulatus, Linn.; LOVEN, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 55.
- †Clypeus altus, McCoy, excentricus, McCoy (sub C. sinuatus), sinuatus, Leske; Woods, Cat. Type Foss. p. 31. C. plotii, Klein, milleri, Wright, Bathonian, Boulonnais; RIGAUX, p. 42.
- †Collyrites bicordata, Des., Oxfordian, Boulonnais; RIGAUX, p. 51. C. ovalis, Leske; Woods, Cat. Type Foss. p. 31.

- "Cataline species: Wiros. Cat. Type Fost, p. 31.
- Copiesans apraeuralisma, Cott., Astartian, Boulonnais, p. 65, C. apraeuralisma. Cott., Kimmeridgian, Boulonnais, pp. 72 & 73, C. apray. Exg. Particulante. Boulonnais, p. 81; Rigaux. C. impress, Suday, see Ecimocophus.
- Transforms Street, J. RES-BROWKE & HARRISON, Q. J. Gool. Sec. Evil. p. 134. and Greecent, Bull. Gool. Soc. Am. iii, p. 107.
- Dienienes sermine. Linn.; Lyves, Bib. Sv. Ak. Handl. xviii (4), No. 1, pp. 31. 34. 51.
- -: Dientema funguideum, ami entertuberculatum, Seeley, see Pseudodiadem.

  D. Vetrugrunnus? scriptum, Seeley, see Diplopodia.
- -Dipimpada penanguna. M::Coy. acriptum [sic] Seeley; Woods, Cat. Type F.m. 2. K.
- -: Lournius spinairies Ag.; Loven, Bih. Sv. Ak. Handl. xviii (4), No. 1. 30. 47, 52. D. marginalis, McCoy, see Holectypus hemisphericus. D. margin. Dence; Woods, Cat. Type Poss. p. 31.
- -Dynamer symmetricus. McCoy, see Collyrites ocalis.
- \* Echemotius, = B-species; Gregory, Bull. Geol. Soc. Am. iii, p. 214. E. prinspagiaria. Say, discussed; Gregory, t. c. p. 105. E. rescuincia, Linn.: Gregory, t. c. p. 107.
- \*: Echanus schmiss: in Econo of N. America; Gregory, Bull. Geol Sig. Am. 33, p. 134. E. econoricus; Gregory, t. c. p. 107.
- Echimorachnius purme. Lamk.; Loves, Bih. Sv. Ak. Handl. xviii (4) No. L. p. 66.
- † ; Echmidrianis priesdechi. Wright, clanicularis, Des., goldfussi, Des. empirus. Des., elimpatus. d'Orh., woodwardi, Wright, quadratu Wright. Bathenian. Boulonnais. p. 42 : E. scutatus, Lam., Corallian Brukennais. p. 59 : E. scutatus. Lam., Astartian, Boulonnais. p. 65 E. perroni. Cott.. Kimmeridgian, Boulonnais, p. 72 : E. haime Wright, Bolonian, Boulonnais, p. 78 : E. rigauxi, Cott., p. 8 brukei. Wright, Portlandian, Boulonnais, p. 85 ; RIGAUX (42).
- Echinocideris nigra, Mol.; Loves, Bih. Sv. Ak. Handl. xviii (4), No. p. 33.
- †: Echinocoms raiguris, d'Orb.: STOLLEY, MT. Min. Inst. Kiel, i, p. 26.

  E. subrotundus, Mantell; Woods, Cat. Type Foss. p. xvi.
- † Echin:corys rulgaris, Breyn., (= E. scutatus); STOLLEY, MT. Min. Ins Kiel, i, p. 284.
- Echinocyumus angulosus, Leske; Loven, Bih. Sv. Ak. Handl. xviii (4 No. 1, p. 69.
- † Echinocyphus impressus, Seeley; Woods, Cat. Type Foes. p. 32.
- Echinometra mathæi, Blainv., p. 33, lucunter, Linn., pp. 43, 44, 51, 52, 55 Loves, Bih. Sv. Ak. Handl. xviii (4), No. 1.
- Echinus acutus, Lamk., esculentus, Linn., elegans, Düb. & Kor., p. 52 microstoma, Wyv. Th., p. 529; Bell, Sci. P. R. Dubl. Soc. (n.s.) vi E. parvituberculatus, Blainv.; Loven, Bih. Sv. Ak. Handl. xviii (4 No. 1, p. 11. E. miliaris, Leske, p. 161, esculentus, Linn., pp. 33, 5 angulosus, Leske, p. 33; Loven, Bih. Sv. Ak. Handl. xviii (4), No.

- †; E. diademata, McCoy, see Stomechinus germinans. E. petallatus, McCoy, see S. gyratus.
- † Enallaster columbianus, Lea, karsteni, de Lor.; GREGORY, Bull. Geol. Soc. Am. iii, p. 103.
- Encope emarginata, Leske; LOVEN, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 65.
- † Eocidaris, abandoned; GREGORY, Bull. Geol. Soc. Am. iii, p. 102.
- † Eupatagus decipiens, Tate; see Macropneustes.
- † Euspatangus, in Eocene of N. America; Gregory, Bull. Geol. Soc. Am. iii, p. 104.
- † Galerites albogalerus, Leske; Loven, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 49.
- † Galeropygus agariciformis, Forbes; Woods, Cat. Type Foss. p. 32.
- †Goniophorus lunatus, Ag., var. minutus, Seeley; Woods, Cat. Type Foss. p. 32.
- † Glypticus hieroglyphicus, Ag., Corallian, Boulonnais; RIGAUX (42), p. 59. Goniocidaris canaliculata, A. Ag.; LOVEN, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 5.
- †Hardouinia, adopted as subgenus of Breynella, by GREGORY, Geol. Mag. (n.s.) iii, ix, p. 436.
- †Hemiaster mccoyi, Seeley; Woods, Cat. Type Foss. p. 32.
- †; Hemicidaris confluens, McCoy; see Acrosalenia sp. H. minor, Ag.; Woods, Cat. Type Foss. p. 32. H. luciensis, d'Ord., legayi, R., langrunensis, Cott., Bathonian, Boulonnais, p. 42, intermedia, Fordes, Oxfordian, Boulonnais, p. 52, intermedia, Fordes, Corallian, Boulonnais, p. 59, hoffmanni, Ag., p. 65, agassizi, Röm., intermedia, Fordes, Astartian, Boulonnais, p. 67, intermedia, Fordes, p. 72, davidsoni, Wright, p. 75, Kimmeridgiau, Boulonnais, purbeckensis, Fordes, Bolonian, Boulonnais, p. 78, davidsoni, brillensis, morinicum, R. S, Portlandian, Boulonnais, pp. 80 & 81; RIGAUX (42).
- †Hemipedina tuberculosa, Wright, Oxfordian, Boulonnais, p. 52, bouchardi, Wright, Portlandian, Boulonnais, p. 81; RIGAUX.
- †; Holaster simplex, Shumard, = H. lævis, Luc.; Gregory, Bull. Geol. Soc. Am. iii, p. 104. H. suborbicularis, Brong.; Woods, Cat. Type Foss. p. 32.
- †; Holectypus depressus, Des., Bathonian, Boulonnais, p. 42, depressus, Phill., Oxfordian, Boulonnais, p. 51, corallinus, d'Orb., Astartian, Boulonnais, pp. 63 & 65, corallinus, d'Orb., Corallian, Boulonnais, p. 59; RIGAUX. H. depressus, Leske; LOVEN, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 51. H. planatus, Roem.; GREGORY, Bull. Geol. Soc. Am. iii, p. 104. H. hemisphæricus, Ag.; Woods, Cat. Type Foss. p. 32.
- † Isopneustes heberti, Maëstrichian of Almaceres, NICKLES, Ann. Hébert. i, p. 202, pl. ix, figg. 12 & 13, n. sp.
- †Laganum decagonale, Lesson, n. var. rictum, Tertiary, W. Australia; Gregory, Geol. Mag. (n.s.), Dec. iii, ix, p. 433. L. decagonale, Blainv.; Loven, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 68.

- Lytechinus semitubercula:us, Val.; LOVEN, Bih. Sv. Ak. Handl. xviii No. 1, p. 33.
- † Macropneustes decipiens, Tate, olim Eupatagus; GREGORY, Geol. 1 (n.s.), Dec. iii, ix, p. 436.
- † Mellita sexforis; GREGORY, Bull. Geol. Soc. Am. iii, p. 107.
- †; Micraster glyphus, Cotteau, p. 255, gibbus, Lam., p. 260; Stoll MT. Min. Inst. Kiel, i.
- †M. haasi, p. 257, gottschei, p. 258, schræderi, Senonian, Schleswig-H stein, Stolley, MT. Min. Inst. Kiel, i, p. 259, n. spp.
- † Monostychia, in Eocene of N. America; GREGORY, Bull. Geol. S. Am. iii, p. 104.
- †Mortonia rogersi, Desor, synonymy discussed; GREGORY, Bull. G. Soc. Am. iii, p. 105.
- †; Nucleolites æqualis, McCoy, clunicularis, Phill., and pyramidalis, McC planulatus, McCoy; Woods, Cat. Type Foss. p. 32.
- †Offaster pilula, Lam., corculum, Goldf.; STOLLEY, MT. Min. It Kiel, i, p. 265.
- †Palæchinus gigas, McCoy, p. 32, intermedius, Keeping, spharic McCoy, p. 33; Woods, Cat. Type Foss.
- †; Pelanechinus: LOVEN, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. P. corallinus, Wright; WOODS, Cat. Type Foss. p. 33.
- †Pedina legayi, Cott, Bathonian, Boulonnais, p. 42, sublevis, Ag, Corlian, Boulonnais, p. 59; RIGAUX.
- † Peltastes wrighti, Desor, p. xvi, hieroglyphica, Keeping, wiltshirei, Seelwrighti, Desor, p. 33; Woods, Cat. Type Foss.
- †; Periarchus, accepted; GREGORY, Bull. Geol. Soc. Am. iii, p. 16 P. altus, Conrad; GREGORY, t. c. p. 105.
- † Perischodomus biserialis, McCoy; Woods, Cat. Type Foss. p. 33.
- Phormosoma placenta, Wyv.-Th.; Bell, Sci. P. R. Dubl. Soc. (n.s.), v p. 526.
- †Pileus hemisphæricus, Desor; Loven, Bih. Sv. Ak. Handl. xviii ( No. 1, p. 51.
- †Plesiospatangide, revised; GREGORY, Q. J. Geol. Soc. xlviii, p. 165.
- †Polycyphus normannus, Des., Bathonian, Boulonnais, p. 42; RIGAUX. †Pseudasterostoma jimenoi, see Asterostoma jimenoi.
- †Pseudocidaris quenstedti, Cott., Oxfordian, Boulonnais, p. 52, thurman Et., Corallian, Boulonnais, p. 59; RIGAUX.
- †; Pseudodiadema depressum, Ag., subcomplanatum, Des., Bathonian, Bolonnais, p. 42, superbum, Des., pellati, Cott., princeps, Des., arovien Des., bipunctatum, Des., Oxfordian, Boulonnais, p. 52, florescens, A hemisphæricum, Ag., Corallian, Boulonnais, p. 59, mamillatum, Rör p. 63, planissimum, Astartian, Boulonnais, p. 67, mamillatum, Rör Kimmeridgian, Boulonnais, p. 75, planissimum, Bolonian, Boulonais, p. 78, sauvagei, Rig., baccatum, R. S., p. 81, thirriæ, Etall., Pollandian, Boulonnais, p. 85; RIGAUX. P. barretti, S. P. Woodvarteri, S. P. Woodw., fungoideum, Seeley, p. 33, intertuberculatu. Seeley, inversum, Seeley, p. 34; Woods, Cat. Type Foss.

- †; Pygaster laganoides, Ag., trigeri, Cott., Bathonian, Boulonnais, p. 42
  umbrella, Astartian, Boulonnais, p. 66, macrocyphus, Wright, Kimmeridgian, Boulonnais, p. 75, gresslyi, Des., Corallian, Boulonnais, p. 59; RIGAUX (42). P. semisulcatus, Phill.; LOVEN, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 51. P. brevifrons, McCoy, see P. semisulcatus. P. semisulcatus, Phill.; Woods, Cat. Type Foss. p. 34. P. sublævis, McCoy, see Galeropygus agariciformis.
- †Pygurus michelini, Cott., Bathonian, Boulonnais, p. 42, blumenbachi, Ag., Corallian, Boulonnais, p. 59, Kimmeridgian, Boulonnais, pp. 70 & 72, and Astartian, Boulonnais, pp. 63 & 65, jurensis, Marc., Astartian, Boulonnais, pp. 66 & 67, Kimmeridgian, Boulonnais, p. 75; RIGAUX (42).
- †; Rhabdocidaris orbignyana, Des., boloniensis, Wright, Astartian, Boulonnais, p. 67, Kimmeridgian, Boulonnais, pp. 69 & 72; RIGAUX. R. thurmanni, de Lor., var. regens, Whidb.; Woods, Cat. Type Foss. p. 34.
- †Rhoechinus irregularis, Keep.; Woods, Cat. Type Foss. p. 34.
- †Rhynchopygus guadaloupensis, Mich., = R. caribbæarum; GREGORY, Bull. Geol. Soc. Am. iii, p. 105.
- Salenia pattersoni, Al. Ag.; LOVEN, Bih. Sv. Ak. Handl. xviii (4), No. 1, p. 34.
- †; Salenia hieroglyphica, Keep., and S. (Hyposalenia) wiltshirei, Seeley, see Peltastes; S. woodwardi, Seeley; Woods, Cat. Type Foss. p. 34.
- †Sarsella, in Eocene of N. America; Gregory, Bull. Geol. Soc. Am. iii, p. 104.
- †Schizaster parkinsoni, Defrance, scillæ, Desmoul.; GREGORY, Bull. Geol. Soc. Am. iii, pp. 105 & 107. S. cuneatus, Greg.; Woods, Cat. Type Foss. p. xvi. S., sp., W. Australia; GREGORY, Geol. Mag. (n.s.) Dec. iii, ix, p. 436.
- †; Scutella striatula, Rem. (pars) = Echinarachnius excentricus; S. rogersi, Morton, discussed; GREGORY, Bull. Geol. Soc. Am. iii, pp. 105 & 107. S. quinquefaria, Say; see Echinanthus.
- †Sismondia marginalis and plana, Conrad, = Monostychia; GREGORY, Bull. Geol. Soc. Am. iii, p. 105.
- Spatangus purpureus, O. F. Müll., raschi, Loven; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 529.
- †Spatangus columbianus, Lea; see Enallaster.
- †Stegaster facki, n. sp., Senonian, Schleswig Holstein; Stolley, MT. Min. Inst. Kiel, i, p. 268.
- †; Stomechinus serratus, Des., Bathonian, Boulonnais, p. 42, S. gyratus, Des., Oxfordian, Boulonnais, p. 52, S. gyratus, Ag., Corallian, Boulonnais, p. 59 (and Woods, Cat. Type Foss. p. 34), S. perlatus, Ag., p. 63, Astartian, Boulonnais, S. perlatus, Des., Astartian, Boulonnais, p. 67, and Kimmeridgian, Boulonnais, p. 73; RIGAUX. †S. germinans, Phill.; Woods, Cat. Type Foss. p. 34.
- Strongylocentrotus droebachiensis, O. F. Müll.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 523, and Loven, Bih. Sv. Ak. Handl. xviii (4), No. 1, pp. 13 & 15.

† Temnechinus; GREGORY, Bull. Geol. Soc. Am. iii, p. 108. Tripneustes gratilla, Linn., p. 33, esculentus, Leske, p. 38; LOVEN, J Sv. Ak. Handl. xviii (4), No. 1.

### 3. ASTEROIDEA.

Asterina gibbosa, Penn.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 525.

Asterias rubens, Linn., murrayi, Bell; Bell, Sci. P. R. Dubl. Soc. (n. vii, p. 525.

A. vulgaris: embryology; Field, Q. J. Micr. Sci. (n. xxxiv, p. 105.

Astropecten sphenoplax, W. Coast of Ireland; Bell, Sci. P. R. D. Soc. (n.s.) vii, p. 522, n. sp.

Astropecten irregularis, Penn.; Bell, Sci. P. R. Dubl. Soc. (n.a.) p. 522.

†; Astropecten orion, Forbes; WILLIAMSON, Proc. Geol. Soc. 1892, p. 1 A. lorioli, and martis, Wright, A. suprajurensis, Sauv., Bolon Boulonnais, p. 78; RIGAUX. A. scarburgensis, Wright, A. hastin Forbes, p. 35, rectus, McCoy; Woods, Cat. Type Foss, p. 35.

Cribrella sanguinolenta, O. F. Müll.; Bell, Sci. P. R. Dubl. Soc. (n.s.)

p. 525.

Culcita veneris, Perr., pp. 73 & 96, coriacea, M. T., pp. 74 & 90, n guinea, M. T., pp. 74 & 77, grex, M. T., pp. 74 & 87, acutispin Bell, pp. 74 & 95, arenosa, Perr., pp. 74 & 92, pentangularis, G p. 74, C. schmideliana, Retz., pp. 75 & 97; HARTLAUB, Notes I Mus. xiv. C. discoidea, Lam.; see C. schmideliana. C. pulverul Perr.; see C. novæ-guinea.

Culcita plana; HARTLAUB, Notes Leyd. Mus. xiv, pp. 74 & 84, Sa Viti, New Guinea, Phillipines, Amboina, Mauritius, n. sp.

†Goniaster cassiope, Kimmeridgian, Boulonnais, p. 72; RIGAUX. G pressæ, Quenst.; Lent (28). G. (Goniodiscus) rectilineus, Mct see Pentagonaster.

Luidia sarsi, Dub. & Kor., ciliaris, Phil.; Bell, Sci. P. R. Dubl. (n.s.) vii, p. 525.

†Monaster, n. subg. of Palæaster; ETHERIDGE, Pal. Mem. N.S.V p. 70.

†Palæaster (Monaster) stutchburii and P. (Monaster) giganteus, Pe Carboniferous, New S. Wales; ETHERIDGE, Pal. Mem. N. v, pp. 73 & 74, n. spp.

†Palæaster (Monaster) clarkei, De Kon., Permo-Carboniferous, Ne Wales; ETHERIDGE, Pal. Mem. N.S.W. v, p. 71. P. eucharis, oral surface; Cole, Bull. Geol. Soc. Am. iii, p. 512. P. hii ruthveni, Forbes, squamatus, Salter; Woods, Cat. Type 2, 25.

† Palæasterina primæva, Forbes, ramseyensis, Hicks; Woods, Cat. 'Foss. p. 35.

† Pentagonaster rectilineus, McCoy; WOODS, Cat. Type Foss. p. 36.

†Plumaster ophiuroides, Wright; Woods, Cat. Type Foss. p. 36.

- Porania pulvillus, O. F. Müll.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 525.
- Psilaster andromeda, M. & T.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 524.
- Randasia spinulosa, Gray, see Culcita coriacea; R. granulata, Gray, see C. pentangularis. B. luzonica, Gray, = Culcita, sp.; HARTLAUB, Notes Leyd. Mus. xiv, p. 77.
- Stichaster roseus, O. F. Müll.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 525.
- †Uraster carinatus, Wright; Woods, Cat. Type Foss. p. 36. U. hirudo, ruthveni, Forbes, see Palæaster; U. primævus, Forbes, see Palæasterina.

#### 4. OPHIUROIDEA.

- New classification of Ophiuroids, based on articulation of arm-ossicles, into Streptophiuræ, Astrophiuræ, Zygophiuræ, and arrangement of genera under these groups according to characters of minor importance; Bell, P. Z. S. 1892.
- Amphiura squamata, M. T.; MACBRIDE, Q. J. Micr. Sci. (n.s.) xxxiv, p. 129 (embryology and anatomy). A. filiformis, O. F. Müll.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 526.
- Astrophiura, new ordinal name; Bell, P. Z. S. 1892, p. 180.
- † Eucladia, H. Woodw., removed from Ophiuroidea; Bell, t. c. p. 182.
- † Onychaster, M. & W., position discussed; Bell, t. c. p. 182.
- Ophiobyrsa hystricis, Lym.; Bell., P. Z. S. 1892, p. 183, and Sci. P. R. Dubl. Soc. (n.s.) vii, p. 526.
- Ophiocoma nigra, O. F. Müll.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 526.
- †Ophioderma carinata, Wright; Woods, Cat. Type Foss. p. 36.
- +Ophiolepis leckenbyi, Wright; Woods, Cat. Type Foss. p. 36.
- Ophiomaza obscura, Ljung.; Bell, P. Z. S. 1892, p. 182.
- Ophioteresis, n. g. for O. elegans, n. sp., Seychelles; Bell, P. Z. S. 1892, p. 179.
- Ophiothrix fragilis, Linn., and leutkeni, Wyv.-Thom.; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 526.
- Ophiura ciliaris, Linn., and albida, Forbes; Bell, Sci. P. R. Dubl. Soc. (n.s.) vii, p. 525.
- +Ophiura salteri, Forbes; see Protaster.
- †; Protaster, Forbes, referred to Streptophiuræ; Bell, P. Z. S. 1892, p. 182; P. petri, Salter, salteri, Forbes, p. 36, sedywicki, Forbes, p. 37; Woods, Cat. Type Foss.
- Streptophiura, new ordinal name; Bell. P. Z. S. 1892, p. 179.
- † Taniaster, Bill., referred to Streptophiura; Bell, t. c. p. 182.
- Zygophiura, new ordinal name; Bell, t. c. p. 180.

#### 5. CRINOIDEA.

- Introduction to anatomy, with special reference to Fossils; BATHER, (5) p. 87.
- †; Actinocrinus, ind. sp., Carboniferous, New S. Wales; ETHERIDGE, Pal. Mem. N.S. W. v, p. 77. A. pulcher, McCoy (? Periechecrinus); WOODS, Cat. Type Foss. p. 37.
- †Actinometra loveni, wurtembergiæ, P. H. Carp.; Woods, Cat. Type Foss. p. 37.
- † Amphoracrinus atlas, olla, McCoy; Woods, Cat. Type Foes. p. 37.
- Antedon bifida, Penn.; Bell, Sci. P. R. Dubl. Soc. (n.s.), vii, p. 522.

  A. rosacea, embryology; Seeliger (44).
- Antedon emendatrix, Mauritius, Bell, Ann. N. H. (6) ix, p. 427, n. sp.
- †; Antedon sp.; Stolley, MT. Min. Inst. Kiel, i, p. 255. A. aspera, Quenst., costata, Goldf., depressa, P. H. Carp., scrobiculata, Goldf.; Woods, Cat. Type Foss. p. 37.
- †; Apiocrinus elegans, Defr., Bathonian, Boulonnais, p. 43, A. roissyanus, d'Orb., Corallian, Boulonnais, p. 59; RIGAUX. A. exutus, McCoy; WOODS, Cat. Type Foss. p. 38.
- † Archaocrinus, OEHLERT, Bull. Soc. Géol. (3) xix, p. 840.
- †Austinocrinus sitteli (columnals only), p. 251, rothpletzi (columnals only), p. 252, meyni (columnals only), Senonian, Schleswig-Holstein, Stolley, MT. Min. Iust. Kiel, i, p. 254, n. spp.
- †Balanocrinus pentagonalis, Goldf.; LENT, p. 624.
- † Botryocrinus pinnulatus, Bather, additional description; BATHER, Ann N. H. (6) ix, p. 192. B. decadactylus (Salter, MS.), Bather; Woods Cat. Type Fors. p. 38.
- †B. quinquelobus, Wenlock Limestone, Dudley, BATHER, Ann. N. H (6) ix, p. 189, n. sp.
- †Bourgueticrinus cylindricus, milleri, politicus, McCoy; WOODS, Cat. Type Foss. p. 38. B. ellipticus, B. sp., Mill.; STOLLEY, MT. Min. Inst Kiel, i, p. 254.
- † Calceocrinus abdominalis, fletcheri, gradatus, Salter, serialis (Austin MS sp.); Woods, Cat. Type Foss. p. 38.
- † Callicrinus beachleri, n. sp., Silurian, Indiana, WACHSMUTH & SPRINGER Am. Geol. x, p. 140.
- †Caryocrinus ornatus, Say; BEACHLER, Am. Geol. ix, p. 408.
- † Cheirocrinus abdominalis, fletcheri, gradatus, serialis, Salter: see Calceo crinus.
- †Cromyocrinus nuciformis, McCoy; Woods, Cat. Type Foss. p. 38.
- †; Clenocrinus sp., Devonian, la Manche; ŒHLERT, Bull. Soc. Géol. (3 xix, p. 835. C. sp., Mid. Devon., Manitoba; WHITEAVES, Contr. Canad. Pal. i, p. 276.
- †Cupressocrinus calyx, McCoy; see Hydreionocrinus; C. impressus, McCoy see H. mccoyanus.
- †; Cyathocrinus, discussed and re-diagnosed; Bather, Ann. N. H. (6 ix, p. 202: American and Gotland species; Bather, t. c. p. 213

- C. distortus, Phill.; see C. planus. C.? Carboniferous, New S. Wales; Etheridge, Pal. Mem. N.S.W. v, p. 81. C. quinquangularis, Phill., referred to Botryocrinus quinquelobus; Bather, Ann. N. H. (6) ix, p. 189. C. arboreus, Salter, nom. nud., referred to Mastigocrinus loreus; Bather, t. c. p. 196. C. acinotubus, Ang.; Bather, t. c. p. 213, and Woods, Cat. Type Foss. p. 39. C. arboreus, scoparius, C. sp. 11, and C. spp., Salter; Woods, Cat. Type Foss. pp. xvi & 39. C. decadactylus, Salter; see Botryocrinus. C. monile; see C. acinotubus; C. nodulosus, Salter, = Taxocrinus sp.; Woods, Cat. Type Foss. p. 39. C. quindecimalis, Salter; see Botryocrinus decadactylus. C. squamiferus, Salter, = Gissocrinus; Woods, Cat. Type Foss. p. 39. C. vallatus, Wenlock Limestone, Dudley; Bather, Ann. N. H. (6) ix, p. 221. C. plunus, Miller, determined; Bather, t. c. p. 202. C. alutaceus (pars), Ang., monile, Salter, nom. nud., mimus, Salter, nom. nud., nodulosus, Salter, nom. nud.; see C. acinotubus.
- † Cyclocrinus variolarius, Seeley; Woods, Cat. Type Foss. p. 39.
- †Dadocrinus kunischi, W. & Sp.; Eck, Z. geol. Ges. xliv, p. 138.
- † Diamenocrinus, n. g., p. 836, for D. jouani, n. sp., Devonian, la Manche, p. 837; ŒHLERT, Bull. Soc. Géol. (3) xix.
- † Dendrocrinus? cambrensis, Hicks; Woods, Cat. Type Foss. p. 40.
- † Dimerocrinus uniformis, Salter; D. multiplex, Salter, = Encrinus speciosus, Ang.; Woods, Cat. Type Foss. p. 40.
- †; Encrinus wagneri, Ben., p. 879, aculeatus, Mey., p. 890; WAGNER, Z. geol. Ges. xliii. E. cassianus, Laube; WOEHRMANN & KOKEN, op. cit. xliv, p. 170. E. gracilis; see Dadocrinus kunischi.
- † Eucalyptocrinus lindahli, n. sp., Silurian, Tennessee, Wachsmuth & Springer, Am. Geol. x, p. 139.
- † Extracrinus fossilis, Blumenbach, dichotomus, McCoy; Woods, Cat. Type Foss, p. 40.
- †; Gissocrinus, British species; BATHER, Ann. N. H. (6) ix, p. 207. G. goniodactylus, Phillips; Woods, Cat. Type Foss. p. 40.
- †Glyptasteridæ, ŒHLERT, Bull. Soc. Géol. (3) xix, pp. 839, et seq.
- †; Glyptocrinus neali, dyeri, baeri; DYCHE, Am. Geol. x, p. 130. G. basalis, McCoy; G. sp. 1, Salter; Woods, Cat. Type Foss. p. 40.
- †; Herpetocrinus; ŒHLERT, Bull. Soc. Géol. (3) xix, pp. 848 & 849. H. fetcheri, Salter; WOODS, Cat. Type Foss. p. 40.
- †; Heterocrinus subcrassus, simplex; DYCHE, Am. Geol. x, p. 130. H.? sp., Ordovician, Piedmont, Virginia; DARTON, Am. J. Sci. xliv, p. 52.
- † Hexacrinida; ŒHLERT, Bull. Soc. Geol. (3) xix, pp. 841, et seq.
- † Hydreionocrinus calyx, McCoy, mccoyanus, De Kon. & Le Hon.; Woods, Cat. Type Foss. p. 40.
- † Hypocrinus mülleri, Beyr.; ROTHPLETZ, Palseontogr. xxxix, p. 74.
- †H. (?) pyriformis, Permian, Timor, Rothpletz, Palmontogr. xxxix, p. 75,
- †Hyptiocrinus, n. g., for H. typus, n. sp., Silurian, Indiana, Wachsmuth & Springer, Am. Geol. x, p. 138.
- † Ichthyocrinus bacchus, mccoyanus, Salter; Woods, Cat. Type Foss. p. 41.

- † Idiocrinus, n. g., p. 135, for I. elongatus, p. 136, I. ventricosus, p. 137, n. spp., Silurian, Indiana; Wachsmuth & Springer, Am. Geol. x.
- †Koninckocrinus agassizi, Seeley; see Torynocrinus canon. K. rugosus, d'Orb.; see Cyclocrinus variolarius.
- † Mariacrinus flabellatus, Salter; see Melocrinus.
- † Mastigocrinus, n. g., p. 195, for M. loreus, n. sp., p. 196, Wenlock Limestone, Dudley; BATHER, Ann. N. H. (6) ix.
- † Melocrinus flabellatus, Salter; Woods, Cat. Type Foss. p. 41.
- † M. roemeri, Silurian, Tennessee, p. 142, oblongus, Silurian, Iudiana and Kentucky, p. 143, parvus, Silurian, Indiana, p. 144, Wachsmuth & Springer, Am. Geol. x, n. spp.
- †; Millericrinus mespiliformis, Schloth., abnormality; Вовни, Z. geol. Ges. xliii, p. 741. M. prutti, Gray; Woods, Cat. Type Foss. p. 41. M. horridus, d'Orb., Oxfordian, Boulonnais; RIGAUX, p. 52.
- † Myelodactylus; ŒHLERT, Bull. Soc. Géol. (3) xix, pp. 848 & 849: see also Herpetocrinus.
- †Ottawacrinus compared with Mastigocrinus; BATHER, Ann. N. H. (6) ix, p. 201.
- †Palæocrinus, discussed; BATHER, Ann. N. H. (6) ix, p. 207.
- † Pentacrinites briareus, Miller; see Extracrinus fossilis, McCoy.
- † Pentacrinus pentagonalis, Goldf., normanianus, Lor., cotteaui, Lor., Oxfordian, Boulonnais, p. 52, P. amblyscalaris, Thurm., Corallian, Boulonnais, p. 59, P. rupellensis, Lor., Astartian, Boulonnais, p. 67; RIGAUX (42). P. carinatus, Roem., bicoronatus, Hag.; STOLLEY, MT. Min. Inst. Kiel, i, p. 249. P. sp., columnals in Jurassic, California; MILLS, Bull. Geol. Soc. Am. iii, p. 428. P. goldfussi, McCoy; Woods, Cat. Type Foss. p. 41. P. dichotomus; see Extracrinus.
- †Phialocrinus re-defined, p. 98, konincki, de Kon., p. 107; ETHERIDGE, Pal. Mem. N.S.W. v.
- †Phialocrinus princeps, p. 110, P. ? nodosus, p. 112, stephensi, p. 113, Permo-Carboniferous, New S. Wales, ETHERIDGE, Pal. Mem. N.S.W. v., n. spp.
- †Periechocrinus indicator, Carboniferous, New S. Wales, ETHERIDGE, Pal. Mem. N.S.W. v, p. 78, n. sp.
- †; P., sp. ind., Carboniferous, New S. Wales; ETHERIDGE, Pal. Mem. N.S.W. v, p. 79. P. limonium, Salter; Woods, Cat. Type Foss, p. 41.
- †Picteticrinus nicoleti, Des., Bathonian, Boulonnais, p. 43, beaugrandi, Lor., Portlandian, Boulonnais, p. 81; RIGAUX.
- † Pisocrinus gemmiformis, S. A. Mill., globosus, Ringueb.; BEACHLER, Am. Geol. ix, p. 408.
- †Platycrinidæ, and other Crinoids, "Entrochi regulares et irregulares," columnals from Permian, Timor, described and figured; ROTHPLETZ, Palæontogr. xxxix, p. 72.
- †; Platycrinus pileatus, Goldf., abnormal; REED, Geol. Mag. (iii) ix, p. 548. P. diadema, megastylus, McCoy, P.? pecten, Salter, P.

- vesiculosus, McCoy; Woods, Cat. Type Foss. p. 41. P. sp., Carboniferous, New S. Wales; ETHERIDGE, Pal. Mem. N.S.W. v, p. 131.
- †; Poteriocrinus?, Carboniferous, New S. Wales; ETHERIDGE, Pal. Mem. N.S.W. v, p. 81. P. cressimanus, McCoy; Woods, Cat. Type Foss. p. 42. P.? sp., Ordovician, Piedmont, Virginia; Darton, Am. J. Sci. xliv, p. 52. P. excavatus, McCoy; see Hydreionocrinus mccoyanus. P. nuciformis, McCoy; see Cromyocrinus. P. longiductylus; see Scaphiocrinus.
- † Reteocrinidæ; ŒHLERT, Bull. Soc. Géol. (3) xix, pp. 839, et seq.
- †Raphanocrinus; ŒHLERT, Bull. Soc. Géol. (3) xix, p. 840.
- †Rhodocrinidæ; ŒHLERT, Bull. Soc. Géol. (3) xix, p. 839.
- †Scaphiocrinus longidactylus, Austin; BATHER, Ann. N. H. (6) ix, p. 203.
- † Schizocrinus, ? sp., Ordovician, Piedmont, Virginia; Darton, Am. J. Sci. xliv, p. 52.
- †Solanocrinus beaugrandi, Lor., Kimmeridgian, Boulonnais; RIGAUX, p. 69.
- †Sphærocrinus discussed; BATHER, Ann. N. H. (6) ix, p. 207.
- †Stephanocrinus osgoodensis, S. A. Mill.; BEACHLER, Am. Geol. ix, p. 408.
- †Symbathocrinus, Carboniferous, New S. Wales; ETHERIDGE, Pal. Mem. N.S.W. v, p. 81.
- †; Taxocrinus nanus, Salter, Porbignii, McCoy, tuberculatus, Miller, Pgranulatus, Salter; Woods, Cat. Type Foss. p. 42. T. longidactylus; see Scaphiocrinus. T. marmoratus, Salter; see T. tuberculatus.
- †; Torynocrinus canon, Seeley; Woods, Cat. Type Foss. p. 43. T. variolarius, Seeley; see Cyclocrinus.
- † Thenarocrinus, referred to the Dendrocrinites; BATHER, Ann. N. H. (6) ix, p. 202.
- † Tribrachiocrinus re-described, p. 82; T. clarkei, McCoy, p. 90; T. corrugatus, Ratte, p. 95; ETHERIDGE, Pal, Mem. N.S.W. v.
- †Tribrachiocrinus ornatus, p. 94, granulatus, p. 97, Permo-Carboniferous, New S. Wales, ETHERIDGE, Pal. Mem. N.S.W. v, n. spp.
- †Zophocrinus howardi, S. A. Mill.; BEACHLER, Am. Geol. ix, p. 408.

### 6. CYSTIDEA.

- † Agelacrinus pileus?, Hall; MILLER & FABER, J. Cincinn. Soc. xv, p. 85.
- † Atelecystis forbesi, De Kon.; Woods, Cat. Type Foss. p. 42.
- † Curyocystites davisi, McCoy; see Echinosphæra.
- Cyclocystoides cincinnatiensis, Ordovician, Cincinnati, MILLER & FABER, J. Cincinn. Soc. xv, p. 84, n. sp.
- †C., sp., MILLER & FABER, J. Cincinn. Soc. xv, p. 85.
- †; Echinosphæra davisi, McCoy; Woods, Cat. Type Foss. p. 43. E. kingi: undescribed species; Noetling (38).

- "Astrophics office 1.14. Mineral Indiana. Ministe & Fabre J. Circle for 3 1 5. The range of the game district be Mossych; a latter of the Latentian 1984.
- Topomortum oficeration quadrificariation Process: Wince Cas Ty From a G.
- Timerana siler. Hadris & Navius & J. Sed. Sec. 1971p.11
- "Treorpetic neutroma Hieles Winde Can Type Form 2 43.
- "Transplie borner, France: Winds, Car. Type France, 43.
- 1 outormus vilucatus panir focusta, Perru : see Lepaizian.

#### T BLAFFORDEA

- "Information vinciniant Confession, McCongr., winderland, var. acutus, McCo Worder, Care, Type Fran. 2. Cd.
- Sermative mas campamalatus. McEry: Words, Cat. Type Fo.
   14. S. weedsmarks, 2000, 100f.: ETHERIESE. Pal. Mem. NS.
   2. 122.
- Montiferrary and antiferrary and ETHERMOSE, Pal. Mem. NSW, p. 123.
- "Petermita sampandatus, McCox: see Grandovinus.
- \*Trendersonn? corporate, 2001. 2011.; ETHERIDGE, Pal. Mem. NS. 7. 3. 123.

# XV. VERMES.

BY

# FLORENCE BUCHANAN, B.Sc.

#### ARRANGEMENT OF RECORD.

A common List of Titles is given, pp. 1-30, and three papers quoted which refer to *Vermes* in general. The Record is then broken up into the following groups, each group being again subdivided into a Biological, a Faunistic, and a Systematic part:—

- I.—Hemichordata, including Balanoglossus, Cephalodiscus, Rhabdopleura, pp. 30, 31.
- II.—Annelida, including Gephyrea, Chætopoda, and Hirudinea, pp. 31-52.
- III.—PLATYHELMINTHES, including CESTODA, TREMATODA, NEMERTINEA, and TURBELLARIA, pp. 53-72.
- IV .- NEMATODA, ACANTHOCEPHALA, and CHÆTOGNATHA, pp. 73-82.
- V.—ROTIFERA, pp. 82-88.
- VI.—MESOZOA, p. 88.

### TITLES.\*

- Anderson, H. H., & Shephard, J. Notes on Victorian Rotifers. P. R. Soc. Vict. (n.s.) iv, pp. 69-80, pls. xii & xiii. Abstr. in J. R. Micr. Soc. 1893, p. 48.
- Andrews, E. A. (1) Notes on the Fauna of Jamaica. Johns Hopk. Univ. Circ. xi, pp. 72-75, Vermes, p. 75.
  - Very brief, not giving the names of species.
- —. (2) Bifurcated Annelids. Am. Nat. xxvi, pp. 725-733, pl. xxi.
- ——. (2) On the Eyes of Polychestous Annelids. J. Morph. vii, No. 2, pp. 169-222, pls. ix-xi.

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

- APATHY, S. (1) Kritische Bemerkungen über das Frenzel'sche Mes zoon Salinella. Biol. Centralbl. xii, pp. 108-123. Transl. in An N. H. ix, pp. 465-482.
- —. (2) Contractile und leitende Primitivfibrillen. MT. z. St. Neap. x, pp. 355-375, pls. xxv & xxvi, 1 woodcut. Studies in Leeches.
- Appellor, A. Om Bergensfjordenes faunistiske præg. Bergens Mt Aarsber. 1891 (2) pp. 4-14.
- Apstein, C. Callizona angelina (Kbg.), Apstein. Festschr. zum 70t Geburtstage Rudolf Leuckarts. Leipsig (Engelmann): 189 pp. 44-47, pl. v.
- ATKINSON, C. F. Note on a Nematode Leaf-Disease. Ins. Life, (1831), p. 31.
- BALDWIN SPENCER. [See SPENCER, W. B.]
- BEDDARD, F. E. (1) Note upon the Encystment of Ælosoma. An N. H. (6) ix, pp. 12-19, 2 woodcuts.
- —... (2) The Earthworms of the Vienna Museum. T. c. pp. 113-13 pl. vii.
  - Schmarda's specimens properly described and most of them renamed
- —... (3) On a new genus of Oligochæta, comprising five new speci provisionally belonging to the Family Ocnerodrilidæ. Op. cit. (6) pp. 74-97, pls. vi & vii.
- ---. (4) Preliminary notice of a new Branchiate Oligochæt. Natur xlv, pp. 109 & 110.
- —. (5) A new Branchiate Oligochæt (Branchiura sowerbyi). Q. Micr. Sci. xxxiii, pp. 325-341, pl. xix.
- —. (6) Researches into the Embryology of the Oligochets. No. On certain points in the development of Acanthodrilus multiporu T. c. pp. 497-540, pls. xxx & xxxi. Abstr. in J. R. Micr. Soc. 189 p. 611.
- —. (7) Two new Genera and some new Species of Earthworm Q. J. Micr. Sci. xxxiv, pp. 243-278.
- —. (8) On the Earthworms collected in Algeria and Tunisia l Dr. Anderson. P. Z. S. 1892, pp. 28-37.
- ——. (9) On some Species of the Genus Perichæta (s. str.). Τ. pp. 153-172, pls. ix & x.
- ——. (10) On some aquatic Oligochætous Worms. T. c. pp. 349-36 2 woodcuts.
- ---. (11) On some Perichætidæ from Japan. Zool. Jahrb. Abth. Syst. vi, pp. 755-766, pl. xxxii.

- Bell, F. J. Remarks upon the habitat of *Bipalium kewense*. P. Z. S. 1892, p. 258.
- BENHAM, W. B. (1) Descriptions of three new Species of Earthworms. T. c. pp. 136-152, pls. vii & viii.
- —. (2) Notes on two Acanthodriloid Earthworms from New Zealand.
  Q. J. Micr. Sci. xxxiii, pp. 289-312, pls. xv & xvi.
- —. (3) A new English Genus of Aquatic Oliyocheta (Sparganophilus) belonging to the family Rhinodrilide. Op. cit. xxxiv, pp. 155-179, pls. xix & xx. Abstr. in J. R. Micr. Soc. 1893, p. 39.
- —. (4) An Earthworm from Ecuador (Rhinodrilus ecuadoriensis, n. sp.). Ann. N. H. (6) ix, pp. 237-246, pl. x.
- —. (5) Note on the occurrence of a freshwater Nemertine in England. Nature, xlvi, p. 611.
  - A species of Tetrastemma.
- ——. (6) British Earthworms. Op. cit. xlvii, p. 102. Supporting Hurst, criticising Friend (4).
- BERANECK, —. L'embryogenie de l'œil des Alciopides. Arch. Sci. Nat. xxviii, pp. 554 & 555.
- \*Beranger-Ferand, —. (1) Sur l'augmentation de fréquence du Tænia en France depuis un demi-siècle. Bull. de l'Acad. de méd. (3) xxvii, p. 112, and in Sém. Médic, 1892, No. 5.
- 2----. (2) Distribution geographique des Tænias de l'homme. Bull. de l'Acad. de méd. pp. 282-304.
- (3) De la ladrerie chez l'homme. Ann. Hyg. publ. 1892, pp. 481-517.
- Bergendal, D. (1) Några anmärkninger om Sveriges Triklader. Œfv. Ak. Förh. 1892, pp. 539-557.
- ——. (2) Kurzer Bericht über eine in Sommer 1890 unternommene zoologische Reise nach Nord Grönland. Bih. Sv. Ak. Handl. xvii, Afd. iv, No. 1. Given more fully as,
- —. (3) Beiträge zur Fauna Grönlands. Act. Lund. 1891-2 (n.s.), iii (iv), 180 pp. 1. Zur Rotatorien Fauna Grönlands, pp. 11-180.
- •——. (4) Gasteroschiza tricantha, Eine neue Gattung und Familie der Räderthiere. Bih. Sv. Ak. Förh. xvii. Also given under the title of "Ehrenberg's Euchlanis Lynceus wieder gefunden," in Act. Lund. 1891-2 (n.s.), iii (v), 2 pp.
- —. (5) Om Uteriporus rulgaris och betydelsen af Uterus hos Trikladerna. Förh. Sk. Naturf. xiv, p. 494, and °Sv. Ak. Handl. 1892, 1st part.

- [Bergendal, D.] (8) Einiges über den Uterus der Tricladen. Festschr. z. 70ten Geburtstage R. Leuckarts. Leipsig: 1892, pp. 310-318, pl. xxxii.
- Bergh, R. S. Kurze Bemerkungen über Herrn Apáthy und die Hirudineen Embryologie. Zool. Anz. xv, pp. 57 & 58.
- BILFINGER, L. Ein Beitrag zur Rotatorienfauna Württembergs. JH. Ver. Württ. xlviii, pp. 107-124.
- BILLET, A. [See GIARD & BILLET.]
- BLANCHARD, R. (1) Description de la Glossiphonia tesselata, O. F. M. Mém. Soc. Zool. v, pp. 56-58, 2 woodcuts.
- ——. (2) Présence de la Glossiphonia tesselata au Chili; description complémentaire de cette Hirudinée. Act. Soc. Chili, ii, pp. 177-187, 2 woodcuts.
- —. (3) Courtes notices sur les Hirudinées. Bull. Soc. Z. Fr. xvii.

  11. Sur la Typhlobdella kovátsi, Diesing, pp. 35-39, 4 woodcuts.

  From examination of type specimen, conclude = Haemopis sanguisuga. 111. Description de la Nephelis atomaria, Carena, pp. 165-172, 5 woodcuts. 1v. Description de la Glossiphonia marginata, O.F.M., pp. 173-178, 2 woodcuts. v. Description de la Glossiphonia sexoculata, Bergm., pp. 178-182, 2 woodcuts. vi. Sur le Branchellio punctatum, Baird, pp. 222 & 223.
- ——. (4) Description de la Xerobdella lecomtei, v. Frauenf. Mém. Soc. Zool. v, pp. 539-553, 9 woodcuts.
- ——. (5) Notices sur les parasites de l'homme, 1ère sér. ; De l'existence et de la prédominance ancienne du *Tænia saginata* dans l'Europe occidentale. Mém. Soc. Biol. (9) iv, pp. 243-258.
- —. (6) Sur la présence de la Trocheta subviridis en Ligurie, et description de cette Hirudinée. Atti Soc. Ligust. iii, pp. 407-435, 8 woodcuts.
- Bles, E. J. Notes on the Plankton observed at Plymouth during June, July, August, and September, 1892. J. Mar. Biol. Ass. (n.s.) ii, pp. 340-343.
- BLOCHMANN, F. (1) Ueber Sommer's sogenannte "plasmatische Längsgefässe" bei *Tænia saginata*, Göze, und *Tænia solium*, L. CB. Bakt. Parasit, xii, pp. 373-379, 3 woodcuts.
- —... (2) Ueber die Enwickelung von Cercarseum aus Helix hortensis zum geschlechtsreifen Distomum. T. c. pp. 649-652.
  - The Distomum is D. caudatum (alimentary canal of Erinaceus).
- ---. (3) Ueber die Anatomie und die verwandtschaftlichen Beziehungen der Brachiopoden. Arch. Ver. Mecklenb. xlvi, 1892, pp. 37-50.
  - Relationship to Phoronis emphasized.

- Bolsius, R. P. H. (1) Sur les organes ciliés des Hirudinées. 'Ann. Soc. Brux. xv, C.R. p. 44.
- ——. (2) Anatomie des organes segmentaires des Hirudinées d'eau douce, d'après les recherches cytologiques. Op. cit. xvi, Mém. pp. 1-70, pls. i & ii.
- Bos, J. RITZEMA. (1) Thierische Schädlinge und Nützlinge fur Ackerbau, Viehzucht, Wald- und Garten-bau; Lebensformen, Vorkommen, Einfluss und die Massregeln zu Vertilgung und Schutz. Berlin (Parcy): 1891, xvi + 876 pp., 477 figg. Reviewed J. R. Agric. Soc. iii, pp. 415-420.
- 200 pages devoted to Thread- and Eel-worms, Tape-worms, and Flukes; good index, and well arranged.
- ——. (2) Aphelenchus olesistus, n. sp., een Nematode, schadelijk aan Begonia's en Varens. Maandbl. Natuurw, 17th year, pp. 101-112, pl. vii.
- ——. (3) L'anguillule de la Tige (Tylenchus devastrix, Kühn) et' les maladies des plantes dues à ce Nematode. Arch. Mus. Teyl. (2) iii, pp. 545-588.
- BOUTAN, L. Voyage dans la Mer Rouge. Rev. Biol. iv.

  Three Polychæts and 1 Nemertine mentioned, but not identified, on p. 222.
- BOVERI, —. Ueber die Entstehung des Gegensatzes zwischen den Geschlechtszellen und den somatischen Zellen bei Ascaris megalocsphala nebst Bemerkungen zur Entwickelungsgeschichte der Nematoden. SB. Ges. Morph. viii, pp. 114-125, 5 figg.
- Brandes, G. (1) Zum feineren Bau der Trematoden. Z. wiss. Zool. liii, pp. 558-577, pl. xxii; also in Ber. Ges. Halle, 1891 (2), pp. 59-84. Abstr. in J. R. Micr. Soc. 1892, p. 372.
  - Criticised by BRAUN (3).
- —. (2) Revision der Monostomiden. CB. Bakt. Parasit. xii, pp. 504-511.
- Braun, M. (1) Ueber Distomum folium, Olf. Op. cit. xi, pp. 461-463,
- —. (2) Ueber Eurycælum sluiteri, Br. T. c. pp. 727-729,
- —. (3) Ueber einige wenig bekannte resp. neue Trematoden, Verh, Deutsch, zool. Ges. 2te Jahresversamml. 1892, pp. 44-52,
- ——. (4) Bronn's Klassen und Ordnungen des Thier-reichs. Bd. IV. Würmer. Continuation, parts 18-27, pp. 562-816, pls. xviii-xxxi, containing the Trematodes, Digenea.
- ——. (5) Verzeichniss von Eingeweidewürmern aus Mecklenburg. Arch. Ver. Mecklenb. xlv, 1891 (2), pp. 97-117.
- BRUNOTTE, C. Deux stations nouvelles du Phreoryctes menkeanus, Hfmst. Rev. Biol. v, pp. 120-122.

- BRYCE, D. (1) On the Macrotrachelous Calidinida. J. Quek. Club (2) v, pp. 15-23, 1 pl.
  - 4 new species.
- ---. (2) On some Moss-dwelling Cuthypnadæ, with descriptions of five new species. Sci. Goss. 1892, pp. 271-275, 11 figg.
- Buchanan, F. Report on Occupation of the British Association Table in the M. B. A. Laboratory at Plymouth. Rep. Brit. Ass. Edinburgh, 1892, pp. 356-360.
- BÜRGER, O. (1) Die Enden des excretorischen Apparates bei den Nemertinen. Z. wiss. Zool. liii, pp. 322-333, pl. xvi.
- —. (2) Zur Kenntnis der Nemertinen des Golfes von Neapel. Vorläufige Mittheilung. Nachr. Ges. Götting. 1892, pp. 137-178.
- Bütschli, O. Ueber den feineren Bau der contractilen Substanz der Muskelzellen von Ascaris nebst Bemerkungen über die Muskelzellen einiger anderer Würmer. Festschr. z. 70ten Geburtstage R. Leuckarts. Leipsig: 1892, pp. 328-336, pl. xxxiv.
- CADIOT, -. [See RAILLIET & CADIOT.]
- Carler, —. Note sur les Œufs et l'Embryon du Bilharzia hæmatoba. C.R. Soc. Biol. pp. 570-576, 4 figg.
- CALANDRUCCIO, S. Descrizione degli embrioni e delle larve della Filaria recondita, Grassi. Atti Acc. Gioen. (4) v, pp. 1-15, 17 woodcuts.
- Calman, W. T. On certain new and rare Rotifers from Forfarshire. Ann. Scot. N. H. 1892, pp. 240-245, pl. viii.
- Camerano, L. (1) Ricerche intorno al parassitismo ed allo svilluppo del Gordius pustulosus, Baird. Atti Acc. Tor. xxvii, pp. 598-607, 1 pl., and Boll. Mus. Zool. Tor. vii, No. 124, 5 pp.
- ——. (2) Intorno ad una species di Gordius (G. æneus, Villot) raccolta dal Sig. G. B. Anselmo in Venezuela, e intorno alle specie di questo genere fin ad ora discritte dell' America meridionale. Ann. Mus. Genov. (2) x, pp. 123-127, 1 fig.
- —... (3) Viaggio di L. Fea in Birmania e regioni vicine, xxvii. Gordii. T. c. pp. 128-131, 2 figg.
- ——. (4) Descrizione di una nuova specie del genere Gordius di Palmeira (Paranà) raccolta dal Dott. G. Franco Grillo. T. c. pp. 965 & 966.
- —. (5) Descrizione di una nuova specie del genere Gordius raccolta nell' isola Engano dall Dott. E. Modigliani. Op. cit. xii, pp. 539-541.
- Camus, F. Note sur la présence de Geophilus submarinus, Gr., et de quelques autres animaux marins sur la côte de Préfaille près Pornie (Loire Insé.). Bull. Soc. Sci. Nat. Ouest France, ii, 1892.
  - Syllis amica and Thulussema neptuni mentioned, pp. 33 & 34.

- CAPARINI, U. Nuove osservazione per servare all' istoria di alcuni parassiti. Com. 2ª. Giorn. Fisiol. Patol. Pisa. Anno 23 (1891), pp. 271-279.
  - 1. Tricocefali nel fegato di una bovina.
  - 2. Cesticerchi embrionali attraversante il fegato di un majale.
- CARSTEN, B. Trichinosis in the Netherlands. Trans. of 7th Internat. Congr. of Hyg. and Demogr., London, 1891 (2) sect. iii, pp. 56 & 57.
- CATHE, J. T. De breede lintworm (Bothriocephalus latus) in Nederland inheemsch. Geneesk. cour. 1891, No. 45.
- CERFONTAINE, P. Contribution à l'étude du système nerveux central du Lombrie terrestré. Bull. Ac. Belg. (3) xxii, pp. 742-752, 2 pls.
- CHARLES, R. H. A contribution on the life-history of the male *Filaria* medinensis, founded on the examination of specimens removed from the abdominal cavity of Man. Sci. Mem. vii, pp. 51-56, 1 pl.
- CHATIN, J. (1) Contribution à l'étude des Elements epidermiques chez les Nematodes. C.R. Soc. Biol. iv, pp. 135 & 136.
- O——. (2) L'Anguillule de la Betterave (Heterodera schachtii). Bull. No. 5, 1891, of Rep. to Fr. Min. of Agric., 50 pp., 9 pls.
- CHICHKOFF, G. D. Recherches sur les Dendrocœles d'eau douce (Triclades). Arch. Biol. xii, pp. 435-568, pls. xv-xx.
- Cholodkowsky, N. Contributions à la Theorie du mésoderme et de la métamerie. Congr. Internat. Zool., 2nd Session, Moscow, 1892, pp. 58-65.

Divides the *Metazoa*, according to method of formation of or absence of coelom, into *Enterocala* (including amongst "Vermes," the Chætognatha and Enteropneusta), *Genitocæla* (including the Annelides), *Acæla* (including the Platyhelminthes and the Mesozoa).

Chun, C. Die Dissogonie. Festschr. z. 70ten Geburtstage R. Leuckarts. Leipsig: 1892.

In a footnote, pp. 100 & 101, it is suggested that this process (reproduction at discontinuous periods in different stages of its development) may occur in Nereids.

- COATS, J. A specimen of the prismatic variety of Tania saginata. Glasg. Med. Journ. 1891, pp. 103-107.
- COLLIN, A. (1) Ueber die Regenwürmer der Umgegend von Berlin. SB. nat. Fr. 1892, pp. 115 & 116.
- ---. (2) Kleine Mittheilungen über Würmer (Bipalium aud Clepsine).
  T. c. p. 164.
- ——. (3) Gephyreen gesammelt von Herrn Oberstabsarzt Dr. Sander auf der Reise S.M.S. "Prinz Adalbert." Arch. f. Nat. lviii, pp. 177-182, pl. xi.

- Collin, G. Sur la fréquence relative des diverses espèces de Tania. Bull. Ac. Méd. xxvii, p. 176.
- COLLORADI, G. La Bilharzia hæmatobia dell' uomo ed i fenomeni morbosi cajionate da essa. Giorn. internaz. Sci. Med. Napoli, xiii (1891), pp. 851-864.
- CONDORELLI-FRANCAVIGLIA, M. (1) Sopra una rara anomalia della Tænia solium. Boll. Soc. Rom. Stud. Zool. i, pp. 31-35, 1 woodcut.
- ——. (2) Sopra una nuova specie di Filaria (F. apapillocephala). T. c. pp. 168-179.
- ——. (3) Sull' identità specifica dell' Echinorhynchus globocaudatus, Zeder, e dell' Echinorhynchus tuba, Budolphi. T. c. pp. 224-232.
- —... (4) A proposito di un grave caso di epistassi prodotta da puntura dell' Hirudo sanguisuga, Bergmann. T. c. pp. 233-241.
- —. (5) Contributo allo studio della *Tienia litterata*. Lo Spallanzani, (2) xx (1891), pp. 384-393, 1 pl.
- CONSALVI, G. (1) Sopra una malattia nuova per le regioni meridionali d'Italia denominata Anchilostomiasi. 2nd Ed. Lanciano, tip. Carabba: 1891, 16mo, 10 pp.
- —. (2) L'anchilostomiasi nelle province meridionali del continente italiano, Risposta alla littera del Dr. F. Follo. T. c. 6 pp. Estr., Riv. Ital. Terap. Igiene, July (1891).
- CORI, C. J. Ueber Anomalien der Segmentierung bei Anneliden und deren Bedeutung fur die Theorie der Metamerie. Z. wiss. Zool. liv, pp. 569-578, pl. xxv.
- Cosmovici, L. C. (1) Quelques remarques sur les Rotifères. Le Nat. xiii, 1891, pp. 65 & 92.

Distribution according to nature of water.

- —. (2) Rotifères Organisation et Faune de la Roumanie. Op. cit. xiv, 1892, pp. 44, 45, 58, 59, 70, 71, & 83.
- —. (3) Ce qu'il faut entendre par "système aquifère, organes segmentaires, organes excréteurs, nephridies." Congr. Internat. Zool. ii, Moscow, pp. 16-40.

Review of the different terms used in different groups of animals, chiefly Vermes. [See Kennel, J. v. (2).]

- CRETY, C. (1) Intorno la struttura delle ventose e di alcune organi tattili nei Distomi. Atti [Rend.] Acc. Rom. (5) i, pp. 21-26, 2 woodcuts. Translated into French in Arch. Ital. Biol. xviii, pp. 380-384.
- —. (2) Intorno al nucleo vitellino dei Trematodi. Atti [Rend.] Acc. Rom. (5) i, pp. 92-97.
- Cuenor, L. Commensaux et Parasites des Echinodermes. Rev. Biol. v, Platyhelminthes, Rotifers, and Polychæts, pp. 8-16, pl. i.
- CUNNINGHAM, J. T. The Preservation of Marine Animals for Zoological purposes. Ess. Nat. vi [Vermes], pp. 124 & 125.

- CURTICE, C. Parasites of Domestic Animals. J. Comp. Med. Vet. Arch. 1892, pp. 223-226.
- Daday, E. v. (1) Die geographische Verbreitung der im Meere lebenden Rotatorien. Math. Nat. Ber. Ung. ix (1891), pp. 55-66. Abstract and criticism in J. B. Micr. Soc. 1892, p. 488.
- ——. (2) Revision der Asplanchna Arten und deren ungarländischen Repräsentanten. Math. Nat. Ber. Ung. ix (1891), pp. 69-89, pls. ii & iii. Abstract and criticism in J. R. Micr. Soc. 1892, p. 794.
- DADAY, JENÖ. A mesőségi tavak mikroskopos faunaja (Microscopic Fauna of the Fields). Math. term. Értes, x, pp. 132-136.
- DALLA-TORRE, K. W. v. Zoologische Mittheilungen: Stephanoceros eichhornii, Ehrb., in Lanser Moor. Ber. Naturw. med. Ver. Innsbrück. 1891, p. viii.
- DEMATEIS, —. Das Austreten der Ascariden bei Fieberbewegungen. CB. Bakt. Parasit. xi, pp. 653-660.
- DENDY, A. (1) On the Australian Land Nemertine (Geonemertes australiensis, n. sp.). P. B. Soc. Vict. iv, pp. 85-122, pls. vii-x.
- —... (2) Description of some Land Planarians from Queensland. T. c. pp. 123-129, pl. xi.
- Carried Some Land Planarians from Tasmania. Rep. Australasian Ass. Advancement of Sci. 1892, Section D 6. Abstr. in J. R. Micr. Soc. 1893, p. 45.
- Despeignes -. [See Lortet & Despeignes.]
- Deurser, —. Zur Entwickelungsgeschichte der Filaria papillosa. Vorläufige Mittheilung. Zool. Auz. xv, pp. 129-131. Abstr. in CB. Bakt. Parasit. xi, p. 771, and in J. R. Micr. Soc. 1892, p. 371.
- °DEWITZ, J. Die Eingeweidewürmer der Haussäugethiere. Leipsig (Parcy): 1892, 8vo, 180 pp., 141 figg.
- DIAMARE, V. Di un nuovo cestode del Gen. Dipylidium, Lt. Boll. Soc. Nat. Napoli (1) vi, 1892, pp. 46-48.
- EHLERS, E. (1) Die Gehörorgane der Arenicolen. Z. wiss. Zool. liii (Suppl.), pp. 217-285, pls. xi-xiv. Abstr. in J. R. Micr. Soc. 1892, p. 480.
- —. (2) Zur Kenntnis von Arenicola marina, L. Nachr. Ges. Götting. 1892, pp. 413-418.
  - Deals with the young stages.
- EPSTEIN, A. Ueber die Uebertragung des menschlichen Spulwurms (Ascaris lumbricoides). Eine Klinisch experimentelle Untersuchung. JB. Kinderheilk. und Phys. Erziehung, xxxiii, 1892, p. 287.
- FAUVEL, A. La faune du Chan-toung. Rev. Quest. Sci. 1892, p. 492.
- FILIPPI, C. DE. Note preliminare sul sistema riproduttore della Tænia botrioceplitis. Boll. Soc. Rom. Stud. Zool. i (1892), pp. 75-79, pl. i.

- Fischer, W. Uebersicht der von Herrn Dr. F. Stuhlmann auf Sansibar und an der gegenüberliegenden Festlandes gesammelten Gephyreen. JB. Hamb. ix, pp. 79-89, 1 pl.
- FLOERICKE, C. Vorläufige Mittheilung über einige anscheinend neue Naidomorphen. Zool. Anz. xv, pp. 468-470.
- FOREL, F. A. Ascaris inflexa trouvé dans un œuf de poule. Bull. Soc. Vaud. xxviii, p. xxiv.
  Only this statement.
- FOWLER, G. H. The Morphology of *Rhabdopleura normani*, Allm. Festschr. z. 70ten, Geburtstage R. Leuckarts, pp. 293-297, pl. xxx; and Abstract in P. R. Soc. lii, pp. 132-134, 3 figg.
- Fowler, —. A curious Parasite. Ent. xxv (1892), p. 247. Found in *Epinephele tithonus*; shown on p. 291 to be a *Gordius*.
- Francaviglia. [See Condorelli-Francaviglia.]
- °FRANCIS, M. Liver Flukes. Bull. Texas Agric. St. xviii (1891), 9 pp., 6 figg.
- ---. [See also STILES (7).]
- François, P. [See Joubin & François.]
- FREAM, W. Trap-plants for Eelworms. J. R. Agric. Soc. (3) iii, pt. 1, pp. 173-176.
- FRENZEL, J. The Mesozoon Salinella. Ann. N. H. ix, pp. 79-84 & 109-111. Translations from German papers. [See Zool. Rec., 1891, Vermes, p. 10, FRENZEL, 1 & 2.]
- FRIEND, H. (1) British Annelids. With especial reference to the Earthworms of Essex. Ess. Nat. v & vi (1891-92).
  - 1. Introduction, Habits, Preserving, &c. Vol. v, pp. 193-196.
  - 2. Division into 3 genera. Terms for description. T.c. pp. 237-241.
  - 3. Allurus tetraëdrus, Eisen. Vol. vi, pp. 30-33.
  - 4. Genus Lumbricus, with 4 species. T. c. pp. 60-65.
  - 5. Genus Allolobophora, group Lumbricoidea. T. c. pp. 107-111.
  - 6. Genus Allolobophora, group Mucida. T. c. pp. 169-174.
  - 7. Genus Allolobophora, group Dendrobæna. T. c. pp. 185-190, 4 figg.
- ——. (2) Studies of British Tree- and Earthworms. J. L. S. Zool. xxiv, pp. 292-315.
- ----. (3) A rare British Earthworm. Nature, xlv, p. 365.

  L. eiseni, Lev.
- ---. (4) New British Earthworms. Op. cit. xlvi, p. 621, 3 figg. Criticised by Hurst & Benham (6).
- ---. (5) The Earthworms of Kent. Field Club, iii, pp. 21-27.
- —. (6) The Earthworms of Northhants. T. c. pp. 83-85, 100-103.
- ---. (7) The Earthworms of Middlesex. Sci. Goss. pp. 193-196.



- [FRIEND, H.] (8) On Hybridity among Worms. Field Club, iii, pp. 147-151, and Naturalist, pp. 301-304.
- —. (9) A new British Worm. Field Club, iii, pp. 167–169, and Sci. Goss. 1892, pp. 242 & 243.

Allurus tetragonurus.

- —. (10) Worm-hunting in Sussex. Sci. Goss. pp. 122-124.
- —. (11) New Yorkshire Earthworms; with an amended list of North British species. Naturalist, 1892, pp. 87-90.
- —. (12) A new Yorkshire Earthworm. T. c. pp. 247-249.
- —. (13) On a new species of Earthworm. P. R. Irish Ac. ii, pp. 402-410, 9 figg.
- He calls it Allolobophora hibernica, but in P.S. refers it to Rosa's A. cantabrica, found at Genoa.
- —. (14) Curiosities of Worm Life. Sci. Goss. 1892, pp. 108 & 109, 7 figg., and p. 161, 2 figg.
- •FRIIS, S. Om forekomsten af trichiner i Danmark. Tidskr. f. Sundhpfl. ii, p. 152.
- FRITSCH, A., & VAVRA, V. Vorläufige Bericht über die Fauna des Unter Pocernitzer und Gatterschlager Teiches. Zool. Anz. xv, p. 27.
- Fusser, —. Permanencia en la Estación de Santander. An. Soc. Esp. xxi [Vermes], Actas, pp. 172 & 175.
  - 2 Platyhelminthes, 6 Polychæts, 1 Phascolosoma
- GARSTANG, W. (1) On some new or rare marine animals recently discovered on the Coast of Devonshire. Rep. Tr. Devon. Ass. xxiv, "Vermes," pp. 381-383.
- —. (2) Marine Invertebrata of Plymouth. J. Mar. Biol. Ass. ii, No. 4, Annelids, p. 335.
- Generalch, A. Majtályog esete. Feregpetek az epiköben. (A case of Liver abscess. Eggs of Ascaris in gall stone). Evkön. Erd. Muz. xvii, pp. 21-29. Translated into German, pp. 220-225.
- Generalcu, G. Adatok a májtályog koroktanához (Contribution to the actiology of Liver abscess). T. c. pp. 8-20. German, pp. 213-219.
- GIARD, A. Sur la distribution geographique du Photodrilus phosphoreus, Dugės, et la taxonomie des Lumbriciens. C.R. Soc. Biol. iii, 1891, p. 252.
- GIARD, A., & BILLET, A. Sur quelques Trématodes parasites des bœufs du Tonkin (Note preliminaire). Op. cit. iv, pp. 613-616.
- GILES, G. M. J. (1) Some observations on the Life History of Sclerostomum tetracanthum, Diesing, and on Schlerutomiasis in equine animals; in connection with a so-called outbreak of "Surra" at Shillong. Sci. Mem. vii, pp. 1-25, 3 pls.
  - 1892. [vol. xxix.]

- [GILES, G. M. J.] (2) On a new Solerostome from the large intestine of Mules. (P.S. to (1).) Sci. Mem. vii, pp. 25-31, 1 pl.
- —. (3) On Nodular Discase of the intestine of Sheep. T. c. pp. 31-45, 1 pl.
- ——. (4) A description of two new Nematodes found in Sheep. T.c. pp. 45-51, 1 pl.
- GOODRICH, E. S. Note on a new Oligochaet (Vermiculus pilosus). Zool. Anz. xv, pp. 474-476, 2 figg.
- \*GRAEFE, A. Extraktion zweier Cysticercen aus einem Auge. Monatsbl. f. Augenheilk., xxx, p. 297.
- GRAFF, L. v. (1) Ueber pelagische Polycladen. Z. wiss. Zool. lv, pp. 189-219, pls. vii-x. Abstr. (without figures) in Verh. deutsch. Zool. Ges. 2ter Jhrg, 1892, pp. 117-121; also Abstr. in J. R. Micr. Soc. 1893, pp. 190 & 191.
- (2) Othelosoma symondeii, Gray. Zool. Anz. xv, pp. 7-9.
- ---. (4) Sur une planaire de la mer des Sargasses (Stylochoplana sargassicola, Mertens). Bull. Soc. Z. Fr. xvii, pp. 146 & 147.
- GRASSI, G., & ROVELLI, G. Ricerche embriologiche sui Cestodi.

  Atti Acc. Gioen. (4) iv, (lxviii), Mem. ii, 108 pp., 4 pls.
- GREENWOOD, M. On retractile cilia in the intestine of Lumbricus terrestris. J. Physiol. xiii, pp. 239-259, pl. ix.
- °GRIFFITHS, A. B. (1) The physiology of the *Invertebrata*. London: Reeve & Co., 8vo, xvi + 477 pp.
  - Circulation of Lumbricus dealt with.
- —... (2) On the blood of the *Invertebrata*. P. R. Soc. Edinb. xix, Annelids, pp. 119-123.
- ——. (3) L'hermérythrine, pigment respiratoire contenu dans le sang de certains Vers. C.R. cxv, pp. 669 & 670.
- GUERNE, J. DE. (1) Sur la dissémination des Hirudinées par les Palmipèdes. C.R. Soc. Biol. iv, pp. 92-95; also in Ann. N. H. x, pp. 117-120.
- —. (2) L'histoire des Némertiens d'eau douce. Leur distribution geographique et leur origine. C.R. Soc. Biol. iv, pp. 360-364; also in Ann. N. H. x, pp. 197-200.

  Answer to DU PLESSIS (1).
- Guerne, J. de, & Richard, J. Sur la faune de l'eau douce l'Islande. C.R. cxiv, pp. 310-313; Bull. Soc. Z. Fr. xvii, pp. 78-89; Ann. N. H. x, No. 58.
  - 2 Rotifers mentioned.

- HALLEZ, P. (1) Une loi embryogénique des Rhabdocælides et des Triclades. C.R. exiv, pp. 1033-1035. Abstr. in J. R. Mier. Soc. 1892, p. 484.
- (2) Sur l'origine vraisemblablement tératologique de deux espèces de Triclades. C.B. cxiv, pp. 1125-1128. Abstr. in J. R. Micr. Soc. 1892, p. 485.
- —. (3) Catalogue des Turbellariés (Rhabdocalides, Triclades, et Polyclades) du Nord de la France et de la côte Boulonnaise recoltés jusqu'à ce jour. Rev. Biol. iv & v [continuation from ii], 12 figg.

Supplement to list of Rhabdocals; classification of Triclades and their affinities with the Alloiocals; Laws of Morphogeny of the Turbellarians, iv, pp. 301-326.

Triclades: --Tribe Maricola, iv, pp. 338-350. Tribe Paludicola, pp. 425-456. Tribe Terricola, v, pp. 92-107.

- —. (4) Classification des Triclades (Note preliminaire). Bull. Soc. Z. Fr. xvii, pp. 106-109. Abstr. in J. R. Micr. Soc. 1893, p. 45.
- HAMANN, O. (1) Zur Entstehung des Exkretionsorganes, der Seitenlinien und der Leibeshöhle der Nematoden. CB. Bakt. Parasit. xi, p. 501-503. Abstr. in J. R. Micr. Soc. 1892, p. 482.
- ——. (2) Die Nemathelminthen. 1. Monographie der Acanthocephalen. Jen. Z. Nat. xxv (1891), 119 pp.
  See Kaiser (1).
- OHASSALL, A., & STILES, C. W. Strongylus rubidus, a new species of Nematode parasitic in Pigs. J. Comp. Med. Vet. Arch. Apr. 1892, 3 pp., 3 figg.
- HASWELL, W. A. (1) Observations on the *Chlorhæmidæ*, with special reference to certain Australian Forms. P. Linn. Soc. N.S.W. vi (1891), pp. 329-356, pls. xxvi-xxviii. Abstr. in J. R. Micr. Soc. 1892, p. 611.
- —... (2) On the Excretory System of Temnocephala. Zool. Anz. xv, pp. 149-151. Abstr. in J. R. Micr. Soc. 1892, p. 486.
- ——. (3) On the Systematic Position and Relationships of the Temnocephaleæ. Abh. Ges. Halle, xvii, pp. 455-460.
- ——. (4) Jottings from the Biological Laboratory of Sydney University. P. Linn. Soc. N.S.W. vii.
  - No. 17. Three Zoological Novelties, pp. 340-342:-
    - The occurrence of a second species of Phoronis in Port Jackson.
    - An Alloiocole Turbellarian inhabiting the underground waters of Canterbury, N.Z.
    - III. A new genus of Temnocephales (= Actinodactylus). A detailed description of this genus in (5) the Macleay Memorial Volume is promised.

- [HASWELL, W. A.] (5) A Monograph of the Temnocephaleα. Maclesy Memorial Vol. of P. Linn. Soc. N.S.W., pt. 1, 1892.
- HATSCHEK, B. System der Anneliden, ein vorläufige Bericht. Lotos, xiii, "1893" (1892), pp. 123-126.
- HENKING, H. Darstellung des Darmkanals von Hirudo. Festschr. z. 70ten Geburtstage R. Leuckarts. Leipsig: 1892, pp. 319-327, pl. xxxiii.
- HERRING, E. Zur Kenntniss der Aleiopiden von Messins. SB. Ak. Wien, pp. 713-768, 6 pls.
- HESSE, R. Ueber das Nervensystem von Ascaris megalocephala. Z. wiss. Zool. liv, pp. 548-568, pls. xxiii & xxiv. Abstr. in J. R. Micr. Soc. 1892, p. 792.
- \*HILLIER, E. A Guinea Worm in the Tongue. Ind. Med. Rec. 1892, p. 79.
- HILLIGER, J. C. Weitere Mittheilung über leuchtende Würmer und Larven. Helios, 9ter Jhrg. (1892), p. [84].
- Hoon, J. (1) Floscularia quadrilobata, n. sp. Int. J. Micr. (3) ii, pp. 26-28, pls. iii & iv.
- ----. (2) Floscularia gosseii: a new Rotifer. T. c. pp. 73-78, pls. vi & vii.
- HORNELL, J. (1) Report on the Polychætous Annelids of the L. M. B. C. District. Fauna of Liverpool Bay, Rep. iii (1892), pp. 126-171, pls. xiii-xv.
- \_\_\_\_. (2) A Strange Commensalism—Sponge and Annelid. Nature, xlvii, p. 78.
  - Leucodore coeca, Oerst., in Microciona plumosa.
- HORST, R. Earthworms from the Malay Archipelago. Zool. Ergebn. einer Reise in Niederländisch Ost-Indien (Ed., Max Weber), iii (?) (1892), pp. 28-77, pls. ii-iv. Abstr. in J. R. Micr. Soc. 1893, p. 38.
- OHUBER, J. C. (1) Bibliographie der klinischen Helminthologie. Heft 3 & 4. Die Darmcestoden der Menschen. München (Lehmann): 1892, 8vo, pp. 65-150.
- ——. (2) Ein Fall von Echinococcus multilocularis der Gallenblase. Deutsch. Arch. Klin. Med. xlviii, p. 432.
- HUBRECHT, A. A. W. The Nephridiopores in the Earthworm. Tijdschr. Nederl. Dierk. Ver. iii, pp. 226-234, pl. xii.
- OHUET, —. Une nouvelle Cercarie (C. pectinata) chez Donax anatinum. J. Anat. Phys. norm. path. xxvii, p. 162.
- HURST, C. H. On a "Supposed New Species of Earthworm," and on Nomenclature of Earthworms. Nature, xlvii, p. 31.

  Criticism of FRIEND (4).

- IMHOF, O. E. (1) Beitrag zur Kenntnis der Lebensverhältnisse der Rotatorien. Ueber marine, brackische, und euryhyaline Rotatorien. Biol. Centralbl. xii, pp. 560-566.
- —. (2) Ueber die Lebensverhältnisse zugefrorener Seen. MT. Aargau. nat. Ges. vi.
  - 3 Rotifers (gen. Anuraea) mentioned, p. 56.
- —. (3) Beiträge zur Fauna der Schweiz; Thierwelt der stehende Gewässer. Die Rotatorien. T. c. pp. 93-110.
- ---. (4) Die zusammensetzung der pelagischen Fauna der Susswasserbecken. Biol. Centralbl. xii, Rotifers, pp. 176-178 & 203.
- JÄGERSKIOLD, L. A. Zwei der Euchlanis lynceus, Ehrenberg, verwandte neue Rotatorien. Zool. Anz. xv, pp. 447-449, 2 figg.
- JAMMES, L. (1) Contributions à l'étude de la couche sous-cuticulaire des Nematodes et particulièrement du genre Ascaris. Ann. Sci. Nat. 59° année (7) Zool. xiii, pp. 321-342, pl. ix.
- Les premières phases du développement de certains vers nématodes. C.R. cxiv, pp. 1555-1557. Critical Abstr. in J. R. Micr. Soc. 1892, p. 613.
- OJANSON, J. L., & TOKISHIGE, H. Filaria immitis und andere bei Hunden in Japan vorkommende Parasiten. MT. Ges. Ostasien's, v, pp. 349-360, 2 pls.
- JOUBIN, L. Quelques indications sur la recherche et la préparation des Némertes. Act. Soc. Chili, ii, pp. 268-270.
- JOUBIN, L., & FRANÇOIS, P. Note sur quelques Némertes de Nomes. Rev. Biol. iv, pp. 161-172, pl. vi. 4 new species.
- JOURDAIN, É. (1) Étude sur les épitheliums sensitifs de quelques vers annelés. Ann. Sci. Nat. (7) xiii, pp. 227-258. Abstr. in J. R. Micr. Soc. 1892, p. 788.
- ——. (2) Les Corps jaunes des Echinodermes et des Géphyriens inermes. C.R. Ass. Fr. Sci. Marseilles, 1891 (2), pt. i, p. 240; pt. ii, pp. 537 & 538.
- ——. (3) De la valeur du mot endothelium en anatomie à propos des cellules à cils vibratiles de la cavité générale des Sipunculiens. Ann. Fac. Sci. Marseille, ii (1892), pp. 43-48, 2 figg. Also in C.R. Soc. Biol. iv, pp. 27-30, where criticised by Malassez, p. 31.
- JOURDAIN, M. S. Sur l'embryogénie des Sagitta. C.R. cxiv, pp. 28 & 29. Also in Ann. N. H. ix, pp. 415 & 416.
- KAFKA, J. Untersuchungen über die Fauna der Gewässer Böhmens. II. Die Fauna der Böhmischen Teiche. Arch. naturw. Landesforsch. Böhmen, viii (2) pp. 1-115.
  - Rotifers, p. 112; Turbellarians, Leeches, Cheetopods, p. 113.

Kaiser, J. (1) Die Acanthocephalen und ihre Entwickelung. Bibl. Zool. Heft vii, 1891-93.

An important monograph on the family. In an appendix some remarks are made on Hamann's Monograph [see Hamann (2)]. The two authors appear to differ in some important points.

- ——. (2) Die Nephridien der Acanthocephalen. CB. Bakt. Parasit. xi, pp. 44–49. Abstr. in J. R. Micr. Soc. 1892, p. 371.
- KALLSTENIUS, E. Eine neue Art der Oligochætengattung Amphichæle, Tauber. Biol. Fören. iv, pp. 42-55, 5 figg.
- KELLICOTT, -. Rotifer Notes. Am. Micr. J. xiii, p. 12.
- Kennel, v. J. (1) Die Insel Trinidad: Ueber die Ableitung der Vertebratenangen von den Augen der Anneliden [see Zool. Bec. 1891, Kennel (1)]. SB. Ges. Dorp. ix (3) 1891 (2).
- ——. (2) Réponses à Questions proposés par M. le Prof. L. Cosmovici.
  2. Fondement naturel d'un système du type de vers. 3. Uniformité de la terminologie des organes sécretoires (? excreteures). Congr. Internat. Zool., Sess. 2, Moscow, pp. 71-73.
- KESWAL, —. A Tubicolar Annelide. J. Bomb. N. H. Soc. vii (1892), pp. 114 & 115.

Short and imperfect description, asking for identification.

Kieffer, J. J. Ueber eine in Gallmücken parasitisch lebende Anguillula. Berl. Ent. Zeit. xxxvi, p. 26, 1 fig.

Living in large numbers in the larva of Asynapta citrina, a Cecidomyid fly.

- KINYOUN, J. J. Echinococcus hominis of the kidneys, liver, and bladder. Ann. Rep. Mar. Hosp. Serv. 1891, pp. 147-149.
- KLAPHAKE, J. Fütterungsversuche mit amerikanischen Trichinen. Z. f. Fleisch u. Milch Hyg. 1892, pp. 152 & 153.
- KNIPOVICHA, H. Otchet ob ekakursii na Solovetzki biologhicheskuyu stantziyu lyetom 1890 ghohda. (Report on a Biological Excursion to Solovetzki.) Trudui St. Petersburg Est. xxii, 1 (1891).

Nemertines, p. 25; Polychets, pp. 25-27; Priapulus, pp. 24 & 40; Balanoglossus, pp. 24, 35, & 40. (In Russian.)

- KOEHLER, —. Les conditions d'existence des organismes pélagiques. Rev. gén. Sci. 3° année, pp. 76-86, figg.
- KOJEVNIKOV, G. La Faune de la mer Baltique orientale et les problèmes des explorations prochaines de cette faune. Congr. Internat. Zool., Sess. 2, Moscow, 1892 (3).

Vermes, pp. 141-145.

- Kraemer, A. (1) Ueber den innern Bau der Taenien der Süsswasserfische (Vorläufige Mittheilung). Zool. Anz. xv, pp. 14-18. [See (2).]
  - —. (2) Beitrage zur Anatomie und Histologie der Cestoden der Süsswasserfische. Z. wiss. Zool. liii, pp. 647-722, pls. xxvii & xxviii. Abstr. in J. B. Micr. Soc. 1892, p. 373.

- Kulagin, N. Ueber einige in europäischen Russland und in Siberien vorkommende Arten von Regenwürmern. Mél. biol. xiii, 1 (1891), pp. 87-96.
- LABBÉ, A. Note sur un nouveau parasite du sang (Trypanomonas danelevskyi). Bull. Soc. Z. Fr. xvi (1891), pp. 229-231.
- Blood of Leeches used as cultivating medium for this and other *Protozoa*.
- LACAZE-DUTHIERS, H. DE. Sur la faune de Banyuls-sur-Mer (Laboratoire Arago). Arch. Z. expér. (2) x, N. & R., p. xiv.
- LANG, A. Ueber die Cercaria von Amphistomum subclavatum. Ber. Ges. Freiburg, vi, pp. 81-89.
- LANGERHANS, —. Ueber regressive Veränderung der Trichinen in ihrer Kapseln. Virchow's Arch. exxx, p. 205.
- LEBRUN, H. Les centrosomes dans l'œuf de l'Ascaris megalocephala.

  Anat. Anz. vii, pp. 627 & 628.
- LENHOSSÉK, M. v. (1) Ursprung, Verlauf und Endigung der sensibelen Nervenfasern bei *Lumbricus*. Arch. mikr. Anat. xxxix, pp. 102-136, pl. v.
- —. (2) Die intraepidermalen Blutgefässe in der Haut des Regenwurmes. Verb. Ges. Basel. x, pp. 84-91.
- LEUCKART, R. (1) Ueber den grossen americanischen Leberegel. CB. Bakt. Parasit. xi, pp. 797-799.
- —... (2) Ueber die Speicheldrüsen der Hirudineen. Ber. Sächs. Ges. vi (1892), pp. 556-558.
- LEVANDER, K. M. (1) Eine neue Pedalion-Art. Zool. Anz. xv, pp. 402-404.
- ---. (2) Mikrofaunistiska anteckningar. Med. Soc. Fenn. xvii (1890-92), pp. 129-143.
- Lewis, —. List of Rotifers found in ponds near Ealing. Rep. Ealing Micr. N. H. Soc. xv, p. xiv.
- OLIEBE, K. TH. Zur Naturgeschichte der Rohrdommel. Ornith. Monatschr. d. Deutsch. Ver. Schutz. Vogelwelt. xvii, No. 12 [p. 323 show Leeches and Earthworms swallowed whole, alive].
- LINDENFELD & PIETRUSZYNSKI. [See NUSBAUM (2).]
- Linstow, von. (1) Ueber Filaria bancrofti, Cobbold. CB. Bakt. Parasit. xii, pp. 88-92, 6 figg.
- \_\_\_\_. (2) Beobachtungen an Vogeltänien. T. c. pp. 501-504, 3 figg.
- —... (3) Beobachtungen an Helmintheularven. Arch. mikr. Anat. xxxix, pp. 325-342, pl. xv.
- ——. (4) Helminthen von Süd-Georgien. Nach der Ausbeute der Deutschen Expedition von 1882-83. JB. Hamb. Anst. ix, pp. 59-77, 3 pls.

- [Linstow, von.] (5) Ueber Mermis nigrescens, Duj. Arch. mikr. Anat. zl, pp. 498-511, pls. zxviii & xxix.
- LINTON, E. (1) Notes on Avian Entozoa. P. U. S. Nat. Mus. xv (1892), pp. 87-113, pls. iv-viii.
- Unfavourably reviewed and summarized by Stiles in CB. Bakt. Parasit. xiii (1893), p. 497, and in Am. Nat xxvi, pp. 1038-1040.
- ——. (3) Notes on a Nematode Parasite from the Chipping Sparrow (Spizilla socialis). T.c. pp. 705-707, 5 figg.
- ——. (4) Notes on the Entozoa of Marine Fishes, with descriptions of New Species. Pt. 111. Acanthocephala. Rep. U. S. Fish. Comm. pt. xvi, 1888 (1892), pp. 523-542, pls. liii-lx.
- —... (5) On the anatomy of Thysanocephalum crispum, Linton, a Parasite of the Tiger Shark. T. c. pp. 543-556, pls. lxi-lxvii.
- <sup>o</sup>Lönnberg, E. (1) Distomum goliath, v. Ben. Verh. Biol. Ver. Stockholm, iii (1891), 14 pp., 1 pl.
- ——. (2) Einige Experimente, Cestoden kunstlich lebend zu erhalten. CB. Bakt. Parasit. xi, pp. 89-92.
- ——. (3) Ueber das Vorkommen des breiten Bandwurmes in Schweden. T. c. pp. 189-192.
- LOOSS, A. (1) Ueber Amphistomum subclavatum, Rud., und seine Entwickelung. Festschr. z. 70ten Geburtstage R. Leuckarts. Leipsig (1892), pp. 147-169, pls. xix & xx.
- ——. (2) Schmarotzerthum in der Thierwelt. Zool. Vorträge hrsg. v. W. Marshall. Leipzig (R. Freese): 1892, 8vo, Heft 9 & 10, 180 pp.
- LORD, J. E. Notes on the genus Distyla, Class Rotifera. Sci. Goss. 1892, pp. 15 & 16.
- LORTET & DESPEIGNES. Vers de terre et Tuberculose. C.R. cxv, p. 66, and Sem. Médic. 1892, No. 5.
- Loz, J. Les Vers de terre agents de fertilisation du sol. Rev. Sci. Nat. appl. 38° ann. (1891), p. 45.
- LUCET, A. [See RAILLET, A., & LUCET, A.]
- LUTZ, A. Zur Lebensgeschichte des Distoma hepaticum. CB. Bakt. Parasit. xi, pp. 783-796, 1 fig.
- McIntosii, W. C. Notes from St. Andrews Marine Laboratory, No. xiii, Ann. N. H. x [Vermes], pp. 103-107, pl. viii, figg. 1-4.
- MAGALHÄES, P. S. DE. (1) Notes d'helminthologie Brésilienne. Bull. Soc. Z. Fr. xvii.
  - 1. Taenia cuneata, v. Linst., pp. 145 & 146.
  - 2. Heterakis brasiliensis, n. sp., of the domestic fowl, pp. 219-221, 1 fig.

- [MAGALHĀES, P. S. DE.] (2) Die Filuria bankrofti, Cobbold, und die Filuria immitis, Leidy. CB. Bakt. Parasit. xii, pp. 511-514, 4 figg.
- MAIER, B. L. Beiträge zur Kenntnis des Hirudineen Auges. Zool. Jahrb. Abth. f. Anat. v, pp. 552-580, pl. xxxvii.
- MAN, J. G. DE. Ueber eine neue in Gallen einer Meeresalge lebende Art der Gattung *Tylenchus*, Bast. Festschr. z. 70ten Geburtstage R. Leuckarts. Leipsig (Engelmann): 1892, pp. 121-126, pl. xvi.
- MANGOLD. —. Ueber den multiloculären Echinococcus und seine Tänie. Berl. klin. Wochenschr. 1892, Nos. 2 & 3. Abstr. in CB. Bakt. Parasit. xi, p. 738, and J. R. Micr. Soc. 1892, p. 619.
- Manson, P. The Geographical Distribution, Pathological Relations, and Life History of Filuria sanguinis hominis diurna, and of F. sang. hom. perstans in connection with preventive medicine. Trans. of the 7th Intern. Congr. Hyg. Demog. London, 1891 (2) sect. i, pp. 79-97, 3 figg.
- MARCIALIS, E. Saggio di un Catalogo metodico dei principali e più comuni animali invertebrati della Sardegna. Boll. Soc. Rom. Stud. Zool. i, pp. 246-282; Vermes, pp. 249-251.
- MARENZELLER, E. v. (1) Die Polychäten der Bremer Expedition nach Ostspitzbergen im Jahre 1889. Zool. Jahrb. Abth. Syst. vi, Heft 2, pp. 397-434, pl. xix.
- ——. (2) Sur une Polynoïde pélagique (Nectochæta grimaldii, n. g. & sp.) recueillie par l'Hirondelle en 1888. Bull. Soc. Z. Fr. xvii, pp. 183-185.
- MARK, E. L. Polychærus caudatus, n. g. & sp. of Turbellarians. (Acæla, fam. Aphanostomida). Festschr. z. Geburtstage R. Leuckarts. Leipsig: 1892, pp. 298-309, pl. xxxi.
  - MASSALONGO, C. Sul elmintocecidio (Tylenchus invalis) dell' Edelweiss. N. Giorn. Bot. Ital. Firenzi. xxiii, 2 (1891), p. 375.
  - MATZDORFF, C. (1) Ursprung des leuchtenden Regenwurmes (Photodrilus phosphoreus). Helios, ix (1891), p. [58].
  - —. (2) Ueber schmarotzende Protozoen. Op. cit. x (1892), pp. 11 & 12.
    - Rhabdostyla arenicolæ on the branchise of Arenicola.
  - —... (3) Commensalisten und Parasiten von Echinodermen. T.c. p. 91. Nematode, Rotifer, Platyhelminths, and a Polychæt.
  - METER, S. D. VAN. The Filaria immitis. Intern. med. Mag. 1892, i, pp. 1060-1064.
  - MICHAELSON, W. (1) Terricolen der Berliner Zoologischen Sammlung (ii). Arch. f. Nat. lviii (ii), pp. 209-261, pl. xiii, and 5 woodcuts.
  - —. (2) Die Lumbriciden Mecklenburgs. Arch. Ver. Mecklenb. xliv, 1890 (1), pp. 48-54.

- [Michaelson, W.] (3) Beachreibung der von Herrn Dr. Fr. Stuhlmann am Victoria Nyanza gesammelten Terricolen. JB. Hamb. Anst. ix, (2), 1891 (92), pp. 29-42, 1 pl.
- —. (4) Polychaeten von Ceylon. T. c. pp. 91-113, 1 pl.
- MONIEZ, R. (1) Notes sur les Helminthes (continued). Rev. Biol. iv. xI. La Gymnorhynchus reptans et sa migration, pp. 150 & 151.
  - XII. Sur un Tétrarhynque nouveau provenant des campagnes de l'Hirondelle (Dibothriorhynchus monticellii), p. 279.
- Monticelli, F. S. (1) Ricerche sulla spermatogenesi nei Trematodi. Month. Int. J. Anat. Hist. ix, pp. 112-118 & 121-149, pls. viii & ix.
- ——. (2) Sul nucleo vitellino delle uova dei Trematodi. Boll. Soc. Nat. Napoli (1) vi, pp. 5-8 (Estr.).
- —. (3) Studii sui Trematodi endoparassiti. Sul genere Notocotyle. T. c. pp. 26-46, pl. i.
- —. (4) Studii sui Trematodi endoparassiti. Dei Monostomum del Box Salpa. Atti Acc. Tor. xxvii, pp. 514-534.
- ---. (5) Studii sui Trematodi endoparassiti. Dei Monostomum cymbium, Diesing, Contribuzione allo studio dei Monostomidæ. Mem. Acc. Tor. 2º ser. xlii, pp. 683-727, 1 pl.
- —... (6) Cotylogaster michaelis, n. g. & sp. Festschr. z. 70ten Geburtstage R. Leuckarts. Leipsig (Engelmann): 1892, pp. 168-214, pls. xxi & xxii.
- —. (7) Nota intorno a due forme di Cestodi. Boll. Mus. Zool. Tor. vii, No. 127, 9 pp., 1 pl.
- —. (8) Sulla cosidetta subcuticola dei Cestodi. Rend. Acc. Nap. (2) vi, anno xxxi, pp. 158-166.
- ——. (9) Appunti sui Cestodaria. Atti Acc. Napoli (2) v, No. 6, pp. 67-78, 4 figg.
- —... (10) Sul genere Bothrimonus, Duvernoy, e proposte per una classificazione dei Cestodi. Monit. Zool. Ital. iii, pp. 100-108.
- MORGAN, T. H. (1) Balanoglossus and Tornaria of New England. Zool. Anz. xv, pp. 456 & 457.
- ——. (2) Growth and Metamorphosis of Tornaria. J. Morph. v, pp. 407-450, pls. xxiv-xxviii.
- ---.. (3) Spiral Modification of Metamerisation. Op. cit. vii, pp. 245-251, 3 figg.
- Moussu, -. [See Railliet & Moussu.]
- •Mudd, H. H. Echinococcus multilocularis of the brain. Am. J. Med. Sci. 1892, pp. 412–422.
- Müller, F. Die Begattung der Clepsinen. Zool. Jahrb. Abth. f. Syst. vi, p. 338.
  - Correction of a statement by WHITMAN.



- Nabias, B. de, & Sabrazés, J. (1) Sur les Embryons de filaire du sang de l'homme. Sem. Médic. 1892, p. 212; CB. Bakt. Parasit. xii, p. 171.
- ---.. (2) Sur les Embryons de la Filaire du sang chez l'homme. C.R. Soc. Biol. iv, pp. 455-461, 1 fig.
- NEUMANN, —. A Treatise on the Parasites and Parasitic Diseases of the Domesticated Animals. English Translation (and Annotation) by G. Flemming, C.B. Loudon (Balliere, Tindall, & Cox): 1892, xxiii + 800 pp., 365 figg.; Review J. Agric. Soc. (3) iii, pp. 397-403.
- NIEL, E. Quelques mots sur les Anguilles. Bull. Soc. Rouen (3) xxvii (1891), pp. 73-75.
- Nusbaum, J. (1) Poglady na stosunki genetyczne pomiędzy Thankowcami i Pierwotniakami z. powodu odkrycia Salinelli (Frenzel). Kosmos, Lemberg, zvii (1892), pp. 233-264, 11 figg.
- Discussion of relationship between *Meta* and *Protozoa*, as affected by the discovery of *Salinella*. Review of all the theories previously advanced. (In Polish.)
- —.. (2) Zur Kenutnis der Wurmfauna und Crustaceenfauna Polens. Biol. Centralbl. xii, pp. 54-57. I. Zur Fauna der Hirudineen; II. der Rhabdoccelen (11 species); III. der Lumbriciden (10 species).
- Abstract from two papers by H. LINDENFELD & J. PIETRUSZYNSKI (in Polish) in the Pamietnik Fizyograficzny, Warsaw: ix (1889), 12 pp., 1 pl. (Leeches); x (1890), 42 pp., 1 coloured pl. & 13 woodcuts. Further details promised by the authors in the same journal.
- OSTERTAG, —. Vermögen Darmtrichinen und wandernde Trichinen auf einen neuen Wirt überzugehen? Z. f. Fleisch- u. Milchhyg. 1892-93, pp. 45-50.
- OTT, H. N. A Study of Stenostoma leucops, O. Schm. J. Morph. vii, pp. 263-304, pls. xiv-xvii.
  - Preliminary account in Zool. Anz. xv, pp. 7-9.
- PARKER, T. J. Exhibition of Brunchellion (? n. sp.), occurring abundantly on <u>Raja</u> nasuta, and of a <u>Dujardinia</u> from Port Chalmers. Tr. N. Z. Inst. xxiv (Proceedings), p. 714.
  - PARONA, C. (1) Elmintologia italiana (Continuation). Boll. scient. xiii, pp. 124-128, xiv, pp. 92-96, 125-127. Bibliography.
  - ——. (2) Di una nuova specie di Echinorhynchus (E. novellæ) parassita di una Chirottero di Porto-Rico. Ann. Mus. Genov. (2) x, pp. 396– 398, 4 figg.
  - PARONA, C., & PERUGIA, A. (1) Contribuzione per una monografia del genere *Microcotyle*. T. c. Res Liguaticae, xiv, pp. 173-220, pls. iii-v.
  - ——. (2) Note sopra Trematodi ectoparassiti. Op. cit. (2) xii, Res Liguatics, xvii, pp. 86-102, pls. ii & iii.

- Parsons, F. A. Notes on Two Rotifers found in Epping Forest. J. Quek. Club, iv, pp. 378-380, pl. xxv, and Am. Micr. J. xiii, pp. 277 & 278.
- PERIASLAVZEFF, S. Dopolneniya k Faunye Chernaglio morya. Trudui Kharkoff Univ. xxv (1891), pp. 235-274, pls. vii & viii.
- A Supplement to the Fauna of the Black Sea. Deals chiefly with Polychests. (In Russian.)
- Perroncito, E. Maladies infectiouses des Animaux qui peuvent être transmises à l'Homme, et reciproquement. Trans. of the 7th Internat. Congr. Hyg. Demogr. London, 1891 (2), Sectn. iii, pp. 189-192.

Tienia mediocanellata and Ankylostoma duodenalis (larvæ).

PERUGIA, A. [See PARONA & PERUGIA.]

Pietruszynski, J. [See Nusbaum (2).]

Plessis, G. Du. (1) Sur les Némertiens du lac de Genève. Zool. Ans. xv, pp. 64-66.

Criticized by VAILLANT and by DE GUERNE (2).

- —... (2) Note sur Tetrastemma lacustris, espéce nouvelle de Némertien d'eau douce. Bull. Soc. Vaud. (3) xxviii, pp. 43-48, pl. iv.
- POUCHET, G. Sur la faune pélagique du Dyrefjord (Islande). C.R. cxiv, pp. 191-193.
- PRESTON, W. N. Mounting Trichina. Am. Micr. J. xiii, p. 197.
- PROUTIO, H. Sur deux Myzostomes parasites de l'Antedon phalangium (Müller). C.R. cxv, pp. 864-849.
- \*Puky, A. Twenty cases of *Echinococcus*. (In Hungarian.) Magyar orvosiarch. 1892, Nos. 3 & 4.
- RAILLIET, A. (1) Les parasites transmissibles des Animaux à l'Homme, envisagés specialement au point de vue de la Prophylaxie. Trans. of the 7th Internat. Congr. Hyg. Demogr., Sectn. iii, pp. 57-87.
- —. (2) Notices parasitologiques. Bull. Soc. Z. Fr. xvii, pp. 110-117:—
  - I. Cysticercus pisiformis à six ventouses, p. 110.
- 11. Téniadé de coloration ardoissée recueilli chez un Lapin de Garonne (Anoplocephala cuniculi, Bl.), p. 111.
- 111. Sur des Téniades échaucrés du Chien (Dipylidium caninum), p. 112.
  - 1v. Dipylidium caninum dans les glandes anales du Chien, p. 113.
- v. Cysticercus tenuicollis chez un chevreau de quatre à six semaines, p. 114.
  - VI. Cysticercus tenuicollis chez l'Orya beisa, p. 115.
- VII. Tania tenuirostris, Rud., chez l'Oie domestique; remarques sur la classification des Cestodes parasites des Oiseaux, p. 115.
  - VIII. Trichocéphale de l'intestin du Furet, p. 116.



- [RAILLIET, A.] (3) Observations sur l'embryon du Gynecophorus hæmatobius, Bilharz. T. c. pp. 161-164.
- ——. (4). Observations sur la Resistance vitale des embryons de quelques Nématodes. C.R. Soc. Biol. iv, pp. 703 & 704.
- —. (5) Sur un Tænia du Pigeon domestique representant une espèce nouvelle (T. delafondi). T. c. pp. 49-53.
- ---. (6). Un cas très ancien de Tænia (Hymenolepis) diminuta chez l'homme. T. c. pp. 894-896.
- (7) Sur la frequence de la Strongylose gastro-intestinale des leporides. Rec. Med. Vét. 1892, No. 8, p. 195-199.
- RAILLIET, A., & CADIOT, —. (1) Essaie de transmission du Strongylus rasorum du Chien au Chien; resultats negatives. C.R. Soc. Biol. iv, pp. 702 & 703.
- —. (2) Strongylose du cœur et du poumon chez un Chien. T. c. pp. 482-486.
- RAILLIET, A., & LUCET, A. (1) Sur le Davainea proglottina, Davaine. Bull. Soc. Z. Fr. xvii, pp. 105 & 106.
- —. (2) Observations et expériences sur quelques helminthes du genre Heterakis, Duj. T. c. pp. 117-120.
- RAILLIET, A., & Moussu, —. La Filaire des boutons hémorrhagiques observées chez l'âne. Découverte du mâle. C.R. Soc. Biol. iv, pp. 545-550.
- RANDOLPH, H. (1) Ein Beitrag zur Kenntniss der Tubificiden. Jen. Z. Nat. xxvii, pp. 463-476, pls. xvii-xix, and Viert. Ges. Zürich, 37ter Jhrg. pp. 145-147.
- ---. (2) The Regeneration of the Tail in Lumbriculus. J. Morph. vii, pp. 317-344, pls. xix-xxii.
- REPIACHOFF, W. O Ghastrulyatzii u Pozvonochn'ikh Zhivota'ikh s zamyechaniyami otnostel'no ghomologhii zarod'ishev'ikh plastov u Metazoa. Zapiski Novoross. Obsch. Estest. xvii (1), pp. 113-137.
- On the Gastrulation of *Vertebrates*, with remarks on the homology of the Germinal layers in the Metazoa. The development of *Nereis*, *Lopadorhynchus*, and *Clepsine* introduced and compared. (In Russian.)
- RETZIUS, G. Biologische Untersuchungen. Neue Folge, iii. Stockholm: 1892, 4to, 68 pp. 1. Das Nervensystem der Lumbriciden, pp. 1-15, pls. i-vi.
- BICHARD, J. Sur la présence d'un Cysticercoïde chez un calauide d'eau douce. Bull. Soc. Z. Fr. xvii, pp. 17 & 18.
- RICHES, T. H. A new British Nemertine (Carinella polymorpha).

  J. Mar. Biol. Ass. (n.s.) ii, pp. 284 & 285.
- RIEHM, G. Ueber die excretorischen Kanäle von Schistocephalus dimorphus. Z. Naturw. lxv, pp. 132-136, pl. ii.

- RITTER, E. (1) Balanoglossus as one of the generalised types in Zoology. Zoe, iii, pp. 187-199 (Errata, p. 282), pl. xxii.
- ---.. (2) Notes on an abnormal Polygordius Larva. Am. Nat. xxvi, pp. 1047-1050, 3 figg.
- RITZEMA-BOS, J. [See Bos, J. RITZEMA.]
- ORIVA, A. Sopra una casa di anguillulosi intestinale. Lo Sperimentale Firenzi, xIvi, pp. 40-69.
- ROHDE, E. (1) Muskel und Nerv. Zool. Beitr. iii.
  - I. Ascaris, pp. 69-106. [See also SB. Ak. Berlin, 1892, pp. 515-526.]
  - Mermis und Amphioxus, pp. 161-182, pls. xxiii-xxv. [See also SB. Ak. Berlin, 1892, pp. 659-664.]
  - III. Gordius, pp. 183-192, pls. xxv & xxvi. [See also SB. Ak. Berlin, 1892, pp. 665-667. "Giebt es Holomyaria?"]
- Rosa, D. (1) Die exotische Terricolen des k. k. naturhistorischen Hofmuseums. Ann. Hofmuseum. Wien, vi (1891), pp. 379-406, pls. xiii & xiv.
  - Results given in Italian in Boll. Mus. Zool. Tor. vii, No. 114 (1892).
- ——. (2) Viaggio di L. Fea in Birmania e regione vicine. xxvi Perichetidi, Seconda Parta. Ann. Mus. Genov. (2) x, pp. 107-122, pl. i.
- —... (3) I Lombrichi raccolti nell' isola di Engano dall Dott. E. Modigliani. Op. cit. (2) xii, pp. 542-548.
- —. (4) Kynotus michaelsenii, n. sp. (Contributio alla morfologia del Geoscolicidæ). Boll. Mus. Zool. Tor. vii, No. 119.
- —... (5) Descrizione dell' Allolobophora festas nuova specie di Lumbricide. T. c. No. 122.
- ——. (6) Descrizione dell' Allolobophora smaragdina nuova specie i Lumbricide. T. c. No. 123.
- —... (7) Megascolex templetonianus, n. sp. T. c. No. 131. Preliminary diagnosis.
- ROSSETER, T. B. On a new Cysticercus and a new Tape Worm. Quek. Club, iv, pp. 361-366, pls. xxii & xxiii (= Echinocotyle reteri, n. g. & sp., Blanch.).
- Rousseler, C. (1) On the best methods of examining Rotifers u the Microscope. Tr. Middlesex Soc. pp. 23-30, 3 figg.
- —... (2) On Notops minor, n. sp. J. Quek. Club (2) iv, pp. 359 & pl. xxiv.
- —. (3) On Two new Rotifers. T. c. pp. 367-370, pl. xxiv; [ar Micr. J. xiii, pp. 273-277.]
- ---. (4) On the Sense of Vision in Rotifers. T. c. pp. 371-2 & 377.
- Povelli, G. [See Grassi G., & Rovelli, G.]

- SAINT JOSEPH, —. Sur la croissance asymétrique chez les Annélides polychètes. C.R. cxv, pp. 887-890.
- SAINT REMY, G. (1) Synopsis des Trématodes Monogénèses (Continuation and Conclusion). Rev. Biol. iv, pp. 136-145, 184-191, 224-230, & 253-261.
- ——. (2) Matériaux pour l'anatomie des Monocotylides. Op. cit. v, pp. 46-52, 2 figg.
- ——. (3) Contribution à l'étude de l'appareil génital chez les Tristomiens. Arch. Biol. xii, pp. 1-56, pls. i & ii, and 3 woodcuts.
- ——. (4) Liste des Trematodes monogénèses recueillis à Roscoff. C.R. Ass. Fr. Sci. 1891 (2), pt. ii, pp. 571-574.
- —. (5) Les Idées actuelles sur le developpement et les relations des Cestodes et des Trematodes. Rev. Gén. Sci., 3° année, pp. 183-188, 10 figg.
- SANGALLI, G. Echinococco ed Œstrus nell' uomo. Rend. Ist. Lombardo, xxv, pp. 1167 & 1168.
- SCHULTÉN, M. W., & HOMÉN, E. A. Fall af Echinococcus i bäckenet och bukhålen. Finska Läkar. Förh. xxxii, p. 358.
- Sekera, E. Einige Bemerkungen über das Wassergefäss-system der Mesostomiden. Zool. Anz. xv, pp. 387 & 388. Criticism of Voigt (2).
- Selti, E. Sulle tenie dell' Hyrax dello Scioa. Genova tip. Ciminago: 1891, 8vo, 12 pp., 1 pl. Estr., Atti Soc. Ligust. ii, 2, 1891.
- SEYFERT, R. Beiträge zur Operation des intralocularen Cysticercus. Inaug. Dissert. Univ. Heidelberg. Leipsig: 1892, 8vo, 1 fasc.
- SHIPLEY, A. E. On Onchnesoma steenstruppii. Q. J. Micr. Sci. xxxiv, pp. 233-250, pl. ix.
- SIMMONS, W. J. The Human Blood Worm (Filaria sanguinis hominis). Sci. Goss. 1892, pp. 11-14, 18 figg.
- SKIMKEVICH, V. Otchet o zaghraníchší poyezdkye 1891 ghoda. Trudui St. Petersburg Est. xxii, 2 (1891), pp. 113-124, 4 figg. (Report on Foreign Travels in the year 1891.)
- 3 species of Balanoglossus. Nerve-cord of Balanoglossus compared with that of Amphioxus.
- SLAUGHTER, R. M. Filaria sanguinis hominis. The Discovery and Prevalence of the Disease in the U. S. A.: report of two new cases. Practise, 1891, pp. 329-335.
- Sonsino, P. (1) The Principal and most Efficacious Means of Preventing the Spread of Entozoal Affections in Man. Trans. of the 7th Internat. Congr. Hyg. Demog. London: 1891 (2), Sect. i, pp. 332-345.
- Gives distribution of various *Entozoa*, according to countries and organs of the body. Synopsis of exotic *Entozoa* in Man, with a table of their geographical distribution.

- [Sonsino, P.] (2) Dei Distomi dello "Zamenis viridoflavus, Lacèp.", e di una fase del ciclo vitale di uno di essi. Atti Soc. Tosc. viii, P.-v. pp. 91-95.
- 6—... (3) Tre casi di malattia da Rabdonema intestinale o Rabdonemiasi. Estr. Supp. Riv. gen. ital. Clinica Med. 1892.
- —. (4) Studi sui parassiti di molluschi di acqua dolce nei distomi di Cairo in Egitto. Festschr. z. 70ten Geburtstage R. Leuckarts. Leipsig (Engelmann): 1892, pp. 134-147, pl. xviii.
- Spencer, W. Baldwin. (1) Land Planarians from Lord Howe Island. Part 1. Description of Species. Tr. B. Soc. Vict. ii, 2, 1891 (1892), pp. 42-51, pls. v & vi.
- ——. (2) A Trip to Queensland in Search of Ceratodus. Vict. Nat. ix, pp. 16-31; Nature, xlvi, p. 306.
  - Land Planarians and Oligochests recorded.
- —. (3) Preliminary Description of Victorian Earthworms. Part I. The Genera Cryptodrilus and Megascolides. P. B. Soc. Vict. (n.s.) iv, pp. 130-156, pls. xiv.-xix.
- SPENGEL, —. Abnormitäten des Geschlechtsapparats von Distomum hepaticum. Verh. deutsch. Zool. Ges. 2te Jahresversamml. 1892, pp. 146 & 147.
- STADELMANN, H. Ueber den anatomischen Bau des Strongylus convolutus, Ostertag, nebst einige Bemerkungen zu seiner Biologie. Arch. f. Nat. lviii, pp. 149-176, pl. x.
- STILES, C. W. (1) Myzomimus scutatus (Müller), Stiles. J. Comp. Med. Vet. Arch. 1892, pp. 65-67, 1 fig. (Preliminary.)
- Subsequent paper on the same animal, with full anatomical details, in Festschr. z. 70ter Geburtstage R. Leuckarts. Leipsig (Englemann): pp. 126-134, 1 pl.
- —. (2) Filaridæ in body cavity of Horse and Cattle. J. Comp. Med. Vet. Arch. 1892, pp. 143-146, 1 fig.
- —. (3) List of Animal Parasites of Cattle. T. c. pp. 346-350.
- —... (4) Two cases of Echinococcus multilocularis in Cattle. T.c. p. 350.
- —. (5) On the presence of Strongylus ostertagi in America. T. c. pp. 147 & 148.
- —. (6) Distoma magnum, Bassi. T. c. pp. 464-466.
- —. (7) A word in regard to Dr. Francis' Distomum texanicum. Am. Vet. Rev. 1892, pp. 732 & 733.
  - Shows that it is D. americanum, Hassall.
- —. (8) Notes sur les Parasites: (13) Tania giardi (Riv.), Moniez; (14) Tania (Moniezia) expansa, Rad.; Moniezia (T.) planissima, n. sp. Bull. Soc. Z. Fr. xvii, pp. 157-159, and C.R. Soc. Biol. iv, pp. 664-666. Full anatomy in Bull. Bur. Anim. Industry promised.

[STILES, C. W.] (9) On the adult Cestodes of Cattle and Sheep (Abstr.).

P. Am. Ass. 41st Meeting. Rochester: 1892.

Discusses relations of *T. giardi* to Blanchard's new genus *Moniezia*. Full paper in Rep. U. S. Dep. Agric. promised.

- —. (10) Criticism of Linton [see Zool. Rec. 1891, Linton (1), (2), (3)]. Am. Nat. xxvi, pp. 65-68.
- —. (11) A method of Killing Nematodes for Microtome Sections. T. c. p. 972.
- ——. [See also HASSALL & STILES.]
- STOLL, —. Zur Zoogeographie der landbewohnenden Wirbelthieren. Viert. Ges. Zürich, xxxvii, pp. 233-274.
- Oligochæts and Nematodes, pp. 248-250. Land Planarians, pp. 253 & 254. Leeches, pp. 254-256.
- Stossich, M. (1) I Distomi nei Mammiferi. Lavoro monografico. Trieste: 1892, 8vo, 42 pp.
- Gives list of 61 species, with various hosts. List of 83 Mammals, with Distomida known to be parasitic in each. Clear and terse.
- ——. (2) I Distomi degli uccelli. Lavoro monografico. Boll. Soc. Adr. xiii, pp. 143-196.

Also with a reciprocal index.

- —... (3) Nuova serie di elminti veneti raccolti dal Dr. P. Alessandro conte Ninni. Glasnik naravosl. Druzt. vi (1891), pp. 216-219, pl. iii.
- —. (4) Osservationi elmintologiche. Op. cit. vii (1892), pp. 64-73, pls. i & ii.
- STRASSEN, O. ZUR. Bradynem rigidum, v. Sieb. Z. wiss. Zool. liv, pp. 653-747, pls. xxix-xxxiii.
- STRODTMANN, S. Die Systematik der Chætognathen und die geographischen Verbreitung der einzelnen Arten im nordatlantischen Ocean. Arch. f. Nat. lviii, pp. 333-377, pls. xvii & xviii. Abstr. in J. R. Micr. Soc. 1892, p. 792.
- STRÖSE, A. Ueber den feineren Bau von Strongylus micrurus. Deutsche Z. Thiermed. xviii, pp. 233-260, 3 pls.
- STUART, J. Search for Balanoglossus and Amphioxus at Marine Laboratory, Port Henderson. J. Inst. Jamaica, i, 2 (1892), pp. 44 & 45.

  New species of Balanoglossus found.
- THOMPSON, J. C. New Rotifer. Tr. Liverp. Biol. Soc. vi (1892), pp. 77-82.
- THOMPSON, P. G. (1) Moss-haunting Rotifers, with descriptions of two new species. Sci. Goss. 1892, pp. 56-61, 8 figg.
- —... (9) Notes on the parasitic tendency of Rotifers of the gonus *Proales*, with an account of a new species. T. c. pp. 219-221, 1 fig.
- TOKISHIGE, H. [See JANSON & TOKISHIGE.]
  1892. [VOL. XXIX.]

- TRYBOM, F. Mermielarver hos Chironomus. Ent. Tidskr. xiii, pp. 81-92.
- Tullberg, T. Ueber Conservierung von Evertebraten in ausgedehntem Zustande. Biol. Fören. iv, pp. 4-9.
  - Results given (in French) in Arch. Z. expér. (2) x, N. & R. p. xi.
- Tuttle, A. H. An interesting case of parasitism (preliminary). P. Am. Ass. 41st Meeting, Bochester, 1892, p. 201.
  - A nematode parasitic in the venom organ of the common Rattlesnake.
- TURNER, C. H. Notes upon Cladocera, Copepoda, Ostracoda, and Rotifera, of Cincinnati, with description of new species. Bull. Denison Univ. vi, 2, pp. 57-74, pls. i & ii. Rotifera, pp. 57-66, 1 pl.
- UDE, H. Ein neues Enchytræiden-Genus. Zool. Anz. xv, pp. 344 & 345.
- VAILLANT, L. Remarques sur les Némertiens d'eau douce. T. c pp. 125 & 126.
  - Criticism of PLESSIS, G. DU. (1).
- VÁVRA, V. [See FRITSCH, A., & VÁVRA, V.]
- VAYSSIÈRE, A. (1) Études sur le Temnocephala, parasite de l'Astacoides madagascariensis. Ann. Fac. Marseille, ii, pp. 77-99, pl. i. Abstr. in C.R. exv, pp. 64 & 65.
  - A new species.
- Vejdowsky, Fr. (1) Ueber die Encystierung von Ælosoma und der Regenwürmer. Zool. Anz. xv, pp. 171-175.
- —. (2) Zur Entwickelungsgeschichte des Nephridial-Apparates von Megascolules australis. Arch. mikr. Anat. xl, pp. 552-562, pl. xxxii.
- VETTER, B. Ueber Rotatorien. SB. Ges. Isis, 1892, p. 4.
  - Relation of Rotifers to Trochosphere larva doubted; no reasons given.
- VIERORDT, —. Ueber das Vorkommen des cystösen Echinococcus in Würtemburg. Med. Corresp. Bl. Würt. 1891, p. 137.
- VILLENEUVE, —. La Bilharsiose in Tunisie. C.R. Ass. Fr. Sci. Marseilles, 1891 (2), pt. i, p. 244, pt. ii, p. 556.
- VILLOT, A. (1) Encore un mot sur la classification des Cestiques. Zool. Anz. xv, pp. 210-212.
  - Maintaining his classification, as opposed to Grassi and Royelli.
- ——. (2) Gordiens de Sumatra. Description de deux espèces nouvelles. Zool. Ergebnisse einer Reise in Niederländisch Ostindien, Bd. ii, Heft. 2, pp. 136-138.
- Vogt, C. Sur le parasitisme transitoire d'une Turbéllariée triclade (Gunda sp.). C.R. Ass. Fr. Sci. 1891 (2), pt. i, pp. 239-240.
- Voigt, W. (1) Die Fortpflanzung von Planaria alpina (Dana). Zool. Auz. xv, pp. 238-241.

- [Voigt, W.] (2) Das Wassergefässsystem von Mesostoma truncatum, O. Sch. Zool. Anz. xv, pp. 247 & 248. Abstr. J. R. Micr. Soc. 1892, p. 616.
- —. (3) Karten zur Verbreitung der Planaria alpina und gonocephala; Polycelis cornuta. SB. niederrhein. Ges. (Naturw. Sekt.) 1892, p. 104.
  - In Siebengebirge, Feldberg, Altkönig in Taunus.
- VONCKEN, —. Empoisonnement par retention de toxines intestinales dû à la présence de vers intestinaux. Arch. Méd. Belge (1892) ii, pp. 8-12.
- WAGNER, FR. v. General observations on Fission and Gemmation in the Animal Kingdom. Ann. N. H. x, pp. 23-54. Transln. from paper in Zool. Jahrb. Abth. f. Anat. iv (1890). [See Zool. Rec. 1890.]
- WALTER, E. Ueber einige Monostomen aus dem Darme einer Schildkröte. Zool. Anz. xv, pp. 448-450. Abstr. in J. R. Micr. Soc. 1892, p. 616.
- WANDOLLECH, B. Zur Embryonalentwickelung des Strongylus paradoxus. Arch. f. Nat. lviii, i, pp. 123-148, pl. ix.
- WARD, H. B. On Nectonema agile, Verrill. Bull. Mus. C. Z. xxiii, No. 3, pp. 135-188, 8 pls.
- WASSILIEFF, E. Turbellaria rhabdocoela okrestnostel Varshavui.

  Tradui obsch. estest. Varsovie (Protok Otyeleniya biologhii) iii
  (1891-2), pp. 12-16.
- WATSON, A. T. The protective device of an Annelid. Nature, xlvi, p. 7. [See Zool. Rec. 1891, WATSON, A. T., Vermes, p. 25.]
  Identification of sp. Sabella saxicava.
- WAWRZIK, F. Ueber das Stützgewebe des Nervensystems der Chætopoden. Zool. Beitr. iii, pp. 107-128, pls. xiv-xix.
- WESTERN, G. Two male Rotifers hitherto undescribed. J. Quek. Club, iv, pp. 374 & 375, pl. xxv (part of), and Am. Micr. J. xiii, pp. 278 & 279.
- WHITMANN, C. O. The Metamerism of Clepsine. Festschr. z. 70ten Geburtstage R. Leuckarts. Leipsig (Engelmann): 1892, pp. 385-395, pls. xxxix & xl.
- Wierzejski, A. (1) Zur Kenntnis der Asplanchna-Arten. Zool. Anz. xv, pp. 345-349, 2 figg.
- —. (2) Crustacés et Rotifères d'eau douce recueillis en Argentine. Bull. Ac. Cracovie, 1892, p. 185.
- —. (3) Rotatoria (wrolki) Galicyi. Resumé. T. c. p. 402. From a book with the same title, and with 3 plates.
- —... (4) Atrochus tentaculatus, n. g. & sp., ein Räderthier ohne Räderorgan. Z. wiss. Zool. lv, pp. 696-712, pl. xxxii.

- WILLACH, —. A Nematode parasitic in nodules in colon of Macacus cynomolgus. Arch. wiss. prakt. Thierheilk. 1891, pp. 340-346.
- WILSON, E. B. The Cell Lineage of Nereis. A contribution to the cytogeny of the Annelid body. J. Morph. vi, pp. 361-480, pls. xiii-xx, and 9 woodcuts.
- WOODWARD, M. F. Description of an Abnormal Earthworm possessing seven pairs of ovaries. P. Z. S. 1892, pp. 184-187, pl. xiii.
- Zacharias, O. Das Vorkommen von Distomencysten betreffend. CB. Bakt. Parasit. xii, pp. 752 & 753.
- ZELINKA, C. Das subcesophageale Ganglion von Rotifern. Verh. deutsch. Zool. Ges. 2te Jahresversamml, 1891 (Exhibition of).
- ZOGRAF, N. (1) Les Cestodes offrent-ils des tissus ectodermiques? Arch. Z. expér. (2) x, pp. 331-343, pl. xiii.
- ——. (2) Note sur l'origine et les parentés des Arthropodes, principalement des Arthropodes trachéates. Congr. Internat. Zool., Moscow, 1892, pp. 278–296, 6 figg.

Relationship to Annelids dealt with.

- ZSCHOKKE, F. (1) Zur Lebensgeschichte des Echinorhynchus proteus, Westrumb. Verh. Ges. Basel, x, pp. 73-83.
- ——. (2) Seltene Parasiten des Menschen. CB. Bakt. Parasit. xii, pp. 497-500.
- ——. (3) Parasiten der Süsswasserfische. In: Die Tier und Pflanzenwelt des Süsswassers. v. Ö. Zacharias. Bd. ii. Leipsig (J. J. Weber): 1892.
  - Zykoff, W. Zur Turbellarienfauna der Umgegend von Moskow. Zool. Anz. xv, pp. 445-447.

## VERMES IN GENERAL.

Theory of mesoderm formation and metamerism; Cholodkowsky.—Discussion of *Vermes*, and especially of the different names given to the Nephridia in the different groups; Cosmovici (3) and v. Kennel (2).

# I.—HEMIOHORDATA.

[Balanoglossus, Cephalodiscus, Rhabdopleura.]

#### A.—BIOLOGICAL.

Balanoglossus, as one of the generalised types in Zoology; RITTER (1).

Morphology of Rhabdopleura, showing its relations to the Hemichordata; FOWLER, G. H.

Comparison of nerve-cord of Balanoglossus with that of Amphioxus; SKIMKEVICH.

Method of obtaining larves of Balanoglossus (improvement on Bateson's); MORGAN (2).

H

#### B.—FAUNISTIC.

Solovetzki: B. merejkowskii; Knipovicha.

United States: Balanoglossus kowalewskii in both Wood's Holl and Newport; Tornaria of New England, ? = that of B. krohnii, Bourne, of the Mediterranean; MORGAN (1).

Jamaica: a new species of Balanoglossus, Andrews (1), STUART. Nightingale I.: Rhabdopleura normanni, Allm.; FOWLER, G. H.

## C.—SYSTEMATIC.

Balanoglossus claviger and minutus, figured; SKIMKEVICH. B. krohnii: its Tornaria in New England? Morgan. B. kowalewskii in New England; Morgan (1). B. merejkowskii compared with Amphioxus; SKIMKEVICH;—from Solovetzki; KNIPOVICHA, pp. 24, 35, & 40. B., n. sp. (or n. spp.), from Jamaica; Andrews (1), STUART.

Rhabdopleura normanni, Allm.: morphology of; Fowler, G. H.

## II.—ANNELIDA.

[Including Gephyrea, Chætopoda, and Hirudinea.]

#### A.—BIOLOGICAL.

## 1. GENERAL.

On the relation of Arthropods (*Tracheata*) to Annelids; ZOGRAF (2) [see also p. 54 of Record, under *Platyhelminthes*: Excretory system; RIEHM].

Anatomy and histology of Branchiura sowerbyi, a new Branchiate Oligochæt; Beddard (5): of Onchnesoma steenstrupi; Shipley: of Sparganophilus tamesis, n. g. & sp., of aquatic Oligochæta; Benham (3): of Australian Chlorhaemidæ; Haswell (1): of Alciopidæ of Messina; Hering: of two Acanthodriloid Earthworms from New Zealand; Benham (2): of the Tubificidæ; Randolph.

Anatomy of Amphichata sannio, n. g. & sp. of Oligochat; Kallsteins: of Victorian Earthworms; Spencer (3): of Russian and Siberian Earthworms; Kulagin.

#### 2. Epidermis.

GOODRICH describes an Oligochæt, with a dense covering of fine hairs or bristles all over. Lenhossék (2) shows by Golgi's methods that the blood vessels penetrate right into the epidermis of the Earthworm, not exclusively in the chitellar region. Lenhossék (1) describes the terminations of sensory nerves in the Earthworms. Jourdain (1) describes them in Polychæts.

ı,

## 3. MUSCULAR SYSTEM.

Absence of spindle muscle, and presence of only one retractor muscle in a Sipunculid; SHIPLEY.

What was formerly taken for jaw muscles in the *Hirudinea* (Gnathobdellidæ) shown to be ducts of salivary glands; LEUCKART (2).

## 4. DIGESTIVE SYSTEM.

Alimentary canal of Hirudo fully described and figured; HENKING.

LEUCKART (2) describes fully the salivary glands of the *Hirudinea* with jaws, in which alone they are present. The duct of each gland (and of these there are about a hundred in *Hirudo*, a dozen in *Aulastoma*) divides, on reaching the root of the tooth, into two halves, which then meet again to open by a short narrow duct in the tooth to the exterior. He concludes that in *Aulastoma* the juice secreted must have some other function than that of preventing coagulation. He shows that besides the salivary glands there are other glands—maxillary glands and unicellular glands—histologically, chemically, and optically distinct from them, but also opening into the month.

BLANCHARD (6) proposes the name "pseudognaths" for the jaws of Nephelids, to distinguish them from those of other Gnathobdellida.

The cells lining the intestine of Lumbricus and their action are described very fully by GREENWOOD. Around and almost covering in the gland-cells, whose secretion serves for the digestion of the food, are other cells ciliated towards the lumen of the intestine, and serving for the absorption of food. Whilst fat is being digested, however, the cilia disappear, and a striated border to the cell appears in their place. The ciliated cells are more evident in the walls of the typhlosole than of the intestine proper.

In a new genus of Oligocheet, allied to *Chætogaster*, it is shown that the lumen of the anterior part of the alimentary canal of the asexually-produced bud, is formed by the fusion of two secondary lateral lumins, on either side of the old alimentary canal; KALLSTENIUS.

#### 5. CORLOM.

JOURDAIN (3) shows that the coelom is lined by a ciliated epithelium in Sipunculids, and refers to instances of the same thing in other animals, vertebrate and invertebrate, concluding therefrom that the word "endothelium," in anatomy, should be done away with. He is criticized by Malassez.

## 6. Connective Tissue.

Yellow bodies in the connective tissue elements of Sipunculids shown by JOURDAIN (2), to be due to granules of a glycogenous matter, serving as reserve food material.

#### 7. CIRCULATORY SYSTEM.

Circulation of Lumbricus; GRIFFITHS (1); and LENHOSSÉK (2), where it is shown that blood vessels penetrate right into the epidermis.

Chemical constitution of the blood in Gephyreans, Chetopods, and Leeches; GRIFFITHS (2). Hermérythrine, a new iron-containing respiratory pigment in the blood of Sipunculids; GRIFFITHS (3).

Vascular system of Chlorhaemidæ; HASWELL (1).

Absence of vascular system in a Sipunculid; SHIPLEY.

## 8. RESPIRATORY SYSTEM.

SHIPLEY maintains that the intestine is the chief respiratory organ in the unarmed Gephyreans, and that the vascular system serves merely to extend the tentacles.

BEDDARD (4, 5) describes a new Oligochet with about fifty pairs of branchies, one pair per segment, at its hinder end. These are simple processes, situated on the middle dorsal and ventral line, not ciliated; the coelom extends into them, and blood vessels lie just beneath the epidermis on each side of each, breaking up into capillaries in the integument.

## 9. EXCRETORY SYSTEM.

Development of nephridia in Megascolides; Vejdowsky (2): in Acanthodrilus multiporus, Beddard (6). In Megascolides functional nephrostomes appear never to be developed at all except in the posterior segments, where the meganephridia afterwards appear; anteriorly the rudiments of them soon disappear and micronephridia are developed straight away, with intercellular efferent ducts, and not at first connected by longitudinal ducts. In Acanthodrilus all the nephridia in the youngest embryos have functional nephrostomes, which all become rudimentary later except for those in segments xi-xiv; they are formed as meganephridia; secondary openings to the exterior are formed later, when the micronephric condition is assumed; the second pair, and then one or two other pairs, become connected with the first to form the "mucous gland;" opening into the stomodæum; the anal nephridia are formed late, and open into the mesenteron, not into the proctodæum; the gonaducts are developed from the nephridial funnels and a small portion of the succeeding tube.

Kallstenius shows that in a new genus of fresh-water Oligocheet the nephridia are fused with the ventral vessel throughout their course.

Bolsius (2) describes the nephridia of fresh-water Leeches, showing that in Nephelis, Clepsine, and Hemiclepsis, there are to each nephridium three distinct unicellular canals, each arising from a separate group of branching intracellular tubules, and uniting with one another only just before opening to the exterior. In Hirudo medicinalis he proves the existence of multicellular glands opening into the intracellular canals. He gives a review of the work hitherto done on the nephridia of Leeches.

He also shows (1) that the so-called ciliated "funnels" have no connection with the nephridia.

Only a single nephridium in a Sipunculid; SHIPLEY.

Excretory organ of Coppingeria (n. g. of Chlorhæmidæ); HASWELL (1) Individual variation in the position of the nephridiopores in Earth worms; HUBRECHT—the conclusion being arrived at that the mega nephric arrangement of nephridia is probably the more primitive: in Leeches; BLANCHARD (6) and (3, 111).

## 10. NERVOUS SYSTEM.

Several very important papers on the Nervous System of Annelids and Nematodes [see Nematode part of Record] throw light on the histology of nervous systems in general. v. Lenhossék (1) and Retzius agree that in Lumbricus the sensory nerve-cells do not lie in the nerve-cord nor in special ganglia, but in the epidermis; from them fibres go to the nervecord dividing there into an ascending and a descending branch, which run on the same side of the body as the nerve enters—through the length of three ganglia according to v. Lenhossek, not quite as far as the next ganglion according to RETZIUS—and so end in the "Punktsubstanz," not uniting with ganglion cells. RETZIUS shows, further, that the motor nerve-cells lie in the ganglia of the ventral cord and send processes which branch into the "Punktsubstanz" as well as nerves to the periphery, which there branch to the muscles; these nerves sometimes arise from ganglion cells of the same side, sometimes from those of the opposite side of the body, so that there is a certain amount of decussation; in each ganglion there is one multipolar cell of unknown function. CERFONTAINE maintains the nervous nature of the giant-fibres in Lumbricus; they are, however, very special nerve fibres resulting from the union of the prolongations of several axis cylinders coming from distinct cells; they give off branches in each ganglion, which connect them either with other elements of the nerve-cord, or with the periphery; the median giant-fibre develops from in front backwards, and the lateral ones from behind forwards, the currents in them are, therefore, going in opposite directions; their function, he concludes, is to connect lengthwise the different parts of the nervous system, and to allow for the simultaneous muscular contraction in the whole extent of the body. APATHY (2), working at Leeches, regards each nerve as a bundle of primitive fibrils surrounding some central protoplasm; these fibrils are supposed to carry the nerve impulse. He is criticized by ROHDE (1, 1).

WAWRZIK discusses the supporting structures of the Nervous System in Chætopods.

## 11. SENSE ORGANS.

JOURDAIN (1) describes sensory nerve-terminations in several Polychæts, situated generally on the appendages, but sometimes on the body-

wall where there are no appendages, consisting sometimes of cells of the Vertebrate olfactory type, sometimes ciliated, and sometimes much like the ordinary epithelial cells.

Eyes: The derivation of the Vertebrate from the Annelid eye; V. KENNEL (1). A Sipunculid, with pigment spots or eyes; Shipley. Eves of Polychata: Andrews (3): the structure of the adult eves of 26 species is described and, for most of them, figured; the development of the eye is given in 5 species; it is found that each eye is a pigmented cup, formed of one layer of retina cells all essentially alike, and consisting each of a peripheral body containing pigment and a central clear rod; some of the cells are much attenuated; the lens is continuous with the cuticle, sometimes also with the retinal cells, and is either secreted by the retinal cells, or formed from the fused central ends of them; the connection with the epidermis persists (except in the Alciopida, where there the development of the eye is exceptional), the eye being formed not as an invagination, but by the elongation and pigmentation of a few epidermis cells, whose central ends become the rods by the withdrawal of the pigment to the opposite end. BERANECK describes the development of the eye in the Alciopida, showing that it is built on quite a different plan to the Arthropod eye. The eyes of certain of the Chlorhaemidæ are described by HASWELL (1): in one genus they are on special processes of the prostomium in the bases of which are optic ganglia which are outgrowths of the brain. MAIER describes the minute structure of the eyes in the Hirudinea, showing that in Hirudo and Aulastomum, while the main branch of the nerve enters the eye posteriorly, there is also a small one entering it anteriorly and ventrally; while in Nephelis, Clepsine, and Piscicola there is only the one branch entering the eye in front. BLANCHARD (6) mentions asymmetry of the eyes as occurring often in Nephelis, but seldom in Hirudo.

Auditory Organs: EHLERS (1) describes very fully the structure of the so-called "auditory organ" in the different species of Arenicola: in A. claparedii there is only an open groove, with no otoliths or neuroepithelium; in A. marina the otocyst, though deep-lying, is still connected with the exterior by a narrow canal, and contains foreign particles; in A. grubei and A. antillensis the sacs are quite closed and the otoliths are intrinsic formations; in no species are there hairs or rods to the cells forming the sac, and the author suggests that they more probably serve as organs of equilibration than as auditory organs; he shows that they have nothing to do with the "Nackenorgane," nor do they correspond to the setal sacs of the parapodia; they may, however, represent modified cirri.

Embryonic sense organs: BEDDARD (6) describes in Acanthodrilus an unpaired sense organ in the young embryo, consisting of a few large cells on one side of the stomodæal aperture; in the advanced embryo he also finds peculiar cells in the epidermis, which he thinks form a sense organ.

#### 12. GENITAL SYSTEM.

BEDDARD (6) shows that there are originally 4 pairs of gonads with their ducts developed in Acanthodrilus in segments x-xiii; and that the one in the 12th segment afterwards degenerates. WOODWARD describes an abnormal Allolobophora with 7 pairs of ovaries. BLANCHARD (4) describes a Leech with two pores on the female segment (xi), one between the 32nd and 33rd, the other between the 35th and 36th annuli; of these, sometimes the one, sometimes the other, is the larger; their significance is not shown. Rosa (1) describes an Earthworm (Paradrilus rosas, Mich.) in which the spermatheces are in communication with the intestine, as well as with the exterior, as in the Enchytræidæ. Goodrich describes a Tubificid, in which the spermatheces open by a median pore in the 10th segment, and in which the sperm-ducts also open by a median atrium.

## 13. Reproduction.

Fertilization of Clepsine: MOLLER. Hybridism in Earthworms; BENHAM (2), FRIEND (8), BEDDARD (11). Asexual reproduction in Amphichaeta, n. g. of Oligochset, Kallstenius. Encystment: [see under "Habits," &c]. Chun suggests that the process to which he gives the name of Dissogony, the ripening of one individual in two different stages of its development, may take place in Nereis. Prouho discusses the different kinds of individuals, with regard to sex, in the Mediterranean Mysostomas, showing that in one species (M. alatum) the sexes are distinct in the young, the males being much smaller than the females, but that then the males develop ovaries, thus becoming hermaphrodite.

## 14. DEVELOPMENT.

The recent researches into the germ-layers of Annelids (Nereis, Lopuclorhynchus, and Clepsine), as affecting the interpretation of the germlayers of Vertebrates; Replachoff. The development of Nereis is very
fully and carefully worked out by Wilson, the whole history of each cell
of the segmentation sphere being given; it is shown that, owing to the
large size of the left posterior macromere, the cells of the segmentation
sphere have a spiral instead of radial arrangement; bilateral symmetry is
only assumed later, after the formation of the mesoblast; the question of
bilateral symmetry in the cleavage of other eggs is discussed, and it is
concluded that in all cases it results from the anticipation of the bilateral
symmetry of the adult; in an appendix the author mentions several other
Polychæts which he has examined, in which he also found spiral segmentation of the ovum, continuing even to a later period.

Development of Arcnicola; EHLERS (2): The young of A. marina is pelagic though not found at the surface; its body consists of 2 regions, the one comprising prostomium, buccal segment, and 20 succeeding double segments, the other numerous, narrow and non-setigerous segments; there

are no branchise. Points in development of Acanthodrilus; BEDDARD (6). Comment on Apáthy's description of the development of Hirudinea [see Zool. Rec. 1891, Vermes, APATHY (2)]; BERGH: provisional larval muscles of Gnathobdellidæ, and larval skin drawn attention to; teloblasts said to have been not clearly interpreted by APATHY.

# 15. METAMERISM AND VARIATIONS THEREIN; ALSO OTHER VARIATIONS.

WHITMAN contributes an important paper on the metamerism of Leeches. He concludes that the head consists merely of a certain number of true metameres (1-6), and these have the same ganglia as the other metameres, only somewhat displaced; also that the typical number of annuli per segment is 3, so that when fewer or more are found either suppression or division of one or more annulus must have taken place. The trunk is formed by two germ bands, which are at first dorsal, but grow round the yolk mass to meet in the mid-ventral line from in front backwards, and segmentation then begins on the ventral surface; segmentation of the head begins in front of the bands and on the dorsal surface only, after the segmentation of the germ-bands on the ventral surface but before their segmentation on the dorsal surface. The metameric sense organs and their nerve supply are very fully described and figured.

Variation in normal type of segmentation for *Trocheta*, owing to fusion of annuli; BLANCHARD (6). Variation in number of annuli in specimens of *Glossiphonia*, from same locality; BLANCHARD (3, IV).

Spiral Metamerism: both Cori and Morgan (3) have independently shown in Cheetopods the frequency with which, in certain genera, normal circular segmentation is replaced by spiral segmentation for a greater or less number of segments. Saint Joseph also shows that a great deal of asymmetry prevails in the Sabellidæ, and it would seem that his cases may also be treated as cases of spiral segmentation. Blanchard (6) mentions the occurrence of the same thing in Leeches.

Other cases of asymmetry: in male genital pore of Oligochests; FRIEND, (14) p. 108, COLLIN, A. (1): in Clitellum; BEDDARD (11): in eyes of Leeches; BLANCHARD (3, v) and (6).

Bifurcation: recorded instances in Chætopods collected; Andrews (2). Tail double, the two halves meeting at the posterior extremity, and separated from one another anteriorly, in an Oligochæt; FRIEND, (14) pp. 108 & 109. Head double, the one half, consisting of two segments and prostomium only, moving like a feeler to the other; FRIEND, (14) p. 161.

Symmetrical variations in position of nephridial and genital pores in Leeches; Blanchard (3, 111) and (6).

Variations in number: an Allolobophora with seven pairs of ovaries though with only the usual one pair of oviducts; WOODWARD.

An abnormal larva of Polygordius; RITTER (2). A larva is described

which behaves very much in the way of a *Phoronis* larva, a bud being given off behind the head, into which the alimentary canal extends, opening to the exterior at its extremity, for a certain length of time. How the original axis becomes the adult axis, and the bud disappears, was not observed, though it is assumed that this happens.

An albino Perichæta monilicystis; MICHAELSEN (1).

# 16. SYNONYMY, CLASSIFICATION, &c.

Synonymy of certain Polychests, including several of Schmarda's species; MICHAELSEN (4): of some northern forms; V. MARENZELLER (1). Schmarda's species of Oligochests renamed on examination of the type specimens; BEDDARD (2).

Classification of Annelids : HATSCHEK.

Value of colours and markings of Leeches in forming new varieties; Nusbaum (2).

[See also Systematic part of Record.]

# 17. Habits, Properties, and Stray Notes.

Temporary encystment of Ælosoma; Beddard (1), Vejdowsky (1). Vejdowsky compares with this the formation of special burrows by certain Lumbricide, and ascribes both sets of phenomena to the rest required after a period of asexual reproductive activity. An Annelid (Leucodora coeca) living in a sponge (Microciona); HORNELL (2). Annelid with tube coiled like the frond of a fern [Zool. Rec. 1891, Vermes, p. 25], shown to be Sabella saxicava; WATSON. Polychæts parasitic on Echinoderms; MATZDORF (3), and CUENOT. Mysostomas parasitic on or in Antedon phalangium; PROUHO. Horse Leech in nasal mucous membrane of man; CONDORELLI (4). Leeches and Earthworms swallowed alive by the Bittern; LIEBE. Clepsine tesseluta in posterior nares of Swan; COLLIN, A. The blood of Leeches as a medium for the cultivation of Huema-Earthworms as fortilizers of the soil; Loz: as the tozoa ; Labbe. carriers of the tubercle bacillus; LORTET & DESPRIGNES. A protozoon (Rhabdostyla arenicola) parasitic on the branchise of Arenicola marina; MATZDORF (2). On the conditions of existence of pelagic organisms, including Polycheets and Sagitta; KOEHLER. Phosphorescence in Earthworms; HILLIGER, GIARD. The dissemination of Leeches by Palmipeds; Guerne (1) and Blanchard (2).

## 18. METHODS.

For nephridia of Leeches; Bolsius (2). For eyes of Leeches; Maier, p. 559.

For preserving Oligocheets and Leeches, paralysing with magnesium chloride and killing with chromic acid recommended; Tullberg. For preserving Leeches and Polycheets; Cunningham. For rearing Polycheet larvee; Bles.

#### B.-FAUNISTIC.

## LAND AND FRESHWATER FORMS.

Note on distribution of Oligochæts and Leeches in general; STOLL.

# a. Palæarctic Region.

British Isles: British Tree- and Earthworms, 6 species of Allolobophora, 5 of Lumbricus recognized; Friend (2). New British Earthworms and summary of old ones, 21 species recognized; Friend (4): criticized by Hurst & Benham (6): Lumbricus eiseni in England; Friend (3). Earthworms of Yorkshire and North Britain; Friend (11, 12): of Northamptonshire; Friend (6): of Essex; Friend (1): of Middlesex; Friend (7): of Sussex; Friend (10): of Kent; Friend (5). A new species of Earthworm from Bangor; Friend (9): from Ireland; Friend (13). Sparganophilus, n. g. of Oligochæt in the Thames; Benham (3). Vermiculus, n. g. of Oligochæt, brackish water, Weymouth; Goodbich.

Western Europe: distribution of *Photodrilus phosphoreus* in; MATZ-DORF (1) and GIARD. *Trochæta subviridis* confined to Western Europe; BLANCHARD (6).

France: Depts. Vosges and Meurthe as two new stations for *Phreorycles menkeanus*, Hffmst.; BRUNOTTE.

Germany.—Mecklenburg Earthworms; MICHAELSEN (2). Earthworms in the neighbourhood of Berlin (3 species Lumbricus, 8 Allolobophora, 1 Criodrilus, 1 Allurus); Collin, A. (1). Altenar: new genus of Enchytræidæ found under moss, Bryodrilus ehlersi; UDE. Saxony: Nephelis atomaria, var. meyeri, n. var.; Blanchard (3, 111).

Austria.—Bohemia: Leeches and Chatopods; KAFKA, FRITSCH & VÁVRA. S. Austria: Xerobdella lecomtei in the mountainous regions of; BLANCHARD (4).

Poland: Leeches and Earthworms of; Nusbaum (2). The colours of the Leeches are shown to vary constantly with the locality, and several new varieties are described.

Finnland: 2 Oligochæts; LEVANDER (2).

Russia and Siberia: Earthworms of; KULAGIN.

China.—Shan-tong: no Earthworms ever seen there; FAUVEL. Amoy: Perichæta aspergillum, Perrier; ROSA (1). Foo-chow: P. sinensis, n. sp., BEDDARD (9).

Japan: 4 species of Perichætidæ of which 3 new; Beddard (11). Another new species of Perichæta and P. sieboldii; Rosa (1). 3 new species of Perichæta and 1 new species of Moniligaster; Michaelsen (1). Liguria: Trochæta subviridis recorded in; Blanchard (6).

Sardinia: 1 Leech (Hamopis vorax) and 2 Oligochests; MARCIALIS: Microscolex modestus; ROSA (1). Sicily: Hormogasta redii; ROSA (5).

Algeria and Tunis: Microscolex algeriensis, n. sp., and Allolobophora

complanatus; BEDDARD (8). Tunis: Allolobophora festæ, n. sp., 1
(5). Tripoli and Egypt: Allolobophora trapezoides, Dugés; Rosa (5
Madeira: Microscolex poultoni, n. sp., BEDDARD (8).

Azores: Perichæta indica; MICHAELSEN (1).

# b. Ethiopian Region.

W. Africa: 4 new species of a new genus, Gordiodrilus, of the fa Ocnerodrilidæ; BEDDARD (3). Togo Land: 5 new species of Benha 1 of Eudrilus; MICHAELSEN (1). Valley of Niger: made fertile Siphonogaster; Loz. Fernando Po: Paradrilus rosae, Mich.; Rosa

Central Africa.—Near Victoria Nyanza: 1 new species of Benha 2 of Pygmaodrilus, 2 of Siphonogaster; MICHAELSEN (2).

S. Africa.—Cape Colony: Microchata belli, n. sp., Benham (1) rappii, Bed., Rosa (1); Allolobophora trapezoides, Rosa (1).

Natal : Microchata papillata, n. sp., BENHAM (1).

Madagascar: Kynotus michaelsenii, n. sp., ROSA (4); K. kelleri, n. MICHAELSEN (1).

Mauritius: Perichata mauritiana, n. sp., Beddard (9). Mahé Island: Perichata pentacystis, n. sp., Rosa (1).

# c. Oriental Region.

Ceylon: Megascolex templetonianus, n. sp., Rosa (7); Typhæus la Rosa, Rosa (1); Perichæta taprobanæ, n. sp., Beddard (9).

Burmah and neighbouring Region: List of Oligocheets, including 4 species of *Perichata*, 1 of *Perionyx*, of which descriptions are giv Rosa (2). Bangkok: *Perionyx excavatus*; Rosa (1). Malacca: a spe of *Urochata*; Rosa (1).

Singapore: Perichata morrisi (?) notes on; BEDDARD (9). Megascarmatus. Bed.; Rosa (1).

Nicobar: Lumbricus rubellus, Hffm., and Allolobophora fætida, St Rosa (1).

Penang: Perichæta morrisi, n. sp., BEDDARD (9).

Sumatra: Desmogaster, Horst; Perichæta longa, n. sp., MICHAELSEN Engano: absence of Perionyx excavatus and Megascolex armatus, prosence of Perichætidæ much like the Indian ones show Indian chara of fauna; 3 new species and 1 new subspecies of Perichæta, and 1 spe of Urochæta (U. corethrura, Müll.) described; Rosa (3).

Banka: Perichæta martensi, n. sp., MICHAELSEN (1).

Java and Borneo: Perichæta racemosa, n. sp., Rosa (1). Java: operculata, n. sp., Rosa (1); P. udekemi, Gr., MICHAELSEN (1). Born P. mandhorensis and Megascolex pictus, n. spp., MICHAELSEN (1).

Malay Archipelago generally: 21 species of Earthworms, amongst we are 8 new species, and 2 new genera (Glyphidrilus and Annadrilus, beloing to the fam. Geoscolecidae), gen. Benhamia, recorded here for the 1 time; Horst.

Philippine Islands.—Luzon: Perichata pulchra, n. sp., Pleinogaster jagori, n. g. & sp.; Samar, Megascolex iris and margaritaceus, n. spp., Pleinogaster samariensis, n. g. & sp., Michaelsen (1). Cebà: Perichata philippina, n. sp., Bosa (1).

# d. Australian Region.

New Guines: Perichata neoguinensis, n. sp., MICHAELSEN (1).

Ard Island: Microchaeta benhami, n. sp., Rosa (1).

Australia: Hæmadipsa in, as well as in Old World; STOLL. Queensland: Perichæta gympiana, n. sp., and Cryptodrilus purpureus; SPENCER (2). N. S. Wales: Sydney, Didymogaster sylvaticus; ROSA (1). Victoria: Earthworms of; SPENCER (3), comprising 11 new species of Cryptodrilus (5 from Gippsland), 10 new species and 1 new variety of Megascolides (2 of them and the variety from Gippsland), and 2 known species of Megascolides.

New Zealand: Neodrilus monocystis, Bed., and Plugiochata punctata, n. g. & sp.; Benham (2).

Tahiti: Perichæta grubei and novaræ, n. spp. (for P. taitensis), Rosa (1).

Marquesas: Perichæta albida, n. sp., MICHAELSEN (1). Hawai: Perichæta hawayana, n. sp., and Allolobophora subrubicunda, Eisen; Rosa (1).

# e. Neotropical Region.

Argentine Republic: Microscolex dubius; ROSA (1).

Chili: Glossiphonia tesselata, and discussion as to how it got there; BLANCHARD (2). Acanthrodrilus platurus, n. sp., MICHAELSEN (1). Chiloe Island: Acanthodrilus pictus?, Mich.; ROSA (1). A. platurus, n. sp., MICHAELSEN (1). Valparaiso: "large phosphorescent Earthworm"; HILLIGER.

Bolivia: Upper reaches (salt) of Pilcomayo R., Kerria halophila, n. g. & sp. of fam. Cryptodrilidæ; BEDDARD (10).

Brazil: Geoscolex maximus; ROSA (1). Tykonus grandis, n. g. & sp., Pontodrilus arenæ, F. Mull.; MICHAELSEN (1). S. Brazil: Cylicobdella; STOLL.

Venezuela: Eudrilus roseus, n. sp., Benhamia bolavi, Mich., Anteus, 3 n. spp.; MICHAELSEN (1).

Ecuador: on Cayambe, at height of 14,000 feet, Rhinodrilus ecuadorensis, n. sp.; BENHAM (4).

Guatemala: Cylicobdella; STOLL.

Ciudad Durango: Benhamia mexicana, n. sp., ROSA (1).

Dominica: Gordiodrilus dominicensis, n. g. & sp., BEDDARD (3).

Barbados: Perichæta barbadensis, n. sp., BEDDARD (9).

Jamaica: peculiar Earthworm on trees at Moneague; Andrews (1).

# f. Nearctic Region.

Bermuda: Perichata bermudensis, n. sp., BEDDARD (9).

Queen Charlotte's Island : Plutellus perrieri, n. sp., BENHAM (1).

## MARINE FORMS.

## a. Arctic Ocean.

Off East Spitzbergen: Polychæts collected by Bremen Expedition, species, with notes on anatomical details of many of them, and with li of synonyms; v. MARENZELLER (1).

White Sea.—Solovetzki: Polychests (pp. 25-27), A. priapulus (pp. & 40); Knipovicha.

## b. N. Atlantic Ocean.

Norway.—N. W. Coast: Onchnesoma steenstrupii, n. sp., Shipl. Bergenfjord: Polychæts and Gephyreans; Appellöf.

Baltic Sea.—E. Baltic: 8 species of Polychæts, 2 Gephyrea Kojevnikov.

North Sea.—St. Andrew's Bay: Clymens ebiensis; McIntosh.

Irish Sea.—Liverpool Bay: 88 species of Polychets, of which 1 n 3 others new to Britain, 2 new varieties; correction of a previous from the same locality by another author; separation of species accord to their vertical range into littoral, deep water, and indefinite forms, a table of the geographical and bathymetrical range of the local P chests; HORNELL (1).

English Channel.—Off the Coast of Devonshire: Staurocepherubrovittatus, Dinophilus tæniatus; Garstang (1): Protodrilus leuckarte Plymouth; Bles: Polychæts of Plymouth, 63 species, of which 5 r. 2? forming new genera, 7 other species new to Britain; Buchan Channel Islands: Sabella saxicava; Watson. Banyuls-sur-Mer: Eugigantea, 3 metres long, recorded; Lacaze-Duthiers.

Bay of Biscay.—Coast of Loire Infe.: 1 Polycheet and 1 armed Gerrean; CAMUS. Santander: 6 species Polycheets and 1 Phascolosomeruser

Mediterranean.—Sardinia: 5 species Polychæts, 2 Gephyreans; M CIALIS. Messina: anatomy of 7 new species of Alciopidæ; Hering.

Black Sea: 17 species of Polychæts, 1 new genus and 6 new speci Périaslavzeff.

Caribbean Sea.—Jamaica: Sipunculids rare; Polychets, no pela forms, Amphinomidae and Sabellidae common, but Eunicidae n Andrews (1).

## c. Indian Ocean.

Zanzibar: 8 species of Gephyreans (3 new ones of Thalassem: Fischer, 1 species Phymosoma; Collin, A. (3).

Red Sea: 3 Polychæts; Boutan.

Ceylon: 14 species of Polychests, including 1 new genus and 4 n species; MICHAELSEN (4).

Indian Coast: a tubicolor annelid; Keswell.

# d. S. Pacific Ocean.

The Australian species of Branchellion; BLANCHARD (3, VI).

Torres Straits.—Darnley Isl.: Coppingeria longisetosa, n. g. & sp. of Chlorhæmidæ; HASWELL (1).

Off Sydney: a second species of *Phoronis* (? = P. psammophila, found at Messina), and remarks on other species; HASWELL (4, 1): two new species of Stylaroides; HASWELL (1).

New Zealand: new species of Branchellion occurring on Raia nasuta; Dujardinea, from Port Chalmers; PARKER.

S. America.—Off Callao: 2 Sipunculids, 1 n. sp.; Collin, A. (3).

#### C.—SYSTEMATIC.

## GENERAL.

HATSCHEK proposes, in a preliminary paper, a new classification of Annelids, dividing them into the 4 Subclasses:—-

- 1. Archiannelida, including the Polygordiida and Dinophilida.
- Chatopoda, including the Protochata (Saccocirrida), Polychata, and Oligochata.
- 3. Hirudinea.
- 4. Echiurida.

With 2 supplementary groups: (5) the Sipunculacea, and (6) the Chætognatha. He gives a further classification of the two first subclasses, deriving all Chætopods (and hence the Hirudinea) from the Archiannelida, through the Protochæta and Spionidæ. He gives separate tables of the relations between some of the different families of Errant Polychæts.

# 1. GEPHYREA (including Phoronis).

## A. PHORONIS.

BLOCHMANN (3) discusses the affinities of the *Brachiopoda*, concluding that they are most nearly related to *Phoronis*. *Phoronis australis*, living in wall of tube of *Cerianthus*, and *P. psammophila*? in Port Jackson; HASWELL (4, 1).

## B. UNARMED GEPHYREANS.

Aspidosiphon cumings, from Zanzibar, remarks on; FISCHER.

Cleosiphon aspergillum, from Zanzibar, remarks on; FISCHER.

Dendrostomum perurianum, n. sp., Callao, Collin, A. (3), anatomy.

Onchnesoma steenstrupis, n. sp., N. W. Coast of Norway, Shipley, anatomy and histology. An Onchnesoma in the Bergenfjord;

Appellof.

1892. [VOL. XXIX.]

Phascolosoma catherina, Fr. Müll., Callao, and P. sanderi, n. sp., locality?, Collin, A. (3), descriptions.

Phymosoma granulatum, from Zanzibar, description; Collin, A. (3).

P. scolops, from Zanzibar, description; FISCHER.

Priapulus caudatus, from Solovetski; KNIPOVICHA.

Sipunculus indicus, Zanzibar, internal anatomy; FISCHER: conclusions arrived at differ from those of SLUITER [see Zool. Rec. 1891, Vermes, p. 22]. S. camanensis, also from Zanzibar, description; FISCHER. Blood of Sipunculus; GRIFFITHS (2): yellow bodies; JOURDAIN (2): coelomic epithelium; JOURDAIN (3).

## C. ARMED GEPHYREANS.


Bonellia viridis, from Sardinia; MARCIALIS.

Thalassema kokotoniense, stuhlmanni, and leptodermon, n. spp., Zauzibar, Fischer, anatomy. T. neptuni, in Bay of Biscay; Camus.

### 2. CHÆTOPODA.

## A. POLYCHETA (including DINOPHILUS and PROTODRILUS).

[See above under "General."] HATSCHEE divides the Polychæta into the two main groups, Cirrifera (including suborders Spiomorpha, Amphinomorpha, and Rapacia with the Myzostomidæ as an appendix) and Acirra (including suborders Drilomorpha, with Sternaspidæ and Ctenodrilidæ as appendices, Terebellomorpha, and Serpulimorpha). He expresses in different trees some of the relations between the different families of Rapacia:—



Aglaurides fulgida, from Ceylon; MICHAELSEN (4).

Alciopidæ, of Messina, full anatomy, histology, and synonomy of 7 species; Herring.

Ampharete grubei, recorded as British; HORNELL (1).

Anaitis: remarks on the genus and on the species, A. lineata, Clp., from which A. maculata, n. sp., is distinguished, although both were included by Claparède under the same name; Périaslavzeff. A. wahlbergi, remarks on genus and species; v. Marenzeller (1), p. 405.

Aphrodite obteta, recorded as British; Buchanan. Blood of Aphrodite; Griffiths (2).

Apomatus globifer, described and figured; v. MARENZELLER (1), p. 430.

Arenicola marina, development; EHLERS (2), HORNELL (1). Auditory organ of different species of Arenicola; EHLERS (1). A Protozoon parasitic on the branchiæ of A. marina; MATZDORF (2). Blood of Arenicola; GRIFFITHS (2).

Aricia norwegica, remarks on previous descriptions; APPELLÖF.

Autolytus alexandri, recorded as British; Hornell (1), = A. verrilli, n. sp., for the (Stephanosyllis ornata) of Verrill, described and figured; v. Marenzeller (1), pp. 416-420. A. prismaticus, and A. sp.? (2 only), from Spitzbergen, descriptions; v. Marenzeller (1), pp. 420 & 422. A. rubovittatus, Clp., Black Sea, remarks on; Périaslavzeff.

Brada inhabilis, Spitzbergen, description and figure; v. MARENZELLER (1), p. 427.

Callizona: detailed description and classification of the genus and its five species, with special anatomy, histology, and figures of C. angelina, n. sp.; APSTEIN.

Capitella multioculata, n. sp., with 6 pairs of eyes, from Black Sea; PÉRIASLAVZEFF.

Chaetopterus tubes containing Crystallogobius; APPELLÖF.

Cirratulus cirratus, recorded from Spitzbergen; v. Marenzeller (1).

C. riridis, with remarks on the genus, and description and figures of the species; Périaslavzeff.

Clymene ebiensis, new description of species; McIntosh.

Coppingeria, n. g. of Chlorhæmidæ with C. longisetosa, n. sp., from Australia, description, anatomy, and figures; HASWELL (1).

Dasychone herdmani, n. sp., Britain; HORNELL (1).

Dinophilus taniatus, recorded on Devonshire Coast; GARSTANG (1).

Drieschia, n. g. of Polynoida, with D. pelagica, n. sp., pelagic, from Ceylon; MICHAELSEN (4), pp. 96-99, description and figures.

Dujardinia: from Port Chalmers; PARKER.

Euchone papillosa, recorded from Spitzbergen; v. MARENZELLER (1).

Eulalia vellifera, recorded from Black Sea; PÉRIASLAVZEFF.

Eunice gigantea, recorded from the English Channel; LACAZE-DUTHIERS.

Eusyllis blomstrandi, description and figure; v. MARENZELLER (1) p. 411.

Exogone naidina, with "lateral buds" on Coast of Devonshire; BUCHANAN.

Gipers Lecenture, Schm., Crylon, description and figure; MICHAELER (4). G. tennista, Gr., from Black Sea; PRELASLAVZETT. Blood: Gipers; GRIFFITES (2).

Haplebranchus: on Const of Devendire; BUCHANAN.

Hormation: species parasitic on Echinodorus; MAYEDORF (3) & CUINO Iphiose spinoses. King. from Coylon, remarks on; MICHAELSEN (4), p. 95. Lapidonatus acardiolopis, Gv., from Coylon, remarks on; MICHAELSEN (4, p. 25.

Lescadore escon, Ocean, Kving in the Sponge Microcione; Hornell (2), Leinies curiopate, young specimen (2), from Ceylon; Michaelsen (4 p. 156.

Magalia, n. sp., and Mysta, n. sp., on Devoushire Coast; BUCHANAN, Magalous: larver found on Devoushire Coast; BUCHANAN, BLES.

Festechete, n. g. of Polymide, with N. grimeldii, n. sp., dredged be the "Hirosdelle," pelagic, and with extremely long ventral size v. MARENTELLER (2).

Nematsuarais seculata, Schm., from Black Sea; PERIASLAVERIT.

Scaletys longicarnic, n. sp., with remarks on the genus; Pérlaslavzer Servis longicirus. Schm., Ceylon, described and figured; Michaelses (1 pp. 92-102. Blood of Noveis; Grippipus (2). Dissogony in Novei; Chux. Cell-lineage of Noveis; Wilson.

Series extenses, found again, and on British Coasts; BUCHANAN.

Oplicious ocumenta, from Spitzbergen, description; v. MARENZELLER (
Oplicystrocks, n. sp., or an allied n. g., Coast of Devonshire; BUCHANA

Pectinaria auricoma and belgics, distinguished from one anothe

HORNELL (1).

Phonocia exilia, Gr., Ceylon; MICHAELSEN (4).

Phyllodocs corniculata, Black Sea; PERIASLAVEEFF. P. macrolepido Schm., Ceylon; MICHAELSEN (4).

Piste, n. sp., or (?) an albed n. g.; BUCHANAN.

Poludora cornuta, Black Sea; PERLASLAVZETT.

Polyophthalmus longisstosus and collaris, n. spp., Ceylon; MICHAELSEN(
Polynos haliati, var. hyema, n. var., Liverpool Bay; HORNELL (1).

(Lagisca) extenuata, recorded as British; HORNELL (1).

Petamilla neglecta, recorded from Spitzbergen; v. MARENZELLER (1). Protodrilus leuchartii, Hatschek, on Devonshire Coast; BLES.

Pterceirrus ceylonicus, n. sp., (? = Phyllod. (Eulalia) macroceros, G Ceylon, described and figured; MICHAELSEN (4).

Saleila and Serpula, blood of; GRIFFITHS (2).

Sabella paroina, var. bicoronata, n. var., Liverpool Bay; HORNELL S. saricara, protective device in; WATSON.

Sabellaria bicornis (Hermella bicornis, Schm.); MICHARLSEN (4).

Sabellides, n. sp., Devonshire Coast; BUCHANAN.

Samytha, n. sp., or an allied n. g., BUCHANAN.

Scalibregma longisetosum, from Spitzbergen; v. MARENZELLER (1).

Scolecolepis: species described and figured, from Spitzbergen; v. MAR ZELLER (1).

- Spio mecznikowianus, Clp., (= S. decoratus, Bobr.) and S. ornatus, n. sp., from the Black Sea; Périaslavzeff.
- "Spiochatopterus typicus", of Liverpool Bay, shown to be really Chatopterus insignis; HORNELL (1).
- Staurocephalus chiaji, Clp., rudolphi, D. Ch., and hyalinus, n. sp., with remarks on the genus and figure of the new species; Périaslavzeff. S. rubrovittatus, on Devonshire Coast; GARSTANG.
- Stylaroides iris, n. sp., Ceylon; MICHAELSEN (4), p. 108, description and S. cinctus and horstii, n. spp., Australia, description, anatomy, and figure, HASWELL (1). S. longisetosus, recorded from Spitzbergen; v. MARENZELLER (1).

Terebella: Blood of; GRIFFITHS (2).

- Trypanosyllis striata, n. sp., Black Sea, described and figured. PÉRIAS-LAVZEFF.
- Typosyllis oerstedti, Spitzbergen, described and figured; v. MARENzeller (1).
- Xenosyllides, n. g., with X. violacea, n. sp., Black Sea, with stiff hairs on each ring of the annulate tentacles and cirri; PÉRIASLAVZEFF.

## B. OLIGOCHETA (including Myzostoma),

- HATSCHEK [see above under "General"] would place the Ctenodrilida, including Ælosoma, with the Polychæts. [Their usual position is retained in this Record.] Schmarda's species of Earthworms renamed: BEDDARD (2).
- Rosa (1) proposes to divide the order Terricolae into the following families:---
  - 1. Moniligastridæ.
  - 2. Lumbricidae.


  - 3. Geoscolicidæ.
    4. Megascolicidæ, with subfams.

    II. Cryptodrilinæ.

    III. Acanthodrilinæ.

    IV. Perichætinæ.

GIARD does not consider the mega- or plecto-nephric condition of the nephridia as important in classification. He expresses the relationships between the different members of the Lumbricomorpha in the following table:-



Lumbricomorpha minora.

Acanthodrilus schmardæ, n. sp., BEDDARD (2). A. pictus?, Mich., from Chiloe, Rosa (1). A. multiporus, development; BEDDARD (6). A. spegazzini, Ross, referred to new genus Kerria; BEDDARD (10). A. platurus, n. sp., Chili, MICHAELSEN, (1) p. 226.

Elosoma, encystment of; BEDDARD (1) and VEJDOWSKY (1). E. nireum serve to connect the genus with the Naidomorpha; BEDDARD (10).

- Allolobophora festæ, n. sp., near Tunis, description and affinities: discussion of other species; Rosa (5). A. smaragdina, n. sp., description, Rosa (6). A. complanata, in Tunisia, Beddard (8); 8 species from neighbourhood of Berlin, Collin, A. (1); 3 exotic species, Rosa (1) A. fatida, spiral segmentation in; Morgan: Gerstfeldt's "L. brevispinus" identified with it; Kulagin. A. cantabrica, n. sp. (Rosa) = "A. hibernica, n. sp.," Friend, Ireland and Genoa; Friend (13) "A. cambrica, n. sp.," Friend (4), shown by Hurst & Benham to be a variety of A. chlorotica, Sav. A. subrubicunda, bockii, profuga, and a new species of A. lactea, in Yorkshire; Friend (11) and (12). Especies referred to genus Dendrobæna; Friend (2). Diagnosis of and classification of genus Allolobophora; Friend (1). A. turgida Eisen: 2 varieties in Poland, and 4 other species; Nusbaum (2).
- Allurus tetragonurus, n. sp., Bangor, FRIEND (9). A. tetraedrus, ness Berlin, Collin, A. (1); in Britain, FRIEND (1); in Poland, Nus BAUM (2).
- Amphichæta, n. g., a primitive form of the family Chætogastridæ, with A sannio, n. sp.; KALLSTENIUS.
- Annudrilus, n. g., of family Geoscolecidæ, with A. quadrangulus, n. sp. anatomy; Horst.
- Anteus heterostichon for Schmarda's "Hypogeon heterostichon," and discus sion of genus; BEDDARD (2). A. brunneus (p. 217), appuni (p. 218) callichatus (p. 220), n. spp., Venezuela, and papillifer (p. 214), n. sp. Brazil, MICHAELSEN (1).
- Benhamia: annae, floresiana, and malayana, n. spp. in Malay Archipelago HORST. B. mexicana, n. sp., Mexico, Rosa (1). B. bolavi, Mich., in Venezuela, MICHAELSEN, (1) p. 225. B. itoliensis, n. sp., by Victori Nyanza, MICHAELSEN (2). B. büttneri (p. 259), gracilis (p. 258) inermis (p. 209), pallida (p. 258), togoënsis (p. 260), n. spp., W. Africa MICHAELSEN (1).
- Branchiura, n. g. of branchiate Oligochæt, with B. sowerbyi, n. sp.; Ben DARD (4) and (5), anatomy and histology.
- Bryodrilus, n. g. of Enchytræidæ, with B. ehlersi, n. sp., found at Altensi UDE.
- Caecaria, n. g. of Naidomorpha, intermediate between Stylaria and Printina, with C. brevirostris, rara, and silesiaca, n. spp.; FLOERICKE.
- Clitellio: genus distinct from Limnodrilus; BEDDARD (10).
- Criodrilus lacuum, near Berlin; Collin, A. (1).
- Cryptodrilus insularis, n. sp., Arù Is., ROSA (1). C. dubius, frenchi, gippe landicus, intermedius, lucasi, macedonensis, minor, narrensis, tanjilessi victoria, willsiensis, n. spp., from Victoria, Spencer (3). C. purpureu in Queensland, Spencer (2).

- Dendrobana rubida, Sav., a new variety of, occurring in Russia and Siberia, with a list of the 7 species found there; Kulagin. Tabular view of 6 British species; Friend (2). D. boeckii, in Poland; Nusbaum (2).
- Dero: a species with 6 pairs contractile "lateral hearts" in segments ▼i-xi; Beddard (10).
- Desmogaster, in Sumatra, HORST.
- Diachata littoralis, n. sp., proposed by BEDDARD, (2) pp. 128-130, for part of Schmarda's Pontoscolex arenicola.
- Embolocephalus, n. g. for Sænuris velutina, Gr., and E. plicatus, n. sp.: anatomy and histology of both species; RANDOLPH.
- Budrilus büttneri, n. sp. (p. 256), W. Africa, E. roseus, n. sp. (p. 224), Venezuela, MICHAELSEN (1).
- Geoscolex maximus, for Schmarda's Lumbricus paucisetis; BEDDARD, (2) pp. 119-121. G. maximus, from Brazil, Rosa (1).
- Glyphidrilus, n. g. of Geoscolecidæ, with G. weberi, n. sp., Malay Archipelago; Horst.
- Gordiodrilus, n. g., provisionally belonging to the family Ocnerodrilide, with G. ditheca, elegans, robustus, and tenuis, n. spp., from West Africa, and G. dominicensis, n. sp., from I. of Dominica; BEDDARD (3).
- (Hypogeon heterostichon, Schm.) = Anteus heterostichon, (H. orthostichon, Schm.) = Megascolides orthostichon; BEDDARD (2).
- Kerria, n. g., intermediate between the Acanthodrilidæ and the Ocnerodrilidæ, with K. halophila, n. sp., found in very salt water in the upper reaches of the Pilcomayo R.; BEDDARD (10).
- Kynotus michaelsenii, n. sp., Madagascar, and remarks on the family Geoscolecidæ; Rosa (4). Criticism of the genus, leading to the conclusion that it is closely allied to the genus Microchaeta; Benham (1).

  K. k. lleri, n. sp., Madagascar, Michaelsen, (1) p. 254.
- Limnodrilus, distinguished from genus Clitellio; BEDDARD (10).
- Lumbricus: digestion, Greenwood: circulation, Griffiths (1): blood, Griffiths (2): intra-epidermal blood vessels, Lenhossék (2): individual variation in position of nephridio pores, Hubrecht: spiral metamerism, Cori: nervous system, Retzius, Cerfontaine: origin, course, and termination of the sensory nerve-fibres, Lenhossék (1). (L. fluviventris, Leuck.), referred to L. puter, Hoffm.; (L. triannularis, Gr.) to L. rubellus, Hoffm.; (L. brevispinus, Gerstf.) to Allolobophora fatida: Kulagin. (L. rubescens), n. sp., England, Friend (2), shown by Hurst, and confirmed by Benham (2), to be L. festivus, Sav.; (L. terrestris), Friend (2), shown by Hurst to be L. herculeus, Sav. Hurst gives reasons for discarding the name L. terrestris altogether. L. eiseni, as a rare British species; Friend (3). 3 species of Lumbricus near Berlin; Collin, A. (1). 2 varietics of L. rubellus in Poland, and 2 other species; Nusbaum (2).
- Megascolex templetonianus, n. sp., Ceylon, Rosa (7). Schmarda's "Perichata cingulata" and "P. brachycycla" referred to genus Megascolex.

and his "P. leucocycla" shown to be M. caruleus, Bourne; Beddard (2). M. iris (p. 244), margaritaceus (p. 245), n. spp., Samar, pictus, n. sp. (p. 246), Borneo; MICHAELSEN (1).

Megascolides australis, development of nephridia in; Vejdowsky (2).

M. attenuatus, cameroni, hulmei, incertus, insignis, manni (with variabilis, n. var.), obscurus, rossus, sinosus, victoriensis, n. spp., Victoria, Spencer (3). M. orthostichon, for "Hypogocon orthostichon," Schmarda; Beddard (2).

Microchæta belli, Cape Colony, and M. papillata, Natal, n. spp., anatomy; Benham (1). M. belli is shown to connect Michaelsen's genus Kerria with Microchæta. M. benhami, n. sp., locality unknown, Rosa, (1) p. 382. M. rappii, anatomy, Rosa (1).

Microscolex algieriensis (pp. 29-32), Algiers, and poultoni, Madeira, n. spp.: anatomy, with discussion of the 2 species; BEDDARD (8).

Moniligaster japonicus, n. sp., Japan, MICHAELSEN, (1) p. 232.

Myzostoma pulvinar and alatum, v. Graff, parasitic on Antedon phalumgium, amendments to v. Graff's description and discussion of the sexes; Prouho.

Nais greef, n. sp., intermediate between N. elinguis and N. barbata FLOERICKE. N. parasitica in Vivipara unicolor in Egypt; SONSINO (4) Neodrilus monocystis, Bed., extended description of; BENHAM (2). Ophidonais reckii, n. sp., FLOERICKE.

Paradrilus rosae, anatomy, Rosa, (1) p. 393.

Perichæta sumatrana, Horst, P. dyeri, sinensis, bermudensis, taprobana morrisi, barbadensis, hesperidum, mauritiana, n. spp., Beddard (9) and discussion of external and internal characters which are import ant in determining species; P. carinensis, bournei, campanuluta peguana, n. spp., from Burmah, Rosa (2); P. rokuko, n. sp., and ? other new species from Japan, BEDDARD (11), all with no setse or clitellar segments, and atria rudimentary. P. aeliana, fasciata, and enganensis, n. spp., with new subspecies P. enganensis tetra, Engano description and anatomy; Rosa (3). P. dubia (Sumatra), and P. minima (Java), n. spp., Horst. P. hawayana (Hawaii), phillip ina (Cebu), operculata (Java), racemosa (Java and Borneo), penta cystis (Mahé), ijimæ, Japan, n. spp., and P. grubei and P. novara n. spp., Tahiti, to replace P. taitensis, which is now destroyed. F pulchra, n. sp. (p. 233), Luzon, hilgendorfi, n. sp. (p. 235), Japan albida, n. sp. (p. 237), Marquesas, longa, n. sp. (p. 239), Sumatra udekemi, Gr. (p. 240), Java, manthorensis, n. sp. (p. 241), Bornec martensi, n. sp. (p. 242), Banka, divergens, n. sp. (p. 243), Japan and P. neoguinensis, n. sp. (p. 229), New Guinea; P. monilicystic n. sp. (from a hothouse in Botanical Gardens, Berlin), P. pallida (p. 227), Brazil, n. sp., MICHAELSEN (1). MICHAELSEN (1) furthe shows (p. 212) that his P. heterochæta from the Azores is only synonym of the widely-spread P. indica. P. gympiana, n. sp., Queens land, Spencer (2). Beddard (2) shows that of Schmarda's Peri chata, only one (P. viridis) really belongs to that genus; of the

- others, P. leucocycla, P. cinguluta, and P. brachycycla, belong to the genus Megascolex; he makes a new species, P. vitiensis, for one of Schmarda's specimens labelled, "Hypogeon sp., aff. H. orthostichon."
- Perionyx excavatus, E. Perrier; Rosa (1) and (2). P. arboricola, n. sp., Burmah; Rosa (2): P. violaceus, n. sp., Java, Sumatra; Horst.
- Photodrilus phosphoreus, geographical distribution and position of; Giard and Matzdorf (1).
- Phreoryctes menkeanus, two new localities in the N. E. of France;
  BRUNOTTE.
- Plagiochata, n. g. of Acanthodrilida with P. punctata, n. sp.; BENHAM (2), anatomy and histology.
- Pleionogaster, n. g. of Perichætidæ, with P. jagori and samariensis, n. spp., Phillipine Is., MICHAELSEN, (1) pp. 247-249.
- Plutellus perrieri, n. sp., Qu. Charlotte's I., BENHAM (1).
- Pristina longiseta, notes on; BEDDARD (10): he concludes that Pristina and Naidonina are no longer to be regarded as distinct species.
- Pontodrilus arenae, Fr. Müll., Brazil; MICHAELSEN, (1) p. 222.
- Pontoscolex arenicola, Schm.; BEDDARD (2) retains this name for some of Schmarda's specimens, referring the others to the genus Diachata, proposing for them D. littoralis, n. sp.
- Pygmæodrilus bukobensis and affinis, n. spp., near Victoria Nyanza, Michaelsen (2).
- Rhinodrilus ecuadorensis, n. sp., from Ecuador at a height of 14,000 feet, Benham (4).
- Siphonogaster fertilizing the valley of the Niger; Loz. S. emini and stuhlmanni, n. spp., and discussion of genus; MICHAELSEN (2).
- Sanuris velutina, referred to new genus Embolocephalus; RANDOLPH.
- Sparganophilus, n. g. of aquatic Oligochest belonging to the family Bhinodrilide, with S. tamesis, n. sp., Goring on Thames: anatomy and histology, and discussion of how conveyed to the above locality; Benham (3).
- Tykonus, n. g., allied to Anteus, with T. grandis, n. sp., Brazil, MICHAEL-SEN, (1) p. 212.
- Urochata corethrura, at Engano; Rosa (3). U. sp. ?, at Malacca; Rosa (1). Abolition of genus; Beddard, (2) p. 126.
- Vermiculus, n. g. of aquatic Oligochæt, with a dense covering of fine hairs or bristles, median atrium of sperm-ducts, and median spermathecal pores. V. pilosus, n. sp., Weymouth Coast; GOODRICH.

#### 3. HIRUDINEA.

BLANCHARD (4) would divide the Hirudinids into the HIRUDININA or fresh-water forms (including Hirudo, Hamopis, Limnatis, Hirudinaria, Macrobdella and Whitmania) and the HEMADIPSINA or terrestrial forms (including Hamadipsa, Moquina and Xerobdella). He discusses (6) the relations between the Nephelids, Nephelis, Dina, and Trocheta.

- Aulastoma: a palmarotic genus; STOLL: minute structure of eyes; MAIER: salivary glands; LEUCKART (2). A. gulo, in Poland, 6 new varieties: maculata, sinuata, grubei, taniata, lithuanica, umbrina; NUSBAUM (2).
- Branchellion punctatum, Baird, and distinction of the 2 other Australian species; BLANCHARD (3, VI). A specimen described by MACDONALD (Tr. L. S. (2) i, 1876, p. 209), from Myliobates, is referred to this species. B. rajæ, ? n. sp., occurring abundantly on Raja nasuta, off New Zealand; Parker.
- Clepsins: a universal genus, Stoll: minute structure of eyes, Maies: salivary glands, Leuckart (2): nephridia, Bolsius (2). C. tesselata, var. marmorata, n. var., Poland, Nusbaum (2): in the posterior nares of Cygnus atratus, and in beak of Haliatus albicella, Collin, A. (2). C. holleneis, n. sp., and its metamerism, Whitman. C. polonica, n. sp., Poland, Nusbaum (2).
- Cylicobdella: confined to tropical new world (S. Brazil and Guatemala);
  Stoll.
- Glossiphenia, splitting up of genus foretold; BLANCHARD (3, IV). G. biocculata and catenigera, geographical distribution: G. marginata, description and distribution; BLANCHARD (3, IV). G. sexoculata, description, BLANCHARD (3, V), G. tesselata, description, synonomy, &c., BLANCHARD (1): in Chili, and its mode of transport thence, with additions to the description of the species, BLANCHARD (2).
- Hæmopis: blood; GRIFFITHS (2). H. sanguisuga, for Diesing's Typhlobdella; BLANCHARD (3, 11).
- Hirudo: minute structure of eyes, MAIER: alimentary canal, HENKING: salivary glands, LEUCKART (2): blood, GRIFFITHS (2). H. sanguisuga in the nasal mucous membrane of man; Condorelli (4). H. medicinalis, 3 varieties in Poland, and 2 intermediate forms between them; Nusbaum (2). Nephridia, Bolsius (2).
- Nephelis: minute structure of eyes; MAIER. N. atomaria, Carena, and meyeri, n. var. of it, description, notes on N. occulata, Bergm., and short description of N. gallica and N. tergestina, n. spp., BLANCHARD (3, III.). N. octocculata, in Poland, and its varieties there, according to locality; Nusbaum (2): 2 new varieties, and 4 new sub. varieties. N. vulgaris, nephridia; Bolsius (2).
- Piscicola: minute structure of eyes, MAIER: salivary glands, LEUCK-ART (2).
- Trocheta subviridis, in Liguria, description and geographical distribution; BLANCHARD (6), and relation of genus to other Nephelids.
- Typhlobdella kovátsi, Diesing, shown to be only a Hæmopis sanguisuga from examination of type specimen; BLANCHARD (3, 111).
- Xerobdella lecomtei, description; BLANCHARD (4). It is shown to be a Hirudinid and not a Nephelid, nearest allied to Hamadipsa, especially to H. japonica, Whitman. Only 4 pairs of eyes to this, the only species known; 2 orifices in female somite, and an infundibuliform pore on 95th ring, with no homologue in other Leeches.

### III.—PLATYHELMINTHES.

#### A.—BIOLOGICAL.

#### 1. GENERAL.

The 4th vol. of Bronn's "Klassen und Ordnungen des Thier-reichs," continued in 1892 by MAX BRAUN (4), gives a very full and up to date account of the digenetic Trematodes.

Bibliography of Platyhelminthes; PARONA (1). Parasitism; Looss (2). Anatomy and histology of Cestodes of freshwater Fish; KRAEMER (2): these are distinguished from those of warm-blooded Vertebrates by the absence of the rostellum, this being often replaced by a 5th small sucker at the front end of the scolex: excretory and reproductive systems also distinctive. Anatomy and histology of Pelagic Polyclads, v. GRAFF (1); of Triclads, CHICHKOFF, HALLEZ (3). Monograph of genus Microcotyle (11 species recognized); PARONA & PERUGIA (1). Anatomy of genus Notocotyle (only 2 species, N. alveatum, Mehlis, and N. verrucosum, Fröhl., recognized); MONTICELLI (3). Anatomy and histology of the Monocotylides, SAINT REMY (2); of Cotylogaster michaelis, n. g. & sp., MONTICELLI (6); of Monostoma, MONTICELLI (4) and (5); of Thysanocephalum, LINTON (5); of Stenostoma leucops, OTT; of Stychoplana sargassicola, v. GRAFF (4); of Temnocephala madagascariens, n. sp., with 12 cephalic digitations, and enumeration and discussion of other species, VAYSSIÈRE (1); of Actinodactylus, n.g. & sp., with 12 tentacles along the lateral margin of the body radiating from it, with ventral suckers as in Temnocephala, and with no eyes, HASWELL (4, III); of Geonemertes australiensis, n. sp., DENDY (1); of Polycharus caudatus (n. g. & sp. of Accelous Turbellarian, fam. Aphanostomida), MARK. Anatomy of Cotyloplana, n. g. of Land Planarian, with suckers on ventral surface close to anterior extremity; SPENCER (1). On a freshwater Alloioccel Planarian, found in two deep wells 100 miles apart, with no pigment and no eyes; HASWELL (4, 11). Anatomy and development of Amphistomum subclavatum; Looss (1). Account of freshwater Nemertines; VAILLANT, and DE GUERNE (2).

## 2. EPIDERMIS.

ZOGRAF (1) and MONTICELLI (8) both show that there is a true, though much modified, ectoderm present in the adult Cestode, which produces the cuticular layer. Zograf shows, further, that this is the remnant of the ectoderm of the 6-hooked embryo which is not thrown off, but only vacuolised when the ciliated envelope is shed. Brandes (1) maintains that there is no true ectoderm in adult Trematodes, except perhaps in Temnocephala; and that the cuticle is formed by an outer glandular layer of parenchyme cells, which he terms "Ectoparenchyme." Brann (3)

criticizes this statement, and shows that, in one Monostomum at least, the so-called "cuticle" is itself a metamorphosed ectoderm. HASWELL (5) shows that the ectoderm is ciliated even in the adult of certain species of *Temnocephala*. Large, oval, refractive concretions are described in the ectoderm of a freshwater Nemertine by DU PLESSIS [see also VAILLANT and DE GUERNE (2)]. Ectoderm of Triclades; CHICHKOFF.

#### 3. MUSCULAR SYSTEM.

Insertion of muscles of parenchyme in Monostomum proteus, Braudes; WALTER.

#### 4. ALIMENTARY CANAL.

Protoplasmic processes of the alimentary epithelium, by which food is taken in, in *Stenostoma*; Ott. Anterior median branch of alimentary canal over the brain absent in 2 species of *Planocera*; v. Graff (1). Multiplicity of pharynx in a Triclad (*Phagocyta*); anastomosis of the two recurrent branches of the alimentary canal in *Dendrocalum*; Hallez (2).

#### 5. CIRCULATORY SYSTEM.

Hæmoglobin in the body liquid of a Turbellarian (Syndesmis echimorum), Cuénot. Vascular system only a specialized part of the excretory system in Geonemertes; DENDY (1). Distinct from the excretory system in Enopla; Bürger (1).

#### 6. Excretory System.

KRAEMER shows that in the Cestodes of Freshwater Fish, the longitudinal vessels of the excretory system all open into a terminal vesicle in the last joint, and by its means open to the exterior posteriorly; anteriorly they communicate with the exterior by means of fine canaliculi, which branch off from a capillary vascular plexus, and lie in the neck and youngest segments. RIEHM shows that in the Cestode Schistocephalus dimorphus there is a pair of apertures to the excretory system in each segment; the whole system branches all over the body, and is continuous from segment to segment; he compares the arrangement to that of the trachese and stigmata of Insects. Walter describes a new organ lying round the excretory pore, and opening into the end part of the efferent duct, just in front of the pore, in the genus Monostomum. HASWELL (2) shows that the excretory system of Temnocephala is intracellular, lying in a few very long and sometimes branching cells; connected with each main duct there is a much-branching system of canaliculi, containing a number of ciliated flames in the substance of the perforated cell which forms the

thick protoplasmic walls of a contractile terminal sac on each side; each sac opens to the exterior anteriorly and dorsally by pores in the same position as in many other Trematodes. DENDY (1) shows that the excretory system of his new Land Nemertine, Geonemertes australiensis, is also intracellular, and that the tubules end in flame cells : he did not succeed in finding apertures to the exterior. BÜRGER (1) shows that in the Enopla the epithelium lining the excretory canals is ciliated, and that the canals end in hollow blind swellings, also one cell layered, with a flame of cilia projecting into the lumen; the vascular system, though close to the excretory system, is not in communication with it. Voigt (2) shows that in Mesostomum truncatum the efferent excretory ducts, instead of opening into the pharynx, open independently on the ventral surface; he would therefore make this species into a new genus. Sekera, however, shows that in other species of Mesostomum, e. g., M. rostratum, the excretory pores are also independent of the pharynx. He also shows that the genus Bothrioplana has excretory organs of the Mesostomid type.

# 7. Organs of Uncertain Function and not in connection with other Systems.

Sommer's "Plasmatic vessels," rediscovered and described in *Tania* saginata and *T. solium*; BLOCHMANN (1). Curious bodies in the anterior part of the segments of *T. expansa* = "interproglottidal" glands; STILES (8). [See also under "Excretory System;" WALTER.]

Unicellular glands of Temnocephala ducts shown to branch in the pro-

toplasm ; HASWELL (2).

### 8. NERVOUS SYSTEM AND SENSE ORGANS.

Complete decentralisation of nervous system in Planocera grubei, n. sp., and approach to it in P. simrothi, n. sp.; v. Graff (1).

Nervous system of *Distomas*; CRETY: suckers shown to be especially well supplied with nerves, and therefore probably to serve as tactile organs. Eyes shown to multiply by division in a Land Nemertine; DENDY (1).

#### 9. GENITAL SYSTEM.

Vitelline nuclei are shown to exist in the ova of Distomum richardii by Monticelli (2), and by Crety (2). Monticelli also finds them in D. veliporum; he concludes that they are produced by the cytoplasm, and probably serve as a centre for the formation of yolk, having no intimate connection with the nucleus, and playing no part in the process of fertilization. Crety compares them to the diffuse granules in the ova of D. megastomum; he regards them as formed endogenously within the cell, and afterwards fusing with its protoplasm; he compares them with the similar structures in the ova of other animals. Spermatogenesis in Trematodes is dealt with by Monticelli (1).

SAINT REMY (3) describes the genital apparatus of the *Tristomida* (including under that head the *Tristomida* proper, the *Monocotylida*, and the *Udonellida*). He suggests that the canal of Laurer of the *Digenia* may be homologized with the vitello-intestinal canal of the *Polystomida*; the vagina, with which it is usually homologized, is in Tristomids developed as a female copulatory organ, either single or paired, but probably originally paired. Braun (3) describes a genital atrium in an *Amphistomum*, and suggests that the large sac present in the genus *Gastrothylax* is an enlargement of such an atrium, and serves as a brood-pouch.

[Abnormalities in the genital apparatus of Distomum hepaticum, see under "Variations."]

The female genital apparatus of Amphilina and Caryophylleus is explained by Monticelli (9). Kraemer (1) and (2) shows that in the Twento of freshwater fishes the female genital aperture is in front of the male; on the female duct are a number of loops, serving as receptacules seminales; the position and form of the yolk-glands is different to what it is in Twentas of warm-blooded Vertebrates. Filippi describes the formation of the ovigerous capsule in T. botriophilis; the eggs appear to be fertilised in the ovary in this species, as there is no uterus or seminal vesicle; he thinks it probable that the proglottids are self-fertilised. Diamare describes the genital apparatus of Dipylidium.

v. Graff (1) confirms Lang's view as to the origin of the gonads from the intestinal epithelium in pelagic Polyclads; he also shows that the uterus is formed from the ectoderm in one species at least; that the accessory vesicle of the 2 genital duct is a seminal vesicle; that there is a female as well as a male copulatory arrangement in two species; that the relative positions of oviduct and shell-gland are different to what they are in non-pelagic Polyclads. The uterus of the Triclads is dealt with fully by Bergendal (8) and (5).

## 10. Reproduction, Development, and Life-History.

Reproduction by transverse fission is described in *Bipalium kewense* by Bergendal (7). Voigt (1) questions the viviparity of *Planaria alpina*, maintained by Zschokke.

Development of Cestodes and Trematodes compared; SAINT RÉMY (5).

GRASSI & ROVELLI discuss fully the embryology and life-history of the Cestodes, comparing them with one another and with other Platyhelminthes; they find that the cysticercoid in Tania elliptica has a simple and direct development; 1 species, T. nana s. murina, has no intermediate host, and thinks that all Tænias originally had but one host. The cysticerci of 8 species of Tania are dealt with by v. Linstow (3). Raillier & Lucet (1) point out that Davainea proglottina must have some other intermediate host than Limax: Giard & Billet that Distomum hepaticum must have some other host than Limasus truncatulus, in Tonkin: Magalhäes that Tania cuneata must have some other intermediate host than Allolobophora fatida in Brasil. Moniez (1, xi)

describes a curious larval appendage, over a metre in length, apparently forming a network in the tissue of its host when protruded, in *Gymnorhynchus reptans*; he compares it to an appendage found on several Cestodes of the type of *T. serrata*.

The cercaria of Amphistomum subclavatum is dealt with by Looss (1), and LANG. LOOSS deals with the whole development, showing how the redia is directly derived from a single organ of the embryo, and the cercaria in the same way from the redia; he regards the development as a metamorphosis, no pædogenesis coming in, and the germinal epithelium of the adult being formed from the elements not used in the development of organs of the embryo, and producing not only the genital organs of the redia and cercaria, but also, in an intermediate stage, an epithelial lining for the body cavity. LANG shows that the cercaria, escaping from Planorbis contortus, becomes first encysted on the skin of the young frog, and that only when this skin is shed and eaten by the frog does the animal reach its final habitat and development continue. The embryo of Gynecophorus hamatobius is dealt with by RAILLIET (3), and by CAHIER; of Gyrodactylus elegans and 5 species of Distoma, by von Linstow (3). The development of Distomum caudatum, by BLOCHMANN (2); a cyst phase of development in D. baraldii, n. sp., by Sonsino (2). history of Distomum hepaticum; LUTZ: it is shown that Limnaus pereger (= L. calmensis) may be an intermediate host.

HALLEZ (1) shows that all the Turbellaria diploblastica (Rhabdoccels and Triclads) have a provisional terminal larval pharynx with no sheath; and that the position of the adult pharynx is determined, after the flattening of the larva on its oral surface, by the relative amount of growth of an anterior and a posterior hemisphere; where the pharynx is posterior (e.g., Triclads and Monotus fuscus) in the adult, stages are gone through in which it is first anterior and then median; the flattening of the body accompanies the backward movement of the pharynx, i.e., the greater amount of movement of the cephalic as compared with the caudal hemisphere.

### 11. VARIATIONS.

CONDORELLI (1) gives examples of different kinds of anomalies in Cestodes, referring especially to a *Tania solium* in which the hooks were absent and replaced by twelve large papillse resembling those of some Nematodes. RAILLIET (2, 1) describes a specimen of *Cysticercus pisiformis* with as many as six suckers. RAILLIET & LUCET (1) mention the variation in number of segments in *Davainea proglottina*. RAILLIET (2, III) records instances of fenestration of segments in *Dipylidium canimum*. WALTER mentions variations in the development of *Monostomum reticulare* from the embryo.

Asymmetry of organs is mentioned by STILES (8) in *Tania giardi*. SPENGEL describes one specimen of *Distomum hepaticum* with no male genital organs, and another with the anterior half of the right vitellarium absent and the duct on the left side only.

## 12. CLASSIFICATION, &c.

Classification of Cestodes, Grassi & Rovelli, Villot (1), Monticel (10). Monticelli (4) proposes the name "Cestodaria," for the simp unjointed monogenetic Cestodes, Gyrocotyle, Amphilina, Caryophylla and Archigetes. Current ideas on the relations between Cestodes a Trematodes: Saint Rent (5). Railliet (2, VII) proposes a classification for Bird Tsenias, dividing them into 4 genera. Stiles (9) d crosses Blanchard's genus Moniezia.

MONTICELLI (6) proposes a new classification of Trematodes, and d cusses the family Aspidebothridæ. The affinities and arrangement of t Monostomidæ is discussed by Monticelli (5), (4), and (3); and Brandes (2). Brandes would exclude the genus Didymozoon from t family; he refers several so-called species of Monostomum to oth genera. The position of Temnocephala is dealt with by Haswell (and by Vayssière (1).

The Nemertines of the Gulf of Naples are classified, and a genealogi tree is given to show their relationships, by BÜRGER (2).

A new classification of Turbellarians is proposed by Hallez (3), and further one of the Triclads, (3) and (4). He discusses (3) the affinit of the Triclads with the Alloicocels and the laws of morphogeny of t Turbellarians generally. Bergendal (1) classifies and discusses t affinities of the Swedish Triclads. v. Graff shows that Weldon's Helodiscus piliger is an accelous Turbellarian.

[For further details, see Systematic part of Record.]

## 13. DISTRIBUTION OF PARASITES ACCORDING TO THEIR HOST:

#### a. General.

Distribution of Cestodes and Trematodes, according to various hos STOSSICH (4); Trematodes and Cestodes of Mecklenburg and their hos BRAUN (5).

#### b. Man.

Tania diminuta, Cysticercus cellullosa, Distomum lanceolatus ZSCHOKKE (2). Echinococcus; SANGALLI, SCHULTÉN, & HOMEN (liver and mesentery), KINYOUN (in kidneys, liver, and bladder); mul locular form in bladder, HUBER (2): in brain, MUDD: in liv MANGOLD. Cysticercus, living in lens of eye; GRAEFE. Synopsis exotic Entozoa of Man; SONSINO (1). Bibliography of hum Cestodes; HUBER (1). Parasites from Man to Animals, and vice vers RAILLIET (1).

#### c. Other Mammals.

Monograph of Distomids in Mammals; Stossich (1). In Domest Animals; Neumann, Dewitz, Curtice. In Cattle and Shee Temias; Stiles (3), Echinococcus multilocularis; Stiles (4). In an

gland of Dog, Dipylidium caninum; RAILLIET (2, IV). In Cat, Dipylidium trinchesi, n. sp., DIAMARE. In Kid and Antelope, Cysticercus tenuicollis; RAILLIET (2, V & VI). In Hyrax, a Tenia; Setti. In Stenorhynchus (Ogmorhinus) leptonyx, Bothriocephalus quadratus, n. sp., v. Linstow (4). In Rabbit, Anoplocephala caniculi; RAILLIET (2, V & VI). In Dolphin, Tania forsteri (= Prosthecocotyle, n. g.); Monticelli (7). In Seal (Cystophora) Bothriocephalus tectus, n. sp., Linstow (4). In Balænoptera, Distomum goliath; Lönnberg (1). In Hedgehog, D. caudatum and Trichosoma exiguum; Blochmann (2).

## d. Birds.

Monograph of Distomids in Birds; STOSSICH (2). Tænias of Birds, notes on; LINTON (1), V. LINSTOW (2): classification of; RAILLIET (2, VII). Revision of Monostomids parasitic in *Grallidæ* and Palmipeds; MONTICELLI (5): of *Notocotyle* in ditto; MONTICELLI (3).

In Duck, Echinocotyle rosseteri, n. g. & sp., with 10 hooks and armed suckers, Rosseter. In Fowl, Davainea proglottina; RAILLIET & LUCET (1). In Pigeon, Tenia delafondi; RAILLIET (5).

## e. Snakes.

Distomum subflavum and baraldii, n. spp., in Zamenis viridiflavus; Sonsino (2).

## f. Fish.

Parasites of Freshwater Fish; ZSCHOKKE (3). Anatomy and histology of Tæniæ of Freshwater Fish; KRAEMER. Fish in which genus Microcotyle found; PARONA & PERUGIA (1). Lamna corumbica with a n. g. & sp. of Cestode, Ceratobothrium xanthocephalum; Monticelli (5). Scorpæna porcus with Bothriocephalus caudatus; MARCIALIS. Coregonus muræna with cysts of a Distomum in wall of heart; ZACHARIAS. Box salpa with 2 species of Monostomum, of which one new; Monticelli (5). Trematodes ectoparasitic on different fish; PARONA & PERUGIA (2). Monogenetic Trematodes of Roscoff, and the fish on which they are found; Saint Rémy (4). Transitory parasitism of a Triclad Turbellarian on the larva of a fish; Vogt.

## g. Molluscs.

Cercariæ in (chiefly) Gasteropods of Egyptian wells; Sonsino (4). A new Cercaria in Limnæus truncatulus, v. Linstow (3): in Donax anatinum (C. pectinata), Huet. Helix hortensis, the intermediate host for Distomum caudatum; Blochmann (2). Limnæus pereger (= L. calmensis), an intermediate host for D. hepaticum; Lutz. Lutz also gives interesting details with regard to the effect of the parasites on the snails, the mortality of the latter, &c.

#### h. Orustaceans.

In Crayfish, a Distomum (? D. nodulosum); LINTON (2). In Astacoides, a n. sp. of Temnocephala; VAYSSIÈRE (1). In branchial cavity of a land Crayfish, Actinodactylus, n. g. of Temnocephaleæ; HASWELL (4). In Cyclops brevicaudatus and crassicornis, the cysticerci of Tenia setigera of the Goose and T. brachycephalu of Machetes, respectively; v. LINSTOW (2). In a Freshwater Calanid (Eurytemora lacinulata) a Cysticercoid resembling the one found by Mrazek in Cyclops agilis (Zool. Rec. 1891, Vermes, p. 19); RICHARD. In Cypris cinerea, the Cysticercus, with 10 hooks and armed suckers, of Echinocotyle rosseteri, n. g. & sp.: ROSSETER.

#### i. Echinoderms.

On Synapta, cysts of Cercaria of Distomum leptosomum. In Ophiurids, Cercaria capriciosa, n. sp. ?; MATZDORF (3), CUÉNOT. On Echiaus, Syndesmis echinorum; CUÉNOT.

## j. Plants.

Cestodes and Trematodes in; Bos (1).

## 14. STRAY MEDICAL NOTES.

Infectious diseases of animals which can be transmitted to Man, and vice versa; Perroncito (Tania mediocanellata chiefly dealt with in this respect); also Railliet (1). Study of T. litterata; Condorelli (5). Prismatic variety of T. saginata; Coats. Increase of Tanias in France during the last half century; Béranger-Ferand (1). Relative frequence of different species of Tania; Collin, G. Bothriocephalus latus in Holland; Cathe. Cysticercus in eye and its removal; Graefe, Seyfert. Morbid phenomena caused by Bilharzia hamatobia; Collorid. Retention of toxines in intestine, and consequent poisoning, by intestinal worms; Voncken. Cases of Echinococcus; Puky. Tania disease in Man; Béranger-Ferand (3).

[See also under 13, b. Man.]

## 15. Methods, &c.

Experiments for keeping Cestodes alive; Lönnberg (2). Methods of preparing Cestodes for histological work; Linton (5), Grassi & Rovelli: of injecting excretory system in Cestodes; Riehm. Dendy (1) recommends the use of chloroform vapour before immersion in spirit for fixing Land Nemertines. Chichroff suggests a new fixing reagent for Turbellarians, similar to Lang's, but with different proportions of the constituents, and with nitric acid added; he finds that he can fix specimens, well extended, in this way. Joubin recommends quick plunging into boiling water for killing Nemertines; Tullberg, paralyzing with MgCl<sub>2</sub> and killing with chromic acid, for Nemertines and Planarians; see also Cunningham.

### B.—FAUNISTIC.

## 1. PARASITIC FORMS.

Geographical distribution of exotic Entozoa of Man; Sonsino: of Tanias of Man; Beranger-Ferand (2).

## a. Palmarctic Region.

Sweden: distribution of Bothriocephalus latus in ; LÖNNBERG (3).

West Europe: distribution of *Tænia saginata* in; BLANCHARD (5). Large American Distomum in Europe; LEUCKART (1).

France: decrease of *T. solium* and increase of *T. mediocanellata* during the last half century, BÉRANGER-FERAND (1). Bois de Boulogne: Cysticercoid in a Freshwater Calanid; RICHARD. Roscoff: Monogenetic Trematodes; SAINT RÉMY (4), including 1 new species on the Sole.

Mecklenburg: Parasitic Worms, 6 genera and species of monogenetic Trematodes, 6 genera and 31 species of digenetic Trematodes, 9 genera and 30 species of Cestodes; Braun (5).

S. Germany, Switzerland, and Austria: Restriction of multilocular Echinococcus to; MANGOLD: in Würtemberg; VIERORDT.

Sardinia: T. solium, T. canurus, and Bothriocephalus angustatus;
MARCIALIS.

Tunis: Bilharzia in; CAHIER, VILLENEUVE.

Egypt: parasites of freshwater Molluscs; Sonsino (4).

## b. Ethiopian Region.

Madagascar: Temnocephala madagascariensis, n. sp., in an Astacoides; geographical distribution of genus discussed; VAYSSIÈRE.

## c. Oriental Region.

Tonkin: Trematode parasites of oxen, Homalogaster poirieri, n. sp., Distomum calomaticum, n. sp., and D. hepaticum, whose intermediate host there is unknown; Giard & Billet.

## d. Australian Region.

Gippsland: Actinodactylus, a new genus of Temnocephalidæ; HASWELL (4, III). Distribution of genus Temnocephala; VAYSSIÈRE.

## e. Neotropical Region.

South Georgia: Bothriocephalus quadratus and tectus, n. spp., and a species of Tunia; V. LINSTOW (4).

Brazil: Tienia cuneata: intermediate host in this region unknown; MAGALHÄES.

## f. Nearctic Region.

Texas: (Distomum texanicum), n. sp., Francis, shown to = F. am canum, Hassal, wh. ? = D. magnum, Bassi; Stiles (6). Distribution large American Distomum; Leuckart.

## 2. FREE-LIVING LAND AND FRESHWATER FORMS.

Geographical distribution of Land Planarians; STOLL.

## a. Palæarctic Region.

Geographical distribution of Freshwater Nemertines; Guerne Rhynchodesmus in Europe; Stoll.

England: Freshwater Nemertine (in the Cherwell); Benham (5). S. Germany: distribution of *Planaria alpina* and *gonoceph* Voigt (3).

Bohemia: Freshwater Turbellarians; KAFKA, FRITSCH & VAVRA.

Poland: 11 species of Rhabdoccels; Nusbaum (2): 7 of the same
6 more species of Rhabdoccels; Wassilieff: 1 species of Vortex, ne
Russia: 7 species of Turbellarians near Moscow not found there bei
ZYKOFF: only one Dendroccel (a Polycelis) there. Solovetzki: Ne

tines; Knipovicha, p. 25. Finland: a Rhabdoccel; Levander (2).

Sweden: Triclads enumerated, and several species discussed, inclu

Polypostia similis, n. g. & sp.; Bergendal (1). Spain: 2 Turbellarians from Santander; FUSSET.

E. Asia : Bipalium ; STOLL.

## b. Ethiopian Region.

Madagascar: Bipalium; STOLL. Temnocephala [see above, u "Parasitic Forms"].

## c. Oriental Region.

Ceylon: Rhynchodesmus; STOLL.

## d. Australian Region.

Distribution of Geonemertes australiensis, n. sp.; DENDY (1).

Queensland: Geoplana regina, n. sp., G. cærulea, Bipalium kew Spencer (2), Dendy (2). Dendy also records 2 other species of Geopland 1 of Rhynchodesmus.

S. Australia: Geoplana fletcheri, and a new variety of it, var. adelaid from Adelaide; Dendy (3).

Tasmania: 4 species of Geoplana, not known to occur there be one of which new; DENDY (3).

New Zealand: in the underground waters near Canterbury, an a occel Turbellarian.

Lord Howe Island: Geoplana absent, and in its place a new genus of Land Planarians, Cotyloplana, with new species, whiteleggi and punctata; 6 new species of Rhynchodesmus; SPENCER (1).

Fiji Isles, Friendly Isles, and Samoa: as original habitat of Bipalium kewense; Bell.

## e. Neotropical Region.

S. America: Bipalium kewense; Collin, A. (2).

Chili, Brazil, Guatemala: Polycladus; Brazil: Geoplana; STOLL.

Jamaica: Freshwater Planarians and Nemertines, 1 Land Planarian; Andrews (1).

## f. Nearctic Region.

Rhynchodesmus in N. America; STOLL.

### 3. MARINE FORMS.

Geographical distribution of Pelagic Polyclads; v. GRAFF (1).

E. Baltic: 17 Turbellarians, 8 Nemertines recorded from, and the problems for further exploration of the fauna dealt with; KOJEVNIKOV.

English Channel: Carinella polymorpha and Cerebratulus aurantiacus on the Coast of Devonshire; GARSTANG (1).

Mediterranean: Nemertines of Gulf of Naples, BÜRGER (2), 65 species of which 28 are new; 1 new genus, Hubrechtia.

Atlantic: Planocera simrothi (between I. of Ascension and Equator) and P. grubei, n. spp.; v. Graff (1). Sargossa Sea: Stylchoplana sargossicola; v. Graff (4) and (1).

S. Indian Ocean: Planocera grubei, n. sp.; v. GRAFF (1).

Pacific Ocean.—Off New Guinea: Stylchoplana sargossicola; v. Graff (4) and (1): Planctoplana challengeri, n. g. & sp.; v. Graff (1). Off Noumea: Eupolia brockii, Cerebratulus caledonicus, bicornis, anas, n. spp., Eunemertes francisca, n. sp.; Joubin & François.

#### C.—SYSTEMATIC.

#### 1. CESTODA.

Affinities with other Platyhelminths discussed by Grassi & Rovelli, who also discuss affinities of Cestodes amongst themselves, criticizing Villot's classification of the cystic forms into Cysticeric, Cysticericidea, and Pseudocystici. Villot maintains his classification.

Monticelli (10) proposes the following classification:—

#### Order CESTODA.

#### I. Subord. ATOMIOSOMA.

Fam. 1. Diplocotylidæ, Montic. Fam. 2. Tricuspidaridæ, Montic. IL Subord. Toxiosoma.

Tr. A. Atrypanorhynchu.

Monossichionia-Fam. 3. Cyathobothride, Montic.

Dissichionia - { Fam. 4. Pseudobothridæ, Montic. Fam. 5. Dibothridæ, Dies. Tetrassichionia - { Fam. 6. Tetrabothridæ, Dies. Fam. 7. Tetracotylidæ, Dies.

Tr. P. Prypanorhyncha.

Fam. 8. Tetrarhynchida.

MONTICELLI (9) proposes name Cestodaria for the genera Gyrocoty Amphilina, Caryophyllaus, and Archigetes, intermediate betwe Trematodes and Cestodes.

### Adult Forms.

Amphilina referred to suborder Cestoduria, and A. liquolidea, anatom Monticelli (9).

Anoplocephala cuniculi: bronze coloured specimen in rabbit; RAILLI (2, 11).

Archigetes referred to suborder Cestodaria; MONTICELLI (9).

Bothrimonus, Duv., synonymy of; Monticelli (10).

Bothriocephalus latus: in Sweden, Lönnberg: in Holland, Cathe. angustatus in Scorpæna porcus, in Sardinia; MARCIALIS. B. qua ratus in Stenorhynchus leptonyx, and B. tectus in Cystophora prob cidea, n. spp., S. Georgia; v. Linstow (4). (B. longicollis) Molin = Bothriotania, n. g.; RAILLIET (2, VII).

Bothriotenia, n. g. for (Bothriocephalus longicollis); BAILLIET (2, VII). Caryophyllæus referred to suborder Cestodaria, and C. tuba, anatom: MONTICELLI (9).

Ceratobothrium, n. g., with C. xanthocephalum, n. sp.; MONTICELLI (7). Davainea proglottina: some other intermediate host than Limax, as variations according to locality; RAILLIET & LUCET (1).

Dibothriorhynchus monticellii, n. sp.; MONIEZ (1, XII).

Dicranotania, n. g. for Bird Tanias; RAILLIET (2, VII).

Dipylidium caninum, with fenestrated segments; RAILLIET (2, 111); the anal glands of the dog; RAILLIET (2, IV). D. trinchesii, n. si from cat : DIAMARE.

Drepanidotania, n. g. for Bird Tenias; RAILLIET (2, VII).

Echinococcus hominis: Kinyonn, Sangalli, Schalten & Hamen, Pury multilocular variety, MUDD, MANGOLD, VIERORDT, HUBER (2 MANGOLD thinks it has a different cysticercus to the unilocula

Echinocotyle, n. g., with E. rosseteri, n. sp. in duck, cysticercus in Cypr cinerea ; ROSSETER.

Epision, n. g., with E. plicatus, n. sp., LINTON (1); but identified wit Tania malleus, Stiles (10).

1

Gymnorhynchus reptans, curious long larval appendage in, and its migration; Moniez (1, xi).

Gyrocotyle referred to suborder Cestodaria; Monticelli (9).

Moniezia: genus discussed and its different species defined; STILES (9).
M. expansa: notes on anatomy, and distinction of species from others with which it is often confused; STILES (8).
M. plunissima, n. sp., STILES (8).

Prosthecocotyle, n. g. for Tania forsteri; MONTICELLI (7).

Schistocephalus dimorphus: excretory canals shown to have a pair of apertures in each segment; RIEHM.

Tania botrioplitis, Gonads; FILIPPI. T. solium; specimen with hooks replaced by 12 large papillæ; CONDORELLI (1). T. saginata: its occurrence and prevalence in W. Europe in olden times; BLANCHARD (5): prismatic variety of; COATS: Sommer's plasmatic vessel in; BLOCHMANN. T. mediocanellata, as transmitted between man and animals; Peroncito. T. of Hyrax; Setti. T. litterata, anatomy; CONDORELLI (5). T. filicollis and T. torulosa, anatomy; KRAEMER (1). T. serpentulus and its hosts; STOSSICH (4). T. setigera (of Anser), eysticercus found in Cyclops brevicaudatus; T. brachycephala (of Machetes), cysticercus found in S. crassicornis; v. Linstow (2). T. malleus: notes on anatomy; v. Linstow (2): identification of Linton's genus (Epision) with it; STILES (10). T. cuneuta, in Brazil, with some other intermediate host than Allolobophora fatida; MAGALHAES (1). T. tenuirostris found in goose; RAILLIET (2, VII). T. compressa, macrocantha, n. spp., in birds; Linton (1). T. delafondi, n. sp., in pigeon; RAILLIET (5). T. rallei, n. sp.; STOSSICH (4). T. sp.? from a water-bird; v. Linstow (4). T. imbutiformis referred to genus Mesocestoides, Vaill.; RAILLIET (2, VII). expansa referred to genus Moniezia; STILES (9). T. giardi, its relations to genus Moniezia; STILES (9): notes on its anatomy; T. lanceolata, Blochm., made type of a new STILES (8). genus Drepanidotænia; T. coronula of a new genus Dicranotænia, RAILLIET (2, VII); T. forsteri of a new genus Prosthecocotyle, Mon-TICELLI (7). T. (Hymenolepis) diminuta, very ancient case in man; RAILLIET (6): notes on anatomy; ZSCHOKKE (2).

Thysanocephalum crispum, anatomy; LINTON (5).

## Cystic Forms.

Cysticercus acanthorhynchæ, bifurcæ, hamanni, integræ, tenuirostris, notes on, and C. lacertæ, n. sp.; v. Linstow (3). C. pisiformis, specimen with 6 suckers; Railliet (2, 1). C. tenuicollis, in a 4-6 weeks' old kid; Railliet (2, v): in the antelope; Railliet (2, vi). C. cellulosæ under the skin of a middle-aged man; Zschokke (2). C. in liver of hog; Caprini (1, II): alive in lens of eye; Graefe Cysticercoid sp.? in a Freshwater Calanid, Eurytemora lacinulata; Richard.

#### 2. TREMATODA.

Monograph of Distomids in Mammals; STOSSICH (1) gives list of 61 species, with their various hosts, and of 83 Mammals, with Distomids known to occur in each.

Monograph of Distantides in Birds; Stossicii (2): carried out on same plan, 64 species.

Histology of Trematodes; Brandes (1). Anatomy and histology, &c., of Digenetic Trematodes; Braun (4).

Synopsis of Monogenetic Trematodes; SAINT REMY (1).

MONTICELLI (6) proposes the following classification of Trematodes:-

Subord. I. ETEROCOTYLEA.

Fam. 1. Temnocephalidæ.

Fam. 2. Tristomidæ.

Fam. 3. Monocotylidæ.

Fam. 4. Polystomida.

Fam. 5. Gyrodactylide.

Subord. II. ASPIDOCOTYLEA.

Fam. 6. Aspidobothridu.

Suborder III. MALACOCOTYLEA.

Fam. 7. Holostomidæ.

Fam. 8. Amphistomida.

Fam. 9. Distomida.

Fam. 10. Didymozoonida.

Fam. 11. Monostomidæ.

He gives a revision of the family Aspidobothridæ, which he would have include the genera Aspidogaster, Aspidocotyle, Platyaspis, Macraspis, and his new genus Cotylogaster.

HASWELL (5) in the Macleay Memorial volume of the Linn. Soc. N.S.W., gives a monograph of the family Temnocephalidæ. In the Abh. Ges. Halle (3) he discusses the position of the family, concluding that, although he considers it to belong to the Trematodes, he could not find fault with anyone classing it as an aberrant family of Rhabdoccels.

BRANDES (2) criticizes Monticelli's classification as far as it affects the Monostomidæ. He shows that the following species said to belong to the genus Monostomum really belong to other genera:—M. liguloideum is an Amphiline; M. squamata and M. spirals belong to the genus Distomum; M. echinostomum is a synonym for D. planicola, and M. hystrix for D. endolabrum; M. cochleariforms is a Gasterostomum; M. cornu he is uncertain about. There are then left 24 species belonging to the genus Monostomum, which he enumerates and groups, Actinodactylus, n. g. of Temnocephaleæ; Haswell (4, 111) & (5).

Amphistomum subclavatum, anatomy and development, Looss: description of its cercaria found in Planorbis contortus, Lang. A. conicum, histology; Brandes (1). A. bothriophoron, n. sp., anatomy, Braun (3). A. scleroporum, in intestine of Chelonia viridis; Walter,

Bilharzia hamutobia, in Tunis, VILLENEUVE, CAHIER: eggs and embryo, CAHIER, RAILLIET (3): morbid phenomena caused by it in man, COLLOBIDI.

Culceostoma elegans and inerme, figured and described; PARONA & PERUGIA (2).

Cotylogaster, n. g. with C. michaelis, n. sp., anatomy; MONTICELLI (6).

Ductylocotyle tuschenbergii, figured and described; PARONA & PERU-GIA (2).

Distomum caudatum, development from cercaria, which found in kidney of Helix hortensis; BLOCHMANN. D. echinatum and endolobium, notes on; v. Linstow (3). Distomum parasitic in Planocera pellucida and simrothi; v. Graff (1). D. folium, anatomy, Braun (1) and (3): he finds it more normal than Zschokke does. D. goliath, v. Ben., from Balanoptera rostrata, description; LÖNNBERG. D. hepaticum, specimen with genital apparatus abnormal, Spengel: life-history, LUTZ: shown to have some other host than Limnœus truncatulus in Tonkin; GIARD & BILLET. D. lanceolatum, in an Arab, in Alexandria; ZSCHOKKE (2). D. leptosomum, cercaria encysted on Holothurians; Cuénot, Matzdorf. D. magnum, Bassi, Stiles (6): Fasciola carnosa, Hassall, identified with it, LEUCKART. D. nodulosum, parasitic in Cray-fish; Linton (2). D. megastomum and richardii, anatomy of suckers described; CRETY (1). D. richardii, Vitelline nucleus in ova, CRETY (2), MONTICELLI (2). D. veliporum, Vitelline nucleus in ova; MONTICELLI (2). D. species?, Cysts found in walls of heart of Coregonus muruena; ZACHARIAS. D. baraldii, n, sp. from Zamenis viridiflavus, and a stage in its development, description; Sonsino (2). D. capriciosa, n. sp. ?, (cercaria only), found in gonads and viscera of Ophiurids; CUÉNOT, MATZDORF. D. calomaticum, n. sp., in Chinese oxen; GIARD & BILLET. D. flexum, n. sp., Linton (1). D. pulicis and sialidis, n. spp., v. Linstow (3). D. subflavum, n. sp., from Zamenis viridiflavum; Sonsino (2). (D. texanicum), n. sp., FRANCIS, = D. americanum, Hassall, STILES (7), which very probably = D. magnum, Bassi. (D. verrucosum), n. sp., Linton (1): but name already taken, Stiles (10).

Dibothrium exile, n. sp., LINTON (1).

Eurycælum sluiteri, Brock: suggestion that belong to genus Apoblema; Braun (2) and (3).

Fasciola carnosa, Hassall, shown to = Distomum magnum, Bassi; Leuckart, Stiles (7).

Gyrodactylus elegans, notes on; v. Linstow (3).

Holostomum variabile, description and fig.; LINTON (1).

Homalogaster poiriei, n. sp., in Chinese oxen; GIARD & BILLET.

Microbothrium (Pseudocotyle) apiculatum, anatomy and histology; SAINT REMY (2) and (3).

Microcotyle: monograph of genus, with list of fishes on which the various species are parasitic, anatomy of 11 species; PARONA & PERUGIA (1). M. salpæ, n. sp., in Salpa; PARONA & PERUGIA (1).

H

- Monostomum: revision of all the species parasitic in Grallids and Palmipeds; these are reduced to 3:—Aavum, mutabile, and surcidonicola; Monticelli (5). M. capitellatum, in Box Salpa, notes on; Monticelli (4). (M. cymbium) = M. favum, its anatomy and affinities; Monticelli (5). M. mutabile, Zed., epidermal layer; Braun (3). M. spinosissimun, St., and stossichianum, n. sp., in Box Salpa, notes on; Monticelli (4). M. trigonocephalum, reticulare, and proteus, in intestine of Chelone viridis, and points in anatomy; Walter. (M. ovatum), Molin., and (attenuatum), Rud., = Notocotyle verrucosum; Monticelli (3). M. alveatum, Mehlis? = young of N. verrucosum; Monticelli (3).
- Notocotyle: historical sketch, anatomy and systematic, of the genus;

  MONTICELLI (3): two species (N. verrucosum and N. alceatum)
  recognized.
- Octobothrium pollachii and chrysophrii on fish at Roscoff; Saint Rémy (4).
- Octocotyle leptogaster and thunning, figured and described; PARONA & PERUGIA (2). O. scombri, discussion of name; PARONA & PERUGIA (2).
- Onchocotyle appendiculata and prenanti on fish at Roscoff; Saint Remy (4).
- Phyllonella soleae, gonads, SAINT RÉMY (3): note on, PARONA & PERUGIA (2).
- Plagiopeltis duplicata, figured and described; PARONA & PERUGIA (2).

  Pseudocotyle squatinæ, notes on; PARONA & PERUGIA (2).
- Temnocephaleæ: monograph of family, Haswell (5): position and relationships, Haswell (3): discussion of genus Temnocephala, geographical distribution of species, and habits, Vayssière: excretory system, Haswell (2). 1. madagascariensis, n. sp., parasitic on Astacoides madagascariensis, anatomy and histology, Vayssière: three other new species, Haswell (5) [See also Actinodactylus, n. g. of Temnocephaleæ; Haswell (4, III) and (5).]
- Tetraonchus van benedenii, additional description and note on habitat: PARONA & PERUGIA (2).
- Tristomum molæ and interruptum, note on occurrence; PARONA & PERUGIA (2).
- Udonella pollachii, note on occurrence; PARONA & PERUGIA (2) & SAINT RÉMY (4). U. n. sp. on Solea vulgaris; SAINT RÉMY (4).

#### CERCABIAS.

Cercaria Limnosa truncatulus, n. sp., v. Linstow (3). C. pectinata, n. sp., from Donax anatinum (its final host probably being a sea bird), anatomy; Huet. C. virax, ? n. sp., found in Cleopatra bulminoides; C. obscura, ? n. sp., found in Limnoca natalensis; C. pleurolophocerca, n. sp., found in Melania tubercolata and Cleopatra bulminoides (tail with transparent, bilateral, expanded membrane, supported by pro-

cesses resembling setse, probably the cercaria of an Amphistomum); C. capsularia, n. sp., in Cleopatra bulminoides; C. pigmentata, of Amphistomum sp.?; C. (? agilis, of Filippi), of Distomum recurratum, v. L., in Physa alexandrina and others; C. distomatosa, sp.?, in Cleopatra bulminoides; C. microcotyle, Fil., C. costata and fusicauda, La Val. St. G., with the different Molluscs in which they are found: Sonsino (4).

#### 3. NEMERTINEA.

Bürger (2) proposes the following classification for the Moditerranean forms:—

Order A. PROTONEMERTINI.

Genus 1. Carinina, with 1 species.

Genus 2. Carinella, with 10 species, of which 4 are new.

Genus 3. Hubrechtia (n. g.), with 1 species.

Order B. MESONEMERTINI.

Genus 4. Cephalothrix, with 5 species, of which 3 are new.

Genus 5. Carinoma, with 1 species.

Order c. METANEMERTINI [= Enopla of M. Schulze. (To be dealt with in a second part of the paper.)]

Order D. HETERONEMERTINI.

Suborder a. AMICRURÆ.

Genus 6. Eupolia, with 4 species.

Genus 7. Valencinia, with 2 species, of which 1 is new.

Genus 8. Lineus, with 14 species, of which 9 are new.

Genus 9. Borlasia, with 2 species, of which 1 is new.

Suborder b. MICRURÆ.

Genus 10. Micrura, with 6 species.

Genus 11. Cerebratulus, with 18 species, of which 10 are new.

Genus 12. Langia, with 1 species.

The relationships are expressed in a genealogical tree.

History of Freshwater Nemertines, their geographical distribution, and their origin; Guerne (2). Notes on Freshwater Nemertines; Valllant.

Nemertines of Noumea; JOUBIN & FRANÇOIS: of the Baltic (8 spp.); KOJEVNIKOV: of Solovetzki; KNIPOVICHA.

(Balanocephalus pellucida), v. Kennel, = Eupolia pellucida, v. Ken.; Bürger (2).

Borlesia immaculata, n. sp., Gulf of Naples, BÜRGER (2).

Carinella polymorpha, on the English Coast; RICHES, GARSTANG (1). C. albida, minuta, nothus, rubicunda, n. spp., Gulf of Naples, BÜRGER (2.). (C. annulata), McInt., = C. macintoshi, n. sp.; BÜRGER (2).

Cephalothriz bipunctata, fragilis, hymenæus, n. spp., Gulf of Naples, Bürger (2).

Cerebratulus aculus, ærugatus, anguillula, aureolus, candida, ferrugineus, fuscoides, lividus, notabilis, simulans, ventrosulcatus, n. spp., Gulf of Naples, Bübger (2). C. anos, bicornis, caledonicus, n. spp., Noumea, Joubin & François. C. aurantiacus, Gr., on Devonshire Coast; Garstang (1)...(C. dohrnii), Hubr., = Lineus dohrnii; (C. geniculatus), Qfg., = L. geniculatus; (C. grubei), Hubr., = L. grubei; (C. delle chiaje), Hubr., = Micrura delle chiaje; (C. tristis), Hubr., = Micrura tristis; Bürger (2). C. aurostriatus, Bürg., in Noumea: notes on; Joubin & François.

Eunemertes francisca, n. sp., Noumea, Joubin & François.

Eupolia brockii, Bürg., in Noumes: notes on; Joubin & François.

Geonemertes australiensis, n. sp., Australia, DENDY (1).

Hubrechtia, n. g. for H. desiderata, v. Ken.; BÜRGER (2).

Lineus coccinus, gelbus, kennelii, lobianki, molochinus, nigricans, parculus rufocaudatus, versicolor, n. spp., Gulf of Naples, BÜRGER (2).

Polis geniculatus), d. Ch. (Cerebrat. genic., Qfg.), = Lineus geniculatu; Bfree (2).

Tetrasterms aquarum dulcium (?), in England; BENHAM. T. lacustria. in Lake of Geneva; DU PLESSIS.

Fairnes Mescs, n. sp., Gulf of Naples, BÜRGER (2).

#### 4. TURBELLARIA.

- HALLEZ 4 and 3 would get rid of the Order Dendrocala altogether, and divide the Turbellarians into the following 2 groups:—
  - Diphibization, including the Rhabdocælida of Graff, and the Tricinis of Lang.
  - 2 Tripl: lastice including the Polyclada of Lang.

He gives in (1) an emb-vonic law determining the position of the pharynx in the adult for all his "Diploblastica," finding that it do pends on the relative amount of growth of the cephalic and caude hemispheres of the larva. He discusses previous classification gives laws of Morphogeny of the Turbellarians, and a furth classification of the Triclads, while discussing their affinities with the Alloioccels in (3). In (4) and in (3), the Triclads are divided in 3 tribes, which, with the families included in each, are seen in the following table:—

Order Tricladu. Tribe 1. MARICOLA.

Fam. i. Otoplanida.

Fam. ii. Procerodida.

Fam. iii. Bdellurida.

Tribe 2. PALUDICOLA.

Fam. iv. Planarida.

Fam. v. Anocelida.

Fam. vi. Dendrocælida.

Tribe 3. TERRICOLA.

Fam. vii. Limacopsida.

Fam. viii. Geoplanida.

Fam. ix. Polycladida.

- BERGENDAL (1) would divide the Swedish Triclads into 2 families:

  A. Uteriporida; B. Planariada (including the subfamilies Planariae, Polycelida, and Gundada).
- Anatomy and histology of Triclads; CHICHKOFF. Abnormalities in Triclads; HALLEZ (2). Uterus of Triclads; BERGENDAL (6). Temporary parasitism of a Triclad; Vogt. Terrestrial Triclads from Queensland; Dendy (2) and Spencer (1): from Tasmania; Dendy (3): from Lord Howe I.; Spencer (1).
- The Pelagic Polyclads are dealt with very fully by v. GRAFF (1). Lang's 9 species are reduced to 3, and 2 new ones are added. The anatomy and geographical distribution of the group is fully discussed.
- Spanish Turbellarians (2 species), Fusset: Bohemian, Fritsch & Vávra (2 + 9 species), Kafka. Turbellarians of the Baltic (17 species), Kojevnikof: of Moscow (8 species), Zykoff: of Warsaw (11 species of Rhabdoccels), Nusbaum; (13 species of Rhabdoccels, of which only 7 the same as in Nusbaum's list), Wassilieff, with notes (in Russian) on the various species.
- Bipalium kewense shown probably to have come originally from the Fiji Is.; Bell: its presence in Queensland; Spencer (2), Dendy (2): in Asia and Madagascar; Stoll: its multiplication by transverse fission; Bergendal (5).
- Castrada radiata, and? a n. sp., near Warsaw, NUSBAUM.
- Cotyloplana, n. g., of Land Planarians, with C. whiteleggi and C. punctata, n. spp., Lord Howe I.; Spencer (1).
- Dendrocælum lacteum and punctatum, affinities discussed; BERGEN-DAL (1). D. nausicaæ: fusion of branches of the alimentary canal; HALLEZ (2).
- Derostoma elongatum, Gr., and typhlops, Vejd., near Warsaw, notes on; Wassilieff.
- Geoplana in Brazil and Tasmania; STOLL. G. corrules and G. regina, n. spp., Queensland, DENDY (2), and SPENCER (2). 4 species in Tasmania, 1 in Adelaide, together with a new variety, G. fletcheri, var. adelaidensis.
- Gunda sp.?, temporarily encysted in the skin of a young mursenid fish; Voot. G. ulvæ, remarks on; BERGENDAL (1).
- Gyrator hermaphroditus, Ehrb., near Warsaw; Nusbaum: remarks on synonymy; Wassilieff.
- (Haplodiscus piliger), Weld., = a species of Convoluta; v. GRAFF (3).
- Macrostomum hystrix, near Warsaw; Nusbaum: notes on; Wassilieff.

  Mesostomum, nephridial system; Voigt (2), Sekera. M. productum,
  near Warsaw, Nusbaum; M. personatum, riridatum, near Warsaw.
- Microstomum lineare, near Warsaw; NUSBAUM: notes on, WASSILIEFF.

NUSBAUM, WASSILIEFF (synonymy discussed).

- Opisthomum pallidum, notes on; WASSILIEFF.
- Olisthanella, name proposed for n. g. (for Mesostomum truncatum); VOIGT (2), but see SEKERA.
- Othelosoma symondsii, anatomy; v. GRAFF (2).
- Phagocyta, multiple pharynx; HALLEZ (2).
- Planaria lactea, montana, polychroa, n. spp., Middle Europe, anatomy and histology; CHICHKOFF. P. alpina, reproduction; Voigt (1).

  P. alpina and gonocephala, distribution; Voigt (3). P. lugubris, polychroa, torra, notes on; Bergendal (1). (P. sargossicola), Martens, = Stylochoplana sargossicola; v. Graff (4).
- Planctoplana, n. g. for P. challengeri, n. sp., New Guinea; v. GRAFF (1), figured and described.
- Planocera pellucida, Mortens, P. grubei, Atlantic and Indian Oceans, mimrothi, between the I. of Ascension and the Equator, n. spp.; v. Graff (1).
- Polycelis cornuta, distribution; Voigt (3). P. cornuta, nigra, and tenuis, synonymy discussed; Bergendal (1).
- Polychaerus, n. g. of Accelous Turbellarians, with P. caudatus, n. sp., MARK.
- Polycladus: distribution of genus in Chili, Brazil, and Guatemala;
- Polypostia, n. g. for P. similis, n. sp., Sweden; BERGENDAL (1) (not described).
- Rhynchodesmus dubius, fasciatus, fistcheri, grandis, laterolineatus, mediolineatus, n. spp., Lord Howe I.; Spencer (1). Distribution of genus (Europe, N. America, Ceylon); Stoll: in Queensland, Dendy (2).
- Stenostoma leucops, anatomy; OTT: occurrence in Warsaw, Nusbaum, Wassilleff.
- Stylochoplana sargossicola, v. Graff, Sargossa Sea and N. Coast of New Guinea, description; v. Graff (4). Distribution, nervous system, Gonads; v. Graff (1).
- Syndesmis echinorum on Echinus; CUENOT.
- Uteriporus vulgaris, BERGENDAL (5), (6), and (1).
- Vortex pictus, sexdentatus, truncatus, and n. sp. (figured), near Warsaw; Wassilieff. V. millportianus, common near Warsaw; Nusbaum.
- An Alloioccel Turbellarian, to which no name is given, found in the underground waters of Canterbury, New Zealand, is briefly described by HASWELL (4, 11).

# IV.—NEMATODA, ACANTHOCEPHLA, and CHÆTOGNATHA.

#### A.—BIOLOGICAL.

#### 1. GENERAL

Monograph of the Acanthocephala; HAMANN (2), KAISER (1). KAISER criticizes some points in HAMANN'S monograph in an appendix.

General account and classification of the Chatognatha; STRODTMANN.

Anatomy and histology of Bradynema rigidum; STRASSEN: of Myzomimus scutatus; STILES (1): of Mermis nigrescens; v. LINSTOW (5): of Strongylus crurus; STROSE: of S. convoluta; STADELMANN: of Tylenchus fucicola, n. sp.; DR MAN: of Filaria apipillocephala, n. sp.; Condorelli (2): of F. hancrofti; v. LINSTOW (1): of Nectonema agile; WARD, pp. 143-177.

### 2. Epidermis.

JAMMES (1) shows that there is at first a well-developed continuous layer of ectoderm cells; that then a number of these take on a protective function and become the "cuticle," while the rest persist in the "granular layer" of the adult, especially in the so-called "nerve regions": the more parasitic the Nematode is, the more ectoderm is converted into cuticle, and the less persists as "granular layer." Chatin also shows that there is originally a cellular ectoderm, at least in *Heterodera*. Condorelli (2) describes a new species of *Filaria*, in which the cuticle is annulate, though the rings are not complete at the sides. Ward describes the double row of lightly attached, immovable bristles, along the dorsal and ventral lines of Nectonema agile. Cuticle of Bradynema; Steasen.

#### 3. MUSCULAR SYSTEM.

Histology of muscle cells of Ascaris and other Nematodes; BÜTSCHLI, ROHDE (1, I): of Mermis, comparing them with those of Amphioxus; ROHDE (1, II). In Gordius, ROHDE (1, III) finds the different stages in the development of the muscle-cells (coelomyarian and holomyarian type) persistent in the adult G. tolosanus; in G. prelii there are only muscle-cells of the coelomyarian type, but these are open towards the sub-cuticle, and not towards the coelom, as in Ascaris.

#### 4. DIGESTIVE SYSTEM.

Rudimentary condition of alimentary canal and absence of functional anus in *Nectonema*; WARD: in *Bradynema*; STRASSEN. Œsophagus of *Ascaris* ectodermal, not entodermal; BOVERI. Formation of stomodeum and proctodeum from the mesoderm in *Bradynema*; STRASSEN.

#### 5. Corlon.

Development in Nematodes; HAMANN (1), JAMMES (2). It is shown that there is no splanchnic mesoblast ever developed, and therefore that the "coelom" of Nematodes is not properly homologous with that of other animals. Development in Sagitta, see under "11. Development."

## 6. VASCUIAR SYSTEM.

A new species of Echinorhynchus (E. hamanni) with lemnisce consisting of four plates, two on each side, each bent in the middle, and the one lying inside the other; wall-less blood-vessels both in them and in the cutis; v. LINSTOW (4).

#### 7. EXCRETORY SYSTEM.

HAMANN (1) maintains that in Nematodes the excretory vessels are mesodermal in origin, and that even in the adult they lie in the connective tissue of the lateral ridges, and not in the lateral ridges themselves. Kaiser (2) has discovered numerous flame-cells in the two disc-shaped bodies, near the genital aperture of *Echinorhyachus gigas*; each flame is composed of a number of parallel thin cilia in constant undulation, and they together form an excretory organ in connection with the three vascular spaces which lie in the walls of the uterine bell, passing from the coelom into the uterus.

### 8. LATERAL LINES.

Their ectodermal origin; Wandolleck: and consequent distinction from the exerctory vessels; Hamann (1). Condorrell (2) describes a new species of *Filaria* with a ringed body and segmented lateral line.

#### 9. NERVOUS SYSTEM AND SENSE ORGANS.

Of Ascaris megalocephala, several new points, Hesse; two kinds of sensory organs in the lips described. Of Nectonema agile, Ward, the brain is placed ventrally, but above it there are four large conical cells, sending processes down to it, which are probably ganglion cells. Hamann (2) maintains that the cephalic ganglion of Echinorhychus is endodermal. Kaiser (1) does not agree with him (pt. 2), and criticizes him (Appendix). Development of the whole of the nervous system, like the granular layer, from a single tissue of neuroepithelial elements in Ascaris; Jammes (1). Histology of nerve in Nematodes, a criticism of Apathy [see Annelid part of Record], and maintaining that there is nothing in nerves corresponding to the contractile fibrils of a muscle-cell, though in both kinds of cells there is spongioplasm; Rohde (1, 1). Nerve of Mermis compared with that of Amphioxus; Rohde (1, 11). Nerve of Gordius; Rohde (1, 111).

## 10. GENITAL SYSTEM.

On the origin of the germ cells, as distinguished from the somatic cells, in Ascaris megalocephala; BOVERI: part of the ectoderm, as well as meso- and endoderm, shown to arise from one of the two first segmentation spheres, and from this the two archigonad cells are developed. Centrosomes in egg of A. megalocephala shown to be present at the very beginning of the development of the egg, before the expulsion of the polar body; LEBRUN.

## 11. DEVELOPMENT AND LIFE-HISTORY.

Segmentation and gastrulation in Strongylus; WANDOLLECK. According to Jammes (2), there is no true gastrula stage in Oxyurus, but a solid planula of ectoderm and mesendoderm, the endoderm remaining solid even after its separation from the mesoderm. Gastrula stage in Echinorhynchus; Hamann (2) & Kaiser (1) take different views.

Formation of coelom in Sagitta shown not to be enterocoelic, as hitherto supposed, but schizocoelic, the pouches of the archenteron only becoming part of the alimentary canal of the adult; JOURDAIN, M. S. Formation of "coelom" in Nematodes; HAMANN (1), JAMMES (2), [see under "Coelom"]: according to both the body cavity arises as a split between mesoderm and endoderm, there being no splanchnic mesoblast. Wandolleck shows the formation of the mesoblast in Strongylus from two pole cells of the endoderm posteriorly. Mesoblast formed from the ectoderm, and give rise to almost all the organs in Bradynema rigidum; Strassen.

WANDOLLECK shows how the different organs are formed from the three germinal layers in *Strongylus*; the nervous system is developed from the ectoderm, the large posterior mesoderm cells persist and form the gonads. In *Oxyurus*, JAMMES (2), the gonads are formed in the middle of the body, the only part where the mesoderm is more than one-cell layered. According to Boveri, in *Ascaris* the gonads are formed from cells which were originally part of the ectoderm. In *Sagitta* they are shown to arise from the mesoblast by JOURDAIN, M. S.

Development of Bradynema rigidum, STRASSEN: of Heterakis, RAILLIET & LUCET (2): of Echinorhynchus, KAISER (1).

Life-history of Filaria papillosa, DEUPSER: of F. recondita, CALAN-DRUCCIO: of Sclerostomum tetracanthum, GILES (1): of Strongylus convolutus, STADELMANN: of Mermis nigrescens, v. LINSTOW (5): of Gordius pustulosus, CAMERANO (1): of Echinorhynchus proteus, ZSCHOKKE [see under "Distribution according to hosts"].

Discovery of male: of Filaria medinensis (living inside the female), Charles: of the Filaria of the "boutons hémorrhagiques" of the ass, RAILLIET & MOUSSU: of Gordius pustulosus, Camerano (1). Discovery

of female of Nectonema agile; WARD.

Descriptions of embryos of Filaria sanguinis hominis, NABL SABRAZÉS: of embryos and larves of F. recondita, CALANDRUCCIO.

Pædogenesis in an *Echinorhynchus*, Hamann (2): criticized, Ka (1). Pædogenesis and protandric hermaphroditism, without heterogin *Bradynema rigidum*; STRASSEN.

## 12. VARIATIONS.

STILES (1) shows that the individual variation amongst par Nematodes is so great that were it not for the complete series of necting links, the extremes might be regarded as distinct species.

#### 13. CLASSIFICATION.

JOURDAIN, M. S., proposes to class the Chatognatha with the Asci and Amphioxus as a group, the "Protovertebrata." STRODTMANN pro a classification for the group CHÆTOGNATHA in itself, recognizing genera: Sagitta, Krohnia, and Spadella; he enumerates the charistics of value in classification, and gives tables for distinguishing species. A classification of the ACANTHOCEPHALA, recognizing families, is given by HAMANN (3), [see Systematic part of Rec The systematic position and relationship of Nectonema agile is disc by WARD, pp. 182–188, and it is concluded that the genus is most n allied to the Gordiida, since it has most of the characteristics v separate this family from the rest of the Nematodes.

### 14. DISTRIBUTION ACCORDING TO HOSTS.

#### a. General.

Distribution of several Nematodes, and of *Echinorhynchus*, according hosts, Stossich (4): of Mecklenburg species (35 Nematodes, an species of *Echinorhynchus*), BRAUN (5).

#### b. Mammals.

Schlerostomids of equine animals, GILES (1): one new specie robustum) in mules, GILES (2). Filaria in ass, RAILLIET & Mot in body cavity of horse and cattle, STILES (2). F. papillosa, in a cavity of horse and ox; Deupser. Trichocephalus, in liver of an Caparini. Filaria in heart and lungs of Chinese horses and c Fauvel. Strongylus of heart and lungs of dog; Railliet & Ca (2). F. immitis and others in Japanese dogs; Janson & Tokisi Unsuccessful attempts to transmit Strongylus vasorum from one de another; Railliet & Cadiot (1). Esophagostoma columbianum, Cu Strongylus colubriformis, and Spiroptera verrucosum, n. spp., in It sheep; Giles (3) and (4). Ascarids in liver of pigs; Orth Strongylus rubidus in stomachs of American pigs, and its effective cavity of the sheet of the strongylus rubidus in stomachs of American pigs, and its effective cavity of the strongylus rubidus in stomachs of American pigs, and its effective cavity of the strongylus rubidus in stomachs of American pigs, and its effective cavity of the strongylus rubidus in stomachs of American pigs, and its effective cavity of the strongylus rubidus in stomachs of American pigs, and its effective cavity of the strongylus rubidus in stomachs of American pigs, and its effective cavity of the strongylus rubidus in stomachs of American pigs, and its effective cavity of the strongylus rubidus in stomachs of the strongylus ru

HASSALL & STILES. Frequency of Strongylus in rabbits; RAILLIET (7). Trichocephalus (2 only) in intestine of ferret, resembling the T. depressiusculus of the dog. Echinorhynchus bullosus, n. sp., in Cystophora proboscidea; v. Linstow (4). E. novellæ, n. sp., in bat; PARONA (2). E. hamanni, n. sp., in Stenorhynchus (Ogmorhinus) leptonyx; v. Linstow (4).

### c. Birds.

Filaria serrata, n. sp., in Circus cyaneus hudsonensis, Echinorhynchus rectus, n. sp., in Larus, sp.?; Linton (1). Trichosoma rubrum, n. sp., in Spizilla socialis; Linton (3). Sclerostomum apiostomum, in Macacus cynomolgus; Willach. Heterakis monticelliana, n. sp., in Otis tarda; Stossich (4). H. perspicillum, in Numida meleagris, H. papillosa, in three other birds; Railliet & Lucet (2). In domestic fowl, H. brasiliensis, n. sp.; Magalhäes (1, 11). In fowl's egg, Ascaris inflexa; Forel.

## d. Reptiles.

In the venom organ of the common rattlesnake, a viviparous species of Nematode; Tuttle. Oxyurus longicollis, Schm., in Testudo nemoralis, development; Jammes (2).

#### e. Fish.

Ten species of *Echinorhynchus*, and the fish they inhabit, including *E. thecatus* and *E. attenuatus*, n. spp. (in *Roccus americanus* and *Accipenser brevirostris* respectively); LINTON (4). Hosts of *E. proteus*, ZSCHOKKE, intermediate hosts shown, one to be a fish, the other a *Gammarus*, when there are two, though either of these apparently will serve alone.

## f. Arthropods.

Filaria recondita in Pulex of dog, its intermediate host shown to be either the Pulex of dog or man, or Rhipocephalus siculus of dog; CALLANDRUCCIO. Gordius pustulosus occurring abundantly in Blaps mucronata, also found in Sphodrus leucophthalmus, Harpalus aneus, several species of Amara, several kinds of spiders, and in Oniscus murarius: only one host shown to be necessary, as the development of the egg to past the larval stage goes on in damp earth; CAMERANO (1). Gordius doriæ, n. sp., in an Acanthodis; CAMERANO (3). A Gordius in Epinephele tithonus; FOWLER. Nectonema agile,? with a Crustacean as intermediate host; WARD. Gammarus? as intermediate host of Echinorhynchus proteus; ZSCHOKKE. Liparis steineni with E. sp. ?, and Notothenia corriceps with E. megarhynchus, n. sp.; V. LINSTOW (4). An Anguillula in larva of Asynapta citrina; KIEFFER. A Mermis larva in Chironomus; TRYBOM. Bradynema rigidum in Aphodius functarius; STRASSEN.

## z. Elikkolome.

A CARTIL COURTER DE FOLINGE MATERIEF (3).

### A. France.

From which among plants and how to destroy them; Bos in Finish many influent spenes of Tyle what and by He many in the means of protecting the finish in the means of protecting the finish many in the means of the partitle in the finish many in the finishes. The many in Massach in the finishes in the many in the finishes. The many in the many in a Fig. A. Nematode Tyle many in a Fig. 12 Massach.

## If Marina Nites.

The first that is if First amplitude him is during and F were E Tro a art a mi perentin Mersin: E gereins and the first later of sempling southest matter form not yet so , which we have the set of F . The speciality  $=F_{0}$  be also form y in the many to the for its intermediate both. Discovery and ;  $\omega_{2}$  =  $\pi$  = 1 which indeed by F > h in the  $T_{i}$ S,  $A_{ij}$ Siath Solvenson a Commune and the di I be object a unithe in and it will a fine up to the Barrier to Arbylist to make the control of the control o Thumbes of Marier Lois Courts Contactington engre in 19 lium 7 Silanguenequen an emen i va liele en vi e e fimil a Itilia Gre on the later of the same of their transmission will in in the strategie auf andere ber Erstein. E till i ville villen till trille. Kliptike. Pro Lilling by 1 se a Trille. Strate. Games-within El con Exerciasaria afort Diversa Avironi: a use of a suspension of the first terms of the

## 11. Promotes and Harrs.

That is enamed if The course is to be a Nation & San a feether of the sand in an inference of the a a reverse theorem is examined a inference species of recolability & loss that community to resect feeders on in a F are a Morsey Regressive changes of Trocking in the mission, and they depend on the little of the fiberarios spin Wall on the 14.

d sales

3

#### 17. METHODS.

For killing Nematodes for sections: corrosive sublimate + 70°/o alcohol + a few drops of 5°/o acetic acid, added while rolling the Nematode between two slides, recommended; STILES (11): for mounting Trichinæ; PRESTON: for staining and preparing embryos of Nematodes; Nabias & Sabrazés, Callandruccio, Wandollek, and Strassen, p. 701: for histology of the adult; STRÖSE, Jammes: for histology of Nectonema; Ward, pp. 136 & 137: of the Acanthocephala; Kaiser (1): for preserving Sagitta; Cunningham.

## B.—FAUNISTIC.

### 1. Parasitic Forms.

[Gordius is placed with the non-parasitic forms.]

General remarks on distribution of Land Nematodes; STOLL.

Geographical distribution of *Filaria sanguinis hominis*; MANSON: its absence in China; FAUVEL. Geographical distribution of *Tylenchus devastatrix*; Bos (3).

Denmark: Trichina, FRIIS. Mecklenburg: 16 genera and 35 species of Nematodes, 13 species of Echinorhynchus; BRAUN (5). S. Italy: Ankylostoma; CONSALVI. Venice: other Nematodes; STOSSICH (3). Trieste: new species of Heterakis; STOSSICH (4).

Japan: Filaria immitis and other parasites of dog; Janson. China: a Filaria fatal in horses and dogs, Ascaris lumbricoides in man; Fauvel. India: Nematodes of sheep and horses; Giles (1), (2), (3), and (4).

Porto Rico: Echinorhynchus novellæ, n. sp., PARONA.

Brazil: Heterakis brasiliensis, n. sp., in fowl; MAGALHAES.

S. Georgia: Ascaris osculata (which hitherto only found in Arctic regions), two other Ascarids, 3 new species and 1 sp.? of Echinorhynchus; v. Linstow (4).

#### 2. Non-parasitic Forms.

N. Atlantic Ocean: geographical distribution of Chatognatha; STRODT-MANN. Devonshire Coast: Spadella cephaloptera; Garstang (1). Bergen: Enoplida; Appelöf.

Bohemia: pond Nematode; Fritsch & Vávra.

Sardinia: Gordius villoti, aquaticus, tolosanus, and pustulosus; MARCIALIS.

Burma: Gordius doriæ, n. [sp., and G. villoti, rosa, and sphærurus; Camerano (3). Sumatra: G. weberi and G. sumatrensis, n. spp., VILLOT (2). Engano: G. modigliani, n. sp., Camerano (5).

S. America,—Venezuela: G. æneus, Camerano (2). Parana (Palmeira): G. paranensis, n. sp., Camerano (4).

S. Georgia: Leptosomatum antarcticum, n. sp., L. sp.? (Q only), Monhystera sp.? (Q only or immature), Oncholaimus sp. (Q only), Euchromadora sp.? (immature only); v. Linstow (4).

#### C.—SYSTEMATIC.

#### 1. NEMATODA, INCLUDING MERMITHIDÆ AND GORDIIDÆ.

Angiostomum macrostomum, notes on larva; V. LINSTOW (3).

Anguillula parasitic in larva of a Cecidomyiid fly; KIEFFER.

Aphelenchus as cause of a loaf disease; ATKINSON. A. olesistus, n. causing disease in Begonias; Bos (2).

Ascaris: subcuticular layer; Jammes (1): muscle and nerve; Ro: (1, 1): muscle, Bötschli: nervous system, Hesse: origin of Gon Boveri: Centrosomes, Lebrun. A. ensicaudata, scombrorum, he Stossich (4). A. scombrorum, n. sp., Stossich (4). A. pi stichi, n. sp., larva, v. Linstow (3). A. osculata, Rud., and 2 o species in S. Georgia; v. Linstow (4). A. inflexa, specimen in hegg; Forel. A. lumbricoides in man, in China; Fauvel.

Bradynema rigidum, v. Sieb., anatomy, histology, &c.; STRASSEN. Dispharagus aduncus, hosts; STOSSICH (4). D. spiralis at Venice, n on; STOSSICH (3).

Euchromadora sp.? (immature) from S. Georgia; v. Linstow (4).

Filaria sanguinis hominis, perstans and diurna; Manson, Simm Slaughter: description of embryos; Nabias & Sabbazés (1) (2): absence in China; Fauvel. F. bancrofti; v. Linstow (distinguished from F. immitis; Megalhaes (2). F. immitis Meter: in dogs of Japan, Janson. Discovery of male of F. meensis; Charles: of another Filaria (in the ass); Raillie Moussu. F. papillosa, life-history; Deupser. F. recondita an immitis (?), description of embryo and larva; Calandruccio. cervina and F. equina, fresh diagnoses; Stiles (2). F. apapiliphala, n. sp., Condorelli (2). F. ephemeridarum and gamn n. spp., v. Linstow (3). F. serrata, n. sp., from intestine of C. cyaneus hudsonius; Linton (1). F. ninni, n. sp., Stossich (3). rigidum), v. Sieb., = Bradynema rigidum, n. g.; Strabsen.

Gordius, muscles and their development in 2 species; ROHDE (1, III)

Gordius parasitic in Epinephele tithonus (2); FOWLER. G. pustul
parasitism and development; CAMERANO (1). G. tolosanus, 1
larva; v. Linstow (3). G. weberi (intermediate between Gor
s. str., and Chordodes) and G. sumatrensis (belonging to Chord
group), n. spp., Sumatra, VILLOT (2). G. modigliani (belonging
Chordodes group), n. sp., Engano, CAMERANO (5). G. paranensis, 1
Palmeira, CAMERANO (4). G. doriæ, n. sp., in an Acanthodis, Bu
CAMERANO (3). G. æneus, Villot, from Venezuela, and
S. American species, anatomy; CAMERANO (2).

Heterakis: remarks on the genus; RAILLIET & LUCET (2)
MAGALHÄES (1, II). H. perspicillum, in small intestine of Nu
meleagris; H. papillosa in three other birds; RAILLIET & LUCET
H. brasiliensis, n. sp., in domestic fowl; MAGALHÄES (1, II); H.
\*inalliana, n. sp., in intestine of Otis tarda; STOSSICH (4).

schachtii, chiefly in its relation to economics; CHATIN, FE

Hystrichis tricolor, in Venice, notes on; Stossich (3).

Leptosomatum antarcticum, n. sp., and a sp.? ( 2 only) from S. Georgia; v. Linstow (4).

Mermis: muscle and nerve compared with that in Amphiozus; ROHDE (1, II). M.nigrescens, anatomy, histology, and life-history; v. LINSTOW (5). M. albicans, larve in Chironomus; TRYBOM. M. crassa, v. L., M. gammari, and M. sialidis, n. spp., larve; v. LINSTOW (3).

Monhystera, sp.? ( ? or immature), from S. Georgia; v. Linstow (4).

Myzomimus scutatus, detailed anatomy, with mention of individual variations; Stiles (1).

Nectonema agile, detailed anatomy, pp. 143-177; discussion on anatomy, pp. 177-182; its relations to Gordius, pp. 182-188; its taxonomy, pp. 137-139; WARD, Bull. Mus. C. Z. xxiii.

Nematodum gamasi, n. sp., larva; v. Linstow (3).

Œsophagostoma columbianum, described and shown to be cause of nodular disease of intestine of Indian sheep; Giles (3).

Oncholaimus, sp.? ( 2 only), from S. Georgia; v. Linstow (4).

Oxyurus longicollis, early stages in development; JAMMES (2).

Rhabdonema, disease caused by; Sonsino (3).

Sclerostomum tetracanthum, life-history; GILES (1). S. robustum, n. sp., in mules, and remarks on the other species in equine animals; GILES (2). S. apiostomum, parasitic in nodules of colon of Macacus cynomolgus; WILLACH.

Spiroptera verrucosa, n. sp., in Indian sheep; GILES (4).

Strongylus convolutus, anatomy and life-history; STADELMANN. S. micrurus, histology; STRÖSE. S. paradoxus, embryonic development; Wandolleck. S. rubidus, in stomach of pigs; Hassell & Stiles. S. vasorum, and other species, in dog; Railliet & Cadiot (1) and (2). S. colubriformis, n. sp., in sheep; Giles (4). S. curticii, n. sp., name proposed for Curtice's S. ventricosus, to distinguish it from Rudolph's; Giles (4).

Syndesmis echinorum, parasitic on Echinus; MATZDORF (3).

Trichocephalus in the intestine of a ferret, Q only found, resembling T. depressiusculus of dog; RAILLIET (2, VIII). T. in liver of an ox; CAPARINI (1, I).

Trichosoma rubrum, n. sp., in thoracic cavity of the Chipping Sparrow; LINTON (3). T. bombinatoris, n. sp., larva; v. LINSTOW (3). T. verrucosum, n. sp., GILES (4), afterwards referred to Spiroptera.

Tylenchus devastatrix, and its ravages on plants; FREAM, Bos (3): its occurrence in a Sphæriaceous plant; NIEL. T. invalis of Edelweiss; Massalongo. T. fucicola, n. sp., parasitic on Fucus; DE MAN.

#### 2. ACANTHOCEPHALA.

Monograph of group; Hamann (2), Kaiser (1): Nephridia; K. Hamann (3) distinguishes 3 families, viz. :—

 Echinorhynchidæ, g. Echinorhynchus, for E. po polymorphus.

- Gigantorhynchidæ, g. Gigantorhynchus, for G. echinodiscus, tæniorides, spira, and ? E. gigas.
- Neorhynchidæ, g. Neorhynchus, for N. claviceps, N. agilis, and others.

Echinorhynchus proteus, development and life-history; ZSCHOKKE (1). E. teres, anatomy and distribution according to hosts; STOSSICH (4). E. lanceolatus, from Venice, notes on; Stossich (3). E. polymorphus and E. proteus, larves; v. LINSTOW (3). E. polymorphus and E. filicollis distinguished; BRAUN (5). E. novellæ, n. sp., from a S. American bat; PARONA (2). E. hamanni, remarkable n. sp. (from Stenorhynchus leptonyx), with anterior part of body apparently serving as a sucker. E. bullosus (from Cytophora probiscoidea), E. megarhynchus (from Notothenia corniceps), n. spp., and E. sp. ? (from Liparis steineni), all from S. Georgia; v. Linstow (4). E. rectus, n. sp., from a lark; LINTON (1). E. thecatus in Roccus americanus, and attenuatus in Accipenser brevirostris, n. spp.; Linton, (4) pp. 528 & 529, with remarks on the following other species in fish. E. acus, p. 525, E. pristis, p. 530, var. tenuicornis, p. 531, E. incrassatus, agilis, serrani, sagittifer, carcharia, proteus, pp. 532-537. (E. globocaudatus), Zeder, = E. tuba, Rud.; Condorelli (3).

#### 3. CHÆTOGNATHA.

Systematic and general monograph, with 2 tables for determining the species; STRODTMANN. 3 genera are distinguished, viz:—Sagitta, with 11 spp.; Krohnia, with 2; and Spadella, with 7.

Development of Sagitta, JOURDAIN, M. S., leading to the suggestion that, with Amphioxus and Assidians, Sagitta would form a group, which may be called the "Protovertebrata."

Spadella cephaloptera, on Devoushire Coast; GARSTANG (1).

## V .- ROTIFERA (including GASTROTRICHA).

#### A.—BIOLOGICAL.

#### 1. GENERAL.

Bergendal (3) gives an important monograph on the Rotifers of Greenland, dealing fairly fully with the anatomy of 82 species, belonging to 42 genera, found there. Although the number of species is large, the number of individuals is comparatively small. The chief characteristic of the fauna is the rarity of the *Rhizota*, except for *Floscularia*, and of the *Tubicolaria*, which are groups with an otherwise almost universal distribution.

WIERZEJSKI (3) gives a general account of the Rotifers of Galicia, dealing with their anatomy, development, &c., with the collecting of

them, and their taxonomy. Cosmovici (2) gives the anatomy of the Roumanian Rotifers, comparing them with those of Galicia.

Anatomy and histology of Atrochus tentaculatus, n. g. & sp. of Rotifer, without a wheel-organ; Wierzejski (4). Description of a new species of Pedalion (P. fennicum), from Finland, with no glandular posterior haired papillæ; Levander. Anatomy and histology of Gastroschiza tricantha, n. g., and type of a new family, of Rotifers, including the Euchlanis lynceus of Ehrenberg; Bergendal (4): of Anapas ovalis, n. g. & sp., belonging to the same family, with carapace like Gastroschiza, but with no foot; Bergendal (4).

#### 2. NERVOUS SYSTEM AND SENSE ORGANS.

Exhibition of subcesophageal ganglion: Zelinka. Sense of vision; Rousselet (4), where it is shown, by experiment, that Rotifers appear to see well even close objects.

## 3. REPRODUCTION, &c.

Remarks on the males and winter eggs of Rotifers; Bergendal, (3) Act. Lund. iii, pp. 130-149. The male of *Notops clavulatus* and of *Triphylus lacustris* found; Western.

## 4. Classification, &c.

VETTER doubts (though no reasons are given) the relationship between Rotifers and the Trochosphere larva. Daday revises the species of Asplanchna, with special reference to the Hungarian forms, concluding that the "reserve jaws," not being part of the real jaw apparatus, are of no importance in classification, but that the external form of both sexes, the shape of the real jaws, the external form of the ovary, and the shell of the winter eggs are of classificatory importance. His species are, however, criticized by Wierzejski (2).

## Навітв, &с.

Occasional parasitism of a species of *Proales* on *Daphnia*; Thompson, P. G. (2). Commensalism of 2 new species, one of *Callidina* and one of *Pterodina*, on *Asellus*; Parsons. *Discopus synapta*, Zel., as an Echinoderm parasite; Cuénot, Matzdorf (3). Moss-haunting Rotifers, 2 new species of *Macrotrachela*; Thompson, P. G. (1).

#### METHODS.

Of collecting; WIERZEJSKI (3): of examining under the microscope; ROUSSELET (1).

#### B.-FAUNISTIC.

Distribution of Rotifers according to saltness of water; Cosmovici (1 & Daday, E. (1). Daday concludes that there are only 28 purely maris species, 32 both marine and lacustrine, 6 also in inland salt lakes; 1 does not deal with the purely freshwater forms; and he is n altogether accurate in the forms whose distribution he does deal wit IMHOF (1) also deals with the distribution of the same kind of Rotife and concludes that of the marine and brackish forms, 9 species occur the Baltic, 25 in the North Sea (23 not yet observed elsewhere), 1 in t Irish Sea, 7 in the Mediterranean; he records 40 species of Euryhyali forms, i.e.: forms which inhabit both sea- and fresh-water. INHOF ( gives a summary of the pelagic fresh-water Rotifers, concluding that the are 29 species, and 1 variety of them; in speaking of the life of from seas, IMHOF (2) mentions 3 species of Rotifers, all belonging to the ger Amuraea. BERGENDAL (3) gives a detailed account of the Rotifer fac of Greenland, comparing it (pp. 150-163) with that of other regio LEVANDER (2) enumerates the Rotifers of the South of Finlar mostly fresh-water ones, but 18 salt-water species occur off Esbo.

#### ARCTOGUEAL REGION.

#### a. Arctic Zone.

Greenland: 82 species and 42 genera, including 3 new genera, 2 n subgenera, each with 1 new species, and 18 other new species; freshwa chiefly, but a few marine; BERGENDAL (3).

Iceland: Pelagic Rotifer of Dyrefjord (Synchata pectinate Poucher. Freshwater Rotifers, 2 species (mentioned only); Guerre & Richard.

#### b. Palæarctic Zone.

Baltic Sea: 9 species, both marine and brackish; IMHOF (1). Baltic: 40 species; KOJEVNIKOV.

Circum-Baltic Region.—Finland: a new species of Pedalic Levander (1). Other Rotifers; Levander (2). Bornholm (?)—"Rön Mölla"—and Stockholm; 2 (or ? 3) species of the new genus Gastroschof which one is probably Ehrenberg's Euchlanis lynceus, and of wh the other would probably serve as type of another new genus, Arap Bergendal (4) & Jägerskiöld.

Norwegian.—Bukkenfjord: a new species of Anuraea; Tho. son, J. C.

North Sea: 25 species, of which 23 not found elsewhere; IMHOF Scotland.—Forfarshire: a new species of Notops, and 2 rare specone of Copeus, and the other Triarthra; Calman. A few miles no to : 2 new species of Floscularia; Hood (1) & (2).

England.—Essex: new species of Macrotrochela; Thompson, P.G. (1). Chingford: & of Triphylus lacustris; Western. Leytonstone: new species of Proales on Daphnia; Thompson, P.G. (2). Epping Forest: new species of Notops, Rousselet (2): 2 new species, Parsons. 3 new species of Macrotrachelous Calidinidæ; Bryce (1). Keston: 2 new species, Rousselet (3). Surrey, Richmond Park: & of Notops clavulatus; Western. Middlesex: List of Rotifers in ponds near Ealing; Lewis. Sussex: new species of Macrotrachela; Thompson, P.G. (1). Isle of Wight: 3 new species; Bryce (1).

Central Europe.—Switzerland: list of Rotifers; IMHOF (3). Tyrol: Stephanoceros eichornii; Dalla-Torre. Württemberg: Rotifer fauna, 117 species; Bilfinger. Bohemia: 4 pelagic pond forms, 2 semipelagic, and 10 littoral; Kafka. Unter-Pocernitz: pond with 9 Rotifers. Gatterschlag: pond with 13 Rotifers and a Gasterotricha. Hungary: 13 species belonging to the genera Asplanchua, Brachionus, and Triarthra; Daday, J. Hungarian representatives of genus Asplanchua, 9 species; Daday, E. (2). Galicia: 2 new genera, each with 1 new species, and 6 other new species; Wierzejski (3). Roumania: 23 species and 15 genera, 3 species new; Cosmovici (2).

Mediterranean: 7 species; Imhof (1). Black Sea: Synchata baltica, Ehr.; Périaslavzeff.

#### c. Nearctic Zone.

United States.—L. Michigan: Rotifer elongatus; Kellicott. Cincinnati: Rotifer fauna, including 6 new species; Turner.

#### NEOTROPICAL REGION.

Argentine: Freshwater Rotifers; WIERZEJSKI (2).

## ETHIOPIAN AND ORIENTAL REGIONS.

[None.]

#### AUSTRALIAN REGION.

Victoria: Rotifer fauna—6 species of Floscularia, of which 1 new; 11 species of the Melicertids, of which 2 (? 3) new, and 1 new variety; 4 Philadissids; 2 Asplanchnids; 3 Synchaetids; 2 Triarthrids; 4 Notomastids; 1 Rattulid; 1 Dinocharid; 2 Salpinids; 1 Euchlanis; 3 Cathypnids, of which 2 new; 1 Colurid, which new; 2 Pterodinids, of which 1 new; 3 Brachionids; 2 Anuraids.

#### C.—SYSTEMATIC.

Actinurus neptunus, Ehr., Roumania, anatomy; Cosmovici (2).
Anapas ovalis, n. g. & sp., Scandinavia; Bergendal (4).

Anuraea cruciformis, n. sp., Bukkenfjord, Thompson, J. C.; 3 species frozen seas, Imhof (2); A. sp.? in ponds of Iceland, DE GUERNI RICHARD. A. cochlearis, var. carinata: notes on distribution Levander, (2) p. 142.

Arthroglena, n. subg. of Diglena, with A. lülkeni, n. sp., Greenlas

Bergendal (3).

Asplanchna: 9 Hungarian species recognized and described, synonymeach given, and also a table for determining the species; Daday (2). Daday, J., also mentions some Hungarian forms. A. herric Guerne, description; Daday (2), Wierzejski (2). A. girodi, Guerdescription; Wierzejski (2): and, under the name of (A. syric Daday (2). A. helvetica in Iceland, de Guerne & Richashown to = (A. kraemeri), Guerne, which = A. priodonta, wild described; Wierzejski (2) and Daday (2). A. sieboldii in H gary; Daday (2): broken up into A. ebbesborni and A. brightu by Wierzejski (2), both of which also described by Daday (2). imhofi), Guerne, belong to either A. ebbesborni or A. brightu Wierzejski (2), and Daday (2). A. triophthalma, Dad., amph Huds., and intermedia, Huds., recognized by Daday (2) as Hunga species, but said not to be good species by Wierzejski (2). A. cinnatus, n. sp., Turner.

Atrochus, n. g., with A. tentaculatus, n. sp., Galicia, WIERZEJSKI anatomy and histology; WIERZEJSKI (4).

Bipalpus, n. g., with B. vesiculosus, n. sp., and including as ano species the Euchlanis lynceus, Ehr.; Wierzejski (3). In a 1 (3) he shows that they = Gasteroschiza flexilis and G. faveol Jägerskiold, respectively.

Brachionus: Hungarian species; DADAY, J. B. n. sp.? (or = B. am ceros, Ehrb., which is a variety of B. pala), Roumania; Cosmo

(2). B. tuberculatus, n. sp., Cincinnati; TURNER.

Callidina: Macrotrachelous forms (i.e., those which when fully extenhave the post anal-portion decidedly shorter than the pre-intestin 21 species recognized, of which 19 British, 4 new species, C. pli (Epping Forest and I. of Wight), lata (I. of Wight and Folkesto spinosa (I. of Wight), and aspersa (Epping Forest), fully descrit BRYCE (1). C. laevis, C. tentaculala, n. spp., and C. sp.?, N. Grland; BERGENDAL (3). C. magna-calcarata, n. sp., commensal Asellus, Epping Forest; PARSONS.

Cathypna, n. sp. (allied to C. luna), Victoria; Anderson & Sheph.

C. leontina, n. sp., Cincinnati, TURNER.

Carlopus similis, n. sp., Galicia, described and figured; WIERZEJSKI Conochilus unicornis, n. sp., Keston, ROUSSELET (3).

Copeus ehrenbergi, Gosse, in Forfarshire; CALMAN.

Diglena? nataus, n. sp., Greenland, BERGENDAL, (3) pp. 94 & 107.

Dinocharis intermedia, n. sp., Greenland, BERGENDAL, (3) pp. 94 & 10 Diops, n. g., of Notommatidæ, with D. marina, n. sp., Greenland; I GENDAL, (3) p. 94.

- Discopus synaptæ, Zel., as an Echinoderm parasite; MATZDORF (3), CUÉNOT.
- Distemma dubia, n. sp, Greenland, BERGENDAL, (3) p. 100.
- Distyla: notes on genus; Lord. D. clara, agilis, inermis, n. spp., British, Bryce (2). D. icthyoura, n. sp., Victoria, Anderson & Shephard.
- Euchlanis parva, n. sp., Keston, Rousselet (3). E. elegans, n. sp., Galicia, description and figure; Wierzejski (3). (E. lynceus), Ehrb., belong to Gasteroschiza, n. g., Bergendal (4), followed by Jägerskiöld: belong to Bipalpus, n. g.; Wierzejski: Bergendal foundit at Rönne-Mölla, Jägerskiöld, near Stockholm, and Wierzejski in Galicia, all independently, and at about the same time.
- Floscularia gossii, n. sp., Stormont Loch, Blairgowrie, occurring on Chara fatida; Hood (1). F. quadrilobata, n. sp., from a pond also a few miles N. of Dundee, attached to Ranunculus and Nitella; Hood (2). F. evansonii, n. sp., Victoria, Anderson & Shephard. F. uniloba, n. sp., Galicia, Wierzejski (3).
- Gastroschiza, n. g. for G. triacantha, n. sp., Rönne Mölla, which allied to or? = (Euchlanis lynceus), Ehrb.; BERGENDAL (4). G. faveolata, n. sp., near Stockholm, and G. triacantha, Berg.; JÄGERSKIÖLD. G. flexilis, n. sp., near Stockholm, JÄGERSKIÖLD, =? Anapas ovalis, n. g. & sp., BERGENDAL (4). [See also under Bipalpus.]
- Gastrotricha, in Bohemia; FRITSCH & VÁVRA.
- Hypopus, n. g. of Notammatida, with H. ritenbenki, n. sp., Greenland; BERGENDAL, (3) p. 45.
- Lucinularia reticulata, n. sp., Victoria; Anderson & Shephard.
- Limnias, sp.?, allied to L. granulosus, Victoria; ANDERSON & SHEPHARD.-Macrotrachela multispinosa and papillosa, n. spp. from moss; Thompson, P. G. (1).
- Mastigocerea capucina, n. sp., Galicia, WIERZEJSKI (3) and (4).
- Metopidia affinis, n. sp., Greenland, BERGENDAL, (3) p. 122. M. elliptica and dentata, n. spp., Cincinnati, Turner. M. ovalis, n. sp., Victoria, Anderson & Shephard.
- Mikrocodides, n. g., with M. dubius, n. sp., Greenland, BERGENDAL, (3) p. 34.
- Monostyla quennerstedti, n. sp., Greenland, Bergendal, (3) p. 118. M. truncata, n. sp., Cincinnati, Turner. M. tentaculata, n. sp., Roumania, Cosmovici (2). M. bifurca and galeata, n. spp., British, Bryce (2).
- " Notholca" ambigua, n. sp., Greenland, BERGENDAL, (3) p. 123.
- Notolca equispinata, n. sp., Roumania, Cosmovici (2).
- Notommata tarda, grönlandica, celer, distincta, longipes, n. spp., Greenland, Bergendal, (3) pp. 55-66.
- Notops clavulatus, & found in Richmond Park; Western. N. minor, n. sp., Epping Forest; Rousselet (2). N. pygmæus, n. sp., Forfarshire, Calman.
- Notostemma, n. subg. of Notommata, with N. makrocephala, affinis, bicarinata, n. spp., Greenland; BERGENDAL, (3) pp. 69-71.
- Ecistes serpentina, Roumania, anatomy; Cosmovici (2). E. wilsoni.

- 1. 45. mil 5.1. 42. d. A. marmilian, Victoria : Asiateus Servicios.
- Factor is now spaces. F famous, from Kalangines 2 only from affecting from F moves standy in their faces are no glandular pr various non-counting popular, and their face incomes processes be no restrictly and the neglecteds are solded in faces; Landscape 1,
- Printer verie Bert, Learner Samons 2.
- Paramos name 1 sp. Schmittel, Bengeniel, (I, p. 5).
- From the relation of the spins-paramia indices of the gain, a summystem of a new spinor from Lance, P. depleneds; Thomas 2 & 2.
- Function was 1 s. Iving manuscrafty is Ludius, Epping Fore Parasse P manuscraft, 1 sp. Galicia, description and figure Virginiant 33. P. Palation, 2 sp. Victoria, Andrews Interfaces.
- Laife samptim u 🗀 Keingut : Killis 1997.
- beginning geinionilism. 2. sp., Generalism. Braternali, (3, p. 111. informa, Button.; Gallien, description, and Agents; Wierzelski (3
- Injuriannya laman, 2. m., Germand, Bracestal, (3) p. 48.
- Transfers. Hungarian species: Danas (7). I. terminalia, Plate, in Fo.
- Institute accurra, 4 franci de Chingland; Western.

#### VL-MESOZOA.

FEZVZZ's papers of last year on Solinello, translated into English Ann. N. H.

NIMEATH (1) gives a summary of the different theories with regard the derivation of the Meto-from the Protocca, and deals especially us Baline Is as a counseting link. APATHY also concludes that the pleasures the Meto- and Protocca, between Trickoplax and Volvox, is sa factorily alled by Salinella.

# XVI. CŒLENTERATA.

BY

## SYDNEY J. HICKSON, M.A., D.Sc.

#### I.—LIST OF PAPERS.\*

- ALCOCK, A. A case of Commensalism between a Gymnoblastic Hydromedusoid (Stylactis minoi, n. sp.) and a Scorpsenoid Fish (Minous inermis). Ann. N. H. x, pp. 207-214, 1 cut.
- Angelis, G. De. Introduzione allo studio degli Antozoi fossili. Riv. Ital. Sci. Nat. xii, pp. 129-132, 141 & 142.
- Anon. The Australian Barrier Reef. Notice of Mr. Saville Kent's book on "The Great Barrier Reef of Australia, 1893," with two photomezzotypes. Nat. Sci. 1892, pp. 646 & 647.
- Antipa, G. Die Lucernariden der Bremer Expedition nach Ostspitzbergen im Jahre 1889. Zool. Jahrb. vi, pp. 377-390, 2 pls. Abstr. in J. R. Micr. Soc. 1892 (v), p. 623.
- Ueber das Vorkommen von rudimentären "Principaltentakeln" bei Lucernariden. T. c. pp. 392 & 393.
- APPELLÖFF, A. Zur Kenntniss der Edwardsien. Bergens Mus. Aarsber. 1891, No. 6, 32 pp., 3 pls.
- BIGELOW, K. P. On a new species of Cassiopea (xamachana) from Jamaica. Zool. Anz. xv, No. 393, pp. 212-214. Abstr. in J. R. Mior. Soc. 1892 (iv), p. 490.
- On the development of the marginal Sense Organs of a Rhizostomatous Medusa (Cassiopea xamachana). Johns Hopk. Univ. Circ. xi, No. 97, pp. 84 & 85. Abstr. in J. R. Micr. Soc. 1892 (iv), pp. 490 & 491.
- On Reproduction by Budding in the Discomeduse. Johns Hopk. Univ. Circ. xi, No. 97, pp. 71 & 72. Abstr. in J. R. Micr. Soc. 1892 (iv), p. 491.

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

- BROOK, G. Preliminary description of (40) new species of Madrepora in the Collection of the British Museum. Aun. N. H. x, pp. 451-465. Abstr. in J. R. Micr. Soc. 1893 (iii), p. 336.
- Brunchorst, J. Die Biologische Meeresstation in Bergen. Bergens Mus. Aarsber. 1890, No. 5.
- CARLGREN, O. Beiträg zur Kenntniss der Edwardsien. Œfv. Ak. Förh. 1892, pp. 451-461, 6 figg. Abstr. in J. R. Micr. Soc. 1893 (ii), p. 195.
- CERFONTAINE, P. Sur l'organisation et le developpement des differentes formes des Anthosoaires. Bull. Ac. Belg. xxii, pp. 128– 148, 1 pl.
- CHUN, C. Die Dissogonie, eine neue Form der geschlechtlichen Zeugung. Leuckart's Festschr. pp. 77-108, 5 pls, 3 figg. in text.
- 15. CLAUS, C. Ueber die Entwickelung des Scyphostoma von Cotylorhiza, Aurelia, und Chrysaora, sowie über die systematische Stellung der Scyphomedusen. Arb. z. Inst. Wien, x, 1, pp. 1-70, 3 pls. Also published separately by Hölder, Vienna, 1892.
- CUNNINGHAM, J. T. On a species of Siphonophore observed at Plymouth. J. Mar. Biol. Ass. ii, 1892, pp. 212-215, 2 figg. Abstr. in J. R. Micr. Soc. 1892 (iv), p. 491.
- DRIESCH, H. Kritische Erörterungen neuerer Beiträge zur theoretischen Morphologie. II. Zur Heteromorphose der Hydroidpolypen. Biol. Centralbl. xii, pp. 545-556, 3 figg. Abstr. in J. B. Micr. Soc. 1892 (vi), p. 799.
- Gerd, W. Zur Frage über die Keimblätterbildung bei den Hydromedusen. Zool. Anz. xv, No. 399, pp. 312-316, 5 figg. Abstr. in J. R. Micr. Soc. 1892 (vi), p. 800.
- GETTE, A. Ueber die Entwickelung von Pelagia noctiluca. SB. Ak. Berlin, 1892, Heft vii, pp. 413-421.
- 20. Gregorio, M. A. de. Intorno a taluni Celenterati mediterranei viventi. Nat. Sicil. xii (3), pp. 74 & 75.
- Gregory, J. W. The Exploration of Coral Reefs by Borings. Nat. Sci. i, No. 1, pp. 50-52.
- GRIEG, J. A. Oversigt over Norges Pennatulider. Bergens Mus. Aarsber. 1891, No. 1, 24 pp., 1 pl.
- 23. Gurley, R. R. The Geological Age of the Graptolite Shales of Arkansas, and New Species of Graptolites. Ann. Rep. Geol. Survey Arkansas, iii (1890), pp. 401-404 & 416-418, 1 pl.
- Häcker, V. Die Furchung des Eies von Aequorea forskalea. Arch. mikr. Anat. xl, pp. 243-263, 2 pls. & 5 figg. in text. Abstr. in J. B. Micr. Soc. 1892 (vi), p. 799.

- Haddon, A. C. The newly-hatched larva of Euphyllia. Sci. P. R. Dublin Soc. vii, pp. 127-136, 1 pl. & 1 fig. in text. Abstr. in J. R. Micr. Soc. 1893 (i), pp. 53 & 54.
- HARTLAUB, C. Zur Kenntniss der Anthomedusen. Nachr. Ges. Götting. 1892 (1), pp. 17-23, 3 figg. in text.
- HICKSON, S. J. Preliminary notes of observations on the anatomy and habits of Alcyonium. P. Cambr. Phil. Soc. 1892 (vii), pt. 6, pp. 305-308. Nature, xlv, No. 1167, p. 455.
- 28. —. Notes on a small collection of *Hydrocorallina*. Sci. P. R. Dublin Soc. vii, pp. 496-510, 3 pls. Abstr. in J. R. Micr. Soc. 1892 (vi), pp. 799-800.
- Note on the female gonophores of *Errina labiata*. Zool. Anz. xv, No. 394, pp. 237 & 238. Abstr. in J. R. Micr. Soc. 1892 (v), p. 623.
- Abstract of a paper entitled, "A Revision of the Genera of the Alcyonaria stolonifera." P. Z. S. 1892, pp. 594-596.
- 31. James, J. H. Manual of the Palsontology of the Cincinnati group: Colenterata. J. Cincinn. Soc. 1892 (xiv), pp. 149-163, 1 fig.
- KIRKPATRICK, R. Reports on the Zoological collections made in Torres Straits by Professor A. C. Haddon. Hydroida and Polyzoa. Sci. P. R. Dubl. Soc. vi, pt. x, pp. 603-626, 4 pls.
- KOCH, G. VON. Kleinere Mittheilungen über Anthozoen.
   Über Kolonien von Bebryce mollis, Phil., welche Cornulariden ähnlich sind.
   Aggregierte Kolonien von Balanophyllia verrucaria, Aut. Morph. JB. xviii, 2 hft., pp. 373-376, and 376-382, with 11 figg. Abstr. in. J. R. Micr. Soc. 1892, pt. 3, p. 377.
- Kleinere Mittheilungen über Korallen. 9. Beobachtung des Wachsens von Clavularia ochracea. Morph. JB. xviii, pp. 605-608, 1 fig. Abstr. in J. R. Micr. Soc. 1892 (vi), p. 799.
- LANG, ALB. Ueber die Knospung bei Hydra und einigen Hydropolypen. Mit einem Vorwort von A. Weismann. Z. wiss. Zool. liv, pp. 365-385, 1 pl. Abstr. in J. R. Micr. Soc. 1892 (vi), pp. 801 & 802.
- Maas, O. Ueber Bau und Entwicklung der Cuninenknospen.
   Zool. Jahrb. v, pp. 271-300, 2 pls. Abstr. in J. R. Micr. Soc. 1893
   (i), pp. 54 & 55.
- NAGEL, W. Der Geschmackssinn der Actinien. Zool. Ans. xv, pp. 334-338. Abstr. in J. R. Micr. Soc. 1893 (i), p. 54.
- 38. ORTMANN, A. Ueber Korallriffe an der ostafricanischen Küste. Zool. Ans. xv, No. 381, pp. 18-20. Translated in Ann. N. H. ix, pp. 339 & 340; copied in Naturwiss. Rundschau, vii, No. 15, p. 196. Abstr. in J. R. Micr. Soc. 1892 (iv), p. 490.

- 2 Merchant A.J. Die Kannleiffe von Dur-er-Salmen und Ung Leit, Janes v. pp. 426–470. I mag. Amer. in J. R. Mer. So. in, p. 224 e. 255.
- 4) Bereit for de voussimilieien Leisungen in der gestnette om antionnen witnend fen Frimes 1909. Arti. v Bei 2 Et 2 pp. 187-173.
- Remark E. Ness and weng belomme Karallen. And Heath an pp. 1—24. 4 pin, and esparately pathilished by F rasen. Hancong.
- 2 to Late 2 Law Kennings for Changington. Aris micn, in 17-222 Type. Lines: in J. R. Mer. Soc. 1892 vi. p.
- 63 Suttera, C. P. Eine genekierkelieine Berichtigung. Die Kotheome win Lechnekolitz. Zwi. Lix. xv. 3g. 256 & 257. Al J. R. Kier. Soc. 1892 [1, 2, 74.
- 55 Tothloven, E. Jahrenbericht für 1999 über die Coelenterate Ausweimen der Spongien und Anthonien. Arch. f. Nat. Iv. 1 Hft. 5, 20, 126–166.
- Die Aksiephen der Flankom Expedition. Ergeb. Pla Exped. 29 pp., 4 pin., and 1 map.
   Warnery, A. See Land, A.
- Willer, V. La étyestion cher les Actinies. Zool. Anz pp. 19-12. Vanderhaeghen. Gand: 1892, 8vo. Extract from See. de Medec. Gand. 1892, pp. 295-305. Abstr. in J. R. Mic 1893 (h., p. 195.
- 42. Zoda, R. Die vitale Methylenblaufärbung bei Hydra. Zool xv. No. 394, pp. 241 & 242.
- Inverso ad alcune particolarità di Struttura dell' I Rend. Ist. Lombardo, xxv, pp. 709-712, 1 pl.

#### II.—MORPHOLOGY.

#### HYDROZOA.

LANG (35) describes the histology of the young buds of Hydra, I drium, and Plumularia, and maintains that the ectoderm, and the ect only, contributes to their formation. When a bud is about to be for the ectoderm of the parent thickens in a certain restricted portion body, by a process of rapid division by fission of the ectoderm After a time some of these ectoderm cells wander through the messive encloderm, and these cells eventually form the encloderm of

bud. The mesoglosa of the parent in the area of the ectoderm thickening then breaks down, and is subsequently replaced by the formation of a new mesoglosa by the young bud.

ZOJA (48) by the use of, comparatively speaking, strong solutions of methyl blue, has been able to clearly demonstrate the ectodermic nervous system of *Hydra*, confirming, in general details, the results arrived at by Schneider. He was able to prove the existence, in the ectoderm, of nerve nodules, from which numerous fibres spring, of nerve cords formed by the junction of several nerve fibres, and complicated networks of fine fibrils and rigid fibrils leading from this nervous system to the nematocysts. These points are illustrated by a coloured plate in the Rendiconti (49).

MAAS (36) describes a stock of Canina buds which he obtained from the enteric cavity of Geryonia. The axis of the stock is not a simple tube, but possesses numerous irregular ramifications. The author is of opinion that a planula is produced sexually, and this settles down in the tissue of the Geryonia, and forms a stock which produces Medusæ by budding, in fact that there is a simple alternation of generations.

DRIESCH (17) gives an account of the different modes of growth in two forms of Sertularella. His investigations, he maintains, prove that there is in the hydroid colony no such substance as idioplasm. A polyp may give rise to another polyp, or a stolon to another stolon or a shoot, according to the physical or mechanical conditions that may be influencing it during its growth.

GERD (18) states that at an early stage in the development of the egg of Bougainvillea, there is a distinct segmentation cavity formed, but that this is soon obliterated by the growth, in a centripetal direction, of the cells of the celoblastula. These cells increase in number by tangential division, and some of them wander into the cavity of the embryo. This process of migration ends in the formation of a solid morula (pseudomorula), and at the time when this is formed the outlines of the cells are very difficult to determine. The mesoglea is formed subsequently, and then the "pseudo-morula" changes into a solid planula.

HICKSON (28) gives an account of a small collection of Hydrocorallina collected by Professor Haddon in Torres Straits. The collection consisted of several specimens, both dried and in spirit, of Millepora murrayi, Distichopora violacea, and a small specimen in spirit of Stylaster gracilis. An account is given of the anatomy of Stylaster gracilis, and there is a short discussion of the relations of the genus Stylaster to Allopora.

Several specimens of Distichopora were examined, and it was found that those of a yellow or orange colour were sterile, those of a purplish brown colour bore male gonangia, and those of a violet colour bore female gonangia. These facts seem to indicate either that the Distichoporas change colour at different stages in their growth, or that the colour is correlated with their sexual condition. In any case, it seems probable that many of the species that have been established on colour characters will have to be abandoned when their anatomy has been examined.

HICKSON (29) states that the wall of the female gonangium of Errina

labiata consists of two layers, one of cetoderm and one of endoder in other Stylasterids previously described.

Hicker (24) gives a description of the early stages of the determent of Equores forskales. The early stages are extremely rette nuclear divisions, up to the sixty-fourth stage, being simultaneously.


#### SCYPHOZOA.

CLAUS (15) gives an account of some new observations on the dement of certain Scyphozoa, and discusses, very fully, the syste position of this group. He found that the 16-armed larva of Cotyl remained unchanged in the Zoological Institution at Vienna and at T all the winter and spring. The first changes that occurred began in when buds were formed, not on the crown, but on the body-wall olarva. Strobilisation began in August, and is limited to the formatione ephyra at a time. In the early Scyphostoma stage, the mesogla clear liquid lamella, composed of a substance very similar to the cor of the endoderm cells. It must be considered to be derived from cells. This is in striking contrast to the origin of the mesoglosa in a maria, where it is derived, according to Kowalewsky and Marion, fro ectoderm.

The inner layer of cells of the proboscis is endodermal. This conthe views of McMurrich, and controverts those of Goette.

Claus states that Goette is also wrong in his account of the dement of the septal funnels. There is no cavity in the septal muscle the funnels do not become the sub-genital pits.

As a general conclusion from these researches, Claus maintains th Scyphomeduse cannot be included in the group of the Anthoroa, correct classification is—



GIETTE (19) finds that, although there is no true Scyphostoma i development of *Pelagia*, the general features of the development of form are very similar to those of *Aurelia* and *Cotylorhiza*. The points of difference are in the absence, in *Pelagia*, of larval tentacle septal muscles.

While the whole inner structure and the course of radiation are the as in other *Meduse*, the rhythm of the formation of the enteric pe and septa is somewhat different, and follows the following course:-

BIGELOW (9) states that, whereas the budding that takes place on any other part of the Scyphostoma than the crown in Aurelia, Cyanæa, and Cotylorhiza is a mere incident in the life-history of the individual, in Cassiopea xamachana the process of budding from the Scyphostoma is an important, if not the chief factor in the perpetuation of the species.

BIGELOW (8) states that Gœtte is wrong in asserting that the rhopalia are formed as evaginations from the under surface of the marginal part of the umbrella independently of the larval tentacles. In Cassiopea xamachana, at any rate, they are formed as modifications of one series of larval tentacles. Otoliths are formed at the base, and the distal portion of the tentacle is absorbed.

ANTIPA (4) found in an example of a very young Craterolophus, between the adradial arms, and therefore in the perradii and interradii, a small true tentacle. This is of importance, he considers, as it proves that the Lucernaridæ are derived from a form which possessed 8 (4 per- and 4 inter-radial) tentacles.

#### ANTHOZOA.

APPELLOFF (6) finds that there is a well-developed nervous system in the ectoderm of the capitulum of *Edwardsia*, but there is no ectodermal musculature. As in Carlgren's tribe *Protanthew*, there are no mesenterial stomata and no acontia. All these and other points show great affinities between *Edwardsia* and the *Protanthew*.

CERFONTAINE (13) discusses the arrangement of the tentacles in Cerianthus membranaceus. Fresh tentacles are constantly being formed. The author also says that if pieces are cut off a polype of Astroides calycularis, polypes arise which never produce a skeleton, but grow nevertheless and reproduce by gemmation.

NAGEL (37) made a series of experiments, proving the existence of a sense of taste in the tentacles of several species of sea anemones. A piece of the flesh of a sardine was greedily seized by the tentacles and swallowed, a piece of blotting paper of the same shape was not touched; a piece of sardine flesh which had been washed in water for some time was seized, but not very energetically, but a piece of blotting paper which had been soaked in the sardine extract was seized as eagerly as the flesh itself.

WILLEM (47) made some experiments on the digestion of sea anemones by feeding them with carminated albumen and the liver of the mussel. He finds that the absorption takes place by the whole of the endoderm and the endoderm only. No grains of carmine or liver globules were ever found in the cells of the mesenterial filaments.

Haddon (25) describes some young larvæ of Euphyllia, 1.5 mm. long. They rested on their aboral surface on the side of the vessel. They had eight complete mesenteries and two incomplete ones, these being the sulco-sulcar laterals. According to Bourne, the adult Euphyllia is radially symmetrical. This larva is obviously bilaterally symmetrical.

HICKSON (27) gives some preliminary notes on the minute anato Alcyonium digitatum. There is no evidence for the existence of a in all the cords of endoderm cells found in the mesoglæa betwee collenters. A nervous system, in the form of a delicate network of and cells, penetrates the mesoglæs. The author also gives an account the mode of retraction of the polyps, and of a series of experime made at Plymouth, proving the tidal rhythm of expansion and controf the polyps in the living state.

Von Koch (33) describes some colonies of Bebryce mollis, which true stolons, like those of the Stolonifera. These stolons, more secrete basally (from the ectoderm) a horny lamella. The authorises aggregated colonies of Balanophyllia verrucaria, which easily be distinguished from those formed in the ordinary wholeding.

Von Koch (34) states that a piece of the stolon of Clau ochracea is constant in growth for twenty days, increasing in length 0.5 mm. per diem, and forming a new polype every sixth day.

STUDER (44) calls attention to a case of reproduction by fission occ in an Alcyonarian (Schizophytum echinatum).

#### CTENOPHORA.

Samassa (42) writes an elaborate and fully illustrated paper of histology of the *Ctenophora*. He was unable to find any definite no system, but he regards the polar bodies, the polar fields, and me bands as forerunners of a nervous system, and capable of perform its physiological functions. He concludes with some remarks in the following street that the *Turbellaria* are derived from the *Ctenopho* 

CHUN (14) has an interesting paper on some of the more recent of his work on the life-history of *Ctenophora*. He defines "Disso as the condition in which a single individual becomes sexually mat two different forms, between which is interposed a stage in which d ration of the sexual products occur.

The particular case discussed is that of *Bolina hydatina*, which forth a sexually mature Cydippe form, followed by a stage in the gonads disappear before the adult characters make their appearance. Dissogony is obviously a form of Pædogenesis, but differs from it interposition of a non-sexual stage between the larval and adult i

The author has also a chapter on the mechanics of the developm the Ctenophora.

#### SYMBIOSIS.

ALCOCK (1) describes a remarkable case of commensalism betwee Scorpsenoid fish, Minous inermis, and the Gymnoblastic hydroid, S minoi, n. sp. The Stylactis is never found without the Minous, Minous without the Stylactis. The association of these two form ide area of the Indian Seas. The Stylactis is usually

on the throat and around the gill opening of the fish, and it seems to afford it additional resemblance to the encrusted rocks among which it lives.

#### CORAL REEFS.

The most important contributions to our knowledge of Coral Reefs during the year 1892, are those of ORTMANN (38). This author states that the reefs of the East African coast, from the coast opposite Zanzibar as far south as Mikiudani, are all fringing reefs. There are no true atolls nor barrier reefs. The coast all along the line has been undergoing gradual elevation, as is proved by numerous remains of old reefs above the sea level.

He found that many corals, such as *Porites, Goniustræa, Caloria*, and *Tubipora*, are able to survive exposure to the sun and air at low tides for hours together.

Some corals, such as *Psammocora*, *Montipora*, and *Lophoseris*, are capable of thriving on a loose gravelly bottom, held together by sea-grass.

In his account of the corals of Dar-es-Salaam, the same author (39 shows that of the 44 species, 12 are widely distributed in the Indo-Pacific region, 23 are Indian species, 2 of which go as far as China, 5 are purely Pacific, and 2 are new.

SLUITER (43) points out that the views as to the origin of coral reefs usually attributed to Chamisso, are really those of Eschscholtz.

GREGORY (21) gives a short account of some of the more recent results obtained by making borings on coral reefs.

The editor of Natural Science (3) gives a short account of Saville Kent's book on the Australian Barrier Reef (1893), illustrated by two excellent photomezzotints of the reef.

#### III.—FAUNISTIC.

BRUNCHORST (11) gives a list of all the Cælenterata known to occur in the neighbourhood of Bergen.

KIREPATRICK (32) gives an account of the *Hydroidea* collected by Professor Haddon in Torres Straits, and a list of all the known species from that district. *Tubularinæ*, 4 species; *Campanularinæ*, 5 species; *Sertularinæ*, 11 species; and *Plumularinæ*, 12 species.

HICKSON (28) gives an account of the *Hydrocorallinæ* collected by Professor Haddon in Torres Straits. 3 species.

ANTIPA (4) describes the *Lucernaridæ* collected in the region of East Spitzbergen. 3 species.

VANHÖFFEN (46) gives an account of the Akalephæ collected by the Plankton Expedition. 9 species, and an account of the distribution of the Cathammata.

ORTMANN (39) gives a list of the corals found upon the reefs in the neighbourhood of Dar-es-Salaam.

THE TWO SQUEETS OF Conference levens and Functions quadra parama was a line of all the Permutalide found on the coast of Norway Leaders of grown in pp. 35-35, a very metal list of the names the spaces of Laborators recurring in different parts of the world.

## IT.—NEW GENERA AND SPECIES.

## HYDROZOA.

#### GTHEORIAGTEA.

Knaparana 'E'; seeciles the following new species:-

Caryne v. Specaryne of Sendrica, p. 606, growing on Cellepora granulas Albany Pranaga. 3 fatha.

Cantiver me raidon, p. 56. growing on Idia printis, Murray I., 15 to!

Zudenierum enfundituisfurme. p. 695, 20 miles N.N.W. of Warrior :

Styleria mine. Alecex (1), p. 212, commensal with the fish Mine mercia a fix Indias Sem.

#### ANTHOMEDUSE.

Form ozz. Harriatz (26), p. 19. Mediterranean. Harriats (26) pt zees against Variety evens in joining Pandaea and Tial The general of the two general are not alike.

#### CALTPROBLASTEA.

Diphana valariformia, Kirkpatrick (32), p. 603, growing on Ayl phenia huxleyi, 20 miles N.N.W. of Warrior I., 51 faths.

#### HYDROCORALLINE.

Millepora tenella, ORTMANN (39), p. 668, Dar-es-Salaam.

#### RHABDOPHORA.

Dicranograptus arkansasensis, Gurley (23), p. 416, Dicellograptus za Arkansas. D. nicholsoni, var. parvangulus (n. var.), id. loc.

Diplograptus trifidus, id. p. 417. Several localities associated with the Dicellograptus fauna. Dictyonid. p. 418, Dicellograptus zone, Arkansas.

#### SIPHONOPHORA.

Muggiaa atlantica, CUNNINGHAM (16), Plymouth. The author remarks that this species may be identical with Haeckel's Muggiaa pyramidalis.

#### SCYPHOZOA.

Vanhöffen (46) disapproves of the division of the Scyphozoa into Octomeralia and Tetrameralia. He divides the Cathammata (of CLAUS) into two divisions:—

Incoronata (Charybdeidæ.)
Lucernariidæ.
Depastridæ.
Tesseridæ.
Coronata (Periphyllidæ.
Ephyropsidæ.

The same author (46) describes the following new species:— Nausithoe clausi, p. 14, Carolines.

Nauphanta vettoris pisani, p. 15, Carolines.

Lucernosa, n. g., which differs from Lucernaria in the more complicated structure of its gonals. It includes four species: L. bathyphila == Lucernaria bathyphila (Haeckel), and the following three which are new, L. waltheri, p. 379, L. kükenthali, p. 386, L. haeckeli, p. 388, all from E. Spitzbergen, 10-35 faths., ANTIPA (4).

Cassiopea xamachana, BIGELOW (7), Jamaica.

## ANTHOZOA.

#### ZOANTHARIA.

#### MADREPORARIA PERFORATA.

Brook (10) adds the following new species of the genus Madrepora:—

M. ambigua, p. 451, Northumberland I. M. arcuata, p. 452, Samoa.

M. armata (= M. spicifera, Dana), M. assimilis (= M. appressa,
Quelch), M. australis, p. 453, Darnley I. and Wreck Bay, Gt.
Barrier Reef. M. bæodactyla (= M. seriata, Brüggemann), M.
bifuria, Java. M. botryodes (= M. gonagra and M. haimei, Brüggemann), p. 454. M. brevicollis, Rodriguez, &c. M. bullata, Port
Denison. M. calamaria (= M. plantaginea and M. acervata, Brüggemann), p. 455. M. clavigera, loc.?. M. cophodactyla, loc.?. M.
coronata, p. 456, Gt. Barrier Reef. M. decipiens, Gt. Barrier Reef.
M. elseyi, N. Australia and Gt. Barrier Reef. M. exilis, p. 457,
Arafura Sea, 10 faths., &c. M. fruticosa, loc.?. M. gemmifera
(= M. seriata, Quelch). M. grandis, Gt. Barrier Reef. M. guppyi,

p. 458, Solomon Is. M. irregularis (= M. alces, Brügge Rodriguez and Macclesfield Bank, 7 or 8 faths. M. kenti, day I. and Gt. Barrier Reef. M. latistella, p. 459, Port D. M. loripes, Gt. Barrier Reef. M. nigra (= M. ehrenbergii, B. M. oligocyathus, p. 460, Mauritius. M. orbicularis, Ceylo patula, Port Denison. M. pectinata, Thursday I. M. rayneri, Fiji. M. recumbens, Gt. Barrier Reef. M. reticulata, Arafu &c. M. sarmentosa, p. 462, Gt. Barrier Reef. M. spectabilis M. squamosa, p. 463, Gt. Barrier Reef. M. syringodes, Gt. Reef, Samoa, &c. M. tenella, p. 464, Macclesfield Bank, 3 faths. M. tizardi (= M. nasuta, B. Smith), M. violacea, Fiji, &c.

ORTMANN (39) describes Madrepora cylindrus and M. horizontalis, Dar-es-Salaam.

Rehberg (41) adds the following species:-

Dendrophyllia japonica, p. 28, Japan.

Canopsammia ramiculosa, p. 29, Japan.

Madrepora incrustans, p. 35, Tonga Tabu. M. repens, p. 36, Mada M. mexicana, p. 38, Mexico. M. dichotoma, p. 39, Pelew. M. pinensis, p. 40, Philippines. M. elliptica, p. 41, Luzon. M. p. 41, Ceylon. M. papillosa, p. 42, Tahiti. M. pelewensis, Pelew.

Turbinaria marmorea, p. 43, Ponapé. T. crispa, p. 44, Tonga. Anacropora spinosa, p. 46, Pelew.

#### PORITDLE.

Porites capricornis, p. 46, Pelew. P. profundus, p. 48, Mad: P. reticulum; ORTMANN (39), Dar-es-Salaam.

## MADREPORARIA IMPERFORATA.

#### TURBINOLIDÆ.

REHBERG (41) describes Heterocyathus pulchellus, p. 8, West Coasttralia. H. oblongatus, p. 9, South China Sea, 16 faths.

#### Oculinida.

Amphihelia adminicularis, p. 10, Japan. Madracis axillaris, p. 10, Fiji, 80 faths.

#### ASTRÆIDÆ.

Pterogyra simplex, p. 14, New Ireland. Mussa crassidentata, p. 16, Pelew Is. Sumphyllia subtilis, p. 16, Fiji.

hyllia stuhlmanni, p. 17, Zanzibar. U. maxima, p. 18, Duke of

Tridacophyllia elongata p. 18, Pelew.

Caeloria elegans, p. 20, N.E. Coast of Australia. C. deltoides, p. 20, E. Coast of Australia. C. australiansis, p. 21, N. E. Coast of Australia.

Hydnophora columellata, p. 21, Yap. H. klunzingeri, p. 22, Samoa.

Clavarina composita, p. 22, Pelew.

Echinopora pellucida, p. 23, Pelew.

Astraosmilia, n.g., allied to Dasyphyllia, but differing from it principally in the long, laterally-connected calyces and the absence of "Collerett." A. connata, p. 664, Dar-es-Salaam; ORTMANN (39).

#### LOPHOSERIDÆ.

- Folioseris, n. g., p. 26: this genus stands close to Pavonia, from which it differs in the arrangement of the calyces, which do not stand in rows. On the other hand, it shows affinities with the genus Mycedium; Rehberg (41).
- F. crispa (= Agaricia crispa, Ehrenberg). F. papyracea (= Pavonia papyracea, Dana).



## XVII. SPONGIÆ.

BY

## R. HANITSCH, Ph.D.

#### I.—LIST OF PUBLICATIONS.\*

- 1. Anonymous. [Note on Sponge Deposits discovered near Island of Pantelleria.] Nature, xlvi, p. 474.
  - APPELLÖFF, A. Om Bergensfjordenes faunistike præg. Bergens Mus. Aarsber. 1891, pp. 1-14.

Sponges, pp. 11, 12, & 14.

- BIDDER, G. Note on the Exertion in Sponges. P. R. Soc. li, No. 313, pp. 474-484, 5 woodcuts.
- On the fiask-shaped Endoderm and Spongoblasts in one of the Keratosa [Cacospongia, sp.]. Op. cit. lii, No. 315, pp. 134-139, 3 woodcuts.
- BRAUN, M. Nachträge und Berichtigungen zum Zoologischen Theil von "Die landeskundliche Literatur über die Grossherzogthümer Mecklenburg," &c. Arch. Ver. Mecklenb. xlv, pp. 87-95.
   Spongiæ, p. 93.
- Buck, E. Einiges über den Bodenseeschwamm. Deutsche Fischerei-Zeitung, xv, pp. 366 & 367.
- Chworostansky, C. Ueber die Zone des Küstenstriches der Solowezki-Inseln. Zool. Anz. xv (1892), pp. 214 & 215.
- 2 Sponges, Suberites glasenapii, Mer., and Halisarca Fr.-schulzei, Mer.
  - 8. CORTI, B. Foraminiferi e Diatomee fossili del pliocene di Castenedolo. Rend. Ist. Lombardo (2) xxv, pp. 991-1012, 1 pl.
    - 1 Sponge, Spongolithis acicularis, Ehrenb., mentioned, p. 1011.

An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

- 11. I Still. 1. The Mannery of the true Nature of the family Technolog. Ioul. Ann. xv 1392, pp. 245 & 245.
  Leady status the penetry over R. v. Leadenfeld of having promotion the family Technolog.
- Français of the American Coloures Heterocale, with proof the theorems of the group, and descriptions of so genera and species. P. R. Sec. Vict. (s.s., v. pp. 69-116.)
  4 new genera : If new species : I new variety.
- 22 Preliminary sensors of Symme polchelle, a new geognesis of Colombian Springes. Op. Ac. iv. pp. 1-6.
  1 new genus. 2 new openios.
- On a new species of Loncosoloxis from the neighbour
   Put: Phillip Heads. Op. od. v, pp. 173-189.
   1 new species.
- Diounizers, L. [Description of Petrodoma schulzei, n. g., Calcarra, supresenting a new order Lithones.] Verh. deuts Sen. i (1992) pp. 143-145.
- 11. Dezema. F. Zode and Wege biologischer Porschung, belet für Hami einer Geristänlichungsmechanik. Jena (G. Fischer für n. 11.2 pp., 5 pär.
- 14. Die Principien der Gerüstbildung bei Rhisopoden, 8 und Erkinsdermen / Fortsetzung . Jen. Z. Nat. xxvi (1 pp. 257-455, 10 pls. ETHERIDGE (29).]
- 17. FANINTZIN, A. Beitrag zur Symbiose von Algen und Mém. Ac. Pétersb. (7) xxxviii, 16 pp.
- 18. Garstang, W. Notes on the Marine Invertebrate F Plymouth for 1892. J. Mar. Biol. Ass. (n.s.) ii, No. 4, pp. 3 1 Sponge, Leucosolenia lacunosa, Bwk.
- 19. Geinitz, H. B. Die Versteinerungen des Herzogthums : Altenburg. MT. Osterlande (n.f.) v, pp. 161-199. Spongiæ, p. 191.
- GRIFFITHS, A. B. The Physiology of the Invertebrata.
   (L. Reeve & Co.): 1892, 8vo, xvi & 477 pp., 81 woodcuts.
   Sponges: pp. 30-32, 79-82, 184, 209, 349, 378.
- 21 HAMITECH, R. Foraminifer or Sponge? Nature, xlvii, I

- [Hanitsch, R.] Foraminifer or Sponge? Nature, xlvii, No. 1219, p. 439.
  - On Neusina agassizi, Goës, a supposed Foraminifer.
- HARTOG, M. M. Some Problems of Reproduction: a Comparative Study of Gametogeny and Protoplasmic Senescence and Rejuvenescence. Q. J. Micr. Sci. xxxiii, pt. 1, pp. 1-79.
   Sponges, pp. 48 & 49.
- 24. HERDMAN, W. A. Fifth Annual Report of the Liverpool Marine Biological Station now on Puffin Island. Proc. Liverpool Biol. Soc. vi, pp. 10-39.
  - Sponges, pp. 26 & 28.
- 25. —. Notes on the Collections made during the Cruise of S.Y. "Argo" up the West Coast of Norway, in July, 1891. T. c. pp. 70-93, 2 pls.
  - Sponges, pp. 88 & 89.
- 26. Hinde, G. J. On Palacosaccus dawsoni, Hinde, a new genus and species of Hexactinellid Sponge from the Quebec group (Ordovician) at Little Métis, Quebec, Canada. Geol. Mag. (3) x, No. 344, pp. 56-58, 1 pl.
  - 1 new genus and species.
- 27. HINDE, G. J., & HOLMES, W. M. On the Sponge-remains in the Lower Tertiary Strata near Oamaru, Otago, New Zealand. J. L. S. xxiv, No. 151, pp. 177-262, 9 pls.
  - 1 new genus, 15 new species, several undetermined.
    - Holmes, W. M. [See Hinde & Holmes (27).]
- 28. HORNELL, J. A Strange Commensalism—Sponge and Annelid. Nature, xlvii, No. 1204, p. 78.
  - Microciona plumosa and Leucodora caca.
- 29. Jack, R. L., & ETHERIDGE, R., Jun. The Geology and Palssontology of Queensland and New Guinea. Brisbane (Beal) and London (Dulau & Co.): 1892, xxx & 768 pp., 68 pls., 1 map.
  - Sponges, pp. 199 & 436-439; 1 new species.
- Kafka, J. Untersuchungen über die Fauna der Gewässer Böhmens.
   Die Fauna der böhmischen Teiche. Arch. naturw. Landesforsch.
   Böhmen, viii, p. 113.
  - Spongillidæ.
- Kellicott, D. S. The Mills Collection of Freshwater Sponges. Bull. Buff. Nat. Club, v, pp. 99-104.
- 32. KOJEVNIKOV, G. La fanne de la mer Baltique orientale et les problèmes des explorations prochaines de cette faune. Congrès internat. de Zoologie, ii, No. 1, p. 140.

- Korschelt, E., & Weltner, W. Die Lebensverhältnisse Dreissensia polymorpha. Naturw. Wochenschrift, vii, pp. 391-39 Spongillidæ, p. 393.
- 84. Lameere, A. Prolégomènes de Zoogénie. Bull. Sci. Fr. Belg. 1 (1891), pp. 399-411.

Systematic position of Sponges, &c.

- 85. LENDENFELD, R. von. Siliceous Spicules of Geodia. J. R. M. Soc. 1892, p. 50. Abstract of paper, Ueber die Kieselnadeln Geodia; Zool. Anz. xiv (1891), pp. 407-409. [Vide Zool. Rec. 18
- Bemerkung über die Homodermidæ. Zool. Anz. xv (18 p. 109.
- Ueber Minchin's Angaben betreffs der Histologie der K schwämme. T. c. pp. 277-279.
- 38. —. Berichtigung. T. c. p. 370.

Reply to Dendy's article, "The Discovery of the true Nature of so-called family *Teichonidæ*," t. c. p. 245. Author states that he arr at his conclusions independently of Dendy.

- Note on Mr. Minchin's paper on Ascetta. Ann. N. H. (6 p. 337.
- Note on Dr. Hinde's Tertiary Sponge-spicules. Op. ci p. 268.
- Bemerkungen über die neuerlich von Dendy beschriebe Kalkschwämme. Biol. Centralbl. xii, pp. 58-60.
- Die Spongien der Adria. 11. Mittheilung. Die Hexacera
   wiss. Zool. liv, hft. 3, pp. 275-315, 1 pl.
- MAAS, O. Die Metamorphoso von Esperia lorenzi, O. S., r Beobachtungen an anderen Schwammlarven. MT. z. Stat. Neal hft. 3, pp. 408-440, 2 pls.
- 44. Die Auffassung des Spongienkörpers und einige neuere Arbiüber Schwämme. Biol. Centralbl. xii, Nos. 18 & 19, pp. 568-Translated in Ann. N. H. (6) x, pp. 399-405: The Interpretatic the Sponge Organism, and some recent works on Sponges. Abs in J. R. Micr. Soc. 1892, p. 803.
- 45. —. [Demonstration of Larvæ of Siliceous Sponges.] \ Deutsch. Zool. Ges. ii, p. 142.
- 46. MARSHALL, W. Spongiologische Beiträge. Festschrift zur Wiederkehr des Geburtstages von Rud. Leuckart. Leipzig (( Winter): 1892, imp. 4to, 36 pp., 8 pp. explanation, 8 pls.
  - 2 new genera, 2 new species.
- 47. MATTHEW, G. F. List of the Fossils found in the Cambrian R in and near St. John. Bull. Nat. Hist. Soc. New Brunswick pp. xi-xxiii.

Sponges, p. xv.

- Minchin, E. A. Dr. von Lendenfeld on the Central Cavity in Euplectella. Ann. N. H. (6) ix, pp. 408 & 409.
- 49. Some points in the Histology of Leucosolenia (Ascetta) clathrus, O. S. Zool. Anz. xv (1892), pp. 180-184, 3 woodcuts.
- 50. ---. Dr. von Lendenfeld als Kritiker. T. c. pp. 415-417.
- 51. —. Note on a Sieve-like Membrane across the Oscula of a Species of Leucosolenia, with some observations on the Histology of the Sponge. Q. J. Micr. Sci. xxxiii, pt. 2, pp. 251-272, 2 pls. Abstract in J. R. Micr. Soc. 1892, ii, pp. 218 & 219.
- The Oscula and Anatomy of Leucosolenia cluthrus, O. S.
   J. Micr. Sci. xxxiii, pt. 4, pp. 477-495, 1 pl.
- Structure of Calcareous Sponges. [Abstract of Minchin's recent papers.] Am. Nat. xxvi, p. 863.
- 54. Munroe, K. Sponge and Spongers of the Florida Reef. Scribner's Magazine, Nov. 1892, pp. 639-649, 6 woodcuts. Popular account of Sponge-fishing.
- 55. Nobre, A. Estudos de Zootomia. Fasc. 2. Porto (Typographia Occidental): 1892, pp. 49-84, 3 pls.

Halichondria caruncula, B.; Cacospongia scalaris, O. S., and Hyalonema lusitanica are described and figured, apparently for practical class purposes.

- NORMAN, A. M. Museum Normanianum; or, a Catalogue of the Invertebrata of Europe, and the Arctic and North Atlantic Oceans. VII & VIII: Spongozoa and Rhizopoda. Durham: 1892, 21 pp. Spongozoa, pp. 5-14.
- 57. OGILVIE, M. M. Contributions to the Geology of the Wengen and St. Cassian Strata in Southern Tyrol. Q. J. Geol. Soc. xlix, pt. 1, pp. 1-78, 14 woodcuts, 5 maps.
  - 22 species of Sponges, including 1 new genus and 1 new species.
- Pearcey, F. G. Foraminifer or Sponge? Nature, xlvii, No. 1217, p. 390.
  - On Neusina agassizi, Goës, a supposed Foraminifer.
- POCTA, P. Ueber Spongien aus der oberen Kreide Frankreichs in dem mineralogischen Museum in Dresden. Mit Vorwort von Dr. H. B. Geinitz. MT. min. geol. Mus. Dresden, xi. Cassel (Th. Fischer): 1892, 4to, 26 pp., 4 pls.
  - 11 new species, 1 new variety.

1892. [vol. xxix.]

- RAUFF, H. Ueber den Bau des Stützskeletes bei den Anomocladinen und Tetracladinen. SB. niederrhein. Ges. xlviii (1891), pp. 33-37.
- 61. —. Ueber eine eigenthümliche Gruppe fossiler Kalkschwämme (Polysteganinæ). T. c. pp. 45-50.

r 9

- 62. THOMEN, J. A. The Study of Animal Life. London (John Mu 1932, 200, xii & 375 pp., 71 woodcuts.
  - Spragie, p. 15%.
- 63. —. Outlines of Zoology. London (Young J. Pentland): êvo, xvi & 641 pp., 32 pks. Spague, pp. 114-13).
- 64. Torsent, E. Sendy of Chonide. J. R. Micr. Soc. 1892, pt. 3, Abstract of Deuxième contribution à l'étude des Clionides. Z. expér. 2 ix (1891). No. 4, pp. 523-554, 1 pl. [Vide Rec. 1291.
- 65. —. Notes histologiques au sujet de Leucosolenia coriacca (! Bwk. Bull. Soc. Z. Fr. xvii (1892), pp. 125-129, 1 woodcut.
- 66. Sur une Éponge du Lac de Tibériade, Potamolepis barrois Rev. Biol. v, No. 3, pp. 85-92, 1 pl.
- 67. Contribution à l'étude des Spongiaires de l'Atlantique Résultats des Campagnes scientifiques accomplies sur son par Albert I., Prince souverain de Monaco, publiés sous sa dir avec le concours du Baron Jul. de Guerne. Fasc. II. Me 1892, gr. 4to, 165 pp., 11 pls, 2 maps gr. fol.

167 species obtained, 1 new family, 3 new subfamilies, 8 new s 58 new species.

- 68. VOSMAER, G. C. J. On the Canal System of the Homocala, 1 the morphological value of the terms osculum and pore in St Tijdschr. Nederl. Dierk. Ver. (2) iii, afl. 3 & 4, pp. 235-242.
- 69. WELTNER, W. Bericht über die Leistungen in der Spong für die Jahre 1885-87. Arch. f. Nat. liv, pp. 176, 227 & 228.
- 70. —. Die Tier- und Pflanzenwelt des Süsswassers. Refera das gleichnamige Werk von O. Zacharias. Naturwissenscha Wochenschrift, 1892, pp. 445 & 446.
- 71. —. Das Vorkommen von Cordylophora lacustris, Allm. bei SB. nat. Fr. 1892, No. 6, pp. 76-80.

Freshwater Sponges, p. 80.

- 72. —. Bemerkungen über den Bau und die Entwicklus Gemmulæ der Spongilliden. Biol. Centralbl. xiii, Nos. 4 & 119-126.
- 73. Metallausgüsse des Canalsystemes eines Süsswasserschwi SB. nat. Fr. 1893, No. 1, pp. 5 & 6, 1 woodcut
- 74. —. Ueber die Autorenbezeichnung von Spongilla erinaceus. pp. 7-13.

One new species.

- 75. —. Spongillenstudien. Arch. f. Nat. 1893, pp. 209-244. Gives the complete literature on Spongillidæ.
  - ---. [See also Korschelt & Weltner (33).]

- WIERZEJSKI, A. Ueber das Vorkommen von Carterius stepanovii, Petr. und Heteromeyenia repens, Potts, in Galizien. Biol. Centralbl. xii, No. 5, pp. 142-145. Abstract in J. R. Micr. Soc. 1892, pt. 3, p. 378.
- Wilson, H. V. Development of Sponges. J. R. Micr. Soc. 1892,
   p. 217. Non-sexual Reproduction in Sponges. Am. Nat. xxvi, pp. 626-628. Abstracts of the paper, by the same author, in J. Morph. v, pp. 511-519. [Vide Zool. Rec. 1891.]
- 78. Zacharias, O. Die biologische Station zu Plön. Zool. Anz. xv (1892), pp. 36-39, 1 woodcut.

The author recommends the station for the study of Freshwater Sponges, which are found there in enormous quantities.

- ZYKOFF, W. Die Entwicklung der Gemmulæ der Ephydatia fluviatilis, Auct. (Vorläufige Mittheilung). Zool. Anz. xv (1892), pp. 95 & 96.
- Die Entwicklung der Gemmulæ bei Ephydatia fluviatilis, Auct. Bull. Mosc. 1892, pp. 1-16, 2 pls. Abstract in J. R. Micr. Soc. 1892, pt. 3, p. 378, and (in Russian language) in Rev. Sci. Nat. St. Pétersb. ii, No. 9, pp. 342-344.
- Entwicklungsgeschichte von Ephydatia mülleri, Lieberk., aus den Gemmulæ. Eine biologisch-embryologische Skizze. Biol. Centralbl. xii, Nos. 23 & 24, pp. 713-716.
- 82. 2—. [Distribution of the Spongillidæ in European Russia. Journal Zool. Departm. & Zool. Museum, Moscow, ii, pp. 27-29.] [Russian?].

# II.—ANATOMY, EMBRYOLOGY, PHYSIOLOGY, BIOLOGY.

## (A.) ANATOMY, HISTOLOGY, MORPHOLOGY, AND PHYLOGENY.

BIDDER (3) describes some observations which he thinks prove that the "Metschnikoff" cells and glandular ectoderm cells of Ascetta have an excretory function. After the Sponge had been left in a solution of indigo-carmine, in sea water, the granules normally present in those cells became replaced by dark-blue granules. From the Metschnikoff cells (which extend throughout the whole thickness of the Sponge) the granules are probably discharged, both into the gastral cavity and on to the ectodermal surface. A completely different set of granules is found in the basal portion of many Sponges (basal spherules of the endoderm)—they are possibly stores of nutritive matter. This basal portion (plinth) may break off, and then it is used as food by wandering cells (generative elements). The life-history of the collar-cells, and of the Metschnikoff cells is now given. The author suggests, as a working hypothesis, that

the granules in the ectodorm and in the Metschnikoff cells of A a nitrogenous excretion, common to the protoplasm in all Sponge varying in their degree of solubility. In some Sponges (Halis excretion forms a slimy covering, in others a horny cuticle, as in Horny Sponges, it has been retained within their bodies until a spongin skeleton.

BILDER (4) is able to show that the ectoderm cells of the Sponges are of the same form and character as those of the July, of T shape, forming roughly hexagonal areas on the surfacentre of which the flask-shaped cell-bodies are attached. The lies at the base of the pendant cell-body. The spongoblasts tinuous with the ectoderm, and are of the same character. To nothing more than the concentrated cuticle of a large number cells, poured out around an intrusive foreign object. This characteristic paper on the histology of Ascetta clathrus, and thinks that theory of a contractile ectoderm not well tenable.

DENDY (11), who last year published the Calcarea Homocæla instalment of his proposed Monograph of the Australian Spunfortunately been prevented, by the reduction of the Go Grant to the Royal Society of Victoria, from continuing his M as intended, and gives now merely a synopsis of the Calcarea E (For a proposed new classification of that group, see Systemat this Record.)

DENDY (12) gives a preliminary account of Synute pulche genus and species of Calcareous Sponges. The Sponge is refrom the fact that it consists of a colony of Sycon individual pletely fused together into a single mass, invested in a common

DODERLEIN (14) describes a most remarkable Calcareous Sponstoma schulzei, n. g. & sp.), representing a new order (Lithon the Sagami Bay, Japan. There are two layers of spicules in th—an outer loose one, composed of triacts and tetracts, and layer which, in the young condition, consists of somewhat tetracts. These latter tetracts soon fuse together, and form a ous network, similar to that of the Lithistidæ. The author e the fact that this Sponge is different to the fossil Pharetronæ the spicules are held together by a foreign substance. It is gretted that no illustrations are given of this curious form.

HINDE & HOLMES (27) describe the Sponge-remains in the Tertiary strata near Oamaru, Otago, New Zealand. The mat sisted exclusively of detached spicules. This was the case not Monaxonid and Tetractinellid Sponges, but also with Lithistid more strange, with Hexactinellid Sponges.

von Lendenfeld (42) gives a detailed description of the He of the Adriatic, viz., Halisarca dujardini, Aphysilla rosea, A. Darwinella aurea. As in former papers, he divides the Silicee subclasses: Triaxonia (with large sack- or pocket-like, simple or



chambers), and Tetraxonia (with small, spherical or oval simple chambers). The Triaxonia comprise two groups: the Hexactinellida (with siliceous skeleton), and the Hexaceratina (with horny or no skeleton). The Hexaceratina consist of three families: (1) Darwinellidae, with horny fibres and horny spicules; (2) Aplysillidae, with horny fibres only; (3) Halisarcidae, without skeleton. This last family is possibly not a natural one, its two genera—Halisarca and Bajulus—are perhaps of different origin.

MAAS (44) discusses the question as to whether the three layers, of which adult Sponges are generally considered to consist, are derived from three or two germinal layers. Recent researches—histological (Topsent. Minchin), and embryological (Schulze)—point towards two germinal layers. The author's observations on the development of Esperia confirm such a theory. There are two layers in the larva of Esperia. The one is made of slender flagellated cells, and covers the anterior and lateral portions of the larva; the other is much more bulky, forms the central and posterior mass, and, even at this stage, contains spicules. Whilst the former of the two layers gives rise only to the endoderm, the latter forms all other tissues of the Sponge. It consists of two kinds of cells in the larval stage. The one kind has a granular protoplasm, and forms, later on, the amœboid wandering cells; the other kind of cells has a homogeneous protoplasm, their nuclei show an intranuclear network; these cells form the "ectoderm" and the "cellules contractiles" of Topsent; and, last of all, the glandular cells, which secrete the spongin. Thus we have two germinal layers, which may be called "Ectomesoderm" and "Endoderm."

MARSHALL (46) describes a new form of spicules from Foliolina lendenfeldii, n. sp., called Gomphostyles, from their nail-like shape. The head of the spicule is flattened out at right angles to its axis. The axis is twisted.

MINCHIN (49) describes the ectoderm of Leucosolenia cluthrus as consisting, in the expanded state, of flattened, non-ciliated cells. The cells, when contracting, assume a mushroom form, the nucleus remaining in proximity to the limiting membrane. The pores originate in certain ectoderm cells, which grow inwards until they join the endoderm cells, thus forming a complete connection between ectoderm and endoderm. These cells then become perforated. The fully-formed pore, therefore, lies in one single cell. Mesoderm and endoderm are also discussed.

MINCHIN (51) describes a sieve-like membrane across the oscula of Leucosolenia (probably L. coriacea), which consists of two layers of cells separated by a small amount of jelly. The outer layer of cells is continuous with the ectoderm, the inner with the endoderm. The sieve-like membrane is probably formed by a breaking through of the gastral cavity to the exterior. The author was not able to see distinct ectoderm cells, and thinks that in the fully-formed Sponge they may degenerate into a cuticle-like structure. In the mesoderm he noticed irregular granular

oder other as commises mentical with Topment's feelinies by Manager Topment - excited property.

Review 52 m incovered any and distinct medic in Locolors, what we increase anymaid in the important on. The sec is that it means it is appeared for recruitive rolls of which is terms. Institute Harmel's hour varieties of the inputs on ments in informations of comments of the same form. The most is many-severed encourage is made by instruction forms.

Later 40 immune his Polympione, a minute group of Lauren. He made that they in not constrainly differ from terms that he woulded Polympione flow is a secondary perfect or restriction, and that he question of flow Springer were of not not appear by stronger, or by may after connecting schools. Later 47 spectrum of the authors of the Laurencholists, opposite of Later, below, and Haute, and manusians that the secondary is the authors of the authority of the secondary operator.

Towers of entered finance; paper Note in a Serve-like some on , and is compared for finance families the ecode appropriate may a small of enterior as in intensit mover experient infentity is temperaturing four sufficiently and Minchia in a Lambert some for the approximate of Lambert are inversed by clinical laws. I operat further says four Minchia in his remarks measured, and remarks are inversed by clinical laws. I operat further says four Minchia in his remarks measured, and remarks of elements, the spherical colle which his function of interior reservoirs. N.R. Minchia was well aware takes, that he measurement in the examination of measurements.

Treat it summinues at impurate memoir to car knowl the Springes of the North Edward, one of the Results of the S Lapsditions of Libert L. Prince of Minutes. The volume is go princery struct. Its chief value has it Systematic and Geog Distribution.

On p. ? Topsens gives a summary of the listolity of Sponge occident comments of one kind of only only (pinneceptes. Sollas are flat. contractile. non-crimical (with a few exceptions, as O blockers. Pierces monologies. de., which are ciliated). The en appears in two forms: in contractile cells (similar to ectoderm cel in chosmocytes. The muscular elements, or myocytes, are possi derived from endoderm. Finally, the mesoderm consists of two l cells: (1) Granular cells, of variable shape, amorboid, and colourle young, pigmented when full-grown, corresponding to Sollas's i and mreencytes, called by Topsent "pigmented digestive cells." have also reproductive function. (2) Non-granular, spherica cells, corresponding to Sollas's chondrencytes, collencytes, and cysticated the corresponding to Tetractinellida. Normally they store up

materials (thesocytes). Finally, as they serve also as connective elements, they correspond, perhaps, in part to Sollas's inocytes. The pigment of Sponges may be in the granular mesoderm cells, or in the flagellated cells. Sollas's chromatocytes, therefore, do not correspond to a special kind of cells. The æsthocytes have probably the function suggested by Sollas.

Vosmaer (68) has a paper on the canal system of Homocæla, and the morphological value of the terms osculum and pore. According to him, the main excurrent tubes of Leucosolenia stipitata, L. tripodifera, of a Sycon, and of Hexactinellida, are homologous, and are the simplest modifications of the original Olynthus-sack without diverticula. The central cavity of these Sponges he calls cloaca, and its aperture osculum. Procts are the openings of canals or lacunæ into the cloaca. Proctions are the numerous small openings in Sponges like Cydonium gigas, where the excurrent canals are covered by a sieve-like membrane. Stomions are the apertures in the dermal membrane of Cydonium. A group of stomions corresponds to one stoma.

### B. EMBRYOLOGY AND ASEXUAL DEVELOPMENT.

Delage (9), who last year gave a short preliminary account of the development of Sponyilla fluviatitis now gives a detailed description of that species, and also of Esperella sordida, and Aplysilla sulfurea. Last year's paper was received with grave doubts by Minchin (Zoological Record 1891) and Maas (see this Record, p. 12), and now Spongologists will probably not be more ready than before to accept an account which, as Delage himself says, is "sans analogue dans les autres classes du règne animal." The larvæ of the above Sponges are almost entirely opaque, and therefore Delage's accounts of the extraordinary movements of the amæboid cells, &c., were based upon sections only. But, although his plates seem faithful, still their interpretation needs confirmation by other workers. In the meantime, Maas's account of the development of Esperella recommends itself far more by its simplicity.

Delage distinguishes four kinds of elements in the larva of Spongilla fluviatilis: (1) ciliated, (2) epidermal, (3) amœboid, (4) intermediate cells. The ciliated cells cover the larva anteriorly and laterally. The epidermal cells are large, and form a discontinuous layer beneath the ciliated cells. [N.B. The author calls these cells "epidermal" throughout the text, but "ectodermal" in the explanation of plates.] The amœboid cells are very large, and constitute the chief mass of the body. Some of them are specialized to form the spicules. Finally, the intermediate cells stand between the epidermal and amœboid cells. The ciliated cells soon loose their cilia, become irregular in shape, separate from each other, and sink beneath the epidermal cells. The epidermal cells now come to lie along the surface and soon form a continuous epidermis. Then the amœboid cells become active, throw out their pseudopodia and capture the former ciliated cells, each being able to incorporate several of them

After that, the amœboid cells retract their pseudopodia and assured nearly spherical shape, having a large nucleus in their centre, and are it about twenty small nuclei, derived from the captured cells. Thes Delage's "groupes polynucléés." Gætte and Maas regarded those a nuclei as yolk-granules. After a pause the "groupes polynucleecome active once more, throw out pseudopodia, move somewhat a from each other, still being connected together by their pseudopodia thus form an irregular mesh-work. Finally, they approach each and arrange themselves in circles, leaving a small cavity in their n Now the cilated cells separate from the "groupes polynucléés," arrange themselves so as to form the lining to those cavities. In way the flagellated chambers are formed. After that, the formatic pores, oscula, &c., is described. The development of Esperella so and Aplysilla sulfurea agrees in its chief features with that of Spon

MAAS (43) describes the metamorphosis of Esperia lorenzi, O. S. begins his account at the stage when the larva is ready to leave the p Sponge. It is then seen to be covered anteriorly and laterally t exceedingly narrow columnar ciliated epithelium. Its posterior pole i ciliated, and there can be made out, only on the closest examination, epithelium continuous with the former. The nuclei of the ciliated are larger than the average diameter of the cells. Accordingly they a bulging out, and, in order to find sufficient space, arrange themselv different levels, thus causing an appearance of several layers of n The internal mass of the larva [N.B. Mass avoids speaking of gen layers.] consists of two kinds of cells. The one kind is usually round a coarse granular protoplasm, and a clear spherical nucleus. The is of a spindle-shaped or amorboid form, has a homogeneous protor and an oblong nucleus with delicate intranuclear network. ment of the spicules is very instructive : it apparently serves a d purpose, (1) to utilise all space as much as possible, (2) to balance the equally when swimming. The megascleres lie in a dense bundle i axis of the larva. Around the posterior end of that bundle, the ro are equally distributed, and, around its anterior end, the toxa. The period is only of short duration (from six to twenty-four hours), which the larva settles down with its anterior pole, and becomes some flattened. Also here, as in the larva, we find two kinds of tissue: cells with small nuclei, apparently derived from the ciliated cells, but lying ventrally and internally; and larger cells, derived from the mass of the larva, lying now dorsally and externally. The young Sp opaque at first, shows after the first day, here and there, lighter s which, in sections, are seen to be subdermal cavities, ciliated chan and exhalant canals, in process of development. In regard t development of the subdermal cavities, Maas agrees with Delage partially with Wilson, but not so in regard to the ciliated chambers. cells with granular contents [groupes polynucléés, Delage] have no to do with the formation of the chambers. Independently of those the small cells, formerly the ciliated cells of the larva, group and ar

themselves so as to form the chambers. Some of them also form the exhalant canals. The two layers of tissue already mentioned, become the ectosome and choanosome of the adult Sponge. The last few pages of the paper are devoted to the further development of the Sponge, including skeleton, tissues, and oscula.

WELTNER (72) has some notes on the structure and development of the gemmules of Spongillidæ. The "Anlage" of the gemmule, and all later stages up to the time when the inner cuticle begins to be formed, consist of naked, amœboid, uninuclear cells. In the fully-developed gemmule, the cells of the inner mass are surrounded by a thin membrane, and show no longer an amœboid movement. Generally, these cells are of two sizes, small uninuclear and large binuclear. Before the contents of the gemmule prepare themselves to leave the capsule, its cells become again amœboid, and then they possess one, two, three, or four nuclei. Weltner says that Zykoff was not able to see any nuclei in the cells of the gemmules, because he stained whole pieces of Sponge instead of the sectionized gemmules, and the staining fluid is never able to penetrate the capsule. Nor did he examine the gemmules in the living condition. Little is known as to how the "Anlage" of the gemmule is formed. There are four possibilities: (1) Either it is formed from a single cell, which has the morphological value of an ovum: in this case the gemmule would be a morula; (2) or it is derived from several precisely similar mesodermal cells; (3) or from several heterogeneous cells of the messderm; (4) or, finally, the gemmule is formed from the cells of (two or three) germinal layers. In this case the gemmule would be a bud.

ZYKOFF (80), after a review of the literature on the subject, gives an account of his own researches on the formation of the gemmules in Ephydatia fluviatilis. Between the ordinary amoeboid cells of the parenchym he noticed cells containing boat-shaped, shining granules. granules are possibly a modification of albumen, and serve as reservestores, and the cells, containing the same, may be called yolk-cells. first appearance of these cells is to be regarded as the first moment in tho development of the gemmule. They, together with similar cells, which, however, have no yolk-granules, begin to show a creeping movement, and at the same time they increase in number. This increase is possibly not caused by division, but by other cells being attracted to the same centre. Contrary to Gotte's statements, neither the canals nor the chambers take any part in the formation of the gemmules. Soon a division of the gemmule into two layers may be noticed: (1) a central mass, consisting of yolk-cells, and a few amœboid cells from the mesenchym; (2) a peripheral layer, consisting of several concentric rows of yolkless cells of the mesenchym, which pass over into the mesenchym of the Sponge. These peripheral cells gradually assume a club-shaped form, the thicker end of the cell lying outwards—and arrange themselves in a single layer. One part of the gemmule remains free from these cells: the hilus. shaped cells are disc-like, flattened out on their inner ends, and there they begin to secrete a chitinous cuticle. The amphidises are formed outside

the cuts-ampai cale, and only afterwards, pushing by analogid coputers returned the same, and assume a califul accompanion. Finally, it dain-amped calls have their position, cours on the estable of the amp them, and form there the second distincts captured. The generale is a fully farmed. The circu-imped calls are gradually absorbed. Lykoffments to use the second mass of the generale as consisting of polygcals. Todoward, we as see model in them (Kornchelt & Heider). On sections, it appeared line a samples, disting, granular substance, with outlines of salls or master. The model are hidden by the yolk. [ Wettness, who fifteen greatly segmenting this latter points.]

Let 197 11 studied the development of Ephydatic milleri, from gustion which had been lying dry for about two years. Rather now the meananest explanation which he gives of the formation of on The young Ephydatic swims upon the surface of the water, its ade being directed apwards, he convex side downwards. Any waters up by it collects near the centre of the convex surface, p flows the excelerate, and family breaks through it. The chambers developed from certain cells of the parenchym cells, which divide and familiary through the chamber thus formed all lies in the neighbourhood of canals, and goes on increasing in size some time, it is periodole that the connection between chamber and c is formed by a breaking through.

# c PHYSIOLOGY AND BIOLOGY.

For the physiology of Sponges, see GRIFFITHS (20) pp. 30-32 & 79 184, 200, 349, & 378.

Hornell (26) describes a case of commensalism of Microciona plun and an Annelld (Lucedora caca).

BIDDER (3, on excretion in Sponges, see Anatomical, p. 7.

#### III.—DISTRIBUTION.

# (A.) FAUNISTIC.

Atlantic, especially Bay of Biscay, Newfoundland, and the  $\Delta z$  Topsent (67).

Portugal: NOBRE (55).

Great Britain: Plymouth; GARSTANG (18): Puffin I. (North Wa HERDMAN (24).

Norway: West Coast; Herdman (25): Fjord of Bergen; Appel (2).

Russia: Solowezki Is. (White Sea); Chwobostansky (17): Sp. wia; Zykoff (82).

Germany: Lakes near Plön, in Holstein, Spongillidæ; ZACHARIAS (78): Spongillidæ in the neighbourhood of Berlin; Weltner (71).

Galizia: Lubien, Carterius stepanovii, Petr., and Heteromeyenia repens Potts. The latter species, of which, however, the gemmulæ only were found, was previously not taken outside North America. Wienzejski (76).

Adriatic: Hexaceratina; LENDENFELD (42).

Palestine: Lake Tiberias, Potamolepis barroisi, n. sp.; TOPSENT (66).

Japan: Sagami Bay, 100 faths., Petrostoma schulzei, n. g. & sp., representing a new order (Lithones) of Calcarea; Döderlein (14).

Australia : Calcarea Heterocæla ; DENDY (11).

# (B.) GROLOGICAL.

Cambrian: St. John, New Brunswick; MATTHEW (47).

Ordovician: Palæosaccus dawsoni, n. g. & sp. (Hexactinellid), from Little Métis, Quebec, Canada; HINDE (26).

Trias: Saxe-Altenburg; Geinitz: St. Cassian, Strata of Southern Tyrol; OGILVIE (57).

Cretaceous (Upper): from Meaulne in the département Maine-et-Loire, France. Lithistidæ only, no Hexactinellida. The absence of the latter can be accounted for by the fact that, amongst recent Siliceous Sponges also, the Lithistidæ live in a different depth (up to 350 faths.) from the Hexactinellidæ (up to 600 faths.). Počta (59).

Tertiary: Lower Tertiary Strata, near Oamaru, Otago, New Zealand. Monaxonida, 70 species; Tetractinellida, 22 species; Lithistida, 7 species; Hexactinellida, 11 species. Many of these Sponges have their nearest existing allies living in the Indian Ocean and in the Australian Seas; but amongst others the genus Guitarra, Carter, has hitherto only been found in the Gulf of Mexico and the North Atlantic. Remarkable is the preponderance of Monaxonida, as generally the proportions have been reversed. In the Oamaru deposit, the delicate spicules of Monaxonida have been preserved equally as well as the larger spicules of the other groups. "The absence of these Sponge-spicules in the older rocks, is rather due to their having perished in the fossilization, than that they did not coexist with those other groups whose remains have been in part preserved." HINDE & HOLMES (27).

#### IV.—SYSTEMATIC AND CLASSIFICATION.

#### 1. CALCAREA.

VON LENDENFELD (41) discusses the Calcarea Homocela described by DENDY (Monograph of the Victorian Sponges, Part I), and substitutes his own nomenclature, viz.:—Grantia labyrinthica remains in Grantia, Lendenfeld; while Leucosolenia asconoides, canata, dubia, stolonifer, and ventricosa are referred to Ascandra; lucasi, pelliculata, pulcherrima, stipitata, and tripodifera to Ascetta; wilsoni to Leucopsis; depressa and protogenes doubtfully to Leucopsis or Ascetta.

DENDY (11) gives the following classification of the *Heterocæla*:—
Order HETEROCŒLA.

Family 1. Leucascidæ.

Genus. Leucascus, n. g.

Family 2. Sycettidæ.

Genera. Sycetta (Hæckel, emend.); Sycon (Risso, emend.) Sycantha, v. Lend.

Family 3. Grantidæ.

Genera. Grantia (Fleming, emend.); Grantiopsis, n. subj Ute (Schmidt, emend.); subg. Synute, Dendy Utella, n. g.; Anamizilla (Poléjaeff, emend. Sycyssa, Hæckel; Leucandra (Hæckel, emend.) Lelapia (Gray, emend.); Leucyssa, Hæckel.

Family 4. Heteropidæ.

Genera. Grantessa (v. Lend., emend.); Heteropia (Cart emend.); Vosmaeropsis, n. g.

Family 5. Amphoriscidæ.

Genera. Heteropegma (Poléjaeff, emend.); Amphoria (Hæckel, emend.); Syculmis (Hæckel, emend Leucilla (Hæckel, emend.).

Grantia vosmaeri, Port Jackson, DENDY, P. R. Soc. Vict. (n.s.), v, p. 88, n. Grantiopsis, n. g., p. 73, for G. cylindrica, n. sp., Port Phillip, p. DENDY, P. R. Soc. Vict. (n.s.), v.

Grantessa hispida, Port Phillip, DENDY, P. R. Soc. Vict. (n.s.), v, p. 1 n. sp.

Leucandra carteri, p. 103, gladiator, p. 101, phillipensis, p. 100, utesi p. 178, Port Phillip, DENDY, P. B. Soc. Vict. (n.s.), v, n. spp.

Leucascus, n. g., p. 72, for L. simplex, p. 77, clavatus, p. 78, n. spp., Phillip, Dendy, P. R. Soc. Vict. (n.s.), v.

Lithones, new order; Döderlein, pp. 143-145. For the reception of single form, Petrostoma schulzei.

Petrostoma, n. g, for P. schulzei, n. sp., 100 faths., Sagami Bay, Ja DÖDERLEIN, Verh. Deutsch. Zool. Ges. ii, pp. 143-145.

Sycon carteri, St. Vincent's Gulf, p. 79, minutum, Port Jackson, p. ensiferum, p. 81, boomerang, p. 82, Port Phillip, giganteum, St. Vince Gulf, p. 84, DENDY, P. R. Soc. Vict. (n.s.) v, n. spp.

S. gelatinosum, var. whiteleggii, p. 84, Port Jackson, Dendy, P. R. Vict. (n.s.) v, n. var.

Synute, n. g. for S. pulchella, n. sp., p. 2, Port Phillip; DENDY, P. R. Vict. (n.s.) iv.

Ute spiculosa, p. 92, spenceri, p. 94, Port Jackson, DENDY, P. R. Soc. (n.s.) iv, n. spp.

Utella, n. g., DENDY, P. R. Soc. Vict. (n.s.) iv, p. 74. Type, Syca hystrix, Hæckel.

Vos naeropsis, n. g., p. 76, for V. depressa, p. 110, wilsoni, p. 111, n. Port Phillip; DENDY, P. R. Soc. Vict. (n.s.) iv.

#### 2. HEXACTINELLIDA.

- Chonelasma schulzei, Topsent, (67) p. 33, pl. iii, fig. 3, & pl. vii, fig. 3, Atlantic, lat. 33° 26′ 33″ N., long. 33° 23′ W., depth 1557 metres, n. sp.
- Hertwigiidæ, n. fam., TOPSENT, (67) p. 25, established for the reception of Hertwigia falcifera, O. S., and Trachycaulus gurlittii, F. E. Schulze. Bhabdodictyum delicatum, O. S., comes close to it.
- Hexactinella grimaldii, TOPSENT, (67) p. 34, pl. ii, figg. 1 & 2, & pl. vii, fig. 1, Atlantic, lat. 38° 33′ 21″ N., long. 30° 28′ 54″ W. (between Pico and San Jorge), depth 1300 metres, n. sp.
- +Palæosaccus, n.g., p. 56, for P. dawsoni, n. sp., Shales of the Quebec Group, p. 57, pl. iv; HINDE.
- Periphragella lusitanica, Topsent, (67) p. 31, pl. vi, fig. 4, & pl. vii, fig. 2, Atlantic, lat. 39° 22′ 48″ N., long. 33° 45′ 30″ W., depth 1384 metres, n. sp.
- †Purisiphonia clarkei, Bowerb., shown to be a Lyssakine Hexactinellid; JACK & ETHERIDGE, (29) pp. 436-439, pl. xix.

# 3. TETRACTINELLIDA (with CHONDROSIDÆ, PLAKINIDÆ, HALISARCA, and OSCARELLA).

- Astrella tuberosa, TOPSENT, (67) p. 44, pl. ii, fig. 4, & pl. viii, fig. 4, Atlantic, lat. 39° 1′ 40″ N., long. 30° 15′ 40″ W., depth 454 metres, n. sp.
- †Astrocladia fratectosa, n. sp., Pocta, MT. min. geol. Mus. Dresden, xi, p. 19, pl. ii, figg. 4 & 5, Upper Cretaceous, Meaulne, France.
- Calthropella simplex, Sollas, var. durissima, n. var.; TOPSENT, (67) p. 43, pl. v, fig. 14, and pl. viii, fig. 9, Atlantic, lat. 39° 1′ 40″ N., long. 30° 15′ 40″ W., depth 454 metres.
- Characella sollasi, n. sp., TOPSENT, (67) p. 40, pl. ii, fig. 3, & pl. viii, fig. 6, Atlantic, lat. 43° 57′ N., long. 9° 27′ W., depth 300 metres; lat. 38° 22′ N., long. 30° 34′ 39″ W., depth 736 metres; and lat. 39° 1′ 40″ N., long. 30° 15′ 40″ W., depth 454 metres.
- †Chenendopora batillacea, p. 7, pl. ii, figg. 1-3, conferta, p. 8, pl. iv, figg. 17-2.), radicata, p. 10, pl. i, figg. 13-16, & pl. ii, fig. 16, C. P scutula, p. 9, pl. iii, figg. 12-14, Upper Cretaceous, Meaulne, France, Pocta, MT. min. geol. Mus. Dresden, xi, n. spp.
- Discodermia ramifera, n. sp., TOPSENT, (67) p. 50, pl. vi, fig. 10, & pl. viii, fig. 1, Atlantic, lat. 38° 23′ 30″ N., long. 30° 20′ 20″ N., depth 318 metres.
- †Ditricenella, n. g. for D. oamaruensis, n. sp., Lower Tertiary, New Zealand, Hinde & Holmes, J. L. S. xxiv, p. 232, pl. xii, figg. 34 & 35.
- Erylus nummulifer, n. sp., Topsent, (67) p. 47, pl. iii, fig. 14, & pl. viii, fig. 5, Atlantic, lat. 38° 22' N., 30° 34' 39" W., depth 736 metres, and lat. 38° 23' 30" N., long. 30° 20' 20" W., depth 318 metres.

- Hulisarca dujardini, redescribed, pp. 292-296, pl. xiii, figg. 25-27, dujardini, var. incrustans, n. var., H. dujardini, var. massa, n. va p. 293, Triest; Lendenfeld, Z. wiss. Zool. liv.
- † ? Isoraphinia gibbosa, n. sp., Pocta, MT. min. geol. Mus. Dresden, p. 11, pl. iii, fig. 1: pl. iv, figg. 4 & 5, Upper Cretaceous, Meaulr France.
- † Jerea clavata, pl. iv, figg. 10-13, J. ? caulis, pl. iii, figg. 8 & 9, p. 1 Upper Cretaceous, Meaulne, France; Pŏcta, t. c.: n. spp.
- †Jereica permiru, n. sp., Pocta, t. c. p. 10, pl. ii, figg. 9-13, Upp Cretaceous, Meaulne, France.
- Pachastrella debilis, n. sp., TOPSENT, (67) p. 42, pl. iii, fig. 8, and pl. vi fig. 8, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 7: metres, and lat. 38° 23' 30" N., long. 30° 20' 20" W., depth 3: metres.
- †Plakina australis, n. sp., HINDE & HOLMES, J. L. S. xxiv, p. 230, pl. x fig. 20, Lower Tertiary, Oamaru, Otago, New Zealand.
- †Polyjerea indistincta, n. sp., POCTA, MT. min. geol. Mus. Dresden, p. 18, pl. iv, figg. 2 & 3, Upper Cretaceous, Meaulne, France.
- †Siphonia piriformis, var. acuta, n. var.; Pŏcta, t. c. p. 13, pl. i figg. 7-9, Upper Cretaceous, Meaulne, France.
- †Spongodiscus tuber, n. sp., Pocta, t. c. p. 21, pl. i, figg. 2-4, Uppe Cretaceous, Meaulne, France.
- Tetilla truncata, n. sp., Topsent, (67) p. 36, pl. viii, fig. 7, Atlantic, ls 43° 40' N., long. 8° 55' W., Coast of Asturia, 134 metres.
- † Triptolemus australis, n. sp., HINDE & HOLMES, J. L. S. xxiv, p. 23 pl. xiii, fig. 34, Lower Tertiary, near Oamaru, Otago, New Zealand
- †Vetulina oamaruensis, n. sp., HINDE & HOLMES, t. c. p. 240, pl. xii figg. 31-33, Lower Tertiary, near Oamaru, Otago, New Zealand.

#### 4. MONAXONIDA (+ Tethyada).

TOPSENT, (67) pp. 63-65, proposes the following classification of the Monaxonida (chiefly after Ridley and Dendy), in which, however, the Spongillida have been left out altogether.

# Order Monaxonida, Ridley & Dendy.

1. Suborder Halichondrina, Vosmaer.

Family 1. Homorrhaphidæ, Ridley & Dendy.

Subfamily 1. Chalinina, Ridley & Dendy.

Subfamily 2. Renierinæ, Ridley & Dendy.

Family 2. Heterorrhaphida, Ridley & Dendy.

Subfamily 1. Phlaodictyina, Carter.

Subfamily 2. Trachyinæ, n. s. f.

Genus Trachya, Carter.

Subfamily 3. Gelliinæ, Ridley & Dendy.

Subfamily 4. Tedaniinæ, Ridley & Dendy.

Subfamily 5. Desmacellinas, Ridley & Dendy.

Genera Desmacella, Schmidt, Biemma, Gray.

Subfamily 6. Hamacanthina, Gray.

Genus Hamacantha, Gray.

Family 3. Desmacidonidæ, Vosmaer.

Subfamily 1. Esperellinæ, Ridley & Dendy.

Genera: Esperella, Vosmaer; Esperiopsis, Carter; Cladorhiza, M. Sars; Axoniderma, Ridley & Dendy; Chondrocladia, W. Thomson; Desmacidon, Bowerbank; Joyeuxia, n. g.; Phelloderma, Ridley & Dendy; Sideroderma, Ridley & Dendy; Histoderma, Carter; Dendoryx, Gray; Forcepia, Carter; Melonanchora, Carter; Yvesia, n. g., Iotrochata, Ridley; Amphilectus, Vosmaer, &c.

Subfamily 2. Ectyoninæ, Ridley & Dendy.

Genera: Myxilla, Schmidt; Pytheas, n. g.; Clathria, Schmidt; Rhaphidophlus, Ehlers; Plumohalichondria, Carter; Stylostichon, n. g.; Microciona, Bowerbank; Hymeraphia, Bowerbank; Rhabderemia, n. g.; Plocamia, Schmidt; Suberotelites, Schmidt; Spanioplon, n. g.; Acurnus, Gray; Echinoclathria, Carter; Agelas, Duchassaing & Michelotti; Echinodictyum, Ridley.

Family 4. Axinellidæ, Ridley & Dendy.

2. Suborder Spintharophora, Sollas (emend).

1. Section Aciculida, n. s.

Family 1. Epallacidæ, new family.

Genera: Epallax, Sollas; Scolopes, Sollas; Dorypleres, Sollas.

Family 2. Stylocordylidæ, new family.

Genus Stylocordyla, W. Thomson.

Family 3. Tethyidæ, Gray.

2. Section Clavulida, n. s.

Family 1. Spirastrellida, Ridley & Dendy.

Genera: Spirastrella, Schmidt; Latrunculia, du Bocage; Hymedesmia, Bowerbank.

Family 2. Suberitidæ, Vosmaer.

Subfamily 1. Suberitine, n. s.

Genera: Suberites, Nardo; Weberella, Vosmaer; Poterion Schlegel; Osculina, Schmidt, &c.

Subfamily 2. Polymastine, n. s.

Genera: Polymastia, Bowerbank; Ridleya, Dendy; Quasillina, Norman; Tentorium, Vosmaer; Trichostemma, M. Sars; Halicnemia, Bowerbank; Tethyspira, Topeent.

Family 3. Clionida, Gray.

Genera: Cliona, Grant; Thoosa, Hancock; Alectona, Carter.

- Apatospongia, n. g. (Suberitidæ?), for A. fullax, n. sp., Australian Waters, MARSHALL, Leuckart Festschr., p. 16, pl. viii, figg. 1-5.
- Artemisina transiens, n. sp., Topsent, (67) p. 94, pl. iv, fig. 8, pl. ix, figg. 7 & 8, Atlantic, lat. 43° 44′ 50″ N., long. 8° 12′ W., depth 135 metres.
- Azinella flustra, n. sp., Topsent, (67) p. 122, pl. ii, fig. 5, & pl. xi, fig. 18, Atlantic, lat. 43° 40′ N., long. 8° 55′ W., depth 134 metres, and lat. 38° 23′ 30″ N., long. 30° 20′ 20″ W., depth 318 metres.
- Biemma grimaldii, p. 81, pl. iv, figg. 1-4, & pl. ix, fig. 14, Atlantic, lat. 38° 23′ 45″ N., long. 30° 51′ 30″ W., depth 927 metres; B. dautzenbergi, p. 83, pl. iii, fig. 5, & pl. ix, fig. 16, Atlantic, lat. 39° 22′ 48″ N., long. 33° 45′ 30″ W., depth 1384 metres, Topsent (67): n. spp.
- Cladocroce, n. g. (Lamelliform Revierinæ), for C. fibrosa, n. sp., p. 72, pl. iii, figg. 1 & 2, Atlantic, lat. 38° 33' 21" N., long. 30° 28' 54" W. (between Pico and San Jorge, Azores), depth 1300 metres; Topsent (67).
- †Cladorhiza haasti, n. sp., HINDE & HOLMES, J. L. S. xxiv, p. 210, pl. x, fig. 35, Lower Tertiary, Oamaru, Otago, New Zealand.
- Dendoryx, Gray. TOPSENT, (67) p. 97, proposes the following definition: "Esperellina generally without a definite shape; megascleres of the ectosome usually diactine (tylote, strongylote, or tornote), rarely monactine, normally smooth; megascleres of the skeleton invariably spined styli." The genus Dendoryx thus defined includes the genera Dendoryx, Gray, Alebion, Iophon, Hastatus, several species of Myxilla, &c. As types of the genus may be regarded: D. incrustans, Gray. D. luciensis, certa, and pectinata, Topsent.
- D. certa, p. 99, pl. x, figg. 7 & 8, Atlantic, lat. 43° 40′ N., long. 8° 55′ W., depth 134 metres; pectinata, p. 100, pl. x, fig. 6, Atlantic, lat. 38° 48′ 30″ N., long. 30° 19′ W., depth 861 metres; Topsent (67): n. spp.
- Desmacella aberrans, n. sp., TOPSENT, (67) p. 85, pl. ii, fig. 7, & pl. ix, fig. 10, Atlantic, lat. 43° 44′ 50″ N., long. 8° 12′ W., depth 135 metros, and lat. 43° 40′ N., long. 8° 55′ W., depth 134 metres.
- Dorypleres incrusturs, n. sp., Topsent, (67) p. 126, pl. ix, fig. 21, Straits of Pico-Fayal (Azores), depth 130 metres.
- Esperiopsis columnata, p. 90, pl. v, fig. 5, & pl. x, fig. 5, Atlantic, lat. 46° 4′ 40″ N., long. 49° 2′ 30″ W., depth 1267 metres; polymorpha, p. 90, pl. vi, figg. 1-3, & pl. x, figg. 1-3, Atlantic, lat. 38° 23′ 45′ N., long. 30° 51′ 30″ W., depth 927 metres, and lat. 38° 22′ N., long. 30° 34′ 39″ W., depth 736 metres; prædita, p. 92, pl. x, fig. 4, Atlantic, lat. 38° 22′ N., long. 30° 34′ 39″ W., depth 736 metres; Topsent (67): n. spp.
- Foliolina lendenfeldii, n. sp., MARSHALL, Leuckart Festschr. p. 2, pl. i, & pl. vii, figg. 1-7c, locality unknown.
- Forcepia versatilis, n. sp., TOPSENT, (67) p. 100, pl. vi, fig. 5, & pl. x, fig. 9, Atlantic, lat. 46° 4′ 40" N., long. 49° 2′ 30" W., depth 1267 metres.
- Telliodes cavicornis, p. 78, pl. iii, figg. 4 & 9, & pl. ix, fig. 12, fayalensis,

- p. 78, pl. iii, fig. 13, & pl. ix, fig. 11, Straits between Pico and Fayal (Azores), depth 130 metres, TOPSENT (67), n. spp.
- Gellius macrosigma, n. sp., Topsent, (67) p. 77, pl. ix, fig. 13, Atlantic, lat. 38° 22' N., 30° 34' 39" W., depth 736 metres.
- †Guitarra carteri, pl. xi, figg. 1-3, intermedia, figg. 4-7, p. 213, Lower Tertiary, near Oamaru, Otago, New Zealand, Hinde & Holmes, J. L. S. xxiv, n. spp.
- Halichondria pachastrelloides, p. 66, pl. ix, fig. 3, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 736 metres, leuconoides, p. 67, pl. v, fig. 4, Atlantic, lat. 38° 23' 30" N., long. 30° 20' 20" W., 318 metres, Torsent (67), n. spp.
- Hamacantha johnsoni, Bow., var. complanata, n. var., pl. ii, fig. 9, johnsoni,
  Bow., var. inelegans, n. var., pl. i, fig. 4, Atlantic, lat. 43° 44′ 50″ N., long.
  8° 12′ W., depth 135 metres; lat. 38° 26′ 25″ N., long. 30° 59′ 10″ W.,
  depth 800 metres, and lat. 38° 22′ N., long. 30° 34′ 39″ W., depth 736
  metres; Topsent, (67) p. 86.
- Hymeraphia tuberosocapitata, p. 113, pl. xi, fig. 6, Atlantic, lat. 38° 22 N., long. 30° 34′ 39″ W., depth 736 metres, minima, p. 114, figg. 2 & 3, Atlantic, lat. 38° 23′ 30″ N., long. 30° 20′ 20″ W., depth 318 metres, geniculata, p. 115, pl. 1, fig. 13, & pl.,xi, figg. 4 & 5, Atlantic, lat. 38° 23′ 30″ N., long. 30° 20′ 20″ W., depth 318 metres, Topsent, (67) n. spp.
- +Iophon hybridus, n. sp., HINDE & HOLMES, J. L. S. xxiv, p. 212, pl. x, fig. 44, Lower Tertiary, Oamaru, New Zealand.
- +Joyeuxia, n. g., p. 93, for J. viridis, n. sp., p. 94, pl. ii, fig. 8, & pl. x, fig. 19, Atlantic, lat. 38° 22′ N., long. 30° 34′ 39″ W., depth 736 metres, and lat. 39° 1′ 40″ N., long. 30° 15′ 40″ W., depth 454 metres; Torsent (67).
- †Lasiocladia? hindei, n. sp., ETHERIDGE, (29) p. 199, pl. xli, figg. 1 & 2. Latrunculia biannulata, p. 127, Atlantic, lat. 38° 22'N., long. 30° 34' 39" W., 736 metres; insignis, p. 128, pl. viii, fig. 11, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 736 metres; Topsent, (67), n. spp.
- †L. oamaruensis, p. 218, pl. xi, fig. 34, obtusa, p. 220, fig. 32, Lower Tertiary, Oamaru, New Zealand, HINDE & HOLMES, J. L. S. xxiv, n. spp.
- †Melonanchora morlandi, n. sp., HINDE & HOLMES, J. L. S. xxiv, p. 200, pl. ix, fig. 38, Lower Tertiary, Oamaru, New Zealand.
- Myzilla pecqueryi, n. sp., Topsent, (67) p. 110, pl. x, fig. 8, Atlantic, lat. 39° 22′ 48″ N., long. 33° 45′ 30″ W., depth 1384 metres.
- Petrosia friabilis, n. sp., Topsent, (67) p. 69, pl. vi, fig. 8, & pl. ix, fig. 4, Atlantic, lat. 43° 40′ N., long. 8° 55′ W., depth 134 metres; lat 43° 57′ N., long. 9° 27′ W., depth 300 metres; lat. 38° 23′ 45″ N., long. 30° 51′ 30″ W., depth 927 metres, Straits between Pico and Fayal (Azores), depth 130 metres.
- Plumohalichondria, Carter. Topsent, (67) p. 111, proposes to break up this genus into two genera, according to whether the ascending fibres are made up of smooth diactine spicules or of spined monactine spicules. The former species are included under the genus Plumo-

halichondria, the latter constitute the new genus Stylostichon (see below). The genus Plumohalichondria is accordingly defined Ectyonina, the main skeleton of which consists of plumose ascending fibres. The spicules of the true main skeleton are smooth diactines. Type, Plumohalichondria mammillata, Carter.

- Polymastina, n. subfam. Suberitida; q.v. Topsent (67).
- Potamolepis barroisi, n. sp., Topsent, Rev. Biol. v, pp. 85-92, 1 pl., Lake Tiberia, depth 8 metres.
- †Pseudohalichondria deformis, p. 214, pl. xi, figg. 8 & 9, camaruensis, p. 215, figg. 12 & 13, Lower Tertiary, Oamaru, New Zealand, Hinde & Holmes, J. L. S. xxiv, n. spp.
- Pytheas, n. g. for P. ater, n. sp., p. 110, pl. i, fig. 1, & pl. xi, fig. 1, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 736 metres, and including Hymedesmia inflatus, Bow.; Topsent (67).
- Raspailia (Syringella) humilis, p. 123, pl. v, fig. 11, & pl. xi, fig. 20,
  Atlantic, lat. 38° 23′ 30″ N., long. 30° 20′ 20″ W., depth 318 metres,
  falcifera, p. 124, pl. vi, fig. 6, & pl. xi, fig. 19, Atlantic, lat. 38°
  33′ 21″ N., long. 30° 28′ 54″ W. (between Pico and San Jorge), depth
  1300 metres, Topsent, (67), n. spp.
- Reniera filholi, n. sp., TOPSENT, (67) p. 70, pl. iv, fig. 7, & pl. ix, fig. 6, Atlantic, lat. 43° 40' N., long. 8° 55' W., depth 134 metres.
- Rhabderemia, n.g., p. 115, for R. guernei, n. sp., p. 116, pl.xi, fig. 7, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 736 metres, including also Microciona pusilla and intexta, Carter; TOPSENT (67).
- Rhisochalina elongata, n. sp., Topsent, (67) p. 75, pl. iv, figg. 5 & 6, & pl. ix, fig. 1, Atlantic, lat. 38° 23′ 30″ N., long. 30° 20′ 20″ W., depth 318 metres.
- Spanioplon, n. g., p. 116, for S. fertile, n. sp., p. 117, pl. xi, fig. 11, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 736 metres, Tor-SENT (67).
- Spheciospongia, n. g. for Spheciospongia (Alcyonium) vesparum, Lamarck, (Suberitidæ), Marshall, Leuckart Festschr. p. 32.
- Spirastrella aculeata, n. sp., Topsent, (67) p. 127, pl. viii, fig. 10, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 736 metres.
- Suberitidæ, redefined by Topsent, (67) p. 59: Clavulida without microscleres derived from an aster. Two subfamilies: (a) Suberitinæ: Suberitidæ, without radiating structure, including the genera Suberites, Weberella, Poterion, Osculina, &c.; (b) Polymastinæ: Suberitidæ, the skeleton of which normally assumes a radiating structure, including the genera Polymastia, Quasillina, Tentorium, Trichostemma, Halicnemia, Tethyspira.
- Suberites capillitium, n. sp., TOPSENT, (67) p. 130, pl. ii, fig. 10, & pl. ix, figg. 19 & 20, Atlantic, lat. 46° 4′ 40″ N., long. 49° 2′ 30″ W., depth 1267 metres.
- Subsrotelides demonstrans, n. sp., Topsent, (67) p. 118, pl. iv, fig. 9, & pl. xi, figg. 15-17, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 736 metres.



- Stylinos, n. g., (= Desmacidonidæ), p. 135, for Desmacidon pannosus, Bwk., incognitus, B, Rhaphiodesma simplicissimum, B., Isodictya uniformis, B., Desmacidon columella, B., and including jullieni, n. sp., Fayal, p. 137, pl. i, fig. 12, & pl. vii, figg. 11 & 12; TOPSENT (67).
- Stylostichon, n. g., p. 111, for Spongia plumosum, Mont.; Myxilla frondosum, R. & D.; Plumohalichondria microcionides, Carter, and dendyi, n. sp., p. 112, pl. ii, fig. 6, pl. xi, figg. 9 & 10, Atlantic, lat. 38° 23′ 30″ N., long. 30° 20′ 20″ W., depth 318 metres, Topsent (67).
- Tedania conuligera, n. sp., TOPSENT, (67) p. 79, pl. i, fig. 16, Atlantic, lat. 46° 50′ 6″ N., long. 50° 11′ 45″ W., depth 155 metres.
- +Thoosa hancocki, n. sp., HINDE & HOLMES, J. L. S. xxiv, p. 222, pl. xi, fig. 41, Lower Tertiary, Oamaru, New Zealand.
- Trachyinæ, n. subfam. (not defined) of the Heterorrhaphidæ, for Trachya, Carter; Topsent, (67) p. 62.
- Trachya hystrix, n. sp., TOPSENT, (67) p. 75, pl. i, figg. 8, 9, 10, & pl. xi, figg. 12, 13, 14, Atlantic, lat. 39° 1′ 40″ N., long. 30° 15′ 40″ W., depth 454 metres, and lat. 38° 23′ 30″ N., long. 30° 20′ 20″ W., depth 318 metres.
- Tragosia hirondellei, n. sp., TOPSENT, (67) p. 120, pl. vi, fig. 7, & pl. ix, fig. 5, Atlantic, lat. 43° 40′ N., long 8° 55" W., depth 134 metres.
- Trinacophora? spissa, n. sp., TOPSENT, (67) p. 124, pl. vi, fig. 12, & pl. ix, fig. 9, Atlantic, lat. 38° 23′ 30″ N., long. 30° 20′ 20″ W., depth 318 metres.
- Trochospongilla horrida, n. sp., Weltner, SB. nat. Fr. 1893, p. 12. Synonym for Non Spongilla erinaceus, Ehrenberg, Spongilla erinaceus, Lieberkühn, &c.
- Yvesia, n. g., p. 102, for 16 species, 8 of which are old, viz., Cribrella elegans, hospitalis, papillosa, O. S., Grayella cyathophora, Carter, Halichondria albula, Bwk., Sclerilla dura, arctica, Hansen, and 8 new, viz., Y. pedunculata, pl. v, fig. 6, & pl. x, fig. 17, Atlantic, lat. 43° 57' N., long. 9° 27' W., depth 300 metres, guernei, pl. iii, fig. 7 & pl. x, fig. 16, p. 105, Atlantic, lat. 43° 40' N., long. 8° 55' W., depth 134 metres, Atlantic, lat. 46° 4′ 40" N., long. 49° 2′ 30" W., depth 1267 metres, fallax, p. 106, pl. vi, fig. 13, & pl. x, fig. 14, Straits of Pico-Fayal (Azores), depth 130 metres, hanseni, p. 106, pl. vi, fig. 9a, & pl. x, fig. 13, Atlantic, lat. 39° 1 40" N., long. 30° 15' 40" W., depth 454 metres, richardi, p. 107, pl. x, fig. 15, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 736 metres, ridleyi, pl. x, fig. 12, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 736 metres, pertusa, pl. iv, fig. 10, & pl. x, fig. 18, p. 107, Atlantic, lat. 39° 1' 40" N., long. 30° 15' 40" W., depth 454 metres, linguifera, p. 108, pl. x, figg. 10 & 11, Atlantic, lat. 38° 22' N., long. 30° 34' 39" W., depth 736 metres, Topsent (67).

#### XVII. SPONGIZ.

#### 5. KERATOSA.

Aplysilla rosea, pp. 290 & 291, sulfurea, pp. 284-290, pl. xii figg. 1-15, Triest and Lesina, redescribed; Lendenfeld, Z. wis Zool, liv.

Darwinella aurea, redescribed; LENDENFELD, t. c. pp. 278-283, pl. xii figg. 16-24, Lesina.

#### INCERTÆ SEDIS.

†Stellispongia, St. Cassian, Tyrol, OGILVIE, p. 48, Q. J. Geol. Soc. xli1 n. sp. (sic.)

†Spongia, n. g. (sic.), OGILVIE, l. c.

No description or illustration is given of these two Sponges.



# XVIII. PROTOZOA.

BY

# R. HANITSCH, Ph.D.

#### I.—LIST OF PUBLICATIONS.\*

- Anonymous. Die neuesten Fortschritte auf dem Gebiete der Peridineen-Forschung. Naturwissenschaftliche Wochenschrift, vii, No. 18, pp. 173-179.
- APÁTHY, S. Kritische Bemerkungen über das Frenzelsche Mesozoon Salinella. Eine biologische Skizze. Biol. Centralbl. zii, pp. 108-123. Abstract in Ann. N. H. (6) vi, p. 465.
- APPELLOFF, A. Om Bergensfjordenes faunistike præg. Bergens Mus. Aarsber. 1891.

Foraminifera, p. 5.

- APSTEIN, C. Ueber das Plankton des Süsswassers. Schr. Nat. Ver. Schleswig, ix, 2, pp. 313-316.
   Protozoa, p. 316.
- BALBIANI, E. G. Nouvelles recherches expérimentales sur la mérotomie des Infusoires ciliés. Ann. Micrograph. 1892, 79 pp., 3 pls. Abstracts in J. R. Micr. Soc. 1892, pp. 803-805, and in Bibliographie Anatomique, i, No. 1, p. 15.
- BARROIS, C. Sur la présence de fossiles dans le terrain azoïque de Bretagne. C.R. cxv, No. 6, pp. 326-328.

Fossil Radiolaria.

 BASSETT-SMITH, P. W. On the Deep-sea Deposits of the Eastern Archipelago. J. R. Micr. Soc. 1892, p. 443.

<sup>\*</sup> An asterisk prefixed to a quotation indicates that the Recorder has not seen the Journal or Work referred to.

- 8. Bertram, -.. Beiträge zur Kenntniss der Sarcosporidien nebst einem Anhange über parasitische Schläuche in der Leibeshöhle von Rotatorien. Zool. Jahrb. v, hft. 3 & 4, pp. 581-604, 3 pls. Abstract in J. R. Micr. Soc. 1893, p. 56.
  - 1 new species.
- 9. BIEDERMANN, R. Ueber die Structur der Tintinnen-Gehäuse. Kiel (Lipsius & Tischer): 1892, 4to, 38 pp., 3 pls.
- 10. BLANC, H. Protistes dragués au fond du lac Léman appartenant au genro Difflugia. Compt. rend. trav. 74 sess. Soc. Helv. Sc. Nat. рр. 55 & 56.
- 11. Étude sur les Protistes du fond du Léman. Arch. Sci. Nat. xxvii, No. 4, p. 472.
- 12. —. Les Difflugies de la faune profonde du Lac Léman. Extrait du Recueil inaugural de l'Université de Lausanne. Lausanne (Viret-Genton): 1892, 4to, 11 pp., 1 pl. Abstract in Bibliographie Anatomique, i, No. 1, p. 16.
  - 1 new species.
- 13. Bles, E. J. Notes on the Plankton observed at Plymouth during June, July, August, and September, 1892. J. Mar. Biol. Ass. (n.s.) ii, No. 4, pp. 340-343.

Dinoflagellata, pp. 341 & 343.

- 14. Borgert, A. Vorbericht über einige Phaeodarien- (Tripyleen) Familien der Plankton-Expedition. Ergebnisse der Plankton Expedition, I. A., pp. 176-184, 1 pl. 5 new species.
- 15. Brandt, K. Ueber Anpassungserscheinungen und Art der Verbreitung von Hochseethieren. In Reisebeschreibung d. Plankton-Expedition, pp. 338-370, 1 map.

Hydrostatic apparatus of Protozoa, pp. 341-344.

16. Braun, M. Nachträge und Berichtigungen zum zoologischen Theil von "Die landeskundliche Literatur über die Grossherzogthümer Mecklenburg," &c. Arch. Ver. Mecklenb, xlv, pp. 87-95. Protozoa, pp. 93-95.

- 17. BÜTSCHLI, O. Untersuchungen über mikroskopische Schäume und das Protoplasma. Verh. deutsch. Zool. Ges. 1891, pp. 14-20. Abstract in J. R. Micr. Soc. 1892, pp. 17 & 18.
- 18. Carter, F. B. Radiolaria: their Life-history and their Classification. Am. Micr. J. xiii, pp. 63 & 64, 111-113, 143-145, 177-180, 216-219.
- 19. Certes, A. Sur la vitalité des germes des organismes microscopiques des eaux douces et salées. C.R. cxiv (1892), pp. 425-428. Abstract in J. R. Micr. Soc. 1892, p. 379.

- 20. CHAPMAN, F. The Foraminifera of the Gault of Folkestone, II. J. R. Micr. Soc. 1892, pp. 319-330, 2 pls.
  - 6 new species, 1 new variety.
- 21. —. The Foraminifera of the Gault of Folkestone, III. J. R. Micr. Soc. 1892, pp. 749-758, 2 pls.
  - 1 new species, 1 new variety.
- 22. Some new forms of Hyaline Foraminifera from the Gault. Geol. Mag. (n.s.) (3) ix, pp. 52-54, 1 pl.
  - 1 new genus, 1 new species, 1 new variety.
- 23. CHASTER, G. W. Foraminifera of Southport. Report of the Southport Society of Natural Science, i (1890-91), pp. 54-72, 1 pl. Abstract in J. R. Micr. Soc. 1892, p. 379.
  - 10 new species.
- 24. CONDORELLI, M., & FIORE, C. DE. Un caso di psorospermosi in un Coccothraustes vulgaris. Boll. Soc. Rom. Studi. Zool. i, pp. 68-74.
- 25. COOKE, J. H. The Marls and Clays of the Maltese Islands. Q. J. Geol. Soc. xlix, pt. 1, pp. 117-128, 4 woodcuts. 31 species of Foraminifera, none new.
- Foraminiferi e Radiolari fossili delle sabbie gialle 26. Corti, B. plioceniche della collina fra Spiechio e Limite sulla sponda destra dell' Arno. Boll. Scient. xiv, Nos. 2 & 3, pp. 61-70, 1 pl. 2 new species, 2 new varieties.
- Foraminiferi e Diatomee fossili del pliocene di Castenedolo. Rend. Ist. Lombardo (2) xxiv, fasc. 15 & 16, pp. 991-1012, 1 pl.
- 28. CUÉNOT, L. Protozoaires commensaux et parasites des Échinodermes (Note préliminaire). Rev. Biol. iii (1891), pp. 285-300, pl. v. Abstract in J. R. Micr. Soc. 1892, p. 54.
  - 4 new species.
- 29. Commensaux et Parasites des Échinodermes (Deuxième Note). Rev. Biol. v, pt. 1 (Oct. 1892), pp. 1-23, 1 pl. Sporozoa, pp. 1-6; Ciliata, pp. 6-8.
- 30. DANGEARD, P. A. La nutrition animale des Péridiniens. Le Botaniste (Dangeard), iii (1892), pp. 7-25, 1 pl.
- 31. —. Note sur un Cryptomonas marin. T. c. p. 32, 1 fig. Abstract in J. R. Micr. Soc. 1892, p. 380. 1 new species.
- 32. Danilewsky, —. Contribution à l'étude de la microbiose malarique. Annales Inst. Pasteur, 1891, p. 578. Abstracts in CB. Bakt. Parasit. xi (1892), pp. 513-515, and in J. R. Micr. Soc. 1892, pp. 626 & 627.
- 33. DAWSON, J. W. Carpenter on Eozoon. Nature, xlv, p. 461.
- 34. On Eosoon. T. c. p. 606.

- Dervieux, E. Studio dei Foraminiferi pliocenici di Villarv Atti Acc. Tor. xxvii, pp. 376-379.
- 36. Dock, G. Studies in the etiology of malarial infection and a haematozoon of Laveran. Medical News, July 19th, Abstracts in CB. Bakt. Parasit. x (1891), p. 254, and in J. B. Soc. 1892, p. 53.
- Döderlein, L. [Land-Foraminiferen von Japan.] Verh. de Zool. Ges. ii (1892), pp. 145 & 146.
   new genus and species.
- S8. DREYER, F. Ziele und Wege biologischer Forschung, beleuch der Hand einer Gerüstbildungsmechanik. Jena (G. Fischer): 8vo, xii & 103 pp., 6 pls.
- Die Principien der Gerüstbildung bei Rhizopoden, Spound Echinodermen. (Fortsetzung.) Jen. Z. Nat. xxvi (n. f pp. 297-468, 10 pls. Abstract in J. R. Micr. Soc. 1892, p. 767
- 40. ENTZ, G. Die elastischen und contractilen Elemente der cellinen. Math. Nat. Ber. Ung. x, hft. 1, pp. 1-48, 3 pls.
- 41. FAGGIOLI, F. De l'action délétère du sang sur les Protistes. Ital. Biol. xvi, fasc. 2 & 3, pp. 276-285.
- Foà, P. Ueber die Krebsparasiten. CB. Bakt. Parasit. xii ( pp. 185-192, 2 pls. Abstract in J. R. Micr. Soc. 1892, pp. 807
- 43. FORNASINI, C. Secondo Contributo alla conoscenza della micro terziaria italiana. Mem. Acc. Bologn. (5) i, (1890), pp. 109-11
- 44. —. Terzo Contributo alla conoscenza della microfauna te italiana. Op. cit. ii, pp. 213-221, 1 pl.
  - On Pliocene forms of Nodosaria obliqua.
- 45. Fox, H., & TEALL, J. J. H. On a Radiolarian Chert from M Island. Q. J. Geol. Soc. xlix, pp. 211-218, 1 pl., 1 map. [F of same paper HINDE, G. J., Note on the Radiolaria in the M Island Chert, pp. 215-218, 1 pl.]
  - 10 species of Radiolaria, partly undetermined, none described as
- FRANCAVIGLIA, M. C., & FIORE, C. DE. [Psorospermium ar Coccothraustes vulgaris.] Boll. Soc. Rom. Stud. Zool. i (1892), 1
   1 fig. Abstract in J. R. Micr. Soc. 1892, p. 495.
- FRENZEL, J. A new form of Trichonymphidæ. J. R. Micr 1892, pp. 52 & 53. Abstract of paper in Arch. mikr. Anat. x: pp. 301-316. [See Zool. Rec. 1891.]
- Remarkable Argentine Protozoa. J. R. Micr. Soc. pp. 219 & 220. Abstract of paper in Z. wiss. Zool. liii, pp. 33-1 pl. [See Zool. Rec. 1891.]
- 49. A multicellular Infusorian-like Animal. Ann. N. H. (pp. 109-111. Abstract of paper in Arch. f. Nat. lviii, pp. (Clear Zool. Rec. 1891.]



- [FRENZEL, J.] Das Mesozoon Salinella. Biol. Centralbl. xi, pp. 577-581. Translated in Ann. N. H. (6) vi, pp. 79-84.
- Ueber einige argentinische Gregarinen. Ein Beitrag zur Organisation und Physiologie der Gregarinen überhaupt. Jen. Z. Nat. xxvii (1892), pp. 233-336, 1 pl. Abstract in J. R. Micr. Soc. 1892, p. 805.
  - 5 new species.
- Untersuchungen über die mikroskopische Fauna Argentiniens.
   I. Theil: Die Protozoen. I. & II. Abtheilung: Die Rhizopoden und Helioamoeben. Bibl. Zool. xii (1892), pp. 1-82, 6 pls. Abstract in J. R. Micr. Soc. 1892, pp. 624 & 625.
  - 12 new genera, 24 new species.
- FRITSCH, A., & VÁVRA, V. Vorläufiger Bericht über die Fauna des Unter- Pocernitzer und Gatterschlager Teiches. Zool. Anz. xv (1892), pp. 26-30.
  - 29 Protozoa obtained; 5 undetermined, none new.
- 54. GEINITZ, E. XIII. Beitrag zur Geologie Mecklenburgs. Weitere Aufschlüsse der Flötzformationen. Arch. Ver. Mecklenb. xlvi, pp. 59-97, 3 pls.
  - Foraminifera, pp. 68 & 69.
- 55. Geinitz, H. B. Die Versteinerungen des Herzogthums Sachsen-Altenburg. MT. Osterlande (n.f.) ▼, pp. 161-199.
   Ehizopoda, pp. 179 & 180.
- 56. Goës, A. On a Peculiar Type of Arenaceous Foraminifer from the American Tropical Pacific: Neusina agassizi. Bull. Mus. C. Z. xxiii, No. 5, pp. 195-197, 1 pl. Abstract in J. R. Micr. Soc. 1893, No. 2, p. 199.
- 57. GOROSCHANKIN, J. Beiträge zur Kenntniss der Morphologie und Systematik der Chlamydomonaden. Bull. Mosc. 1891, pp. 101-142, 3 pls. Abstract in J. R. Micr. Soc. 1892, p. 54. 3 new species.
- GRASSI, B., & FELETTI, R. Contribuzione alla studio dei parassiti malarici. 5 l. e. a. (1892), 4to, 80 pp., 1 pl. [From Zool. Anz. 1892.]
- Grassi, B. Conclusioni d'una Memoria sulla società dei Termiti.
   Atti Acc. Pontif. Lincei (5) i, (1892), pp. 33-36. Abstract in J. R.
   Micr. Soc. 1892, p. 472.
  - Parasitic Protozoa in Termites. 3 new genera, 5 new species.
- GREEFF, R. Terricolous Amaba. J. R. Micr. Soc. 1892, p. 51.
   Abstract of paper in MT. Ges. Marb., Feb. 1891. [See Zool. Rec. 1891.]
- Organization of Amaba. J. R. Micr. Soc. 1892, p. 51. Abstract of paper in Biol. Centralbl. xi, p. 599. [See Zool. Rec. 1891.]

- Criber America. III. Mittheilung. SB. Ges. Marb. pp. 22-43. and Birl. Centralbl. xii, pp. 373-384. Abstract i Micr. Soc. 1992, p. 625.
- GEENET, J. W. The Tudor Specimen of Econom. Nature pp. 486 & 487.
- GETTYTHE. A. R. The Physiology of the Invertebrata. 1
   (L. Reeve & Co.): 1892, 8vo. 477 + xvi pp., 81 woodcuts.
   Processes. pp. 23-30. 79. 119, 120, 183, 184, 208, 246-248, 295, 29
   326. 275-377, & 404-468.
- GETHER, A. Eine Mittheilung über Kernvermehrung und 8 merkildung bei Süsswamer-Rhisopoden. Ber. Ges. Freibt pp. 114–114, 1 pl.
- Ernzellige Zwerge. In Festschrift zum 70. Geburtst Leuchart's, pp. 74-76, 2 woodcuts.

Abnormally small specimens of Steater curuleus and S. polymor,

- 63. GTERNZ, J. DE, & RICHARD, J. Sur la faune des eaux do FIslande. C.R. pp. 310-313. Abstract in Ann. N. H. (6) x, 1 Protonoon: Ceratium longicorne, Perty.
- HANTTSCH. R. Foraminifer or Sponge? Nature, xlvii, No. p. 365.
- 7). —. Foraminifer or Sponge? Op. cit. No. 1219, p. 439. On Kenina agessizi, Goës, a supposed Foraminifer.
- CHEHIR, P. [Hæmatozoon of Malaria.] J. R. Micr. Soc p. 389. Abstract of "Separate copy, sent by author, no d place of publication, 4to, 27 & ii pp., 9 pla."
- 72. c—. [New Cholera Microbe.] Indian Medical Gazette ( Supplement) April, 1892, 7 pp., 11 pls. Abstract in J. R. Mi 1892, p. 495.
- 73. HENNEGUY, F., & THÉLOHAN, P. Sur un Sporozoaire para muscles des Crustacés décapodes. C.R. cxiv (1892), pp. 155 Abstract in Ann. N. H. (6) x, pp. 342-344, and in J. R. Mi 1892, p. 626.
- 74. HERDMAN, W. A. Notes on the Collections made during the of S.Y. "Argo," up the West Coast of Norway, in July P. Liverp. Biol. Soc. vi, pp. 70-93, 2 pls.
  Francision and S. 88
  - Foraminifera, pp. 86–88.
- Hinde, G. J. Note on a Radiolarian Rock from Fanny Ba Darwin, Australia. Q. J. Geol. Soc. xlix, pp. 221-226, 1 pl. 11 new species.

- 76. HOBNE, J. On the Contact Metamorphism of the Radiolarian Chert in the Lower Silurian Rocks along the margin of the Loch Doon Granite. Rep. Brit. Ass. 1892, p. 712.
- 77. Hosius, A. Beiträge zur Kenntniss der Foraminiferen-Fauna des Miocans. Verh. Ver. Rheinl. xlix, pt. i, pp. 148-197, 2 pls. Many undetermined, none described as new.
- 78. Imhor, O. E. Notizie sulla Fauna pelagica della laguna di Venezia. Neptunia, i, No. 1. 1. Gennaio: 1891, pp. 1-4. Several Protozoa. 1 new species.
- 79. JAKOWLEW, W. Opisanie nyeskol'kikh vidov myelov'ikh Foraminifer. Description de quelques espèces des Foraminifères crétacés. Trav. Soc. Univ. Kharkow, xxiv, pp. 341-361, 3 pls. 1 new genus, 2 new species.
- 80. JENSEN, P. Ueber den Geotropismus niederer Organismen. Arch. ges. Phys. liii, hft. 9 & 10, pp. 428-480, 13 woodcuts.
- 81. Methode der Beobachtung und Vivisection von Infusorien in Gelatinelösung. Biol. Centralbl. xii, Nos. 18 & 19, pp. 556-560.
- 82. KLEBS, G. Flagellatenstudien. Theil I. Z. wiss. Zool. lv, hft. 2, pp. 265-351, 4 pls. Theil II. t. c. hft. 3, pp. 353-445, 2 pls. Abstract in J. R. Micr. Soc. 1893, No. 2, pp. 196 & 197.
  - 8 new families, 9 new genera, 42 new species.
- 83. KOROTNEFF, A. Myxosporidium bryozoides. Z. wiss, Zool, liii, pp. 591-596, 1 pl. Abstract in J. R. Micr. Soc. 1892, p. 379.
- 84. Kruse. W. Der gegenwärtige Stand unserer Kenntnisse von den parasitären Protozoen. Hyg. Rundschau, 1892, pp. 357-380 & 453-485.
- 85. Lameere, A. Prolégomènes de Zoogenie. Bull. Sci. Fr. Belg. xxiii (1891), pp. 399-411.
  - Systematic position of Protozoa, &c.
- Ueber Periodizität im Auftreten und in der 86. LAUTERBORN, R. Fortpflanzung einiger pelagischen Organismen des Rheines und seiner Altwasser. Zeitschrift für Fischerei, &c., i (1893), No. 2, pp. 81-95, 4 woodcuts, and in Verh. Ver. Heidelb. (n.f.), v, hft. 1, pp. 1-15.
- 87. LAVERAN, A. Du paludisme et son hématozoaire. Paris (G. Masson): 1891, 8vo, 300 pp. Abstract in CB. Bakt. Parasit. xi (1892), pp. 510-512, and in J. R. Micr. Soc. 1892, p. 626.
- 88. Léger, L. Recherches sur les Grégarines. Rev. Sci. xlix, pp. 660 & 661.
- 89. LEVANDER, K. M. Notiz über die Täfelung der Schalenmembran des Glenodinium cinctum, Ehbg. Zool. Ans. xv (1892), pp. 405-408, 4 woodcuts. Abstract in J. R. Micr. Soc. 1893, No. 2, p. 197.

- LT from E M | Verrescribes der wihrend des Sommers Erstelle des Arch. Ver. Meckle
  - ... Tested Some new ones will be described in a fate
- District E. Notice of the Commence of Protozoon Remember of Commond Fisher in Ohio. Bull. U.S. Fisher in Chem. 1986.
- La La Fa Francia delle sopre dulli di Genova e dinto
- 1 and J. L. Lossening Lamppoist. No. 8, Sci. Goss., F. Program of the Unit. No. 8 in Gossil June, 1862.
- 4. L. C. A. F. Tambiert mossini fi. Paulio nell' A. A. Gress, Am. So. Mod. http://dx.
  - F-1.- 111: 1:T.
  - F manuferi pliceena il l'assellativato e Lugago Francia i Placetani Bell. Son geol Itali xi, fasc. I, pp.
  - I for E F. Le mouvement des êtres morresopiques as a constitution protons. C.B. exiv.pp. (8) 5-29 ( Abstraction to to constitution.
- 7 A. . . E. L. form a formulation lells marks she at a concept violation of Viterio, B.H. Six geal Ital. 2 (2017).
- The model of F. List of the Fishels found in the Cambridge and results of Total Edition Not. Histo Soc. Brunswick, min.
  - ·.. . --
- (1) The first of E. North to street in momente de M. Sone thank to Inter Flatter 18 (1) 178. Abstracts in CB. Bakes of the first fill E. Mars Son 1881, pp. 617 & 628.
  - The American Control
  - A construction of the Gregarines of Holes Construction of Holes Construction of the Gregarines of the G
  - Differences
  - A main of F classification of Species. J. R. Micr. of the species Act. Acc. Rom. (4) vii. pp. S. I. of E. of the control of the species act.
- Now Harmond in regardless. J. R. Mich Soc. 1communication of Latte Ann Ram. (4) vii. pp. 229-2 Local Res. (5).
  - Piene erikung liella Senetenia octopiana. Atti I

- 104. [Mingazzini, P.] Classificazione dei Coccidi e delle Gregarine. Atti Acc. Pontif. Lincei (5) i, pp. 68-75. Abstract in J. R. Micr. Soc. 1892, p. 380.
- Contributo alla conoscenza dei Coccidi. Atti Acc. Pontif. Lincei (5) i, pp. 175-181. Abstract in J. R. Micr. Soc. 1892, p. 495.
- 106. —. Nuove specie di Sporozoi. Atti Acc. Pontif. Lincei (5) i, pp. 396-402, 4 figg. Abstract in J. R. Micr. Soc. 1892, p. 806.
  2 new genera; 2 new species.
- 107. MITTER, J. Beitrag zur Kenntniss des Balantidium coli im menschlichen Darmkanale. Inaugural Dissertation. Kiel: 1891. Abstracts in CB. Bakt. Parasit. xii (1892), p. 111, and in J. R. Micr. Soc. 1892, p. 625.
- 108. MRÁZEK, A. O tak zvané Monocystis tenax, Stein. SB. Böhm. Ges. 1892, pp. 67-76, 2 woodcuts.
- No résumé in German. The author endeavours to prove that the species belongs to the Flagellata.
- 109. MURRAY, J., & IRVINE, R. On Silica and Siliceous Remains of Organisms in Modern Seas. P. R. Soc. Edinb. xviii, pp. 229-250.
- 110. Noll, F. C. Die Ernährungsweise des Trichosphærium sieboldii, Schn. Zool. Anz. xv (1892), pp. 209 & 210. Abstract in J. R. Micr. Soc. 1892, p. 805.
- 111. NORMAN, A. M. Museum Normanianum; or, a Catalogue of the Invertebrata of Europe, and the Arctic and North Atlantic Oceans. VII & VIII: Spongozoa and Rhizopoda. Durham: 1892, 8vo, 21 pp. Rhizopoda, pp. 14-21.
- OSBORN, H. L. The Protozoa—a Phylum of the Animal Kingdom considered biologically. Am. Micr. J. xiii, No. 10, pp. 233-243, 1 pl.
- 113. Peach, B. N. On a widespread Radiolarian Chert of Arenig Age from the Southern Uplands of Scotland. Rep. Brit. Ass. 1892, pp. 711 & 712.
- 114. Pearcey, F. G. Foraminifer or Sponge? Nature, xlvii, No. 1217, p. 390.
  - On Neusina agassizi, Goës, a supposed Foraminifer.
- 115. Perner, J. Předběžný Kritický Seznam Foraminifer z Březenských Vrstev. Résumé, Kritisches Verzeichniss der Foraminiferen aus den Priesener Schichten der böhmischen Kreideformation. SB. Böhm. Ges. 1892, pp. 34-53.
  - 1 new species.
- 116. —... Ueber die Foraminiferen des böhmischen Cenomans. (Résumé des böhmischen Textes.) Palseontographica Bohemise, i, pp. 49-65, 10 pls.
  - 28 new species.

- 117. PFEIFFER, R. Beiträge zur Protozoen-Forschung. 1. He Coccidien-Krankheit der Kaninchen. Berlin (A. Hirs-1892, 8vo, 24 pp., 12 pls.
- 118. —. Vergleichende Untersuchungen über Schwärmspungen bei den Coccidieninfektionen und bei Inte Fortschritte d. Medicin, 1890, pp. 939-951. Abstracts in (Parasit. xii (1892), pp. 109 & 110, and in J. R. Micr. Soc. 806 & 807.
- 119. —. Ueber einige neue Formen von Miescherschen Semit Mikro-, Myxo-, und Sarkosporidieninhalt. Virchow's exxii, pp. 552-573, 1 pl. Abstracts in CB. Bakt. Parasit. x pp. 110 & 111, and in J. R. Micr. Soc. 1892, p. 807.
- 120. —... Die Protozoen als Krankheitserreger sowie der Ze Zellkernparasitismus derselben bei nicht bakteriellen II krankheiten des Menschen. 2 Aufl. Jena (G. Fische Lex. 8vo, vi & 216 pp., 91 woodcuts. Abstracts in CB. Bakt xii (1892), pp. 168-171, and in J. R. Micr. Soc. 1892, p. 806
- 121. PODWYSSOZKI, W., & SAWTSCHENKO, J. Ueber Parasiti Carcinomen nebst Beschreibung einiger in den Car sohwülsten schmarotzenden Sporozoen. CB. Bakt. Parasit. 1 pp. 493-500, 532-538, & 559-563, 2 pls. Abstract in J. B. 1 1892, p. 629.
- 122. POLLARD, E. C. A New Sporozoon in Amphioxus. Q. Sci. xxxiv, pt. 3, pp. 311-316, 1 pl. Species not named.
  - POUCHET, G. Cinquième contribution à l'histoire des Péri Peridinium pseudonoctiluca, Pouchet. J. de l'Anat. Phy pp. 143-150, 1 pl.
  - 124. \*RAILLIET, A. [On Myoxsporidia.] Bull. Soc xv (1890), pp. 192-198.
  - 125. RAILLIET, A., & LUCET, A. Note sur que' encore peu étudiées. Bull. Soc. Z. Fr. 2 new species.
  - 126. RHUMBLER, L. Eisenkiesabl körper von Foraminiferer Schultze's u. A. Vorla-1892, No. 12, pp. 419-4
  - 127. ROBERTSON, D. Island of Skyrpp. 239-242

None new

- 128. ROSENBERG, —. Ein Befund von Psorospermien (Sarcosporidien) im Herzmuskel des Menschen. Zeitschr. f. Hygiene, xi, No. 3. Abstracts in CB. Bakt. Parasit. xi (1892), p. 739, and in J. R. Micr. Soc. 1892, pp. 628 & 629.
  - 1 new species.
- 129. Rüst, J. Beiträge zur Kenntniss der fossilen Radiolarien aus Gesteinen der Trias und der palæozoischen Schichten. Palæontogr. xxxviii, pp. 107-200, 25 pls.
  - 3 new genera; 244 new species.
- 130. RUFFER, M. A., & WALKER, J. H. On some Parasitic Protozous found in Cancerous Tumours. J. Path. Bact. i, No. 2 (Oct. 1892), pp. 198-215, 3 pls.
- 131. SAWTSCHENKO, G. Weitere Untersuchungen über schmarotzende Sporozoen in den Krebsgeschwülsten. CB. Bakt. Parasit. xii (1892), pp. 17-28, 1 pl. Abstract in J. R. Micr. Soc. 1892, p. 628.
- 132. SCHACKO, G. Foraminiferen und Ostracoden aus der Kreide von Moltzow. Arch. Ver. Mecklenb. xlv (1891), pp. 155-160, 1 pl. 4 new species of Foraminifera.
- 133. SCHEWIAKOFF, W. Ueber die geographische Verbreitung der Süsswasser-Protozoen. Verh. Ver. Heidelb. (n.f.) iv, hft. 5, pp. 544-567.
  17 new genera; 36 new species.
- 134. Schilling, A. J. Die Süsswasser-Peridineen. Inaugural-Dissertation. Marburg (R. Friedrich): 1891, 8vo, 81 pp., 3 pls.
- 8 new species. [The Recorder for last year, although he mentions this paper, was unable to see it, and to enumerate the new species.]
- 135. —. Untersuchungen über die thierische Lebensweise einiger Peridineen. Ber. Deutsch. Bot. Ges. ix (1891), pp. 199-208, 1 pl. Abstract in J. R. Micr. Soc. 1892, p. 52.
  - 1 new species.
- 136. Schlumberger, C. Note sur le Ramulina grimaldii. Mém. Soc. Zool. iv, pp. 509-511, 1 pl.
  - 1 new species.
- Bévision des Biloculines des grands fonds. Mém. Soc. Zool. iv, pp. 542-579, 4 pls., 46 woodcuts.
  - 8 new species.
- 138. —... Note préliminaire sur les Foraminifères dragués par S. A. le Prince Albert de Monaco. Mém. Soc. Zool. v, pp. 207-212, 1 pl., 5 woodcuts.
  - 2 new species.
- 139. SCHUBERG, A. Bemerkungen zu den 'Untersuchungen' des Herrn Dr. Angelo Fiorentini über die Protozoen des Wiederkäuermagens CB. Bakt. Parasit. xi, pp. 280–283.

- 140. [SCHUBERG, A.] Ueber einige Organisationsverhältnisse d fusorien des Wiederkäuermagens. SB. Ges. Würzb. 1891, pp 137. Abstract in J. R. Micr. Soc. 1892, p. 494.
- 141. —. Ueber Coccidien des Mäusedarms. SB. Ges. Würzb. pp. 65-72.
- 142. SCHÜTT, F. Ueber Organisationsverhältnisse des Plasm der Peridineen. SB. Ak. Berlin, 1892, pp. 377-384, 1 pl., MT. Ak. Berlin, 1892, pp. 165-171, 1 pl. Abstract in J. R. Soc. 1892, p. 805.
- 143. SEVERI, A. Gregarinosi polmonale in infante nato-morto. R. Accad. Medic. Genova, vii (1892), fasc. 2, 8 pp. Abstract Riforma Medica, 1892, No. 80, and in CB. Bakt. Parasit. xii (p. 262, and in J. R. Micr. Soc. 1892, p. 806.
- 144. SIMMONS, W. J. Some Stalked Actinophyrans. Sci. Goss pp. 124-127, 9 figg. Abstract in J. R. Micr. Soc. 1892, p. 494
- 145. SMILEY, C. W. A beautiful Rhizopod. Am. Micr. J. xiii, 1 54, 8 woodcuts.
- 146. Solger, B. Parasitic Protozoa. J. R. Micr. Soc. 1892, Abstract of paper in MT. Vorpomm. xxii (1890), pp. 99-102. Zool Rec. 1891.]
  - 1 new genus; 1 new species.
- 147. Sollas, W. J. Supposed Radiolarian Remains from the S Howth. [Title only.] Rep. Brit. Ass. 1892, p. 723.
- 148. —... Supposed Radiolarian Remains from the Culdaff Lim [Title only.] *Ibid.*
- SOUDAKEWITSCH, —. Recherches sur le parasitisme intrace et intranucléaire chez l'homme. Ann. Inst. Pasteur, 1892,
   pls. Abstracts in CB. Bakt. Parasit. xii (1892), p. 39,
   J. R. Micr. Soc. 1892, p. 627.
- 150. Tellini, A. Istruzioni per la raccolta, preparazione e cozione dei Foraminiferi viventi e fossili. Riv. Ital. Sci. N (1892), No. 1, pp. 4-7.
- 151. Thélohan, P. Note sur la Glugea microspora. C.B. So. (9) iv (1892), 3 pp.
- 152. —. Sur quelques nouvelles Coccidies parasites des parasites de parasites de parasites de parasites de parasit
- 153. —. Myxosporidies de la vésicule biliaire des Poissons. I nouvelles. C.R. oxv, No. 22, pp. 961-964, and No. 24, pp. 1094. Abstract in J. R. Micr. Soc. 1893, No. 2, pp. 198 & 199 1 new genus; 5 new species.

- 154. [Thélohan, P.] Sur les Myxosporidies. C.R. Soc. Philom. Paris, xx, Août, 1892, pp. 2 & 3.
- THOMAS, B. Notes on Amaba and Infusoria. I. The Amaba. Sci. Goss. March, 1892, pp. 52 & 53, 1 woodcut. II. Flagellate Infusoria. Op. cit. April, 1892, pp. 81 & 82, 1 woodcut, May, 1892, pp. 99-103, 4 woodcuts, June, 1892, pp. 135-138, 5 woodcuts, July, 1892, pp. 147-149, 3 woodcuts, August, 1892, pp. 172-176, 5 woodcuts, October 1892, pp. 231-233, 4 woodcuts.
- 156. TOPSENT, E. Sur un nouveau Rhizopode marin (Pontomyxa flava, n. g. & sp.). C.R. cxiv, pp. 774 & 775. Abstracts in Rev. Sci. xlix, p. 475, and in J. R. Micr. Soc. 1892, pp. 379 & 380.
- 157. Vejdovsky, F. Sur une Thuricola d'eau douce. Congrès Internat. Zool. Sess. ii, No. 1, pp. 52-57. Abstract in J. R. Micr. Soc. 1893, No. 2, p. 190.
  - 1 new species.
- 158. ——. Sur la Monodontophyra, nouvelle espèce d'Opalinide. Congrès Internat. Zool. Sess. ii, No. 1, pp. 24-31.
  1 new species.
- 159. VERBEEK, R. D. M. Vorloopig bericht over nummulieten, orbitoiden en alveolinen van Java, en over den ouderdom der gesteenten waarin zij optreden. Nat. Tijdschr. Nederl. Ind. li, pp. 101-138, 1 pl.
  - 1 new species.
- 160. Verworn, M. Die physiologische Bedeutung des Zellkerns. Arch. ges. Phys. li, pp. 1-118, pls. i-vi.
- 161. VINE, G. R. Notes on *Polyzoa* found at Cleethorpes. In 1875, 1879, and 1882. Naturalist, 1892, pp. 5-11.
  - 1 Foraminifer mentioned (Planorbulina lobatula).
- 162. Weltner, W. Ueber Myxosporidiensporen in den Eiern von Esox lucius. SB. nat. Fr. 1892, No. 4, pp. 28-36, 16 woodcuts.
- 163. Zacharias, O. Vorläufiger Bericht über die Thätigkeit der Biologischen Station zu Plön. Zool. Anz. xv (1892), pp. 457-460.
  2 new species, several undetermined.
- 164. —... Ueber eine Ichthyophthirius-Art aus den Aquarien der Biologischen Station zu Plön. In Festschrift zum 70. Geburtetage
  - 1 new species.
- 165. ZOPF, W. Beiträge zur Physiologie und Morphologie niederer Organismen. 1. Heft. Leipzig (A. Felix): 1892, 8vo, pp. iv & 97, 3 pls.
  - 1892. [vol. xxix.]

R. Leuckart's, pp. 289-292, 1 pl.

# II.—ANATOMY, BIOLOGY.

#### 1. GENERAL MORPHOLOGY.

BÜTSCHLI (17) gives an account of the various opinions regar structure of protoplasm, and states his own views on the subject.

DREYER (39) discusses the tetraxial type of the skeleton in t cystina (also in Sponges and in Echinodermata), and offers mexplanations.

Entz (40) gives a detailed account of the elastic and concents of the Vorticellina, especially of the large forms, umbellaria, L., and Zoothamnium arbuscula, Eh. Covering the ethere is a delicate membrane (pellicula) which has a double f (1) it serves as attachment for the contractile elements; and (elastic, it acts as antagonist to the same. The contractile elements of two systems, an outer and an inner one, each again being of two layers. Thus there are altogether four layers of myor outer circular, an outer longitudinal, an inner circular, and an intudinal. These layers are now described in their most minut. The paper is illustrated by a series of very high-power drawings

FRENZEL (47-52) gives elaborate papers on Argentine Protoz cially on Rhizopoda, Helioamaba, and Sporozoa.

GREEFF (63) describes a thin, very flexible, resistent, and chitinous membrane, which encloses the body of *Amaba fluide* interrupted by an opening through which the pseudopodial: ("Zottenanhang") projects.

GRUBER (67) describes abnormally small specimens of Stentor and S. polymorphus. The former had only a single nuclear joint

KLEBS (82) describes the peripheral portion of the *Flagella* "periblast" is the outermost plasmic layer, and is always Outside this there is a secondary envelope ("Hülle"), which absent.

LEVANDER (89) describes the cuticular shell of Glenodinium Ehrenb., as being composed of a number of polygonal plates, as dinium (tabulatum).

RUST (129) makes important contributions to our knowledge *Badiolaria*, with exhaustive illustrations.

SCHLUMBERGER (137) has studied the Deep Sea *Biloculines*, cl means of median sections, and describes a kind of dimorphism (t term in a sense slightly different from its customary meaning). species may be distinguished two forms, A and B. In the form B chamber (microsphère) is extremely small, and is succeeded by n small chambers. The form A begins at once with a large (megasphère).

SCHÜTT (142) has studied the structure of *Peridinidæ*. He distitute concentric layers of plasma: an outer layer (which consists two layers, one hyaline and the other granular), and an inner man

is not granular, and contains nucleus, sap-chambers, and various vacuoles.

Schilling (134) has a paper on the structure of the freshwater Peridinidæ.

#### 2. REPRODUCTION.

GREEFF (63) asserts that he has never noticed reproduction by means of binary fission, either in Amaba fluids or in any other Amaba. He describes phenomena which remind one of the formation of spores.

GRUBER (66) describes the formation of spores in Arcella vulgaris and Lecythium hyalinum.

KLEBS (82) says that in studying the Flagellata (from which he excludes the Volvocina), he never has seen a sexual process of reproduction. All reproduce asexually and by longitudinal fission (except Oxyrrhis marina, which divides transversely). Still, he thinks it possible that forms along the border line, allied to Ciliata or Volvocina, may yet be discovered which reproduce sexually.

SCHUBERG (140) has studied the reproduction of Dasytricha ruminantium, and of other Ciliata, living in the rumen and reticulum of Ruminantia.

SCHUBERG (141) describes the formation of spores of Coccidia in the stomach of a mouse.

SCHILLING (134) discusses the reproduction of the *Peridinidæ*. The rule is vegetative reproduction, and doubtful are the accounts of copulation and conjugation.

#### 3. PHYSIOLOGY.

BALBIANI (5) has been experimenting on the artificial division of *Ciliata*. In merozoites which contain the whole, or part, of a nucleus, the regeneration is rapid and complete.

BRANDT (15) describes the hydrostatic apparatus of some Protozoa.

CERTES (19) describes his experiments on the vitality of germs of microscopic organisms. Cultivation of freshwater sediments yields by far more life (*Protozoa* and a few *Metazoa*) than that of marine sediments.

Dangeard (30) discusses the animal-like nutrition of some Peridinide.

FRENZEL (51) gives an account of the chemistry of *Gregarinida*. He distinguishes the following substances: Protoëlastin, Alveolin, Paralveolin, neutral fats, Albuminoids, Protocollagen, Paraglycogen, Pyxinin, Antienzym, Morulin, Paramorulin, Nuclein, Nuclear Sap, Cell Sap.

GREEFF (63) gives an account of the movement and nutrition of Amaba.

GRIFFITHS (65) treats of *Protozoa* in the following pages: 23-30, 79, 119-123, 183, 184, 208, 246-248, 295, 296, 346-348, 375-377, & 404-408.

JENSEN (80) speaks of the Geotropism of Protozoa.

LAUTERBORN (86) gives some interesting sketches on the periodicity

in the appearance and reproduction of polagic organisms in the Rhits "Altwasser." Besides Alga, Rotatoria, Crustacea, Mollusca, he of Dinobryon, Ceratium hirundinella, and other Protozoa.

Noll (110) gives an account of his observations of *Trichospisieboldii* feeding on Diatoms along the glass walls of his aquariu reproducing rapidly.

RHUMBLER (126) shows that the so-called "Budspores" ("Keimk of Foraminifera are foreign bodies. They may be (1) either I (Cocconeis?), or (2) granules of excretory matter (?), or (3) prod decomposition of the animal. He has studied especially this thi of bodies. They are of a black colour, show sometimes a distinct talline structure, and consist of iron-pyrite. This agrees with th ments of geologists (e.g., Justus Roth) that a great part of th pyrites are of organic origin.

SCHILLING (135) discusses the animal-like nutrition of some *Per* Verworn (160) treats of the physiological significance of the respecially of that of *Protozoa*.

Protozoa and Human Pathology: Sporozoa (or other Protozoa be demonstrated in numerous cases of Cancer. See papers b Metschnikoff, Pfeiffer (120), Podwyssozki, Ruffer & W Sawtschenko, and Soudakewitsch. For Malaria and Hamato Danilewsky, Dock, Grassi & Feletti, Hehir, and Laveral various other diseases, in their connections with Protozoa, see pa Kruse, Mitter, Pfeiffer, Rosenberg, and Severi.

# III.—DISTRIBUTION.

#### A. FAUNISTIC.

Argentinia.—Rhizopoda, Heliamaba, Gregarinida; FRENZEL.

Atlantic.—BORGERT treats of the geographical distribution *Phæodaria*, collected during the Plankton-Expedition. Richer the Irminger Sea and the Labrador current; exceedingly poor 1 gossa Sea, probably on account of its very high temperature poorer were the Florida, and the East- and West-Greenland a Foraminifera: SCHLUMBERGER.

Iceland.—Freshwater Protozoa; GUERNE & RICHARD.

Great Britain.—I. of Skye: Foraminifera; ROBERTSON.—Sou Foraminifera; CHASTER.—Cleethorpes: 1 Foraminifer; VINE mouth: Dinoflagellata; BLES.—Plymouth: Sporozoa; MINCHIN Norman — West Coast: Foraminifera: HERDWAY. Roman

Norway.—West Coast: Foraminifera; Herdman:—Bergen: infera; Appellöff.

France.—Banyuls: Rhizopoda; TOPSENT.

Germany.—Holstein: Freshwater Protozoa; Zacharias.—Le (90) gives a list of freshwater Protozoa collected near Rostock, in lenberg, viz., 12 Rhizopoda, 4 Heliozoa, 45 Flagellata, 4 Dinofla Ciliata.

Belgium.—Ostend: Marine Amæbæ; GREEFF.

Bohemia.—Protozoa; FRITSCII & VÁVRA.

Switzerland.—Basle: Flagellata; KLEBS: Dinoflagellata; SCHILLING.

-Lake of Geneva: Difflugia and Protozoa generally; BLANC.

Italy.—Genoa: Freshwater Protozoa; Longiii.—Sporozoa, Mingazzini.

-Lagoons of Venice: Protozoa, IMHOF.

India.—Calcutta: Clathrulina elegans and Hedriocystis pellucida; SIMMONS: C. elegans was first found in St. Petersburgh, afterwards in Central and Western Europe, and in North America.

Malay Archipelago.—Freshwater Protozoa; SCHEWIAKOFF.

Japan.—Sand Foraminifera; DÖDERLEIN.

Eastern Archipelago.—Deep-sea deposits; BASSETT-SMITH.

Australia.—Freshwater Protozoa; Schewiakoff.

New Zealand.—Freshwater Protozoa; Schewiakoff.

Sandwich Islands.—Freshwater Protozoa; Schewiakoff.

#### B. GEOLOGICAL.

Rüst examined all rocks from the Trias and Palæozoic strata, which were accessible to him, for *Radiolaria*. The Limestones of Palæozoic strata and Trias contained only exceedingly few *Radiolaria*, and these in a condition not suitable for identification. Exceedingly rich were numerous siliceous limestones, siliceous schists, hornstones, jasper, adinole, and wetzschiefer. The skeletons, too, were here in splendid condition.

Laurentian or Precambrian.—Bretagne: Radiolaria allied to the Monosphæridæ. The oldest organic remains of France and probably of the world. Barrois.

Cambrian.—St. John, New Brunswick: Species of Monadites and Radiolarites; MATTHEW.

Ordovician (?).—Mullion I. (off Lizard Peninsula, Cornwall): Radiolarian Chert; Fox & Teall.

Silurian.—Cabrières, Languedoc, France: Radiolaria, Rüst.—Scotland: Radiolaria, Peach.

Devonian.—Schebenholz, Hartz Mountains, and South Ural: Radiolaria, Rüst.

Carboniferous.—Hartz Mountains, Bükk Mountains in Hungary, and Sicily: Radiolaria, Rüst.

Permian.—Nowgorod: Radiolaria, RUST.

Trias.—Astura, Felső Eörs in Bakony and Röthelstein, near Aussee, in Tyrol: Radiolaria, Rüst.—Saxe-Altenburg: Rhizopoda, H. B. GEINITZ.

Cretaceous.—Gault of Folkestone: Foraminifera, CHAPMAN.—Moltzow in Mecklenburg: Foraminifera, SCHACKO. — Priesen, Kamajk, near Časlau, and Gangberg, near Kuttenberg, Bohemia: Foraminifera, PERNER.—Russia: Foraminifera, JAKOWLEW.

Eocene.—Java: Foraminifera, VERBEEK.

Oligocene (Upper).—Maliss, in Mecklenburg: Foraminifera, E. Gein Miocene.—Java, Sumatra, and Nias: Foraminifera, Verbeek.—Kui Mühle, near Dingden, Germany: Foraminifera, Hosius.—Apenni near Modena: Foraminifera, Malagoli.—(Miocene?) Radiolarian R from Fanny Bay, Port Darwin, Australia; Hinde.

Pliocene.—Near River Arno, Italy: Foraminifera and Radiole Corti.—Castellarquato and Lugagno, near Piacenza: Foramini, MALAGOLI.—Italy: Foraminifera, FORROSINI.—Villarvernia, Italy: F minifera, DERVIEUX.—Volcanic tuf of Viterbo, Italy: Foramini, MARIANI.

# IV.—SYSTEMATIC.

# A. -GYMNOMYXA.

#### Class 3.—LOBOSA.

- Amaba fluida, Gruber, pp. 24-41, crystalligera, pp. 41-43, Oste redescribed by GREEFF, SB. Ges. Marb. 1892.
- A. hercules, p. 24, pl. iii, figg. 10, 11, & 17, pellucida, p. 29, pl. i, figg. 3
  Argentinia, FRENZEL, Bibl. Zool. xii, n. spp.
- Diffugia lemani, Lake Leman; Blanc (12), p. 5, pl. ii, figg. 13-15, n. Guttulidium, n. g. with G. tinctum, & sp., Argentinia; Frenzel, Bibl. 2 xii, p. 1, pl. iii, figg. 7 & 8.
- Limulina, n. g. with L. unica, & sp., Argentinia; FRENZEL, Bibl. 2 xii, p. 41, pl. iii, fig. 9.
- Mastigamæbu schulzei, Argentinia; FRENZEL, Bibl. Zool. xii, p. 49, 1 figg. 1-14.
- Mastigella, n. g. with M. polymastix, & sp., Argentinia; FRENZEL, 1 Zool. xii, p. 38, pl. ii, fig. 3.
- Mastigina, n. g. with M. chlamys, p. 42, pl. iv, figg. 3-7, paramylon, ppl. ii, fig. 7, n. spp., Argentinia, Frenzel, Bibl. Zool. xii.
- Micromastix, n. g. with M. januarii, & sp., Argentinia; FRENZEL, Zool. xii, p. 37, pl. ii, fig. 8.
- Pontomyxa, n. g. with P. flava, & sp. Banyuls, France, on Microco. sabatieri; TOPSENT, C.R. cxiv, p. 774.
- Saccamæba, n. g. with S. punctata, p. 2, pl. iii, figg. 5 & 6, lucens, pl. i, fig. 11, magna, p. 8, pl. iii, fig. 19, cubica, p. 13, pl. iv, figg. & 13, morula, p. 14, pl. i, fig. 10, renacuajo, p. 16, pl. i, figg. 7 cirrifera, p. 11, pl. ii, fig. 6, n. spp., Argentinia, FRENZEL, J Zool. xii.
- Tricholimax, n. g. with T. hylae, & sp., Argentinia; FRENZEL, Bibl. 2 xii, p. 35, pl. iii, figg. 2-4.

#### Class 5.—HELIOZOA.

- Actinophrys alveolata, Sydney, SCHEWIAKOFF, Verh. Ver. Heidelb. (n.s. iv, p. 549, n. sp.
- Astrodisculus araneiformis, Melbourne, Schewiakoff, Verh. Ver. Heidelb. (n.s.) iv, p. 549, n. sp.
- Eleorhanis arenosa, Argentinia, FRENZEL, Bibl. Zool. xii, p. 72, pl. vi, fig. 12, n. sp.
- Estrella, n. g. with E. aureola, p. 74, pl. vi, fig. 16, socialis, p. 75, pl. vi, figg. 5 & 7, n. spp., Argentinia, FRENZEL, Bibl. Zool. xii.
- Heliosphærium, n. g. with H. aster, p. 76, pl. vi, figg. 10 & 11, and pl. x, fig. 4, polyedricum, p. 79, pl. vi, figg. 6, 9, & 17, and pl. x, fig. 5, n. spp., Argentinia, Frenzel, Bibl. Zool. xii.
- Lithosphærella, n. g. with L. compacta, & sp., Argentinia, FRENZEL, Bibl. Zool. xii, p. 73, pl. vi, figg. 13 & 15.
- Monobia solitaria, Sandwich Is., Oahu, Schewiakoff, Verh. Ver. Heidelb. (n.s.) iv, p. 548, n. sp.
- Nuclearella, n. g. with N. variabilis, & sp., Argentinia, FRENZEL, Bibl. Zool. xii, p. 63, pl. i, figg. 1 & 2, pl. ii, figg. 1, 2, & 10, and pl. iv, fig. 16.
- Nuclearia polypodia, Malay Archipelago, Bali, Schewiakoff, Verh, Ver. Heidelb. (n.s.) iv, p. 548, n. sp.
- Nuclearina, n. g. with N. leuckarti, & sp., Argentinia, FRENZEL, Bibl. Zool. xii, p. 59, pl. vi, figg. 4, 8, & 18.

### Class 6.—RETICULARIA.

- Amphicoryne parasitica, SCHLUMBERGER, Mém. Soc. Zool. v, p. 211, pl. vii, figg. 10-12, Azores, depth 130 metres, n. sp.
- Biloculina sarsi, p. 553, pl. ix, figg. 55-59, woodcuts 10-12, North Sea, 2000 faths.; labiata, p. 556, pl. ix, figg. 60-62, woodcuts 13 & 14, Mediterranean, 555 metres; bradyi, p. 557, pl. x, figg. 63-71, woodcuts 15-19, Atlantic, Bay of Biscay, 1850 metres; vespertilio, p. 561, pl. x, figg. 74-76, woodcuts 20-22, Bay of Biscay, 1850 metres; fischeri, p. 563, pl. xi, figg. 77 & 78, woodcuts 23-25, Bay of Biscay, 1850 metres; milne-edwardsi, p. 567, pl. xi, figg. 79 & 80, woodcuts 29 & 30; pisum, p. 569, pl. xi, figg. 81-83, woodcut 31, Mediterranean; anomala, p. 569, pl. xi, figg. 84-86, and pl. xii, fig. 101, woodcuts 32-34, Mediterranean, 555 metres; Schlumberger, Mém. Soc. Zool. iv: n. spp.
- †Bolivinopsis, n. g. and B. capitata, Cretaceous, Russia; JAKOWLEW, Trav. Soc. Univ. Karkow, xxiv, p. 349, n. sp.
- † Bulimina brevicona, p. 54, pl. iv, figg. 1 a, b, inflata, p. 55, pl. iii, figg. 4 a, b, c, conoidea, p. 55, pl. iii, figg. 5 a, b, depressa, p. 55, pl. iii, figg. 3 a, b, Cenoman, Kamajk, Bohemia, Perner, Palsontographica Bohemis, i: n. spp.
- †B. presli, Reuss, var. sabulosa, n. var.; Chapman, J. R. Micr. Soc. 1892, p. 755, pl. xii, fig. 5, Gault of Folkestone.

Discorbina minutissima, Shore mud of Southport, CHASTER, Rep. So

port Soc. i, p. 65, pl. i, fig. 15, n. sp.

† Frondicularia fritschi, p. 58, pl. vii, figg. 1 a, b, c, Cenoman, Kan Bohemia, coronata, p. 58, pl. vii, figg. 2 a, b, Cenoman, Gangl Bohemia, linea, p. 59, pl. vii, figg. 3 a, b, parallela, pl. 59, pl. vii, fig. 4, b, Kamajk, obsoleta, p. 59, pl. vii, figg. 6 a, b, conica, p. 59, pl. figg. 7 a, b, bohemica, p. 60, pl. vii, fig. 4, quadrigona, p. 60, pl. vii, fig. a, b, incerta (n. sp. ?), p. 60, pl. vii, figg. 10 a, b, gracilis, p. 60, pl. fig. 9, lanceolata, p. 60, pl. vii, figg. 12 a, b, foliacea, p. 61, pl. figg. 13 a, b, acutangula, p. 61, pl. vii, fig. 11, Kamajk, Per Palseontographica Bohemise, i: n. spp.

†Gaudryina dispansa, Gault of Folkestone, CHAPMAN, J. R. Micr. 1892, p. 753, pl. xi, figg. 10 a, b, n. sp.

- †Haplophragmium elegans, n. sp., p. 322, pl. v, fig. 10, H. latidorse Born, var. papillosa, n. var., p. 323, pl. v, fig. 13, Gault of Festone; Chapman, J. R. Micr. Soc. 1892.
- †Haplostiche oligostegia, PERNER, Palæontographica Bohemiæ, i, I pl. ii, figg. 13 & 14, Cenoman, Gangberg, Bohemia; H. sherbora Gault of Folkestone, CHAPMAN, J. R. Micr. Soc. 1892, p. 325, p figg. 6-8: n. spp.

Lagena millettii, p. 61, pl. i, fig. 10, falcata, p. 61, pl. i, fig. 7, depr p. 62, pl. i, fig. 5, protea, p. 62, pl. i, fig. 14, Shore mud, South Chaster, Rep. Southport Soc. i, n. spp.

Lingulina herdmani, Shore mud, Southport, CHASTER, Rep. South Soc. i, p. 63, pl. i, fig. 9, n. sp.

†Lituola cylindrica, p. 52, pl. ii, figg. 7-12, Cenoman, Kamajk, Bohe globigerinoides, p. 52, pl. ii, figg. 15-17, Perner, Palseontograf Bohemise, i, n. spp.

†Miliolina maggii, Pliocene, River Arno, Corti, p. 66, fig. 1a,

scient. xiv, n. sp.

† Nodosaria longispina, Cretaceous, Russia, Jakowlew, Trav. Soc. I Kharkow, p. 343; N. mayeri, Cretaceous, Bohemia, Perner, Böhm. Ges. 1892, p. 43; cryptostegia, p. 56, pl. vi, fig. 2, Cenor Kamajk, Bohemia, clarata, p. 56, fig. 4, subnodosa, p. 57, fig pseudaffinis, p. 57, figg. 6 a, b, bohemica, p. 57, figg. 7-9, diver, p. 57, fig. 11, Kamajk, Perner, Palseontographica Bohen n. spp.

† Nummulites javanus, Java, Verbeek, Nat. Tijdschr. Nederl. Inc

p. 105, pl. i, figg. 1-3, n. sp.

†Polymorphina orbignii, Zbor. sp., cercicornis, n. var., Gault of Fostone; Chapman, Geol. Mag. (n.s.) (3) ix, p. 54, pl. ii, figg. 5 & 6 Psammonyx, n. g. with P. vulcanicus, & sp., Sagami Bay, Japan, 100

faths.; Döderlein, Verh. deutsch. Zool. Ges. ii, p. 145.

Pulvinulina nitidula, Shore mud, Southport, Chaster, Rep. South Soc. i, p. 66, pl. i, fig. 17, n. sp.

H

- Ramulina grimaldii, Fosse de Fayal, between Fayal and Pico (Azores), 130 metres, Schlumberger, Mém. Soc. Zool. iv, p. 509, pl. v, n. sp.
- †Reophax lageniformis, p. 319, pl. v, figg. 1 a, b, folkestoniensis, p. 321, pl. v, figg. 6 a, b, cylindruceu, p. 321, pl. v, figg. 7 a, b, Gault of Folkestone, Chapman, J. R. Micr. Soc. 1892, n. spp.
- Rheophax scotti, Shore mud, Southport, Chaster, Rep. Southport Soc. i, p. 57, pl. i, fig. 1, n. sp.
- †Rotalia eroluta, Pliocene, River Arno, Corti, Boll. scient. xiv, p. 67, figg. 2 a, n. sp.
- †Spirillina minima, p. 159, pl. i, fig. 4, trochiformis, p. 159, pl. i, fig. 3, Cretacoous, Helle Mühle near Moltzow, Mecklenburg; Schacko, Arch. Ver. Mecklenb. xlv.
- †Spiroculina, n. sp., Schacko, Arch. Ver. Mecklenb. xlv, p. 158, pl. i, fig. 1, Cretaceous, Helle Mühle near Moltzow, Mecklenburg.
- Textularia fusiformis, Shore mud, Southport, CHASTER, Rep. Southport Soc. i, p. 58, pl. i, fig. 3, n. sp.
- †T. brevicona, p. 54, pl. ix, fig. 12, parallela, p. 54, pl. ix, fig. 13, Cenoman, Kamajk, Bohemia, Perner, SB. Böhm. Ges. 1892, n. spp.
- Trichosphærium sieboldii, Schn., belongs, according to GREEFF, Zool. Anz. xv, (1892) pp. 62-64, to the monothalamous Foruminifera.
- Triloculina aspergillum, Azores, Schlumberger, Mém. Soc. Zool. v, p. 207, pl. viii, figg. 8 & 9, 1 woodcut, n. sp.
- † Trochammina concava, Gault of Folkestone, Chapman, J. R. Micr. Soc. 1892, p. 327, pl. vi, figg. 14 a, b; T. serpuloides, Cretaceous, Helle Mühle near Moltzow, Mecklenburg, Schacko, Arch. Ver. Mecklenb. xlv, p. 159, pl. 1, fig. 5: n. spp.
- † Vitriwebbina, n. g. with V. sollasi, & sp., Gault of Folkestone; Charman, Geol. Mag. (3) ix, p. 53, pl. ii, figg. 1-3.

#### Class 7.—RADIOLARIA.

- †Acantosphæra entactinia, p. 147, pl. xiii, fig. 1, Carboniferous, Hartz Mountains, A. macracantha, Silurian, Cabrières, p. 147, pl. xiii, fig. 2, Rüst, Palmontogr. xxxviii, n. spp.
- †Acrosphæra herzynica, Devonian, Schæbenholz, p. 140, pl. ix, fig. 3, A. glitzii, Carboniferous, Hartz Mountains, p. 140, pl. ix, fig. 2, Rüst, Palæontogr. xxxviii, n. spp.
- †Actinomma schæbenholzianum, Carboniferous, Hartz Mountains, Rüst, Palæontogr. xxxviii, p. 148, pl. xiv, fig. 5, n. sp.
- † Amphibrachium inaquale, p. 170, pl. xxiii, fig. 10, A. bacillum, p. 169, pl. xxiii, fig. 9, A. pulchellum, p. 169, pl. xxiii, fig. 8, Devonian, South Ural, A. devoniense, Carboniferous, South Ural, p. 169, pl. xx, fig. 8, A. desecatum, Carboniferous, Sicily, p. 169, pl. xx, fig. 9, Rüst, Palmontogr. xxxviii; † A. crassum, pl. v, fig. 9, truncatum, fig. 10, fragile, fig. 11, p. 223, Fanny Bay, Port Darwin, Hinde, Q. J. Geol. Soc. xlix: n. spp.
- † Amphymenium alienum, pl. xxiv, fig. 2, A. krautii, fig. 1, Devonian, Schebenholz, Rüst, Palsontogr, xxxviii, p. 170, n. spp

- \*Archelicom loss, Carboniferous, Hartz Mountains, Röst, Paleont xxxviii, p. 186. pl. xxii. fig. 5, n. sp.
- \*Aprilipen Impicubacia. Carbaniferom, Sicily, Röst, Paleoni xxxvii. p. 152 pl. xvi. fig. 6, n. sp.
- Communication, Atlantic, Boussert, Ergeb. Plankt. Exp. p. pl. vi. fig. 4, 2, 49.
- \*Coryamphores enfruerinate, pl. vii, fig. 3. C. jejana, fig. 1, C. macracti fig. 4. C. magnes, fig. 2, p. 136. Carboniferous, Hartz Mountains pyrman, p. 135. pl. vi. fig. 13, Carboniferous, Hartz Mountains from J. R. 187. Palmontogy, xxxviii, n. spp.
- \*Carpuspisers graddecisis, Devonian, Schubenholz, Röst, Palaent xxxxii, p. 130. pl. ix. fig. 1. n. sp.
- †Correction infundibulum, Carboniferous, Sicily, p. 184, pl. xxvii, 10, C. silpopressa, Permina, Nougorod, p. 183, fig. 9, RCsr, Palson xxxvii. 2. app.
- †Concilipas direculpura. Silurian. Cabrières. p. 151, pl. xv, fig. 5. C. e formes, Devonian, Schubenholz, p. 152, pl. xvi, fig. 1, C. pusilla, T. B. tchelstein, pl. xvi, fig. 5, C. ercolata, pl. xv, fig. 10, C. citrus, f. p. 151, Devonian. South Ural. C. multiplex, Carboniferous, I. Mountains. pl. xvi, fig. 3, C. reticulosa, Carboniferous, Hartz M. tains. fig. 2, C. cypridina, pl. xv, fig. 9, p. 152, C. rectipora, fig. curvatopicata, fig. 7. Devonian, South Ural, C. perovalia, Carl ferous. Hartz Mountains, pl. xvi, fig. 4, p. 151, Rüst, Palson xxxvii. n. 550.
- \*Constante intermedius, pl. xix, f.g. 19, C. nummularis, fig. 9, p. 162, boniferous, Harts Mountains, C. primordialis, Silurian, Cabri p. 161, fig. 8, Rfst. Paleontogr. xxxviii, n. spp.
- +Crossphere autorana, Trias, Astura, pl. vi, fig. 7, C. bakonyana, T Fels: E'rs. Bakony, fig. 4, C. carbonica, fig. 2, C. ingens, fig. 8, boniferous, Hartz Mountains. C. castanea, Devonian, Schæben Hartz Mountains, fig. 5, C. wralensis, fig. 3, C. rossica, fig. 9, p. C. apiaria, fig. 6, p. 135. Devonian, South Ural, C. macropora, f p. 134. Silurian, Cabrières, Röst, Palæontogr. xxxviii, n. spp.
- Challengeron rottenburgi, pl. vi. fig. 1, neptuni, p. 180, figg. 2 & 3, Irmi Sea, Atlantic, BORGERT, Ergeb. Plankt. Exp., n. spp.
- †Cornulanus cornu-raccinum, Trias, Röthelstein, pl. xxvi, fig. 1 dilatata, fig. 12. C. macropora, fig. 13, Carboniferous, Sicily, I Palæontogr. xxxviii. p. 180, n. spp.
- †Cromyolruppa prunulina, Carboniferous, Sicily, Röst, Palseon xxxviii, p. 157, pl. xviii, fig. 5, n. sp.
- †Cromyomma grandærum, Silurian, Cabrières, Rüst, Palæontogr. xx: p. 149, pl. xiv, fig. 7, n. sp.
- † Cromyosphæra alternans, Carboniferous, Sicily, pl. viii, fig. 7, C. dis Devonian, South Ural, fig. 6, C. eminens, fig. 8, p. 139, C. frequens, C. radiata, Carboniferous, Hartz Mountains, fig. 5, C. petsc. Carboniferous, Ural, pl. viii, fig. 3, C. prisca, Silurian, Cabr fig. 4, p. 138, Rüst, Palsontogr. xxxviii, n. spp.

- †Crucidiscus præcursor, Silurian, Cabrières, Rüst, Palæontogr. xxxviii, p. 163, pl. xx, fig. 6, n. sp.
- †Cyphanta piscis, p. 160, pl. xix, fig. 5, C. quiniseriata, p. 161, fig. 6, Devonian, South Ural, Rüst, Palmontogr. xxxviii, n. spp.
- †Cyrtoculpis serieporata, pl. xxvi, fig. 16, C. obtusa, fig. 15, Devonian, South Ural, C. prima, fig. 14, Carboniferous, Hartz Mountains, Rüst, Palæontogr. xxxviii, p. 180, n. spp.
- † Dictyocephalus ventricosus, Carboniferous, Sicily, Rüst, Palmontogr. xxxviii, p. 182, pl. xxvii, fig. 5, n. sp.
- † Dictyomitra magnifica, pl. xxix, fig. 1, D. micropora, pl. xxviii, fig. 11 & pl. xxix, fig. 2, p. 188, Carboniferous, Sicily, D. pumilio, Devonian, Schæbenholz, pl. xxviii, fig. 10, p. 187, D. sasinana, Siliceous Limestone, Astura, p. 188, pl. xxix, fig. 4, Rüst, Palæontogr. xxxviii; D. australis, pl. v, fig. 14, triangularis, pl. v, fig. 15, Fanny Bay, Port Darwin, Hinde, p. 224, Q. J. Geol. Soc. xlix: n. spp.
- † Dictyophimus dubius, Carboniferous, Sicily, RUST, Palseontogr. xxxviii, p. 181, pl. xxvii, fig. 3, n. sp.
- † Dizonium laeve, Carboniferous, Sicily, Rüst, Palæontogr. xxxviii, p. 174, pl. xxv, fig. 8, n. sp.
- †Druppatractus dictyococcus, Carboniferous, Hartz Mountains, Rüst, Palæontogr. xxxviii, p. 158, pl. xviii, fig. 9, n. sp.
- †Druppula amygdalina, Carboniferous, Petschora, p. 155, pl. xvii, fig. 7, D. angustiporata, pl. xviii, fig. 1, D. cornus, fig. 2, p. 156, Carboniferous, Bükk Mountains, D. andreana, pl. xvii, fig. 5, p. 155, D. berberis, fig. 11, D. crucifera, fig. 10, p. 156, D. endechinata, fig. 9, p. 155, D. fxata, Carboniferous, Sicily, fig. 12, p. 156, D. pomatia, Trias, Astura, fig. 6, D. cembra, Devonian, South Ural, fig. 8, p. 155, D. silurica, Silurian, Langenstriegis, fig. 4, p. 154, Rüst, Palæontogr. xxxviii, n. spp.
- †Dystympanium carbonicum, Carboniferous, Sicily, Rüst, Palæontogr. xxxviii, p. 177, pl. xi, fig. 9, n. sp.
- † Echinomma oligacanthum, Carboniferous, Sicily, Rüst, Palæontogr. xxxviii, p. 149, pl. xiv, fig. 6, n. sp.
- † Ellipsidium aculeatum, p. 152, pl. xvi, fig. 7, E. spinosum, p. 153, pl. xvii, fig. 1, Carboniferous, Hartz Mountains, Rüst, Palæontogr. xxxviii, n. spp.
- † Ellipsostylus obliquus, Carboniferous, Sicily, Rüst, Palæontogr. xxxviiis, p. 153, pl. xvi, fig. 11, n. sp.
- † Ellipsoxiphus hystrix, Carboniferous, Hartz Mountains, pl. xvi, fig. 10, E. procerus, Carboniferous, Lerbach, fig. 9, Rüst, Palæontogr. xxxviii, p. 153, n. sp.
- Euphysetta lucani, Atlantic, BORGERT, Ergeb. Plankt. Exp., p. 182, pl. vi, fig. 8, n. sp.
- † Halicalyptra aculsata, Carboniferous, Petschora, Rüst, Paleontogr. xxxviii, p. 179, pl. xxvi, fig. 7, n. sp.
- † Halicapsa megapora, Carboniferous, Sicily, Rüst, Palssontogr. xxxviii, p. 181, pl. xxvii, fig. 2, n. sp.

- \*Hallmanne operature, Carboniferent, Sielly, pl. xiv, fig. 2, H. phorum. Carboniferent, Hartz Mouxtains, pl. xiii, fig. 4, Röst, Palmontoge, xxxviii. 2, app.
- \*Helmborns ornouress. Sikrika, Cabrieres, pl. xxii, fig. 1. H. orto Carinomierross. Hartz Mountains, fig. 2. River, p. 165, Pales xxxvii. 2. opp.
- "He monast uniformies. Carboniferous, Hartz Mountains, pl. riv H. romen. Silvina. Cabrieros, Sy. 3. Rist, p. 184, Paleo Exercis. 2. 199.
- "Heinsphert berbeum, pl. xir, fig. 3. H. lýrslé, pl. xiv, fig. 1. Carbonierom, Harta Mannaian, Rier, Paleontogr. xxxviii,
- \*Hersing van ofene Carismiferens, Bükk Mountains, Rüst, Pales axxvii. 2. 122, pl. axiv. fig. 9. n. sp.
- \*Henrimeia jainenouse. Devraian. Schubenholz, pl. xii. fig. 7, H. Carimuiferous. Secty. fig. 5, p. 146, Riss. Paleontogr. xxxviii,
- \*Lospiners deconicus. A. Devonian. South Ural, pl. vii. fig. 5, L. Silmian. Cadrieres. Sg. 4, p. 126. River, Palmontogr. xxxviii, 1
- \*Lehenura elevirum Silminn Cabrières, Rist, Palmontogr. 1 2. 154 31. xvii. for 3. n. sp.
- \*Lokarecas ironiques. pi. xviii, fig. 7. L. perforatus, p. 158, pl fig. 3. Carboniferous. Secily. L. mojeri, Carboniferous, Hartz mins, fig. 4. p. 157. Rfsr. Paleoutogr. xxxviii, n. spp.
- \*Leisiss éficies. Carboniferous. Hartz Mountains, Rêst, Palas xxxvin. a. 175. pl. xxvi. Sgg. 1 & 2. n. sp.
- \*Leiner mye inseem. pl. xxix, fig. 6. L. techernytechevii, fig. 8, L. techernytechevii, fig. 8, L. techernytechevii, fig. 8, P. techernytechevii, fig. 8, P. techernytechevii, fig. 10, pyramia, L. syberrospheriou, fig. 9. L. techertoma, p. 189, fig. 10, Carboni Seely, L. techer, Carboniferous, Bükk Mountains, p. 190, 1 Rist. Palmontogr. xxxviii: L. fusiformia, Port Darwin, 1 Q. J. Geol. Soc. xiix, p. 225, pl. v. fig. 16.: n. spp.
- †Lubergelis marroccess. Devonian, Schwbenholz, pl. xxii, fig. 4, L. Carboniferous, Hartz Mountains, pl. xxii, fig. 3, p. 166. Palmontogr. xxxviii; L. eride. Fanny Bay. Port Darwin, J. Q. J. Geol. Soc. xlix. p. 223, pl. v. fig. 8, : n. spp.
- \*Labonespiller kipoleris, Carboniferous, Hartz Mountains, p. 154.

  fig. 12, L. kezzezetkus, Silurian, Langenstriegis, fig. 8, p. 1

  stekenorth, Devonian, Schwbenholz, p. 154, pl. xvii, fig. 2,
  Palmentogr, xxxviii, n. spp.
- \*Laisuites forficula, Siliceous Limestone, Röthelstein, Rüst, Palse xxxviii, p. 188, pl. xxix, fig. 3, n. sp.
- \*Leident-dee mea lieadi. Carboniferous, Sicily, Rüst, Paleontogr. 1 p. 187. pl. xxviii, fig. 9, n. sp.
- \*\*Glostophera ecklostatets. Silurian, Cabrières, Rüst, Palse xxxviii, p. 140, pl. ix, fig. 4, n. sp.
- †Port dram primiticum, Carboniferous, Būkk Mountains, 27. xxxviii, p. 172, pl. xxiv, fig. 8, n. sp.

- † Peripanartus deficiens, Devonian, South Ural, Rüsr, Palmontogr. xxxviii, p. 161, pl. xix, fig. 7, n. sp.
- †Phænoscenium excentricum, Carboniferous, Hartz Mountains, Rüst, Palæontogr. xxxviii, p. 179, pl. xxvi, figg. 8-10, n. sp.
- †Phorticium macropylium, Silurian, Langenstriegis, Rüst, Palæontogr. xxxviii, p. 175, pl. xxvi, figg. 4 & 5, n. sp.
- †Porodiscus cabrierensis, Silurian, Cabrières, pl. xxiii, fig. 3, P. intricatus, p. 168, fig. 5, P. rossicus, pl. xxii, fig. 6, p. 168, Devonian, South Ural, P. paronæ, pl. xxiii, fig. 2, P. parvulus, pl. xxii, fig. 7, p. 167, P. subspiralis, p. 168, pl. xxiii, fig. 4, Muschelkalk, Felsö Eörs, P. triassicus, Muschelkalk, Röthelstein, p. 167, fig. 1, Rust, Palæontogr. xxxviii, n. spp.
- † Prismatium paradictyum, Carboniferous, Sicily, Rüst, Palssontogr. xxxviii, p. 178, pl. x, fig. 10, n. sp.
- †Prunulum armeniacum, Devonian, Schæbenholz, pl. xviii, fig. 4, P. murrayi, Carboniferous, Hartz Mountains, fig. 3, p. 157, Rüst, Palæontogr. xxxviii, n. spp.
- † Pterocanium tenuecostatum, Carboniferous, Sicily, Rüst, Palmontogr. xxxviii, p. 183, pl. xxvii, fig. 8, n. sp.
- † Rhizoplegma betulinum, Carboniferous, Sicily, Rüst, Palsontogr. xxxviii, p. 150, pl. xv, fig. 2, n. sp.
- † Rhodosphæra crucifera, Carboniferous, Hartz Mountains, pl. vii, fig. 11, R. idonea, fig. 9, p. 137, R. pulchra, p. 138, pl. viii, fig. 1, Carboniferous, Sicily, R. devoniensis, Devonian Schæbenholz, p. 137, pl. vii, fig. 10, Rüst, Palæontogr. xxxviii, n. spp.
- †Rhopalastrum carbonicum, Carboniferous, Hartz Mountains, p. 170, pl. xxiv. fig. 3, R. giganteum, fig. 6, R. isselii, fig. 5, R. siculum, fig. 4, p. 171, Carboniferous, Sicily, Rüst, Palæontogr. xxxviii, n. spp.
- †Rhopalodictyum astrictum, Devonian, Schæbenholz, Rüst, Palæontogr. xxxviii, p. 174, pl. xxv, fig. 7, n. sp.
- †Sethocapsa obstipa, Devonian, South Ural, p. 182, pl. xxvii, fig. 6, S. occlusiva, Muschelkalk, Felső Eörs, p. 183, fig. 7, Rüst, Palæontogr. xxxviii, n. spp.
- †Sethocyrtis excisa, Carboniferous, Hartz Mountains, Rüst, Palssontogr. xxxviii, p. 182, pl. xxvii, fig. 4, n. sp.
- †Sethostaurus exsculptus, Silurian, Cabrières, Rüst, Palmontogr. xxxviii, p. 165, pl. xxi, fig. 7, n. sp.
- † Sphæropyle, n. g., Dreyer, p. 140, with † S. dreyeri, p. 141, pl. ix, fig. 6, S. lævis, p. 140, fig. 5, Devonian, South Ural, S. entostoma, fig. 9, S. eurystoma, fig. 8, Carboniferous, Sicily, S. multangularis, Carboniferous, Hartz Mountains, fig. 7, p. 141, Rüst, Palæontogr. xxxviii, n. spp.
- †Spirocapsa, n. g. for †S. singularis, n. sp., Carboniferous, Sicily; RUST, Palmontogr. xxxviii, p. 192, pl. xxx, fig. 13.
- †Spironium hæckelii, Carboniferous, Hartz Mountains, Rüst, Palsontogr. xxxviii, p. 175, pl. xxvi, fig. 3, n. sp.

- Spangarrana nonstigua Sincian, Cabrieras, pl. xix. fig. 6, 6, 4 open. Anthonolleman. Scaly. pl. xix. fig. 4, p. 169, Rist, Palaxerii. 1, 499.
- "Igaameye's suradicipus. Carbonidarson. Harra Mountains Passanage excerii. p. 18'... ps. xxv2. fig. 1. s. sp.
- \* Igangodienyas tropatrum, Derminia, Santh Ural, Röst, Palazzeili, p. 140, pl. viii, fq. 10, n. sp.
- Theory discuss descriptions described and the second section of the se
- Symposium urudiame, 3 190, 31. xix. fig. 5. S. trimilista, fig. Curamiferous, Harry Monacuus, R\*\*\*, Princescopy, xxxvii
- "insequence symmetries. Piet Durwin, Haven, Q. J. Gool. 5 p. ma. pl. = fig. 13, p. 69.
- Thankrightere demices. Minerhelleule, Feins Edes, Röst, Pa exerti. 3. II. 31. ext in 3. a.m.
- The manner of the last of the
- \* hampmorums tainder pous. Carboniferous. Hertz Mountain ig. 1. f. repasseus. Carboniferous, Sicily. pl. aviii, fig. 1 2. fr. Frimmung. axaviii. 1. sp.
- "Spanyraghers minne. Dreminn, South Ural, Rist, Palmontogr 2. 150. 24. 27, fug. 3 & 4, 2, sp.
- "Francourage recomments, Carboniferous, Hartz Mountains Promotories recoming, 1773, pl. xxv. fig. 4, e. sp.
- \*\*iponoprarieina elempation Carinotiferous, Sietly, Röst, Pa xxxviii, 2, 173, 36, xxv fig. i, 2, 59.
- Theoremse Sciences. Terminal Schabenholz, pl. aviii, fig. 12, 1 former, fig. 11, p. 150. Section Lie. Sig. 10, p. 158, Carbo Socily, River, Pulsonness, account a so.
- #Steen on them outpouls. Carboniferous. Stelly, pl. nii, fig. 4. planes. fig. 3. S. roph.giorean. fig. 5. Carboniferous, Hartz M B.(et. p. 145. Palgeonogy. exercit. a. sep...
- \*Some alementa accione at Carbonifectus, Sicily, Röst, Pal xxxxxii p. 171, pl. xxiv. fig. 7, m. sp.
- \*State of trade & authority, Trian Felni Elen, pl. xi, fig. 6, S. mac fig. 1, campain, pl. xii, fig. 1, Carboniferous, Sicily, spinigera, ferous, Hartz Mountains, pl. xi, fig. 4, p. 144, S. micropora, Cabrutess, p. 143, fig. 3, Röst, Palsontogr, xxxviii, m. spp.
- †Steorologelides a coloni, Carboniferous, Hartz Mountains, Rüst vogt. xxxviii, p. 144. pl. xii, fg. 2. n. sp.
- †Someophara frazilis, pl. x. fig. 4, S. quadrangularis, fig. 5, Cabrieres, setispina, Carboniferous, Petschora, fig. 6, Rüst Palæontogr, xxxviii, n. s; p.
- †Staurostylus tenuispinus. Carboniferous, Hartz Mountains, pl.; ziphophorus, Carboniferous, Petschora, fig. 1, Rüst, p. 143, togr. xxxviii, n. spp.

- †Stichocapsa anguillula, pl. xxx, fig. 11, S. elpenor, p. 192, fig. 10, S. tænia, pl. xxx, fig. 12, p. 192, Devonian, South Ural, S. biceps, fig. 7, S. citriformis, fig. 8, S. ellipsoides, fig. 6, S. gravida, fig. 1, pupa, fig. 9, Carboniferous, Sicily, S. bükkiana, Carboniferous, Bükk Mountains, fig. 5, p. 191, Rüst, Palseontogr.xxxviii; S. pinguis, pl. v, fig. 17, S. chrysalis, pl. v, fig. 18, p. 225, Port Darwin, Hinde, Q. J. Geol. Soc. xlix: n. spp.
- †Stichopera aculeata, Carboniferous, Sicily, Rüst, Palsontogr. xxxviii, p. 186, pl. xxviii, fig. 7, n. sp.
- †Stigmosphæra mira, pl. vi, fig. 11, S. rothpletzii, fig. 10, S. suspecta, fig. 12, Carboniferous, Hartz Mountains, Rüst, Palæontogr. xxxviii, p. 135, n. spp.
- †Stigmosphærostylus, n. g. for S. notabilis, n. sp., Carboniferous, Hartz Mountains, Rüst, Palæontogr. xxxviii, p. 142, pl. x, fig. 2.
- †Stilodictya polaris, p. 169, pl. xxv, fig. 1, S. zittelii, p. 168, pl. xxiii, figg. 6 & 7, Devonian, Schæbenholz, Büst, Palæontogr. xxxviii, n. spp.
- † Stylodiscus dictyosus, Carboniferous, Hartz Mountains, Rüst, Palæontogr. xxxviii, p. 162, pl. xx, fig. 1, n. sp.
- †Stylosphæra acuta, Carboniferous, Hartz Mountains, Rüst, Palæontogr. xxxviii, p. 142, pl. x, fig. 3, n. sp.
- †Stylotrochus dolfussi, Carboniferous, Sicily, Rüst, Palmontogr. xxxviii, p. 173, pl. xxv, fig. 5, n. sp.
- † Tetrapylonium ovatum, Carboniferous, Sicily, Rüst, Palæontogr. xxxviii, p. 174, pl. xxv, fig. 9, n. sp.
- † Thecosphæra sexactis, Carboniferous, Hartz Mountains, pl. vii, fig. 8, P. sicula, Carboniferous, Sicily, fig. 7, Rüst, p. 137, Palæontogr. xxxviii, n. spp.
- † Theocampe bulbosa, Carboniferous, Sicily, Rüst, Palmontogr. xxxviii, p. 185, pl. xxvii, fig. 12, n. sp.
- † Theocyrtis apis, Muschelkalk, Astura, Rüst, Palssontogr. xxxviii, p. 184, pl. xxvii, fig. 11, n. sp.
- † Theodiscus brachyacanthus, pl. xx, fig. 2, T. convexus, fig. 3, Carboniferous, Hartz Mountains, T. gigus, fig. 4, T. planus, fig. 5, Silurian, Cabrières, Rüst, p. 163, Palæontogr. xxxviii, n. spp.
- †Theosyringium præcox, Carboniferous, Sicily, Rüst, Palmontogr. xxxviii, p. 184, pl. xxviii, fig. 1, n. sp.
- † Tricolocapsa abdominalis, pl. xxviii, fig. 4, T. thoracica, fig. 3, p. 186, Carboniferous, Sicily, T. anceps, fig. 6, T. grandæra, fig. 5, p. 185, Carboniferous, Hartz Mountains, Rüst, Palæontogr. xxxviii, n. spp.
- †Tripilidium dissociatum, Carboniferous, Hartz Mountains, Rüst, Palseontogr. xxxviii, p. 178, pl. xxvi, fig. 6, n. sp.
- † Trissocircus primævus, Carboniferous, Sicily, pl. xi, fig. 7, T. quadratus, Devonian, South Ural, pl. x, fig. 9, p. 177, Rüst, Palssontogr. xxxviii, n. spp.
- † Trochodiscus convexus, p. 164, pl. xx, fig. 7, T. longispinus, pl. xxi, fig. 4, p. 165, T. nicholsoni, fig. 3, T. proavus, fig. 1, T. serrula, fig. 2, Carboniferous, Hartz Mountains, T. recurvispinus, Devonian, Scheeben-

- holz, fig. 5, *T. diversispinus*, Silurian, Cabrières, fig. 6, p. 164 Palssontogr. xxxviii, n. spp.
- Tuscarora nationalis, Irminger Sea, Atlantic, 600 metres, BORGERT, Plankt. Exp. p. 182, pl. vi, fig. 7, n. sp.
- †Xiphosphæra macrostyla, Devonian, Hartz Mountains, Rüst, Pala xxxviii, p. 141, pl. ix, fig. 10, n. sp.
- †Xiphostylus ardea, Carboniferous, Hartz Mountains, pl. x, fig. asper, Silurian, Cabrières, pl. ix, fig. 11, p. 142, Rüst, Palst xxxviii, n. spp.
- †Zonodiscus dentatus, Silurian, Cabrières, pl. xix, fig. 11, Z. ma. Carboniferous, Hartz Mountains, fig. 12, Rüst, p. 162, Palæxxxviii, n. spp.
- † Zygocircus priscus, Carboniferous, Hartz Mountains, pl. x, fiş simplicissimus, Carboniferous, Sicily, fig. 7, Rüst, p. 176, Palæ xxxviii, n. spp.
- † Zygosterhanus quadrupes, Carboniferous, Sicily, Rüst, Palse xxxviii, p. 177, pl. xi, fig. 8, n. spp.

### B.—CORTICATA.

### Class 1.—SPOROZOA.

- For a revised classification of the group, see MINGAZZINI (104). Blanchardia, n. g. with B. cypricola, n. sp., parasitic in Cypris ca
  - Solger, p. 99, MT. Vorpomm. xxii.
- Ceratomyxa agilis, pp. 962 & 963, in Trygon vulgaris, Concarneau, diculata, pp. 963 & 964, in Lophius piscatorius, Roscoff, a p. 1091, in Motella tricirrata, Thélohan, C.R. cxv, n. spp.
- Coccidium cruciatum, p. 136, in the liver of Caranx trachurus, Conciminatum, p. 137, in liver, spleen, and kidney of Tench, The J. de l'Anat. Phys. 1892; C. tenellum, in intestinal epithel Fowl, truncatum, p. 249, in kidney of Fowl, RAILLIET & p. 248, Bull. Soc. Z. Fr. xvi: n. spp.
- Cretya neapolitana, in the pyloric appendages of Sphyrana vi Mingazzini, p. 400, fig. 2, Atti Acc. Pontif. Lincei (5), 1, n. s
- Gonobia colubri, in vas deferens and testis of Zamenis virid MINGAZZINI, p. 396, fig. 1, Atti Acc. Pontif. Lincei (5), I, n. 1
- Gregarina satiræ, p. 234, pl. viii, figg. 1-15, in intestine of Statira w. Bl., bergi, p. 286, pl. viii, figg. 16-19, in intestine of Coruficollis, panchloræ, p. 299, pl. viii, figg. 20, in intestine of Panexoleata, blaberæ, p. 300, pl. viii, figg. 21-33, in intestine of I claraziana, Córdoba, Argentinia, FRENZEL, Jen. Z. Nat. xxvi irregularis, Minchin, p. 291, pl. xxvii, Q. J. Micr. Sci. xx bloodvessels of Holothuria nigra, at Plymouth: n. spp.
- Myxidium incurvatum; Thelohan, C.R. exv, p. 1093, in 1 tricirrata.



- Pyxinia crystalligera, in intestine of Dermestes rulpinus, Fabr., and peruvianus, Cateln., Córdoba, Argentinia, Frenzel, p. 314, pl. viii, figg. 34-50, Jen. Z. Nat. xxvii, n. spp.
- Sarcocystis hominis, ROSENBERG, in human heart muscle, S. platydactili, BERTRAM, Zool. Jahrb. v, p. 584, pl. xxxviii, figg. 2-4, in muscle fibres of Platydactylus facetanus, Ald., s. P. mauritanicus, L., from Menorca, n. spp.
- Sphæromyra, n. g. with S. balbianii, n. sp., Thélohan, C.R. cxv, p. 1093, in Motella tricirrata,

#### Class 2.--FLAGELLATA.

KLEBS, Z. wiss. Zool. lv, p. 278, gives the following definition of the group: Flagellata are lower organisms which possess a usually sharply defined uninuclear protoplasmic body, the periplast of which is either a simple cortical layer, or a differentiated plasmic membrane. They are in motion for the greatest part of their life, or at least they always remain capable of such. They all have a specially formed anterior end bearing one or more flagella, and one or more pulsating vacuoles. Reproduction takes place by simple longitudinal fission mostly during the flagellate state, sometimes when at rest. All are able to form resting cysts for a shorter or longer period.

### He proposes the following classification :-

- 1. PROTOMASTIGINA. Families: Rhizomastigina, Monadina, Bikacida, Craspemonadina, Spongomonadina, Amphimonadina, Bodonina.
- 2. POLYMASTIGINA. Families: Trimastigina, Tetramitina, Distomata.
- 3. Euglenoidina. Families: Euglenida, Astasiida, Peranemida.
- 4. CHLOROMONADINA.
- 5. CHROMOMONADINA. Families: Chrysomonadina, Cryptomonadina.
- Anisonema variabile, p. 385, pl. xvii, figg. 5 a, b, striatum, p. 386, figg. 14 a, b, ovale, p. 386, figg. 6 b, c, Basle, Klebs, Z. wiss. Zool. lv, p. spp.
- Astasiida, n. fam., Klebs, p. 356, Z. wiss. Zool. lv.
- Astasiodes lagenula, Malay Archipelago, Bali, Schewiakoff, p. 551, Verh. Ver. Heidelb. (n.s.) iv, n. sp.
- Bodo minimus, p. 310, pl. xiii, figg. 7 a-d, edax, p. 312, figg. 8 a-e, celer, p. 313, figg. 6 a-c, mutabilis, p. 317, pl. xiv, figg. 2 a-c, repens, p. 317, figg. 1 a-c, Basle, Klebs, Z. wiss. Zool. lv, n. spp.
- Chlamydomonas de-baryand, near Moscow, p. 106, pl. i, figg. 9-12, kuteinikowi, p. 117, pl. ii, figg. 9-13, reticulata, p. 124, pl. iii, figg. 1-9, Lichobory and Butyrki, Russia, Goroschankin, Bull. Mosc. 1891, n. spp.
- Chloromonadina, n. fam., Klebs, p. 391, Z. wiss. Zool. lv.
- Chromomonadina, n. fam., KLEBS, p. 394, Z. wiss. Zool. lv.
- Chromulina batalini, Sydney, p. 550, Schewiakoff, Verh. Ver. Heidelb.

7

- 1.2. (# ) 2 servena Bada K.226, 3. 916 pl. xviii, by 6d. Zon. (# ) 2. 100.
- Chrystaness, 1 g. vin C. ruboma, 1 sp. Books, Kiros, p. 497, p. 497, p. 497, p. 497, p. 497, p. 1 sp. 1 see Z. von Zui, 3r.
- They mercent at Z. with the reference at up., Roole, Killin, p. 413, p. 443, p. 445, T. wie. Zoni, by.
- Chaptery: Election Spinery, St. T. Villen, p. 509, Verh. Ver. 1 14, 17, 1, 80.
- Cryptomonus mercus, Luc-sur-Mer. Dan-valad. Le Botaniste, iii, p. 3 4g. 2l., 1. sp.
- Le margan 1922. 3. 3. 3. 3. 122. 3. 3. rediate, p. 3.1. figg. 3. 3. rediate, p. 3.1. figg. 3. 3. Reside. Kless. Z. wiss. Z. 1. 1939.
- Danieryna und sin s.n. Banie, K. 238, p. 414, pl. xviii, figg. 10 a, b, Zon. ir. 2. 19.
- Designate, a. fam., K1228, p. 229, Z. was. Zool, lv.
- Easternam subspecies. Baile. K.1224, p. 399, pl. xvii, figg. 15 a-c, Zork 17, 2, sp.
- Englem e capita. Gazannita, New Zealand, Schewiakoff, p. 55 Vec. Heideld, R.A., iv. R. sp.
- Engineering, z. z. with E. wordt, n. sp., Basile, KLERS, p. 367. fizz. 2 s-d, Z. wise. Zool. lv.
- Heteropeus speck. Bask, KLESS, p. 373. pl. xvii, fig. 12, Z. wis lv. z. sp.
- Heonatta crassa. pl. xv. figg. 9 a, b, fesiformis, pl. xvi. figg p. 339, parties pl. xv. figg. 6 a, b, name, figg. 8 a, b, p. 334 Kieze, Z. wim. Zool, lv. n. spp.
- Hoomangote. n. g., with H. elongatum, n. sp., parasitic in Terifag. 4; Grassi, p. 36, Atti Acc. Pontif. Lincei (5) i.
- Jania. n. g., with J. annectens, n. sp., parasitic in Calotermes fle Gravi, p. 35. Atti Acc. Pontif. Lincei (5) i.
- Marsujurguster. n. g.. with M. striata, n. sp., Oahu, Schewiakoff Verh. Ver. Heidelb. (n.s.) iv.
- Mastigamæba būtschlii. p. 298, pl. xiv, figg. 23 a, b, invertens, p. 297 figg. 1 a-c, Basle, Klebs, Z. wiss. Zool. lv, n. spp.
- Mastigosphæra, n. g., with M. gobii, Tarawera, New Zealand, Sc κορρ, p. 552, Verh. Ver. Heidelb. (n.s.) iv.
- Maupasia, n. g., with M. paradora, n. sp., Hawaii, Kilauea, 46 Schewiakoff, p. 552, Verh. Ver. Heidelb. (n.s.) iv.
- Microjania, n. g., with M. hexamitoides, n. sp., parasitic in Tern fugus, Grassi, p. 36, Atti Acc. Pontif. Lincei (5) i.
- Monas obliqua, New Zealand, Taupo Lake, Schewiakoff, p. 55 Ver. Heidelb. (n.s.) iv, n. sp.
- Monocercomonas termitis, parasitic in Termes lucifugus and Ce flucicollis, Grassi, p. 36, Atti Acc. Pontif. Lincei (5) i, n. sp.
- Ochromonas mutabilis, pl. xviii, figg. 2 & 3 a, b, crenata, pl. x 4 a-e, p. 411, Basle, Klebs, Z. wiss. Zool. lv, n, spp.

- Oikomonas excavata, Sydney, Schewiakoff, p. 549, Verh. Ver. Heidelb. (n.s.) iv, n. sp.
- Peranemida, n. fam., Klebs, p. 362, Z. wiss. Zool. lv.
- Petalomonas sexlobata, Basle, Klebs, p. 383, pl. xiv, figg. 19, a, b, Z. wiss. Zool. lv, n. sp.
- Phyllomitus amylophugus, Basle, Klebs, p. 321, pl. xiv, figg. 6 a-e, Z. wiss. Zool. lv, n. sp.
- Phyllomonas, n. g., with P. contorta, n. sp., Basle, Klebs, p. 305, pl. xiii, figg. 11 a-d, Z. wiss. Zool. lv.
- Pyrsonympha flugellata, parasitic in Termes lucifugus, GRASSI, p. 36, Atti Acc. Pontif. Lincei (5) i, n. sp.
- Rhynchomonas, n. g., with R. nasuta, Stokes, n. sp., Basle, p. 320, pl. xiv, figg. 7 a-d; KLEBS, Z. wiss. Zool. lv.
- Spironema, n.g., with S. multiciliatum, n. sp., Basle; Klebs, p. 350, pl. xvi, figg. 9 a-c, Z. wiss. Zool. lv.
- Stephanoon, n. g, with S. askenasii, Melbourne, Schewiakoff, Verh. Ver. Heidelb. (n.s.) iv.
- Streptomonas, n. g., p. 307, with S. cordata, Perty., n. sp., Basle, p. 307, pl. xiii, figg. 10 a, b; Klebs, Z. wiss. Zool. lv.
- Thylakomonas, n. g., with T. compressa, n. sp., Sydney, Schewiakoff, p. 550, Verh. Ver. Heidelb. (n.s.) iv.
- Trepomonas rotans, p. 349, pl. xvi, figg. 4 a-c, steinii, p. 349, figg. 5 a-d, Basle, Klebs, Z. wiss. Zool. lv, n. spp.
- Trigonomonas, n. g., with T. compressa, n. sp., Basle, Klebs, p. 330, pl. xv, figg. 5 α-g, Z. wiss. Zool. lv.
- Urophagus, n. g., with U. rostratus, Stein, n. sp., Basle, p. 341, Klebs, Z. wise. Zool. lv.
- Xunthodiscus, n. g., with X. vacillans, n. sp., Gippsland, Victoria, Schewiakoff, p. 551, Verh. Ver. Heidelb. (n.s.) iv.

### Class 3.—DINOFLAGELLATA.

- Glenodinium uliginosum, Brennet, in Baden, p. 64, pl. iii, fig. 16, neglectum, Baden, p. 65, pl. iii, fig. 17, cornifax, p. 66, pl. iii, fig. 18, SCHILLING (134); G. edax, Basle, pp. 206 & 207, pl. x, figg. 23 & 24, SCHILLING, Ber. Deutsch. Bot. Ges. ix: n. spp.
- Gymnodinium palustre, p. 58, pl. iii, fig. 11, carinatum, p. 59, fig. 12, paradoxum, p. 59, fig. 13, hyalinum, p. 60, fig. 14, pusillum, p. 60, fig. 15, SCHILLING (134), n. spp.

### Class 5.—CILIATA.

- Astylozoon pyriforme, New Zealand, Schewiakoff, p. 561, Verh. Ver. Heidelb. (n.s.) iv, n. sp.
- Balantiophorus elongatus, Gray's Peak (8700 feet) in Colorado, p. 558, bursaria, Bali I., p. 558, Schewiakoff, Verh. Ver. Heidelb. (n.s.), iv, n. spp.
- Blepharostoma, n. g., with B. glaucoma, n. sp., Sydney; SCHEWIAKOFF, p. 554, Verh. Ver. Heidelb. (n.s.) iv.

Codonella radix, Lagoons of Venice, IMHOF, p. 4, Neptunia, i, n. sp Cranotheridium, n. g., with C. taniatum, n. sp., New Zoaland, Sch KOFF, p. 554, Verh. Ver. Heidelb. (n.s.), iv.

Dichilum, n. g., with D. cuneiforme, n. sp., Sydney, Schewiakoff, Verh. Ver. Heidelb. (n.s.), iv.

Glaucoma setosa, p. 554, reniformis, p. 555, Sydney, colpidium, Nevland, p. 555, Schewiakoff, Verh. Ver. Heidelb. (n.s.), iv, n. 1
Holophrya simplex, Oahu and Hawaii, Schewiakoff, p. 553, Verl

Heidelb. (n.s.) iv, n. sp.

Ichthyophthirius cryptostomus, Plön, Holstein, Zacharias, (164); pl. i, n. sp.

Meseres, n. g., cordiformis, M. stentor, n. spp., New Zealand, Sch Koff, Verh. Ver. Heidelb. (n.s.) iv.

Monochilum, n. g., with M. frontatum, n. sp.; Oahu, Schewiakoff, ; Verh. Ver. Heidelb. (n.s.) iv.

Plagiocampa, n. g., with P. mutabile, n. sp., Sydney, Schewis p. 557, Verh. Ver. Heidelb. (n.s.) iv.

Rhabdostyla arenaria, from Synapta adhærens, Roscoff, Cuenor, Biol. iii, p. 290, pl. v, fig. 2, n. sp. 4

Stegochilum, n. g., with S. fusiforme, n. sp., Taupo Lake, N. Z., Sch Koff, p. 556, Verh. Ver. Heidelb. (n.s.) iv.

Strobilidium, n. g., with S. adhærens, n. sp., New Zealand, SCHEWIA p. 559, Verh. Ver. Heidelb. (n.s.) iv.

Tetrastyla, n. g., with T. oblonga, n. sp., New Zealand, SCHEWIA p. 561, Verh. Ver. Heidelb. (n.s.) iv.

Thylakidium, n. g., with T. truncatum, n. sp., Sydney, Schewia p. 559, Verh. Ver. Heidelb. (n.s.) iv.

Trichodina synaptæ, p. 289, pl. v, fig. 1, antedonis, p. 290, fig. 2, both Synapta adhærens, Roscoff, Cuénot, Rev. Biol. iii, n. spp.

Uronema orale, Sydney, SCHEWIAKOFF, p. 556, Verh. Ver. Heidelb. iv, n. sp.

Urotricha furcata, Oahu, p. 553, globosa, New Zealand, p. 553, Schikoff, Verh. Ver. Heidelb. (n.s.) iv, n. spp.

Vorticella amphiuræ, fig. 4, from Amphiura squamata, Naples, Cu p. 292, pl. v, Rev. Biol. iii, n. sp.

### INCERTÆ SEDIS.

Limnophrya elegans, Zacharias, Zool. Anz. xv, p. 459, n. sp. [n. g.] not described.

Neusina, n. g., with N. agassizi, n. sp., from the American Tr Pacific; Göes, Bull. Mus. C. Z. xxiii, p. 195. He supposes it to arenaceous Foraminifer: but as the description and illustration this form are almost identical with those given by HÆCKEL for Sin phyllum zonarium, one of his deep-sea Keratosa, we have to further research, before assigning to this species a definite plathe system.

### INDEX TO NAMES

OF

# NEW OR NOT PREVIOUSLY RECORDED GENERA AND SUBGENERA

### MENTIONED IN VOLUME XXIX,

INCLUDING NEW NAMES FOR GENERA PREVIOUSLY ESTABLISHED.\*

Abantis, Fairmaire, Ins. 133 Ablabophis, Boulenger, Rept. 28 Abrachius, Roelofs, Ins. 146 Abrisa, Kirby, Ins. 226 Abuzaida, Keyserling, Arachn. 21 Acanthogobio, Herzenstein, Pisces Acanthomyrmex, Emery, Ins. 196 Acanthoprion, Pictet and Saussure, Ins. 327 Acanthosphinx, Aurivillius, Ins. Achætoneura, Brauer and Bergenstamm, Ins. 299 Achelura, Kirby, Ins. 229 Achlæna, Karsch, Ins. 326 Achrysis, Sėmenow, Ins. 201 Aciagrion, Selys, Ins. 322 Acinopterus, Duzee, Ins. 315 Acrocœlum, Cossmann, Moll. 76 Acrophylictis, Cossmann, Moll. 78 Acrostigma, Emery, Ins. 83 Acrotrema, Cossmann, Moll. 69 Actenioides, Warren, Ins. 274 Actinodactylus, Haswell, Vermes Actinoxia, Simon, Arachn. 26 Adeloides, Warren, Ins. 274 Adelphomyia, Bergroth, Ins. 285 Adixoa, Hampson, Ins. 228 Admetula, Cossmann, Moll. 72 Æbutina, Simon, Arachn. 32 Ælianus, Jacoby, Ins. 171 Æmilia, Kirby, Ins. 229 Aeromyrma, Forel, Ins. 196 Aeronautes, Hartert, Aves 44 Aeschrodomus, Pilsbry, Moll. 63 Agada, Fairmaire, Ins. 160 Agastillus, Reitter, Ins. 91 Agathostola, Simon, Arachn. 26 Aglaops, Warren, Ins. 274
Ahlia, Jordan and Davis, Pisces Aigialosaurus, Gorjanovic - Kram-berger, Rept. 25 Ailuravus, Rütimeyer, Mamm. 31 Aleurodicus, Douglas and Morgan. Ins. 317 Allomorpha, Jacoby, Ins. 171 Alophogaster, Hampson, Ins. 230 Alsomyia, Brauer and Bergenstamm. Ins. 298

The total number of names in this Index is 1520, distributed as follows:—
Mammalia, 31; Aves, 37; Reptilia and Batrachia, 44; Pisces, 34; Tunicata, 3
Mollusca, 145; Brachiopoda, 27; Bryozoa, 2; Crustacsu, 37; Arachnida, 120;
Myriopoda, 2; Insecta, 916; Echinodermata, 7; Vermes, 42; Cælenterata, 3;
Spongiæ, 19; Protozoa, 51. The corresponding total last year was 1399, and ten
years ago, in 1882, 1015.

Alsopsyche, Brauer and Bergenstamm, Ins. 298 Amalocichla, De Vis, Aves 59 Amaurocichla, Sharpe, Aves 59 Ambigatus, Fairmaire, Ins. 134 Amelora, Meyrick, Ins. 262 Ametrosomus, Tepper, Ins. 327 Ammospermophilus, Merriam, Mamm. 34 Amphichæta, Kallstenius, Vermes Amphilochoides, Sars, Crust. 25 Amphithemis, Selys, Ins. 322 Amplogladius, Cossmann, Moll. 75 Anagonia, Brauer and Bergenstamm, Ins. 299 Anamastax, Brauer and Bergenstumm, Ins. 299 Anapas, Bergendal, Vermes 85 Anaptycha, Brunner, Ins. 325 Anartomorpha, Alphéraky, Ins. 251 Anaspilus, Kolbe, Ins. 119 Anatinomma, Bates, Ins. 160 Anaxidia, Swinhoe, Ins. 230 Andrimus, Casey, Ins. 140 Anelaphinis, Kolbe, Ins. 119 Anelasma, Cossmann, Moll. 67 Aneogmena, Brauer and Bergenstamm, Ins. 301 Anephilus, Faust, Ins. 146 Aneuretus, Emery, Ins. 196 Aneurychilus, Cossmann, Moll. 75 Angularia, Woehrmann and Koken, Moll. 76 Anhibernia, Staudinger, Ins. 262 Annadrilus, Horst, Vermes 48 Anodonthyla, Müller, Rept. 38 Anoedopoda, Karsch, Ins. 327 Anomolocaris, Whiteares, Crust. 29 Anomostictis, Warren, Ins. 274 Antha, Staudinger, Ins. 251 Anthiphula, Jacoby, Ins. 171 Anthocoptes, Nalepa, Arachn. 37 Anthocrypta, Warren, Ins. 274 Anthrenoptera, Swinhoe, Ins. 228 Anthyria, Swinhoe, Ins. 263 Anticypella, Meyrick, Ins. 263 Aon, Neumoegen, Ins. 230 Apatospongia, Marshall, Spong. 20 Aphanesthes, Kolbe, Ins. 119 Aphanolimnas, Sharpe, Aves 30 Aphrogeneia, Gumppenberg, Ins. 263 Apiomidas, Coquillett, Ins. 288 Apiotoma, Cossmann, Moll. 71 Aplographe, Warren, Ins. 274 Aporoctena, Meyrick, Ins. 263 Aporozoum, Germain, Ins. 107 Apostichus, Simon, Arach. 26

Apotomogonius, Mon 303 Apotrepus, Casey, Ius. Apteronomus, Tepper, Apteroscirtus, Karsch, Aræocera, Hampson, I Aræogyia, Hampson, I Archæopneustes, Greg Archæotrogon, Miln Aves 46

Archondess, Moore, It Arenoxenus, Bates, In Argopistoides, Jacoby, Argyrocheila, Staudin, Argyrocottus, Herzen

Argyrophylax, Brauer stamm, Ins. 299
Ariagona, Krauss, Ins. Arietellus, Giesbrecht, Aristolebia, Bates, Ins. Arntros, Schaus, Ins. Arrenopus, Brauer

Armenopus, Brauer

stamm, Ins. 300 Arrhinidia, Brauer a stamm, Ins. 301 Arsinotus, Berg, Ins. 3 Artemidora, Meyrick, Arthroglena, Bergen 86

Artiora, Meyrick, Ins. Artobius, Winge, Mar Aschistophleps, Hamp Asmonax, Gorham, In Aspatha, Sharpe, Ave. Astræosmilia, Ortmani Astralites, Whiteares. Astrapephora, Alphére Atasca, Swinhoe, Ins. : Atelolathys, Simon, A Ateuchetus, Bedel, In Atomophora, Alphéra: Atrochus, Wierzejski, Atropharista, Townsen Atrotophopalpus, To 294

Aucheromyia, Brauer stamm, Ins. 301
Autarcus, Senna, Ins. Autocosmia, Warren, Autolethrus, Sémenow Auximus, Simon, Arac Aventina, Staudinger, Aventiola, Staudinger, Avocettina, Jordan Pisces 31

Avus, Ortmann, Crust

Axestoleus, Bates, Ius. 160 Aysha, Keyserling, Arachn. 21 Azarelius, Fairmaire, Ius. 134 Azygophleps, Hampson, Ins. 232

Bactromyia, Brauer and Bergenstamm, Ins. 298 Bactrotheca, Novak, Moll. 70 Baglivia, Brusina, Moll. 78 Barroisella, Hall and Clarke, Brach. 5 Basenius, Kolbe, Ins. 169 Bathippus, Forel, Arachn. 15 Bathycranium, Strobl, Ins. 290 Bathymedon, Surs, Crust. 26 Batodon, Marsh, Mamm. 53 Bellota, Peckham, Arachn. 15 Benthobia, Dall, Moll. 72 Benthonella, Dall, Moll. 78 Bessophora, Meyrick, Ins. 263 Bifidoscala, Cossmann, Moll. 83 Billingsella, Hall and Clarke, Brach. 6 Bindax, Thorell, Arachn. 15 Bipalpus, Wierzejski, Vermes 86 Blanchardia, Solger, Prot. 28 Blaptocampus, Thomson, Ins. 202 Blenniophidium, Boulenger, Pisces Blepharidopsis, Brauer and Bergenstamm, Ins. 298 Blepharostoma, Schewiakoff, Prot. Blepharucha, Warren, Ins. 274 Bocus, Peckham, Arachn. 15 Bolbodimyia, Bigot, Ins. 286 Bolivinopsis, Jakowlew, Prot. 19 Bolomyia, Brauer and Bergenstamm, Ins. 299 Bonatea, Druce, Ins. 264 Boophilus, Curtice, Arachn. 37 Borophagus, Cope, Mamm. 29 Bothriocyrtum, Simon, Arachn. 26 Bothriotænia, Railliet, Vermes 64 Brachyceræa, Röder, Ins. 293 Brachychalcinus, Boulenger, Pisces 28 Brachycolia, Meade, Ins. 291 Brachymylus, Woodward, Pisces 35 Brachypelma, Simon, Arachn. 27 Bramus, Pomel, Mamm. 35 Branchiura, Beddard, Vermes 48 Bremeria, Alphéraky, Ins. 232 Brochopeplus, Pictet and Saussure, Ins. 328 Bryodrilus, Ude, Vermes 48 Bryophilina, Staudinger, Ins. 252 Bubastodes, Blackburn, Ins. 123

Bucolinus, Blackburn, Ins. 183 Bulimactæon, Cossmann, Moll. 69 Buttikoferia, Roelofs, Ins. 148 Byrrhomorpha, Blackburn, Ins. 114

Caddo, Banks, Arachn. 34 Cæcaria, Floericke, Vermes 48 Calapnita, Simon, Arachn. 25 Caletes, Redtenbacher, Ins. 331 Callarge, Leech, Ins. 219 Callartona, Hampson, Ins. 233 Calleva, Simon, Arachn. 19 Callipareius, Senna, Ins. 169 Calliphona, Krauss, Ins. 328 Callosila, Saussure, Ins. 189 Callotaria, Palmer, Mamm. 30 Calocerinus, Howard, Ins. 206 Calybia, Kirby, Ins. 233 Camerania, Giglio-Tos, Ins. 290 Camptochilus, Hampson, Ins. 233 Camptomastix, Warren, Ins. 274 Camptostichomma, Karsch, Arachn 14 Canimartes, Cope, Mamm. 30 Canistes, Casey, Ins. 148 Capnodes, Rostock, Ins. 321 Carcinodon, Scott, Mamm. 32 Carcinopisthius, Kolbe, Ins. 169 Carphonotus, Casey, Ins. 148 Carrhotus, Thorell, Arachn. 15 Cassena, Weise, Ins. 172 Catacheta, Brauer and Bergen-stamm, Ins. 298 Catagonia, Brauer and Bergenstamm, Ins. 299 Catapaguroides, Edwards and Bouvier, Crust. 21 Catapastus, Casey, Ins. 148 Cataponus, Desbrochers, Ins. 148 Catargynnis, Röber, Ins. 219 Cenchromorphus, Fairmaire, Ins. Centrinites, Casey, Ins. 149 Centrinogyna, Casey, Ins. 149 Centrinopus, Casey, Ins. 149 Centrodontus, Goding, Ins. 315 Cephis, Ragonot, Ius. 274 Cerania, Weise, Ins. 172 Ceratobothrium, Monticelli, Vermes Ceratoconcha, Gorjanovic - Kramberger, Moll. 91 Ceratocorema, Hampson, Ins. 228 Ceratonema, Hampson, Ins. 233 Ceratotheca, Novák, Moll. 70 Cestophorus, Faust, Ins. 149 Chestodemoticus, Brauer and Bergenstamm, Ins. 301

Cheetogedia, Brauer and Bergenstamm, Ins. 299 Chætoglossa, Townsend, Ins. 294 Chætomyia, Brauer and Bergenstamm, Ins. 298 Chætophthalmus, Brauer and Bergenstamm, Ins. 300 Chætoprocta, Brauer and Bergenstamm, Ins. 299 Chalcinopelecus, Holmberg, Pisces Chalcocelis, Hampson, Ins. 233 Chaleponotus, Casey, Ins. 149 Chalioides, Hampson, Ins. 233 Chamops, Marsh, Rept. 23 Chaperina, Mocquard, Rept. 38 Chara, Staudinger, Ins. 253 Characta, Redtenbacher, Ins. 328 Charisia, Champion, Ins. 161 Cheloctonus, Pocock, Arachn. 14 Chenuala, Swinhoe, Ins. 233 Chevallieria, Cossmann, Moll. 78 Chilobrachys, Karsch, Arachn. 27 Chipota, Hulst, Ins. 274 Chiridius, Giesbrecht, Crust. 30 Chirostylus, Ortmann, Crust. 20 Chlænioctenus, Bates, Ins. 92 Chlamys, Beneden, Crust. 30 Chlidonoptera, Karsch, Ins. 326 Chloracris, Pictet and Saussure, Ins. 328 Chlorampulex, Saussure, Ins. 190 Chlorochropsis, Rothschild, Ins. 214 Chlorotettix, Duzee, Ins. 315 Chlorotribonia, Pictet and Saussure, Ins. 328 Chonopectus, Hall and Clarke, Brach. 6 Chonostrophia, Hall and Clarke, Brach. 6 Choranesa, Moore, Ins. 219 Choristoneura, Rübsaamen, Ins. 284 Choristostigma, Warren, Ins. 275 Hall and Clarke, Christiania, Brach. 6 Chrotoma, Casey, Ins. 161 Chrysamœba, Klebs, Prot. 30 Chrysartona, Swinhoe, Ins. 234 Chrysilla, Thorell, Arachn. 15 Chrysococcus, Klebs, Prot. 30 Chrysoctenis, Meyrick, Ins. 264 Chrysodasia, Reuter, Ins. 311 Cirsochilus, Cossmann, Moll. 85 Cirsope, Cossmann, Moll. 77 Cladocroce, Topsent, Spong. 20 Clanodon, Scott, Mamm. 31 Claudia, Hartert, Aves 44 Clavipalpula, Staudinger, Ins. 253

Cleionychia, Ulrich, Moll. 94 Cleonidius, Casey, Ins. 149 Clistomorpha, Townsend, Ins. 2 Cnephaliodes, Brauer and Ber stamm, Ins. 299 Cnephidia, Ragonot, Ins. 275 Cnidia, Gumppenberg, Ins. 265 Codiosoma, Kirby, Ins. 234 Coleolissus, Bates, Ins. 92 Colomyia, Kieffer, Ins. 284 Columbellisipho, Cossmann, Mol Columbia, Eigenmann, Pisces 3 Coniophis, Marsh, Rept. 30 Conogaster, Brauer and Berg stamm, Ins. 298 Conosara, Meyrick, Ins. 265 Conradtia, Kolbe, Ins. 119 Copa, Weise, Ins. 173 Cophixalus, Beettger, Rept. 38 Coppingeria, Haswell, Vermes 4 Coptochetus, Cossmann, Moll. 7 Coracocephalus, Mik, Ins. 290 Cormopus, Kolbe, Ins. 169 Cornucaprina, Futterer, Moll. 91 Coronimyia, Townsend, Ins. 294 Corynotrichius, Kolbe, Ins. 119 Cosmocreon, Warren, Ins. 275 Cosmodesmus, Haase, Ins. 211 Cossulinus, Kirby, Ins. 234 Cotylogaster, Monticelli, Vermes Cotyloplana, Spencer, Vermes 7 Cranotheridium, Schewiakoff, P 32Cratorrhagus, Simon, Arachn. 2 Crecoides, Shufeldt, Aves 30 Cremnophila, Ragonot, Ins. 275 Crenigomphus, Selys, Ins. 323 Crenilabium, Cossmann, Moll. ( Crinocraspeda, Hampson, Ins. 2. Criomacha, Meyrick, Ins. 205 Crocomela, Kirby, Ins. 235 Crommium, Cossmann, Moll. 8: Crossiura, Nicéville, Ins. 225 Cryptomeigenia, Brauer and . genstamm, Ins. 298 Cryptopone, Emery, Ins. 198 Ctenagenia, Saussure, Ins. 190 Ctenolobus, Desbrochers, Ins. 1 Ctenophorocera, Brauer and genstamm, Ins. 299 Curvatula, Staudinger, Ins. 253 Cyanoplax, Pilsbry, Moll. 88 Cyclocranium, Poll, Ins. 162 Cyclodostomia, Sacco, Moll. 83 Cyclomias, Desbrochers, Ins. 15 Cyclopteroides, Garman, Pisces Cyclosattus, Casey, Ins. 135 Cycloscymnus, Blackburn, Ins.

Cyclotynaspis, Montandon, Ins. 308 Cylindrococcus, Maskell, Ins. 318 Cylindromus, Aurivillius, Ins. 150 Cylloproctus, Faust, Ins. 150 Cyniscella, Cossmann, Moll. 85 Cyphotarsis, Jacoby, Ins. 174 Cystiphora, Kieffer, Ins. 284

Dactylophysus, Fleutiaux, Ins. 127 Dactylopteryx, Karsch, Ins. 326 Dactylorhyncha, Hampson, Ins. 235 Dæochæta, Townsend, Ins. 294 Dalium, Dall, Moll. 82 Dalmanella, Hall and Clarke, Brach. 6 Dalterus, Fairmaire, Ins. 162 Daramus, Fairmaire, Ins. 162 Dasyphleps, Karsch, Ins. 328 Dasypteroma, Standinger, Ins. 265 Delagrangeus, Pic, Ins. 162 Deltopecten, Jack and Etheridge, Moll. 94 Dendrophleps, Hampson, Ins. 235 Dermatæstrus, Brauer, Ins. 293 Descanso, Peckham, Arachn. 15 Desmoglyptus, Casey, Ins. 150 Desvoidia, Meade, Ins. 294 Dexiophana, Brauer and Bergen-stamm, Ins. 300 Diacme, Warren, Ins. 275 Diaglyptus, Pilsbry, Moll. 63 Dialytostoma, Cossmann, Moll. Diamenocrinus, Oehlert, Ech. 19 Diaphlebus, Karsch, Ins. 328 Diaphorapteryx, Forbes, Aves 30 Diaphorus, Faust, Ins. 150 Diastrophilus, Faust, Ins. 150 Dibolocelus, Bedel, Ins. 100 Dichelomyia, Rübsaamen, Ins. 284 Dichilum, Schewiakoff, Prot. 32 Dichotis, Warren, Ins. 275 Dicloniscus, Cope, Rept. 24 Dicranopsyra, Dohrn, Ins. 328 Dicranotænia, Railliet, Vermes 64 Didymostoma, Warren, Ins. 275 Dillwynella, Dall, Moll. 85 Dimeropus, Cope, Rept. 24 Dimorphomyrmex, André, Ins. 198 Dinarella, Bittner, Brach. 8 Dinocleus, Casey, Ins. 151 Dinorthis, Hall and Clarke, Brach. Dionana, Moore, Ins. 219

Diops, Bergendal, Vermes 86

Diploa, Kolbe, Ins. 120

Dioptrophorus, Faust, Ins. 151 Diphaulacosoma, Jacoby, Ins. 174

Diptychus, Cossmann, Moll. 83 Discalma, Meyrick, Ins. 265 Dischiropus, Karsch, Arachn. 15 Discinopsis, Hall and Clarke, Brach. 5 Distantella, Saussure, Ins. 202 Ditrimella, Hinde and Holmes, Spong. 17 Dixoa, Hampson, Ins. 236 Dolichostethus, Kolbe, Ins. 120 Doliodus, Traquair, Pisces 37 Dolothymus, Thorell, Arachn. 25 Dorachosa, Distant, Ins. 313 Drepanidotænia, Railliet, Vermes Drepanula, Röber, Ins. 221 Drexelia, McCook, Arachn. 22 Drieschia, Michaelsen, Vermes 45 Dromicodryas, Boulenger, Rept. 27 Drucea, Kirby, Ins. 236 Drymoptila, Meyrick, Ins. 265 Dynamopterus, Milne - Edwards, Aves 46 Dyscerus, Faust, Ins. 151 Dystrichothorax, Blackburn, Ins.

Eccopteroma, Standinger, Ins. 254 Eccoptometopus, Péringuey, Ins. Echinocotyle, Rosseter, Vermes 64 Echnomolaus, Bates, Ins. 93 Echtrusia, Lefèrre, Ins. 175 Ectatostica, Šimon, Arachn. 23 Ectinochilus, Cossmann, Moll. 75 Ectophylla, Allen, Mamm. 27 Ectorisma, Tate, Moll. 90 Ectrocta, Hampson, Ins. 236 Eilica, Keyserling, Arachn. 20 Elmoptera, Redtenbacher, Ins. 328 Elaphrocnemus, Milne - Edwards, ∆ves 29 Elasmocnema, Karsch, Ins. 306 Elginia, Newton, Rept. 35 Ellipsodon, Scott, Mamm. 31 Elsa, Honrath, 1ns. 214 Embolocephalus, Randolph, Vermes Emertonius, Peckham, Arachn, 16 Emphanopteryx, Townsend, Ins. 295Emporomyia, Brauer and Bergenstamm, Ins. 300 Encaumaptera, Hampson, Ins. 236 Encentra, Redtenbacher, Ins. 329 Encyocrates, Simon, Arachn. 27 Endopachychilus, Cossmann, Moll.

į

Enicoderus, Péringuey, Ins. 151 Entomella, Cossmann, Moll. 87 Entomope, Cossmann, Moll. 77 Eonius, Tepper, Ins. 329 Eopleurotoma, Cossmanu, Moll. 71 Epalxis, Cossmann, Moll. 71 Epelyx, Blackburn, Ins. 93 Ephebopus, Simon, Arachn. 27 Ephemeroidea, Hampson, Ius. 236 Ephippithytoidea, Tepper, Ins. 329 Ephoria, Meyrick, Ins. 266 Epichriacus, Scott, Mamm. 31 Epicoinoneus, Senna, Ins. 169 Epicompsa, Meyrick, Ins. 266 Epimydia, Staudinger, Ius. 236 Epineura, Brauer and Bergenstamm, Ins. 301 Epision, Stiles, Vermes 64 Epocilla, Thorell, Arachn. 16 Epoecus, Emery, Ins. 198 Eremocossus, Hampson, Ins. 237 Eremoxenus, Sémenow, Ins. 169 Erica, Peckham, Arachn. 16 Erimacrus, Benedict, Crust. 17 Eriphioides, Kirby, Ins. 237 Eristus, Blackburn, Ins. 151 Erithionyx, Blackburn, Ins. 184 Erythrandra, Brauer and Bergenstamm, Ins. 300 Erythrolophus, Swinhoe, Ins. 266 Erythronychia, Brauer and Bergenstamm, Ins. 300 Erythrophaia, Staudinger, Ins. 254 Estrella, Frenzel, Prot. 19 Eucasta, Dall, Moll. 85 Euceromyia, Townsend, Ins. 295 Eucirrichthys, Perugia, Pisces 28 Eucuephalia, Townsend, Ins. 295 Euctenospila, Warren, Ins. 275 Eudiplura, Simon, Arachn. 27 Eudoxazus, Kolbe, Ins. 120 Eudranes, Sharp, Ins. 103 Eufischeria, Brauer and Bergenstamm, Ins. 300 Euglenopsis, Klebs, Prot. 30 Eulasinus, Sharp, Ins. 103 Eulasiona, Townsend, Ins. 295 Eumacronychia, Townsend, Ins. 295 Eumegalodon, Brongniart, Ins. 329 Eumelepta, Jacoby, Ins. 175 Eumera, Staudinger, Ins. 267 Eumyothyria, Townsend, Ins. 295 Eunotosaurus, Seeley, Rept. 34 Eunyssobia, Casey, Ins. 151 Eupachnoda, Kolbe, Ins. 120 Eupegylis, Duvicier, Ins. 114 Euphorocera, Townsend, Ius. 295 Euponera, Forel, Ins. 198

Euposaurus, Lortet, Rept. Eupsychellus, Röber, Ins. Eurhinospiza, Oustalet, A Euromoia, Staudinger, Ins. Euryaptus, Bates, Ins. 94 Euryetisus, Gavino, Crust Euscopolia, Townsend, Ins. Eusphinetus, Emery, Ins. Eutelesthes, Kolbe, Ins. 15 Eutettix, Duzee, Ins. 316 Euthyprosopa, Townsend, Eversmanella, Jakowelev, I Exechestoma, Cossmann, I Exophthalmida, Faust, In

Fanchonia, Werner, Rept. Felderia, Kirby, Ins. 237 Feltria, Kanike, Arachn. Fergana, Staudinger, Ins. Filholornis, Milne-Edwar 29 Fluda, Peckham, Arachn. Fohlia, Lohnanu, Tun 7

Fluda, Peckham, Arachu. Fohlia, Lohmann, Tun. 7 Folioseris, Rehberg, Cœl. 1 Forelius, Emery, Ins. 198 Frenchella, Blackburn, Ins. Frenchia, Maskell, Ins. 311 Furcillus, Cope, Rept. 24

Gædiopsis, Brauer and stamm, Ins. 299 Galerusoma, Jacoby, Ins. 1 Gardinia, Kirby, Ins. 238 Gargantua, Kirby, Ins. 22' Gasponia, Fairmaire, Ins. Gastroschiza, Bergendal, 87

Gavialinum, Lortet, Rept. Geikia, Newton, Rept. 35 Gelotia, Thorell, Arachn. Gemeneta, Kursch, Ins. 33 Geosesarma, De Man, Cru Geosomus, Faust, Ins. 152 Geothauma, Crosse, Moll. Geranopterus, Milne-Edwa 41

Ginglymia, Townsend, Ins. Gitanopsis, Sars, Crust. 2t Glauconoë, Warren, Ins. 2 Glaucophana, Brauer and stamm, Ins. 299 Glyphidrilus, Horst. Verm.

Glyphidrilus, Horst, Verm Glyptobaris, Casey, Ins. 17. Glyptostyla, Dall, Moll. 7. Gnopharmia, Staudinger, I Goë, Hampson, Ins. 238 Goëtia, Karsch, Ins. 329 Goldia, Keyserling, Arachn Gonanticlea, Swinhoe, Ins. 267 Goniocera, Brauer and Bergenstamm, Ins. 299 Gonionotophis, Boulenger, Rept. 28 Goniops, Aldrich, Ius. 286 Gordiichthys, Jordan and Davis, Pisces 31 Gordiodrilus, Beddard, Vermes 49 Gordonia, Newton, Rept. 35 Gorgonichthys, Claypole, Pisces 31 Grammostola, Simon, Arachn. 27 Grandipatula, Cossmann, Moll. 62 Grantiopsis, Dendy, Spong. 16 Granulolabium, Cossmann, Moll. Gravenreuthia, Karsch, Ins. 329 Grouvelleus, Guillebeau, Ins. 106 Grumia, Alphéraky, Ins. 255 Grypnus, Desbrochers, Ins. 152 Gurna, Swinhor, Ins. 238 Guttulidium, Frenzel, Prot. 18 Gymnodexia, Brauer and Bergen-stamm, Ins. 300 Gymnodiptychus, Herzenstein, Pisces 27 Gymnolæmus, Ogilvie-Grant, Aves Gymnoprosopa, Townsend, Ins. 295 Gymnoscymnus, Blackburn, Ins. Gynæphila, Staudinger, Ins. 260 Gytha, Keyserling, Arachn. 20 Hadjina, Staudinger, Ins. 255 Hadropopillia, Kraatz, Ins. 118 Hadrosticta, Kraatz, Ins. 120 Hallicarnia, Kirby, Ins. 238 Hallina, Winchell and Schuchert, Brach. 8 Haloplia, Snellen, Ins. 238 Handhirschiella, Montandon, Ins. Haploclastus, Simon, Arachn. 27 Haplopelma, Simon, Arachn. 28 Haploscincus, Cope, Rept. 24 Haporema, Fairmaire, Ins. 136 Harima, Moore, Ins. 219 Harmilla, Aurivillius, Ins. 217 Harpaliscus, Bates, Ins. 94

Hedleya, Cox, Moll. 81 Helectrus, Guillebeau, Ins. 106 Helenoconcha, Pilsbry, Moll. 63

Heligmomerus, Simon, Arachn. 28

Heliosphærium, Frenzel, Prot. 19

Heliocausus, Kohl, Ins. 191

Hemiaulax, Bates, Ins. 94

Hemichlænius, Bates, Ins. 94

Hemiconus, Cossmann, Moll. 71

Hemimeria, Saussure, Ins. 191 Hemipimpla, Saussure, Ins. 202 Hemipleurotoma, Cossmann, Moll. Hemipodiulus, Verhoeff, Myr. 6 Hemipogonius, Saussure, Ins. 191 Hemisalius, Saussure, Ins. 191 Hemithrixion, Brauer and Bergenstamm, Ins. 300 Hemyloticus, Jacoby, Ins. 176 Henschia, Lethierry, Ins. 316 Hepatica, Standinger, Ins. 255 Hephthocara, Alcock, Pisces 23 Hepialiscus, Hampson, Ins. 239 Heræma, Staudinger, Ins. 255 Herbertella, Hall and Clarke, Brach. 6 Hermatobates, Carpenter, Ins. 309 Hermosa, Peckham, Arachu. 16 Hesperarion, Simroth, Moll. 59 Hesperobaris, Casey, Ins. 152 Heteroæcium, Hincks, Bry. 4 Heterocharmus, Pocock, Arachn. Heterorthis, Hall and Clarke. Brach. 6 Heylaertsia, Hampson, Ins. 239 Hildala, Swinhoe, Ins. 239 Histima, Fairmaire, Ins. 137 Hollandia, Butler, Ins. 261 Holocryptis, Lucas, Ins. 256 Hologyra, Woehrmann and Koken, Moll. 84 Holomastigotes, Grassi, Prot. 30 Hommophleus, Gahan, Ins. 163 Homalochilus, Cossmann, Moll. 86 Homocoplacis, Simon, Arachn. 28 Homoneura, Wulp, Ins. 301 Homostola, Simon, Arachu. 28 Hoplacerus, Jacoby, Ius. 176 Hopliterrhynchus, Senna, Ins. 169 Hoplopisthius, Senna, Ins. 169 Hoplozonitis, Blackburn, Ins. 144 Hordeulima, Sacco, Moll. 83 Hormambogaster, Karsch, Hubrechtia, Bürger, Vermes 70 Hyaloplagia, Warren, Ins. 276 Hyalorista, Warren, Ins. 276 Hydrogamasus, Berlese, Arachn. Hylaspoides, Durivier, Ins. 176 Hymenomaga, Karsch, Ius. 304 Hypatia, Kirby, Ins. 239 Hyperephanus, Senna, Ins. 169 Hyperlophus, Douglas - Ogilby, Pisces 30 Hyphantospiza, Reichenow, Aves 53

Hyphantrophaga, Townsend, Ins. Hypochlorosis, Röber, Ins. 222 Hypocyrema, Blackburn, Ins. 184 Hypolamprus, Hampson, Ins. 239 Hypopomyrmex, Emery, Ins. 83 Hypopus, Bergendal, Vermes 87 Hypostigmodera, Blackburn, Ins. 124 Hypotachina, Brauer and Bergenstamm, Ins. 299 Hypotomorpha, Rebel, Ins. 276 Hypsicorypha, Krauss, Ins. 326 Hypsipleura, Woehrmann and Koken, Moll. 76 Hypsobia, Heude, Moll. 79 Hyptiocrinus, Wachsmuth and Springer, Ech. 19 Hyracops, Marsh, Mamm. 39 Hysterocrates, Simon, Arachn. 28

Ichneumenoptera, Hampson, Ins. Ichniotherium, Pohlig, Rept. 41 Ideobisium, Balzan, Arachn. 34 Ideoblothrus, Balzan, Arachn. 34 Idiobates, Casey, Ins. 137 Idiocrinus, Wachsmuth and Springer, Ech. 20. Idiopholis, Boulenger, Rept. 29 Idiostethus, Casey, Ins. 152 Idonauton, Swinhoe, Ins. 240 Ilyophis, Jordan and Davis, Pisces Iniopsis, Lydekker, Mamm. 50 Iola, Peckham, Arachn. 16 Iphidosoma, Berlese, Arachn. 35 Iphthimorhinus, Roelofs, Ins. 152 Ischnias, Berg, Ins. 311 Ischnocryptus, Kriechbaumer, Ins. Ischnopopillia, Kraatz, Ins. 118 Isoclerus, Lewis, Ins. 131 Isomma, Selys, Ins. 323 Isoplatys, Montandon, Ins. 304 Ispaterus, Fairmaire, Ins. 163 Ityocephala, Redtenbacher. Ins. 329

Jænia, Grassi, Prot. 30 Jermakia, Jakowlew, Ins. 209 Jheringia, Keyserling, Arachu. 25 Jothochilus, Kolhe, Ins. 120 Joyeuxia, Topsent, Spong. 21

Kanakia, Distant, Ins. 313 diana, Distant, Ins. 314

Katosira, Woehrmann am Moll. 76 Kayserella, Hall and Brach. 6 Kerrata, Moore, Ins. 219 Kerria, Beddard, Vermes Keyserlingella, Peckham, 16 Kirrodesa, Moore, Ins. 220

Kirrodesa, Moore, Ins. 220 Kittlitzia, Hartlaub, Aves Konbirella, Duvivier, Ins. Kyrtolitha, Staudinger, Ins

Lachnomma, Townsend, In Lævidentalium, Cossman: 89

Lagarosia, Wulp, Ins. 301 Lampropelma, Simon, Ara Lamprophthalma, Portschiv 302

Larradomorpha, Stadelma 206

Larropsis, Patton, Ins. 191 Lasiogaster, Gahan, Ins. 16 Lasiogma, Meyrick, Ins. 26 Lasionycta, Aurivillius, Ins. Latirulus, Cossmann, Moll. Lembeja, Distant, Ins. 313 Lepidodexia, Brauer and stamm, Ins. 300

Lepidopterus, Pohlig, Pisca Lepidozona, Pilsbry, Moll. Leproscirtus, Karsch, Ins. Leptella, Hall and Clarke, Leptodema, Carlini, Ins. 31 Leptorrhamphus, Ambroset 32

Leptostylus, Faust, Ins. 1; Leptotachina, Brauer and stamm, Ins. 299 Leptothrix, Hadaerts, Inc.

Leptothrix, Heylaerts, Ins. Lethrotrypes, Jacobsohn, In Lethrulus, Semenow, Ins. 1 Leucascus, Dendy, Spong. Leucocythere, Kaufman, C Leucodrepana, Hampson, I Leucomyia, Brauer and stamm, Ins. 300

stamm, Ins. 300
Libania, Thorell, Arachn. 2
Libytherium, Pomel, Mam
Limeria, Staudinger, Ins. 2
Limnophrya, Zucharias, Pr
Limulina, Frenzel, Prot. 18
Lindstræmella, Hall and
Brach. 5

Linocerocoris, Karsch, Ins. Linonotus, Casey, Ins. 153 Liotrichius, Kolbe, Ins. 120

Liparops, Garman, Pisces 22 Liparopsis, Hampson, Ins. 241 Litaneutria, Saussure, Ins. 326 Lithoglyptes, Aurivillius, Crust. 32 Lithophasia, Staudinger, Ins. 256 Lithosphærella, Frenzel, Prot. 19 Litochroides, Guillebeau, Ins. 106 Lobantale, Cossmann, Moll. 89 Lobocara, Bergroth, Ins. 309 Lobostoma, Fairmaire, Ins. 107 Lochetica, Kriechbaumer, Ins. 203 Longula, Staudinger, Ins. 268 Lophosoma, Swinhoe, Ins. 241 Loxocreon, Warren, Ins. 276 Loxoptyxis, Cossmann, Moll. 83 Lucernosa, Antipa, Col. 11 Lycognathophis, Boulenger, Rept. 28 Lycosella, Thorell, Arachn. 23 Lyctosoma, Lewis, Ins. 131

Maccoyella, Jack and Etheridge, Moll. 94 Macrocrambus, Kirby, Ins. 241 Macrodostomia, Sacco, Moll. 83 Macrohelodes, Blackburn, Ins. 129 Macrolabis, Kieffer, Ins. 284 Macromeigenia, Brauer and Bergenstamm, Ins. 298 Macromphalina, Cossmann, Moll. Macronola, Kirby, Ins. 241 Macropalpus, Meunier, Ins. 296 Macropisthodon, Boulenger, Rept. Macroplectra, Hampson, Ins. 241 Macropophis, Boulenger, Rept. 27 Macrotarsipus, Hampson, Ins. 228 Macrotrypeta, Portschinsky, Ins. 302 Madarellus, Casey, Ins. 153 Magulla, Simon, Arachn. 28 Maizania, Bourguignat, Moll. 81 Maltharchus, Weise, Ins. 130 Mandarella, Duvivier, Ins. 177 Mandarinia, Leech, Ins. 220 Marænaspis, Karsch, Ins. 310 Marconia, Bourquignat, Moll. 57 Marengo, Peckham, Arachn. 17 Margineulima, Cossmann, Moll. 83 Marsupiogaster, Schewiakoff, Prot. Martella, Peckham, Arachn. 17 Masiphya, Brauer and Bergenstamm, Ins. 298 Masipoda, Brauer and Bergenstamm,

Ins. 299

Masistylum, Brauer, Ins. 296 Mastigella, Frenzel, Prot. 18 Mastigina, Frenzel, Prot. 18 Mastigocrinus, Bather, Ech. 20 Mastigosphæra, Schewiakoff, Prot. Masupha, Distant, Ins. 313 Maulia, Blackburn, Ins. 164 Maupasia, Schewiakoff, Prot. 30 Maxia, Kirby, Ins. 241 Mecinonota, Kraatz, Ins. 121 Medoriopsis, Cossmann, Moll. 77 Megaceramis, Hampson, Ins. 241 Megalochlora, Meyrick, Ins. 269 Megametopon, Alphéraky, Ins.  $2\overline{69}$ Megatyloma, Cossmann, Moll. 87 Mekla, Swinhoe, Ins. 242 Melia, Neumoegen, Ins. 242 Melichrus, Brenske, Ins. 116 Mellevillia, Cossmann, Moll. 75 Melliera, Saussure, Ins. 326 Merismopteria, Jack and Etheridge, Moll. 94 Mesa, Saussure, Ins. 192 Mesarmadillo, Dollfuss, Crust. 28 Mesercs, Schewiakoff, Prot. 32 Mesocetus, Moreno, Mamm. 51 Mesochæta, Brauer and Bergenstamm, Ins. 299 Mesocrabro, Verhoeff, Ins. 192 Mesomycterus, Cope, Rept. 24 Mesopteryx, Hutton, Aves 25 Mesothyris, Warren, Ins. 276 Mesotrypes, Jacobsohn, Ins. 112 Metabacetus, Bates, Ins. 95 Metalopha, Staudinger, Ins. 257 Metapteryx, De Vis, Aves 25 Metaschalis, Hampson, Ins. 242 Metopella, Sars, Crust. 27 Metopodia, Brauer and Bergenstamm, Ins. 300 Metopotoms, Casey, Ins. 154 Metosamia, Druce, Ins. 242 Micractis, Warren, Ins. 276 Microbaris, Casey, Ins. 154 Microclæuodon, Scott, Mamm. 31 Microcrabro, Saussure, Ins. 192 Microcreagris, Balzan, Arachn. 34 Microcyllus, Thorell, Arachn. 32 Microjænia, Grassi, Prot. 30 Micromastix, Frenzel, Prot. 18 Micromerus, Guillebeau, Ins. 106 Micromphalina, Cossmann, Moll. 82 Micromyobia, Brauer and Bergen-stamm, Ins. 301

Microphana, Brauer and Bergenstamm, Ins. 299

284Mydrothauma, Butler, Ins. 242 Myinodes, Meyrick, Ins. 269 Myiophasia, Brauer and Bergenstamm, Ins. 300 Myrmoplasta, Gerstaecker, Ins. 307 Mysticoptera, Meyrick, Ins. 269 Myxexorista, Brauer and Bergenstamm, Ius. 299 Myxodexia, Brauer and Bergenstamm, Ins. 300

;

:

Microprion, Pictet and Saussure, Ins. 329 Microsebus, Kolbe, Ins. 169 Microtachina, Mik, Ins. 296 Mictodoca, Meyrick, Ins. 269 Mikrocodides, Bergendal, Vermes 87 Mila, Aurivillius, Ins. 242 Milione, Thorell, Arachn. 31 Mimastroides, Jacoby, Ins. 178 Mimoschinia, Warren, Ins. 276 Mimudea, Warren, Ins. 276 Minthodexia, Brauer and Bergen-stamm, Ins. 300 Mioacris, Pictet and Saussure, Ins. 330 Miomerocerus, Karsch, Ins. 310 Mionyx, Sanssure, Ins. 326 Miosaga, Saussure, Ins. 330 Mnesampela, Meyrick, Ins. 269 Mobianus, Giesbrecht, Crust. 31 Mochlotona, Meyrick, Ins. 269 Monacanthonyx, Bates, Ins. 95 Monarthropalpus, Rübsaamen, Ins. Monaster, Etheridge, Ech. 16 Monaulax, Nalepa, Arachu. 38 Monochilum, Schewiakoff, Prot. 32 Monocirsus, Cossmann, Moll. 75 Monocladum, Pic, Ins. 164 Monocona, Warren, Ins. 276 Monoculopsis, Sars, Crust. 26 Monophyaspis, Cope, Rept. 24 Monostola, Alphéraky, Ins. 257 Monotropia, Kirby, Ins. 242 Mormonomyia, Brauer and Bergenstamm, Ins. 301 Musæus, Thorell, Arachn. 32 Muscopteryx, Townsend, Ins. 296

Nanochlanius, Bates, Ins. 95 Narceus, Thorell, Arachn. 32 Newrasa, Staudinger, Ins. 242 Necrastur, De Vis, Aves 38 Nectochæta, Marenzeller, Vermes 46 Negalius, Casey, Ins. 144 Nemacerota, Hampson, Ins. 250 Nematopagurus, Edwards and Bouvier, Crust. 22 Nemetis, Moore, Ins. 220

Nemorilloides, Brauer ar stamm, Ins. 299 Nemosinus, Faust, Ins. 1 Neobagrus, Bellotti, Pisce Neoberlesia, Berlese, Arac Neobubastes, Blackburn, Neocalliphora, Brauer ar stamm, Ins. 301 Neocardia, Sowerby, Moll Neoclerus, Lewis, Ins. 13 Neoctodon, Bedel, Ins. 11 Neohydnus, Gorham, Ins. Neolesbia, Salvin, Aves 4 Neolythria, Alpheraky, In Neomintho, Brauer and stamm, Ins. 299 Neoteristis, Meyrick, Ins. Neotocerus, Duvivier, Ins. Neotractocera, Townsend,

Moll. 82 Nesace, Kirby, Ins. 243 Neusina, Göes, Prot. 32 Newmania, Swinhoe, Ins. Nicentrus, Casey, Ins. 154 Nicylla, Thorell, Arachn. Ninguta, Moore, Ins. 220 Niphograpta, Warren, Ins. Nirina, Weise, Ins. 178 Nocticola, Bolivar, Ins. 32 Nodepus, Gorham, Ins. 13 Norrisella, Cossmann, Mo Nossa, Kirby, Ins. 243 Notaspis, Warren, Ins. 27 Noteraula, Meyrick, Ins. 2 Notobrachypterus, Blackl 108

Neritaria, Wochrmann an

Notocetus, Moreno, Mami Notostemma, Bergendal, V Nuclearella, Frenzel, Prot. Nuclearina, Frenzel, Prot Nydia, Thorell, Arachn. 1 Nyoma, Duvivier, Ins. 165 Nyssonotus, Casey, Ins. 1:

Oberthueria, Kirby, Ins. 2 Oberthuria, Staudinger, II Ocala, Hulst, Ins. 277 Ocnerogyia, Staudinger, In Octocella, Helm, Bry. 4 Octoplasia, Brenske, Ins. Ecophiceus, Simon, Aracl Œdonia, Kirby, Ins. 244 Œhlertella, Hall and Brach. 6

Œnospila, Swinhoe, Ins. 2' Ogovia, Holland, Ins. 257 Ojedana, Distant, Ins. 306

Olenochæta, Townsend, Ins. 296 Oligodon, Amalizky, Moll. 93 Oligolochus, Casey, Ins. 154 Olisthanella, Voigt, Vermes 72 Ollochirus, Cope, Rept. 24 Omorphina, Alphéraky, Ins. 257 Oncodermus, Senna, Ins. 170 Oncopus, Cope, Rept. 24 Onychocryptus, Karsch, Arachn. 20 Onychora, Meyrick, Ins. 270 Oomorphidius, Casey, Ins. 154 Oopterinus, Casey, Ins. 154 Opetiopteryx, Blackburn, Ins. 154 Ophioteresis, Bell, Ech. 17 Ophoniscus, Bates, Ins. 96 Orcevia, Thorell, Arachn. 17 Orinonotus, Reuter, Ins. 311 Orna, Kirby, Ins. 244 Ornithoctonus, Pocock, Arachu. 28 Orophila, Krauss, Ins. 330 Orphnœcus, Simon, Arachn. 28 Orthaulaca, Weise, Ins. 179 Orthidium. Hall and Brach. 7 Orthochilus, Cossmann, Moll. 75 Orthocnemus, Milne-Edwards, Aves Orthocraspeda, Hampson, Ins. 244 Ortholophus, Oglivie-Grant, Aves 42 Orthoptychus, Futterer, Moll. 91 Orthotheca, Novák, Moll. 70 Orthotichia, Hall & Clarke, Brach. 7 Otaulax, Cossmann, Moll. 86 Otocryptus, Candèze, Ins. 128 Otomyrmex, Forel, Ins. 199 Oulobophora, Staudinger, Ins. 270 Ovengua, Distant, Ins. 307 Ovulactæon, Dall, Moll. 69 Oxyacrum, Cossmann, Moll. 71 Oxygoniola, Horn, Ins. 90 Oxyplax, Hampson, Ins. 244 Oxypleurites, Nalepu, Arachn. 38 Oxyrhabdium, Boulenger, Rept. 28 Oxysdonsaurus, Ambrosetti, Rept. 32 Oxytachina, Brauer and Bergenstamm, Ins. 300

Pachnephoptrus, Reitter, Ins. 179
Pachydasys, Butler, Ins. 258
Pachygnatha, Weise, Ins. 179
Pachygraphia, Brauer and Bergenstamm, Ins. 300
Pachyma, Weymer, Ins. 244
Pachymerola, Bates, Ins. 165
Pachymylus, Woodward, Pisces 35
Pachypalpa, Weise, Ins. 179
Pachypodagrus, Tepper, Ins. 330
Pagiophlœus, Faust, Ins. 155

Palæocasuarius, Forbes, Aves 25 Palæochrysopila, Meunier, Ins. 83 Palæocorax, Forbes, Aves 49 Palæoctenis, Meyrick, Ins. 270 Palæomutela, Amalizky, Moll. 93 Palaeonarica, Kittl, Moll. 85 Palæopelagus, De Vis, Aves 34 Palæosaccus, Hinde, Spong. 17 Palæotetrix, Shufeldt, Aves 27 Palembus, Casey, Ins. 138 Palpifer, Hampson, Ins. 244 Pandeus, Keyserling, Arachn. 25 Panthauma, Staudinger, Ins. 258 Pantorhytes, Faust, Ins. 155 Papua, Röber, Ins. 223 Paradidyma, Brauer and Bergenstamm, Ins. 300 Paradoria, Brauer and Bergenstumm, Ins. 299 Paradoxodon, Scott, Mamm. 32 Paragedia, Brauer and Bergenstamm, Ins. 299 Paragona, Staudinger, Ins. 258 Paragymnomma, Brauer and Bergenstamm, Ins. 301 Parahypocheta, Brauer and Bergenstamm, Ins. 299 Paralæa, Meyrick, Ins. 270 Paralispe, Brauer and Bergenstamm, Ins. 299 Paralucilia, Brauer and Bergenstamm, Ins. 301 Parameleus, Faust, Ins. 155 Paramesochæta, Brauer and Bergenstamm, Ins. 299 Paramorinia, Brauer and Bergenstamm, Ins. 300 Paramphibolia, Brauer and Bergenstamm, Ins. 301 Paramythia, De Vis, Aves 50 Paraneæra, Brauer and Bergen. stamm, Ins. 299 Parauemia, Heyden, Ins. 138 Parapallene, Carpenter, Crust. 33 Paraphilernus, Desbrochers, Ins. 155 Paraphysa, Simon, Arachn. 28 Paraplagia, Brauer and Bergenstamm, Ins. 299 Paraplinthus, Faust, Ins. 155 Pararicia, Brauer and Bergenstamm, Ins. 301 Pararistias, Robertson, Crust. 25 Parasarcophila, Brauer and Bergenstamm, Ins. 300 Parasetigena, Brauer and Bergen-stamm, Ins. 299

Paratachina, Brauer and Bergen-

stamm, Ins. 300

Paratænia, Kerremans, Ins. 125 Parathrylea, Durivier, Ins. 180 Paratrixa, Braver and Bergenstamm, Ins. 300 Paratryphera, Brauer and Bergenstamm, Ins. 298 Pardaliscoides, Stebbing, Crust. 27 Parcediceros, Sars, Crust. 26 Parthanosaurus, Skuph : Rept. 34 Passiena, Thorell, Arachn. 24 Patala, Moore, Ins. 220 Patanocnema, Karsch, Ins. 305 Patulopsis, Suter, Moll. 62 Pavieia, Brongniart, Ins. 165 Pegada, Moore, Ins. 220 Pelagia, Gumppenberg, Ins. 270 Pelecostoma, Reibisch, Moll. 63 Pelioderma, Bergroth, Ins. 305 Pelobrochis, Lucas, Ins. 244 Peltorhynchus, Thorell, Arachn. 32 Peltura, Perugia, Pisces 26 Pentachemus, Howard, Ins. 207 Pentacodon, Scott, Mamm. 31 Pentarthrinus, Casey, Ins. 155 Penthima, Kraatz, Ins. 121 Penthoplophora, Tepper, Ins. 330 Percolestus, Sloane, Ins. 96 Periaulax, Cossmann, Moll. 86 Periculodes, Sars, Crust. 26 Perissobasis, Reuter, Ins. 311 Peristichia, Dall, Moll. 84 Perrierella, Chevreux and Bouvier, Crust. 25 Petinops, Braner and Bergenstamm, Ins. 299 Petrostoma, Döderlein, Spong. 16 Pexomyia, Brauer and Bergenstamm, Ins. 298 Phænoderus, Péringuey, Ins. 156 Phalacra, Staudinger, Ins. 258 Pharcidella, Dall, Moll. 84 Pharmacophagus, Haase, Ins. 213 Phasiophana, Brauer and Bergenstamm, Ins. 301 Philænia, Kirby, Ins. 245 Philagria, Kirby, Ins. 245 Philiris, Röber, Ins. 223 Philudoria, Kirby, Ins. 245 Phitrus, Karsch, Ins. 326 Phlebohecta, Hampson, Ins. 245 Phorculus, Cossmann, Moll. 86 Phricta, Redtenbacher, Ins. 330 Phrissœcia, Simon, Arachn. 28 Phthorarcha, Meyrick, Ins. 271 Phyllomantis, Saussure, Ins. 326 Phyllomonas, K/ebs, Prot. 31 Phyllotribonia, Pictet and Saussure, Ins. 330

Phyllozelus, Pictet and Saussui Ins. 330 Physetonotus, Reuter, Ins. 311 Piarosoma, Hampson, Ins. 245 Piasus, Jacoby, Ins. 180 Pilemostoma, Desbrochers, Ins. 1 Pimeliopsis, Champion, Ins. 138 Pinnaspis, Cockerell, Ins. 318 Pirga, Aurivillius, Ins. 245 Placetron, Schalfeef, Crust. 22 Placilla, Moore, Ins. 220 Plæsiomys, Hall and Clarke, Brack Plagiobothrus, Karsch, Arachn. 2 Plagiocampa, Schewiakoff, Prot. ? Plagiochæta, Benham, Vermes 5. Plagiomima, Brauer and Berge stamm, Ins. 301 Plagiopholis, Boulenger, Rept. 27 Plagiprospherysa, Townsend, Ins. 2 Planadecta, Simon, Arachu. 28 Planctoplana, Graff, Vermes 72 Planolocha, Meyrick, Ins. 271 Platamartus, Reitter, Ins. 108 Platychilina, Woehrmann and Koken, Moll. 85 Platydasys, Butler, Ins. 258 Platylister, Lewis, Ins. 105 Platyopisthen, Roelofs, Ins. 156 Platytatus, Bergroth, Ins. 305 Platyzygæna, Swinhoe, Ins. 245 Plecteulota, Moellendorff, Moll. 6 Plectorthis, Hall and Brach, 7 Pleionogaster, Michaelsen, Verm 15 Plesiobaris, Casey, Ins. 156 Plesiocerithium, Cossmann, Moll. Plethocardia, Ulrich, Moll. 92 Pleurobranchillus, Bergh, Moll. 7 Pleuronota, Kraatz, Ins. 121 Plusilla, Standinger, Ins. 258 Podohuenia, Gavino, Crust. 17 Podotachina, Brauer and Berge stamm, Ins. 299 Pœcilogramma, Butler, Ins. 259 Pœcilomallus, Bates, Ins. 166 Peccilopsyra, Dohrn, Ins. 330 Pœcilosticta, Kraatz, Ins. 118 Polemistus, Saussure, Ins. 193 Polichus, Thorell, Arachn. 17 Polychærus, Mark, Vermes 72 Polycirsus, Cossmann, Moll. 79 Polymazus, Gerstaecker, Ins. 310 Polymierus, Faust, Ins. 156 Polynesia, Swinhoe, Ins. 271 Polypostia, Bergendal, Vermes 72 Polysyncraton, Nott, Tun. 6 Polytoechia, Hall & Clarke, Brach Pontomyxa, Topsent, Prot. 18 Pontosaurus, Gorjanovic - Kramberger, Rept. 25 Porphyriornis, Allen, Aves 31 Porrhothele, Simon, Arachn. 28 Porzanula, Frohawk, Aves 30 Potamanax, Pilsbry, Moll. 77 Præstilbia, Staudinger, Ins. 259 Preptos, Schaus, Ins. 246 Prionopaltys, Warren, Ins. 277 Prisma, Simroth, Moll. 66 Pristoneura, Reuter, Ins. 312 Proalligator, Ambrosetti, Rept. 32 Proboloptera, Meyrick, Ins. 271 Procampta, Holland, Ins. 225 Proctammodes, Blackburn, Ins. 113 Procyclotus, Fischer, Moll. 81 Prodasycnemis, Warren, Ins. 277 Progenetta, Déperet, Mamm. 29 Prolasius, Forel, Ins. 200 Promecotarsus, Casey, Ins. 156 Promilesia, Arribalzaga, Ins. 292 Propodinema, Ambrosetti, Rept. 23 Prorhynchops, Brauer and Bergenstamm, Ins. 300 Prosenoides, Braver and Bergenstamm, Ins. 300 Prosheliomyia, Brauer and Bergenstamm, Ins. 300 Prosopophora, Douglas, Ins. 318 Prosthecocotyle, Monticelli, Vermes Prosthenodon, Cossmann, Moll. 77 Protadia, Simon, Arachn. 19 Proteinidium, Bates, Ins. 166 Protochriacus, Scott, Mamm. 31 ?Protocimex, Moberg, Ins. 83 Protodus, Woodward, Pisces 37 Protogonodon, Scott, Mamm. 32 Protopaussus, Gestro, Ins. 103 Protorthis, Hall and Clarke, Brach. 7 Provipera, Kinkelin, Rept. 30 Proxyrodes, Blackburn, Ins. 156 Psalidopus, Wood - Mason Alcock, Crust. 24 Psammonyx, Döderlein, Prot. 20 Pselligmus, Simon, Arachn. 28 Pseudachlamys, Duvivier, Ins. 132 Pseudaclees, Faust, Ins. 156 Pseudaccenites, Kriechbaumer, Ins. 204 Pseudanchonus, Faust, Ins. 157 Pseudanthersea, Weymer, Ins. 246 Pseudantiora, Kirby, Ins. 246 Pseudaphelia, Kirby, Ins. 246 Pseudargynnis, Karsch, Ins. 218 Pseudarthromerus, Karsch, Arachn. 1892. [VOL. XXIX.]

Pseudasbolia, Kirby, Ins. 246 Pseudathyrma, Butler, Ins. 259 Pseudatractocera, Townsend, Ins. Pseudavicula, Juck and Etheridge, Moll. 94 Pseuderchia, Kirby, Ins. 246 Pseudhypsa, Kirby, Ins. 246 Pseudobarynotus, Desbrochers, Ins. Pseudocassida, Desbrochers, Ins. 180 Pseudochthonius, Balzan, Arachu. 34 Pseudocrania, Weise, Ins. 180 Pseudodexia, Brauer and Bergenstamm, Ins. 300 Pseudodiabrotica, Jacoby, Ins. 180 Pseudodiloma, Cossmann, Moll. 86 Pseudodinera, Brauer and Bergen-stamm, Ins. 300 Pseudofossarus, Koken, Moll. 78 Woehrmann and Pseudogermaria, Brauer and Bergenstamm, Ins. 299 Pseudoglessula, Bættger, Moll. 66 Pseudoheteronyx, Blackburn, Ins. Pseudoides, Jacoby, Ins. 181 Pseudojana, Hampson, Ins. 246 Pseudolatirus, Cossmann, Moll. 73 Pseudomalaia, Kraatz, Ins. 118 Pseudomaniola, Röber, Ins. 22) Pseudomicra, Butler, Ins. 259 Pseudomyothyria, Townsend, Ins. 297 Pseudoparlatoria, Cockerell, Ins. 319 Pseudopholis, Duvivier, Ins. 116 Pseudopiomera, Jacoby, Ins. 181 Pseudorhynchocyon, Filhol, Mamm. Pseudoscalites, Kittl, Moll. 75 Pseudoscopelus, Lütken, Pisces 23 Pseudotagma, Staudinger, Ins. 271 Pseudoviviania, Brauer and Bergenstamm, Ins. 298 Psiloderces, Simon, Arachn. 23 Psilosticha, Meyrick, Ins. 271 Psomus, Casey, Ins. 157 Pterygotheca, Novák, Moll. 70 Ptilodegeeria, Brauer and Bergenstamm, Ins. 300 Ptilolæmus, Ogilvie-Grant, Aves 42 Ptilotachina, Brauer and Bergenstamm, Ins. 299 Ptiloxena, Chapman, Aves 51 Ptycheulimella, Sacco, Moll. 84 Puer, Ortmann, Crust. 22 Pungtungia, Herzenstein, Pisces 27

Pustularia, Woehrmann and Koken, Moll. 77
Putlia, Moore. Ins. 220
Pycnobaria, Casey, Ins. 157
Pygetera, Kirby, Ins. 246
Pyracoderus, Guillebeau, Ins. 107
Pyrgodesmus, Pocock, Myr. 5
Pyrgolampros, Sacco, Moll. 84
Pytheas, Topsent, Spong. 22

Quassiremus, Jordan and Davis, Pisces 31

Rabula, Jordan and Davis, Pisces 31 Rachias, Simon, Arachn. 29 Raddea, Alpheraky, Ins. 259 Radulphius, Keyserling, Arachn. 20 Radsiella, Pilsbry, Moll. 89 Ræcius, Simon, Arachn. 33 Rafinesquina, Hall and Clarke. Brach. 7 Rancora, Smith, Ins. 259 Rangbia, Moore, Ins. 220 Rezia, Kirby, Ins. 246 Rhabderemia, Topsent, Spong. 22 Rhabdops, Boulenger, Rept. 27 Rhadinomerus, Faust, Ins. 157 Rhammatopoda, Redtenbacher, Ins. 330 Rhamphocolus, Casey, Ins. 157 Rhechosticha, Simon, Arachn. 29 Rhectocraspeda, Warren, Ins. 277 Rhicnodus, Fairmaire, Ins. 138 Rhinochrosis, Desbrochers, Ins. 157 Rhinomacquartia, Brauer and Bergenstamm, Ins. 300 Rhinoscopelus, Lütken, Pisces 26 Rhizorhina, Hausen, Crust. 31 Rhochmogaster, Karsch, Ins. 310 Rhodacanthis, Rothschild, Aves 52 Rhodareas, Kirby, Ins. 246 Rhodometra, Meyrick, Ins. 271 Rhopalomyia, Rübsaamen, Ins. 284 Rhoptria, Gumppenberg, Ins. 271 Rhynchomonas, Klebs, Prot. 31 Rhyparobia, Krauss, Ins. 326 Rhyparonotus, Faust, Ins. 157 Rivularia, Heude, Moll. 79 Roemerella, Hall and Clarke. Brach. 6 Rotania, Giesbrecht, Crust. 31 Rues, Casey, Ins. 138 Rusidrina, Staudinger, Ins. 259

Saccamœba, Frenzel, Prot. 18 Sagdellina, Cossmann, Moll. 62 Samuza, Keyserling, Arachn. 21

Rybakowia, Jacobsohn, Ins. 181

Sarcoclista, Townsend, Ins. 297 Sarcodexia, Townsend, Ins. 297 Sarcodidemnoides, Oka and Wille Tun. 6 Sarcomacronychia, Townsend, Is Sarcotachinella, Townsend, Ins. 2 Sarinda, Peckham, Arachn. 18 Sarothromyia, Brauer and Berge stamm, Ins. 300 Saurorhynchus, Reis, Pisces 33 Scalidognathus, Karsch, Arachn. Scelolethrus, Sémenow, Ius. 113 Schiosia, Boehm, Moll. 91 Schistomysis, Norman, Crust. 24 Schistosalius, Saussure, Ins. 194 Schizonotus, Casey, Ins. 157 Schizonotus, Reuter, Ins. 312 Scioglyptis, Meyrick, Ins. 271 Sciton, Blackburn, Ins. 117 Sclerocyphon, Blackburn, Ins. 129 Scodiomima, Staudinger, Ins. 272 Scotincecus, Simon, Arachn. 29 Scutotribonia, Pictet and Saussur Ins. 331

Sapinella, Kirby, Ins. 247

Sarax, Simon, Arachn. 14

Scytalichthys, Jordan and Davi Pisces 31 Sebastea, Kirby, Ins. 247 Seidlitzia, Desbrochers, Ins. 157 Selenomyia, Brauer and Bergel stamm, Ins. 300

Scymnomorpha, Blackburn, Ins. 18

Sematocera, Durrant, Ins. 281
Sematocera, Durrant, Ins. 281
Semiactwon, Cossmann, Moll. 69
Semiauricula, Cossmann, Moll. 67
Semichnoodes, Weise, Ins. 185
Semiterebellum, Cossmann, Moll. 75
Semiomima, Warren, Ins. 278
Semora, Peckham, Arachn. 18
Senex, Ortmann, Crust. 22
Sepidiopsis, Gestro, Ins. 138
Sepiomus, Aurivillius, Ins. 158
Sericocoris, Karsch, Ins. 308
Sericoplaga, Warren, Ins. 278
Sesiophaga, Brauer and Berge stamm, Ins. 301
Sidemia Standinger Ing. 259

stamm, Ins. 301
Sidemia, Staudinger, Ins. 259
Sigaretopsis, Cossmann, Moll. 82
Siglophora, Butler, Ins. 247
Silvanopsis, Grouvelle, Ins. 109
Simophorus, Faust, Ins. 158
Simopone, Forel, Ins. 200
Sinchula, Moore, Ins. 221
Sipalolasma, Simon, Arachn. 29
Siphoclytia, Townsend, Ins. 298

Siphophyto, Townsend, Ins. 298 Somabrachys, Kirby, Ins. 247 Somatrichia, Kirby, Ins. 247 Sophronopterus, Faust, Ins. 158 Spanioplon, Topsent, Spong. 22 Sparganophilus, Benham, Vermes 51 Sphærarthra, Weise, Ins. 181 Sphærococcus, Maskell, Ins. 319 Sphæromyxa, Thélohan, Prot. 29 Sphæropyle, Rüet, Prot. 25 Spheciospongia, Marshall, Spong. 22 Sphecodoptera, Hampson, Ins. 228 Spheniscosomus, Kruatz, Ins. 129 Sphragifera, Staudinger, Ins. 259 Spica, Sacco, Moll. 84 Spilopopillia, Kraatz, Ins. 118 Spinipalpa, Alphéraky, Ins. 259 Spinoza, Lewis, Ins. 132 Spintharina, Sémenow, Ins. 201 Spirocapsa, Rüst, Prot. 25 Spironema, Klebs, Prot. 31 Spirulirostrina, Canavari, Moll. 54 Spongia, Ogilvie, Spong. 24 Stasimopus, Simon, Arachn. 29 Stathmorrhopa, Meyrick, Ins. 272 Stegochilum, Schewiakoff, Prot. 32 Stenagenia, Saussure, Ins. 194 Stenancylus, Casey, Ins. 158 Stenochirus, Karsch, Arachn. 14 Stenochora, Warren, Ins. 278 Stenomeles, Warren, Ins. 278 Stenophylloides, Hampson, Ins. 248 Stenopyga, Karach, Ins. 326 Stenovalgus, Kolbe, Ins. 121 Stenygrocercus, Simon, Arachn. 29 Stephanocrates, Kolbe, Ins. 121 Stephanoon, Schewiakoff, Prot. 31 Sterrhochrotus, Simon, Arachn. 29 Sthenaropoda, Karsch, Ins. 331 Stibaroma, Meyrick, Ins. 272 Stichobasis, Kirby, Ins. 248 Sticthippus, Scudder, Ins. 332 Stictobaris, Casey, Ins. 158 Stigmosphærostylus, Rüst, Prot. 27 Stilbina, Stuudinger, Ins. 259 Streptomonas, Klebs, Prot. 31 Streptopteron, Swinhoe, Ins. 272 Strioturbonilla, Sacco, Moll. 84 Strobilidium, Schewiakoff, Prot. 32 Strobilæstrus, Brauer, Ins. 293 Stylinos, Topsent, Spong. 23 Stylogymnomyia, Brauer and Bergenstamm, Ins. 301 Styloneuria, Brauer and Bergenstamm, Ins. 300 Stylostichon, Topsent, Spong. 23 Sulcosubularia, Sacco, Moll. 84 Sulcoturbonilla, Sacco, Moll. 84

Sumnius, Weise, Ins. 185 Suteria, Pilsbry, Moll. 60 Sveltella, Cossmann, Moll. 72 Symmachia, Brenske, Ins. 117 Synapticola, Voigt, Crust. 31 Synarmadillo, Dollfuss, Crust. 28 Synchelidium, Sars, Crust. 26 Synegiodes, Swinhoe, Ins. 272 Syngonorthus, Butter, Ins. 272 Syntomoides, Hampson, Ins. 248 Synute, Dendy, Spong. 16 Systenope, Cossmann, Moll. 72

Tachinomima, Brauer and Bergen-

stamm, Ins. 300 Tachinomyia, Townsend, Ins. 298 Tachinophyto, Townsend, Ins. 298 Tachinoptera, Brauer and Bergenstamm, Ins. 298 Tachyornis, Milne-Edwards, Aves Tactoderus, Fuirmaire, Ins. 139 and *errutu* Tamphana, Schaus, Ins. 248 Tanaophysa, Warren, Ins. 278 Tantura, Kirby, Ins. 248 Tanyzonus, Marshall, Ins. 207
Tapinattus, Thorell, Arachn. 18
Tapinopus, Milne-Edwards, Aves Tectariopsis, Cossmann, Moll. 86 Tedinus, Casey, Ins. 141 Tegonotus, Nalepa, Arachn. 38 Telacodon, Marsh, Mamm. 53 Tenagogonus, Gerstaecker, Ins. 309 Tenesis, Duvivier, Ins. 139 Tephromyia, Brauer and Bergenstamm, Ins. 300 Terastiodes, Warren, Ins. 278 Terpnosia, Distant, Ins. 313 Teslasena, Fleutiaux, Ins. 129 Tetracladia, Howard, Ins. 207 Tetraclænodon, Scott, Mamm. 31 Tetragrapha, Brauer and Bergen-stamm, Ins. 299 Tetralepis, Bættger, Rept. 28 Tetraphleps, Hampson, Ins. 249 Tetrastyla, Schewiakoff, Prot. 32 Tetreurytarsa, Horn, Ins. 90 Teucer, Kirby, Ins. 249 Thalassohelix, Pilsbry, Moll. 69 Theclinesthes, Röber, Ins. 224 Thecocyrtella, Bittner, Brach. 7 Thelychæta, Brauer and Bergenstamm, Ins. 301 Thelymyia, Brauer and Bergenstamm, Ins. 299 Thesalia, Casey, Ins. 167

Thlæodon, Cope, Mamm. 54 Thoracites, Brauer and Bergenstamm, Ins. 300 Thrinaconyx, Saussure, Ins. 326 Thylakidium, Schewiakoff, Prot. 32 Thylakomonas, Schewiakoff, Prot.31 Thysanomyia, Brauer and Bergen-stamm, Ins. 299 Tiaporus, Cope, Rept. 23 Tiarella, Cossmann, Moll. 75 Tibetobia, Frivaldszky, Ins. 167 Ticera, Swinhoe, Ins. 249 Tigidia, Simon, Arachn. 29 Timus, Péringuey, Ins. 159 Tineomima, Staudinger, Ins. 281 Tmesiphantes, Simon, Arachn. 29 Tolypaspis, Schmidt, Pisces 35 Tomasina, Hall and Clurke, Brach. 6 Tortisipho, Cossmann, Moll. 73 Tosaura, Swinhoe, Ins 273 Totua, Keyserling, Arachn. 31 Toxoides, Hampson, Ins. 250 Trabaloides, Kirby, Ins. 249 Trachyschenium, Cossmann, Moll.75 Trachyzulpha, Dohrn, Ins. 331 Transvaalia, Distant, Ins. 308 Trematobolus, Matthew, Brach. 6 Trepobaris, Casey, Ins. 159 Tretospira, Woehrmann and Koken, Moll. 78 Triacetelus, Bates, Ins. 167 Trichæta, Swinhoe, Ins. 249 Trichauchenia, Warren, Ins. 278 Trichocarenum, Blackburn, Ins. 98 Trichocerota, Hampson, Ins. 228 Tricholimax, Frenzel, Prot. 18 Trichoreus, Blackburn, Ins. 185 Trichromia, Fairmaire, Ins. 167 Tricosemeia, Holland, Ins. 226 Tridentulus, Cope, Rept. 24 Trigonomonas, Klebs, Prot. 31 Trigonosomus, Faust, Ins. 159 Trilophomys, Depéret, Mamm. 35 Triplophleps, Hampson, Ins. 249 Trirhinopholis, Boulenger, Rept. 28 Trischistognatha, Warren, Ins. 279 Tristania, Kirby, Ins. 249 Tritomicrus, Fairmaire, Ins. 167 Trixoclista, Townsend, Ins. 298 Trophonina, Röber, Ins. 221 Turritodostomia, Sacco, Moll. 84 Tykonus, Michaelsen, Vermes 51 Tylochilus, Cossmann, Moll. 76 Tylopteryx, Hutton, Aves 25 Tympanoptera, Pictet and Saussure, Ins. 331 Typhlotriton, Steineger, Rept. 40

Uhlunga, Distant, Ins. 305
Undularia, Woehrmann and Kok
Moll. 77
Urgulania, Thorell, Arachn. 26
Urophagus, Klebs, Prot. 31
Usambius, Kolbe, Ins. 170
Utella, Dendy, Spong. 16

Valenus, Casey, Ins. 168 Venefica, Jordan and Davis, Pisces Vermiculus, Goodrich, Vermes 51 Virgo, Staudinger, Ins. 260 Viridonia, Bothschild, Aves 54 Vitalis, Fairmaire, Ins. 170 Vitrinoides, Simroth, Moll. 58 Vitriwebbins, Chapman, Prot. 21 Vosmaeropsis, Dendy, Spong. 16

Wateletia, Cossmann, Moll. 75 Weltina, Piersig, Arachn. 36

Xanthodiscus, Schewiakoff, Prot. Xenesthis, Simon, Arachn. 29 Xenogenius, Kolbe, Ins. 122 Xenomys, Merriam, Mamm. 35 Xenomystax, Jordan and Dau Pisces 31 Xenorthrius, Gorham, Ins. 132

Xenosyllides, Périaslavzeff, Vern 47 Xerospermophilus, Merriam, Mam

33 Xiphopheromyia, Bigot, Ins. 292 Xylomoia, Staudinger, Ins. 260 Xylotachina, Brauer and Berge

Yvesia, Topsent, Spong. 23

stamm, Ins. 299

Zamacra, Meyrick, Ins. 273 Zametopias, Thorell, Arachn. 32 Zatapina, Karsch, Arachn. 26 Zebeeba, Kirby, Ins. 249 Zeloticus, Péringuey, Ins. 98 Zeuzerodes, Pagenstecher, Ins. 279 Ziælas, Fairmaire, Ins. 139 Zonochroa, Brauer and Berge stamm, İns. 301 Zosteromyia, Brauer and Berge stamm, Ins. 300 Zschokkea, Kænike, Arachn. 36 Zuniga, Peckham, Arachn. 18 Zutulba, Kirby, Ins. 250 Zygænopsis, Swinhos, Ins. 250 Zygopleura, Woehrmann and Kok Moll. 77

### LIST OF THE PUBLICATIONS

OF THE

## ZOOLOGICAL SOCIETY OF LONDON.

THE scientific publications of the Zoological Society of London are of two kinds—"Proceedings," published in an octavo form, and "Transactions," in quarto.

According to the present arrangements, the "Proceedings" contain not only notices of all business transacted at the scientific meetings, but also all the papers read at such meetings and recommended to be published in the "Proceedings" by the Committee of Publication. A large number of coloured plates and engravings are attached to each annual volume of the "Proceedings," to illustrate the new or otherwise remarkable species of animals described in them. Amongst such illustrations, figures of the new or rare species acquired in a living state for the Society's Gardens are often given.

The "Proceedings" for each year are issued in four parts, on the first of the months of June, August, October, and April, the part published in April completing the volume for

the preceding year.

The "Transactions" contain such of the more important communications made to the scientific meetings of the Society as, on account of the nature of the plates required to illustrate them, are better adapted for publication in the quarto form.

They are issued at irregular intervals.

Fellows and Corresponding Members, upon payment of a Subscription of One Guinea before the day of the Anniversary Meeting in each year, are entitled to receive all the Society's Publications for the year. They are likewise entitled to purchase the Publications of the Society at 25 per cent. less than the price charged for them to the Public. A further reduction of 25 per cent. is made upon purchases of Publications issued prior to 1871, if they exceed the value of five pounds.

The following is a complete list of the publications of the Society already issued. They may be obtained at the Society's Office (3 Hanover Square, W.), at Messrs. Longmans', the Society's publishers (Paternoster Row, E.C.), or through any bookseller.

[August, 1893.]

TRANSACTIONS C	F THE ZOOLOGIC	AL SOCIETY OF LO
	4to. 12 vois. and 1	Price to Pri Fellows, I
		£ s. d. £
	g 59 Plates (1833-35)	
" <u>II</u> ., "	71 , (1835–41)	,
" III., "	63 " (1842–49) 77 " (1851–62)	,
", IV., ", ", ", ", ", ", ", ", ", ", ", ", ",	67 ' (1969-86)	
VI	92 ", (1866–69	, , , , , , , , , , , , , , , , , , , ,
", VII., ",	73 " (1869–72)	/ **** 22
" VIII., "	82 , (1872-74)	$() \dots 9 8 3 \dots 12$
IX	99 ,, (1875–77	$1 \cdots 12 1 6 \cdots 16$
", X., ",	95 ,, (1877–79	
Index, Vols. IX.	(1833-79	
VII.	ning 97 Plates (1880–85 65 (1886–90	,
~ <b>∀III</b> D+ 1	" A " (Tan 180	
D+ 9	", 6 ", (Apr. 189	.,
″ ″ D. 9	" 6 " (Oct. 189	
" " D+ ∧	" 1 " (Apr.189	2) 0 4 6 0
TD4 K	" 9 " (Feb. 189	
" " Pt. 6	" 4 " (June 189	
" " Pt. 7	" 6 " (Aug. 189	93) 0 9 0 <b>0</b>
PROCEEDINGS O	OF THE COMMIT	TEE OF SCIENC
CORRESPONI	_	
LONDON. 8v	o. 2 vols.	Price to Price Fellows. Pul
Part I. 1830-	-31. 1 vol. 8vo	4s. 6d 6e
" II. 1832.		4s. 6d 6e
		AT COCHEW OF TA
PROCEEDINGS O		
8vo.	<del>-</del>	(First Series.)
	Price to Price to the Fellows. Public.	Price t Fellow
Part I. 1833, 1 vol. 8ve		IX. 1841. 1 vol. 8vo. 4s. 6c
" II. 1834. "	4s. 6d 6s. ,	X. 1842. " 4s. 6
" III. 1835. "	$4s. 6d. \ldots 6s.$ ,	XI. 1843. ,, 4s. 6e
" IV. 1836. "	4s. 6d 6s. ,,	XII. 1844. " 4s. 6
" V. 1837. "		KIII. 1845. " 4s. 6
" VI. 1838. "		XIV. 1846. ", 4s. 6c
" VII. 1889. "	4s. 6d 6s.† ,	XV. 1847. " 4s. 6.
" VIII. 1840. "	4s. 6d 6s.†   Index :	1830–1847. " 4s. 6.
8vo.	13 vols. and Index.	(Second Series.)
	Letterpress only.	With Plates cole
	Price to Price to th	
	Fellows. Public.	Fellows. $\pounds$ s. $d$ .
Part XVI. 1848. 1 v	ol. 8vo. 4s. 6d 6s.	$\ldots \ldots$ $\tilde{1}$ $\overset{\circ}{0}$ $\overset{\circ}{8}$ $\ldots$
YVII 1840	,, 4s. 6d 6s.	1 0 8
", XVIII. 1850.		1 8 6
" XIX. 1851.	"	0 15 9
" XX. 1852.	$,, \qquad 4s. \ 6d. \qquad \qquad 6s.$	0 15 9
" XXI. 1853.	" 4s. 6d 6s.	0 18 0
" XXII. 1854.	$,, \qquad 4s. \ 6d. \qquad 6s.$	$\dots \dots 0 19 6 \dots$
" XXIII. 1855. XXIV 1856	,, 48. 6d 6s	$egin{array}{cccccccccccccccccccccccccccccccccccc$
" XXIV. 1856. " XXV. 1857.	4s. 6d 6s. $4s. 6d 6s.$	1 0 0
" YYVT 1959	" An Ra Ro	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
" XXVII. 1859.	" A. R.J. Q.	111 6
XXVIII. 1860.	,, 4s. 0d 6s.	8 III
1848-1860.	,, 48. 6d 6s.	
* No perfect co		† Out c

# PROCEEDINGS OF THE SCIENTIFIC MEETINGS OF THE LOOLOGICAL SOCIETY OF LONDON. 8vo. 30 vols. and 3 Indices.

Lette	erpress only.	With Plat	es uncoloured.	With Pl	ates coloured.
Price to Fellows.	Price to the Public.	Price to Fellows.	Price to the Public.	Price to Fellows.	Price to the Public.
1861 4s. 6d.	вв	9s.	12s	33s. 9d.	45s.
1862 4s. 6d.	вы	9 <b>s.</b>	128	33s. 9d.	458.
1863 4s. 6d.	вв	9 <i>s</i> .	128	33s. 9d.	45s.
1864 4s. 6d.		98.	128.†	33s. 9d.	45s.
1865 4s. 6d.			128	33s. 9d.	458.
1866 4s. Ud.	6s. <b>*</b>		12s.†	33s, 9d.	458.
			12s.*	33s. 9d.	458.
			128	33s. 9d.	458.
			, 12s	33s. 9d.	458.
			128	33s. 9d.	45s.
Index, 1861-187	70	4s. 6d.	68.		
			128.*	33s. 9d.	45s.
			12s.*	33s. 9d.	45e.†
	• • • • • • • • • • • •		128	33s. 9d.	45s.
			128.†	36s.	48s.†
	• • • • • • • • • • • • • • • • • • • •	• •	128	36s.	48.
	• • • • • • • • • • •		128	36s.	48.
	• • • • • • • • • • • • • • • • • • • •		128	36s.	48.
20,0	• • • • • • • • • • • • • • • • • • • •		128	36s.	48.
	•••••		128	36s.	48.
			128	36s.	48s.
Index, 1871–188			вя.	00	40
	• • • • • • • • • • • • • • • • • • • •		128	36s.	48s.
		_ `	128	30s.	48s.
	• • • • • • • • • • • • • • • • • • • •		128	36s.	48.
	• • • • • • • • • • • • • • • • • • • •	•	128	36s.	48s.
	• • • • • • • • • • • • • • • • • • • •		128	36s.	48.
	• • • • • • • • • • • • • • • • • • • •		128	36*.	48s.
	• • • • • • • • • • • • • • • • • • • •		128	36s.	48s.
		_	128	36s.	48s.
1000	•••••		128	36s.	48s.
	20	98.	12s.	36s.	48s.
Index, 1881–189	30	4s. 6d.	68.		

<sup>\*</sup> No perfect copies in stock.

## PROCEEDINGS OF THE GENERAL MEETINGS FOR SCIENTIFIC BUSINESS OF THE ZOOLOGICAL SOCIETY OF LONDON.

	Fellows.	
1891	. <b>3</b> 6s	48s.
1892		
1808, part 1 (Jan. & Feb.)	. 98	12s.
1893, , 2 (Mar. & Apr.)	. 9s	12s.
1893, " 3 (May & June)	. 9s	12s.

### LISTS OF THE ANIMALS IN THE SOCIETY'S GARDENS.

List of Vertebrated Animals Living in the Gardens of the Zoological Society of London. (First Edition.) 8vo. 1862. Price 1s. 6d.

List of Vertebrated Animals Living in the Gardens of the Zoological Society of London. (Second Edition.) 8vo. 1863. Price 1s. 6d.

List of Vertebrated Animals Living in the Gardens of the Zoological Society of London. (Third Edition.) 8vo. 1865. Price 1s. 6d.

List of Vertebrated Animals Living in the Gardens of the Zoological Society of London. (Fourth Edition.) 8vo. 1868. Price 1s. 6d.

<sup>†</sup> Out of print.

Revised List of the Vertebrated Animals now or lately Living in t Gardens of the Zoological Society of London. (Fifth Edition 8vo. 1872. Price:

Revised List of the Vertebrated Animals now or lately Living in t Gardens of the Zoological Society of London.—Supplementations Additions received in 1872, 1873, and 1874. 8 1875.

List of the Vertebrated Animals now or lately Living in the Garde of the Zoological Society of London. (Sixth Edition.) Close 8vo. 1877. Price 3s. 6

List of the Vertebrated Animals now or lately Living in the Garde of the Zoological Society of London. (Seventh Edition.) Clot 8vo. 1879. Price 3s. 6

List of the Vertebrated Animals now or lately Living in the Garde of the Zoological Society of London.—First Supplement, co taining Additions received in 1879. 8vo. 1880. Price 1s. 6

List of the Vertebrated Animals now or lately Living in the Garde of the Zoological Society of London. (Eighth Edition.) Clot 8vo. 1883.

Price 3s. 6

### THE ZOOLOGICAL RECORD.

The Zoological Record for the years 1864-1885. Twenty-tv volumes. Price £5 10

The Zoological Record for 1886; being Volume the Twenty-thi of the Record of Zoological Literature. Edited by Frank | Beddard, M.A., F.Z.S., Prosector and Davis Lecturer to tl Zoological Society of London. Cloth, 8vo. 1887. Price 10s., Ne

The Zoological Record for 1887; being Volume the Twenty-four of the Record of Zoological Literature. Edited by Frank I Beddard, M.A., F.Z.S., Prosector and Davis Lecturer to the Zoological Society of London. Cloth, 8vo. 1888. Price 10s., No.

The Zoological Record for 1888; being Volume the Twenty-fift of the Record of Zoological Literature. Edited by Frank 1 Beddard, M.A., F.Z.S., Prosector and Davis Lecturer to the Zoological Society of London. Cloth, 8vo. 1890. Price 10s., Ne

The Zoological Record for 1889; being Volume the Twenty-sixt of the Record of Zoological Literature. Edited by Frank 1 Beddard, M.A., F.Z.S., Prosector and Davis Lecturer to the Zoological Society of London. Cloth, 8vo. 1890. Price 10s., Ne

The Zoological Record for 1890; being Volume the Twenty-sevent of the Record of Zoological Literature. Edited by Frank 1 Beddard, M.A., F.Z.S., Prosector and Davis Lecturer to th Zoological Society of London. Cloth, 8vo. 1892. Price 10

The Zoological Record for 1891; being Volume the Twenty-eighth of the Record of Zoological Literature. Edited by DAVID SHAR. Esq., M.D., F.R.S., F.Z.S. Cloth, 8vo. 1892. Price 30


The Zoological Record for 1892; being Volume the Twenty-ninth of the Record of Zoological Literature. Edited by David Share Esq., M.D., F.R.S., F.Z.S. Cloth, 8vo. 1893. Price 30


Catalogue of the Library of the Zoological Society of London (Fourth Edition.) Cloth, 8vo. 1887. Price 4

These publications may be obtained at the Societ's Offic (3 Hanover Square, W.), at Messrs. Longmans' (Paternoster Ro E.C.), or through any bookseller.

•

•





STANFORD UNIVERSITY LIBRARIES STANFORD, CALIFORNIA 94305

